# A SIMPLE CONSTRUCTION OF THE DYNAMICAL  $\Phi_3^4$  MODEL

AUKOSH JAGANNATH AND NICOLAS PERKOWSKI

ABSTRACT. The  $\Phi_3^4$  equation is a singular stochastic PDE with important applications in mathematical physics. Its solution usually requires advanced mathematical theories like regularity structures or paracontrolled distributions, and even local well-posedness is highly nontrivial. Here we propose a multiplicative transformation to reduce the periodic  $\Phi_3^4$  equation to a well-posed random PDE. This leads to a simple and elementary proof of global well-posedness, which only relies on Schauder estimates, the maximum principle, and basic estimates for paraproducts, and in particular does not need regularity structures or paracontrolled distributions.

### 1. INTRODUCTION

In this note we prove the global-in-time existence and uniqueness of solutions to the  $\Phi_3^4$ equation, which is formally written as

<span id="page-0-0"></span>
$$
(\partial_t - \Delta + 1)\phi = -\phi^3 + \xi, \qquad (t, x) \in \mathbb{R}_+ \times \mathbb{T}^3,
$$
\n
$$
(1.1)
$$

where  $\xi$  is a space-time white noise on  $\mathbb{R} \times \mathbb{T}^3$  and  $\mathbb{T}^3 = (\mathbb{R}/\mathbb{Z})^3$  is the three-dimensional torus. The  $\Phi_3^4$  equation is one of the most prominent singular SPDEs and it has at least three important applications: Its invariant measure is given by the  $\Phi_3^4$  measure from quantum field theory, and as predicted by the stochastic quantization approach of [\[PW81\]](#page-15-0) we can use the SPDE to give a dynamic construction of this (highly nontrivial) measure [\[GH21\]](#page-14-0). The  $\Phi_3^4$  equation is also a universal model for 3d interface coexistence models near bifurcations [\[HX18,](#page-14-1) [FG19\]](#page-14-2), and it is expected to describe the dynamics of 3d ferromagnets close to their critical temperature [\[MW17a\]](#page-14-3).

The rigorous interpretation and local well-posedness of the  $\Phi_3^4$  equation had been an open problem for a long time, because the solution  $\phi(t)$  is, for each  $t > 0$ , a distribution in the space variable and therefore the interpretation of the term  $-\phi^3$  is unclear. With the development of new mathematical theories such as regularity structures [\[Hai14\]](#page-14-4) or paracontrolled distributions [\[GIP15,](#page-14-5) [CC18\]](#page-14-6), for example, the rigorous construction of  $-\phi^3$  became possible and this lead to a local existence-uniqueness result. Because of the nonlinearity of the equation, global existence is more subtle and it was first shown by Mourrat and Weber [\[MW17b\]](#page-14-7), who even proved a "coming down from infinity" estimate, i.e., a bound for  $\phi(t)$  at  $t > 0$  which is independent of the initial condition. In recent years simpler proofs have been obtained, which even allow to solve the equation on  $\mathbb{R}_+ \times \mathbb{R}^3$  instead of  $\mathbb{R}_+ \times \mathbb{T}^3$ ; see [\[GH19,](#page-14-8) [MW20\]](#page-14-9).

All of these results (even local existence) require sophisticated mathematical tools. Here we suggest a sort of inverse Cole-Hopf transform, previously employed in the context of linear equations for example in [\[HL15,](#page-14-10) [GP17\]](#page-14-11), which leads to a short and relatively elementary proof of global well-posedness of the periodic  $\Phi_3^4$  equation. In particular, we do not need paracontrolled distributions or regularity structures.

In the next section we introduce the transform and we prove the local well-posedness of the transformed equation. Section [3](#page-4-0) contains the proof of global well-posedness, which is similar to the arguments by Gubinelli and Hofmanová for the  $\Phi_2^4$  equation [\[GH19\]](#page-14-8); roughly speaking, the transform reduces the singularity of the equation and brings  $\Phi_3^4$  to the same level of difficulty as the much easier  $\Phi_2^4$  equation. In Section [4](#page-10-0) we discuss possible extensions of our results.

*Notation:* In the following, for two functions we say that  $f \leq_a g$  if there is a constant  $C(a) > 0$  depending only on a such that  $f \leq C(a)g$ . We say that  $f \simeq g$  if  $f \lesssim g$  and  $g \lesssim f$ .

# 2. Local Solutions

<span id="page-1-1"></span>We begin by proving the local well-posedness of the  $\Phi_3^4$  equation. Recall that the  $\Phi_3^4$ equation is to be understood in a renormalized sense. For this reasons, in place of [\(1.1\)](#page-0-0), the equation is sometimes written formally as  $(\partial_t - \Delta + 1)\phi = -\phi^3 + \infty \cdot \phi + \xi$ , where the  $\infty$ denotes a divergent counter-term.

More precisely, let  $\mathcal{L} = \partial_t - \Delta + 1$  and let Z be stationary such that

$$
\mathcal{L}Z=\xi,
$$

for a two-sided space-time white noise  $\xi$  on  $\mathbb{R}_+ \times \mathbb{T}^3$  and let  $\mathcal{C}^{\alpha}$  denote the Hölder-Besov space of regularity  $\alpha$ . (See [\(2.2\)](#page-2-0) below for a precise definition of this space.) Stated rigorously, our goal is to understand the sequence of solutions to

$$
\begin{cases} (\partial_t - \Delta + 1)\phi_\delta = -\phi_\delta^3 + (3a_\delta - 3b_\delta)\phi_\delta + \xi_\delta, & (t, x) \in \mathbb{R}_+ \times \mathbb{T}^3, \\ \phi_\delta(0, x) = Z_\delta(0) + \phi_0^\sharp, & (t, x) \in \mathbb{R} \end{cases}
$$

where  $\phi_0^{\sharp} \in C^{3/2-\epsilon}$  for some sufficiently small  $\epsilon > 0$ ,  $\xi_{\delta}$  is a suitable mollification of  $\xi$ , and  $a_{\delta}, b_{\delta}$  are (deterministic) sequences of scalars which diverge for  $\delta \to 0$ ; also,  $\mathcal{L}$  is stationary with  $\mathcal{L}Z_{\delta} = \xi_{\delta}$ . Our goal is to understand the limit  $\phi = \lim_{\delta \to 0} \phi_{\delta}$ , which we call the renormalization limit.

Since the renormalization procedure is present at every step of our analysis, we will suppress the dependencies on  $\delta$  for the sake of readability. For example, we will simply write that we seek the solution to the equation

<span id="page-1-0"></span>
$$
\begin{cases} (\partial_t - \Delta + 1) \phi = -\phi^3 + (3a - 3b)\phi + \xi, & (t, x) \in \mathbb{R}_+ \times \mathbb{T}^3, \\ \phi(0, x) = Z(0) + \phi_0^{\sharp}. \end{cases}
$$
(2.1)

Here and in the following, we use the convention that whenever we write an equation or inequality, the expression is to be understood as holding for all  $\delta$ , with any unquantified constants appearing in bounds being independent of  $\delta$ .

In order to develop solutions to this equation, we need to understand certain terms built from Z which we will refer to as the stochastic tree terms. To define these terms, we recall the following. Let  $\mathscr{S}'$  denote the space of tempered distributions on  $\mathbb{T}^3$ , and let  $\Delta_j$  denote the usual Littlewood-Payley operators [\[BCD11\]](#page-14-12). Consider the sum  $\Delta_{\leq j} = \sum_{i \leq j} \Delta_i$ , and define  $\Delta_{\geq j}, \Delta_{\geq j}$ , and  $\Delta_{\leq j}$  similarly. For  $u, v \in \mathscr{S}'$ , consider the paraproduct

$$
u \otimes v = \sum_{j \ge -1} \Delta_{\le j-2} u \Delta_j v
$$

		$\frac{1}{2} \alpha_{\tau}$ $\left[-\frac{1}{2} - \epsilon \right] - 1 - \epsilon$ $\frac{1}{2} - \epsilon$ $\left[1 - \epsilon \right] - \epsilon$		

<span id="page-2-1"></span>TABLE 1. Regularity of stochastic objects.

and the resonant product

$$
u \odot v = \sum_{i,j:|i-j| \leq 1} \Delta_i u \Delta_j v.
$$

We denote their sum by  $u \preccurlyeq v = u \otimes v + u \odot v$ .

We consider the following nonlinear terms built from  $Z$ :

$$
\llbracket Z^2 \rrbracket := Z^2 - a, \quad \mathcal{L}Z^{\mathbf{V}} = \llbracket Z^3 \rrbracket := Z^3 - 3aZ, \qquad \mathcal{L}Z^{\mathbf{V}} = \llbracket Z^2 \rrbracket,
$$
  

$$
Z^{\mathbf{V}} \odot Z, \qquad Z^{\mathbf{V}} \odot \llbracket Z^2 \rrbracket - \frac{b}{3}, \qquad |\nabla Z^{\mathbf{V}}|^2 - \frac{b}{3}, \qquad Z^{\mathbf{V}} \odot \llbracket Z^2 \rrbracket - bZ,
$$

where  $Z^{\vee}$  and  $Z^{\Psi}$  have initial condition 0:  $Z^{\vee}(0) = Z^{\Psi}(0) = 0$ . We remind the reader that these terms all implicitly depend on  $\delta$ , e.g., the terms  $Z, a, b$  should be understood as  $Z_{\delta}, a_{\delta}, b_{\delta}$ . (For fixed  $\delta > 0$  the terms a and b in the above expressions are the same as those in  $(2.1).$  $(2.1).$ 

Using Gaussian analysis, it can be shown that each of these terms converges as  $\delta \to 0$ to a limit that does not depend on the specific mollification used for  $\xi_{\delta}$ . As the existence of the limits is implicit in any solution theory to the  $\Phi_3^4$  equation, we do not prove this here. Instead, we refer the reader to [\[CC18\]](#page-14-6) or, for a pedagogical exposition, [\[MWX17\]](#page-15-1). While the term  $|\nabla Z^{\vee}|^2 - \frac{b}{3}$  $\frac{b}{3}$  does not appear in the references, it is (basically) equivalent to  $Z^{\vee} \odot [Z^2] - \frac{b}{3}$  $\frac{b}{3}$ ; for the convenience of the reader, we discuss this in the appendix. The regularities of these terms are given in Table [1,](#page-2-1) which should be read as  $Z_{\delta} \to Z \in C_T C^{-\frac{1}{2}-\epsilon}$ ,  $\llbracket Z_{\delta}^2 \rrbracket \to \llbracket Z^2 \rrbracket \in C_T C^{-1-\epsilon}$ , etc., and we can take any  $\epsilon > 0$ .<br>We work with the following function spaces:

We work with the following function spaces:

<span id="page-2-0"></span>
$$
||u||_{\mathcal{C}^{\alpha}} = ||(2^{j\alpha}||\Delta_j u||_{L^{\infty}})_{j\geq -1}||_{\ell^{\infty}},
$$
\n(2.2)

for  $\alpha \in \mathbb{R}$  and  $C_T C^{\alpha} := C([0,T], \mathcal{C}^{\alpha})$  with  $||u||_{C_T \mathcal{C}^{\alpha}} := \max_{t \in [0,T]} ||u(t)||_{\mathcal{C}^{\alpha}}$ .

The usual approach to solving the  $\Phi_3^4$  equation starts by considering  $v = \phi - Z + Z^{\Psi}$ . Then we obtain a new equation for v, which is better behaved than the equation for  $\phi$  because by subtracting  $Z - Z^{\mathbf{v}}$  we removed the most singular terms. However, the equation for v is still ill-posed because it involves the singular product  $-3v\llbracket Z^2 \rrbracket$ . To deal with this product,<br>provious works used requierity structures or paracentrolled distributions. previous works used regularity structures or paracontrolled distributions.

Here we propose an alternative approach, by observing that the singular product  $-3v\llbracket Z^2 \rrbracket$ <br>n be removed with a multiplicative transform: We consider لا<br>ا can be removed with a multiplicative transform: We consider

$$
v = e^{3Z^{\vee}} \left( \phi - Z + Z^{\Psi} \right).
$$

Then, formally,  $\phi$  solves [\(2.1\)](#page-1-0) if and only if v solves

$$
\mathcal{L}v = \left(-3Z^{\vee} + 9\left(|\nabla Z^{\vee}|^2 - \frac{b}{3}\right)\right)v - 6(\nabla Z^{\vee} \cdot \nabla v - be^{3Z^{\vee}}Z^{\Psi}) \n+ e^{3Z^{\vee}} \left[3Z^{\Psi}\llbracket Z^2\rrbracket - 3b\left(Z + Z^{\Psi}\right) - 3Z\left(e^{-3Z^{\Psi}}v - Z^{\Psi}\right)^2 - \left(e^{-3Z^{\Psi}}v - Z^{\Psi}\right)^3\right],
$$
\n(2.3)

with initial data  $v(0) = \phi(0) - Z(0) = \phi_0^{\sharp}$  $\frac{1}{0}$ . This equation is still ill-posed, because of the singular term  $3e^{3Z\vee}Z^{\vee}[[Z^2]]$  on the right hand side. To remove it, we consider

<span id="page-3-0"></span>
$$
\mathcal{L}Y = 3e^{3Z^{\vee}}(Z^{\mathbf{v}}[[Z^2]] - b(Z + Z^{\mathbf{v}})), \quad Y(0) = 0.
$$

We show in the appendix that  $Y \in \bigcap_{\epsilon > 0} C_T \mathcal{C}^{1-\epsilon}$  can be constructed as a continuous function of the stochastic tree terms described above. Let then  $u = v - Y$ , which solves

$$
\mathcal{L}u = \left(-3Z^{\vee} + 9\left(|\nabla Z^{\vee}|^2 - \frac{b}{3}\right)\right)(u+Y) - 6\nabla Z^{\vee} \cdot \nabla u - 6(\nabla Z^{\vee} \cdot \nabla Y - be^{3Z^{\vee}}Z^{\vee})
$$
  

$$
-e^{3Z^{\vee}}\left[3Z\left(e^{-3Z^{\vee}}(u+Y) - Z^{\vee}\right)^2 + \left(e^{-3Z^{\vee}}(u+Y) - Z^{\vee}\right)^3\right],
$$
\n(2.4)

with initial condition  $u(0) = v(0) = \phi_0^{\sharp}$  $\frac{1}{0}$ . We show in the appendix that all the explicit (renormalized) nonlinear functions of the stochastic tree terms that appear on the right hand side can be constructed as continuous functions of the given trees. Expanding the nonlinearities, we can then rewrite the equation for  $u$  as

<span id="page-3-1"></span>
$$
\mathcal{L}u = -6\nabla Z^{\vee} \cdot \nabla u - e^{-6Z^{\vee}} u^3 + Z_2 u^2 + Z_1 u + Z_0, \tag{2.5}
$$

for some  $Z_0, Z_1$ , and  $Z_2$  that are in  $C_T \mathcal{C}^{-1/2-\epsilon}$  for every  $T, \epsilon > 0$ .

After these preparations, the following local existence/uniqueness result is easy. For the remainder of the paper, we fix  $\epsilon > 0$  which is small enough so that all of the following arguments work ( $\epsilon < 1/6$  is sufficient).

<span id="page-3-4"></span>**Theorem 2.1.** Assume that  $u(0) = \phi_0^{\sharp} \in C^{3/2-\epsilon}$ . There exists  $T^* > 0$  such that for all  $T < T^*$  the equation [\(2.4\)](#page-3-0) has a unique solution  $u \in C_T C^{3/2-\epsilon}$ , and this solution depends continuously on  $\mathbb{Z} = (Z^{\vee}, Z_0, Z_1, Z_2) \in \bigcap_{\epsilon > 0} (C_T \mathcal{C}^{-\epsilon'} \times (C_T \mathcal{C}^{-1/2 - \epsilon'})^3)$  and  $\phi_0^{\sharp} \in \mathcal{C}^{3/2 - \epsilon}$ . Thus, in particular, its limit for  $\delta \to 0$  exists and is independent of the specific mollification used for  $\xi_{\delta}$ . Moreover, if  $T^* < \infty$ , then  $\lim_{t \uparrow T^*} ||u(t)||_{\mathcal{C}^{3/2-\epsilon}} = \infty$ .

Before we get to the proof, recall that for  $\alpha + \beta > 0$ , the map  $(u, v) \mapsto uv$  extends to a bounded linear form on  $\mathcal{C}^{\alpha} \times \mathcal{C}^{\beta} \to \mathcal{C}^{\alpha \wedge \beta}$ , see [\[BCD11\]](#page-14-12). In particular,

<span id="page-3-3"></span>
$$
||uv||_{\mathcal{C}^{\alpha \wedge \beta}} \le ||u||_{\mathcal{C}^{\alpha}} \cdot ||v||_{\mathcal{C}^{\beta}}.
$$
\n
$$
(2.6)
$$

Moreover,  $\nabla$  is a bounded linear operator from  $\mathcal{C}^{\alpha}$  to  $\mathcal{C}^{\alpha-1}$  for any  $\alpha \in \mathbb{R}$  [\[BCD11\]](#page-14-12).

Also, let  $(P_t)_{t>0}$  be the semigroup generated by  $\mathcal L$ . Recall the following estimate for  $(P_t)_{t\geq0}$  [\[GIP15,](#page-14-5) [GH19\]](#page-14-8): for  $\gamma\geq0$  and  $\alpha\in\mathbb{R}$ 

<span id="page-3-2"></span>
$$
||P_t w||_{\mathcal{C}^{\alpha+\gamma}} \lesssim e^{-t} t^{-\gamma/2} ||w||_{\mathcal{C}^{\alpha}} \leq t^{-\gamma/2} ||w||_{\mathcal{C}^{\alpha}}, \qquad t \geq 0. \tag{2.7}
$$

*Proof.* We work with the formulation [\(2.5\)](#page-3-1). The regularity of  $\nabla Z^{\vee}$  is  $C^{-\epsilon}$  and therefore  $-6\nabla Z^{\vee} \cdot \nabla u$  is well-defined if  $u \in C^{\alpha}$  for  $\alpha > 1 + \epsilon$ . Similarly,  $Z_1, Z_2$  have regularity  $\mathcal{C}^{-1/2-\epsilon/2}$ , and therefore the remaining products are well-defined if  $u \in \mathcal{C}^{\alpha}$  for  $\alpha > 1/2 + \epsilon/2$ . In that case, the right hand side of [\(2.5\)](#page-3-1) will be in  $C^{-1/2-\epsilon/2}$ . By Schauder estimates, we then expect that we can take  $\alpha = 3/2 - \epsilon$ , and thus the PDE is locally well-posed. We make this precise by a standard Picard iteration argument as follows.

Consider the map

$$
F(u)(t) = P_t \phi_0^{\sharp} + \int_0^t P_{t-s} \left( -6\nabla Z^{\vee} \cdot \nabla u - e^{-6Z^{\vee}} u^3 + Z_2 u^2 + Z_1 u + Z_0 \right)(s) ds.
$$

Applying the semigroup estimate [\(2.7\)](#page-3-2) with  $\alpha = -1/2 - \epsilon/2$  and  $\gamma = 2 - \epsilon/2$ , we have for any  $u, v \in C_T C^{3/2 - \epsilon}$ ,

$$
||F(v) - F(u)||_{C_T C^{3/2-\epsilon}}
$$
  
\n
$$
\lesssim T^{\epsilon/4} \cdot ||-6\nabla Z^{\vee} \cdot \nabla (v - u) - e^{-6Z^{\vee}} (v^3 - u^3) + Z_2(v^2 - u^2) + Z_1(v - u)||_{C_T C^{-\frac{1}{2} - \frac{\epsilon}{2}}}
$$
  
\n
$$
\lesssim_{\mathbb{Z}} T^{\epsilon/4} (1 + ||v||_{C_T C^{3/2-\epsilon}}^2 + ||u||_{C_T C^{3/2-\epsilon}}^2) ||v - u||_{C_T C^{3/2-\epsilon}},
$$

where in the second line we used the product estimate [\(2.6\)](#page-3-3). Let  $M = 2||\phi_0^{\sharp}||_{\mathcal{C}^{3/2-\epsilon}}$ . Then for  $T > 0$  sufficiently small (depending only on M and  $\mathbb{Z}$ ), the map F is a contraction on the ball of radius 2M in  $C_T \mathcal{C}^{3/2-\epsilon}$ . The fixed point is then our desired solution on [0, T]. By iterating this construction on  $[T, T + T']$ , for  $T' > 0$  possibly smaller than T, etc., we can extend the solution up to a possibly finite explosion time  $T^*$ .

The continuous dependence on the data is shown with similar arguments: For  $\mathbb{Z}, \mathbb{Z}$  and  $\phi_0^{\sharp}$  $\phi_0^{\sharp}$ ,  $\tilde{\phi}_0^{\sharp}$ , we consider the fixed points  $u, \tilde{u}$  of the maps  $F_{\mathbb{Z}, \phi_0^{\sharp}}$ ,  $F_{\mathbb{Z}, \tilde{\phi}_1^0}$  and write  $u - \tilde{u} = F_{\mathbb{Z}, \phi_0^{\sharp}}(u) F_{\tilde{Z},\tilde{\phi}_{\sharp}^{0}}(\tilde{u})$  to bound  $u-\tilde{u}$  in terms of  $\mathbb{Z}-\tilde{\mathbb{Z}}$  and  $\phi_{0}^{\sharp}-\tilde{\phi}_{0}^{\sharp}$ . — Первый процесс в серверності процесс в серверності процесс в серверності процесс в серверності процесс в с<br>Процесс в серверності процесс в серверності процесс в серверності процесс в серверності процесс в серверності

#### 3. Global Solutions: The Gubinelli-Hofmanova Approach

<span id="page-4-0"></span>The goal of this section is to prove the following theorem:

<span id="page-4-1"></span>**Theorem 3.1.** Let u,  $T^*$  and  $\epsilon$  be as in Theorem [2.1](#page-3-4) and let  $T < T^* \wedge 1$ . There exists a constant  $C > 0$  that depends only on  $\phi_0^{\sharp}$  and  $\mathbb{Z} = (Z^{\vee}, Z_0, Z_1, Z_2)$  but not on T, such that

$$
||u||_{C_T\mathcal{C}^{3/2-\epsilon}} \leq C.
$$

In particular,  $T^* = \infty$ .

We remind the reader that here (and in the subsequent)  $Z_0, Z_1, Z_2$  are as in [\(2.5\)](#page-3-1).

For readability, in the remainder of this section we will suppress the dependence of norms on time when it is clear from context, and we simply write  $||u||_{\alpha} = ||u||_{C_T C^{\alpha}}$ , with the convention that  $||u||_0 = ||u||_{C_T L^{\infty}}$ .

3.1. A Paraproduct decomposition. For  $n \geq 1$  to be determined later, let us consider the coupled system of equations

$$
(\mathcal{L} + 6\nabla Z^{\vee} \otimes \nabla) u_1 = U_1(u_1, u_2)
$$
  

$$
(\mathcal{L} + 6\nabla Z^{\vee} \cdot \nabla) u_2 = -e^{-6Z^{\vee}} u_2^3 + U_2(u_1, u_2),
$$

with initial conditions  $u_1(0) = u(0) = \phi_0^{\sharp}$  $_{0}^{\sharp}$  and  $u_{2}(0)=0$ , where

$$
U_1(u_1, u_2) = u^2 \otimes \Delta_{>2n} Z_2 + u \otimes \Delta_{>n} Z_1 + Z_0 \tag{3.1}
$$

<span id="page-5-1"></span>
$$
U_2(u_1, u_2) = -6\nabla u_1 \succcurlyeq \nabla Z^{\vee} - e^{-6Z^{\vee}} (u^3 - u_2^3) \tag{3.2}
$$

$$
+ u^2 \otimes \Delta_{\leq 2n} Z_2 + u \otimes \Delta_{\leq n} Z_1 + u^2 \succcurlyeq Z_2 + u \succcurlyeq Z_1,
$$
 for  $u := u_1 + u_2$ . Note here that we have implicitly suppressed the dependence of  $u_1$  and  $u_2$  on n. The idea is that  $U_1$  collects all the singular contributions, while  $U_2$  is more regular and

on n. The idea is that  $U_1$  collects all the singular contributions, while  $U_2$  is more regular and in fact in  $L^{\infty}$ , so that we can apply the maximum principle to deal with it. Note that  $u_1 + u_2$ solves the equation [\(2.5\)](#page-3-1), so by the uniqueness proved in Theorem [2.1](#page-3-4) we have  $u_1 + u_2 = u$ . We begin by estimating  $U_1$  and  $U_2$ :

<span id="page-5-0"></span>**Lemma 3.2.** For any  $\delta \geq 0$ ,  $n \geq 1$ ,  $\epsilon' > 0$ , we have

$$
||U_1||_{-1/2-\epsilon-\delta} \lesssim \mathbb{Z} \left(1 + 2^{-n\delta}||u_1 + u_2||_0\right)^2.
$$
  

$$
||U_2||_{-1/2-\epsilon} \lesssim \mathbb{Z} \sum_{i=1}^3 ||u_1||_0^i \cdot ||u_2||_0^{3-i} + ||u_1||_{1+\epsilon'} + ||u_1 + u_2||_{1/2+\epsilon'} (1 + ||u_1 + u_2||_0).
$$

Furthermore, for any  $\kappa > 1/2$ ,  $n \ge 1$ ,  $\epsilon' > 0$ 

$$
||U_2||_0 \lesssim \sum_{i=1}^3 ||u_1||_0^i \cdot ||u_2||_0^{3-i} + ||u_1||_{1+\epsilon'} + ||u_1 + u_2||_{1/2+\epsilon'} (1 + ||u_1 + u_2||_0)
$$
  
+  $2^{n\kappa} ||u_1 + u_2||_0 + 2^{2n\kappa} ||u_1 + u_2||_0^2$ .

Before turning to the proof of this result, we first recall the following useful estimate on Littlewood-Payley blocks which follows directly from the definition: For any  $\alpha \in \mathbb{R}$ ,  $f \in \mathcal{C}^{\alpha}$ and  $\delta \geq 0$ ,

<span id="page-5-3"></span>
$$
\|\Delta_{>n}f\|_{\mathcal{C}^{\alpha-\delta}} \lesssim 2^{-n\delta} \|f\|_{\mathcal{C}^{\alpha}} \qquad \|\Delta_{\leq n}f\|_{\mathcal{C}^{\alpha+\delta}} \lesssim 2^{n\delta} \|f\|_{\mathcal{C}^{\alpha}}.\tag{3.3}
$$

We will also require the following result, which we will refer to as the paraproduct estimates.

<span id="page-5-2"></span>**Lemma 3.3** (Paraproduct Estimates, [\[BCD11\]](#page-14-12)). For all  $\beta \in \mathbb{R}$ ,  $\alpha < 0$ , and  $u, v \in \mathscr{S}'$ , the paraproduct satisfies

$$
||u \otimes v||_{\mathcal{C}^{\beta}} \lesssim ||u||_{L^{\infty}} ||v||_{\mathcal{C}^{\beta}}
$$
  

$$
||u \otimes v||_{\mathcal{C}^{\alpha+\beta}} \lesssim ||u||_{\mathcal{C}^{\alpha}} ||v||_{\mathcal{C}^{\beta}}.
$$

Furthermore, if  $\alpha + \beta > 0$ , then the resonant product satisfies

 $\lesssim_{\mathbb{Z},\epsilon} (1+2^{-n\delta}||u_1+u_2||_0)^2,$ 

 $||u \odot v||_{\mathcal{C}^{\alpha+\beta}} \lesssim ||u||_{\mathcal{C}^{\alpha}} ||v||_{\mathcal{C}^{\beta}},$ 

and if  $\beta > 0$  also

$$
||u \odot v||_{\mathcal{C}^{\beta}} \lesssim ||u||_{L^{\infty}} ||v||_{\mathcal{C}^{\beta}}.
$$

With these estimates in hand, we may now prove the above lemma.

**Proof of Lemma [3.2](#page-5-0)**. Let us begin with the first bound, for  $U_1 = u^2 \otimes \Delta_{\geq 2n} Z_2 + u \otimes \Delta_{\geq 2n} Z_1 + Z_2$ . By (3.1) and the paraproduct estimates (Lemma 3.3)  $\Delta_{>n}Z_1 + Z_0$ . By [\(3.1\)](#page-5-1) and the paraproduct estimates (Lemma [3.3\)](#page-5-2),  $||U_1||_{-1/2-\epsilon-\delta} \lesssim ||u_1+u_2||_0^2 \cdot ||\Delta_{>2n} Z_2||_{-1/2-\epsilon-\delta} + ||u_1+u_2||_0 \cdot ||\Delta_{>n} Z_1||_{-1/2-\epsilon-\delta} + ||Z_0||_{-1/2-\epsilon-\delta}$  where in the second line we used the first estimate from [\(3.3\)](#page-5-3).

Let us now turn to the bound for  $U_2$ . Since  $Z_2, Z_1 \in C_T C^{-1/2-\epsilon'/2}$ , we may apply the paraproduct estimates and the embedding  $C_T \mathcal{C}^{\epsilon'/2} \subset C_T L^{\infty}$  to find

$$
||u^2 \succcurlyeq Z_2||_0 + ||u \succcurlyeq Z_1||_0 \lesssim_{\mathbb{Z}} ||u^2||_{1/2+\epsilon'} + ||u||_{1/2+\epsilon'} \lesssim ||u||_{1/2+\epsilon'} (||u||_0+1).
$$

The gradient term is upper bounded by  $\|\nabla u_1 \succcurlyeq \nabla Z^{\vee}\|_0 \lesssim_{\mathbb{Z}} \|u_1\|_{1+\epsilon'}$ . The third order term is bounded trivially. Evidently, it remains to bound the terms with the block sums of the type  $\Delta_{\leq k}$ . To this end, we apply [\(3.3\)](#page-5-3) with  $\kappa' \in (1/2, (1/2 + \epsilon) \wedge \kappa)$  and  $\delta \geq 0$  with  $\delta \neq \kappa'$ :

 $||u^2 \otimes \Delta_{\leq 2n} Z_2||_{-\kappa'+\delta} + ||u \otimes \Delta_{\leq n} Z_1||_{-\kappa'+\delta} \lesssim_{\mathbb{Z}} 2^{2n\delta} ||u^2||_0 + 2^{n\delta} ||u||_0.$ 

Choosing  $\delta = 0$  and bounding  $||u^2||_0 \le ||u||_0^2$  yields the first bound (since  $-\kappa' > -1/2 - \epsilon$ ). Choosing  $\delta = \kappa$  yields the other (since  $\kappa - \kappa' > 0$ ).

3.2. Controlling  $u_1$  and  $u_2$ . With this in hand, we now turn to estimating the solutions  $u_1$ and  $u_2$  themselves. In particular, we prove the following corollary of the preceding lemma.

<span id="page-6-0"></span>**Corollary 3.4.** If  $n \geq 0$  is such that  $2^{-n(3/2-2\epsilon)} (\|u_1 + u_2\|_0 \vee 1) \simeq 1$ , then we have for all  $\alpha \in [0, 3/2 - \epsilon]$ 

<span id="page-6-2"></span>
$$
||u_1||_{\alpha} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^{\frac{2\alpha}{3/2-\epsilon}} \tag{3.4}
$$

<span id="page-6-6"></span><span id="page-6-5"></span>
$$
||u_2||_0 \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_{1/2+\epsilon'}^{1/2}
$$
\n(3.5)

$$
||u_2||_{3/2-\epsilon} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^3. \tag{3.6}
$$

Before turning to its proof, let us observe the following Schauder estimate.

<span id="page-6-8"></span>**Lemma 3.5** (Schauder estimates). Suppose that  $\alpha < 1$  and  $b, c \in C_T C^{-\alpha}$ . For any  $f \in$  $C_T \mathcal{C}^{2-\alpha}$  we have

<span id="page-6-1"></span>
$$
||f||_{2-\alpha} \lesssim_{b,c} ||(\mathcal{L} - b \otimes \nabla - c \otimes)f||_{-\alpha} + ||f(0)||_{\mathcal{C}^{2-\alpha}}.
$$
\n(3.7)

If  $b \in C_T C^{-\alpha+1}$ , then the same estimate holds for all  $\alpha < 2$ . If  $\alpha < 1/2$ , then we also have

<span id="page-6-7"></span>
$$
||f||_{2-\alpha} \lesssim_{b,c} ||(\mathcal{L} - b \cdot \nabla - c)f||_{-\alpha} + ||f(0)||_{\mathcal{C}^{2-\alpha}}.
$$
 (3.8)

It will also be helpful to note the following maximum principle.

<span id="page-6-9"></span>**Lemma 3.6** (Maximum principle). Let  $b \in C_T C^{-\alpha}$  for  $\alpha < 1/2$  and  $\Xi, g \in C_T C_b$  and  $f(0, \cdot) \in C_b$ , where  $C_b$  is the space of continuous bounded functions, equipped with the supremum norm. Then

<span id="page-6-4"></span>
$$
||f||_0 \lesssim_{b,\Xi} ||(\mathcal{L} - b \cdot \nabla f) f + e^{\Xi} f^3||_0^{1/3} + ||f(0)||_{L^{\infty}}.
$$
 (3.9)

The proof of these results follow by standard arguments and they are deferred to the end of Section [3.](#page-4-0)

We will also need the following interpolation estimate: for  $f \in \mathscr{S}'$  and any  $0 \leq \alpha \leq \beta$ ,

<span id="page-6-3"></span>
$$
||f||_{\mathcal{C}^{\alpha}} = \sup_{j} 2^{j\alpha} ||\Delta_j f||_{L^{\infty}} \leq \sup_{j} \left( 2^{j\beta} ||\Delta_j f||_{L^{\infty}} \right)^{\frac{\alpha}{\beta}} \sup_{j} ||\Delta_j f||_{L^{\infty}}^{1-\frac{\alpha}{\beta}} \lesssim ||f||_{\mathcal{C}^{\beta}}^{\frac{\alpha}{\beta}} ||f||_{L^{\infty}}^{1-\frac{\alpha}{\beta}}.
$$
 (3.10)

**Proof of Corollary [3.4](#page-6-0)**. Let us begin with the estimate for  $u_1$ . By the Schauder estimate [\(3.7\)](#page-6-1) with  $b = -6\nabla Z^{\vee} \in \bigcap_{\epsilon' > 0} C_T \mathcal{C}^{-\epsilon'}$  and  $c = 0$  and the assumption  $u(0) = \phi_0^{\sharp}$  $_0^{\sharp}$ , we have for  $\delta \in [0, 3/2 - \epsilon) \setminus \{1/2 - \epsilon\}$ :

$$
||u_1||_{3/2-\epsilon-\delta}\lesssim_{\mathbb{Z},\phi_0^\sharp}1+||U_1(u_1,u_2)||_{-1/2-\epsilon-\delta}.
$$

Indeed, for  $\delta \in [0, 1/2 - \epsilon)$  the claim follows from the first version of [\(3.7\)](#page-6-1) (with  $\alpha =$  $1/2 + \epsilon + \delta < 1$ ), while for  $\delta \in (1/2 - \epsilon, 3/2 - \epsilon)$  we apply the second version of  $(3.7)$  (with  $\alpha = 1/2 + \epsilon + \delta \in (1, 2)$ .

By the first estimate of Lemma [3.2,](#page-5-0) we have for  $\delta \geq 0$ 

$$
||U_1(u_1, u_2)||_{-1/2-\epsilon-\delta} \lesssim_{\mathbb{Z}} (1 + 2^{-n\delta}||u_1 + u_2||_0)^2
$$

Choosing  $\delta = 3/2 - 2\epsilon$ , we see that the  $L^{\infty}$  bound for  $u_1$  ([\(3.4\)](#page-6-2) with  $\alpha = 0$ ) follows by our choice of  $n \geq 1$ . Now that we know that  $||u_1||_0 \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1$ , we apply the above estimates once more, this time with  $\delta = 0$ , to obtain  $||u_1||_{3/2-\epsilon} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^2$ , i.e. [\(3.4\)](#page-6-2) with  $\alpha = 3/2 - \epsilon$ . The general version of [\(3.4\)](#page-6-2) then follows from the interpolation estimate [\(3.10\)](#page-6-3).

Now for the estimates for  $u_2$ . By the maximum principle [\(3.9\)](#page-6-4) and the assumption  $u_2(0)$  =  $\theta$ ,

$$
||u_2||_0 \lesssim_{\mathbb{Z}} ||U_2(u_1, u_2)||_0^{1/3}
$$

.

By the third estimate from Lemma [3.2](#page-5-0) and the bounds on  $u_1$  that we just derived, we have that for any  $\kappa > 1/2$  and  $\epsilon' > 0$ ,

$$
||u_2||_0 \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^{1/3} + ||u_2||_0^{2/3} + ||u_1||_{1+\epsilon'}^{1/3} + ||u_1 + u_2||_{1/2+\epsilon'}^{1/3} \left(1 + ||u_1 + u_2||_0^{1/3}\right) + 2^{n\kappa/3}||u_1 + u_2||_0^{1/3} + 2^{2n\kappa/3}||u_1 + u_2||_0^{2/3} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^{1/3} + ||u_2||_0^{2/3} + ||u_2||_0^{\frac{2(1+\epsilon')}{3(3/2-\epsilon)}} + (||u_2||_{1/2+\epsilon'} + ||u_2||_0^{\frac{2(1/2+\epsilon')}{3/2-\epsilon}})^{1/3} \left(1 + ||u_2||_0^{1/3}\right) + 2^{n\kappa/3}||u_2||_0^{1/3} + 2^{2n\kappa/3}||u_2||_0^{2/3}.
$$

By Young's inequality for products (with weights), we may subsume the powers of  $||u_2||_0$  on the left hand side. Combining this with our assumption on  $n$  yields

$$
||u_2||_0 \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_{1/2+\epsilon'}^{1/3} \left(1 + ||u_2||_0^{1/3}\right) + 2^{2n\kappa/3} ||u_2||_0^{2/3}
$$
  

$$
\lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_{1/2+\epsilon'}^{1/3} \left(1 + ||u_2||_0^{1/3}\right) + ||u_2||_0^{\frac{2\kappa}{3(3/2-2\epsilon)} + \frac{2}{3}},
$$

for any  $\kappa > 1/2$ . If we take  $\kappa \in (1/2, (3/2 - 2\epsilon)/2)$ , we may again apply Young's inequality to subsume the remaining factors of  $||u_2||_0$  on the left hand side to obtain

<span id="page-7-0"></span>
$$
||u_2||_0 \lesssim \mathbb{Z} 1 + ||u_2||_{1/2+\epsilon'}^{1/2},
$$

which is  $(3.5)$ .

It remains to prove [\(3.6\)](#page-6-6). We begin as in item 1. By the Schauder estimate [\(3.8\)](#page-6-7) for  $\mathcal{L} + 6\nabla Z^{\vee} \cdot \nabla$  and the assumption  $u_2(0) = 0$ ,

$$
||u_2||_{3/2-\epsilon} \lesssim \mathbb{Z} \left[ -e^{-6Z^{\prime \prime}} u_2^3 + U_2 \right]_{-1/2-\epsilon} \lesssim \mathbb{Z} \left[ |u_2| \right]_0^3 + \left[ |U_2| \right]_{-1/2-\epsilon}.
$$
 (3.11)

Recall by Lemma [3.2](#page-5-0) and [\(3.4\)](#page-6-2), we have that

<span id="page-8-0"></span>
$$
||U_2||_{-1/2+\delta} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^3 + ||u_2||_0^{\frac{2(1+\epsilon')}{3/2-\epsilon}} + (||u_2||_0^{\frac{2(1/2+\epsilon')}{3/2-\epsilon}} + ||u_2||_{1/2+\epsilon'}) \left(1 + ||u_2||_0\right) \tag{3.12}
$$
  

$$
\lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^3 + ||u_2||_{1/2+\epsilon'}^{3/2}.
$$

Now we apply again the interpolation estimate [\(3.10\)](#page-6-3) and obtain

$$
||u_2||_{1/2+\epsilon'}^{3/2} \lesssim \left(||u_2||_{3/2-\epsilon}^{\frac{1/2+\epsilon'}{3/2-\epsilon}}||u_2||_0^{1-\frac{1/2+\epsilon'}{3/2-\epsilon}}\right)^{3/2} = ||u_2||_{3/2-\epsilon}^{\frac{1}{2}+\epsilon''} \cdot ||u_2||_0^{1-\epsilon''},
$$

for  $\epsilon'' = \epsilon' + \epsilon \frac{1/2 + \epsilon'}{3/2 - \epsilon}$  $\frac{1}{3/2-\epsilon}$ . Combining this with [\(3.11\)](#page-7-0)-[\(3.12\)](#page-8-0), yields upon applying Young's inequality and re-arranging again,

$$
||u_2||_{3/2-\epsilon} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^3
$$
 as desired.  $\square$ 

3.3. Proof of Theorem [3.1.](#page-4-1) By Corollary [3.4,](#page-6-0) we have  $||u_1||_{3/2-\epsilon} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^2$ . It remains to show that  $||u_2||_{3/2-\epsilon} \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1$ . By the interpolation estimate  $(3.10)$ 

$$
||u_2||_{1/2+\epsilon'} \lesssim ||u_2||_{3/2-\epsilon}^{\frac{1}{3}+\frac{2}{3}\epsilon''} ||u_2||_0^{\frac{2}{3}-\frac{2}{3}\epsilon''},
$$

for  $\epsilon'' = \epsilon' + \epsilon \frac{1/2 + \epsilon'}{3/2 - \epsilon}$  $\frac{3}{2}$  as in the proof of Corollary [3.4.](#page-6-0) Plugging this into [\(3.5\)](#page-6-5), we see that for  $\gamma = \frac{\epsilon^{\prime\prime}}{3}$  $\frac{1}{3}$ ,

$$
||u_2||_0 \lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_{3/2-\epsilon}^{\frac{1}{6}+\gamma} ||u_2||_0^{\frac{1}{3}-\gamma}
$$
  

$$
\lesssim 1 + ||u_2||_{3/2-\epsilon}^{\frac{1}{4}+\frac{3}{2}\gamma} + ||u_2||_0^{1-3\gamma}
$$
  

$$
\lesssim_{\mathbb{Z},\phi_0^{\sharp}} 1 + ||u_2||_0^{\frac{3}{4}+\frac{9}{2}\gamma} + ||u_2||_0^{1-3\gamma},
$$

where in the last inequality, we applied [\(3.6\)](#page-6-6). As these exponents are less than 1 for  $\epsilon, \epsilon'$ (and thus  $\gamma$ ) sufficiently small, Young's inequality yields

$$
||u_2||_0 \lesssim_{\mathbb{Z},\phi_0^\sharp} 1.
$$

Applying this to [\(3.4\)](#page-6-2) and [\(3.6\)](#page-6-6) then yields the desired bound on  $u_1$  and  $u_2$ .

To complete the proof, we still have to justify that  $T^* = \infty$ . But since we know that  $||u||_{C_T\mathcal{C}^{3/2-\epsilon}} \leq C$  for any  $T < T^* \wedge 1$ , we must have  $T^* > 1$ . Then we can repeat the same argument on the interval [1, 2] to see that in fact  $T^* > 2$ , and so on.

# 3.4. Proof of Schauder estimates and maximum principle.

**Proof of Lemma [3.5](#page-6-8)**. We first show [\(3.7\)](#page-6-1) for  $b \in C_T C^{-\alpha}$  and  $\alpha < 1$ , and then indicate how to adapt the argument to obtain [\(3.7\)](#page-6-1) for  $b \in C_T C^{-\alpha+1}$  and  $\alpha < 2$ , as well as [\(3.8\)](#page-6-7). By the Schauder estimates for L, we have for  $\tau \in [0, T]$ , for  $\epsilon \in [0, 2]$ , and for  $\epsilon' \geq 0$  small enough so that  $2 - \alpha - \epsilon' > 1$  (and thus  $\|\nabla f\|_{C_{\tau}L^{\infty}} \lesssim \|f\|_{C_{\tau}C^{2-\alpha-\epsilon'}}$ ):

$$
||f||_{C_{\tau}C^{2-\alpha-\epsilon}} \lesssim \tau^{\epsilon/2} ||\mathcal{L}f||_{C_{\tau}C^{-\alpha}} + ||f(0)||_{C^{2-\alpha-\epsilon}}\leq \tau^{\epsilon/2} ||(\mathcal{L} - b \otimes \nabla - c \otimes)f||_{C_{\tau}C^{-\alpha}} + \tau^{\epsilon/2} ||b \otimes \nabla f||_{C_{\tau}C^{-\alpha}}+ \tau^{\epsilon/2} ||c \otimes f||_{C_{\tau}C^{-\alpha}} + ||f(0)||_{C^{2-\alpha-\epsilon}}\leq \tau^{\epsilon/2} ||(\mathcal{L} - b \otimes \nabla - c \otimes)f||_{C_{\tau}C^{-\alpha}} + \tau^{\epsilon/2} (||b||_{C_{T}C^{-\alpha}} + ||c||_{C_{T}C^{-\alpha}}) ||f||_{C_{\tau}C^{2-\alpha-\epsilon'}}+ ||f(0)||_{C^{2-\alpha-\epsilon}}.
$$
\n(3.13)

We choose  $\epsilon = \epsilon' > 0$  and  $\tau > 0$  small enough so that with the implicit constant C on the right hand side  $C\tau^{\epsilon/2}(\|b\|_{C_T\mathcal{C}^{-\alpha}} + \|c\|_{C_T\mathcal{C}^{-\alpha}}) \leq \frac{1}{2}$  $\frac{1}{2}$ . In that way we obtain

$$
||f||_{C_{\tau}C^{2-\alpha-\epsilon}} \lesssim \tau^{\epsilon/2} ||(\mathcal{L} - b \otimes \nabla - c \otimes) f||_{C_{\tau}C^{-\alpha}} + ||f(0)||_{C^{2-\alpha-\epsilon}}.
$$

As  $\tau$  can be chosen independently of  $f(0)$  and as  $||f(\tau)||_{C^{2-\alpha-\epsilon}} \lesssim ||f||_{C_{\tau}C^{2-\alpha-\epsilon}}$ , we can now iterate this on  $[\tau, 2\tau]$ ,  $[2\tau, 3\tau]$ , ..., and obtain

<span id="page-9-0"></span>
$$
||f||_{C_{\tau}C^{2-\alpha-\epsilon}} \lesssim ||(\mathcal{L} - b \otimes \nabla - c \otimes)f||_{C_TC^{-\alpha}} + ||f(0)||_{C^{2-\alpha}}.
$$

To obtain the bound for  $\epsilon = 0$  we simply apply [\(3.13\)](#page-9-0) once more, this time for  $\tau = T$  and  $\epsilon = 0$  and with the same  $\epsilon' > 0$  as before.

If  $b \in C_T C^{-\alpha+1}$  for  $\alpha < 2$ , then we only use the  $C_{\tau} L^{\infty}$  regularity of f and obtain  $||b \otimes$ <br> $||b||_{\alpha}$ ,  $||c||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $||c||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $||d||_{\alpha}$ ,  $\nabla f||_{C_{\tau}C^{-\alpha}} \leq ||b||_{C_{\tau}C^{-\alpha+1}} ||f||_{C_{\tau}L^{\infty}}$ . Since  $2 - \alpha > 0$  and thus  $C_{\tau}C^{2-\alpha-\epsilon} \subset C_{\tau}L^{\infty}$  for all sufficiently small  $\epsilon > 0$ , we can then use the same argument as before.

Nearly the same arguments also work for [\(3.8\)](#page-6-7), except that now we have to make sure that the products  $b \cdot \nabla f$  and  $c \nabla f$  are well-defined. This is the case if  $-\alpha + 2 - \alpha - 1 > 0$ , i.e., if  $\alpha < 1/2$ .

**Proof of Lemma [3.6](#page-6-9).** We assume first that  $b, \Xi, q, f(0, \cdot)$  are smooth functions, so that  $f \in C^{1,2}$  is a classical solution, and we write

$$
(\partial_t - \Delta - b \cdot \nabla)f = -e^{\Xi}f^3 + g.
$$

By compactness, f takes its minimum and maximum on  $[0, T] \times \mathbb{T}^3$ . Let  $f(t, x)$  be a local maximum. If  $t = 0$ , then  $|f(t, x)| \leq \|f(0, \cdot)\|_{L^{\infty}}$ . If  $t \in (0, T)$ , then  $\partial_t f(t, x) = \partial_t f(t, x) = 0$ , while  $\Delta f(t, x) \leq 0$ . Therefore

$$
-e^{\Xi(t,x)}f^3(t,x) + g(t,x) = (\partial_t - \Delta - b \cdot \nabla)f(t,x) \geq 0,
$$

and thus

$$
f(t,x) \leqslant \left(\frac{g(t,x)}{e^{\Xi(t,x)}}\right)^{1/3} \leqslant \frac{\left\|g\right\|_{C_TL^\infty}^{1/3}}{e^{-\left\|\Xi\right\|_{C_TL^\infty}/3}}.
$$

If f attains its maximum in  $(T, x)$ , then we must have

$$
\partial_t f(T, x) \geq 0
$$

because otherwise  $f(t, x) > f(T, x)$  for some  $t < T$ . Furthermore, x is a local maximum of  $f(T, \cdot)$  so that we still have  $\partial_i f(T, x) = 0$  and  $\Delta f(T, x) \leq 0$ . Thus, we have again  $(\partial_t - \Delta - b \cdot \nabla) f(t, x) \geq 0$  and we get the same estimate as before.

To control the minimum, it suffices to note that minima of f are maxima of  $-f$  and that  $(\partial_t - \Delta - b \cdot \nabla)(-f) = -e^{\Xi}(-f)^3 - g.$ 

Our estimate does not depend on b at all, and it only depends on  $\Xi$  through its  $L^{\infty}$ norm. For  $b \in C_T C^{-\alpha}$  with  $\alpha < 1/2$  it follows from our Schauder estimates that if we approximate  $b_n \to b$  in  $C_T \mathcal{C}^{-\alpha-}$  and  $\Xi_n, g_n \to \Xi$  in  $C_T L^{\infty}$  and  $f_n(0, \cdot) \to f(0, \cdot)$  in  $L^{\infty}$  with  $b_n, \Xi_n, g_n, f_n(0, \cdot)$  smooth, then the solutions  $f_n$  to

$$
(\partial_t - \Delta - b_n \cdot \nabla) f_n = -e^{\Xi_n} f^3 + g_n, \qquad f_n(0) = f_n(0, \cdot),
$$

converge locally uniformly to f, and thus the estimate also holds for f.  $\Box$ 

#### 4. Possible extensions

<span id="page-10-0"></span>In recent years there have been many works dedicated to the  $\Phi_3^4$  equation, and by now we know much more than global existence and uniqueness of solutions on the torus. We expect that with the multiplicative transform some proofs of more refined properties can be simplified. To keep the paper short and accessible we do not attempt this here, but let us sketch some possible extensions of our results:

- Initial condition: We could easily extend the local solution theory to  $\phi_0^{\sharp} \in C^{-\alpha}$  by working in spaces  $C([0,T], \mathcal{C}^{-\alpha}) \cap \mathcal{M}_T^{\gamma} \mathcal{C}^{1+\epsilon'}$  with  $||u||_{\mathcal{M}_T^{\gamma} \mathcal{C}^{1+\epsilon'}} = \sup_{t \in [0,T]} t^{\gamma} ||u(t)||_{\mathcal{C}^{1+\epsilon'}},$ for  $\gamma = \frac{1+\epsilon'+\alpha}{2}$  $\frac{e^t + \alpha}{2}$ , see [\[GP17,](#page-14-11) Section 6] for details. To control the nonlinearity  $u^3$  while keeping the singularity integrable in time we need  $\alpha < 2/3$ . Since  $Z(0) \in C^{-1/2-\epsilon}$ , we can thus take any element of  $C^{-2/3+\epsilon}$  as initial condition and the special form  $\phi_0 = Z(0) + \phi_0^{\sharp}$  $\frac{1}{0}$  is not necessary.
- More singular noise: For applications in quantum field theory we are only interested in the invariant measure on  $\mathbb{T}^3$  (or on  $\mathbb{R}^3$  if we consider the equation on the whole space). Therefore, the physically meaningful dimension would be  $1 + 3 = 4$ , i.e. the  $\Phi_4^4$  equation on  $\mathbb{R}_+ \times \mathbb{T}^4$  resp.  $\mathbb{R}_+ \times \mathbb{R}^4$ . (Here the lower index in  $\Phi_d^k$  indicates the space dimension in the equation, the upper index  $k$  means that the leading order nonlinearity is  $\phi^{k-1}$ . Recall that in the stochastic quantization approach [\[PW81\]](#page-15-0), the "stochastic time",  $\mathbb{R}_+$ , is an extra dimension and does not correspond to "real" time.) But in  $d = 4$  the space-time white noise is more singular and the  $\Phi_4^4$  equation is *critical* in the sense of Hairer [\[Hai14\]](#page-14-4), which roughly means that the solution  $u$  to our transformed equation is expected to have the same regularity as  $\phi$ . There are currently no mathematical tools for the  $\Phi_4^4$  equation. But it is possible to treat a proxy for "4 –  $\epsilon$  dimensions", by considering a slightly regularized noise in  $d = 4$ [\[BCCH21,](#page-14-13) [CMW19\]](#page-14-14).

Our approach works if the regularity of  $-6\nabla Z^{\vee}$  is bigger than  $-1/2$ , and also bigger than  $-1$  minus the minimum regularity of  $Z_0, Z_1, Z_2$ . For more singular noise it might be possible to further improve the regularity of  $Z_0, Z_1$  by additional additive and multiplicative transformations. But  $Z_2$  will never be more regular than  $Z$ . In " $d = 4 - 2\epsilon$ " we have  $Z \in C_T C^{-1+\epsilon}$  and  $\nabla Z^{\vee} \in C_T C^{-1+2\epsilon}$ . Therefore, we need  $\epsilon > 1/3$ . Beyond that it might be possible to additionally use a Zvonkin transform as in [\[ZZZ20\]](#page-15-2) to remove the term  $-6\nabla Z^{\vee}\cdot\nabla u$ . But this becomes quite complicated, and for very singular noise it seems easier to use the arguments of [\[CMW19\]](#page-14-14).

• Coming down from infinity: By adapting Lemma 2.7 in [\[MW20\]](#page-14-9) we could derive an alternative version of the maximum principle:

$$
||f(t)||_{L^{\infty}} \lesssim_{b,\Xi} \max\left\{ t^{-1/2}, ||(\mathcal{L} - b \cdot \nabla f) f + e^{\Xi} f^{3}||_{0}^{1/3} \right\},\tag{4.1}
$$

where the right hand side is independent of  $f(0)$ . With this it should be possible to control  $u(1)$  independently of the initial condition. As in [\[MW17b\]](#page-14-7) this should lead to a proof of existence of stationary measures for  $\phi$  via the Krylov-Bogoliubov method.

- Existence of moments: The constant C in Theorem [3.1](#page-4-1) depends on  $e^{3Z^{\vee}}$  and on  $e^{-6Z^{\vee}}$ . Since  $Z^{\vee}$  is a second order polynomial of a Gaussian process, it is therefore not even clear if  $\mathbb{E}[\|u\|_{C_1C^{3/2-\epsilon}}] < \infty$ , nor if after transforming back we could get  $\mathbb{E}[\|\phi\|_{C_1C^{-1/2-\epsilon}}]<\infty$ . While it is actually known that  $\phi$  has stretched exponential moments [\[MW20\]](#page-14-9) and, at least at stationarity, even  $\mathbb{E}[\exp(\lambda \langle \phi(t), \eta \rangle^4)] < \infty$  for suitable test functions  $\eta$  and sufficiently small  $\lambda > 0$  [\[HS21\]](#page-14-15). To try recovering such results with our approach, we would have to slightly change the multiplicative transformation and multiply with  $e^{3\Delta_{>m}Z^{\vee}}$  instead, where m is large enough so that  $\|\Delta_{>m}Z^{\vee}\|_{C_T\mathcal{C}^{\alpha}} \simeq 1$  for a suitable  $\alpha < 1 - \epsilon$ .
- Extension to the whole space: To solve the equation on  $\mathbb{R}_+ \times \mathbb{R}^3$  we would have to use weighted spaces. Then we have to be careful when implementing the exponential transform, because  $e^{3Z^{\vee}(x)}$  grows too fast as  $|x| \to \infty$ . We should therefore apply [\[GH19,](#page-14-8) Lemma 2.4] to decompose  $Z^{\vee} = Z^{\vee} > +Z^{\vee} <$ , where  $Z^{\vee} > \in C_T C^{1-\epsilon}$  (without weight) and  $Z^{\vee}$  is in a weighted space but has regularity better than 1. And then we use the multiplicative transform with  $e^{3Z\check{}}$ . After that we expect that the construction of solutions on  $\mathbb{R}_+ \times \mathbb{R}^3$  can be handled similarly as for the  $\Phi_2^4$  equation on  $\mathbb{R}_+ \times \mathbb{R}^2$  in [\[GH19\]](#page-14-8).

# Appendix A. Some nonlinear functions of the noise

Let  $\mathcal{T} = (Z, \llbracket Z^2 \rrbracket, Z^{\vee}, Z^{\vee}, Z^{\vee} \odot Z, Z^{\vee} \odot \llbracket Z^2 \rrbracket - \frac{b}{3}$ <br>rulerities given in Section 2. In the preceding was  $\frac{b}{3}, |\nabla Z^{\vee}|^2 - \frac{b}{3}$  $\frac{b}{3}$ ,  $Z^{\Psi} \odot [Z^2] - bZ$ , with the regularities given in Section [2.](#page-1-1) In the preceding, we used that the trees  $\mathcal T$  are well-defined after renormalization and that  $Y$  and a certain product involving its derivative are continuous functions of these trees and, in particular, are well-behaved under renormalization. For the convenience of the reader, we collect these results here.

**Lemma A.1.** If  $\xi$  is a white noise on  $\mathbb{R} \times \mathbb{T}^3$ , then all terms in  $\mathcal{T}$  are well-defined as (sums of ) iterated stochastic integrals, and they have the regularities given in Table [1.](#page-2-1)

*Proof.* This is shown in [\[CC18,](#page-14-6) [MWX17\]](#page-15-1), except that there the term  $|\nabla Z^{\vee}|^2 - \frac{b}{3}$  does not  $\frac{3}{3}$  about the paraproduct is always well-defined, the existence of this term is appear. Note that, since the paraproduct is always well-defined, the existence of this term is equivalent to the existence of the renormalized resonant product  $\sum_i \partial_i Z^{\vee} \odot \partial_i Z^{\vee} - \frac{b}{3}$  $\frac{b}{3}$ . As a space-time distribution, this latter term is equivalent to  $Z^{\vee} \odot [Z^2] - \frac{b}{3}$ <br>Leibniz's rule to  $\mathcal{L}(Z^{\vee} \odot Z^{\vee})$  we obtain  $\frac{b}{3}$ : Indeed, by applying Leibniz's rule to  $\mathcal{L}(Z^{\vee} \odot Z^{\vee})$  we obtain

$$
\sum_{i} \partial_{i} Z^{\vee} \odot \partial_{i} Z^{\vee} - \frac{b}{3} = Z^{\vee} \odot \mathcal{L} Z^{\vee} - \frac{b}{3} - \mathcal{L} (Z^{\vee} \odot Z^{\vee}) - Z^{\vee} \odot Z^{\vee}
$$

$$
= Z^{\vee} \odot [Z^{2}] - \frac{b}{3} - \mathcal{L} (Z^{\vee} \odot Z^{\vee}) - Z^{\vee} \odot Z^{\vee}.
$$

Since  $Z^{\vee}$  has positive regularity, all the terms on the right hand side are well-defined. However, it is not clear if  $\mathcal{L}(Z^{\vee} \odot Z^{\vee})$  is in a  $C_T C^{\alpha}$  space and we can only treat it as a space time distribution. space-time distribution.

Therefore, we need to use stochastic arguments to construct  $\sum_i \partial_i Z^{\vee} \odot \partial_i Z^{\vee} - \frac{b}{3}$ <br>cin Leibniz's rule, this time for  $\Delta$ , we see that construction  $\sum_i \partial_i Z^{\vee} \odot \partial_i Z^{\vee} = b$  $\frac{b}{3}$ . Using again Leibniz's rule, this time for  $\Delta$ , we see that constructing  $\sum_i \partial_i Z^{\vee} \odot \partial_i Z^{\vee} - \frac{b}{3}$  $\frac{b}{3} \in C_T C^{-\epsilon}$ is equivalent to constructing  $Z^{\vee} \odot \Delta Z^{\vee} - \frac{b}{3}$  $\frac{b}{3}$ . But this is nearly the same as  $Z^{\vee} \odot [Z^2] - \frac{b}{3}$  $\frac{b}{3}$ , except that we replace  $[\![Z^2]\!](t)$  by  $\int_0^t \Delta P_{t-s}[\![Z^2]\!](s)ds$ . The kernel  $\int_0^t \Delta P_{t-s}ds$  neither gains<br>nor loses regularity, and therefore (possibly) the same argument as in [CC18, Section 4.5] or nor loses regularity, and therefore (nearly) the same argument as in [\[CC18,](#page-14-6) Section 4.5] or [\[MWX17,](#page-15-1) Section 4.2] show that  $Z^{\vee} \odot \Delta Z^{\vee} - \frac{b}{3}$  $\frac{b}{3} \in C_T C^{-\epsilon}$ .

To prove the desired continuity estimates for  $Y$  and a certain product involving its derivative, we need some tools from paracontrolled distributions. First we recall the paralineariza-tion theorem (see [\[BCD11\]](#page-14-12) or [\[GIP15,](#page-14-5) Lemma 2.6]) which says that for  $0 < \alpha < 1$ ,  $F \in C^2$ and  $f \in \mathcal{C}^{\alpha}$  we have

<span id="page-12-0"></span>
$$
F(f) - F'(f) \otimes f \in \mathcal{C}^{2\alpha}.
$$
\n(A.1)

Next, we recall Bony's paramultiplication bound [\[Bon81,](#page-14-16) Theorem 2.3] which says that if  $f, g \in \mathcal{C}^{\alpha}$  with non-integer  $\alpha > 0$  and  $h \in \mathcal{C}^{\beta}$ , then

<span id="page-12-2"></span>
$$
||f \otimes (g \otimes h) - (fg) \otimes h||_{\alpha + \beta} \lesssim ||f||_{\alpha} ||g||_{\alpha} ||h||_{\beta}.
$$
 (A.2)

Next, let us recall the well-known fact that if we let  $Jf(t) = \int_0^t P_{t-s} f ds$ , then J approximately commutes with paraproducts in the sense that for all  $\alpha \in (0,1)$ ,  $\beta \in \mathbb{R}$  and  $\delta < 2$ 

<span id="page-12-3"></span>
$$
||J(f \otimes g) - f \otimes Jg||_{C_T C^{\alpha+\beta+\delta}} \lesssim (||f||_{C_T C^{\alpha}} + ||f||_{C_T^{\frac{\alpha}{2}} L^{\infty}}) ||g||_{C_T C^{\beta}},
$$
(A.3)

where  $||f||_{C_T^{\frac{\alpha}{2}}L^{\infty}} = \sup_{0 \le s < t \le T} \frac{||f(t) - f(s)||_{L^{\infty}}}{|t - s|^{\frac{\alpha}{2}}}$  $\frac{1-1}{|t-s|^{\frac{\alpha}{2}}}$ . (While to the best of our knowledge this exact bound does not appear in the literature, it is a folklore result; see, e.g., [\[Per14,](#page-15-3) Lemma 5.5.7] or [\[KP20,](#page-14-17) Lemma A.1] for very similar estimates from which this follows.) Finally, we recall the commutator estimate [\[GIP15,](#page-14-5) Lemma 2.4] which says that

<span id="page-12-1"></span>
$$
C(f, g, h) = (f \otimes g) \odot h - f(g \odot h)
$$
\n(A.4)

is a bounded trilinear map from  $\mathcal{C}^{\alpha} \times \mathcal{C}^{\beta} \times \mathcal{C}^{\gamma}$  to  $\mathcal{C}^{\alpha+\beta+\gamma}$ , provided that  $\alpha \in (0,1)$  and  $\beta+\gamma < 0$ while  $\alpha + \beta + \gamma > 0$ . We now show that Y and a certain product involving its derivative are continuous in  $\mathcal{T}$ . (And consequently, they are well-defined in the renormalization limit.)

# Lemma A.2. Let

$$
\mathcal{L}Y = 3e^{3Z^{\mathcal{V}}}(Z^{\mathcal{V}}[[Z^2]] - b(Z + Z^{\mathcal{V}})), \quad Y(0) = 0.
$$

Then  $Y \in \bigcap_{\epsilon > 0} C_T \mathcal{C}^{1-\epsilon}$  is a continuous function of  $\mathcal{T}$ . Moreover,

$$
\nabla Z^{\vee} \cdot \nabla Y - be^{3Z^{\vee}} Z^{\Psi} \in \bigcap_{\epsilon > 0} C_T \mathcal{C}^{-2\epsilon}
$$

is also a continuous function of  $\mathcal T$ .

Proof. We split up

$$
Z^{\mathbf{V}}[[Z^2]] - b(Z + Z^{\mathbf{V}}) = (Z^{\mathbf{V}} \odot [[Z^2]] - bZ) + (Z^{\mathbf{V}} \otimes [[Z^2]] - bZ^{\mathbf{V}}) + Z^{\mathbf{V}} \otimes [[Z^2]].
$$

By assumption,  $(Z^{\mathbf{V}} \odot [Z^2] - bZ) \in C_T C^{-\frac{1}{2} - \epsilon}$ , and since  $Z^{\mathbf{V}} \in C_T C^{1-\epsilon}$  and the regularity is preserved under the smooth function  $\exp(\cdot)$ , the product  $3e^{3Z^{\vee}}(Z^{\Psi}\odot [Z^2]-bZ)$  is well-<br>defined and in  $C\mathscr{L}^{-\frac{1}{2}-\epsilon}$ . Similarly,  $Z^{\Psi}\odot [Z^2] = C\mathscr{L}^{-\frac{1}{2}-\epsilon}$  by the program dust extinct as and defined and in  $C_T C^{-\frac{1}{2}-\epsilon}$ . Similarly,  $Z^{\Psi} \otimes [Z^2] \in C_T C^{-\frac{1}{2}-\epsilon}$  by the paraproduct estimates, and therefore also  $3e^{3Z^{\vee}}Z^{\Psi} \otimes [Z^2] \in C_T C^{-\frac{1}{2}-\epsilon}$ . The most complicated term is the one involving  $Z^{\Psi} \otimes [Z^2]$ . This parameter is of course well defined, but its regularity is only  $C-C^{-1-\epsilon}$  $Z^{\mathbf{V}} \otimes [Z^2]$ . This paraproduct is of course well-defined, but its regularity is only  $C_T \mathcal{C}^{-1-\epsilon}$ and therefore the resonant product  $3e^{3Z^{\vee}} \odot (Z^{\Psi} \otimes [Z^2])$  is not defined.<br>Thus we need to use the more refined tools from personal use described.

Thus we need to use the more refined tools from paracalculus described above to construct this product as a continuous function of  $\mathcal T$ . First note that by the paralinearization theorem [\(A.1\)](#page-12-0) we have  $e^{3Z^{\vee}} - e^{3Z^{\vee}} \otimes 3Z^{\vee} \in C_T C^{2-2\epsilon}$ , and therefore this difference can be multiplied<br>with  $Z^{\Psi} \otimes \mathbb{Z}^{2\mathbb{I}}$ . Then, by the commutator estimate (A.4), we have that with  $Z^{\mathbf{V}} \otimes [Z^2]$ . Then, by the commutator estimate [\(A.4\)](#page-12-1), we have that

$$
3e^{3Z\checkmark} \odot (Z^{\Psi} \otimes \llbracket Z^{2} \rrbracket) - 3e^{3Z\checkmark} bZ^{\Psi} = (3e^{3Z\checkmark} - 9e^{3Z\checkmark} \otimes Z^{\checkmark}) \odot (Z^{\Psi} \otimes \llbracket Z^{2} \rrbracket) + 9C(e^{3Z\checkmark}, Z^{\checkmark}, Z^{\checkmark}, Z^{\Psi} \otimes \llbracket Z^{2} \rrbracket) + 9e^{3Z\checkmark} (Z^{\checkmark} \odot (Z^{\Psi} \otimes \llbracket Z^{2} \rrbracket)) - 3e^{3Z\checkmark} bZ^{\Psi} = C_T C^{1-3\epsilon} + 9e^{3Z\checkmark} \left( C(Z^{\Psi}, \llbracket Z^{2} \rrbracket, Z^{\checkmark}) + Z^{\Psi} \left( (\llbracket Z^{2} \rrbracket \odot Z^{\checkmark}) - \frac{b}{3} \right) \right),
$$

where on the right hand side we used the notation  $C_T \mathcal{C}^{1-3\epsilon}$  to denote a term of this regularity which is given as a continuous function of the data. The commutator on the right hand side has positive regularity, so it no longer poses a problem. The term  $(\llbracket Z^2 \rrbracket \odot Z^{\vee}) - \frac{b}{3}$  $\frac{b}{3}$  is in  $C_T C^{-\epsilon}$ by assumption, so we can multiply it with  $Z^{\mathbf{V}} \in C_T C^{\frac{1}{2}-\epsilon}$  and the product is in  $C_T C^{-\epsilon}$ . Then we can multiply this product with  $e^{3Z^{\vee}} \in C_T C^{1-\epsilon}$ , and therefore the remaining expression on the right hand side is well-defined and in  $C_T C^{-\epsilon} \subset C_T C^{-\frac{1}{2}-\epsilon}$ .

Next, we consider the renormalized product  $\nabla Z^{\vee} \cdot \nabla Y - be^{3Z^{\vee}} Z^{\Psi}$ . The idea is to write Y as a paraproduct plus a more regular remainder, and to apply the commutator estimate for C as above. By the previous considerations and the paraproduct estimates,  $\mathcal{L}Y - 3e^{3Z}$  $\overline{a}$  $(Z^{\mathbf{v}} \otimes [Z^2]) \in C_T C^{-\frac{1}{2}-\epsilon}$ . By Bony's paramultiplication bound [\(A.2\)](#page-12-2), we moreover have

$$
\|3e^{3Z^{\vee}}\otimes (Z^{\vee}\otimes \llbracket Z^{2}\rrbracket) - (3e^{3Z^{\vee}}Z^{\vee})\otimes \llbracket Z^{2}\rrbracket\|_{C_{T}C^{-3\epsilon}} \lesssim \|3e^{3Z^{\vee}}\|_{C_{T}C^{\frac{1}{2}-\epsilon}}\|Z^{\vee}\|_{C_{T}C^{\frac{1}{2}-\epsilon}}\|\llbracket Z^{2}\rrbracket\|_{C_{T}C^{-1-\epsilon}},
$$

so that  $\mathcal{L}Y - (3e^{3Z^{\vee}}Z^{\vee}) \otimes [Z^2] \in C_T C^{-\frac{1}{2}-\epsilon}$ . By the Schauder estimates for  $\mathcal{L}$  we deduce that  $Y - J((3e^{3Z^{\vee}}Z^{\vee}) \otimes [Z^2]) \in C_T C^{\frac{3}{2}-\epsilon}$ . Therefore, the product  $\nabla Z^{\vee} \cdot \nabla (Y - J((3e^{3Z^{\vee}}Z^{\vee}) \otimes$  $[\![Z^2]\!]$ ))  $\in C_T C^{-\epsilon}$  is well-defined. To deal with the term  $\nabla Z^{\vee} \cdot \nabla J((3e^{3Z^{\vee}}Z^{\vee}) \otimes [\![Z^2]\!])$ ,<br>we need to use the commutator estimate (A.3). To apply this however we first need we need to use the commutator estimate  $(A.3)$ . To apply this, however, we first need time regularity of  $3e^{3Z\vee}Z^{\Psi}$ . For  $Z^{\Psi}$  it follows from the estimate (1.11) in [\[MWX17\]](#page-15-1) that  $\|Z^{\mathbf{\Psi}}\|_{C^{\frac{1}{4}-\frac{\epsilon}{2}}_T L^\infty}$  $< \infty$ . For  $Z^{\vee}$  it follows from the Schauder estimates in [\[GH19,](#page-14-8) Lemma 2.10] that  $\|\dot{Z}^{\vee}\|_{C_T^{\frac{1}{2}-\frac{\epsilon}{2}}L^{\infty}} \lesssim \|\llbracket Z^2 \rrbracket\|_{C_T C^{-1-\epsilon}} < \infty$ . Therefore, using that  $J\llbracket Z^2 \rrbracket = Z^{\vee}$ , we can apply [\(A.3\)](#page-12-3) to get

$$
J((3e^{3Z^{\mathcal{V}}}Z^{\mathcal{V}})\otimes[[Z^2]])-(3e^{3Z^{\mathcal{V}}}Z^{\mathcal{V}})\otimes Z^{\mathcal{V}}\in C_T\mathcal{C}^{\frac{3}{2}-3\epsilon},
$$

and it remains to make sense of  $\nabla[(3e^{3Z^{\vee}}Z^{\Psi}) \otimes Z^{\vee}] \cdot \nabla Z^{\vee}$ . By Leibniz's rule and the paraproduct estimates we have  $\nabla[(3e^{3Z^{\mathcal{V}}}Z^{\mathcal{V}})\otimes Z^{\mathcal{V}}] - (3e^{3Z^{\mathcal{V}}}Z^{\mathcal{V}})\otimes \nabla Z^{\mathcal{V}} \in C_T\mathcal{C}^{\frac{1}{2}-2\epsilon}$ . Then we apply the commutator estimate for  $C$  to reduce the last ill-defined product to

$$
\sum_{i} ((3e^{3Z^{\vee}} Z^{\mathbf{v}}) \otimes \partial_i Z^{\vee}) \odot \partial_i Z^{\vee} - be^{3Z^{\vee}} Z^{\mathbf{v}}
$$
  
= 
$$
\sum_{i} C(3e^{3Z^{\vee}} Z^{\mathbf{v}}, \partial_i Z^{\vee}, \partial_i Z^{\vee}) + 3e^{3Z^{\vee}} Z^{\mathbf{v}} (\sum_{i} (\partial_i Z^{\vee} \odot \partial_i Z^{\vee}) - \frac{b}{3}).
$$

Now it suffices to note that  $\sum_i (\partial_i Z^{\vee} \odot \partial_i Z^{\vee}) - \frac{b}{3} = |\nabla Z^{\vee}|^2 - \frac{b}{3} - 2 \sum_i (\partial_i Z^{\vee} \otimes \partial_i Z^{\vee})$  and that the paraproducts are well-defined and in  $C_T C^{-2\epsilon}$ . that the paraproducts are well-defined and in  $C_T \tilde{C}^{-2\epsilon}$ 

#### Acknowledgements

The first author gratefully acknowledges support from NSERC, cette recherche a été financée par CRSNG [RGPIN-2020-04597, DGECR-2020-00199]. The second author gratefully acknowledges financial support by the DFG via Research Unit FOR2402.

#### **REFERENCES**

- <span id="page-14-13"></span>[BCCH21] Y. Bruned, A. Chandra, I. Chevyrev, and M. Hairer, Renormalising SPDEs in regularity structures, J. Eur. Math. Soc. (JEMS) 23 (2021), no. 3, 869-947. MR 4210726
- <span id="page-14-12"></span>[BCD11] Hajer Bahouri, Jean-Yves Chemin, and Raphael Danchin, Fourier analysis and nonlinear partial differential equations, Springer, 2011.
- <span id="page-14-16"></span>[Bon81] Jean-Michel Bony, Calcul symbolique et propagation des singularites pour les équations aux dérivées partielles non linéaires, Ann. Sci. Éc. Norm. Supér. (4)  $14$  (1981), 209–246.
- <span id="page-14-6"></span>[CC18] Rémi Catellier and Khalil Chouk, *Paracontrolled distributions and the 3-dimensional stochastic* quantization equation, Ann. Probab.  $46$  (2018), no. 5, 2621–2679.
- <span id="page-14-14"></span>[CMW19] Ajay Chandra, Augustin Moinat, and Hendrik Weber, A priori bounds for the  $\phi^4$  equation in the full sub-critical regime, arXiv preprint arXiv:1910.13854 (2019).
- <span id="page-14-2"></span>[FG19] M. Furlan and M. Gubinelli, Weak universality for a class of 3d stochastic reaction-diffusion models, Probab. Theory Related Fields 173 (2019), no. 3-4, 1099–1164. MR 3936152
- <span id="page-14-8"></span>[GH19] Massimiliano Gubinelli and Martina Hofmanová, Global solutions to elliptic and parabolic  $\Phi^4$ models in Euclidean space, Comm. Math. Phys. 368 (2019), no. 3, 1201–1266. MR 3951704
- <span id="page-14-0"></span>[GH21]  $\qquad \qquad \qquad$ , A PDE construction of the Euclidean  $\phi_3^4$  quantum field theory, Comm. Math. Phys. 384 (2021), no. 1, 1–75. MR 4252872
- <span id="page-14-5"></span>[GIP15] Massimiliano Gubinelli, Peter Imkeller, and Nicolas Perkowski, Paracontrolled distributions and singular PDEs, Forum of Mathematics, Pi 3 (2015), no. e6.
- <span id="page-14-11"></span>[GP17] Massimiliano Gubinelli and Nicolas Perkowski, KPZ reloaded, Comm. Math. Phys. 349 (2017), no. 1, 165–269.
- <span id="page-14-4"></span>[Hai14] Martin Hairer, A theory of regularity structures, Invent. Math. 198 (2014), no. 2, 269–504.
- <span id="page-14-10"></span>[HL15] Martin Hairer and Cyril Labbé, A simple construction of the continuum parabolic Anderson model on  $\mathbb{R}^2$ , Electron. Commun. Probab. 20 (2015), no. 43, 11. MR 3358965
- <span id="page-14-15"></span>[HS21] Martin Hairer and Rhys Steele, *The*  $\phi_3^4$  *measure has sub-Gaussian tails*, arXiv preprint arXiv:2102.11685 (2021).
- <span id="page-14-1"></span>[HX18] Martin Hairer and Weijun Xu, Large-scale behavior of three-dimensional continuous phase coexistence models, Comm. Pure Appl. Math. 71 (2018), no. 4, 688–746. MR 3772400
- <span id="page-14-17"></span>[KP20] Helena Kremp and Nicolas Perkowski, *Multidimensional SDE with distributional drift and Lévy* noise, arXiv preprint arXiv:2008.05222 (2020).
- <span id="page-14-3"></span>[MW17a] Jean-Christophe Mourrat and Hendrik Weber, Convergence of the two-dimensional dynamic *Ising-Kac model to*  $\phi_2^4$ , Comm. Pure Appl. Math. **70** (2017), no. 4, 717–812.
- <span id="page-14-7"></span>[MW17b]  $\ldots$ , The dynamic  $\phi_3^4$  model comes down from infinity, Comm. Math. Phys. **356** (2017), no. 3, 673–753.
- <span id="page-14-9"></span>[MW20] Augustin Moinat and Hendrik Weber, Space-time localisation for the dynamic  $\Phi_3^4$  model, Comm. Pure Appl. Math. 73 (2020), no. 12, 2519–2555. MR 4164267
- <span id="page-15-1"></span>[MWX17] Jean-Christophe Mourrat, Hendrik Weber, and Weijun Xu, Construction of  $\Phi_3^4$  diagrams for pedestrians, From particle systems to partial differential equations, Springer Proc. Math. Stat., vol. 209, Springer, Cham, 2017, pp. 1–46. MR 3746744
- <span id="page-15-3"></span>[Per14] Nicolas Simon Perkowski, Studies of robustness in stochastic analysis and mathematical finance, Ph.D. thesis, Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Fakultät II, 2014.
- <span id="page-15-0"></span>[PW81] G. Parisi and Yong Shi Wu, Perturbation theory without gauge fixing, Sci. Sinica 24 (1981), no. 4, 483–496. MR 626795
- <span id="page-15-2"></span>[ZZZ20] Xicheng Zhang, Rongchan Zhu, and Xiangchan Zhu, Singular HJB equations with applications to KPZ on the real line, arXiv preprint arXiv:2007.06783 (2020).

(Aukosh Jagannath) Department of Statistics and Actuarial Sciences, University of Wa-TERLOO

Email address: a.jagannath@uwaterloo.ca

(Nicolas Perkowsi) INSTITUT FÜR MATHEMATIK, FREIE UNIVERSITÄT BERLIN Email address: perkowski@math.fu-berlin.de