L^q-REGULARITY FOR NONLINEAR ELLIPTIC EQUATIONS WITH SCHRÖDINGER-TYPE LOWER ORDER TERMS

MIKYOUNG LEE AND JIHOON OK

ABSTRACT. We consider nonlinear elliptic equations of the *p*-Laplacian type with lower order terms which involve nonnegative potentials satisfying a reverse Hölder type condition. Then we obtain interior and boundary L^q estimates for the gradient of weak solutions and the lower order terms, independently, under sharp regularity conditions on the coefficients and the boundaries. In particular, the proof in this paper does not employ Fefferman-Phong type inequalities which are essential tools in the linear cases in [3, 47].

1. INTRODUCTION

In this paper we study L^q -regularity theory for the following nonlinear equations of the *p*-Laplacian type with lower order terms:

(1.1)
$$\begin{cases} -\operatorname{div} \mathbf{a}(x, Du) + V|u|^{s-2}u &= -\operatorname{div}\left(|F|^{p-2}F\right) & \text{in} \quad \Omega, \\ u &= 0 & \text{on} \quad \partial\Omega \end{cases}$$

where $1 , <math>1 < s < p^*$ (see (2.1) with $\gamma = p$), Ω is a bounded and open set in \mathbb{R}^n with $n \ge 2$, $V : \mathbb{R}^n \to \mathbb{R}$ is nonnegative and called a *potential*, and $F \in L^p(\Omega, \mathbb{R}^n)$. The nonlinearity $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ is assumed to be a vectorvalued Carathéodory function (i.e., \mathbf{a} is measurable in the *x*-variable and continuous in the ξ -variable) of *p*-Laplacian type whose prototype is

(1.2)
$$\mathbf{a}(x,\xi) = (A(x)\xi \cdot \xi)^{\frac{p-2}{2}}A(x)\xi,$$

where $A: \mathbb{R}^n \to \mathbb{R}^{n^2}$ is an $n \times n$ matrix satisfying that

$$u|\eta|^2 \le A(x)\eta \cdot \eta \text{ and } |A(x)| \le L, \quad x,\eta \in \mathbb{R}^n,$$

for some $0 < \nu \leq L$.

For equations (1.1) with identically zero potential function, i.e. $V \equiv 0$, the L^q -regularity theory has been extensively studied after pioneering work of Calderón and Zygmund in [14] where L^q -regularity estimates for Poisson equations $-\Delta u = f$ or div F were proved. In particular, Byun and Wang established the global L^q -regularity for the linear equations with Bounded Mean Oscillation(BMO) coefficients in Reifenberg flat domains [12]. We also refer to [39, 19] and references therein for more L^q -regularity results of the linear equations. With regard to nonlinear equations of the p-Laplacian type, Iwaniec [28] first obtained L^q -regularity estimates for

²⁰¹⁰ Mathematics Subject Classification. Primary 35J10, 35J92; Secondary 35J25, 35B65.

Key words and phrases. Schrödinger operator; L^q -estimates; nonnegative potential; p-Laplacian.

M. Lee was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea Government (NRF-2019R1F1A1061295). J. Ok was supported by the National Research Foundation of Korea funded by the Korean Government (NRF-2017R1C1B2010328).

the simplest case $\mathbf{a}(x,\xi) = |\xi|^{p-2}\xi$, and Caffarelli and Peral [13] considered general $\mathbf{a}(x,\xi)$ that can be discontinuous for x-variable, see also [11, 17, 34, 32] for further results. Based on those works, L^q -regularity theory has been actively developed for various equations generalized from the *p*-Laplace equations, for instance, parabolic equations of the *p*-Laplacian [2, 9, 5], elliptic equations with nonstandard growth [1, 8, 10, 16].

On the other hand, in the case $V \neq 0$, Shen [47] obtained various L^q estimates for the following linear elliptic equation:

(1.3)
$$(-\Delta + V)u = -\Delta u + Vu = \operatorname{div} F \text{ (or } f)$$

i.e. with the right hand side in divergence form or in nondivergence form. Here, the linear operator $-\Delta + V$ is called the Schrödinger operator since the above equations were motivated by the (normalized) Schrödinger equation

$$iu_t = -\Delta u + Vu.$$

In particular, in [47], it is shown that if the potential function V satisfies the reverse Hölder type condition in (2.2) (i.e., $V \in \mathcal{B}_{\gamma}$) for some $\gamma \geq \frac{n}{2}$, then the following estimates hold:

(1.4)
$$\begin{cases} \|Du\|_{L^{2q}(\mathbb{R}^n)} \le c \|F\|_{L^{2q}(\mathbb{R}^n)}, & \frac{(\gamma^*)'}{2} \le q \le \frac{\gamma^*}{2} \ (\frac{1}{2} < q < \infty \text{ if } \gamma \ge n), \\ \|V^{\frac{1}{2}}|u\|\|_{L^{2q}(\mathbb{R}^n)} \le c \|F\|_{L^{2q}(\mathbb{R}^n)}, & \frac{(\gamma^*)'}{2} \le q \le \gamma \ (\frac{1}{2} < q < \gamma \text{ if } \gamma \ge n), \end{cases}$$

where the constants c > 0 depend only on n, q, γ , and b_{γ} . Later, Auscher and Ben Ali [3] extended the range of γ such that $\gamma > 1$ and proved the first estimate in (1.4) whenever $\frac{1}{2} < q \leq \max\{\frac{\gamma^*}{2}, \gamma\}$ and the second one whenever $\frac{1}{2} < q \leq \gamma$, by improving the techniques used in [47] and applying the regularity results for the equation (1.3). In addition, Shen's results in [47] have been extended to linear equations with variable coefficients by Bramanti, Brandolini, Harboure and Viviani [7] and Pan and Tang [42]. We further refer to [15, 18, 29, 30, 45, 46, 33, 52] and references therein for regularity theory relevant to the Schrödinger type elliptic equations.

From the equation (1.3), it is natural to consider the following semi-linear equation:

(1.5)
$$-\Delta u + V|u|^{s-2}u = \operatorname{div} F \quad (\text{or } f),$$

where $1 < s < 2^*$. For the basic theory of semi-linear elliptic equations, we refer to, for instance, [6] and references therein. The equation (1.5) is the Euler-Lagrange equation of the following energy functional:

$$v \in W^{1,2}(\Omega) \quad \mapsto \quad \int_{\Omega} |Dv|^2 + V|v|^s + F \cdot Dv \, dx.$$

We also note that time-independent inhomogeneous nonlinear Schrödinger(INLS) equations reduce to the equation (1.5) with $F \equiv 0$. We refer to [20, 22, 31, 38, 44] and references therein for INLS equations and [4, 24, 36, 50] for their physical applications. For the equations (1.5), however, in our best knowledge, no systematic L^q regularity result has been reported.

Our main equation (1.1) is a generalized version of (1.5). Indeed, the equation (1.1) with (1.2) is the Euler-Lagrange equation of

$$v \in W^{1,p}(\Omega) \quad \mapsto \quad \int_{\Omega} (A(x)Dv \cdot Dv)^{\frac{p}{2}} + V|v|^s + |F|^{p-2}F \cdot Dv \, dx, \quad 1$$

Recently, for the equation (1.1) in the case s = p with $V \in \mathcal{B}_{\gamma}$ and $\frac{n}{p} < \gamma < n$, the authors [35] derived the following local and global L^q estimates for $1 \le q \le \frac{\gamma^*(p-1)}{p}$:

$$\int_{B_r} \left[|Du|^p + V|u|^p \chi_{\{q \le \gamma\}} \right]^q dx \le c \left(\int_{B_{2r}} |Du|^p + V|u|^p dx \right)^q + c \int_{\Omega_{2r}} |F|^{pq} dx$$

with $B_{2r} \Subset \Omega$ and $r \leq R_0$, and

$$\|Du\|_{L^{pq}(\Omega)} + \|V^{\frac{1}{p}}|u|\|_{L^{pq}(\Omega)}\chi_{\{q\leq\gamma\}} \le c_0\|F\|_{L^{pq}(\Omega)}$$

Here, $\chi_{\{q \leq \gamma\}} := 1$ or 0 when $q \leq \gamma$ or $q > \gamma$, respectively, and c_0 and R_0 depend on $\|V\|_{L^{\gamma}(\Omega)}$. We note that the ranges of γ and q are extended from the ones in (1.4) when p = 2. (In fact, a naturally extended range could be $\frac{(\gamma^*)'(p-1)}{p} < q \leq \frac{\gamma^*(p-1)}{p}$, but the case q < 1 is a famous open problem even when $V \equiv 0$, see [28].) However, these results do not cover the ones in [3]. In addition, the above resulting estimates are not sharp since R_0 and c_0 depend on $\|V\|_{L^{\gamma}}$, whereas the estimates for the linear case in [3, 47] are independent of $\|V\|_{L^{\gamma}}$. Moreover, the estimates were derived for $|Du|^p$ and $V|u|^p$ considered together. We also refer to [51] for regularity estimates in the Lorentz spaces.

In this paper, we establish L^q estimates for Du and $V|u|^s$ with possibly sharp ranges of the exponents γ and q. In particular, we deal with the estimates for Duand $V|u|^s$, separately, and find both local and global estimates that are independent of $||V||_{L^{\gamma}}$. Our results extend the known linear regularity results, especially in [3] to nonlinear setting. Furthermore, a BMO, possibly discontinuous, nonlinearity **a** for x-variable and non-smooth domain Ω that beyonds the Lipschitz category are considered as our regularity assumptions.

The main difficulty is that we cannot take advantage of the techniques for linear operators that were used in [47, 3], since we deal with nonlinear problems. Instead, we apply various estimates and techniques used in the study of the regularity theory to our problems. For instance, for (1.1) with $F \equiv 0$, we employ L^{∞} estimates and Calderón-Zygmund estimates, and use an iteration argument. As a consequence, additionally using the reverse Hölder condition of the potential function V, we obtain reverse Hölder type inequalities for $|Dh|^p$ and $V|h|^s$, where h is a weak solution to a localized equation of (1.1) with $F \equiv 0$ (see Theorems 3.6 and 3.7). In particular, we stress that we do not make use of Fefferman-Phong inequality in [21] or its variation that plays an important role in the proofs of L^q estimates in [3, 35, 47] and in turn, the approach in this paper is simpler than eariler ones. Moreover, we obtain comparison estimates for the gradient of solutions and the lower order terms separately (see Lemma 4.1). Therefore, we can handle them independently in the final proof of L^q estimates to discover better resulting estimates.

The remaining is organized as follows. In the next section, we state our main results. Section 3 contains various regularity estimates for the homogenous equations with auxiliary lemmas. Lastly, we prove our main results in Section 4.

2. Main result

2.1. **Preliminaries.** We start with standard notation and definitions. We write $B_r(y)$ for the open ball in \mathbb{R}^n with center $y \in \mathbb{R}^n$ and radius r > 0. We denote $\Omega_r(y) = B_r(y) \cap \Omega$ and $\partial_w \Omega_r(y) = B_r(y) \cap \partial \Omega$. For the sake of simplicity, we write $B_r = B_r(0)$ and $\Omega_r = \Omega_r(0)$. For a measurable function $g: U \to \mathbb{R}$ with $U \subset \mathbb{R}^n$,

we define

$$(g)_U := \int_U g \, dx = \frac{1}{|U|} \int_U g \, dx,$$

$$g_+ := \max\{g, 0\} \quad \text{and} \quad g_- := (-g)_+ = \max\{-g, 0\}.$$

For $g \in W^{1,p}(\Omega_r(y))$, the boundary condition "g = 0 on $\partial_w \Omega_r$ " means the zero

For $g \in W^{1,p}(\Omega_r(y))$, the boundary condition "g = 0 on $\mathcal{O}_{\mathbf{w}}\Omega_r$ " means the zer extension of g to $B_r(y)$ is in $W^{1,p}(B_r(y))$. We also define

(2.1)
$$\gamma^* := \begin{cases} \frac{n\gamma}{n-\gamma}, & \text{when } 1 < \gamma < n, \\ \infty, & \text{when } \gamma \ge n. \end{cases}$$

We say that a nonnegative function $V : \mathbb{R}^n \to [0, \infty)$ belongs to \mathcal{B}_{γ} for some $\gamma > 1$ if $V \in L^{\gamma}_{loc}(\mathbb{R}^n)$ and there exists a constant $b_{\gamma} > 0$ such that the reverse Hölder inequality

(2.2)
$$\left(\frac{1}{|B|}\int_{B}V^{\gamma} dx\right)^{\frac{1}{\gamma}} \leq b_{\gamma}\left(\frac{1}{|B|}\int_{B}V dx\right)$$

holds for every ball B in \mathbb{R}^n . This \mathcal{B}_{γ} class, which is a wide class including all nonnegative polynomials, was introduced independently by Muckenhoupt [37] and Gehring [23] in the study of weighted norm inequalities and quasi-conformal mapping, respectively. One notable example of this element is $V(x) = |x|^{-n/\gamma}$ which actually belongs to the $\mathcal{B}_{\tilde{\gamma}}$ class for all $\tilde{\gamma} < \gamma$. Moreover, the \mathcal{B}_{γ} class is strongly connected to the Muckenhoupt classes A_p . We say that a nonnegative function $w \in L^1_{\text{loc}}(\mathbb{R}^n)$ is in the A_p class for some $1 \leq p < \infty$, denoted by $w \in A_p$, if and only if

$$[w]_p := \sup_B \left(\oint_B w \, dx \right) \left(\oint_B w^{-\frac{1}{p-1}} \, dx \right)^{p-1} < \infty,$$

where the supremum is taken over all balls $B \subset \mathbb{R}^n$. Then, we have the following equivalent relations:

(2.3)
$$V \in A_p \text{ for some } p > 1 \iff V \in \mathcal{B}_{\gamma} \text{ for some } \gamma > 1$$

Here, if we consider only the left arrow, the constant p and $[V]_p$ are determined by γ and b_{γ} . We refer to [26, Theorem 9.3.3] for its proof and more details on properties and relations of those classes.

2.2. Main result. We introduce the main result in this paper.

We first recall the equation (1.1) with the basic setting in the first paragraph in Introduction, Section 1. The nonnegative potential $V : \mathbb{R}^n \to [0, \infty)$ satisfies that $V \in L^{\gamma_0}(\Omega)$, where

(2.4)
$$\gamma_0 := \begin{cases} \frac{np}{np-s(n-p)}, & \text{when } 1 n. \end{cases}$$

Note that $\gamma_0 = \frac{n}{p}$ if $s = p \in (1, n)$. The nonlinearity $\mathbf{a}(x, \xi)$ is assumed that $\mathbf{a}(x, \cdot) \in C^1(\mathbb{R}^n \setminus \{0\}, \mathbb{R}^n)$ for each $x \in \Omega$ and satisfies the following growth and ellipticity conditions:

(2.5)
$$|\mathbf{a}(x,\xi)| + |D_{\xi}\mathbf{a}(x,\xi)||\xi| \le L|\xi|^{p-1}$$

and

(2.6)
$$D_{\xi}\mathbf{a}(x,\xi)\,\eta\cdot\eta\geq\nu|\eta|^2|\xi|^{p-2}$$

for almost all $x \in \mathbb{R}^n$ and any $\xi, \eta \in \mathbb{R}^n (\xi \neq 0)$ and for some constants L, ν with $0 < \nu \leq 1 \leq L$. We remark that the condition (2.6) implies the monotonicity condition:

(2.7)
$$(\mathbf{a}(x,\xi) - \mathbf{a}(x,\eta)) \cdot (\xi - \eta) \ge c(p,\nu) \left(|\xi|^2 + |\eta|^2 \right)^{\frac{p-2}{2}} |\xi - \eta|^2$$

for any $\xi, \eta \in \mathbb{R}^n$ and a.e. $x \in \mathbb{R}^n$. Under the above setting, we say that $u \in W_0^{1,p}(\Omega)$ is a weak solution to the problem (1.1) if

$$\int_{\Omega} \mathbf{a}(x, Du) \cdot D\varphi \, dx + \int_{\Omega} V|u|^{s-2} u\varphi \, dx = \int_{\Omega} |F|^{p-2} F \cdot D\varphi \, dx$$

holds for any $\varphi \in W_0^{1,p}(\Omega)$. Under the above setting, the existence and the uniqueness of the weak solution of (1.1) follow from the theory of nonlinear functional analysis, see for instance [48, Chapter 2].

The following two definitions are related to our main assumptions imposed on the nonlinearlity \mathbf{a} and the domain Ω .

Definition 2.1. We say that $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ is (δ, R) -vanishing if

$$\sup_{0<\rho\leq R} \sup_{y\in\mathbb{R}^n} \oint_{B_{\rho}(y)} \left|\Theta\left(\mathbf{a}, B_{\rho}(y)\right)(x)\right| \, dx \leq \delta,$$

where

$$\Theta\left(\mathbf{a}, B_{\rho}(y)\right)(x) := \sup_{\xi \in \mathbb{R}^n \setminus \{0\}} \frac{\left|\mathbf{a}(x, \xi) - (\mathbf{a}(\cdot, \xi))_{B_{\rho}(y)}\right|}{|\xi|^{p-1}}$$

and

$$(\mathbf{a}(\cdot,\xi))_{B_{\rho}(y)} := \int_{B_{\rho}(y)} \mathbf{a}(x,\xi) \, dx.$$

The above definition implies that the map $x \mapsto \mathbf{a}(x,\xi)/|\xi|^{p-1}$ is a locally BMO function with the BMO semi-norm which is less than or equals to δ for all $\xi \in \mathbb{R}^n$. Then we note that the nonlinearity \mathbf{a} can be discontinuous for the *x*-variable. In particular, in the model case in (1.2), Definition 2.1 implies that $A(\cdot)$ is a locally BMO function.

Definition 2.2. Given $\delta \in (0, \frac{1}{8})$ and R > 0, we say that Ω is a (δ, R) -Reifenberg flat domain if for every $x \in \partial \Omega$ and every $\rho \in (0, R]$, there exists a coordinate system $\{y_1, y_2, \ldots, y_n\}$ which may depend on ρ and x, such that in this coordinate system x = 0 and that

$$B_{\rho}(0) \cap \{y_n > \delta\rho\} \subset B_{\rho}(0) \cap \Omega \subset B_{\rho}(0) \cap \{y_n > -\delta\rho\}.$$

In the above definition, δ is usually supposed to be less than $\frac{1}{8}$. This number follows from the Sobolev embedding, see for instance [49]. In this paper, however, it is not important since we will deal with sufficiently small δ . We remark that the Lipschitz domains with the Lipschitz constant which is less than or equal to δ belong to the class of (δ, R) -Reifenberg flat domains for some R > 0. In addition, the (δ, R) -Reifenberg flat domain Ω has the following measure density condition:

(2.8)
$$\sup_{0<\rho\leq R} \sup_{y\in\overline{\Omega}} \frac{|B_{\rho}(y)|}{|\Omega\cap B_{\rho}(y)|} \le \left(\frac{2}{1-\delta}\right)^n \le \left(\frac{16}{7}\right)^n$$

We refer to [12, 41, 43, 49] for more details on the Reifenberg flat domains and their applications.

.

Now we present the main results in this paper. The first result is local L^q estimates in both interior and boundary regions.

Theorem 2.3. Let $1 , <math>1 < s < p^*$, γ_0 be from (2.4), $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ satisfy (2.5) and (2.6), $V : \mathbb{R}^n \to [0, \infty)$ with $V \in L^{\gamma_0}(\Omega)$, and $F \in L^p(\Omega, \mathbb{R}^n)$. Suppose that the constants p, γ , q satisfy specific conditions given below and that $V \in \mathcal{B}_{\gamma}$. There exists a small $\delta = \delta(n, p, L, \nu, \gamma) > 0$ such that if \mathbf{a} is (δ, R) -vanishing, Ω is a (δ, R) -Reifenberg flat domain for some $R \in (0, 1)$, and $u \in W_0^{1, p}(\Omega)$ is a weak solution to (1.1), then for any $x_0 \in \overline{\Omega}$ and $r \in (0, \frac{R}{2}]$ we have the following estimates:

(1) If
$$1 , $\tilde{\gamma} < \gamma < \infty$ where$$

(2.9)
$$\tilde{\gamma} := \max\left\{1, \frac{np}{np - n + p}\right\} = \left\{\begin{array}{cc} \frac{np}{np - n + p} & \text{if } 1
$$and \ 1 < q < \frac{\gamma^*(p-1)}{p},$$$$

(2.10)
$$\int_{\Omega_r(x_0)} |Du|^{pq} \, dx \le c \left(\int_{\Omega_{2r}(x_0)} |Du|^p \, dx \right)^q + c \int_{\Omega_{2r}(x_0)} |F|^{pq} \, dx.$$

(2) If
$$p \ge 2$$
, $1 < \gamma < \infty$, and $1 < q < \gamma$

(2.11)
$$\int_{\Omega_r(x_0)} [V|u|^s]^q \, dx \le c \left(\int_{\Omega_{2r}(x_0)} V|u|^s \, dx \right)^q + c \int_{\Omega_{2r}(x_0)} |F|^{pq} \, dx.$$

(3) If
$$1 , $\frac{n}{p} \le \gamma < \infty$ and $1 < q < \gamma$$$

(2.12)
$$\int_{\Omega_r(x_0)} [V|u|^s]^q \, dx \le c \left(\int_{\Omega_{2r}(x_0)} |Du|^p + V|u|^s \, dx \right)^q + c \int_{\Omega_{2r}(x_0)} |F|^{pq} \, dx.$$

Here the constants c > 0 depend on $n, p, L, \nu, s, \gamma, b_{\gamma}, q$.

Regarding the ranges of γ and q, we will discuss in Remarks 2.6 and 2.7 below.

Remark 2.4. In the above theorem, we obtain local L^q estimates for $|Du|^p$ without the lower order term $V|u|^s$, and for $V|u|^s$ without the term $|Du|^p$ when p > 2. However, when $1 , the local <math>L^q$ estimates for $V|u|^s$ involve $|Du|^p$, which follows from the approximation lemma, Lemma 4.1. As a consequence, the lower bound of γ in (3) has to be chosen as not 1 but $\frac{n}{p}$. This choice insures $\frac{\gamma^*(p-1)}{p} \ge \gamma$, hence we have $|Dh_i|^p + V|h_i|^s \in L^{\gamma}(\Omega_{5\rho_i}(y_i))$ in (4.16) with Case 3 in the proof of Theorem 2.3.

The second result is global L^q estimates. As a consequence of Theorem 2.3, we obtain the following global estimates by using the standard covering argument in the proof of [35, Corollary 2.6], hence we omit the proof.

Theorem 2.5. Let $1 , <math>1 < s < p^*$, γ_0 be from (2.4), $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ satisfy (2.5) and (2.6), $V : \mathbb{R}^n \to [0, \infty)$ with $V \in L^{\gamma_0}(\Omega)$, and $F \in L^p(\Omega, \mathbb{R}^n)$. Suppose that the constants p, γ, q satisfy specific conditions given below and that $V \in \mathcal{B}_{\gamma}$. There exists a small $\delta = \delta(n, p, L, \nu, \gamma) \in (0, \frac{1}{8})$ such that if \mathbf{a} is (δ, R) -vanishing, Ω is a (δ, R) -Reifenberg flat domain for some $R \in (0, 1)$, and $u \in W_0^{1, p}(\Omega)$ is a weak solution to (1.1), then we have the following estimates: (1) If $1 , <math>\tilde{\gamma} < \gamma < \infty$ where $\tilde{\gamma}$ is given in (2.9), and $1 < q < \frac{\gamma^*(p-1)}{p}$, then we have

$$\|Du\|_{L^{pq}(\Omega)} \le c \left(\frac{\operatorname{diam}(\Omega)}{R}\right)^{n(q-1)} \|F\|_{L^{pq}(\Omega)}.$$

(2) If $p \ge 2$, $1 < \gamma < \infty$ and $1 < q < \gamma$, or if $1 , <math>\frac{n}{p} \le \gamma < \infty$ and $1 < q < \gamma$, then we have

$$\|V^{\frac{1}{p}}|u|^{\frac{s}{p}}\|_{L^{pq}(\Omega)} \le c \left(\frac{\operatorname{diam}(\Omega)}{R}\right)^{n(q-1)} \|F\|_{L^{pq}(\Omega)}.$$

Here the constants c > 0 depend on $n, p, L, \nu, s, \gamma, b_{\gamma}, q$.

Remark 2.6. If $V \in \mathcal{B}_{\gamma}$, then V belongs to the $\mathcal{B}_{\gamma+\epsilon}$ class for some small $\epsilon > 0$ from the self improving property of the \mathcal{B}_{γ} class (see [23]). Therefore, by considering $\gamma + \epsilon$ instead of γ in Theorems 2.3 and 2.5, the ranges of q can be extended to $q \in [1, \frac{\gamma^*(p-1)}{p}]$ and $q \in [1, \gamma]$, respectively. Note that the case q = 1 is trivial. On the other hand, as mentioned in Introduction, L^q estimates with q < 1 is an open problem even in the case $V \equiv 0$.

Remark 2.7. We further comment on the ranges of γ and q for the L^q estimates for $|Du|^p$ in Theorem 2.3 (1) and Theorem 2.5 (1).

- (i) Suppose p < n. Then the range of γ is $(\frac{np}{np-n+p}, \infty)$. On the other hand, for the linear case with p = 2, we see from [3] that it is $(1, \infty)$. Therefore, the case $\gamma \in (1, \frac{np}{np-n+p})$ seems missing in our results. However, if $\gamma < \frac{np}{np-n+p}$, we have $\frac{\gamma^*(p-1)}{p} < 1$ and the L^q estimate with q < 1 is the open problem mentioned above. Therefore, our range is best for now.
- (ii) We note that $\frac{\gamma^*(p-1)}{p} < \gamma$ when $\gamma < \frac{n}{p}$. For the linear case with p = 2, in [3], the range of q is $(\frac{1}{2}, \gamma)$ when $\gamma < \frac{n}{2}$. Therefore, the range of q in our results could be extended to $1 < q < \gamma$ when $\gamma < \frac{n}{p}$.

Remark 2.8. From Remark 4.4, we see that the estimates in Theorem 2.3 (2) and Theorem 2.5 (2) when $p \ge 2$ still hold without the (δ, R) -vanishing condition of **a** and the (δ, R) -Reifenberg flat condition of Ω . Moreover, in Theorem 2.3 (1) and Theorem 2.5 (1), the dependence γ of δ is replaced with q, when $\gamma \ge n$.

3. Estimates for homogenous equations

In this section, we prove various regularity estimates for weak solutions to localized equations of (1.1) with $F \equiv 0$.

We start by recalling interior and boundary Calderón-Zygmund estimates for *p*-Laplace type elliptic equations. In particular, we consider non-divergence data. For the following results, we refer to, for instance, [34].

Lemma 3.1. Let $1 and <math>\tilde{\gamma} < \gamma < n$ where $\tilde{\gamma}$ is given in (2.9). There exists a small $\delta = \delta(n, p, L, \nu, \gamma) \in (0, \frac{1}{8})$ so that if **a** is (δ, R) -vanishing and Ω is a (δ, R) -Reifenberg flat domain for some $R \in (0, 1)$, then for any $x_0 \in \overline{\Omega}$, $r \in (0, \frac{R}{2}]$ and for any weak solution $h \in W^{1,p}(\Omega_{2r}(x_0))$ to

$$\begin{cases} -\operatorname{div} \mathbf{a}(x, Dh) = f & in & \Omega_{2r}(x_0), \\ h = 0 & on & \partial_{w}\Omega_{2r}(x_0) \text{ if } B_{2r}(x_0) \not\subset \Omega. \end{cases}$$

with $f \in L^{\gamma}(\Omega_{2r}(x_0))$, we have

$$\left(\oint_{\Omega_r(x_0)} |Dh|^{\gamma^*(p-1)} \, dx \right)^{\frac{p}{\gamma^*(p-1)}} \le c \oint_{\Omega_{2r}(x_0)} |Dh|^p \, dx + c \left(\oint_{\Omega_{2r}(x_0)} |rf|^\gamma \, dx \right)^{\frac{p}{\gamma(p-1)}}$$

for some $c = c(n, p, L, \nu, \gamma) > 0$.

We need the standard iteration lemma whose proof can be found in, for instance, [27].

Lemma 3.2. Let $g : [a,b] \to \mathbb{R}$ be a bounded nonnegative function. Suppose that for any τ_1, τ_2 with $0 < a \le \tau_1 < \tau_2 \le b$,

$$g(\tau_1) \le \tau g(\tau_2) + \frac{C_1}{(\tau_2 - \tau_1)^{\beta}} + C_2$$

where $C_1, C_2 \ge 0, \beta > 0$ and $0 \le \tau < 1$. Then we have

$$g(\tau_1) \le c\left(\frac{C_1}{(\tau_2 - \tau_1)^\beta} + C_2\right)$$

for some constant $c = c(\beta, \tau) > 0$.

Now , we consider the following homogeneous Dirichlet problems of the following type

(3.1)
$$\begin{cases} -\operatorname{div} \mathbf{a}(x, Dh) + V|h|^{s-2}h &= 0 \quad \text{in } \Omega_r, \\ h &= 0 \quad \text{on } \partial_{\mathbf{w}}\Omega_r, \text{ if } B_r \not\subset \Omega, \end{cases}$$

where $1 , <math>1 < s < p^*$, $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ satisfies

(3.2)
$$|\mathbf{a}(x,\xi)| \le L|\xi|^{p-1} \text{ and } \mathbf{a}(x,\xi) \cdot \xi \ge \nu|\xi|^p, \quad x,\xi \in \mathbb{R}^n,$$

for some $0 < \nu \leq L$, and $V : \mathbb{R}^n \to [0,\infty)$ does $V \in L^{\gamma_0}(\Omega)$. Note that the assumptions (2.5) and (2.6) imply (3.2).

We shall need the following Caccioppoli type estimates. For simplicity, we use both the plus-minus sign \pm and the minus-plus sign \mp , and they are all linked, that is, we take all upper signs or all lower signs.

Lemma 3.3. (Caccioppoli estimates) Under the setting above, let $h \in W^{1,p}(\Omega_r)$ be a weak solution to (3.1). Then for every $k \ge 0$, every $B_{\rho}(y) \subset B_r$ with $B_{\rho}(y) \cap \Omega_r \ne \emptyset$, and every $\nu \in (0,1)$, we have (3.3)

$$\int_{\Omega_{\nu\rho}(y)} |D(h\mp k)_{\pm}|^p \, dx + \int_{\Omega_{\nu\rho}(y)} V|h|^{s-2} h_{\pm}(h\mp k)_{\pm} \, dx \le c \int_{\Omega_{\rho}(y)} \left[\frac{(h\mp k)_{\pm}}{(1-\nu)r} \right]^p \, dx$$

for some $c\tilde{\gamma} < \gamma < n$, where $\tilde{\gamma}$ is given in (2.9). onstant $c = c(n, p, L, \nu) > 0$.

Proof. We take $\pm (h \mp k)_{\pm} \eta^p$ as a test function in the weak form of (3.1), where $\eta \in C_0^{\infty}(B_{\rho}(y))$ such that $0 \leq \eta \leq 1, \eta \equiv 1$ on $B_{\nu\rho}(y)$, and $|D\eta| \leq \frac{c(n)}{(1-\nu)r}$. Note that $\pm (h \mp k)_{\pm} \eta^p = 0$ on $\partial_w \Omega_r$ since h = 0 on $\partial_w \Omega_r$ and $k \geq 0$. Then we have

$$\int_{\Omega_{\rho}(y)} \mathbf{a}(x, Dh) \cdot D[\pm (h \mp k)_{\pm} \eta^{p}] \, dx + \int_{\Omega_{\rho}(y)} V|h|^{s-2} h[\pm (h \mp k)_{\pm} \eta^{p}] \, dx = 0.$$

Note that since $k \ge 0$, $Dh = \pm D(h \mp k)_{\pm}$ and $\pm h = h_{\pm}$ when $(h \mp k)_{\pm} > 0$. Moreover, applying (3.2), we find

$$\mathbf{a}(x,Dh) \cdot D[\pm(h\mp k)_{\pm}\eta^{p}] = \pm \mathbf{a}(x,\pm D(h\mp k)_{\pm}) \cdot [\eta^{p}D(h\mp k)_{\pm} + (h\mp k)_{\pm}D\eta^{p}]$$

$$\geq \nu\eta^{p}|D(h\mp k)_{\pm}|^{p} - c\eta^{p-1}|D(h\mp k)_{\pm}|^{p-1}\frac{(h\mp k)_{\pm}}{(1-\nu)r}.$$

Therefore, by the above results and the second inequality in (3.2), we have

$$\begin{split} \nu \int_{\Omega_{\rho}(y)} \eta^{p} |D(h \mp k)_{\pm}|^{p} \, dx + \int_{\Omega_{\rho}(y)} V|h|^{s-2} h_{\pm}(h \mp k)_{\pm} \eta^{p} \, dx \\ &\leq c \int_{\Omega_{\rho}(y)} \eta^{p-1} |D(h \mp k)_{\pm}|^{p-1} \frac{(h \mp k)_{\pm}}{(1-\nu)r} \, dx. \end{split}$$

Using Young's inequality and the properties of η , we have the estimate (3.3).

Next, we prove the local boundedness of the weak solutions to (3.1) with $L^{\infty} - L_w^q$ type estimates.

Lemma 3.4. Under the assumptions of Lemma 3.3, we have that $h \in L^{\infty}_{loc}(B_r)$, where we extend h to B_r by 0. Moreover, for every $0 < r_1 < r_2 \leq r$, every q > 0 and every weight $w \in A_t$ with $t \geq 1$,

(3.4)
$$\|h\|_{L^{\infty}(\Omega_{r_{1}})} \leq c \left(\frac{r_{2}}{r_{2}-r_{1}}\right)^{\frac{nt}{q}} \left(\frac{1}{w(B_{r_{2}})} \int_{\Omega_{r_{2}}} |h|^{q} w \, dx\right)^{\frac{1}{q}}$$
$$= c \left(\frac{r_{2}}{r_{2}-r_{1}}\right)^{\frac{nt}{q}} \left(\frac{1}{(w)_{B_{r_{2}}}} \frac{1}{|B_{r_{2}}|} \int_{\Omega_{r_{2}}} |h|^{q} w \, dx\right)^{\frac{1}{q}}$$

for some constant $c = c(n, p, L, \nu, q, t, [w]_t) > 0$, where $w(B_{r_2}) := \int_{B_{r_2}} w \, dx$.

Proof. Since $V \ge 0$, we have from (3.3) that for every $B_{\rho}(y) \subset B_r$, $k \ge 0$, and $\nu \in (0, 1)$,

$$\int_{\Omega_{\nu\rho}(y)} |D(h\mp k)_{\pm}|^p \, dx \le c \int_{\Omega_{\rho}(y)} \left[\frac{(h\mp k)_{\pm}}{(1-\nu)r} \right]^p \, dx.$$

This implies that h belongs to the De Giorgi class in [25, Chapter 7]. Therefore, in view of [25, Chapter 7.2], we deduce that

(3.5)
$$\|h\|_{L^{\infty}(\Omega_{r_1})} \le c \left(\frac{r_2}{r_2 - r_1}\right)^{\frac{n}{q}} \left(\frac{1}{|B_{r_2}|} \int_{\Omega_{r_2}} |h|^q \, dx\right)^{\frac{1}{q}}$$

for every q > 0, which is the desired estimate (3.4) with the trivial weight $w \equiv 1$. Since $w \in A_t$, we note from [26, Proposition 9.1.5 (8)] that for every $B_{\rho} \subset \mathbb{R}^n$ and every $f \in L^t_w(B_{\rho})$,

1

$$\int_{B_{\rho}} |f| \, dx \le [w]_t \left(\frac{1}{w(B_{\rho})} \int_{B_{\rho}} |f|^t w \, dx \right)^{\frac{1}{t}}$$

Therefore, plugging this inequality with $f = |h|^{q/t}$ and $B_{\rho} = B_{r_2}$ into (3.5) replacing q by q/t, we obtain (3.4).

Using the above results, we deduce the following lemma, which allows us to change the integrangd $V|h|^s$ to $|Dh|^p$ later.

Lemma 3.5. Under the assumptions of Lemma 3.3, let $h \in W^{1,p}(\Omega_r)$ be a weak solution to (3.1). If $V \in B_{\gamma}$ for some $\gamma > 1$, then

$$(3.6) r^{\frac{p}{p-1}}(V)_{B_{r/2}}^{\frac{p}{s(p-1)}} \left(\frac{1}{|B_{r/2}|} \int_{\Omega_{r/2}} V|h|^s \, dx\right)^{\frac{p(s-1)}{s(p-1)}} \le \frac{c}{|B_r|} \int_{\Omega_r} |Dh|^p \, dx$$

for some $c = c(n, p, L, \nu, s, \gamma, b_{\gamma}) > 0$.

Proof. We first note from (2.3) that $V \in A_t$ with $t \ge 1$ and $[V]_t$ depending on γ and b_{γ} . We extend h to B_r by 0. Let $r/2 \le r_1 < r_2 \le r$ be arbitrary, $r_3 := \frac{r_1+r_2}{2}$, and $\eta \in C_0^{\infty}(B_{r_3})$ with $\eta \ge 0$, $\eta \equiv 1$ in B_{r_1} and $|D\eta| \le c/(r_2 - r_1)$. We take $h\eta^p$ as a test function in the weak form of (3.1). Then, in the same argument as in the proof of Lemma 3.3 we have

$$\nu \int_{B_{r_3}} |Dh|^p \eta^p \, dx + \int_{B_{r_3}} V|h|^s \eta^p \, dx \le \frac{c}{r_2 - r_1} \int_{B_{r_3}} \eta^{p-1} |Dh|^{p-1} |h| \, dx.$$

Applying Hölder's inequality, it follows that

$$\int_{B_{r_1}} V|h|^s \, dx \le \frac{c}{r_2 - r_1} \left(\int_{B_{r_3}} |Dh|^p \, dx \right)^{\frac{p-1}{p}} \left(\int_{B_{r_3}} |h|^p \, dx \right)^{\frac{1}{p}}.$$

We multiply the both sides with

$$M := \left[r^p(V)_{B_{r/2}}^{\frac{p}{s}} \left(\oint_{B_{r/2}} V|h|^s \, dx \right)^{\frac{s-p}{s}} \right]^{\frac{1}{p-1}}$$

to obtain

$$\begin{split} M \int_{B_{r_1}} V|h|^s \, dx &\leq \frac{c \, r M^{\frac{1}{p}}}{r_2 - r_1} \left(\int_{B_{r_3}} |Dh|^p \, dx \right)^{\frac{p-1}{p}} \\ & \times \left[\underbrace{(V)_{B_{r/2}}^{\frac{p}{s}} \left(\oint_{B_{r/2}} V|h|^s \, dx \right)^{\frac{s-p}{s}} \int_{B_{r_3}} |h|^p \, dx}_{=:I} \right]^{\frac{1}{p}}. \end{split}$$

(Here, we assume that $(V|h|^s)_{B_{r/2}} > 0$. If the average is zero, the estimate (3.6) is trivial.) We now estimate *I*. By (3.4) with (q, r_1, w) in place of (s, r_3, V) ,

$$\begin{split} I &\leq |B_r|(V)_{B_{r/2}}^{\frac{p}{s}} \left(\oint_{B_{r/2}} V|h|^s \, dx \right)^{\frac{s-p}{s}} \|h\|_{L^{\infty}(B_{r_3})}^p \\ &\leq c|B_r| \left(\frac{r}{r_2 - r_1}\right)^{\frac{npt}{s}} (V)_{B_{r/2}}^{\frac{p}{s}} \left(\oint_{B_{r/2}} V|h|^s \, dx \right)^{\frac{s-p}{s}} \left(\frac{1}{(V)_{B_{r_2}}} \oint_{B_{r_2}} V|h|^s \, dx \right)^{\frac{p}{s}} \\ &\leq c \left(\frac{r}{r_2 - r_1}\right)^{\frac{npt}{s}} \int_{B_{r_2}} V|h|^s \, dx. \end{split}$$

Inserting this into the preceding estimate and using Young's inequality, we have

$$\begin{split} M \int_{B_{r_1}} V|h|^s \, dx &\leq c \left(\frac{r}{r_2 - r_1}\right)^{1 + \frac{nt}{s}} \left(\int_{B_r} |Dh|^p \, dx\right)^{\frac{p-1}{p}} \left[M \int_{B_{r_2}} V|h|^s \, dx\right]^{\frac{1}{p}} \\ &\leq \frac{1}{2} M \int_{B_{r_2}} V|h|^s \, dx + c \left(\frac{r}{r_2 - r_1}\right)^{(1 + \frac{nt}{s})\frac{p}{p-1}} \int_{B_r} |Dh|^p \, dx. \end{split}$$

Theorefore, by applying Lemma 3.2 and recalling the definition of M, we get the conclusion.

Finally, we derive the following two reverse Hölder type higher integrability results for homogeneous equations.

Theorem 3.6. Let $1 , <math>1 < s < p^*$, γ_0 be from (2.4), $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ satisfy (3.2), and $V : \mathbb{R}^n \to [0, \infty)$ do $V \in L^{\gamma_0}(\Omega)$ and $V \in \mathcal{B}_{\gamma}$ for some $\gamma > 1$. If $h \in W^{1,p}(\Omega_{2r}(x_0))$ is a weak solution to

$$\begin{cases} -\operatorname{div} \mathbf{a}(x, Dh) + V|h|^{s-2}h &= 0 \quad in \quad \Omega_{2r}(x_0), \\ h &= 0 \quad on \quad \partial_{\mathbf{w}}\Omega_{2r}(x_0) \quad if \quad B_{2r}(x_0) \not\subset \Omega_{2r}(x_0) \end{cases}$$

where $x_0 \in \overline{\Omega}$, then we have

$$\left(\oint_{\Omega_r(x_0)} [V|h|^s]^\gamma \ dx\right)^{\frac{1}{\gamma}} \le c \oint_{\Omega_{2r}(x_0)} V|h|^s \ dx$$

for some $c = c(n, p, \nu, L, s, \gamma, b_{\gamma}) > 0.$

Proof. Note that, by (2.3), $V \in A_t$ with $t \ge 1$ and $[V]_t$ depending on γ and b_{γ} . Then, the desired estimate directly follows from $V \in \mathcal{B}_{\gamma}$ and Lemma 3.4 with q = s and w = V as

$$\left(\frac{1}{|B_r|} \int_{\Omega_r} [V|h|^s]^\gamma \, dx\right)^{\frac{1}{\gamma}} \le b_\gamma(V)_{B_r} \|h\|_{L^{\infty}(\Omega_r)}^s \le \frac{c}{|B_{2r}|} \int_{\Omega_{2r}} V|h|^s \, dx. \qquad \Box$$

Theorem 3.7. Let $1 , <math>1 < s < p^*$, γ_0 be from (2.4), $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ satisfy (2.5) and (2.6), and $V : \mathbb{R}^n \to [0, \infty)$ do $V \in L^{\gamma_0}(\Omega)$ and $V \in \mathcal{B}_{\gamma}$ for some γ satisfying $\tilde{\gamma} < \gamma < n$, where $\tilde{\gamma}$ is given in (2.9). There exists a small $\delta = \delta(n, p, L, \nu, \gamma) > 0$ such that the following holds: if \mathbf{a} is (δ, R) -vanishing, Ω is a (δ, R) -Reifenberg flat domain for some R > 0, and $h \in W^{1,p}(\Omega_{8r}(x_0))$ is a weak solution to

$$\begin{cases} -\operatorname{div} \mathbf{a}(x, Dh) + V|h|^{s-2}h = 0 & in \quad \Omega_{8r}(x_0), \\ h = 0 & on \quad \partial_{w}\Omega_{8r}(x_0) & if \quad B_{8r}(x_0) \not \subset \Omega_{8r}(x_0) \end{cases}$$

where $x_0 \in \overline{\Omega}$ and $8r \leq R$, then we have

$$\left(\oint_{\Omega_r(x_0)} |Dh|^{\gamma^*(p-1)} dx\right)^{\frac{p}{\gamma^*(p-1)}} \le c \oint_{\Omega_{8r}(x_0)} |Dh|^p dx$$

for some $c = c(n, p, \nu, L, s, \gamma, b_{\gamma}) > 0$.

Proof. For simplicity, we write $\Omega_{\rho} = \Omega_{\rho}(x_0)$ with $\rho > 0$. We extend h by 0 to $B_{16r} \setminus \Omega$ if it is nonempty. From the facts that $h \in L^{\infty}(B_{2r})$ and $V \in L^{\gamma}(\Omega_{2r})$, we

see that $V|h|^{s-2}h \in L^{\gamma}(\Omega_{2r})$. Therefore, applying Lemma 3.1 with $f = V|h|^{s-2}h$, we have

(3.7)
$$\begin{pmatrix} \int_{\Omega_r} |Dh|^{\gamma^*(p-1)} dx \end{pmatrix}^{\frac{p}{\gamma^*(p-1)}} \\ \leq c \int_{\Omega_{2r}} |Dh|^p dx + c \left(\int_{\Omega_{2r}} \left[rV|h|^{s-1} \right]^{\gamma} dx \right)^{\frac{p}{\gamma(p-1)}}.$$

We now estimate the second integral on the right hand side of (3.7). Since $V \in \mathcal{B}_{\gamma}$ and $|\Omega_{\rho}| \approx |B_{\rho}|$ with $0 < \rho < R$, see (2.8),

$$\left(\int_{\Omega_{2r}} [rV|h|^{s-1}]^{\gamma} dx\right)^{\frac{p}{\gamma(p-1)}} \le b_{\gamma} r^{\frac{p}{p-1}} (V)^{\frac{p}{p-1}}_{B_{2r}} \|h\|_{L^{\infty}(B_{2r})}^{\frac{p(s-1)}{p-1}}$$

Moreover, applying Lemma 3.4, replacing (r, r_1, r_2, w) by (4r, 2r, 4r, V), and Lemma 3.5, replacing r with 8r

$$\begin{split} \left(\oint_{\Omega_{2r}} [rV|h|^{s-1}]^{\gamma} \, dx \right)^{\frac{p}{\gamma(p-1)}} &\leq b_{\gamma} r^{\frac{p}{p-1}} (V)_{B_{2r}}^{\frac{p}{p-1}} \|h\|_{L^{\infty}(B_{2r})}^{\frac{p}{p-1}} \\ &\leq cr^{\frac{p}{p-1}} (V)_{B_{4r}}^{\frac{p}{p-1}} \left(\frac{1}{(V)}_{B_{4r}} f_{B_{4r}} V|h|^{s} \, dx \right)^{\frac{p(s-1)}{s(p-1)}} \\ &\leq cr^{\frac{p}{p-1}} (V)_{B_{4r}}^{\frac{p}{s(p-1)}} \left(f_{B_{4r}} V|h|^{s} \, dx \right)^{\frac{p(s-1)}{s(p-1)}} \\ &\leq c f_{\Omega_{8r}} |Dh|^{p} \, dx. \end{split}$$

Plugging the preceding estimate into (3.7) we have the desired estimate.

Remark 3.8. In the above theorems, it is possible that $\gamma < \gamma_0$. In this case, since $V \in L^{\gamma_0}$, we have $V|h|^s \in L^{\gamma_0}(\Omega_r(x_0))$ in Theorem 3.6 and $Dh \in L^{\gamma_0^*(p-1)}(\Omega_r(x_0), \mathbb{R}^n)$ in Theorem 3.7. However, the reverse Hölder type estimates can be obtained with the exponent γ , by the assumption that $V \in \mathcal{B}_{\gamma}$.

4. L^q estimates

We are now ready to prove our main results.

4.1. Comparison. We start with the following comparison lemma.

Lemma 4.1. Let $1 , <math>1 < s < p^*$, γ_0 be from (2.4), $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ satisfy (2.5) and (2.6), $V : \mathbb{R}^n \to [0, \infty)$ do $V \in L^{\gamma_0}(\Omega)$, and $F \in L^p(\Omega, \mathbb{R}^n)$. If $u \in W_0^{1,p}(\Omega)$ is the weak solution to (1.1), and $h \in W^{1,p}(\Omega_{8r})$ is the weak solution to

(4.1)
$$\begin{cases} -\operatorname{div} \mathbf{a}(x, Dh) + V|h|^{s-2}h &= 0 \quad in \quad \Omega_{8r}, \\ h &= u \quad on \quad \partial\Omega_{8r}, \end{cases}$$

where $\Omega_{8r} = \Omega_{8r}(x_0)$ with $x_0 = \overline{\Omega}$, then we have the following estimates: (i) (Energy estimates)

(4.2)
$$\int_{\Omega_{8r}} |Dh|^p \, dx \le c \int_{\Omega_{8r}} |Du|^p \, dx + c \int_{\Omega_{8r}} |F|^p \, dx.$$

(ii) If
$$p \ge 2$$
,
(4.3)
$$\int_{\Omega_{8r}} |Du - Dh|^p \, dx \le c \int_{\Omega_{8r}} |F|^p \, dx,$$
and for every $\epsilon \in (0, 1)$,

(4.4)
$$\int_{\Omega_{8r}} V|u-h|^s \, dx \le \epsilon \int_{\Omega_{8r}} V|u|^s \, dx + c(\epsilon) \int_{\Omega_{8r}} |F|^p \, dx.$$
(iii) If $1 , for every $\epsilon \in (0, 1)$,$

(4.5)
$$\int_{\Omega_{8r}} |Du - Dh|^p \, dx \le \epsilon \oint_{\Omega_{8r}} |Du|^p \, dx + c(\epsilon) \int_{\Omega_{8r}} |F|^p \, dx,$$

(4.6)
$$\int_{\Omega_{8r}} V|u-h|^s \, dx \le \epsilon \int_{\Omega_{8r}} |Du|^p + V|u|^s \, dx + c(\epsilon) \int_{\Omega_{8r}} |F|^p \, dx$$

Here, c > 0 depends on n, p, L, ν and s, and $c(\epsilon) > 0$ does on n, p, L, ν, s and ϵ .

Proof. We test the equations (1.1) and (4.1) with the test function $\varphi := u - h$ in order to discover

(4.7)
$$\int_{\Omega_{8r}} (\mathbf{a}(x, Du) - \mathbf{a}(x, Dh)) \cdot (Du - Dh) \, dx + \int_{\Omega_{8r}} V\left(|u|^{s-2}u - |h|^{s-2}h\right) \cdot (u - h) \, dx = \int_{\Omega_{8r}} |F|^{p-2} F \cdot (Du - Dh) \, dx.$$

We recall the monotonicity conditions (2.7) and

(4.8) $(|\xi_2|^{s-2}\xi_2 - |\xi_1|^{s-2}\xi_1) \cdot (\xi_2 - \xi_1) \ge c(s)(|\xi_1|^2 + |\xi_2|^2)^{\frac{s-2}{2}} |\xi_2 - \xi_1|^2, \quad \xi_1, \xi_2 \in \mathbb{R}.$ Then we see that the two terms on the left hand side of (4.7) are nonnegative. We first prove (i). By (4.7), (2.7) and (2.5), we infer

$$\begin{split} &\int_{\Omega_{8r}} |Dh|^p \, dx \leq c \int_{\Omega_{8r}} \mathbf{a}(x, Dh) \cdot Dh \, dx \\ &\leq c \int_{\Omega_{8r}} |\mathbf{a}(x, Du)| (|Du| + |Dh|) + |\mathbf{a}(x, Dh)| |Du| + |F|^{p-1} (|Du| + |Dh|) \, dx \\ &\leq c \int_{\Omega_{8r}} |Du|^p + |Du|^{p-1} |Dh| + |Dh|^{p-1} |Du| + |F|^{p-1} (|Du| + |Dh|) \, dx. \end{split}$$

Therefore, applying Young's inequality, we obtain (4.2).

We next prove (ii). Note that in this case the exponent s satisfies either s < 2 or $s \ge 2$. Using the monotonicity conditions (2.7) and (4.8) together with the fact that $2 \le p$, we derive from (4.7) that

$$\int_{\Omega_{8r}} |Du - Dh|^p \, dx + \int_{\Omega_{8r}} V(|u|^2 + |h|^2)^{\frac{s-2}{2}} |u - h|^2 \, dx \le c \int_{\Omega_{8r}} |F|^{p-1} |Du - Dh| \, dx,$$
which together Young's inequality implies

which together Young's inequality implies

$$\int_{\Omega_{8r}} |Du - Dh|^p \, dx + \int_{\Omega_{8r}} V(|u|^2 + |h|^2)^{\frac{s-2}{2}} |u - h|^2 \, dx \le c \int_{\Omega_{8r}} |F|^p \, dx.$$

Therefore, we obtain (4.3). On the other hand, by Young's inequality again, we have that for any $\epsilon \in (0, 1)$,

$$\int_{\Omega_{8r}} V|u-h|^s \, dx \le \epsilon \int_{\Omega_{8r}} V|u|^s + V|h|^s \, dx + c(\epsilon) \int_{\Omega_{8r}} |F|^p \, dx,$$

from which, by choosing ϵ so small we first have

$$\int_{\Omega_{8r}} V|h|^s \, dx \le c \int_{\Omega_{8r}} V|u|^s \, dx + c \int_{\Omega_{8r}} |F|^p \, dx.$$

Inserting this into the previous estimate yields (4.4).

Lastly we prove (iii). Applying the monotonicity conditions (2.7) and (4.8) and Young's inequality to (4.7) yields that for any $\kappa_1 \in (0, 1)$

$$\int_{\Omega_{8r}} \left(|Du|^2 + |Dh|^2 \right)^{\frac{p-2}{2}} |Du - Dh|^2 \, dx + \int_{\Omega_{8r}} V \left(|u|^2 + |h|^2 \right)^{\frac{s-2}{2}} |u - h|^2 \, dx$$

$$\leq c \int_{\Omega_{8r}} |F|^{p-1} |Du - Dh| \, dx \leq \kappa_1 \int_{\Omega_{8r}} |Du - Dh|^p \, dx + c(\kappa_1) \int_{\Omega_{8r}} |F|^p \, dx.$$

Since $1 , then, by Young's inequality again, we have that for any <math>\kappa_2 \in (0, 1)$,

$$\begin{aligned} |\xi_2 - \xi_1|^p &= |\xi_2 - \xi_1|^p \left(|\xi_2|^2 + |\xi_1|^2 \right)^{\frac{p(p-2)}{4}} \left(|\xi_2|^2 + |\xi_1|^2 \right)^{\frac{p(2-p)}{4}} \\ &\leq \kappa_2 \left(|\xi_2|^2 + |\xi_1|^2 \right)^{\frac{p}{2}} + c(\kappa_2) \left(|\xi_2|^2 + |\xi_1|^2 \right)^{\frac{p-2}{2}} |\xi_2 - \xi_1|^2, \end{aligned}$$

for all $\xi_1, \xi_2 \in \mathbb{R}^d$ with $d \in \mathbb{N}$. Therefore, using the above results we have

$$\int_{\Omega_{8r}} |Du - Dh|^p dx \le c\kappa_2 \int_{\Omega_{8r}} |Du|^p + |Dh|^p dx$$
$$+ \kappa_1 c(\kappa_2) \int_{\Omega_{8r}} |Du - Dh|^p dx + c(\kappa_1) c(\kappa_2) \int_{\Omega_{8r}} |F|^p dx,$$

which implies (4.5) by choosing κ_1 and κ_2 such that $c\kappa_2 = \frac{\epsilon}{2}$ and $\kappa_1 c(\kappa_2) = \frac{1}{2}$ and applying (4.2). In addition, we also have

$$\int_{\Omega_{8r}} |Du - Dh|^p dx + \int_{\Omega_{8r}} V|u - h|^s dx$$

$$\leq c\kappa_2 \int_{\Omega_{8r}} |Du|^p + |Dh|^p + V|u|^s + V|h|^s dx$$

$$+ \kappa_1 c(\kappa_2) \int_{\Omega_{8r}} |Du - Dh|^p dx + c(\kappa_1) c(\kappa_2) \int_{\Omega_{8r}} |F|^p dx.$$

Then we first choose κ_1 and κ_2 so small to get

(4.9)
$$\int_{\Omega_{8r}} V|h|^s \, dx \le c \int_{\Omega_{8r}} V|u|^s + |Du|^p + |F|^p \, dx,$$

where we used (4.2). Inserting this into the previous estimate, we have

$$\begin{split} \int_{\Omega_{8r}} |Du - Dh|^p \, dx + \int_{\Omega_{8r}} V|u - h|^s \, dx \\ &\leq c\kappa_2 \int_{\Omega_{8r}} |Du|^p + V|u|^s + |F|^p \, dx \\ &+ \kappa_1 c(\kappa_2) \int_{\Omega_{8r}} |Du - Dh|^p \, dx + c(\kappa_1) c(\kappa_2) \int_{\Omega_{8r}} |F|^p \, dx \end{split}$$

Therefore, again choosing κ_1 and κ_2 so small that $c\kappa_2 = \epsilon$ and $\kappa_1 c(\kappa_2) = \frac{1}{2}$, we obtain (4.6).

Remark 4.2. In the above lemma, by using (4.4) and (4.9), we also obtain

$$\int_{\Omega_{8r}} V|h|^s \, dx \le c \int_{\Omega_{8r}} |Du|^p \chi_{\{1$$

4.2. **Proof of Theorem 2.3.** We prove the theorem by using we the approach introduced by Mingione in [2, 40]. The proof goes in four steps.

Step 1. Setting.

Assume that $\mathbf{a} : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}^n$ is (δ, R) -vanishing and Ω is (δ, R) -Reifenberg flat for some R > 0, where $\delta \in (0, 1)$ will be chosen sufficiently small later in Step 3 (see Remark 4.4). Fix any $x_0 \in \overline{\Omega}$ and r > 0 satisfying $r \leq \frac{R}{2}$. Assume that $V \in \mathcal{B}_{\gamma}$, where the range of $\gamma > 1$ is given in (1)–(3) of Theorem 2.3.

We prove the estimates (2.10)-(2.12) at one time by denoting the function $\Phi(v; x)$ and constant γ_1 differently as follows:

Case 1. Estimation of (2.10): Let $\gamma \in (\tilde{\gamma}, \infty)$ with $\tilde{\gamma}$ given in (2.9). We fix any $q \in (1, \frac{\gamma^*(p-1)}{p})$ with γ^* in (2.1), and denote by

$$\Phi(v;x) := |Dv(x)|^p \text{ and } \gamma_1 := \begin{cases} \frac{\gamma^*(p-1)}{p} & \text{if } \gamma < n, \\ \max\{q+1, \left(\frac{3}{2}\right)^* \frac{p-1}{p}\} & \text{if } \gamma \ge n. \end{cases}$$

In Case 1, we note that, when $\gamma \geq n$, one can find the constant $\gamma_2 \in [\frac{3}{2}, n)$ such that $\gamma_1 = \frac{\gamma_2^*(p-1)}{p}$. Moreover, it is clear that $V \in \mathcal{B}_{\gamma_2}$ since $\gamma_2 < n \leq \gamma$. **Case 2. Estimation of** (2.11): Let $\gamma \in (1, \infty)$. We fix any $q \in (1, \gamma)$, and denote by

$$\Phi(v; x) := V(x)|v(x)|^s \quad \text{and} \quad \gamma_1 := \gamma.$$

Case 3. Estimation of (2.12): Let $\gamma \in [\frac{n}{p}, \infty)$. We fix any $q \in (1, \gamma)$, and denote by

$$\Phi(v;x) := |Dv(x)|^p + V|v(x)|^s \quad \text{and} \quad \gamma_1 := \gamma.$$

We note that $p \ge 2$ in Case 2 and 1 in Case 3.

With $\Phi(v; x)$ denoted in above and the weak solution u to (1.1), we define

$$E(\lambda,\rho) := \{ x \in \Omega_{\rho} : \Phi(u;x) > \lambda \}, \quad \lambda > 0$$

and

(4.10)
$$\lambda_0 := \oint_{\Omega_{2r}} \Phi(u; x) \, dx + \frac{1}{\delta_1} \oint_{\Omega_{2r}} |F|^p \, dx,$$

where $\delta_1 \in (0, 1)$ will be chosen sufficiently small later in Step 4.

Finally, fix any τ_1, τ_2 with $1 \leq \tau_1 < \tau_2 \leq 2$. Note that $\Omega_r \subset \Omega_{\tau_1 r} \subset \Omega_{\tau_2 r} \subset \Omega_{2r}$.

Step 2. Covering argument.

We consider $\lambda > 0$ large enough so that

(4.11)
$$\lambda > \alpha \lambda_0$$
, where $\alpha := \left(\frac{16}{7}\right)^n \left(\frac{80}{\tau_2 - \tau_1}\right)^n$.

Note that $\Omega_{\rho}(y) \subset \Omega_{2r}$ for any $y \in E(\lambda, \tau_1 r)$ and any $\rho \in (0, (\tau_2 - \tau_1) r]$. Then from the measure density condition (2.8) and the definition of λ_0 given in (4.10), we infer that

$$\begin{aligned} \int_{\Omega_{\rho}(y)} \Phi(u;x) \, dx &+ \frac{1}{\delta_1} \int_{\Omega_{\rho}(y)} |F|^p \, dx \le \frac{|\Omega_{2r}|}{|\Omega_{\rho}(y)|} \lambda_0 \le \left(\frac{16}{7}\right)^n \left(\frac{2r}{\rho}\right)^n \, \lambda_0 \\ &\le \alpha \, \lambda_0 < \lambda, \end{aligned}$$

provided that

$$\frac{(\tau_2 - \tau_1) r}{40} \le \rho \le (\tau_2 - \tau_1) r.$$

On the other hand, it follows from Lebesgue's differentiation theorem that for almost every $y \in E(\lambda, \tau_1 r)$,

$$\lim_{\rho \to 0} \left(\oint_{\Omega_{\rho}(y)} \Phi(u; x) \, dx + \frac{1}{\delta_1} \oint_{\Omega_{\rho}(y)} |F|^p \, dx \right) > \lambda.$$

Therefore the above inequalities and the continuity of the integral with respect to the measure of the domain yield that for almost every $y \in E(\lambda, \tau_1 r)$, there exists

$$\rho_y = \rho(y) \in \left(0, \frac{(\tau_2 - \tau_1)r}{40}\right)$$

such that

$$\int_{\Omega_{\rho_y}(y)} \Phi(u;x) \, dx + \frac{1}{\delta_1} \oint_{\Omega_{\rho_y}(y)} |F|^p \, dx = \lambda,$$

and for any $\rho \in (\rho_y, (\tau_2 - \tau_1)r]$ there holds

$$\int_{\Omega_{\rho}(y)} \Phi(u;x) \, dx + \frac{1}{\delta_1} \int_{\Omega_{\rho}(y)} |F|^p \, dx < \lambda$$

As a consequence, Vitali's covering theorem implies the following:

Lemma 4.3. Given $\lambda > \alpha \lambda_0$, there exists a disjoint family of $\{\Omega_{\rho_i}(y^i)\}_{i=1}^{\infty}$ with $y^i \in E(\lambda, \tau_1 r)$ and $\rho_i \in \left(0, \frac{(\tau_2 - \tau_1) r}{40}\right)$ such that

$$E(\lambda, \tau_1 r) \subset \bigcup_{i=1}^{\infty} \Omega_{5\rho_i}(y^i),$$

(4.12)
$$\int_{\Omega_{\rho_i}(y^i)} \Phi(u;x) \, dx + \frac{1}{\delta_1} \int_{\Omega_{\rho_i}(y^i)} |F|^p \, dx = \lambda$$

and for any $\rho \in (\rho_i, (\tau_2 - \tau_1) r]$,

(4.13)
$$\int_{\Omega_{\rho}(y^{i})} \Phi(u;x) \, dx + \frac{1}{\delta_{1}} \oint_{\Omega_{\rho}(y^{i})} |F|^{p} \, dx < \lambda.$$

Furthermore, according to Lemma 4.3 we infer

$$\begin{split} &|\Omega_{\rho_{i}}(y^{i})| = \frac{1}{\lambda} \left(\int_{\Omega_{\rho_{i}}(y^{i})} \Phi(u;x) \, dx + \frac{1}{\delta_{1}} \int_{\Omega_{\rho_{i}}(y^{i})} |F|^{p} \, dx \right) \\ &\leq \frac{1}{\lambda} \left(\int_{\Omega_{\rho_{i}}(y^{i}) \cap \{\Phi(u;x) > \frac{\lambda}{4}\}} \Phi(u;x) \, dx + \frac{1}{\delta_{1}} \int_{\Omega_{\rho_{i}}(y^{i}) \cap \{|F|^{p} > \frac{\delta_{1}\lambda}{4}\}} |F|^{p} \, dx + \frac{\lambda}{2} \left| \Omega_{\rho_{i}}(y^{i}) \right| \right) \\ &= \frac{1}{2} \left| \Omega_{\rho_{i}}(y^{i}) \right| + \frac{1}{\lambda} \left(\int_{\Omega_{\rho_{i}}(y^{i}) \cap \{\Phi(u;x) > \frac{\lambda}{4}\}} \Phi(u;x) \, dx + \frac{1}{\delta_{1}} \int_{\Omega_{\rho_{i}}(y^{i}) \cap \{|F|^{p} > \frac{\delta_{1}\lambda}{4}\}} |F|^{p} \, dx \right) \end{split}$$

and so (4.14)

$$\left|\Omega_{\rho_i}(y^i)\right| \leq \frac{2}{\lambda} \bigg(\int_{\Omega_{\rho_i}(y^i) \cap \{\Phi(u;x) > \frac{\lambda}{4}\}} \Phi(u;x) \, dx + \frac{1}{\delta_1} \int_{\Omega_{\rho_i}(y^i) \cap \{|F|^p > \frac{\delta_1\lambda}{4}\}} |F|^p \, dx \bigg).$$

Step 3. Comparison estimates.

We note from (4.13) in Lemma 4.3 that

$$\int_{\Omega_{40\rho_i}(y^i)} \Phi(u;x) \, dx + \frac{1}{\delta_1} \oint_{\Omega_{40\rho_i}(y^i)} |F|^p \, dx < \lambda.$$

Applying Lemma 4.1, we have that for any $\epsilon \in (0,1)$, there exists a small $\delta_1 = \delta_1(\epsilon, n, p, L, \nu, s) \in (0,1)$ such that

(4.15)
$$\int_{\Omega_{40\rho_i}(y^i)} \Phi(u-h_i;x) \, dx \le \epsilon \int_{\Omega_{40\rho_i}(y^i)} \Phi(u;x) \, dx + c(\epsilon) \int_{\Omega_{40\rho_i}(y^i)} |F|^p \, dx$$
$$\le \epsilon \lambda + c(\epsilon) \delta_1 \lambda \le 2\epsilon \lambda.$$

(In fact, when $p \ge 2$ in **Case 1**, $c(\epsilon)$ in (4.15) does not depend on ϵ .) Furthermore, recalling the definition of γ_1 in **Cases 1–3** and applying Theorems 3.6 and 3.7 with $\gamma = \gamma_1$, we have

(4.16)
$$\left(\oint_{\Omega_{5\rho_i}(y^i)} \Phi(h_i; x)^{\gamma_1} dx \right)^{\frac{1}{\gamma_1}} \leq c \oint_{\Omega_{40\rho_i}(y^i)} \Phi(h_i; x) dx \\ \leq c \oint_{\Omega_{40\rho_i}(y^i)} \Phi(u; x) + |F|^p dx \leq c\lambda,$$

where $h_i \in W^{1,p}(\Omega_{20\rho_i}(y^i))$ is the unique weak solution to

$$\begin{cases} -\operatorname{div} \mathbf{a}(x, Dh_i) + V|h_i|^{s-2}h_i &= 0 \quad \text{in} \quad \Omega_{40\rho_i}(y^i), \\ h_i &= u \quad \text{on} \quad \partial\Omega_{40\rho_i}(y^i). \end{cases}$$

Remark 4.4. At this stage, the constant δ is fixed as the one in Theorem 3.7 with $\gamma = \gamma_1$. Therefore, δ depends on n, p, L, ν, γ when $\gamma < n$ or n, p, L, ν, q when $\gamma \geq n$. Moreover, in Case 2, Theorem 3.7 is not used, but only Theorem 3.6 is. Hence the (δ, R) -vanishing smallness assumptions on **a** and Ω are not needed in this case.

Let $y \in \Omega_{5\rho_i}(y^i)$ such that $\Phi(u; y) > K\lambda$, where constant $K \ge 1$ will be chosen later. We then note that

$$\Phi(u;y) \le 2^{\max\{p,s\}-1} [\Phi(u-h_i;y) + \Phi(h_i;y)].$$

Here, we need to consider the two cases:

(i)
$$\Phi(h_i; y) \le \Phi(u - h_i; y)$$
, (ii) $\Phi(h_i; y) > \Phi(u - h_i; y)$.

For the case (i), it is clear that

$$\Phi(u;y) \le 2^{\max\{p,s\}} \Phi(u-h_i;y).$$

For the case (ii), we see that

$$K\lambda < \Phi(u; y) \le 2^{\max\{p, s\}} \Phi(h_i; y),$$

from which, it follows that

$$\Phi(u;y) \le 2^{\max\{p,s\}} \Phi(h_i;y) \left[\frac{2^{\max\{p,s\}} \Phi(h_i;y)}{K\lambda}\right]^{\gamma_1-1} = \frac{2^{\gamma_1 \max\{p,s\}}}{(K\lambda)^{\gamma_1-1}} \Phi(h_i;y)^{\gamma_1}.$$

For the both cases (i) and (ii), we finally obtain that

$$\Phi(u;y) \le 2^{\max\{p,s\}} \Phi(u-h_i;y) + \frac{2^{\gamma_1 \max\{p,s\}}}{(K\lambda)^{\gamma_1-1}} \Phi(h_i;y)^{\gamma_1}$$

for any $y \in \Omega_{5\rho_i}(y^i)$ such that $\Phi(u; y) > K\lambda$. Then we apply (4.15) and (4.16) to discover

$$\begin{split} &\int_{\Omega_{5\rho_i}(y^i)\cap E(K\lambda,\tau_2r)} \Phi(u;x) \, dx \\ &\leq c \int_{\Omega_{5\rho_i}(y^i)} \Phi(u-h_i;x) \, dx + \frac{c}{(K\lambda)^{\gamma_1-1}} \int_{\Omega_{5\rho_i}(y^i)} \Phi(h_i;x)^{\gamma_1} \, dx \\ &\leq c \left(\epsilon\lambda + \frac{\lambda^{\gamma_1}}{(K\lambda)^{\gamma_1-1}}\right) \left|\Omega_{5\rho_i}(y^i)\right| \\ &\leq c\lambda \left(\epsilon + K^{1-\gamma_1}\right) \left|\Omega_{\rho_i}(y^i)\right| = c\tilde{\epsilon}\lambda \left|\Omega_{\rho_i}(y^i)\right| \end{split}$$

for some constant $c = c(n, p, L, \nu, s, \gamma, b_{\gamma}) > 0$, where (4.17) $\tilde{\epsilon} := \epsilon + K^{1-\gamma_1}$.

Inserting (4.14) into the previous estimate, we conclude that

$$\int_{\Omega_{5\rho_i}(y^i)\cap E(K\lambda,\tau_2r)} \Phi(u;x) dx$$

$$\leq c\tilde{\epsilon} \left(\int_{\Omega_{\rho_i}(y^i)\cap\left\{\Phi(u;x)>\frac{\lambda}{4}\right\}} \Phi(u;x) dx + \frac{1}{\delta_1} \int_{\Omega_{\rho_i}(y^i)\cap\left\{|F|^p>\frac{\delta_1\lambda}{4}\right\}} |F|^p dx \right).$$

From Lemma 4.3, we note that $\Omega_{\rho_i}(y^i)$ is mutually disjoint and

$$E(K\lambda, \tau_1 r) \subset E(\lambda, \tau_1 r) \subset \bigcup_{i=1}^{\infty} \Omega_{5\rho_i}(y^i) \subset \Omega_{\tau_2 r},$$

since $K \geq 1$. Then we obtain that

(4.18)
$$\int_{E(K\lambda,\tau_1r)} \Phi(u;x) \, dx \leq \sum_{i=1}^{\infty} \int_{\Omega_{5\rho_i}(y^i) \cap E(K\lambda,\tau_1r)} \Phi(u;x) \, dx$$
$$\leq c\tilde{\epsilon} \left(\int_{\Omega_{\tau_2r} \cap \left\{ \Phi(u;x) > \frac{\lambda}{4} \right\}} \Phi(u;x) \, dx + \frac{1}{\delta_1} \int_{\Omega_{\tau_2r} \cap \left\{ |F|^p > \frac{\delta_1\lambda}{4} \right\}} |F|^p \, dx \right)$$

for some constant $c = c(n, p, \gamma, L, \nu, b_{\gamma}) > 0$.

Step 4. Proof of (2.10)-(2.12).

We are now ready to conclude the proof via Fubini's theorem with a truncation argument. For k > 0, let us define

$$\Phi_k(u;x) := \min\left\{\Phi(u;x),k\right\},\,$$

and consider the upper level set with respect to Φ_k as

$$E_k(\tilde{\lambda}, \rho) := \left\{ y \in \Omega_\rho : \Phi_k(u; y) > \tilde{\lambda} \right\} \text{ for } \tilde{\lambda}, \rho > 0.$$

Then since $E_k(\tilde{\lambda}, \rho) = \emptyset$ when $k \leq \tilde{\lambda}$ and $E_k(\tilde{\lambda}, \rho) = E(\tilde{\lambda}, \rho)$ when $k > \tilde{\lambda}$, it follows from (4.18) that

$$\int_{E_k(K\lambda,\tau_1r)} \Phi(u;x) \, dx \le c \tilde{\epsilon} \bigg(\int_{E_k\left(\frac{\lambda}{4},\tau_2r\right)} \Phi(u;x) \, dx + \frac{1}{\delta_1} \int_{\Omega_{\tau_2r} \cap \left\{ |F|^p > \frac{\delta_1\lambda}{4} \right\}} |F|^p \, dx \bigg).$$

By multiplying both sides by λ^{q-2} and integrating with respect to λ over $(\alpha\lambda_0, \infty)$, we then have that

$$(4.19)$$

$$I_{0} := \int_{\alpha\lambda_{0}}^{\infty} \lambda^{q-2} \int_{E_{k}(K\lambda,\tau_{1}r)} \Phi(u;x) \, dx d\lambda$$

$$\leq c\tilde{\epsilon} \left(\int_{\alpha\lambda_{0}}^{\infty} \lambda^{q-2} \int_{E_{k}\left(\frac{\lambda}{4},\tau_{2}r\right)} \Phi(u;x) \, dx d\lambda + \int_{\alpha\lambda_{0}}^{\infty} \lambda^{q-2} \int_{\Omega_{\tau_{2}r} \cap \left\{\frac{|F|^{p}}{\delta_{1}} > \frac{\lambda}{4}\right\}} \frac{|F|^{p}}{\delta_{1}} \, dx d\lambda \right)$$

$$=: c\tilde{\epsilon}(I_{1} + I_{2}).$$

Here, by virtue of Fubini's theorem, we derive that

$$I_{0} = \int_{E_{k}(K\alpha\lambda_{0},\tau_{1}r)} \Phi(u;x) \left(\int_{\alpha\lambda_{0}}^{\Phi_{k}(u;x)/K} \lambda^{q-2} d\lambda \right) dx$$
$$= \frac{1}{q-1} \left\{ \int_{E_{k}(K\alpha\lambda_{0},\tau_{1}r)} \Phi(u;x) \left[\frac{\Phi_{k}(u;x)}{K} \right]^{q-1} dx - (\alpha\lambda_{0})^{q-1} \int_{E_{k}(K\alpha\lambda_{0},\tau_{1}r)} \Phi(u;x) dx \right\},$$

and

$$I_{1} = \int_{E_{k}\left(\frac{\alpha\lambda_{0}}{4},\tau_{2}r\right)} \Phi(u;x) \left(\int_{\alpha\lambda_{0}}^{4\Phi_{k}(u;x)} \lambda^{q-2} d\lambda\right) dx$$

$$\leq \frac{1}{q-1} \int_{E_{k}\left(\frac{\alpha\lambda_{0}}{4},\tau_{2}r\right)} \Phi(u;x) \left[4\Phi_{k}(u;x)\right]^{q-1} dx$$

$$\leq \frac{4^{q-1}}{q-1} \int_{\Omega_{\tau_{2}r}} \Phi(u;x) \Phi_{k}(u;x)^{q-1} dx.$$

Similarly, we obtain that

$$I_{2} = \int_{\Omega_{\tau_{2}r} \cap \left\{\frac{|F|^{p}}{\delta_{1}} > \frac{\alpha\lambda_{0}}{4}\right\}} \frac{|F|^{p}}{\delta_{1}} \int_{\alpha\lambda_{0}}^{4|F|^{p}/\delta_{1}} \lambda^{q-2} d\lambda dx$$
$$\leq \frac{1}{q-1} \int_{\Omega_{\tau_{2}r} \cap \left\{\frac{|F|^{p}}{\delta_{1}} > \frac{\alpha\lambda_{0}}{4}\right\}} \frac{|F|^{p}}{\delta_{1}} \left[\frac{4|F|^{p}}{\delta_{1}}\right]^{q-1} dx$$
$$\leq \frac{4^{q-1}}{q-1} \int_{\Omega_{\tau_{2}r}} \left[\frac{|F|^{p}}{\delta_{1}}\right]^{q} dx.$$

Therefore we insert the previous estimates for I_0 , I_1 , I_2 into (4.19) to discover

$$\begin{split} &\int_{E_k(K\alpha\lambda_0,\tau_1r)} \Phi(u;x)\Phi_k(u;x)^{q-1} dx \\ &\leq (K\alpha\lambda_0)^{q-1} \int_{\Omega_{\tau_1r}} \Phi(u;x) dx \\ &+ c\tilde{\epsilon}K^{q-1} \bigg(\int_{\Omega_{\tau_2r}} \Phi(u;x)\Phi_k(u;x)^{q-1} dx + \int_{\Omega_{\tau_2r}} \bigg[\frac{|F|^p}{\delta_1} \bigg]^q dx \bigg). \end{split}$$

We also note that

$$\int_{\Omega_{\tau_1 r} \setminus E_k(K\alpha\lambda_0, \tau_1 r)} \Phi(u; x) \Phi_k(u; x)^{q-1} \, dx \le (K\alpha\lambda_0)^{q-1} \int_{\Omega_{\tau_1 r}} \Phi(u; x) \, dx.$$

In turn, by the last two estimates we obtain

$$\begin{split} &\int_{\Omega_{\tau_1 r}} \Phi(u; x) \Phi_k(u; x)^{q-1} \, dx \\ &\leq (K \alpha \lambda_0)^{q-1} \int_{\Omega_{\tau_1 r}} \Phi(u; x) \, dx \\ &+ c_2 \tilde{\epsilon} K^{q-1} \bigg(\int_{\Omega_{\tau_2 r}} \Phi(u; x) \Phi_k(u; x)^{q-1} \, dx + \int_{\Omega_{\tau_2 r}} \left[\frac{|F|^p}{\delta_1} \right]^q \, dx \bigg) \end{split}$$

for some $c_2 = c_2(n, p, L, \nu, s, \gamma, b_{\gamma}, q) > 0$. At this stage, we recall the definition of $\tilde{\epsilon}$ given in (4.17), and then take large K > 1 and small $\epsilon \in (0, 1)$ depending on $n, p, L, \nu, s, \gamma, b_{\gamma}, q$ such that

$$K \ge (4c_2)^{\frac{1}{\gamma_1 - q}}$$
 and $\epsilon \le \frac{1}{4c_2 K^{q-1}}$,

hence $\delta_1 = \delta_1(n, p, L, \nu, \gamma, b_{\gamma}, q) \in (0, 1)$ is finally determined. Recalling the definition of α in (4.11) we consequently obtain

$$\int_{\Omega_{\tau_1 r}} \Phi(u; x) \Phi_k(u; x)^{q-1} dx$$

$$\leq \frac{1}{2} \int_{\Omega_{\tau_2 r}} \Phi(u; x) \Phi_k(u; x)^{q-1} dx + \frac{c\lambda_0^{q-1}}{(\tau_2 - \tau_1)^n} \int_{\Omega_{2r}} \Phi(u; x) dx + c \int_{\Omega_{2r}} |F|^{pq} dx.$$

Then we apply Lemma 3.2 to discover

$$\int_{\Omega_r} \Phi(u;x) \Phi_k(u;x)^{q-1} \, dx \le c\lambda_0^{q-1} \int_{\Omega_{2r}} \Phi(u;x) \, dx + c \int_{\Omega_{2r}} |F|^{pq} \, dx$$

for any k > 0. Finally, by virtue of Lebesgue's monotone convergence theorem, Hölder's inequality and Young's inequality together with the definition of λ_0 in (4.10), we derive that

$$\begin{aligned} \oint_{\Omega_r} \Phi(u;x)^q \, dx &= \lim_{k \to \infty} \oint_{\Omega_r} \Phi(u;x) \Phi_k(u;x)^{q-1} \, dx \\ &\leq c \lambda_0^{q-1} \oint_{\Omega_{2r}} \Phi(u;x) \, dx + c \oint_{\Omega_{2r}} |F|^{pq} \, dx \\ &\leq c \left(\oint_{\Omega_{2r}} \Phi(u;x) \, dx \right)^q + c \oint_{\Omega_{2r}} |F|^{pq} \, dx. \end{aligned}$$

This implies the desired estimates (2.10)–(2.12), by recalling the definition of $\Phi(u; x)$ in **Cases 1–3**.

References

- 1. Acerbi, E. and Mingione, G., Gradient estimates for the p(x)-Laplacean system, J. Reine Angew. Math. **584** (2005), 117–148.
- Acerbi, E. and Mingione, G., Gradient estimates for a class of parabolic systems, Duke Math. J., 136 (2007), 285–320.

- Auscher, P. and Ben Ali, B. Maximal inequalities and Riesz transform estimates on L^p spaces for Schrödinger operators with nonnegative potentials, Ann. Inst. Fourier (Grenoble) 57 (2007), no. 6, 1975–2013.
- Bergé, L., Soliton stability versus collapse, Phys. Rev. E (3) 62 (2000), no. 3, part A, R3071– R3074.
- Bögelein, V., Global Calderón-Zygmund theory for nonlinear parabolic systems, Calc. Var. Partial Differential Equations 51 (2014), no. 3-4, 555–596.
- Brézis, H. and Strauss, W. A. Semi-linear second-order elliptic equations in L¹, J. Math. Soc. Japan 25 (1973), 565–590.
- Bramanti, M., Brandolini, L., Harboure, E. and Viviani, B., Global W^{2,p} estimates for nondivergence elliptic operators with potentials satisfying a reverse Hölder condition, Ann. Mat. Pura Appl. (4) **191** (2012), no. 2, 339–362.
- Byun, S. and Oh, J., Global gradient estimates for non-uniformly elliptic equations, Calc. Var. Partial Differential Equations 56 (2017), no. 2, Paper No. 46, 36 pp.
- Byun, S., Ok, J. and Ryu, S., Global gradient estimates for general nonlinear parabolic equations in nonsmooth domains, J. Differential Equations 254 (2013), no. 11, 4290–4326.
- 10. Byun, S., Ok, J. and Ryu, S., Global gradient estimates for elliptic equations of p(x)-Laplacian type with BMO nonlinearity, J. Reine Angew. Math. **715** (2016), 1–38.
- Byun, S., Palagachev, D. K. and Shin, P., Global Sobolev regularity for general elliptic equations of p-Laplacian type, Calc. Var. Partial Differential Equations 57 (2018), no. 5, Paper No. 135, 19 pp.
- Byun, S. and Wang, L., *Elliptic equations with BMO coefficients in Reifenberg domains*, Comm. Pure Appl. Math. 57 (2004), no. 10, 1283–1310.
- Caffarelli L. A. and Peral I., On W^{1,p} estimates for elliptic equations in divergence form, Comm. Pure Appl. Math. 51 (1998), no. 1, 1–21.
- Calderon A. P. and Zygmund A., On the existence of certain singular integrals Acta Math. 88 (1952), 85–139.
- Chiarenza, F., Fabes, E. and Garofalo, N., Harnack's inequality for Schrödinger operators and the continuity of solutions, Proc. Amer. Math. Soc. 98 (1986), no. 3, 415–425.
- Colombo, M. and Mingione, G., Calderón-Zygmund estimates and non-uniformly elliptic operators, J. Funct. Anal. 270 (2016), no. 4, 1416–1478.
- DiBenedetto, E. and Manfredi, J., On the higher integrability of the gradient of weak solutions of certain degenerate elliptic systems, Amer. J. Math. 115 (1993), no. 5, 1107–1134.
- Di Fazio, G., Hölder-continuity of solutions for some Schrödinger equations, Rend. Sem. Mat. Univ. Padova 79 (1988), 173–183.
- Di Fazio, G. L^p estimates for divergence form elliptic equations with discontinuous coefficients, Boll. Un. Mat. Ital. A (7) 10 (1996), no. 2, 409–420.
- Dinh, V. D., Energy scattering for a class of the defocusing inhomogeneous nonlinear Schrödinger equation, J. Evol. Equ. 19 (2019), no. 2, 411–434.
- Fefferman, C., The uncertainty principle, Bull. Amer. Math. Soc. (N.S.) 9 (1983), no. 2, 129–206.
- Fibich, G. and Wang, X., Stability of solitary waves for nonlinear Schrödinger equations with inhomogeneous nonlinearities, Phys. D 175 (2003), no. 1-2, 96–108.
- Gehring, F.W., The L^p-integrability of the partial derivatives of a quasiconformal mapping, Acta Math. 130 (1973), 265–277.
- Gill, T. S., Optical guiding of laser beam in nonuniform plasma, Pramana J. Phys. 55 (2000), no. 5–6, 835–842.
- Giusti, E., Direct methods in the calculus of variations, World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- Grafakos, L., Modern Fourier analysis, Second edition. Graduate Texts in Mathematics, 250. Springer, New York, 2009.
- Han, Q. and Lin, F., *Elliptic Partial Differential Equations*, Courant Lecture Notes in Mathematics, 1, New York University, Courant Institute of Mathematical Sciences, New York, American Mathematical Society, Providence, RI, 1997.
- Iwaniec, T., Projections onto gradient fields and L^p-estimates for degenerated elliptic operators, Studia Math. 75 (1983), no. 3, 293–312.
- Kato, T., Schrödinger operators with singular potentials, Israel J. Math. 13 (1972), 135–148 (1973).

- Kato, T., L^p-theory of Schrödinger operators with a singular potential. Aspects of positivity in functional analysis (Tübingen, 1985), 63–78, North-Holland Math. Stud., 122, Notas Mat., 108, North-Holland, Amsterdam, 1986.
- Kim, J., Lee, Y. and Seo, I., On well-posedness for the inhomogeneous nonlinear Schrödinger equation in the critical case, J. Differential Equations 280 (2021), 179–202.
- Kinnunen, J. and Zhou, S., A local estimate for nonlinear equations with discontinuous coefficients, Comm. Partial Differential Equations 24 (1999), no. 11-12, 2043–2068.
- Kurata, K., Continuity and Harnack's inequality for solutions of elliptic partial differential equations of second order, Indiana Univ. Math. J. 43 (1994), no. 2, 411–440.
- Lee, M. and Ok, J. Nonlinear Calderón-Zygmund theory involving dual data, Revista Matemática Iberoamericana, 35 (2019), no. 4, 1053–1078.
- Lee, M. and Ok, J. Interior and boundary W^{1,q}-estimates for quasi-linear elliptic equations of Schrödinger type, J. Differential Equations 269 (2020), no. 5, 4406–4439.
- Liu, C. S. and Tripathi, V. K., Laser guiding in an axially nonuniform plasma channel, Phys. Plasmas 1 (1994), no. 9, 3100–3103.
- Muckenhoupt, B., Weighted norm inequalities for the Hardy maximal function, Trans. Amer. Math. Soc. 165 (1972), 207–226.
- 38. Merle, F., Nonexistence of minimal blow-up solutions of equations $iu_t = \Delta u k(x) |u|^{4/N} u$ in \mathbb{R}^N , Ann. Inst. H. Poincaré Phys. Théor. 64 (1996), no. 1, 33–85.
- Meyers, N. G. An L^p-estimate for the gradient of solutions of second order elliptic divergence equations Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) 17 (1963), 189–206.
- Mingione, G., The Calderón-Zygmund theory for elliptic problems with measure data, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 6 (2007), no. 2, 195–261.
- Palagachev, D.K. and Softova, L., The Calderón-Zygmund property for quasilinear divergence form equations over Reifenberg flat domains, Nonlinear Anal. 74 (2011) no. 5, 1721–1730.
- Pan, G. and Tang, L., Solvability for Schrödinger equations with discontinuous coefficients, J. Funct. Anal. 270 (2016) no. 1, 88–133.
- Reifenberg, E.R., Solution of the Plateau Problem for m-dimensional surfaces of varying topological type Acta Math. 104 (1960) 1–92.
- Raphaöl, P. and Szeftel, J., Existence and uniqueness of minimal blow-up solutions to an inhomogeneous mass critical NLS, J. Amer. Math. Soc. 24 (2011), no. 2, 471–546.
- Semenov, Y. A., Schrödinger operators with L^p_{loc}-potentials, Comm. Math. Phys. 53 (1977), no. 3, 277–284.
- Shen, Z., On the Neumann problem for Schrödinger operators in Lipschitz domains, Indiana Univ. Math. J. 43 (1994), no. 1, 143–176.
- Shen, Z., L^p estimates for Schrödinger operators with certain potentials, Ann. Inst. Fourier 45 (1995), no. 2, 513–546.
- Showalter, R. E., Monotone operators in Banach Space and Nonlinear Partial Differential Equations, Mathematical Surveys and Monographs, Vol. 49, American Mathematical Society, Providence, RI, 1997.
- Toro, T., Doubling and flatness: geometry of measures, Notices Amer. Math. Soc. 44 (1997), no. 9, 1087–1094.
- Towers, I. and Malomed, B. A., Stable (2+1)-dimensional solitons in a layered medium with sign-alternating Kerr nonlinearity, J. Opt. Soc. Amer. B Opt. Phys. 19 (2002), no. 3, 537–543.
- Tran, M., Nguyen, T. and Nguyen, G., Lorentz gradient estimates for a class of elliptic p-Laplacian equations with a Schrödinger term, J. Math. Anal. Appl. 496 (2021), no. 1, Paper No. 124806, 18 pp.
- Zhong, J. The Sobolev estimates for some Schrödinger type operators, Math. Sci. Res. Hot-Line 3 (1999), no. 8, 1–48.

MIKYOUNG LEE, DEPARTMENT OF MATHEMATICS, PUSAN NATIONAL UNIVERSITY, BUSAN 46241, REPUBLIC OF KOREA

Email address: mikyounglee@pusan.ac.kr

JIHOON OK, DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 04107, REPUBLIC OF KOREA

Email address: jihoonok@sogang.ac.kr