

Stoichiometric and dynamical autocatalysis for diluted chemical reaction networks

J eremie Unterberger^a, Philippe Nghe^b

^aInstitut Elie Cartan,¹ Universit e de Lorraine,
B.P. 239, F – 54506 Vandœuvre-l es-Nancy Cedex, France
jeremie.unterberger@univ-lorraine.fr

^bUMR CNRS-ESPCI Chimie Biologie Innovation 8231,
ESPCI Paris, Universit e Paris Sciences Lettres,
10 rue Vauquelin, 75005 Paris, France
philippe.nghe@espci.psl.eu

Autocatalysis underlies the ability of chemical and biochemical systems to replicate. Recently, Blokhuis et al. [8] gave a *stoichiometric* definition of autocatalysis for reaction networks, stating the existence of a combination of reactions such that the balance for all autocatalytic species is strictly positive, and investigated minimal autocatalytic networks, called *autocatalytic cores*. By contrast, spontaneous autocatalysis – namely, exponential amplification of all species internal to a reaction network, starting from a *diluted regime*, i.e. low concentrations – is a *dynamical* property.

We introduce here a topological condition (Top) for autocatalysis, namely: restricting the reaction network description to highly diluted species, we assume existence of a strongly connected component possessing at least one reaction with multiple products (including multiple copies of a single species). We find this condition to be necessary and sufficient for stoichiometric autocatalysis. When degradation reactions have small enough rates, the topological condition further ensures *dynamical autocatalysis*, characterized by a strictly positive Lyapunov exponent giving the instantaneous exponential growth rate of the system.

The proof is generally based on the study of auxiliary Markov chains. We provide as examples general autocatalytic cores of Type I and Type III in the typology of [8]. In a companion article [30], Lyapunov exponents and the behavior in the growth regime are studied quantitatively beyond the present diluted regime .

Keywords: autocatalysis, chemical reaction networks, origin of life, Lyapunov exponent, growth rate, continuous-time Markov chains

MSC Classification (2020): 34D08, 60J20, 80A32, 92C42, 92C45, 92E20

¹Laboratoire associ e au CNRS UMR 7502.

Contents

1	Introduction	2
1.1	Our main result in a nutshell, and contexts for applications	2
1.2	Linearized dynamics of reaction networks	4
1.3	Diluted regime and statement of condition (Top)	9
2	A motivating example: the simplest autocatalytic core	13
3	(Top) characterizes stoichiometric autocatalysis in diluted networks	15
4	(Top) implies dynamical autocatalysis for dilute networks	24
5	Perspectives	28
6	Supplementary information	30
6.1	Presentation of type I cycles	30
6.2	Presentation of type III cycles	31
6.3	Autocatalysis from the stoichiometric matrix	32
6.4	Linearized time-evolution generator for reaction networks	33
6.5	Generalized Markov generators	34

1 Introduction

1.1 Our main result in a nutshell, and contexts for applications

The chemical mechanism that epitomizes the ability of living systems to reproduce themselves is autocatalysis, namely, catalysis brought about by one of the products of the reactions. Autocatalysis must have been present from the early stages of the origin of life, from primitive forms of metabolism [2], to autocatalytic sets based on the first catalytic biopolymers [21] and the emergence of sustained template-based replication of nucleic acids [11]. Diverse artificial autocatalytic systems have been implemented in the laboratory [18]), and remnants of ancestral autocatalytic networks may be found in extant metabolic networks [23]. These examples reveal the diversity of autocatalytic mechanisms and chemistries. However, the stoichiometry of autocatalytic has been characterized only recently [8] and we still lack a systematic understanding of dynamical conditions for autocatalysis [6] which limits our ability to conceive plausible prebiotic scenarios [20].

To fill this gap, it is necessary to investigate how autocatalysis may emerge in complex mixtures. This would help us understand the appearance of self-sustaining reactions in messy prebiotic mixtures [10], and interpret experiments that search for such reactions [7]. Identifying autocatalytic systems is also critical to explain the appearance of Darwinian

evolution, from complex mixtures [10], to autocatalytic sets [19] and ultimately template-based replication [25], a path which comprises multiple transitions and can be studied experimentally in RNA reaction networks [3],[4].

The focus here is on *spontaneous* autocatalysis in chemical reaction networks, namely, exponential amplification of a set of species with low initial concentrations. This requires that certain other species, from which the network feeds, are provided in sufficiently large quantities in the environment. These resource species, sometimes called the 'food-set', may be constantly supplied from a large reservoir or external fluxes, or may be the products of reactions that already self-sustain in the milieu [9].

Our main result, Theorem 3.1, gives a general condition, denoted (Top), for spontaneous autocatalysis to be possible in a *stoichiometric*, respectively *dynamical* sense, understood as the existence of, respectively: combinations of reactions that lead to an increase of every autocatalytic species, and instantaneous growth of the dynamical system associated with the reaction network. Our result holds provided that the reaction set satisfies the formal conditions stated in [8]: (i) *autonomy*: reactions should possess at least one reactant and one product; (ii) *non-ambiguity*: a species cannot be both a reactant and a product of the same reaction. Point (i) ensures that concentrations do not increase merely due to reactions that only consume species from the environment. Said differently, it ensures that any concentration increase depends on the presence of another autocatalytic species, as required by the definition of autocatalysis [8]. Point (ii) imposes a formal choice of coarse-graining in the description of the reaction network. This choice ensures that catalytic steps can be distinguished at the level of the stoichiometric matrix as the catalysts then appear in the stoichiometry (as shown in [8]). Note that such a choice implies no restriction of generality, as it is always possible to introduce additional reaction intermediates in the description so that (ii) is respected [8].

Given the above conventions, verifying autocatalysis consists in isolating subsets of reactions that obey the topological criteria below (Fig. 1.1):

1. Retain only species that are initially absent or rare and discard from the description those that are abundant (the environment).
2. Dismiss reactions that have more than one reactant among the absent or rare species.
3. In the resulting network, identify strongly connected components which possess at least one reaction with multiple products within the component, including the case of multiple copies of a single species.

Strongly connected components are defined as subgraphs in which any pair of vertices (species) are connected by a chain of reactions. Successful verification of the steps above implies *stoichiometric autocatalysis*, independently of the reaction rates. It further implies

dynamical autocatalysis for sufficiently small degradation rates, as characterized by an exponential increase of every species in the component assuming initially low concentrations, at least in the early phase of the dynamics.

Here is a brief outline of the article. In the remainder of this section 1, we introduce the formal framework for reaction networks. A simple motivating example is presented in section 2. Sections 3 and 4 are the core of the article. We state our main result (Theorem 3.1) in section 3, and discuss equivalence of stoichiometric autocatalysis with the topological property (Top). In section 4, we prove that this property implies dynamical autocatalysis in the diluted regime when degradation reactions have small enough rates. We present perspectives for future work in section 5. Finally, section 6 provides supplementary information for the main text: a presentation of type I and type III cycles, and mathematical concepts and results used in the article, based on general Markov theory.

1.2 Linearized dynamics of reaction networks

We now introduce the framework of the present article, which is a mathematical and physical elaboration on the recent theoretical work [8] by Blokhuis, Lacoste and Nghe on autocatalysis in chemical reaction networks, the main conclusions of which we first recapitulate.

The general setting is that of open reaction networks, see e.g. [15] and references within. Chemical species fall into two categories: *dynamical* (or *non-chemostatted*) species, whose concentrations vary over time according to kinetic (or stochastic, if present in small number) equations, as opposed to *chemostatted* species, whose concentrations are fixed (or large w.r. to dynamical species, so that their concentrations may be considered as almost constant). Chemostatted species influence rates, but are not included into the stoichiometric matrix (see below), therefore they need not even be specified when dealing with stoichiometry alone. In [8], only *autonomous* networks are considered, i.e. every reaction – save *degradation reactions* – is supposed to have at least one (dynamical) reactant and at least one (dynamical) product. Degradation reactions $A \rightarrow \emptyset$ are natural in a biological setting; they play a major rôle in the story, but are not taken explicitly into account in the network.

The authors of [8] further insist on the necessity of writing reactions in an *unambiguous* form, i.e. in such a way that no chemical species can be both a reactant and a product of a reaction. For instance, this avoids reactions to be written as $A + E \rightleftharpoons B + E$ where the catalyst E appears on both sides, thus cancels from the total stoichiometric balance. Instead, the reaction should be written in two steps $A + E \rightleftharpoons EA \rightleftharpoons E + B$ which formally ensures that E appears in the stoichiometric balance and ultimately makes it





Figure 1: Species c and j (gray squares) are initially abundant in the environment, thus can be safely ignored. The reaction $g + d \rightarrow h + f$ (dashed) has multiple reactants that are initially rare or absent, thus has a negligible rate compared to others and is discarded from the description. In the remaining graph, the set $\{a, b, d, e, g, i\}$ forms a strongly connected component (SCC), as there exists a directed path between any two of its members. Species h and f (dashed gray circles) are not part of the SCC. The SCC comprises a reaction ($e \rightarrow a + d$) with multiple products. Thus, the SCC is stoichiometrically autocatalytic (note that it is actually a Type III autocatalytic core according to [8], see Supplementary Information 6.2). Furthermore, it is dynamically autocatalytic provided degradation rates of the species of the SCC are sufficiently small.

possible to recognize the catalytic cycle associated with enzyme E in the structure of the stoichiometric matrix (see [8] for details).

The authors then introduce a *stoichiometric criterion for autocatalysis* that depends only on the stoichiometric matrix \mathbb{S} , a matrix with columns indexed by reactions (other than degradation reactions) $(\mathcal{R}_1, \dots, \mathcal{R}_N)$, and rows indexed by the set $\mathcal{S} = \{A_1, \dots, A_{|\mathcal{S}|}\}$ of dynamical chemical species. Recall that, by definition, each column of \mathbb{S} corresponds to the stoichiometry of a given reaction



that is, $\mathbb{S}_{j,\mathcal{R}} = -\sum_{\ell=1}^n s_\ell \delta_{i_\ell,j} + \sum_{\ell'=1}^{n'} s'_{\ell'} \delta_{i'_{\ell'},j}$ (note that the coefficients of \mathbb{S} depend on the choice of an orientation for every reaction). Namely, they require that there should exist

- (i) a choice of orientations for reactions, and
- (ii) a positive reaction vector $c \in (\mathbb{R}_+)^N$ such that $\mathbb{S}c > 0$.

This means that the reaction obtained by taking the linear combination $\sum_{\mathcal{R}} c_{\mathcal{R}} \mathcal{R}$ strictly increases the number of molecules of all species in \mathcal{S} ; in other terms, the chemical balance $(\mathbb{S}c)_i$ for species i is > 0 . The main rationale for this condition is Gordan's theorem [1], which states that (ii) holds if and only if there is no mass-like conservation law, i.e. there exists no linear combination $\sum n_i [A_i]$ with positive coefficients $\mathbf{n} = (n_i)_{i \in \mathcal{S}} > 0$ such that $\mathbf{n} \cdot \mathbb{S} = 0$, i.e. preserved under all reactions.

The authors go on to give a classification of all autocatalytic *cores*, that is of all *minimal autonomous sub-networks* satisfying the above criteria, into 5 types I-V. (Types I and III are presented in Suppl. Info.)

Among foremost questions raised by this new classification, let us single out the two following:

- (A) *Are stoichiometrically autocatalytic networks able to replicate? Conversely, are chemical networks capable of replication stoichiometrically autocatalytic?*
- (B) (If the answer to (A) is: yes, and assuming some natural form for the rates, in particular, for mass-action rates.) *Under which conditions over the concentrations and the rates does an autocatalytic network indeed replicate? If it does, can one estimate its replication rate?*

To be specific, *kinetic rates will always be assumed to be mass-action rates*. Then this work presents an essentially complete answer to question (A) in a specific regime which we

call *diluted regime*, where all concentrations of dynamical species are *low*, and assuming that there are *no degradation reactions*, or that these have sufficiently small rates. The companion article [30], on the other hand, presents a detailed case study for question (B) for a broad class of autocatalytic cores in a large part of the growth regime, well beyond the diluted regime, and in presence of degradation reactions; it rests on the notations and concepts introduced here, which are therefore presented in great generality.

Partial answers to questions (A) and (B) are already available in [8]; they are based on self-consistent equations for survival probability, and are therefore rather given in the framework of stochastic networks, assuming only a few molecules are initially present. Generally speaking, survival criteria are given in a form akin to that given by King [22]. In [30], it is proved that this regime has connections to the diluted regime in the kinetic framework studied here, in the case when the Lyapunov exponent is zero. On the other hand, our formalism makes it possible to make predictions for the Lyapunov exponent far from the latter regime, as developed in the companion article [30].

Despite the fact that kinetic equations are not linear in general, our work is largely based on the study of the time-evolution of linear evolution models of the type

$$\frac{d[A_i]}{dt} = \sum_j M_{ij}[A_j] \quad (2)$$

with negative diagonal coefficients $M_{ii} \leq 0$ and positive off-diagonal coefficients $M_{ij} \geq 0, i \neq j$, which are found in different contexts in the literature. Note that these equations are formally similar to linear mutation-selection models, where off-diagonal coefficients are interpreted as mutation rates, and selection rates related to diagonal coefficients; see e.g. [11], [24]. Since we are mainly inspired by Markov techniques, we speak here of M as *generalized adjoint Markov generator*, see Supp. Info. Indeed, when the sum of coefficients on any column is zero, the total concentration $\sum_{i=1}^{|S|} [A_i]$ is a constant. Normalizing it to one, (2) yields a master equation, namely, the time-evolution of a probability measure. On the other hand, the case $|M_{j,j}| > \sum_{i \neq j} M_{ij}$ yields the time-evolution of a sub-Markov process, i.e. a Markov process with killing rates $a_j = |M_{j,j}| - \sum_{i \neq j} M_{ij}$. The case when some a_j are negative – indicating ‘source’ terms – is not standard in probability theory, but remains mathematically valid. Indeed, whatever the sign of a_j , the *Feynman-Kac formula* yields the solution to (2) in terms of a *sum over paths with transition rates $w_{i \rightarrow j}$ proportional to M_{ji}* (mind the index transposition due to the fact that M is a backward generator). Thinking in terms of kinetic networks (and in spite of the fact that these are assumed to be written as autonomous systems), a *positive killing rate a_j* is associated to a *degradation reaction $A_j \rightarrow \emptyset$* , whereas a *negative killing rate* is associated to an *inverse creation reaction $\emptyset \rightarrow A_j$* . (Since chemostatted species are left out of the equations, the latter, seemingly creation-ex-nihilo, reaction should be thought of really as $A' \rightarrow A_j + A''$, where A', A'' are chemostatted species).

Markov generators come out by *linearizing* the kinetic equations. Formally, the time-evolution of concentrations may be expressed in terms of the stoichiometric matrix and the current vector $J = (J_i)_{i=1,\dots,n}$,

$$\frac{d[A]}{dt} = \mathbb{S}J. \quad (3)$$

Linearizing around given concentrations $([A_i])_{i=1,\dots,|S|}$, one gets for infinitesimal variations $[A] \rightarrow [A] + A$ (mind the notation without square brackets for *variations*)

$$\frac{dA}{dt} = \mathbb{S}J_{lin}([A], A) \quad (4)$$

where $J_{lin,i}([A], A) = \sum_{\ell} J_{lin,i}^{\ell}([A])A_{\ell}$ is linear in the variations. Letting

$$M([A]) := \mathbb{S}J_{lin}([A]), \quad (5)$$

we get the linear system

$$\frac{dA}{dt} = M([A])A. \quad (6)$$

The matrix $M([A])$ is sometimes (but not always) a generalized Markov generator. A case for which $M([A])$ is indeed a generalized Markov generator is when *each reaction has exactly one reactant*, so that its rate is linear in its concentration: the reaction $A_1 \xrightarrow{k_+} s_1B_1 + \dots + s_nB_n$, $n \geq 1$ makes the following additive contribution to $M([A])$,

$$\begin{array}{c} A_1 \\ A_1 \\ B_1 \\ \vdots \\ B_n \\ 0 \\ \vdots \\ 0 \end{array} \left[\begin{array}{ccc} & & \\ -k_+ & & \\ s_1k_+ & & \\ \vdots & & \\ s_nk_+ & 0 & \\ 0 & & \\ \vdots & & \\ 0 & & \end{array} \right]$$

An important particular case is that of a 1-1 reaction $A_1 \xrightarrow{k_+} B_1$; the contribution to $M([A])$ is then simply

$$\begin{array}{c} A_1 \\ A_1 \\ B_1 \\ 0 \\ \vdots \\ 0 \end{array} \left[\begin{array}{ccc} & & \\ -k_+ & & \\ k_+ & & \\ 0 & 0 & \\ \vdots & & \\ 0 & & \end{array} \right]$$

for which the sum of coefficients on the A_1 -column is zero, in coherence with probability preservation.

Interestingly, autocatalytic cores, as proved in [8], satisfy the latter condition – except that the stoichiometry is more general, allowing for reactions of type $sA \rightarrow s_1B_1 + \dots + s_nB_n$, $s \geq 1$. This only turns the top coefficient $M_{A,A}$ into $-sk_+$. The associated killing rate for species A is $(1 - (s_1 + \dots + s_n))k_+$, it is ≤ 0 for reactions of the type $A \xrightarrow{k_+} s_1B_1 + \dots + s_nB_n$, $n \geq 1$ (but not necessarily when $s \neq 1$).

Going one step further, we note that reactions with ≥ 2 reactants have a vanishing rate in the limit when concentrations go to zero. In that limit, furthermore, all killing rates are ≤ 0 . We call this the *zero concentration limit* of networks. In this limit, where the linearized time evolution generator involves only mutations and creation reactions, it is easily conceived that autocatalysis should hold in any reasonable sense. *The adjacency graph associated to the generalized Markov generator $M([A] = 0)$ (see section 3 for its precise construction) is denoted $G(\mathbb{S})$; it depends only on the stoichiometric matrix.*

1.3 Diluted regime and statement of condition (Top)

The present study is devoted to **diluted networks**. These are systems with *low*, but nonzero, concentrations, for which reactions with ≥ 2 reactants exist but have low rate compared to the others. The physical picture is that of a system of reactions of three types:

- (i) reversible reactions, with linear rates, involving *one* reactant and *one* product,



- (ii) irreversible forward reactions involving *one* reactant and *several* products, with linear rates,



such reactions are totally irreversible in the zero concentration limit;

- (iii) and, possibly, the reverse reactions associated to the reactions in (ii),



with nonlinear, but low (compared to (i) and (ii)) or zero reaction rates.

Degradation reactions (which are non-autonomous) may also be included. **Degradation-less diluted networks** are diluted networks in which degradation reactions are either absent or have small enough rates.

In that setting, it is natural to approximate reactions (iii) by their linearizations, which have in any case a small rate compared to reactions of type (i) or (ii). We do not get in general a generalized adjoint Markov generator, because off-diagonal coefficients of $M([A])$ are not necessarily positive. (They *are* positive when reverse reactions are strictly of the form $mA_{i'} \rightarrow A_i$ with $m \geq 2$; see Suppl. Info. for examples and general statements). Even in that case, however, the Feynman-Kac formula holds (see Suppl. Info.), so our general strategy works.

Generally speaking, reactions such as (7) or (8) may be represented in the form of a *hypergraph* called *hypergraph associated to \mathbb{S}* (Anderson 2019, §2), with 'pitchforks' connecting A_i to $A_{i'}$ by s'_ℓ arrows in the case of a one-to-several irreversible reaction. Under these conditions, a natural quantity characterizing the replication rate is the **Lyapunov exponent** $\lambda_{max} \equiv \lambda_{max}(M([A]))$, by definition

$$\lambda_{max} := \max\{\text{Re}(\lambda) \mid \lambda \text{ eigenvalue of } M([A])\}. \quad (10)$$

Under our hypotheses, it can be proved (using Perron-Frobenius theorem) that λ_{max} is an eigenvalue of $M([A])$ with multiplicity 1, and that an eigenvector associated to λ_{max} can be chosen in such a way that all its coordinates are > 0 . When λ_{max} is positive, it characterizes the onset of the exponential growth regime of the system, namely, for small initial concentrations,

$$\frac{[A_j](t)}{\max_i ([A_i](t=0))} \approx e^{\lambda_{max} t} \quad (11)$$

for all species $A_j \in \mathcal{S}$, for not-too-large time values t . When $\lambda_{max} > 0$, we say that the reaction network is **(strongly) autocatalytic in the dynamical sense** – or **(strongly) dynamically autocatalytic** –. Generalizing (in order to include the case of reducible networks, see below), we say that the reaction network is **weakly autocatalytic in the dynamical sense** (or **weakly dynamically autocatalytic**) if $\lambda_{max} > 0$ is an eigenvalue of $M([A])$ (multiplicity can be arbitrary, and Jordan blocks associated to λ_{max} may be non-trivial), and an eigenvector v associated to λ_{max} may be chosen in such a way that all its coordinates are ≥ 0 . Then (11) remains valid for all species A_j such that $v_j > 0$.

Turning now to the stoichiometric side, in coherence with the above discussion, we neglect altogether reverse reactions (iii): we say that *a diluted network is stoichiometrically autocatalytic if there exists a positive reaction vector c such that $\mathbb{S}c > 0$, where the orientation of reversible, type (i) reactions is arbitrary, but forward reactions (ii) are given positive orientation*.

We can now state our main result (see Theorem 3.1 below); recall that $G(\mathbb{S})$ is the adjacency graph associated to the generalized Markov matrix $M([A] = 0)$. Thinking

of it as if it were the graph of a conventional Markov matrix, we decompose it into classes (see Suppl. Info. §6.5), with probability flow flowing downstream from minimal to maximal classes. Letting \mathcal{C} be one of the classes, we now define its set of **internal reactions**. If $\mathcal{R} : A \rightarrow s_1 A'_1 + \dots + s_n A'_n$ is an irreversible reaction such that $A \in \mathcal{C}$, and $\{i = 1, \dots, n \mid A'_i \in \mathcal{C}\} = \{1, \dots, n'\}$ for some $n' \geq 1$, then we introduce the **truncated reaction** $\mathcal{R}_{\mathcal{C}} : A \rightarrow s_1 A'_1 + \dots + s_{n'} A'_{n'}$. Internal reactions of \mathcal{C} are then: (i) reversible reactions $A \rightleftharpoons A'$ with $A, A' \in \mathcal{C}$; (ii) truncated reactions $\mathcal{R}_{\mathcal{C}} : A \rightarrow s_1 A'_1 + \dots + s_{n'} A'_{n'}$ as above, with $A, A'_1, \dots, A'_{n'} \in \mathcal{C}$. Note that, if $n' = 1$, we obtain a reaction of a new type: an *irreversible 1-1 reaction*.

What we prove is the following:

1. *diluted networks are stoichiometrically autocatalytic if and only if the following topological condition on the hypergraph associated to \mathbb{S} holds,*

(Top) : **all minimal classes of $G(\mathbb{S})$ contain at least one internal one-to-several irreversible reaction,**

i.e. each minimal class \mathcal{C} contains a truncated reaction $\mathcal{R}_{\mathcal{C}} : A \rightarrow s_1 A'_1 + \dots + s_{n'} A'_{n'}$ with $s := \sum_{i=1}^{n'} s_i \geq 2$;

2. *the latter topological condition implies weak dynamical autocatalysis in the diluted regime, i.e. for small enough concentrations, if there are no degradation reactions, or, more generally, if their rates are small enough. Strong dynamical autocatalysis holds under the same conditions if the network is irreducible.*


When $G(\mathbb{S})$ is *irreducible*, condition 1. states that at least one irreversible forward reaction (8) must be present in the system for stoichiometric autocatalysis to hold – an obvious

statement in view of Gordan's theorem since $\mathbf{n} \cdot \mathbb{S} = 0$ if $\mathbf{n} = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}$ and only 1 – 1

reversible reactions (7) are present in the system –, and most importantly, that this is a sufficient condition. The presence of irreversible forward reactions is also necessary for dynamical autocatalysis to hold – otherwise only mutation-like coefficients are present, kinetic equations are those of a conventional Markov system, and then it is known that all generator eigenvalues have ≤ 0 real part –, but 2. states again that this is a sufficient condition.

When $G(\mathbb{S})$ is *reducible*, it is easy to see that the condition (Top) on minimal classes in 1. is necessary. Namely, let \mathcal{C} be one of the minimal classes, and $\mathbb{S}_{\mathcal{C}}$ the stoichiometric matrix associated with species in \mathcal{C} and internal reactions of \mathcal{C} . If all internal reactions are 1-1, then the same argument based on Gordan's theorem implies that, for every positive reaction vector $c_{\mathcal{C}}$, the balance $(\mathbb{S}_{\mathcal{C}} c_{\mathcal{C}})_i$ is ≤ 0 for at least one of the species $i \in \mathcal{C}$.

Now, including *external* (for \mathcal{C}) reactions $\mathcal{R} : A_i \rightarrow s_1 A'_1 + \dots + s_n A'_n$ with $A_i \in \mathcal{C}$ and $A'_1, \dots, A'_n \notin \mathcal{C}$ can only worsen the balance for species A_i , while other reactions $A' \rightarrow s_1 A'_1 + \dots + s_n A'_n$ with $A' \notin \mathcal{C}$ – and therefore, $A'_1, \dots, A'_n \notin \mathcal{C}$ also since \mathcal{C} is minimal – do not change it. Thus, for example, the network with hypergraph



featuring two reversible reactions $A_0 \rightleftharpoons A_1$, $A_1 \rightleftharpoons A_2$ (in blue) and an irreversible reaction $\mathcal{R} : A_0 \rightarrow A_2 + A'_0$ (in red) coupling the minimal class $\mathcal{C} = \{A_0, A_1, A_2\}$ to another class $\mathcal{C}' = \{A'_0\}$, does not satisfy (Top), because the truncated irreversible reaction $\mathcal{R}_{\mathcal{C}} : A_0 \rightarrow A_2$ is 1-1.

Looking closely at the (generalized) eigenspace associated to λ_{max} in the reducible case, one realizes that a complete discussion of the nature of dynamical autocatalysis (e.g. multiplicity of λ_{max} , and support of an associated eigenvector v , i.e. the set of species A_j such that $v_j > 0$) cannot rely only on topological considerations; see detailed example p. 26. In addition, the presence of reverse reactions (9), even at negligible rates from a biological or chemical point of view, modifies the above characteristics. Let us explain this subtlety on two minimal examples. Assume there are two classes $\mathcal{C}, \mathcal{C}'$ with probabilistic flow flowing out of \mathcal{C} into \mathcal{C}' ,




and that, as a first case, (i) *only the maximal (downstream) class \mathcal{C}' contains an internal irreversible reaction* (so that (Top) is not satisfied); then (excluding reverse reactions going upstream from \mathcal{C}' to \mathcal{C}) the network restricted to \mathcal{C}' is irreducible and (strongly) dynamically autocatalytic; thus the whole network is weakly dynamically autocatalytic. In this case however, whatever \mathcal{C} reactants present in the solution disappear exponentially in time in favor of species in \mathcal{C}' . Now imagine choosing one of the reactions (8) connecting \mathcal{C} to \mathcal{C}' and adding the reverse reaction with a negligible rate $O(\varepsilon)$, $0 < \varepsilon \ll 1$. This makes the network irreducible, implying strong dynamical autocatalysis, while perturbing only slightly λ_{max} . The associated positive eigenvector $v = v(\varepsilon)$ – unique up to normalization – will have nonzero but very small coefficients along \mathcal{C} , making it probably difficult in practice to observe exponential increase of the corresponding species. Next, consider as a second case the possibility that (ii) *(Top) is satisfied, so that the network restricted to \mathcal{C} (i.e. suppressing all \mathcal{C}' -products of reactions with reactant in \mathcal{C}) is autocatalytic, but \mathcal{C}' contains no irreversible reaction, hence is not autocatalytic*. Then the network (as proved in section 4) is already strongly autocatalytic in itself.

Dismissing these subtleties and summarizing, our result may be rephrased as follows: *stoichiometric autocatalysis implies dynamical autocatalysis* in our diluted regime and, in absence of degradation reactions; and (at least in the case of irreducible networks), it may be said that the converse is also true.

Remark. As follows from the above, (Top) depends on the hypergraph associated to \mathbb{S} , not only on the graph $G(\mathbb{S})$. However, the main tools in the proof are based on properties of $G(\mathbb{S})$.

2 A motivating example: the simplest autocatalytic core

We treat in this section the simplest type I autocatalytic core in the classification of [8]. It involves two chemostatted species (A, A'), which may be thought of as a redox or energy carrier (ATP/ADP) couple, or as fuel and waste [16]; two dynamical species (B, B_1); and two reactions



Autocatalysis is made possible by the duplication reaction $B_1 \xrightarrow{\nu_+} 2B + A'$. We also include degradation reactions



The degradationless diluted regime which is the main topic of the article is defined by

- (i) (low concentrations) $[B], [B_1] \ll 1$. Kinetic equations lack any reference concentration or volume to produce adimensional quantities, and chemostatted quantities $[A], [A']$ are not limited, so (by simple rescaling of the concentrations) this criterion is equivalent to

$$\nu_- [B] \ll 1. \quad (14)$$

In other words, the reverse of the duplication reaction is rate-limited.

- (ii) (no degradation) $a_0, a_1 = 0$. Our analysis actually extends (by perturbation) to low enough degradation rates,

$$a_0, a_1 \ll 1. \quad (15)$$

Kinetic equations are:

$$\left(\frac{d}{dt} + a_0\right)[B] = 2\nu_+[B_1] - (k_{on}[A][B] - k_{off}[B_1]) - 2\nu_-[B]^2[A'] \quad (16)$$

$$\left(\frac{d}{dt} + a_1\right)[B_1] = -\nu_+[B_1] + (k_{on}[A][B] - k_{off}[B_1]) + \nu_-[B]^2[A'] \quad (17)$$

When $\nu_- = 0$, these equations are linear, otherwise we linearize around $([B], [B_1])$, and find the system

$$\frac{d}{dt} \begin{pmatrix} B \\ B_1 \end{pmatrix} = M \begin{pmatrix} B \\ B_1 \end{pmatrix} \quad (18)$$

where (B, B_1) is an infinitesimal variation around $([B], [B_1])$, and

$$M = \begin{bmatrix} -k_{on}[A] - 4\nu_-[A'][B] - a_0 & k_{off} + 2\nu_+ \\ k_{on}[A] + 2\nu_-[A'][B] & -k_{off} - \nu_+ - a_1 \end{bmatrix} \quad (19)$$

Note that off-diagonal elements of M are > 0 , so that, by the Perron-Frobenius theorem, the spectrum of M consists of two complex numbers $\lambda_{max}(M), \lambda_{min}(M)$ with $\lambda_{max}(M)$ real, and $\lambda_{max}(M) > \text{Re } \lambda_{min}(M)$. Furthermore, M has an eigenvector for the eigenvalue $\lambda_{max}(M)$ with positive coefficients. Write $M = \begin{bmatrix} -a & b \\ c & -d \end{bmatrix}$. Explicit computations actually produce two real numbers,

$$\lambda_{max}(M) = \frac{1}{2} \left(-(a+d) + \sqrt{(a+d)^2 - 4 \det(M)} \right) = \frac{1}{2} \left(-(a+d) + \sqrt{(a-d)^2 + 4bc} \right) \quad (20)$$

$$\text{and } \lambda_{min}(M) = \frac{1}{2} \left(-(a+d) - \sqrt{(a-d)^2 + 4bc} \right).$$

Lemma 2.1 (see [14]) Let $M = \begin{bmatrix} -a & b \\ c & -d \end{bmatrix}$ ($a, b, c, d > 0$) and $\lambda_{max} = \lambda_{max}(M)$ the eigenvalue of M with largest real part. Then the following alternative holds,

- (i) If $\det(M) = ad - bc < 0$, then $\lambda_{max} > 0$;
- (ii) if $\det(M) = 0$, then $\lambda_{max} = 0$;
- (iii) if $\det(M) > 0$, then $\lambda_{max} < 0$.

Autocatalysis is then equivalent to the condition $\det(M) < 0$. We now check that, in the degradationless diluted regime defined by (14, 15),

$$\det(M) = -k_{on}[A]\nu_+ + O(\nu_-[B]) + O(a_0) + O(a_1) < 0. \quad (21)$$

Going beyond this particular regime, autocatalysis is not the rule. Let us consider two specific cases:

- (i) (no reverse reaction) We neglect reverse reactions by setting $k_{off} = 0$ and $\nu_- = 0$. Then

$$\det(M) = -2k_{on}[A]\nu_+ + (k_{on}[A] + a_0)(\nu_+ + a_1) < 0 \quad (22)$$

if and only if (see King's criterion [22]) the product of the specificities of positively oriented reactions along the replication cycle $B \rightarrow B_1 \rightarrow 2B$ is larger than $\frac{1}{2}$,

$$\frac{k_{on}[A]}{k_{on}[A] + a_0} \frac{\nu_+}{\nu_+ + a_1} > \frac{1}{2}. \quad (23)$$

- (ii) (no degradation) We assume here that $a_0 = a_1 = 0$. Then

$$\det(M) = -k_{on}[A]\nu_+ + 2k_{off}\nu_-[A'][B] < 0 \quad (24)$$

if and only if

$$[B] < [B]_{max} := \frac{k_{on}[A]}{k_{off}} \times \frac{\nu_+}{2\nu_-[A']}, \quad (25)$$

or equivalently,

$$\frac{k_{on}[A]}{k_{off}} \times \frac{\nu_+}{2\nu_-[A'][B]} > 1, \quad (26)$$

a criterion somewhat analogous to King's criterion, but featuring the ratio (product of forward reaction rates)/(product of reverse reaction rates).

3 (Top) characterizes stoichiometric autocatalysis in diluted networks

We reconsider in this section the stoichiometric autocatalysis criterion of Blokhuis-Lacoste-Nghe [8] in the case when all concentrations are *low*. Under such circumstances, reactions involving > 1 reactants have very small rate. It is therefore reasonable to discard them from the beginning when dealing with the stoichiometric definition of autocatalysis.

This leads us to introduce a subclass of stoichiometrically autocatalytic networks, which we call **diluted stoichiometrically autocatalytic networks**. Consider a reaction network with species set \mathcal{S} and reaction set $\{1, \dots, N\}$, choose an orientation for each reaction, characterizing *forward* reactions, by opposition to *reverse* reactions. Degradation reactions possibly exist, but are not included in the reaction set, and play no rôle in the discussion. Choosing some arbitrary ordering of reactions, we get a stoichiometric matrix S . Then we require the following conditions:

- (i) the reaction network is unambiguous and autonomous;

- (ii) there exists a positive reaction vector $c \in (\mathbb{R}_+)^N$ such that $\mathbb{S}c > 0$;
- (iii) **each reaction has only one reactant, and its stoichiometry is 1.**

The last condition (iii) restricts the class introduced in [8]. Recall however (as already mentioned) that all autocatalytic cores satisfy partially this condition, in the sense that all reactions have exactly one reactant (with arbitrary stoichiometry, however).


Diluted networks, associated graph. Removing assumption (ii), we get the definition of a **diluted network**, which is the general class of topological networks of interest in the present work. We associate to such a reaction network its **split graph** (or simply **graph**) $G(\mathbb{S})$, which depends only on the stoichiometric matrix, and corresponds mathematically to the linearization of the kinetic network in the zero concentration limit. It may be defined topologically as follows: (direct) reactions of the type $A \rightarrow s_1 B_1 + \dots + s_n B_n$ ($n \geq 1, s_1, \dots, s_n \in \mathbb{N}^*$) such that $s_1 + \dots + s_n \geq 2$, i.e. with > 1 products, are totally irreversible in the limit of vanishing concentrations, therefore they contribute to $G(\mathbb{S})$ irreversible arrows




upon **splitting** the reaction into reactions with unique products. On the other hand, forward reactions of the type $A \rightarrow B$ with only one product are reversible; therefore, they contribute to $G(\mathbb{S})$ reversible arrows $A \rightleftharpoons B$. In case of multiple arrows $A \rightarrow B$, we only keep one, in order not to have multiple edges from A to B . This happens if there are several competing irreversible reactions $A \rightarrow sB + s_2 B_2 + \dots + s_n B_n$, $A \rightarrow s'B + s'_2 B'_2 + \dots + s'_n B'_n$, or if irreversible reactions $A \rightarrow sB + \dots$ and a reversible reaction $A \rightleftharpoons B$ coexist. We always assume that $G(\mathbb{S})$ is *connected* (otherwise one can reduce the analysis to each of the subsystems defined by the connected components).

Having a graph instead of a hypergraph with pitchforks connecting several reactants and several products (see below, and examples in §5.1 and 5.2) is a major simplification. To be precise, we note that $G(\mathbb{S})$ is sometimes not quite enough to characterize stoichiometric autocatalysis: in case an irreversible reaction $A \rightarrow sB + \dots$ and the reversible reaction $A \rightleftharpoons B$ coexist (so that A and B are in the same class \mathcal{C} , see below), the graph $G(\mathbb{S})$ by itself does not keep track of the existence of the irreversible reaction. Then we keep the memory of the irreversible transition $A \rightarrow B$ by saying that \mathcal{C} contains an *internal irreversible reaction*. In case $A \rightarrow B + \dots$ is not in competition with a reversible reaction, but A and B are in the same class thanks to the presence of an irreversible reaction $B \rightarrow A + \dots$, both split reactions $A \rightarrow B$ and $B \rightarrow A$ are considered as internal irreversible reactions. A simple way to summarize these rules is to decide that *reversible arrows are painted blue, irreversible arrows are painted red, and red prevails*. Thus we get a graph with *two-colored edges*. This is sometimes useful, but still not enough to define our topological condition (Top) when the graph is not irreducible (see §1.3). Classes are defined below without taking the color of the arrows into account.


Classes. Upon linearizing the time-evolution equations, while neglecting reverse reactions, one obtains a *generalized Markov matrix* (see Suppl. Info.) M with graph $G(\mathbb{S})$. This justifies resorting to the usual description of G in terms of communicating classes, connected by irreversible arrows. Arrows define a partial order of classes, with $\mathcal{C}' > \mathcal{C}$ if there is a path from \mathcal{C} to \mathcal{C}' , i.e. if \mathcal{C}' is downstream of \mathcal{C} . In Suppl. Info. (§6.5), the reader will find several examples worked out in details: cores of type I and III,




and the " $A_1A_2A_3 \rightarrow B_1B_2B_3$ " autocatalytic kinetic reaction network, and its graph $G_{(123) \rightarrow (1'2'3')}$, where (A_1, A_2, A_3) , resp. (B_1, B_2, B_3) are encoded by indices $(1, 2, 3)$, resp. $(1', 2', 3')$:



Note that the stoichiometry is not indicated, nor is it important in the analysis that follows, once understood that irreversible arrows come from splitting reactions with > 1 products. As a matter of fact, Type (I) cores (B_0, \dots, B_n) have originally a "pitchfork" reaction $B_n \rightarrow 2B_0$



Type (III) cores, involving species A_i , $i = 0, \dots, n$, B'_i , $i' = 0', \dots, n'$, B''_i , $i'' = 0'', \dots, n''$, have originally a pitchfork



Only one-sided arrows indicate the location of the original hypergraph pitchforks. The one-sided arrow $n \rightarrow 0$ in (I) indicates any reaction $B_n \rightarrow mB_0$ with $m = 2, 3, \dots$. The one-sided arrows $A_n \rightarrow B''_{0''}, A_n \rightarrow B'_{0'}$ come either from $A_n \rightarrow s''B''_{0''} + s'B'_{0'}$, $s', s'' = 1, 2, \dots$ or from $(A_n \rightarrow m''B''_{0''}, A_n \rightarrow m'B'_{0'})$, $m', m'' = 2, 3, \dots$, or from a combination of these.

All cores are irreducible. The " $A_1A_2A_3 \rightarrow B_1B_2B_3$ " network, on the other hand, has two classes, $\mathcal{C} = (1, 2, 3)$ and $\mathcal{C}' = (1', 2', 3')$, with \mathcal{C}' downstream of \mathcal{C} . The partial ordering defines in particular minimal (upstream) and maximal (downstream) classes; here, \mathcal{C} is minimal, and \mathcal{C}' is maximal.

Our main result is the following:

Theorem 3.1 *1. A diluted network with stoichiometric matrix \mathbb{S} is a diluted stoichiometrically autocatalytic network if and only if the following topological condition (Top) of the hypergraph associated to \mathbb{S} is satisfied:*

(Top) : each of the minimal classes of $G(\mathbb{S})$ contains at least one internal one-to-several irreversible reaction.

2. A diluted network satisfying (Top) is weakly autocatalytic in the dynamical sense if there are no degradation reactions, or, more generally, if their rates are small enough. Furthermore, it is strongly autocatalytic in the dynamical sense under the same conditions if the network is irreducible.

We prove in the rest of the section the first part of the Theorem, concerning *stoichiometric autocatalysis*; the second part will be proved in the next section.

Stoichiometric autocatalysis, at least in the case of an irreducible network, can be proven quite simply by playing directly with the columns of the stoichiometric matrix \mathbb{S} ; see Suppl. Info. 6.3. Instead, we provide here a general demonstration using properties of $G(\mathbb{S})$. Though a little more involved, it has the advantage of exploiting the properties of an underlying *auxiliary* Markov chain, which will also play a major rôle in §4. In the case of a reducible network, arguments rely on the class decomposition of the graph $G(\mathbb{S})$.

We have already proved in the Introduction that (Top) is necessary for a diluted network to be autocatalytic. So the interesting part is to show that (Top) is a *sufficient condition* for autocatalysis. We split the proof into several points. The general idea is to construct an explicit reaction vector c which depends on the choice of a kinetic rate for each reaction, and is a perturbation of the stationary flow vector for an auxiliary Markov chain.

The chemical balance for species A_k associated to a reaction $\mathcal{R} : s_1A_{i_1} + \dots + s_nA_{i_n} \rightarrow s'_1A_{i'_1} + \dots + s'_{n'}A_{i'_{n'}}$ will be denoted $\delta_{\mathcal{R}}[A_k] = -\sum_j s_j\delta_{k,i_j} + \sum_{j'} s'_{j'}\delta_{k,i'_{j'}}$. Then the total

chemical balance for species A_k associated to the combination of reactions represented by the reaction vector c is $\delta[A_k] = \sum_{\mathcal{R}} c_{\mathcal{R}} \delta_{\mathcal{R}}[A_k]$.

A. (stationary flows for split graph). Theorem 3.1 (1) is obtained by perturbation from the following remark. One can define an **auxiliary Markov chain** $(\tilde{X}(t))_{t \geq 0}$ (a conventional, continuous-time Markov chain, i.e. with vanishing killing rates) with transition rates $\tilde{k}_{i \rightarrow j}$ obtained by superposing the following transitions:

- (i) Reversible transitions with rates $k_{i \rightarrow j}, k_{j \rightarrow i}$ are associated to 1-1 reactions of the type $\mathcal{R}_{i \rightarrow j} : A_i \xrightarrow{k_{i \rightarrow j}} A_j, \mathcal{R}_{j \rightarrow i} : A_j \xrightarrow{k_{j \rightarrow i}} A_i$;
- (ii) Irreversible transitions with rates $s_j k_i^+$ are associated to split irreversible 1-1 reactions $\tilde{\mathcal{R}} : A_i \xrightarrow{s_j k_i^+} A_j, j = i_1, \dots, i_n$ coming from the one-to-several irreversible reaction $A_i \xrightarrow{k_i^+} s_{i_1} A_{i_1} + \dots + s_{i_n} A_{i_n}$.

The associated adjoint Markov generator is obtained by summing matrices with only two non-vanishing coefficients as on p.8; then the sum of coefficients on any column is zero, which ensures probability preservation. In other words, $\tilde{k}_{i \rightarrow j} = \sum_{\tilde{\mathcal{R}} : A_i \rightarrow A_j} \tilde{k}_{i \rightarrow j}(\tilde{\mathcal{R}})$, where, depending on the split reaction $\tilde{\mathcal{R}} : A_i \rightarrow A_j$, one has defined: $\tilde{k}_{i \rightarrow j}(\tilde{\mathcal{R}}) = k_{i \rightarrow j}$ (one-to-one reaction) or $s_j k_i^+$ (split forward reaction $A_i \rightarrow A_j$ coming from a one-to-several reaction $A_i \rightarrow s_j A_j + \dots$) or 0 (excluded reverse reaction).

Assume the graph $G(\mathbb{S})$ is irreducible. Then the auxiliary Markov chain $(\tilde{X}(t))_{t \geq 0}$ is irreducible; it reproduces correctly the transition rates of the kinetic network from A_i to A_{i_ℓ} for irreversible transitions (ii), but *increases* the exit rate from A_i , since $\frac{d[A_i]}{dt} = -s k_i^+[A_i]$ (by probability conservation) with $s = \sum_{\ell} s_{i_\ell} \geq 2$ for the Markov chain, as compared to $\frac{d[A_i]}{dt} = -k_i^+[A_i]$ for the kinetic network. The auxiliary Markov chain admits exactly one stationary probability measure $\mu = (\mu_i)_{i=1, \dots, |\mathbb{S}|}$. Define $\tilde{c}_{i \rightarrow j}(\tilde{\mathcal{R}}) := \begin{cases} \mu_i \tilde{k}_{i \rightarrow j}(\tilde{\mathcal{R}}) & \text{if } \tilde{\mathcal{R}} : i \rightarrow j \\ 0 & \text{else} \end{cases}$, and let $\tilde{c}_{i \rightarrow j} = \sum_{\tilde{\mathcal{R}} : i \rightarrow j} \tilde{c}_{i \rightarrow j}(\tilde{\mathcal{R}}) = \mu_i \tilde{k}_{i \rightarrow j}$ be the stationary flow along the edges. Then the antisymmetrized quantity $\tilde{J}_{i \rightarrow j} := \sum_{\tilde{\mathcal{R}}} \tilde{J}_{i \rightarrow j}(\tilde{\mathcal{R}}) \equiv \sum_{\tilde{\mathcal{R}}} \left\{ \tilde{c}_{i \rightarrow j}(\tilde{\mathcal{R}}) - \tilde{c}_{j \rightarrow i}(\tilde{\mathcal{R}}) \right\} = \tilde{c}_{i \rightarrow j} - \tilde{c}_{j \rightarrow i}$ is the associated current, and *the total current for species i vanishes by stationarity, i.e. hence*

$$\sum_j \tilde{J}_{i \rightarrow j} = 0. \quad (28)$$

Choice of the reaction vector c . Going back to the initial network, *we now define* $c(\mathcal{R}) := \tilde{c}_{i \rightarrow j}(\mathcal{R}) = \mu_i k_{i \rightarrow j}$ for the reversible one-to-one reaction $\mathcal{R} : A_i \xrightarrow{k_{i \rightarrow j}} A_j$, and

$c(\mathcal{R}) := \mu_i k_i^+$ for the irreversible one-to-several reaction $\mathcal{R} : A_i \xrightarrow{k_i^+} s_{i_1} A_{i_1} + \dots + s_{i_n} A_{i_n}$. Note that (for convenience) we have chosen to accept both orientations for reversible 1-1 reactions; this is equivalent to choosing an orientation for each of them and letting $c(\mathcal{R}_{(i,j)}) := c(\mathcal{R}_{i \rightarrow j}) - c(\mathcal{R}_{j \rightarrow i})$ (the orientation may be chosen in such a way that $c(\mathcal{R}_{(i,j)}) \geq 0$). The chemical balance $\delta[A_i] := (\mathbb{S}c)_i$ for species i is obtained by summing

$$-\tilde{c}_{i \rightarrow j}(\mathcal{R}) + \tilde{c}_{j \rightarrow i}(\mathcal{R}) = -\tilde{J}_{i \rightarrow j}(\mathcal{R}) \quad (29)$$

for a reversible one-to-one reaction \mathcal{R} connecting i and j ,

$$-c(\mathcal{R}) = -\mu_i k_i^+ \quad (30)$$

for the reactant of a one-to-several reaction $\mathcal{R} : A_i \rightarrow \dots$, and

$$+s_i \mu_j k_j^+ = +\tilde{c}_{j \rightarrow i}(\tilde{\mathcal{R}}) \quad (31)$$

for products of a one-to-several reaction $\mathcal{R} : A_j \rightarrow s_i A_i + \dots$, split into several 1-1 reactions including $\tilde{\mathcal{R}} : A_j \rightarrow s_i A_i$. By construction, we obtain

$$\delta[A_i] = -\sum_j \tilde{J}_{i \rightarrow j} \quad (32)$$

if species i is *not* the reactant of a one-to-several reaction; thus, in that case, $\delta[A_i] = 0$. If, on the other hand, i is the reactant of a one-to-several reaction $A_i \xrightarrow{k_i^+} s_{i_1} A_{i_1} + \dots + s_{i_n} A_{i_n}$ with associated split reactions $\tilde{\mathcal{R}}_{i,i_\ell} : A_i \rightarrow A_{i_\ell}$, then the associated balance for $[A_i]$ is

$$-\mu_i k_i^+ > -\sum_\ell \tilde{c}_{i \rightarrow i_\ell}(\tilde{\mathcal{R}}_{i,i_\ell}) = -\mu_i s k_i^+ \quad (33)$$

with $s = \sum_\ell s_{i_\ell} > 1$. Comparing with the above stationarity equation (28), we may conclude: *our choice for the vector c yields a strictly positive balance for reactants of a one-to-several reaction, and zero balance for all other species.*

Remark. If the graph is over-connected, i.e. if reversible 1-1, or one-to-several irreversible, reactions can be removed without breaking irreducibility, then the auxiliary Markov chain may be defined while leaving them out, yielding another simpler set of coefficients $c_{\mathcal{R}}$ that vanish for left-out reactions.

B. (irreducible networks). The reaction vector c constructed in **A.** is not quite satisfactory yet. We now turn to a *perturbation argument for irreducible networks*, ensuring


that there exist vectors $\delta c^q = \begin{pmatrix} (\delta c^q)_{\mathcal{R}_1} \\ \vdots \\ (\delta c^q)_{\mathcal{R}_N} \end{pmatrix}$, $q = 1, 2, \dots$ vanishing for q large enough such

that $\mathbb{S}(c + \sum_{q \geq 1} \varepsilon^q \delta c^q) > 0$ for all small enough $\varepsilon > 0$. By hypothesis, there exists at least one irreversible reaction. Choose one, $\mathcal{R}_0 : A_0 \xrightarrow{k_0^+} s_{i_1} A_{i_1} + \dots + s_{i_n} A_{i_n}$, and define $c \equiv c(\mu)$ as in the previous paragraph, μ being the stationary probability measure for \tilde{X} . Since the balance for $[A_0]$ is > 0 , we can tilt μ by a small amount in direction 0, i.e. replace μ_0 by $\mu_0 + \varepsilon$, while keeping $\delta[A_0] > 0$. This is equivalent to saying that $c \rightsquigarrow c + \varepsilon \delta c^1$, with $\delta c^1(\mathcal{R}_0) = k_0^+$, yielding $(c + \varepsilon \delta c^1)(\mathcal{R}_0) = (\mu_0 + \varepsilon)k_0^+$; similarly, $\delta c^1(\mathcal{R}) = k_{0 \rightarrow j}$, resp. $(k'_0)^+$ for all other possible reactions $A_0 \xrightarrow{k_{0 \rightarrow j}} A_j$ or $A_0 \xrightarrow{(k'_0)^+} s'_{i'_1} A_{i'_1} + \dots + s'_{i'_{n'}} A_{i'_{n'}}$ with reactant A_0 ; and $\delta c^1(\mathcal{R}') = 0$ for all other reactions. But then $\delta[A_{i_\ell}]$ is shifted by $+\varepsilon s_{i_\ell} k_0^+$, and possibly other positive coefficients ($+\varepsilon k_{0 \rightarrow i_\ell}$ or $+\varepsilon s'_{i'_\ell} (k'_0)^+$), so the balance for species 0 and for all products of \mathcal{R}_0 is now > 0 ; more precisely, $\delta[A_0]$ is of order ε^0 , while $\delta[A_{i_\ell}]$, $\ell = 1, \dots, n$ – and similarly, the balance for all products of reactions with reactant A_0 – are of order ε^1 .

We now let $\mathcal{S}_0 := \{0\}$, define $\mathcal{S}_1 \subset \mathcal{S}$ to be made up of 0, together with all products of reactions having 0 as reactant, and consider products of reactions having as reactant one of the elements of the set $\mathcal{S}_1 \setminus \mathcal{S}_0$. Since the graph is irreducible, the corresponding set of reactions can be empty only if $\mathcal{S}_1 = \mathcal{S}$. If this is not the case, tilt μ by a small uniform amount in all directions indexed by the set $\mathcal{S}_1 \setminus \mathcal{S}_0$, i.e. replace μ_i by $\mu_i + \varepsilon^2$ for all $i \in \mathcal{S}_1 \setminus \mathcal{S}_0$. For convenience, we reindex the set of species so that $\mathcal{S}_1 \setminus \mathcal{S}_0 = \{1, \dots\}$.

Choosing one of the above reactions, either one-to-several $\mathcal{R}_1 : A_1 \xrightarrow{k_1^+} s_{i_1} A_{i_1} + \dots + s_{i_n} A_{i_n}$ or one-to-one, $\mathcal{R}_{1 \rightarrow j} : A_i \xrightarrow{k_{i \rightarrow j}} A_j$, this is equivalent to saying that $c \rightsquigarrow c + \varepsilon \delta c^1 + \varepsilon^2 \delta c^2$, with $\delta c^2(\mathcal{R}_1) = k_1^+$, resp. $\delta c^2(\mathcal{R}_{1 \rightarrow j}) = k_{1 \rightarrow j}$. We thus shift $\delta[A_i]$, $i \in \mathcal{S}_1 \setminus \mathcal{S}_0$, by $-O(\varepsilon^2)$, and simultaneously $\delta[A_{i'}]$ (i' ranging in the set of products of reactions having as reactant one of the elements of $\mathcal{S}_1 \setminus \mathcal{S}_0$, including possibly species in \mathcal{S}_1) by $+O(\varepsilon^2)$. The $\delta[A_i]$ were of order ε^0 , resp. ε^1 at previous step for $i \in \mathcal{S}^0$, resp. $\mathcal{S}^1 \setminus \mathcal{S}^0$; the ε^2 -corrections do not change these orders, but ensure that now $\delta[A_i]$, $i \in \mathcal{S}_2 \setminus \mathcal{S}_1$ are of order ε^2 , where $\mathcal{S}_2 \setminus \mathcal{S}_1$ is the set of new products. We stop the induction in q as soon as we have exhausted all species, i.e. the maximum index q is the minimum index such that $\mathcal{S}_q = \mathcal{S}$.

A simple example. Consider the network with species A_0, A_1, A_2 , reversible 1-1 reactions $A_0 \rightleftharpoons A_2$ and $A_1 \rightleftharpoons A_2$, and a single irreversible one-to-several reaction $\mathcal{R}_0 : A_0 \rightarrow A_1 + A_2$ with $s = 2$. The graph is



following the convention that "red prevails". The network is irreducible. Choose all rates

to be equal to 1. Then the adjoint Markov generator of the auxiliary chain is

$$\begin{pmatrix} -2 & & \\ 1 & & \\ 1 & & \end{pmatrix} + \begin{pmatrix} -1 & 1 & \\ & & \\ 1 & -1 & \end{pmatrix} + \begin{pmatrix} & & \\ -1 & 1 & \\ & & \\ 1 & -1 & \end{pmatrix} = \begin{pmatrix} -3 & 0 & 1 \\ 1 & -1 & 1 \\ 2 & 1 & -2 \end{pmatrix}. \quad (34)$$

Stationary measures are multiples of $\mu := \begin{pmatrix} 1 \\ 4 \\ 3 \end{pmatrix}$. Stationary flows are $\tilde{c}_{0 \rightarrow 1} = 1, \tilde{c}_{0 \rightarrow 2} =$

2; $\tilde{c}_{1 \rightarrow 0} = 0, \tilde{c}_{1 \rightarrow 2} = 4; \tilde{c}_{2 \rightarrow 0} = 3, \tilde{c}_{2 \rightarrow 1} = 3$, and then stationary currents are $\tilde{J}_{0 \rightarrow 1} = 1, \tilde{J}_{0 \rightarrow 2} = -1, \tilde{J}_{1 \rightarrow 2} = 1$. Following our construction, we choose for reaction vector c with $c(\mathcal{R}_0) = 1$ and $c(\mathcal{R}_{0 \rightarrow 2}) = 1, c(\mathcal{R}_{2 \rightarrow 0}) = 3, c(\mathcal{R}_{1 \rightarrow 2}) = 4, c(\mathcal{R}_{2 \rightarrow 1}) = 3$. Then


$$\delta[A_1] = c(\mathcal{R}_0) - c(\mathcal{R}_{1 \rightarrow 2}) + c(\mathcal{R}_{2 \rightarrow 1}) = 0; \quad (35)$$

similarly, $\delta[A_2] = 0$; and $\delta[A_0] = -c(\mathcal{R}_0) - c(\mathcal{R}_{0 \rightarrow 2}) + c(\mathcal{R}_{2 \rightarrow 0}) = 1$, which can be identified with $(s-1)\mu_0 k_0^+$ using the notations of the proof. We perturb it by a one-step construction since $\mathcal{S}_1 = \{0, 1, 2\}$: we replace c by $c + \varepsilon \delta c^1$ with $\delta c^1(\mathcal{R}_0) = \delta c^1(\mathcal{R}_{0 \rightarrow 1}) = \delta c^1(\mathcal{R}_{0 \rightarrow 2})$. Thus the perturbed balance $\delta[A_0] = 1 - \varepsilon, \delta[A_1] = +\varepsilon, \delta[A_2] = +2\varepsilon$ is > 0 for all species as soon as $0 < \varepsilon < 1$.

C. (reducible networks). We must finally adapt the above argument to the case of a reducible network. To have a picture in mind, the reader may think of the "contracted graph"

$$\mathcal{T}_{(123) \rightarrow (1'2'3')} = \begin{array}{c} \mathcal{C} \\ \downarrow \\ \mathcal{C}' \end{array}$$

of the " $A_1 A_2 A_3 \rightarrow B_1 B_2 B_3$ " network (see §6.5), or, for a more general example,



In both examples here, there is a unique minimal class, \mathcal{C} , and a unique maximal class, \mathcal{C}' . Note that arrows go downwards, defining a probability flow from minimal classes to maximal classes. We define the *height* $h(\mathcal{C}'')$ of a class \mathcal{C}'' to be the minimal distance on the contracted graph from a minimal class to it. Here e.g. $h(\mathcal{C}) = 0, h(\mathcal{C}') = 1$ on our first example, and $h(\mathcal{C}) = 0, h(\mathcal{C}_1) = h(\mathcal{C}_2) = 1, h(\mathcal{C}') = 2$ on our second example. Our

proof is by induction on the maximal height h_{max} . The case $h_{max} = 0$ has been solved in **B.**, so we assume $h_{max} \geq 1$.

The argument goes as follows. Consider a minimal class \mathcal{C} connected downwards to $\mathcal{C}_1, \dots, \mathcal{C}_m$. A reaction \mathcal{R} is *internal* to \mathcal{C} if its reactant and all its products belong to \mathcal{C} ; one then writes $\mathcal{R} : \mathcal{C} \rightarrow \mathcal{C}$. On the other hand, irreversible arrows from \mathcal{C} to \mathcal{C}_i represent split irreversible reactions $\tilde{\mathcal{R}} : A \rightarrow A_i$ with $A \in \mathcal{C}$ and $A_i \in \mathcal{C}_i$, coming from the linearization of a one-to-several reaction $\mathcal{R} : A \rightarrow s_1 A'_1 + \dots + s_n A'_n$. There are two cases:

- (i) (purely external reaction) either $A'_i, i = 1, \dots, n$ all belong to $\uplus_{j=1}^m \mathcal{C}_j$, so that all split reactions $\tilde{\mathcal{R}} : A \rightarrow A'_i$ are external;
- (ii) (mixed reaction) or one of the A'_i belongs to \mathcal{C} , so that $\tilde{\mathcal{R}} : A \rightarrow A'_i$ is an internal irreversible reaction of \mathcal{C} .

The second case is called a mixed case because some of the A'_i belong to \mathcal{C} , and some do not, hence the one-to-several reaction \mathcal{R} is neither internal nor external. Now, if there is no mixed reaction with reactant in \mathcal{C} , we can extract from the set of reactions those which are internal to \mathcal{C} , and build the \mathcal{C} -valued auxiliary Markov chain $(\tilde{X}_{\mathcal{C}}(t))_{t \geq 0}$ as in **A.** with set of transitions associated to those internal reactions. The construction in **A.** and **B.** yields a positive vector $c_{\mathcal{C}} = (c_{\mathcal{R}})_{\mathcal{R} : \mathcal{C} \rightarrow \mathcal{C}}$ such that the associated chemical balance for all species in \mathcal{C} is > 0 .

Considering now the case of a mixed reaction $\mathcal{R} : A \rightarrow s_1 A'_1 + \dots + s_n A'_n$ with $A'_1, \dots, A'_{n'} \in \mathcal{C}, (A'_\ell)_{\ell > n'} \in \uplus_{j=1}^m \mathcal{C}_j$, we split it for our purposes into a truncated internal reaction $\mathcal{R}_{\mathcal{C}} : A \rightarrow s_1 A'_1 + \dots + s_{n'} A'_{n'}$, and $n - n'$ external split reactions $\tilde{\mathcal{R}} : A \rightarrow A'_i, i = n' + 1, \dots, n$. Joining truncated internal reactions $\mathcal{R}_{\mathcal{C}}$ to the set of internal reactions, one proceeds as in the previous paragraph, and obtains a positive vector $c_{\mathcal{C}} = (c(\mathcal{R}))_{\mathcal{R} : \mathcal{C} \rightarrow \mathcal{C}}$, where now $\mathcal{R} : \mathcal{C} \rightarrow \mathcal{C}$ represents the set of all (truncated or not) reactions internal to \mathcal{C} , such that the associated chemical balance for all species in \mathcal{C} is > 0 .

We proceed similarly for all minimal classes.

Consider now a height 1 class \mathcal{C}_1 . Start as in the previous paragraph by constructing a \mathcal{C}_1 -valued auxiliary Markov chain with set of transitions associated to the (truncated or not) reactions internal to \mathcal{C}_1 . Proceed similarly for all classes of height 1. Using the construction in **A.** and coupling with the height 0 class reaction vectors obtained in the previous step, one obtains a reaction vector $c = (c_0, c_1)$ such that $c_0(\mathcal{R}) > 0$, resp. $c_1(\mathcal{R}) > 0$ iff \mathcal{R} is (truncated or not) internal to a height 0, resp. 1 class, and the associated balance is > 0 , resp. ≥ 0 , for species belonging to height 0, resp. height 1 classes.

We now adapt the perturbation argument of **B.** First, if $\mathcal{R} : A \rightarrow \dots, A$ belonging to a minimal class \mathcal{C} , is of mixed type, we redefine $c(\mathcal{R}) = c_{\mathcal{C}}(\mathcal{R}_{\mathcal{C}})$. Choosing a class \mathcal{C}' of

height 1, we now explain how to obtain a strictly positive balance for species in \mathcal{C}' . There are two cases:

- (i) (purely external case) Assume that all reactions $\mathcal{R} : A \rightarrow s'A' + \dots$, such that $A' \in \mathcal{C}'$ and A in a class of height 0, are purely external, so none of these have been taken into account previously in the auxiliary Markov chains. The balance associated to such reactions is strictly negative for the reactant A , and strictly positive for products, including A' . Choosing a small enough coefficient $c(\mathcal{R})$ for them, the net balance for height 0 species remains > 0 , and we get a strictly positive balance for A' .
- (ii) (mixed case) Assume there exists a mixed reaction $\mathcal{R} : A \rightarrow (s_1A'_1 + \dots + s_{n'}A'_{n'}) + A' + \dots$, with $A, A'_1, \dots, A'_{n'}$ in a height 0 class \mathcal{C} , and $A' \in \mathcal{C}'$. This reaction has already been taken into account, by construction $c_{\mathcal{C}}(\mathcal{R}_{\mathcal{C}}) > 0$. Replacing the truncated internal reaction $\mathcal{R}_{\mathcal{C}}$ by \mathcal{R} only increases the balance for external species, including A' .


In both cases, one has obtained a positive balance for at least one species in each height 1 class, which can be considered as a local influx. One may now modify the construction in **B.** by simply using the local influx (instead of the positive balance due to an internal irreversible reaction) to perturb c_1 , and obtains a positive vector c' such that $c'(\mathcal{R}) = c(\mathcal{R})$ if the reactant of \mathcal{R} belongs to a height 0 class, and the balance associated to c' is > 0 for species belonging to classes of height ≤ 1 .

Proceeding by induction on $h \leq h_{max}$ and using reactions connecting classes of height $h - 1$ to classes of height h , we get the result. \square


4 (Top) implies dynamical autocatalysis for dilute networks

We show here the second part of Theorem 3.1, and prove spontaneous autocatalysis (i.e. exponential amplification of some species starting from an arbitrary initial condition with low concentrations).

The following notations are used. Reversible 1-1 reactions (for which some arbitrary orientation is chosen) are denoted



Forward, irreversible split reactions coming from a reaction



are denoted

$$\tilde{\mathcal{R}}_{i,j_\ell}^{for} : A_i \xrightarrow{s_j k_i^+} A_{j_\ell} \quad (38)$$

Combining all these reactions defines (see section 3 **A.**) an auxiliary Markov chain $(\tilde{X}(t))_{t \geq 0}$, whose adjoint generator we denote \tilde{M} . On the other hand, the linearized time-evolution generator of the reaction network containing all reversible 1-1 reactions and forward, irreversible reactions (*excluding* possible degradation reactions) is called M . It is a generalized adjoint Markov generator; we shall use the path representation of resolvents of \tilde{M} and M introduced in Suppl. Info. (§6.5).

Choose a set of degradation rates $(\alpha_i)_{i \in \mathcal{S}} > 0$ – we remind the reader that M itself involves by assumption no degradation reaction. Discrete-time transition rates are

$$w(\alpha)_{i \rightarrow j} := \frac{(M_\alpha)_{ji}}{|(M_\alpha)_{i,i}|} = \frac{M_{ji}}{|M_{i,i}| + \alpha_i} \quad (39)$$

for $M_\alpha := M - \alpha$, and similarly

$$\tilde{w}(\alpha)_{i \rightarrow j} := \frac{(\tilde{M}_\alpha)_{ji}}{|(\tilde{M}_\alpha)_{i,i}|} = \frac{\tilde{M}_{ji}}{|\tilde{M}_{i,i}| + \alpha_i} \quad (40)$$

for $\tilde{M}_\alpha := \tilde{M} - \alpha$.

The general purpose of this section is to prove that a diluted network satisfying the topological hypothesis (Top) of Theorem 3.1 is weakly dynamically autocatalytic, provided it is degradationless, or degradation reactions have small enough rates. Furthermore, we shall be able to prove strong dynamical autocatalysis in some cases, including the irreducible case.

A. Irreducible case. We assume here that the split graph $G(\mathbb{S})$ is irreducible, and prove strong dynamical autocatalysis. Define M as above (or replace M by $M - \beta$, where $(\beta_i)_{i \in \mathcal{S}}$ is a set of small enough degradation rates). For any $\alpha \geq 0$, let $R(\alpha)$ be its resolvent, with coefficients in $[0, +\infty]$ given by the path representation (68); in Suppl. Info., it is proved that positivity of the Lyapunov exponent of M is equivalent to having

$$(R(\alpha))_{i,j} = +\infty \quad (41)$$

for some (or all) $i, j \in \mathcal{S}$ and some $\alpha > 0$. Then this condition implies dynamical autocatalysis for degradation rates $< \alpha$. In turn, Lemma 6.2 and the discussion below give quantitative criteria for spontaneous autocatalysis. So let us prove (41).

By hypothesis, there exists at least one forward irreversible reaction as in (37); reindexing, we assume that $i = 0$ and $j_\ell = \ell, \ell = 1, \dots, n$. Choose a set of degradation rates $(\alpha_i)_{i \in \mathcal{S}} > 0$. The generalized adjoint Markov generator $M - \alpha$ and the adjoint sub-Markov generator $\tilde{M} - \alpha$ have same off-diagonal coefficients, but diagonal coefficients of

M are larger than those of \tilde{M} . Namely (decomposing M into a sum of contributions by individual split reactions, see §6.4), $M(\mathcal{R}) = \tilde{M}(\mathcal{R})$ if \mathcal{R} is reversible, while

$$\sum_{\ell} M(\tilde{\mathcal{R}}_{i,j_{\ell}}^{for})_{i,i} = -k_i^+ > \sum_{\ell} \tilde{M}(\tilde{\mathcal{R}}_{i,j_{\ell}}^{for})_{i,i} = -(s_1 + \dots + s_n)k_i^+ \quad (42)$$

for a forward irreversible reaction. Now

$$\begin{aligned} M_{i,i} &= \sum_{\mathcal{R} \text{ reversible}} M(\mathcal{R})_{i,i} + \sum_{\tilde{\mathcal{R}}^{for} \text{ irreversible}} M(\tilde{\mathcal{R}})_{i,i} \\ &\geq \sum_{\mathcal{R} \text{ reversible}} \tilde{M}(\mathcal{R})_{i,i} + \sum_{\tilde{\mathcal{R}}^{for} \text{ irreversible}} \tilde{M}(\tilde{\mathcal{R}})_{i,i} \end{aligned} \quad (43)$$

The inequality is strict for $i = 0$. It follows: $\tilde{w}(\alpha)_{i \rightarrow j} \leq w(\alpha)_{i \rightarrow j}$, and in particular, $\tilde{w}(\alpha)_{0 \rightarrow j} < w(\alpha)_{0 \rightarrow j}$ if $j = 1, \dots, n$.

Then

$$(R(\alpha))_{0,0} = \frac{1}{|(M_{\alpha})_{0,0}|} \sum_{p \geq 0} (f(\alpha)_{0 \rightarrow 0})^p \quad (44)$$

where $f(\alpha)_{0 \rightarrow 0}$ is the total weight of *excursions* from 0 to 0, computed using transition rates $w(\alpha)$, namely, $f(\alpha)_{0 \rightarrow 0} = \sum_{\ell \geq 1} \sum_{0=x_0 \rightarrow x_1 \rightarrow \dots \rightarrow x_{\ell} \rightarrow 0=x_{\ell+1}} \prod_{k=0}^{\ell} w(\alpha)_{x_k \rightarrow x_{k+1}}$, where the sum is restricted to paths $(x_k)_{1 \leq k \leq \ell}$ of length ≥ 1 in $\mathcal{S} \setminus \{0\}$. Summing over all possible first steps, we get

$$f(\alpha)_{0 \rightarrow 0} = \sum_{i \neq 0} w(\alpha)_{0 \rightarrow i} f(\alpha)_{i \rightarrow 0}, \quad (45)$$

where $f(\alpha)_{i \rightarrow 0}$ is the total weight of paths in $\mathcal{S} \setminus \{0\}$ issued from i , with a final additional step leading back to 0. In turn, using again the path representation, we see that $f(\alpha)_{i \rightarrow 0}$ may be written as an infinite series whose coefficients are products of transition rates $w(\alpha)$.

Similarly, one may define $\tilde{f}(\alpha)_{0 \rightarrow 0} = \sum_{i \neq 0} \tilde{w}(\alpha)_{0 \rightarrow i} \tilde{f}(\alpha)_{i \rightarrow 0}$, where $\tilde{f}(\alpha)_{i \rightarrow 0}$ is the same sum as $f(\alpha)_{i \rightarrow 0}$, but with transition rates $w(\alpha)$ replaced by $\tilde{w}(\alpha)$.

When $\alpha = 0$, $\tilde{f}(0)_{0 \rightarrow 0}$ is simply the probability for the true (i.e. probability-preserving) Markov chain \tilde{X} to get back to 0. Irreducible Markov chains with finite state space are recurrent, so $\tilde{f}(0)_{0 \rightarrow 0} = 1$. Now $w(\alpha) \geq \tilde{w}(\alpha)$ (implying $f(\alpha)_{i \rightarrow 0} \geq \tilde{f}(\alpha)_{i \rightarrow 0}$) and $w(\alpha)_{0 \rightarrow i} > \tilde{w}(\alpha)_{0 \rightarrow i}$, hence (by a simple continuity argument w. r. to α) $f(\alpha)_{0 \rightarrow 0} > 1$ for $\alpha > 0$ small enough, implying

$$(R(\alpha))_{0,0} = +\infty. \quad (46)$$

B. Reducible case. We start with a one-parameter family of examples to show the variety of autocatalytic behaviors (see p. 13). Let $\mathcal{C} = \{A_1, A_2\}$ and $\mathcal{C}' = \{B\}$ be

two classes with probabilistic flow flowing from \mathcal{C} into \mathcal{C}' , and (in the (A_1, A_2, B) -basis)

$$M := \begin{bmatrix} -1 & 2 & 0 \\ 1 & -1 & 0 \\ 0 & 1 & m-1 \end{bmatrix} \text{ corresponding to the reaction network}$$



with $m > 0$. The Lyapunov exponent of $M|_{\mathcal{C}} = \begin{bmatrix} -1 & 2 \\ 1 & -1 \end{bmatrix}$ is $\lambda_{max} := \sqrt{2} - 1$, and

$v_{\mathcal{C}} := \begin{bmatrix} \sqrt{2} \\ 1 \end{bmatrix}$ is an associated positive eigenvector; that of $M|_{\mathcal{C}'} = [m-1]$ is $m-1$. The maximum (Lyapunov) eigenvalue of M is $\max(\lambda_{max}, m-1)$. There are three cases, depending on the spectral parameter m :

1. if $\lambda_{max} > m-1$, then $v_{\mathcal{C}}$ can be extended into a positive Lyapunov eigenvector for M , implying strong dynamical autocatalysis;

2. if $\lambda_{max} = m-1$ (resonant case), then this is not possible (the associated Jordan block is not trivial). Instead, one gets the downstream Lyapunov eigenvector $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.

Thus dynamical autocatalysis holds only in the weak sense;

3. if $\lambda_{max} < m-1$, then $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ is again a downstream Lyapunov eigenvector, and dynamical autocatalysis holds only in the weak sense.

Our proof of weak autocatalysis encompasses all cases without addressing such spectral considerations. It follows from our argument in **A.** through an elementary perturbation argument. Namely, replace the above matrix M by $M(\varepsilon) := M + \varepsilon J$, where $\varepsilon > 0$ is a small parameter, and J is an off-diagonal matrix with non-negative coefficients. If J has enough nonzero coefficients, then $M(\varepsilon)$ will be irreducible. (Assuming all concentrations of all species are > 0 , this may e.g. be achieved by including also some split reverse reactions coming from reverse reactions $s_1 A_{j_1} + s_2 A_{j_2} + \dots + s_n A_{j_n} \xrightarrow{k_i^-} A_i$ connecting classes upwards,



with $k_{j_\ell \rightarrow i}^- = k_i^- s_{j_\ell} [A_{j_\ell}]^{s_{j_\ell}-1} \prod_{\ell' \neq \ell} [A_{j_{\ell'}}]^{s_{\ell'}}$, see §6.4).

Now $M(\varepsilon)$ is an irreducible generalized Markov matrix. The Perron-Frobenius theorem implies that $\lambda_{max}(\varepsilon) := \lambda_{max}(M(\varepsilon))$ has multiplicity 1, and that there exists a unique associated eigenvector $v(\varepsilon) = (v_i(\varepsilon))_{i \in \mathcal{S}}$ such that $v_i(\varepsilon) > 0$ for all i , and $\sum_{i \in \mathcal{S}} v_i(\varepsilon) = 1$.

Following the arguments in **A.**, one sees that the addition of εJ modifies transition rates $w(\alpha)$ only by $\pm O(\varepsilon)$, thus $\lambda_{max}(\varepsilon) \geq \lambda > 0$ for some constant λ uniformly in ε if ε is small enough. Following the compactness argument ([29], Th. 6.10.), one proves the existence of a limiting eigenvector v such that $Mv = \lambda_{max}v$, where v is the limit of some subsequence $(v(\varepsilon_k))_{k=1,2,\dots}$ with $\varepsilon_k \rightarrow 0$, and $\lambda_{max} := \lim_{\varepsilon \rightarrow 0} \lambda_{max}(\varepsilon) \geq \lambda > 0$. By continuity, $\sum_{i \in \mathcal{S}} v_i = 1$ so that v is normalized, but coefficients $v_i, i \in \mathcal{S}$ are only non-negative in general. Thus we obtain weak dynamical autocatalysis.

Let us briefly discuss specific hypotheses (generalizing case 1. above) under which strong dynamical autocatalysis holds. First, the Perron block decomposition (see [29]) of M implies that its spectrum $\Sigma(M)$ is the union of the spectra $(\Sigma(M|_{\mathcal{C}}))_{\mathcal{C}}$ of its restrictions to all classes \mathcal{C} . We assume (i) that there exists a minimal class \mathcal{C} from which all classes can be attained from it following the probabilistic flow (i.e. following arrows downward); (ii) that $\lambda_{max}(M|_{\mathcal{C}}) = \lambda_{max} := \max(\Sigma(M))$ is the maximum of all Lyapunov exponents of all classes, more precisely, $\lambda_{max}(M|_{\mathcal{C}'}) < \lambda_{max}$ for all $\mathcal{C}' \neq \mathcal{C}$. By the above compactness argument, there exists a nonzero eigenvector $v \geq 0$ such that $Mv = \lambda_{max}v$. Restricting to \mathcal{C} , we get a positive Lyapunov eigenvector $v_{\mathcal{C}} = (v_i)_{i \in \mathcal{C}} > 0$ for $M|_{\mathcal{C}}$. Hypothesis (i) then implies that $e^{tM}v > 0$ for all $t > 0$. Now $e^{tM}v = e^{t\lambda_{max}}v$, so that $v > 0$ is positive, and strong autocatalysis is proven. To be concrete (as a final remark), exponential growth (76) will hold for all components for time values $t > \tau$, where $\tau > 0$ is a homogeneization time as in Lemma 6.2 (ii) (see discussion below the Lemma).

5 Perspectives

We have introduced in our main result, Theorem 3.1, a condition (Top) that provides a topological characterization of autocatalysis in the dilute regime (i.e. for low concentrations). This characterization is complete in the limit of negligible degradation rates: indeed, in this case, (Top) is necessary and sufficient for autocatalysis both in the stoichiometric and in the dynamical sense (at least, for irreducible networks). We have furthermore shown that, in this limit, an infinitesimal amount of any species participating in the autocatalytic network ensures the onset of dynamical autocatalysis (see Lemma 6.2). In practice, this means that autocatalytic amplification can start spontaneously upon the rare appearance of a single autocatalyst. Interestingly, these conclusions directly apply to the universal minimal autocatalytic networks (autocatalytic cores) found in any autocatalytic system [8], as they all respect (Top).

A first excursion out of this well-understood regime consists in including significant degradation reactions. Then condition (Top) remains necessary and sufficient for stoichiometric autocatalysis, but is only necessary for dynamical autocatalysis. Determining viability

thresholds, i.e. maximum combinations of degradation rates which allow dynamical autocatalysis, is critical for the design of autocatalytic reaction networks and in origin of life studies [20]. Using branching processes, viability thresholds were determined for autocatalytic cores in the stochastic regime where only a few molecules are present [8]. Specifically, it was shown there that a *single* molecule survives with positive probability if and only if a certain inequality involving kinetic and degradation rates is satisfied. A next step of the treatment presented here will be to characterize viability thresholds allowing positivity of the Lyapunov exponent, and understand the relationship between the continuous and stochastic treatments of the viability thresholds. Notably, a conclusion of the stochastic treatment is that a multiplicity of internal catalytic cycles within the autocatalytic network favors survival (equivalently, allows larger degradation rates). It is tempting to speculate that this conclusion should apply as well to viability thresholds in the kinetic limit, as derived from the study of Lyapunov exponents.

Another direction for generalization is to go beyond the diluted regime. Away from it, Lyapunov exponents characterize stability in the neighborhood of stationary points other than the zero concentration limit, including equilibrium for networks satisfying detailed balance and growth modes for systems with dilution rate ensuring constant total concentration (as in [12]). However, in all generality, there is not necessarily a direct relationship between positivity of the Lyapunov exponent (growth rate) of the *linearized* system and the growth of the original nonlinear dynamical system.

In a companion paper [30], we discuss all these points using an approach based on the analysis of §6.4 and §6.5. We obtain the following tentative conclusions, valid in the non-diluted regime: (i) Topology *and* thermodynamics together inform about autocatalysis; (ii) Estimating the Lyapunov exponent is (despite objections raised in the previous paragraph) a useful 'proxy' allowing quantitative estimates of the growth rate. General quantitative statements include: the computation of certain autocatalytic thresholds in the diluted regime; and estimates for Lyapunov exponents depending essentially on the topology of the network and on thermodynamics *for arbitrary concentrations*. We also show on examples that the curves of 'proxy' dynamical systems based on the above estimates compare well to the curves obtained by numerical integration, over a surprisingly large range of growth regimes.


The approach developed here and in our companion paper is a promising one for the investigation of more complex networks. Indeed, it shows that partial knowledge based on topology and thermodynamics informs on dynamics, independently of the knowledge of reaction rate constants, which is generally missing. A particularly important question is to understand the conditions for the existence of multiple growth modes that could support rudimentary forms of Darwinian evolution [17]. Together with threshold estimates, this may allow us to build scenarios for the emergence of evolution during the origin of life [20].

6 Supplementary information

For simplicity, the reaction networks we consider in this appendix contain only *forward* reactions with *one* reactant,



and the corresponding *reverse* reactions,





All reaction networks discussed in the article are of this type.

This section is organized as follows. We start by presenting Type I and Type III cycles; contrary to [8], we assume stoichiometry 1 for reactants, in conformity with (49). Extended results for these cycles are presented in the companion paper [30]. A short argument for Theorem 3.1 (1) (characterization of stoichiometric autocatalysis) is presented in §6.3 in the case of an irreducible network. Then (§6.4), we write down explicit formulas for the linearized time-evolution generator M of a reaction network. Finally, we present in §6.5 a "theory in a nutshell" for generalized adjoint Markov generators, extending results known from Markov chain theory; see in particular Lemma 6.1 and 6.2.

6.1 Presentation of type I cycles

We consider in this subsection type I cycles of arbitrary length in the Blokhuis-Lacoste-Nghe classification,






The simple model studied in section 2 corresponds to the special case $n = 1$. Chemostatted species – to be thought of e.g. as redox/energy carrier couple as in section 2 – are (A, A') (in red). Dynamical species are (B, B_1, \dots, B_n) . Leaving out chemostatted species, we have a cycle $B \rightleftharpoons B_1 \rightleftharpoons B_2 \rightleftharpoons \dots \rightleftharpoons B_n \rightleftharpoons 2B$.


We have assumed trivial $1 \leftrightarrow 1$ stoichiometry for all reactions along the cycle, except for the *duplication reaction* $(\mathcal{R}_n) : B_n \rightarrow 2B$ closing the cycle. There is nothing special about stoichiometry 2. The extension to the case when $(\mathcal{R}_n) : B_n \rightarrow mB$ with arbitrary $m = 2, 3, \dots$ is straightforward.

Type I cycles are autocatalytic in the stoichiometric sense, as seen by choosing any reaction vector $c = (c_0, \dots, c_n)$ such that $c_0 > c_1 > \dots > c_n > c_0/2 > 0$. On the other hand, choosing $c = (1, \dots, 1)$ yields the coarse-grained duplication reaction for species B



6.2 Presentation of type III cycles





Chemostatted species (in red) are (A, A', A'') . Dynamical species are $(A_i)_{0 \leq i \leq n}, (B'_i)_{0 \leq i \leq n'}, (B''_i)_{0 \leq i \leq n''}$.

We have trivial $1 \leftrightarrow 1$ stoichiometry for all reactions along the two cycles, and $1 \rightarrow 1 + 1$ for the *pitchfork reaction* $A_n \rightarrow B'_0 + B''_0$. Choosing a positive reaction vector such that $c_0 > \dots > c_n, c'_0 > \dots > c'_{n'}, c''_0 > \dots > c''_{n''}$, and $c_n > \max(c'_0, c''_0), c'_{n'} + c''_{n''} > c_0$, one obtains a positive balance for all species. Choosing instead $c = (1, \dots, 1)$ yields the coarse-grained duplication reaction for species A_0



6.3 Autocatalysis from the stoichiometric matrix

We give here a short argument for Theorem 3.1 (1) in the irreducible case.

We consider an irreducible component of the reaction network with n species verifying (Top): every reaction has exactly one reactant and at least one reaction has ≥ 2 products or a product with a stoichiometry strictly ≥ 2 . Correspondingly, each column j of the stoichiometric matrix \mathbb{S} possesses a coefficient $s_{ij} = -1$ and otherwise positive coefficients such that

$$\sum_{i=1}^n s_{ij} \geq 0. \tag{56}$$

Additionally, there is a column index k such that $\sum_{i=1}^n s_{ik} > 0$.

We want to show that there exists a reaction vector $c > 0$ such that $\mathbb{S}c > 0$, i.e. $(\mathbb{S}c)_i > 0$ for all i . For this, it is sufficient to show that $Mc' > 0$ for a certain reaction vector $c' > 0$, where M is a matrix whose columns are positive linear combinations of those of \mathbb{S} , as constructed below.

Let $J(i) = \{j \mid s_{ij} = -1\}$ be the set of reactions having species i as reactant, and N_i the cardinal of $J(i)$. As the network is irreducible, $N_i \geq 1$ for all i . Let $n := |\mathcal{S}|$. Denoting C_M^j , resp. C_S^j the j -th column of M , resp. \mathbb{S} , we let, for $j = 1, \dots, n$:

$$C_M^j := \frac{1}{N_j} \sum_{j' \in J(j)} C_S^{j'}.$$

By construction, $M \equiv A - I$, where A is square and non-negative, i.e. $A_{ij} \geq 0$ for all i, j . Given that the network is strongly connected, A is irreducible. Stoichiometric hypotheses (56) impose $\sum_{i=1}^n A_{ij} \geq 1$ for every j and $\sum_{i=1}^n A_{ik} > 1$ for a certain k . By the Perron-Frobenius theorem, the largest eigenvalue λ of A is positive and associated with an eigenvector $c' > 0$. We have:

$$\lambda \sum_{i=1}^n c'_i = \sum_{i=1}^n \left(\sum_{j=1}^n A_{ij} c'_j \right) = \sum_{j=1}^n \left(\sum_{i=1}^n A_{ij} \right) c'_j > \sum_{j=1}^n c'_j$$

This implies $\lambda > 1$. Hence c' is a positive eigenvector of $M = A - I$ with eigenvalue $\lambda - 1 > 0$.

6.4 Linearized time-evolution generator for reaction networks

The linearized time-evolution generator $M = M([A])$ of a reaction network has been defined in eq. (3)–(6). The current $J_{\mathcal{R}} = k_+[A_i]$ associated to a forward reaction \mathcal{R} as in (49) is straightforwardly linearized to $J_{lin, \mathcal{R}}(A) = k_+ A_i$. Considering now a reverse reaction (50), the reverse current is $k_- \prod_{\ell=1}^n [A_{i'_\ell}]^{s_\ell}$, yielding a linearized current

$$J_{lin} := \sum_{\ell} J_{lin}^{\ell} A_{i'_\ell}, \quad J_{lin}^{\ell} := k_- s_{\ell} \left(\prod_{\ell' \neq \ell} [A_{i'_{\ell'}}]^{s_{\ell'}} \right) [A_{i'_\ell}]^{s_{\ell}-1}. \quad (57)$$

The coefficients of the matrix M are obtained by summing individual matrices $M(\mathcal{R})$ associated to linearized forward reactions $\mathcal{R} : A_i \xrightarrow{k_+} s_1 A_{i'_1} + \dots + s_n A_{i'_n}$,

$$\frac{dA_i}{dt} = -k_+ A_i; \quad \frac{dA_{i'_\ell}}{dt} = s_{\ell} k_+ A_i, \quad \ell = 1, \dots, n \quad (58)$$

and matrices $\sum_{\ell} M(\bar{\mathcal{R}}, \ell)$ associated to linearized reverse reactions $\bar{\mathcal{R}}$ (see (57))

$$\frac{dA_i}{dt} = J_{lin}^{\ell} A_{i'_{\ell}}; \quad \frac{dA_{i'_{\ell}}}{dt} = -s_{\ell'} J_{lin}^{\ell} A_{i'_{\ell}}, \quad \ell' = 1, \dots, n. \quad (59)$$

If \mathcal{R} is a forward reaction, the corresponding contribution $M(\mathcal{R})$ to M is (see below (6)) a generalized adjoint Markov generator with *negative* killing rate $a_i(\mathcal{R}) = k_+(1 - \sum_{\ell} s_{\ell})$, which vanishes precisely in the case of a reversible reaction $A_i \rightleftharpoons A_j$.

Consider now a *reverse* reaction $\bar{\mathcal{R}}$. The matrix $M(\bar{\mathcal{R}}, \ell)$ is *not* a generalized adjoint Markov generator if $n \geq 2$, because of the *probability leak currents* $-s_{\ell'} J_{lin}^{\ell} A_{i'_{\ell}} = (M(\bar{\mathcal{R}}, \ell))_{i'_{\ell'}, i'_{\ell}}$ from state $i'_{\ell} \neq i'_{\ell'}$; also, it features ≥ 0 killing rates $a_{i'_{\ell}}(\bar{\mathcal{R}}, \ell) = J_{lin}^{\ell} [s_{\ell} - 1]$, computed without considering probability leak currents, considered as external non-diagonal terms without probabilistic interpretation. The reverse reaction $\bar{\mathcal{R}} : B''_0 + B'_0 \xrightarrow{\nu_-} A_n$ in type III cores has $n = 2$, and does exhibit leak currents. Matrices $M(\bar{\mathcal{R}}, 0')$, resp. $M(\bar{\mathcal{R}}, 0'')$, are identified with the two columns of the matrix

$$M(\bar{\mathcal{R}}) := M(\bar{\mathcal{R}}, 0') + M(\bar{\mathcal{R}}, 0'') = \begin{bmatrix} 0 & \nu_-[B''_0] & \nu_-[B'_0] \\ 0 & -\nu_-[B''_0] & -\nu_-[B'_0] \\ 0 & -\nu_-[B''_0] & -\nu_-[B'_0] \end{bmatrix}$$

Reverse reactions putting into contact $n \geq 2$ different species produce negative off-diagonal coefficients, here emphasized in red. If the resulting matrix $M = \sum_{\mathcal{R}} M(\mathcal{R}) + \sum_{\bar{\mathcal{R}}, \ell} M(\bar{\mathcal{R}}, \ell)$ has negative off-diagonal coefficients, it cannot be interpreted as a generalized Markov generator, therefore Lemma 6.1 below (allowing easy upper bounds for the Lyapunov exponent) does not hold.

6.5 Generalized Markov generators

A central notion in this article is that of *generalized Markov processes*, i.e. discrete- or continuous-time Markov processes which are not necessarily probability-preserving; a general introductory reference is ([26], chap. 2). Let $\mathcal{S} = \{1, \dots, n\}$ be a finite state space. Then an $n \times n$ matrix M is a *generalized adjoint Markov generator* if


- (i) diagonal coefficients $M_{i,i}$, $1 \leq i \leq n$ are < 0 ;
- (ii) and off-diagonal coefficients $M_{i,j}$, $i \neq j$ are ≥ 0 .

If $\sum_{i=1}^n M_{i,j} = 0$ for all j , then coefficients of the transposed matrix M^t sum up to zero on each line, so that M^t is a conventional probability-preserving Markov generator: $\mathbf{1}^t = (1 \ \dots \ 1)$ is a left-eigenvector of M with eigenvalue 0. The probability measure


$\mu(t)$ of the process at time t is $e^{tM}\mu(0)$, solution of the master equation $\frac{d}{dt}\mu = M\mu$; probability preservation means that $\sum_i \mu_i(t) = 1$ for all time. Assume, more generally, that $a_j := |M_{j,j}| - \sum_{i \neq j} M_{i,j} \geq 0$, then the total probability $\sum_i \mu_i(t)$ is a decreasing function of time, and a_j can be interpreted as a killing rate (biologically, a degradation rate) in state j . For a finite set of states, there is no obstacle in considering the case when killing rates a_j can have either sign. We discuss the associated random process $(X(t))_{t \geq 0}$ later on; by definition $X(t) \in \mathcal{S}$ has transition rate $M_{i,j}$ from j to i . By construction, $\mu_i(t) \equiv \mathbb{P}[X(t) = i] = (e^{tM}\mu(0))_i$, generalizing the above master equation, where \mathbb{P} is a (non-normalized) measure on trajectories. Probabilistic tools give an intuitive access to the resolvent in terms of trajectories of the Markov process, from which we derive a characterization and properties of the Lyapunov exponent.


Communicating classes, irreducibility. Example of the ” $A_1A_2A_3 \rightarrow B_1B_2B_3$ ” autocatalytic reaction network. Let M be a generalized adjoint Markov generator on $\mathcal{S} = \{1, \dots, n\}$. The matrix M defines a graph $G(M)$ with vertex state \mathcal{S} and oriented edge set $\mathcal{E}(M)$: a pair $e = (x \rightarrow y)$, $x \neq y$ is an edge if $M_{y,x} > 0$; the probability flow follows edges. Following standard terminology in Markov chains, we say that $x \neq y$ communicate (which we denote $x \sim y$) if there exists a path from x to y and a path from y to x , namely, a chain of edges $(x \rightarrow x_1), (x_1 \rightarrow x_2), \dots, (x_n \rightarrow y)$ and a chain of edges $(y \rightarrow \tilde{x}_1), (\tilde{x}_1 \rightarrow \tilde{x}_2), \dots, (\tilde{x}_{n'} \rightarrow x)$ with $n, n' \geq 0$. Letting also $x \sim x$ for all x , this defines equivalence classes called *communicating classes*. M is said to be *irreducible* if there is only one class.

M is clearly reducible if the graph is not connected, but this means that we are dealing with several independent systems, an uninteresting situation. We may assume instead that the graph $G(M)$ is always connected. On the other hand, there exist connected graphs which are not irreducible, for instance the graph of the ” $A_1A_2A_3 \rightarrow B_1B_2B_3$ ” autocatalytic reaction network (also discussed in section 3), a graph on the set $\mathcal{S} = \{1, 2, 3, 1', 2', 3'\}$,




The associated reaction network is






(with arbitrary transition rates) whose graph coincides in the zero-concentration limit with $G_{(123) \rightarrow (1'2'3')}$ through the state identification $(A_1, A_2, A_3, B_1, B_2, B_3) \leftrightarrow (1, 2, 3, 1', 2', 3')$. Adding all the forward reactions (\rightarrow) with coefficients $c_{\mathcal{R}_1} = 5, c_{\mathcal{R}_2} = 4, c_{\mathcal{R}_3} = 3; c_{\mathcal{R}_{1,1'}} = \frac{1}{2}; c_{\mathcal{R}_{1'}} = \frac{5}{6}, c_{\mathcal{R}_{2'}} = \frac{4}{6}, c_{\mathcal{R}_{3'}} = \frac{3}{6}$ yields the result



Thus this network is autocatalytic in the stoichiometric sense.

Partial order, minimal classes, maximal classes. Generalizing the above example, one sees that, by shrinking communicating classes to single points, one reduces the oriented graph $G(M)$ to an oriented "contracted" graph $\mathcal{T}(M)$ which has no loops. (Mind that the associated *unoriented* graph may have loops, so that $\mathcal{T}(M)$ is not necessarily a topological tree.) It is possible to represent this graph with edge arrows going downwards, e.g. in the above example,



with $\mathcal{C} = \{1, 2, 3\}, \mathcal{C}' = \{1', 2', 3'\}$. Note that the graph would become irreducible if (as discussed above) one added the reverse arrow $1' \rightarrow 1$ corresponding to the reverse reaction $2B_1 \rightarrow A_1$. We get a partial order on the set of classes by letting $\mathcal{C}' \succ \mathcal{C}$ if there is a \mathcal{T} -path downstream (i.e. following the probability flow) from \mathcal{C} to \mathcal{C}' . Maximal (downstream) elements (here \mathcal{C}') are called *closed classes*, because they have no outgoing arrows: one cannot escape from them. Minimal (upstream) classes (here \mathcal{C}), on the other hand, have no ingoing arrows.

Autocatalysis in the stoichiometric sense. Let us now discuss the connection between reaction networks and graphs. Consider a reaction network with species set $\mathcal{S} = \{A_1, \dots, A_{|\mathcal{S}|}\}$, forward reaction set $\{1, \dots, N\}$ and positive, mass-action reaction rates for both forward and reverse reactions. We are particularly interested in the limit of small concentrations, so we distinguish:

- (i) reversible reactions $A_i \rightleftharpoons A_j$ ($i \neq j$);

(ii) and irreversible reactions $A_i \xrightleftharpoons[k_-]{k_+} s_1 A_{i'_1} + \dots + s_n A_{i'_n}$, with $n \geq 1$, $s_i \in \mathbb{N}^*$, $\sum_{i=1}^n s_i > 1$.

Let us construct the graph associated to the linearized time-evolution generator M in the zero-concentration limit; note that the graph actually depends only on the stoichiometry matrix \mathbb{S} , not on the rates, so we can call it $G(\mathbb{S})$. As discussed in §3, in case of multiple arrow $i \rightarrow j$, we keep only one.

(i) Reversible reactions $A_i \rightleftharpoons A_j$ contribute to $G(\mathbb{S})$ two arrows $i \rightarrow j$ and $j \rightarrow i$.

(ii) Forward reactions $\mathcal{R} : A_i \xrightarrow{k_+} s_1 A_{i'_1} + \dots + s_n A_{i'_n}$ ($\sum_i s_i > 1$) contribute to $G(\mathbb{S})$ an arrow $i \rightarrow i'_\ell$ for each $\ell = 1, \dots, n$. Reversible reactions $A_i \rightleftharpoons A_j$ decompose into two forward reactions $A_i \rightarrow A_j$ and $A_j \rightarrow A_i$.

On the other hand, reverse reactions $\bar{\mathcal{R}} : s_1 A_{i'_1} + \dots + s_n A_{i'_n} \xrightarrow{k_-} A_i$, with $\sum_i s_i > 1$ contribute no arrow.

Case of the "A₁A₂A₃ → B₁B₂B₃" autocatalytic kinetic reaction network.

The linearized evolution generator is a sum of 7 matrices, one per reaction (provided paired generators associated to forward/reverse reversible reactions $i \rightleftharpoons j$, $i' \rightleftharpoons j'$ are summed together), $M = \sum_i M(\mathcal{R}_i) + M(\mathcal{R}_{11'}) + \sum_{i'} M(\mathcal{R}_{i'})$, with

$$M(\mathcal{R}_1) = \begin{array}{c} \begin{array}{ccc} 1 & 2 & 3 \end{array} \\ \left[\begin{array}{ccc|ccc} -k_{12}^+ & k_{12}^- & & & & \\ k_{12}^+ & -k_{12}^- & & 0 & & \\ \hline & & & & & \\ & 0 & & 0 & & \end{array} \right] \end{array}$$

and similarly for the four other reversible generators $A_2 \rightleftharpoons A_3$, $B_i \rightleftharpoons B_j$; these are probability preserving adjoint Markov generators since the sum of coefficients on any column is zero;

$$M(\mathcal{R}_3) = \begin{array}{c} \begin{array}{ccc|ccc} & 1 & 2 & 3 & 1' & 2' & 3' \\ \hline & & & 2k_{31}^+ & & & \\ & & & -k_{31}^+ & & 0 & \\ \hline & & 0 & & & 0 & \end{array} \end{array}$$

which is an adjoint Markov generator with negative killing rate; and

$$M(\mathcal{R}_{11'}) + M(\bar{\mathcal{R}}_{11'}) = \begin{array}{c} \begin{array}{ccc|ccc} & 1 & 2 & 3 & 1' & 2' & 3' \\ \hline & & & & -k_{11'}^+ & & 2[B_1]k_{11'}^- \\ & & & & 2k_{11'}^+ & & -4[B_1]k_{11'}^- \\ \hline & & & & & & \end{array} \end{array}$$

for the irreversible reaction $\mathcal{R}_{11'}$ coupling \mathcal{C}_1 to $\mathcal{C}_{1'}$, to which one has added the (red) reverse reaction, with rate proportional to the low concentration $[B_1]$, absent in the zero-concentration limit.

Path measure for generalized Markov generators (see [26], chap. 2). When killing rates vanish, one has a probability law \mathbb{P} on trajectories $(X(t))_{0 \leq t \leq T}$: letting $t_1 = 0$, $(T_k)_{k \geq 2} \leq T$ be the jumping times,

$$\begin{aligned} & \mathbb{P}[T_k = t_k + dt_k, X(T_k) = x_k, 2 \leq k \leq \ell \mid X(t_1) = x_1] \\ &= \left[\prod_{k=1}^{\ell-1} \left(e^{(t_{k+1}-t_k)M_{x_k, x_k}} dt_{k+1} \times M_{x_{k+1}, x_k} \right) \right] \times e^{(T-t_\ell)M_{x_\ell, x_\ell}} \end{aligned} \quad (64)$$

Integrating over the jumping times, one obtains the law of the trajectories $\underline{X} = (\underline{X}_k)_{k \geq 0}$ of the underlying "skeleton" discrete-time Markov chain,

$$\mathbb{P}[\underline{X}_k = x_k, 2 \leq k \leq \ell \mid \underline{X}_1 = x_1] = \prod_{k=1}^{\ell-1} w_{x_k \rightarrow x_{k+1}} \quad (65)$$

with transition rates

$$w_{i \rightarrow j} := \frac{M_{ji}}{|M_{i,i}|}, \quad i \neq j \quad (66)$$

featuring the Markov generator M^t . We generalize to arbitrary killing rates and use (64), (65) as a definition for an unnormalized measure \mathbb{P} over trajectories.

A path representation of the resolvent. When killing rates vanish, $\sum_{j \neq i} w_{i \rightarrow j} = 1$, and $w_{i \rightarrow j}$ are simply the transition rates of the underlying skeleton discrete-time Markov chain; by extension, the coefficients $w_{i \rightarrow j}$ will be called *transition rates* in the general case. In chemical terms, $w_{i \rightarrow j}$ measures the *specificity* of the reaction $i \rightarrow j$. Let $\alpha := \text{diag}((\alpha_i)_{1 \leq i \leq |S|})$ a positive diagonal matrix, and $M_\alpha := M - \alpha$. Then

$$(R(\alpha))_{i,j} := \int_0^{+\infty} dt (e^{tM_\alpha})_{i,j} \in [0, +\infty] \quad (67)$$

defines a matrix with positive coefficients, which can be computed as a sum over backward paths $i = x_1 \rightarrow x_2 \rightarrow \dots \rightarrow x_{\ell-1} \rightarrow x_\ell = j$ of arbitrary length $\ell \geq 0$,

$$(R(\alpha))_{i,j} = \sum_{\ell \geq 0} \sum_{x_2, \dots, x_{\ell-1} \in \mathcal{S}} \left(\prod_{k=1}^{\ell-1} w(\alpha)_{x_{k+1} \rightarrow x_k} \right) \times \frac{1}{|M_{j,j}| + \alpha_j}. \quad (68)$$

where

$$w(\alpha)_{x_{k+1} \rightarrow x_k} := \frac{M_{x_k, x_{k+1}}}{|M_{x_k, x_k}| + \alpha_{x_k}}. \quad (69)$$

When finite, $(R(\alpha))_{i,j} < \infty$ are the coefficients of the resolvent $(\alpha - M)^{-1} = (-M_\alpha)^{-1}$; see e.g. [28], chap. III, or [26], §4.2 for an introduction in connection to potential theory.

Proof. The $\ell = 0$ contribution is non-zero only if $i = j$, in which case it corresponds to the integral $\int_0^{+\infty} dt e^{t(M_\alpha)_{j,j}} = \frac{1}{|M_{j,j}| + \alpha_j}$. Splitting M_α into $(M_\alpha)_{diag} + (M_\alpha)_{off}$, where $(M_\alpha)_{diag}$, resp. $(M_\alpha)_{off} = M_{off}$, is its diagonal part, resp. its off-diagonal (jump) part, and expanding the exponential $(e^{tM_\alpha})_{i,j}$ using the Feynman-Kac (or Trotter product) formula, one obtains a sum over trajectories $(x(t'))_{0 \leq t' \leq t}$ such that $x|_{[t_k, t_{k+1})} = x_k$, $k = 1, \dots, \ell - 1$, and $x|_{[t_\ell, t]} = j$, with $0 = t_1 < t_2 < \dots < t_\ell < t$. Integrating over t , one obtains multiple integrals

$$\begin{aligned} & \left(\int_0^{+\infty} dt_2 e^{t_2(M_\alpha)_{x_1, x_1}} M_{x_1, x_2} \right) \left(\int_{t_2}^{+\infty} dt_3 e^{(t_3 - t_2)(M_\alpha)_{x_2, x_2}} M_{x_2, x_3} \right) \dots \\ & \left(\int_{t_{\ell-1}}^{+\infty} dt_\ell e^{(t_\ell - t_{\ell-1})(M_\alpha)_{x_{\ell-1}, x_{\ell-1}}} M_{x_{\ell-1}, x_\ell} \right) \times \int_{t_\ell}^{+\infty} dt e^{(t - t_\ell)(M_\alpha)_{x_\ell, x_\ell}}, \end{aligned} \quad (70)$$

yielding (68).

Lemma 6.1 (Properties of the resolvent for M irreducible) We assume that M is irreducible. Then:

- (i) Coefficient functions $\alpha \mapsto (R(\alpha))_{i,j}$ are decreasing, namely, if $\alpha \leq \alpha'$, i.e. $\alpha_i \leq \alpha'_i$ for all i , then $R(\alpha') \leq R(\alpha)$;
- (ii) let $t \mapsto \alpha(t)$ ($t \geq 0$) be an increasing function of time, i.e. $(t' \leq t) \Rightarrow (\alpha(t') \leq \alpha(t))$; then there exists a transition time $t = t_0$ (possibly, $t_0 = 0$) such that all coefficients of $R(\alpha(t))$ are $< \infty$ if $t > t_0$, and all coefficients of $R(\alpha(t))$ are ∞ if $0 \leq t < t_0$. If $t > t_0$, then $R(\alpha(t)) = (-M_{\alpha(t)})^{-1}$.
- (iii) (Lyapunov exponent) Let $\lambda_{max} := \max\{\operatorname{Re}(\lambda) \mid \lambda \text{ eigenvalue of } M\}$ be the Lyapunov exponent of M . If one lets $\alpha(t) := t\operatorname{Id}$, and the transition time t_0 is > 0 , then $\lambda_{max} = t_0$. Conversely, if $t_0 = 0$, then $\lambda_{max} \leq 0$.
- (iv) (positivity criterion for Lyapunov exponent) assume $R(\alpha) = +\infty$ for some $\alpha \geq 0$ which is not identically zero, then $\lambda_{max} > 0$.

Proof. For (ii) we need only remark that $(\exists i, j, (R(\alpha))_{i,j} = +\infty) \Rightarrow (\forall i, j, (R(\alpha))_{i,j} = +\infty)$. Namely, let i', j' be indices; M being irreducible, there exists a backward path from i' to i , and a backward path from j to j' ; sandwiching $(R(\alpha))_{i,j}$ – which is the sum of the weights of all backward paths from i to j – between them, one gets $(R(\alpha))_{i',j'} = +\infty$.

Let now $\alpha(t) = t\operatorname{Id}$. If $t > \max(0, \lambda_{max})$, then (as can be proved by standard arguments using e.g. Jordan's form for M) there exists some constant $c > 0$ s.t. for all $\tau > 0$, $\|e^{\tau M_{\alpha(t)}}\| = O(e^{-c\tau})$ ($\|\cdot\|$ being any norm), hence $R(\alpha(t))_{ij} < \infty$ for all i, j . Conversely, if $R(\alpha(t))_{ij} < \infty$ for all i, j , then $\lambda - M$ is invertible if $\operatorname{Re} \lambda \geq \alpha(t)$, as follows from the path representation (68). This implies (iii).

Discussing finally (iv), assume that $R(\alpha) = +\infty$ with $\alpha_i > 0$. Let $\mathcal{W}_i(\alpha)$ be the weight of *excursions* from i , i.e. the total weight of all backward paths $i = x_1 \rightarrow x_2 \rightarrow \dots \rightarrow x_{\ell-1} \rightarrow x_{\ell} = i$ such that $x_2, \dots, x_{\ell-1} \neq i$. Then

$$(R(\alpha))_{i,i} = \left(\sum_{n=0}^{+\infty} (\mathcal{W}_i(\alpha))^n \right) \times \frac{1}{|M_{i,i}| + \alpha_i} \quad (71)$$

hence

$$(R(\alpha) = +\infty) \Rightarrow (\mathcal{W}_i(\alpha) \geq 1). \quad (72)$$

The function $\alpha \mapsto \mathcal{W}_i(\alpha)$ is a strictly decreasing function, so $\mathcal{W}_i(\alpha/2) > 1$. This strict inequality remains true (by continuity) in a neighborhood Ω of $\alpha/2$ in \mathbb{R}_+^S , and (by monotony) for all α' such that $\alpha' \leq \alpha''$ for some $\alpha'' \in \Omega$. In particular, for t small enough, $R(t\operatorname{Id}) = +\infty$, so that $\lambda_{max} > 0$. \square

To finish with, we study in some details the onset of the exponential growth using the Perron-Frobenius theorem.

Lemma 6.2 (i) *If λ is an eigenvalue of M and $\text{Re } \lambda = \lambda_{max}$, then $\lambda = \lambda_{max}$. In particular, λ_{max} is an eigenvalue of M . Furthermore, the multiplicity of λ_{max} is 1, and there exists an associated eigenvector with > 0 coefficients.*

(ii) *Let $\tau > 0$. There exist two constants $c = c(\tau), C = C(\tau) > 0$ such that, for every nonzero initial concentration vector v with ≥ 0 coefficients, and for every $t > \tau$,*

$$c \left(\max_i v_i \right) e^{\lambda_{max} t} \leq \left(e^{tM} v \right)_i \leq C \left(\max_i v_i \right) e^{\lambda_{max} t}. \quad (73)$$

The upper bound (73) holds uniformly in τ , but the lower bound degenerates as $\tau \rightarrow 0$ (because $(e^{tM} v)_i \rightarrow_{t \rightarrow 0} v_i$ can vanish). The *homogenization time* τ is discussed below.

Proof.

- (i) This is a consequence of the Perron-Frobenius theorem, see e.g. [29], since (for $C > 0$ large enough constant) $M + C\text{Id}$ has positive coefficients and is irreducible.
- (ii) Fix the eigenvector v_{max} associated to the maximal eigenvalue λ_{max} by requiring that $v_{max,i} > 0$ for all i and $\|v_{max}\|_\infty := \max_i v_{max,i} = 1$. The upper bound follows by standard computations from splitting v into $v_{//} + w$, where $v_{//}$ is the linear projection of v onto the one-dimensional eigenspace $\mathbb{R}v_{max}$ parallel to the sum of all other generalized eigenspaces. For the lower bound, we note that $(e^{\tau M})_{ij} > 0$ for all indices i, j and $\tau > 0$. Fix some (small) instant $\tau > 0$; there exists then $c > 0$ such that that

$$(e^{\tau M} v)_i \geq c \|v\|_\infty v_{max,i}. \quad (74)$$

Let $t \geq \tau$. Since the matrix $e^{(t-\tau)M}$ has positive coefficient, we get $(e^{tM} v)_i \geq c \|v\|_\infty \left(e^{(t-\tau)M} v_{max} \right)_i = c \|v\|_\infty e^{(t-\tau)\lambda_{max}} v_{max,i}$.

□

For applications, we are mostly interested in the onset of the exponential growth regime, and may assume that $\lambda_{max} > 0$. Let $M := M([A] = 0)$ be the generalized Markov generator obtained by linearizing the kinetic equations at zero concentrations. By definition, $\frac{d[A]}{dt} = M[A] + O([A]^2)$. Hence it follows from the above Corollary that, for all $i \in \mathcal{S}$ and $t > \tau$,

$$\frac{[A_i](t)}{\max_j ([A_j](t = 0))} \approx e^{\lambda_{max} t} \quad (75)$$

for time values t such that $\max_i [A_i](t = 0) \times e^{\lambda_{max} t}$ is small enough (depending on kinetic rates), where $a \approx b$ ($a, b > 0$) means: $ca < b < Ca$ for constants $c, C > 0$ independent of t, i and $[A](t = 0)$, but depending on the homogenization time τ . Eq. (75) may be regarded as a mathematical expression for *spontaneous autocatalysis*. The homogenization time τ should be chosen as small as possible in order for (74) to hold for a not too small constant c , with $M = M([A] = 0)$.

References

- [1] Borwein J., Adrian S. (2006). *Convex analysis and nonlinear optimization*, Springer.
- [2] Preiner M., Xavier J. C., Vieira A. D. N., Kleinermanns K., Allen J. F., Martin W. F. (2019). *Catalysts, autocatalysis and the origin of metabolism*, Interface Focus **9** (6), 20190072.
- [3] Arsène S., Ameta S., Lehman N., Griffiths A. D. and Nghe P. (2018). *Coupled catabolism and anabolism in autocatalytic RNA sets*, Nucleic Acids Research **46** (18), 9660-9666.
- [4] Ameta S., Arsène S., Foulon S., Saudemont B., Clifton B. E., Griffiths A. D. and Nghe P. (2021). *Darwinian properties and their trade-offs in autocatalytic RNA networks*, Nat. Commun. **12**, 842.
- [5] Andersen J. L., Flamm C., Merkle D., Stadler P. F. (2019). *Chemical Transformation Motifs – Modelling Pathways as Integer Hyperflows*, EEE/ACM Transactions on Computational Biology and Bioinformatics **16** (2), 510-523.
- [6] Andersen J. L., Flamm C., Merkle D., Stadler P. F. (2020). *Defining autocatalysis in chemical reaction networks*, J. Syst. Chem. **8**, 121-133.
- [7] Vincent L., Berg M., Krismer M., Saghafi S. T., Cosby J., Sankari T., Vetsigian H. K., Cleaves H. J., Baum D. A. (2019). *Chemical ecosystem selection on mineral surfaces reveals long-term dynamics consistent with the spontaneous emergence of mutual catalysis*, Life **9**(4), 80.
- [8] Blokhuis A., Lacoste D., Nghe P. (2020). *Universal motifs and the diversity of autocatalytic systems*, PNAS **117** (41), 25230-25236.
- [9] Fontana W., Buss L. W. (1994). *"The arrival of the fittest": Toward a theory of biological organization*, Bulletin of Mathematical Biology **56**(1), 1-64.
- [10] Danger G., d'Hendecourt V., Pascal R. (2020). *On the conditions for mimicking natural selection in chemical systems*, Nature Reviews Chemistry **4**(2), 102-109.
- [11] Eigen M. (1971). *Selforganization of matter and the evolution of biological macromolecules*, Naturwissenschaften **58**, 465.
- [12] Eigen M., Schuster P. (1979). *The hypercycle. A principle of natural self-organization*, Springer.
- [13] Eigen M., McCaskill J., Schuster P. (1989). *The molecular quasi-species*, Adv. Chem. Phys. **75**, 149.

- [14] Sarkar S. and England. J. L. (2019) *Design of conditions for self-replication*, Phys Rev E **100**, 022414 .
- [15] Esposito M. (2016). *Nonequilibrium Thermodynamics of Chemical Reaction Networks: Wisdom from Stochastic Thermodynamics*, Phys. Rev. X **6**, 041064.
- [16] Penocchio E., Rao R. , Esposito M. (2019). *Thermodynamic efficiency in dissipative chemistry*, Nat. Commun. 0:3865.
- [17] Vasas V. , Fernando C. , Santos M. , Kauffman S., Szathmary E. (2011). *Evolution before genes*, Biology direct **7(1)** 1-14.
- [18] Hanopolskyi A. I., Smaliak V. A., Novichko A. I.v, Semenov S. N. (2021). *Autocatalysis: Kinetics, Mechanisms and Design*, ChemSystemsChem, 3(1), e2000026.
- [19] Hordijk W., Steel M., Kauffman S. (2012). *The structure of autocatalytic sets: Evolvability, enablement, and emergence*, Acta biotheoretica, **60(4)**, 379-392.
- [20] C. Jeancolas, C. Malaterre, P. Nghe (2020). *Thresholds in origin of life scenarios*, Iscience **23(11)**, 101756.
- [21] Kauffman S. A. (1986). *Autocatalytic sets of proteins. Journal of theoretical biology*, **119(1)**, 1-24.
- [22] King G. A. M. (1982). *Recycling, reproduction, and life’s origins*, BioSystems **15**, 89–97.
- [23] Kun A., Papp B., Szathmary E. (2008). *Computational identification of obligatorily autocatalytic replicators embedded in metabolic networks*, Genome biology, **9(3)**, 1-11.
- [24] Kussell E., Leibler S. (2005). *Phenotypic Diversity, Population Growth, and Information in Fluctuating Environments*, Science **309**.
- [25] Nghe P. (2015). *Prebiotic network evolution: six key parameters*, Molecular BioSystems **11-12**, 3206-3217.
- [26] Norris J. R. (1997) *Markov chains*, Cambridge University Press.
- [27] Z. Peng, A. M. Plum, P. Gagrani, D. A. Baum (2020). *An ecological framework for the analysis of prebiotic chemical reaction networks*, J. Theor. Biol. **507**, 110451.
- [28] Revuz D., Yor M. (1999). *Continuous martingales and Brownian motion*, Springer.
- [29] Stewart G. W., Sun J.-G. (1990). *Matrix perturbation theory*, Academic Press.
- [30] Unterberger J. (2021). *Dynamical autocatalysis for autocatalytic cores*, in preparation.