

An Explanation of Mellin's 1921 Paper

Wayne M. Lawton¹

¹Department of the Theory of Functions, Institute of Mathematics and Computer Science,
Siberian Federal University, Krasnoyarsk, Russian Federation.
wlawton50@gmail.com

Abstract

In 1921 Mellin published a Comptes Rendu paper computing the principal root of the polynomial $Z^n + x_1 Z^{n_1} + \dots + x_p Z^{n_p} - 1$ using hypergeometric functions of its coefficients x_1, \dots, x_p . He used an integral transform nowadays bearing his name. Slightly over three pages, the paper is written in French in a terse style befitting the language. Unable to find an elementary explanation on the web or in a textbook, we wrote this expository article to make Mellin's landmark result accessible to interested people who are not experts in hypergeometric functions and complex analysis.

2020 Mathematics Subject Classification:32-03;12-08;33C70 ¹

1 Principal Solution Z and Mellin Transform of Z^α

In the opening paragraph of his paper, Mellin says that it summarizes research he undertook years ago and was prompted by notes [2] of Richard Birkeland, a Norwegian mathematician known for his contributions to the theory of algebraic equations. The 7 numbered equations in Mellin's paper and this paper coincide. We prove each of them.

$$Z^n + x_1 Z^{n_1} + \dots + x_p Z^{n_p} - 1 = 0 \quad (1)$$

For integers $p \geq 1$, $0 < n_p < \dots < n_1 < n$, we define a principal solution to be an analytic function $Z(x_1, \dots, x_p)$ on $[0, \infty)^p$ satisfying (1) and $Z(0, \dots, 0) = 1$.

Lemma 1 *If a principal solution $Z(x_1, \dots, x_p)$ of (1) exists, it is unique.*

Proof For $r > 0$ define $D_r := \{z \in \mathbb{C} : |z| < r\}$. For sufficiently small r a principal solution $Z(x_1, \dots, x_p)$ of (1) extends to give a holomorphic solution $\tilde{Z}(x_1, \dots, x_p)$ of (1) for (x_1, \dots, x_p) in the polydisc D_r^p . Therefore, for every n -th root of unity ϵ , the function $Z_\epsilon(x_1, \dots, x_p) := \epsilon \tilde{Z}(\epsilon^{n_1} x_1, \dots, \epsilon^{n_p} x_p)$ is a holomorphic solution of (1) in D_r^p . Since a polynomial of degree n can have at most n distinct roots, these n distinct functions describe all holomorphic solutions of (1) on D_r^p . The conclusion follows since the restrictions of $Z(x, \dots, x_p)$ and $Z_1(x_1, \dots, x_p)$ to $[0, r)^p$ are equal and analytic. Define $\Xi := \{\xi := (\xi_1, \dots, \xi_p) : \xi_1 + \dots + \xi_p \in (-\infty, -1]\}$, $W(\xi) := 1 + \xi_1 + \dots + \xi_p$, and the holomorphic function $\Psi := (x_1, \dots, x_p) : \mathbb{C}^p \setminus \Xi \rightarrow \mathbb{C}^p$ by

$$x_i(\xi) := \xi_i W(\xi_1, \dots, \xi_p)^{\frac{n_i}{n} - 1}, \quad i = 1, \dots, p \quad (2)$$

where $W^{\frac{1}{n}} : \mathbb{C}^p \setminus \Xi \rightarrow \mathbb{C}$ is the branch of $\sqrt[n]{W}$ that satisfies $W^{\frac{1}{n}}(0, \dots, 0) = 1$.

Lemma 2 *If x_1, \dots, x_p are defined by (2), then $Z := W^{-\frac{1}{n}}$ satisfies (1).*

Proof Follows by substitution.

¹This work is supported by the Krasnoyarsk Mathematical Center and financed by the Ministry of Science and Higher Education of the Russian Federation in the framework of the establishment and development of Regional Centers for Mathematics Research and Education (Agreement No. 075-02-2020-1534/1).

Lemma 3 *The restriction $\Psi : [0, \infty)^p \rightarrow [0, \infty)^p$ is a bijection. Furthermore, the Jacobian of $\Psi : \mathbb{C}^p \setminus \Xi \rightarrow \mathbb{C}^p$ satisfies*

$$\frac{\partial(x_1, \dots, x_p)}{\partial(\xi_1, \dots, \xi_p)} = \left(1 + \sum_{k=1}^p \xi_k\right)^{\frac{n_1 + \dots + n_p}{n} - p - 1} \left(1 + \frac{1}{n} \sum_{k=1}^p n_k \xi_k\right). \quad (3)$$

The restriction $\Psi : [0, \infty)^p \rightarrow [0, \infty)^p$ and its inverse $\Psi^{-1} : [0, \infty)^p \rightarrow [0, \infty)^p$ are analytic. The principal solution of (1) is $Z := W^{-\frac{1}{n}} \circ \Psi^{-1} : [0, \infty)^p \rightarrow [1, \infty)$.

Proof The first assertion is Proposition 1. Define the Kronecker symbol

$$\delta_{i,j} = \begin{cases} 1, & \text{if } i = j \\ 0 & \text{if } i \neq j. \end{cases}$$

Since the matrix

$$\frac{\partial \Psi_i}{\partial \xi_j} = W^{\frac{n_i}{n} - 1} \left(\delta_{i,j} + \xi_i \left(\frac{n_i}{n} - 1 \right) W^{-1} \right),$$

is a product of two matrices, it follows that

$$\frac{\partial(x_1, \dots, x_p)}{\partial(\xi_1, \dots, \xi_p)} := \det \frac{\partial \Psi_i}{\partial \xi_j} = W^{\frac{\sum_{k=1}^p n_k}{n} - p} \det \left(\delta_{i,j} + \xi_i \left(\frac{n_i}{n} - 1 \right) W^{-1} \right).$$

Proposition 2 implies that

$$\det \left(\delta_{i,j} + \xi_i W^{-1} \right) = 1 + \sum_{k=1}^p \xi_k \left(\frac{n_k}{n} - 1 \right) W^{-1} = W^{-1} \left(1 + \frac{1}{n} \sum_{k=1}^p n_k \xi_k \right)$$

and concludes the proof of the second assertion. Since the Jacobian of Ψ is nonzero and holomorphic on $\mathbb{C}^p \setminus \Xi$ and $[0, \infty)^p \subset \mathbb{C}^p \setminus \Xi$, assertion three follows from the inverse function theorem for holomorphic functions ([9], p. 40, Theorem 2). The fourth assertion then follows from Lemma 2.

Remark 1 Let $n = 2, p = 1$. Then $\Psi : \mathbb{C} \setminus (-\infty, -1] \rightarrow \mathbb{C} \setminus i((-\infty, -2] \cup [2, \infty))$ is a holomorphic bijection with inverse

$$\Psi^{-1}(z) = -1 + \left(\frac{z}{2} + \sqrt{1 + \left(\frac{z}{2} \right)^2} \right)^2.$$

Letting $\xi_1 = -1 + e^{s+it}$ with $s \in \mathbb{R}$ and $-\pi < t < \pi$ gives $\Psi(\xi_1) = 2 \sinh \frac{s+it}{2} = u + iv$ where $u = 2 \sinh \frac{s}{2} \cosh \frac{t}{2}$ and $v = 2 \cosh \frac{s}{2} \sinh \frac{t}{2}$. For $t = 0$ this gives the curve $v = 0$. For fixed $t \neq 0$ this gives the branch of the hyperbola described by the equation $\left(\frac{v}{\sinh t/2} \right)^2 - \left(\frac{u}{\cosh t/2} \right)^2 = 4$ with $tv \geq 0$.

Question 1 For $n \geq 3$ and $p \geq 1$, is Ψ injective and what is its image?

Following (3) Mellin says that using the known formula, derived in Proposition 2, in combination with (2) and (3), one can deduce the following result:

Lemma 4 *The principal solution of (1) satisfies*

$$\int_0^\infty \dots \int_0^\infty Z(x_1, \dots, x_p)^\alpha x_1^{u_1-1} \dots x_p^{u_p-1} dx_1 \dots dx_p = \frac{\alpha}{n} \frac{\Gamma(u) \Gamma(u_1) \dots \Gamma(u_p)}{\Gamma(u + u_1 + \dots + u_p + 1)}, \quad (4)$$

where $\alpha > 0$, and the real parts of $u := \frac{\alpha}{n} - \frac{n_1}{n} u_1 - \dots - \frac{n_p}{n} u_p$, and u_1, \dots, u_p are positive.

Proof Equation (2) and Jensen's inequality implies

$$W^{(u_1 n_1 + \dots + u_p n_p)/n} > p^{u_1 + \dots + u_p} x_1^{u_1} \dots x_p^{u_p}.$$

Therefore, since $Z = W^{-\frac{1}{n}}$ and $\alpha > u_1 n_1 + \dots + u_p n_p$, the integral I in (4) exists. Proposition 3 implies that $F : [0, \infty)^p \rightarrow [0, \infty)^p$ is a bijective diffeomorphism, so I can

be expressed as an integral over $[0, \infty)^p$ with respect to the variables ξ_1, \dots, ξ_p . Hence (2) and (3) give

$$I = \int_0^\infty \cdots \int_0^\infty \frac{(1 + \frac{n_1}{n}\xi_1 + \cdots + \frac{n_p}{n}\xi_p)\xi_1^{u_1-1} \cdots \xi_p^{u_p-1}}{W^\omega} d\xi_1 \cdots d\xi_p$$

where $\omega := u + u_1 + \cdots + u_p + 1$. Proposition (2) implies that $I + I_0 + I_1 + \cdots + I_p$ where

$$I_0 = \frac{\Gamma(\omega - u_1 - \cdots - u_p)\Gamma(u_1) \cdots \Gamma(u_p)}{\Gamma(\omega)} = \frac{\Gamma(u+1)\Gamma(u_1) \cdots \Gamma(u_p)}{\Gamma(\omega)},$$

and

$$I_i = \frac{n_i}{n} \frac{\Gamma(\omega - u_1 - \cdots - u_p - 1)\Gamma(u_1) \cdots \Gamma(u_i + 1) \cdots \Gamma(u_p)}{\Gamma(\omega)} = \frac{n_i u_i}{nu} I_0.$$

We conclude the proof by observing that

$$I = \left(1 + \frac{1}{u} \sum_{i=1}^p \frac{n_i u_i}{n}\right) I_0 = \frac{\alpha}{nu} I_0 = \frac{\alpha}{n} \frac{\Gamma(u+1)}{u} \frac{\Gamma(u_1) \cdots \Gamma(u_p)}{\Gamma(\omega)}.$$

2 Computing Z^α from its Mellin Transform

Mellin continues: “The law of reciprocity relating to the integrals of this species, demonstrated by us in a previous work [5], allows us to invert as follows (4):”

$$Z(x_1, \dots, x_p)^\alpha =$$

$$\frac{1}{(2\pi i)^p} \int_{a_1-i\infty}^{a_1+i\infty} \cdots \int_{a_p-i\infty}^{a_p+i\infty} \frac{\alpha}{n} \frac{\Gamma(u)\Gamma(u_1) \cdots \Gamma(u_p)}{\Gamma(u+u_1+\cdots+u_p+1)} x_1^{-u_1} \cdots x_p^{-u_p} du_1 \cdots du_p, \quad (5)$$

$$\alpha - n_1 a_1 - \cdots - n_p a_p > 0, \quad a_s > 0, \quad u := \frac{\alpha}{n} - \frac{n_1}{n} u_1 - \cdots - \frac{n_p}{n} u_p$$

$$u + u_1 + \cdots + u_p = \frac{\alpha}{n} + \frac{n_1}{n} u_1 + \cdots + \frac{n_p}{n} u_p, \quad n_s := n - n_s.$$

“This formula constitutes our solution of (1). It supposes that

$$-\frac{n_s \pi}{2n} < \arg(x_s) < \frac{n_s \pi}{2n}$$

but we can extend our domain of validity by suitably deforming the integration paths.”

The right side of (4) is the Mellin transform of $Z(x_1, \dots, x_p)^\alpha$ and (5) represents $Z(x_1, \dots, x_p)^\alpha$ by the inverse Mellin transform. Lacking the luxury of accessing [5] we refer the reader to the derivation of the inverse Mellin transform via the Fourier transform by Debnath and Bhatta in section 8.1 of [3]. They address the univariate case, but the extension to multivariate case is straightforward.

3 Generalized Hypergeometric Functions

Let n be a positive integer and $f_1, \dots, f_p, g_1, \dots, g_s : \mathbb{C}^p \rightarrow \mathbb{C}$ be entire functions and $F : \mathbb{C}^p \rightarrow \mathbb{C}$ satisfy the following system of functional equations

$$F(u_1, \dots, u_s + n, \dots, u_p) = \frac{f_s(u_1, \dots, u_p)}{g_s(u_1, \dots, u_p)} F(u_1, \dots, u_p), \quad s = 1, \dots, p. \quad (6)$$

and such that the following integral converges and does not change when we move the (vertical) integration paths for each u_s n units to the right:

$$y(x_1, \dots, x_p) = \frac{1}{(2\pi i)^p} \int_{u_1} \cdots \int_{u_p}, F(u_1, \dots, u_p) x_1^{-u_1} \cdots x_p^{-u_p}, \quad (x_1, \dots, x_p) \in [0, \infty)^p.$$

Mellin cites a result [5] that y satisfies the following system of partial differential equations

$$f_s \left(-x_1 \frac{\partial}{\partial x_1}, \dots, -x_p \frac{\partial}{\partial x_p} \right) y = g_s \left(-x_1 \frac{\partial}{\partial x_1}, \dots, -x_p \frac{\partial}{\partial x_p} \right) s_s^n y, \quad s = 1, \dots, p. \quad (7)$$

Remark 2 (7) is a system of PDE's of finite order iff f_s and g_s are polynomials.

Mellin calls solutions of (6) hypergeometric type if the factors of f_s, g_s have the form

$$c_1 u_1 + \dots + c_p u_p + a$$

where each c_i is a rational real number.

Remark 3 Polynomials are not hypergeometric functions or series as defined in [4, 10] but they are of hypergeometric type since $\prod_{k=0}^{m-1} (x \frac{d}{dx} - k) = x^m \frac{d^m}{dx^m}$.

Formally (7) follows from (6) since the functions $x_s^{-u_s}$ are eigenfunctions with eigenvalue $-u_s$ of the Euler operator $x_s \frac{\partial}{\partial x_s}$, $s = 1, \dots, p$. Clearly the function

$$F(u_1, \dots, u_p) = \frac{\alpha}{n} \frac{\Gamma(u) \Gamma(u_1) \dots \Gamma(u_p)}{\Gamma(u + u_1 + \dots + u_p + 1)}$$

satisfies (6) where f_s and g_s have the form above, hence $Z(u_1, \dots, u_p)^\alpha$ is of hypergeometric type whenever $\alpha > n_1 n$.

Question 2 How large must α be to ensure that Z^α is a solution of (1) of hypergeometric type? The condition $\alpha > nn_1$ is sufficient but not necessary because for $n = 1, p = 1$ the root $Z(x_1) := -\frac{x_1}{2} + \sqrt{1 + (x_1/2)^2}$ is of hypergeometric type. Semusheva and Tsikh [8] proved this fact directly by deriving the following Mellin–Barnes integral representation

$$Z(x_1) = \frac{1}{4\pi i} \int_{\Re z = \frac{1}{2}} \frac{\Gamma(z) \Gamma((1+z)/2)}{\Gamma((3+z)/2)} x_1^z dz.$$

Remark 4 For a comprehensive development of Mellin's solution of (3) and systems of differential equations of hypergeometric type see ([1], Chapitre V).

4 Appendix: Crucial Propositions

We prove results required to derive the equations in Mellin's paper.

Let $e_1 := (1, 0, \dots, 0), e_2 := (0, 1, \dots, 0), \dots, e_p = (0, \dots, 0, 1)$ be the standard basis for \mathbb{R}^p . For $s > 0$ let H_s denote the convex hull of $\{se_i : i = 1, \dots, p\}$ and let K_s denote the convex hull of $\{s(1+s)^{\frac{n_i}{n}-1} e_i : i = 1, \dots, p\}$.

Proposition 1 $\Psi : [0, \infty)^p \rightarrow [0, \infty)^p$ is a bijection.

Proof For every $s > 0$ define the linear map $L_s : \mathbb{R}^p \rightarrow \mathbb{R}^p$ by

$$(L_s y)_i := y_i (1+s)^{\frac{n_i}{n}-1}, \quad i = 1, \dots, p.$$

If $(\xi_1, \dots, \xi_p) \in H_s$, then $W(\xi_1, \dots, \xi_p) = 1+s$ so Equation 2 implies that

$$\Psi(\xi_1, \dots, \xi_p) = L_s(\xi_1, \dots, \xi_p) = \sum_{i=1}^p \frac{\xi_i}{s} s(1+s)^{\frac{n_i}{n}-1} e_i \in K_s,$$

hence the restriction $\Psi : H_s \rightarrow K_s$ is a linear bijection. Clearly $\Psi(0, \dots, 0) = (0, \dots, 0)$ and $[0, \infty)^p$ is a disjoint union of $\{(0, \dots, 0)\}$ and the sets $H_s, s > 0$. Since for $i = 1, \dots, p$, the function $s(1+s)^{\frac{n_i}{n}-1}$ is increasing, it follows that $[0, \infty)^p$ is a disjoint union of $\{(0, \dots, 0)\}$ and the sets $K_s, s > 0$. This concludes the proof.

Proposition 2 Let $p \geq 1$, y_1, \dots, y_p be indeterminates, and define the matrix

$$M(y_1, \dots, y_p) := \begin{bmatrix} 1 + y_1 & y_1 & \cdots & y_1 & y_1 \\ y_2 & 1 + y_2 & \cdots & y_2 & y_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ y_{p-1} & y_{p-1} & \cdots & 1 + y_{p+1} & y_{p+1} \\ y_p & y_p & \cdots & y_p & 1 + y_p \end{bmatrix}.$$

Then

$$\det M(y_1, \dots, y_p) = 1 + y_1 + \cdots + y_p.$$

Proof For $i = 1, \dots, p$ let $\partial_i := \frac{\partial}{\partial y_i}$. Clearly $\det M(y_1, \dots, y_p)$ is a polynomial with constant term $\det M(0, \dots, 0) = \det I_p = 1$ and of degree at most 1 in each variable y_1, \dots, y_p so for $i = 1, \dots, p$

$$\partial_i^2 \det M(y_1, \dots, y_p) = 0.$$

Let $M_i(y_1, \dots, y_p)$ denote the matrix obtained from $M(y_1, \dots, y_p)$ by replacing each entry in its i -th row with 1 and for $j \neq i$ let $M_{i,j}(y_1, \dots, y_p)$ denote the matrix obtained from $M(y_1, \dots, y_p)$ by replacing each entry in its i -row and its j -row by 1. Since a determinant of a matrix is a linear function of each of its row vectors,

$$\partial_i \det M(y_1, \dots, y_p) := \det M_i(y_1, \dots, y_p)$$

and for $j \neq i$

$$\partial_j \partial_i \det M(y_1, \dots, y_p) := \det M_{i,j}(y_1, \dots, y_p) = 0$$

since $M_{i,j}(y_1, \dots, y_p)$ has two identical rows. Therefore Taylor's expansion gives

$$\det M(y_1, \dots, y_p) = 1 + \sum_{i=1}^p y_i \partial_i \det M(0, \dots, 0) = 1 + \sum_{i=1}^p y_i \det M_i(0, \dots, 0).$$

It suffices to prove $\det M_i(0, \dots, 0) = 1$, $i = 1, \dots, p$. This follows since

$$M_i(0, \dots, 0) = \begin{bmatrix} 1 & 0 & \cdots & 0 & \cdots & 0 & 0 \\ 0 & \ddots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 1 & 0 & 0 & \cdots & 0 \\ 1 & \cdots & 1 & 1 & 1 & \cdots & 1 \\ 0 & \cdots & 0 & 0 & 1 & \cdots & 0 \\ 0 & \ddots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & \cdots & 0 & \cdots & 0 & 1 \end{bmatrix}.$$

Remark 5 Nilsson and Passare ([7], Example 1) used basic calculus to compute the following integrals for $\Re u_i \in (0, 1)$, $i = 1, 2$.

$$\int_0^\infty \int_0^\infty \frac{\xi_1^{u_1-1} \xi_2^{u_2-1}}{1 + \xi_1 + \xi_2} d\xi_1 d\xi_2 = \frac{\Gamma(u_1) \Gamma(u_2) \Gamma(1 - u_1 - u_2)}{\Gamma(1)}$$

The following result uses exterior calculus to extend their computation.

Proposition 3 For every positive integer p , complex u_1, \dots, u_p satisfying $\Re u_i > 0$, and $\omega > \max\{\Re u_i\}$ the integral below converges and satisfies the stated identity.

$$\int_0^\infty \cdots \int_0^\infty \frac{\xi_1^{u_1-1} \cdots \xi_p^{u_p-1}}{(1 + \xi_1 + \cdots + \xi_p)^\omega} d\xi_1 \cdots d\xi_p = \frac{\Gamma(u_1) \cdots \Gamma(u_p) \Gamma(\omega - u_1 - \cdots - u_p)}{\Gamma(\omega)}.$$

Proof Multiplying both sides of the asserted identity in the lemma by

$$\Gamma(\omega) = \int_0^\infty x^{\omega-1} e^{-x} dx$$

gives the equivalent identity $I_1(u_1, \dots, u_p, \omega) = \Gamma(u_1) \cdots \Gamma(u_p) \Gamma(\omega - u_1 - \cdots - u_p)$ where

$$I_1(u_1, \dots, u_p, \omega) := \int_0^\infty \cdots \int_0^\infty \frac{\xi_1^{u_1-1} \cdots \xi_p^{u_p-1}}{(1 + \xi_1 + \cdots + \xi_p)^\omega} x^{\omega-1} e^{-x} d\xi_1 \wedge \cdots \wedge d\xi_p \wedge dx.$$

Here we express the volume form using exterior products to implement the following change of variables: $y := x/(1 + \xi_1 + \dots + \xi_p)$ and $z_i := y \xi_i, i = 1, \dots, p$. Then $I_1(\xi_1, \dots, \xi_p, x) = I_2(z_1, \dots, z_p, y)$ where

$$I_2(z_1, \dots, z_p, y) := \int_0^\infty \dots \int_0^\infty z_1^{u_1-1} e^{-z_1} \dots z_p^{u_p-1} e^{-z_p} y^{\omega-u_1-\dots-u_p-1} e^{-y} \theta$$

where

$$\theta = \frac{y^p}{1 + \xi_1 + \dots + \xi_p} d\xi_1 \wedge \dots \wedge d\xi_p \wedge dx$$

In order to finish the proof it suffices to prove that $dz_1 \wedge \dots \wedge dz_p \wedge dy = \theta$. Since $dz_i = \xi_i dy + y d\xi_i$ and $dy \wedge dy = 0$ it follows that

$$dz_1 \wedge \dots \wedge dz_p \wedge dy = y^p d\xi_1 \wedge \dots \wedge d\xi_p \wedge dy.$$

Substitute

$$dy = \frac{1}{1 + \xi_1 + \dots + \xi_p} dx - \frac{x}{(1 + \xi_1 + \dots + \xi_p)^2} \left(\sum_{k=1}^p d\xi_k \right)$$

and use the fact that $d\xi_i \wedge d\xi_i = 0$ to obtain

$$dz_1 \wedge \dots \wedge dz_p \wedge dy = \frac{y^p}{1 + \xi_1 + \dots + \xi_p} d\xi_1 \wedge \dots \wedge d\xi_p \wedge dx = \theta$$

which concludes the proof.

Acknowledgments We thank Elaine Wong of the Johann Radon Institute for Computational and Applied Mathematics (RICAM), Linz for her meticulous proofreading of and corrections to this article.

References

- [1] G. Belardinelli, *Fonctions hypergéométriques de plusieurs variables et résolution analytique des équations algébriques générales*. Mémoires des sciences mathématiques, no. 145, Gauthier-Villars, Paris, 1960.
- [2] M. R. Birkeland, *Ein allgemeiner Satz über algebraische Gleichungen*. Annales Academiae Scientiarum Fennicae, ser. A, t. 7, (1915).
- [3] L. Debnath and D. Bhatta, *Integral Transforms and Their Applications, 2nd edition.*, Chapman & Hall/CRC, New York, 2007.
- [4] M. Kauers and P. Paule, *The Concrete Tetrahedron, Symbolic Sums, Recurrence Equations, Generating Functions, Asymptotic Estimates*. Springer, Vienna, 2011.
- [5] H. J. Mellin, *Zur Theorie zweier allgemeiner Klassen bestimmter Integrale*. 75 S. Ibid, Bd. 22 (1896).
- [6] H. J. Mellin, *Résolution de l'équation algébrique générale à l'aide de la fonction gamma*. C. R. Acad. Sci. Paris Sér. I Math. 172 (1921) 658–661.
- [7] L. Nilsson and M. Passare, *Mellin transforms of multivariate rational functions*. Journal of Geometric Analysis, 23 (2013) 24–46. arXiv:1010.5060
- [8] A. Y. Semusheva and A. K. Tsikh, *Continuation on Mellin's research on solving algebraic equations*. Complex analysis and differential operators. To the 150th anniversary of S. V. Kovalevskaya. Executive editor A. K. Tsikh, Krasnoyarsk State University, Krasnoyarsk, 2000. 196 p. (Russian)
- [9] B. V. Shabat, *Introduction to Complex Analysis, Part II Functions of Several Variables*. American Mathematical Society, Providence, Rhode Island, 1992.
- [10] L. J. Slater, *Generalized Hypergeometric Functions*. Cambridge University Press, 1960.