Analyticity of the energy in an Ising spin glass with correlated disorder

Hidetoshi Nishimori

Institute of Innovative Research, Tokyo Institute of Technology, Yokohama 226-8503, Japan Graduate School of Information Sciences, Tohoku University, Sendai 980-8579, Japan RIKEN Interdisciplinary Theoretical and Mathematical Sciences Program (iTHEMS), Wako, Saitama 351-0198, Japan

The average energy of the Ising spin glass is known to have no singularity along a special line in the phase diagram although there exists a critical point on the line. This result on the model with uncorrelated disorder is generalized to the case with correlated disorder. For a class of correlations in disorder that suppress frustration, we show that the average energy in a subspace of the phase diagram is expressed as the expectation value of a local gauge variable of the Z_2 gauge Higgs model, from which we prove that the average energy has no singularity although the subspace is likely to have a phase transition on it. Though it is difficult to obtain an explicit expression of the energy in contrast to the case of uncorrelated disorder, an exact closed-form expression of a physical quantity related to the energy is derived in three dimensions using a duality relation. Identities and inequalities are proved for the specific heat and correlation functions.

I. INTRODUCTION

The problem of spin glass is one of the most challenging topics in statistical physics [1, 2]. Only a very limited number of exact solutions are known so far, among which the Parisi solution of the mean-field-type Sherrington-Kirkpartick model [3] stands out as a distinguished achievement [4, 5]. Another rare example is the exact solution for the average energy of the Ising spin glass in finite dimensions on a special line in the phase diagram [1, 6, 7]. The resulting expression of the energy is written as a simple hyperbolic tangent of the inverse temperature without any singularity. Since a multicritical point (transition point) lies on this special line in the phase diagram, it is highly non-trivial and counter-intuitive that the exact average energy is an analytic function. Other physical quantities such as the specific heat and magnetic susceptibility are expected to have a singularity at the phase transition, but nothing is known so far exactly or rigorously on those quantities in finite-dimensional models except for the existence of a phase transition on the line as concluded from the existence of ferromagnetic order in the low-temperature side of the line [8] (see also [9] for the continuous-variable case).

Those results concern the Edwards-Anderson model [10], in which disorder of a given interaction bond distributes independently of disorder of other bonds. In real spin glass materials, disorder exists mostly in sites, not in bonds [11]. Edwards and Anderson assumed that properties of spin glasses will be independent of the type of disorder and proposed their model with uncorrelated disorder. One of the important differences between site and bond disorder is the existence or absence of correlation of disorder among nearby bond variables. For example, if the position of a magnetic atom in a metal is affected by disorder, all interactions between this and other atoms are changed in a correlated manner. It therefore makes sense to study how correlations in disorder would affect the system properties.

There have been several attempts to study the effects of correlation in disorder [12–15]. It has generally been observed that correlations, if weak, do not significantly modify the system properties qualitatively. These studies exploit numerical methods or mean-field-type models, and it is desirable to establish analytical results in finite-dimensional systems.

We introduce a model with correlations of a specific type in disorder, which makes it possible to reduce the expression of the average energy to the expectation value of a local variable in the Z_2 gauge Higgs model [16]. Based on this reduction, we prove that the average energy is an analytic function in a subspace of the phase diagram, a generalization of the special line in the case of uncorrelated disorder. It is difficult to derive a closed-form exact solution for the average energy when correlations are introduced. Nevertheless, using self duality, we show in the case of three spatial dimensions that a simple formula holds for a physical quantity related to the energy. We also derive several non-trivial identities and inequalities for the specific heat and correlation functions, generalizing the results for uncorrelated disorder. Also discussed are more complex types of correlations in disorder.

In Sec. II, we define the model and analyze its properties. Conclusion is given in Sec. III, and some technical details are described in Appendixes.

II. ISING SPIN GLASS WITH CORRELATED DISORDER

We first define the model of correlated disorder and then study the properties of the corresponding Ising spin glass, in particular the behavior of the average energy in a subspace of the phase diagram.

A. Model of correlated disorder

As mentioned in the Introduction, it is reasonable to study a spin-glass system with short-range correlations in disorder variables. It is likely that site disorder has a different degree of frustration than in the case of uncorrelated bond disorder, a prominent example of which is the Mattis model [17] without frustration but with site-dependent quenched disorder $\xi_i = \pm 1$,

$$H_{\text{Mattis}} = -J \sum_{\langle ij \rangle} \xi_i \xi_j S_i S_j, \tag{1}$$

where $S_i(\pm 1)$ is the Ising variable at site i, and $\langle ij \rangle$ runs over interacting spin pairs on a lattice with interaction strength J. An example with site disorder but with frustration is the Hopfield model [18–21],

$$H_{\text{Hopfield}} = -\frac{1}{N} \sum_{i,j} J_{ij} S_i S_j, \tag{2}$$

where N is the number of sites, and the interaction is composed of site-dependent quenched variables $\{\xi_i^{\mu} = \pm 1\}$,

$$J_{ij} = \sum_{\mu=1}^{r} \xi_i^{\mu} \xi_j^{\mu}. \tag{3}$$

This is a model with site-dependent disorder, in which correlation exists between disorder of different bonds J_{ij} and J_{jk} sharing a site index j. When r = 1, the model reduces to the Mattis model of Eq. (1) without frustration, and in the limit of $r \to \infty$, the model becomes equivalent to the Sherrington-Kirkpatrick model with uncorrelated bond disorder with strong frustration according to the central limit theorem. The Hopfield model thus interpolates the unfrustrated and fully frustrated cases by the parameter r.

These examples motivate us to introduce a system with correlated disorder in bond variables with a parameter to control the degree of frustration. We therefore propose and analyze the following probability distribution of bond-assigned disorder variables $\{\tau_{ij} = \pm 1\}$ with control parameters K_1 and K_2 ,

$$P(\tau, K_1, K_2) = \frac{1}{Z_{\tau}(K_1, K_2)} \exp\left(K_1 \sum_{\langle ij \rangle} \tau_{ij} + K_2 \sum_{\square} \tau_{\square}\right), \tag{4}$$

where $Z_{\tau}(K_1, K_2)$ is the normalization factor (or the partition function of disorder variables)

$$Z_{\tau}(K_1, K_2) = \sum_{\{\tau_{ij} = \pm 1\}} \exp\left(K_1 \sum_{\langle ij \rangle} \tau_{ij} + K_2 \sum_{\square} \tau_{\square}\right), \tag{5}$$

and τ_{\square} is the product of four disorder variables around a unit plaquette denoted by \square on the d-dimensional hypercubic lattice, $\tau_{\square} = \tau_{ij}\tau_{jk}\tau_{kl}\tau_{li}$ [22]. This is a generalization of the conventional $\pm J$ model $(K_2 = 0)$ to the case with correlation in disorder $(K_2 > 0)$. The above specific type of correlation was motivated by the following reasons. (i) Reference [14] studied the case of just the second term $(K_1 = 0, K_2 > 0)$ and measured critical exponents in three dimension by numerical simulations. We are interested in the interplay between the first (K_1) and second (K_2) terms in the above more general model. (ii) The above form allows us to directly control the degree of frustration by the coefficient K_2 . (iii) The K_2 term is gauge invariant as will be discussed below, which is essential for our theory in the following sections to be applicable. Notice that the exponent in the probability distribution Eq. (4) is of the same form as the action of the Z_2 gauge Higgs model [16], and this analogy facilitates our analysis below.

The Hamiltonian of the Ising spin glass is

$$H = -K \sum_{\langle ij \rangle} \tau_{ij} S_i S_j, \tag{6}$$

where K is the dimensionless coupling constant (the inverse temperature K = 1/T). We study the properties of this Hamiltonian with the quenched disorder variables $\{\tau_{ij}\}$ chosen from the probability distribution of Eq. (4). Possible generalization of the probability distribution will be discussed at the end of this section.

Average energy in a subspace of the phase diagram

1. Expression of the average energy

We first show that the average energy can be expressed in a simple formula in terms of the \mathbb{Z}_2 gauge Higgs model in a subspace of the phase diagram.

The average energy E of the model defined in the preceding section is written as

$$E(K, K_1, K_2) = \frac{1}{Z_{\tau}(K_1, K_2)} \sum_{\{\tau_{ij} = \pm 1\}} e^{K_1 \sum \tau_{ij} + K_2 \sum \tau_{\square}} \langle H \rangle_K, \tag{7}$$

where H is the Hamiltonian of Eq. (6) and the angular brackets denote the thermal average,

$$\langle H \rangle_K = -\frac{1}{Z_s(K)} \frac{\partial}{\partial K} Z_s(K)$$
 (8)

with the partition function of the Ising model.

$$Z_s(K) = \sum_{\{S_i = \pm 1\}} e^{K \sum \tau_{ij} S_i S_j}.$$
(9)

Notice that the constant multiplicative factor K is dropped for simplicity in Eq. (8) of the energy.

Following the standard prescription [1, 6], we apply the gauge transformation

$$S_i \to S_i \sigma_i, \, \tau_{ij} \to \tau_{ij} \sigma_i \sigma_j \, (\sigma_i = \pm 1) \, (\forall i, j)$$
 (10)

to Eqs. (7) and (9). The thermal average $\langle H \rangle_K$ is gauge invariant, and only the term in the exponent of Eq. (7) with coefficient K_1 is affected. We sum the resulting expression over all possible values of $\{\sigma_i\}$, and divide the expression by 2^N , and find

$$-E(K, K_1, K_2) = \frac{1}{2^N Z_{\tau}(K_1, K_2)} \sum_{\{\tau_{ij} = \pm 1\}} \sum_{\{\sigma_i = \pm 1\}} e^{K_1 \sum \tau_{ij} \sigma_i \sigma_j + K_2 \sum \tau_{\square}} \frac{\partial_K \sum_{\{S_i\}} e^{K \sum \tau_{ij} S_i S_j}}{\sum_{\{S_i\}} e^{K \sum \tau_{ij} S_i S_j}}.$$
 (11)

Since τ_{\square} is gauge invariant and $\{\sigma_i\}$ appears only in the term with coefficient K_1 , the factor in the numerator $\sum_{\{\sigma_i\}} e^{K_1 \sum \tau_{ij} \sigma_i \sigma_j}$ cancels with the denominator $Z_s = \sum_{\{S_i\}} e^{K \sum \tau_{ij} S_i S_j}$ when $K = K_1$ and we are left with a simplified expression

$$-E(K, K, K_{2}) = \frac{1}{2^{N} Z_{\tau}(K, K_{2})} \sum_{\{\tau_{ij}\}} e^{K_{2} \sum \tau_{\square}} \frac{\partial}{\partial K} \sum_{\{S_{i}\}} e^{K \sum \tau_{ij} S_{i} S_{j}}$$

$$= \frac{1}{2^{N} Z_{\tau}(K, K_{2})} \sum_{\{S_{i}\}} \sum_{\{\tau_{ij}\}} \left(\sum_{\langle ij \rangle} \tau_{ij} S_{i} S_{j} \right) e^{K \sum \tau_{ij} S_{i} S_{j} + K_{2} \sum \tau_{\square}}$$

$$= \frac{1}{Z_{\tau}(K, K_{2})} \sum_{\langle ij \rangle} \left(\sum_{\{\tau_{ij}\}} \tau_{ij} \right) e^{K \sum \tau_{ij} + K_{2} \sum \tau_{\square}},$$

$$(12)$$

where we have applied the gauge transformation $\tau_{ij} \to \tau_{ij} S_i S_j$ to the second line to derive the third line. In the case of uncorrelated distribution of disorder $(K_2 = 0)$, the condition $K = K_1$ defines a line in the K- K_1 phase diagram, sometimes called the Nishimori line [9, 23-38]. It is known that the low-temperature side of this line $(K = K_1 \gg 1)$ lies in the ferromagnetic phase in two and higher dimensions [8, 39] and that the spin glass phase exists away from the line [1, 6]. The line generally goes across a multicritical point where the paramagnetic, ferromagnetic and spin glass phases meet in three and higher dimensions [1, 40]. Our present model with correlation in disorder has an additional parameter K_2 to control the degree of correlation, and the condition $K = K_1$ defines a subspace in the three-dimensional phase diagram drawn in terms of K_1, K_2 , and K.

The expression of the average energy in Eq. (12) in the subspace $K = K_1$ can be interpreted as the expectation value of the local bond variable τ_{ij} summed over $\langle ij \rangle$ for the Z_2 gauge Higgs model [16] [41]. The condition $K = K_1$ has greatly simplified the expression of the average energy, leaving only the expectation value of a single τ_{ij} summed up over $\langle ij \rangle$.

The average energy of Eq. (12) is an analytic function of $K(=K_1)$ for sufficiently small $K_2(>0)$:

Theorem 1. The average energy $E(K, K, K_2)$ of Eq. (12) is analytic in K for sufficiently small K_2 in the thermodynamic limit.

This result is a known property of the Z_2 lattice gauge theory, see e.g. Ref. [42] for a theoretical analysis and Refs. [43–47] for numerical studies. Since the argument presented in Ref. [42] is fairly sketchy, we provide a more formal proof in Appendix A.

When $K_2 = 0$ (uncorrelated disorder), the analyticity is trivial. Equation (12) immediately gives a simple formula

$$E(K, K, 0) = -N_B \tanh K,\tag{13}$$

where N_B is the number of bonds. Nevertheless, the line defined by $K = K_1$ in the K- K_1 phase diagram has a transition point on it since the low-temperature side $(K \gg 1)$ is in the ferromagnetic phase in two and higher dimensions [8] and the high-temperature side is trivially in the paramagnetic phase. Therefore the absence of singularity in E(K, K, 0) is highly non-trivial, a consequence of delicate balance of the probability weight $P(\tau, K_1, 0)$ and the partition function $Z_s(K)$ in Eq. (11). Theorem 1 claims that this non-trivial analyticity remains valid even after the introduction of correlation in disorder. This may look rather trivial mathematically, but we believe it to be highly non-trivial physically.

Although it is difficult to prove that the subspace $K=K_1$ has a transition point (critical point) on it when $K_2>0$, in contrast to the case $K_2=0$ where it is proven [8, 9], we expect that the physical properties of the system are unlikely to change dramatically by the introduction of small but finite K_2 as suggested by the continuation of analyticity of the average energy as in Theorem 1. An indirect indication of the stability of the low-temperature ($K=K_1\gg 1$) ferromagnetic phase after the introduction of K_2 is provided by small- K_2 perturbation of the distribution function of a single-bond variable defined by

$$P(l, K_1, K_2) = \sum_{\{\tau_{ij} = \pm 1\}} \delta_{l,\tau_{ij}} P(\tau, K_1, K_2) \quad (l = \pm 1),$$
(14)

where δ is Kronecker's delta. We define the probability for a bond to be positive (ferromagnetic) p_+ by

$$P(l, K_1, K_2) = p_+ \, \delta_{l,1} + (1 - p_+) \, \delta_{l,-1}. \tag{15}$$

Then, as shown in Appendix B, p_+ is found by first-order perturbation in $\tanh K_2$ as

$$p_{+} = p \left(1 + 2c(1-p)(2p-1)^{3} \tanh K_{2} + \mathcal{O}(\tanh K_{2})^{2} \right), \tag{16}$$

where c is a positive constant and p is the probability of $\tau_{ij}=1$ when $K_2=0$ (i.e., p_+ for $K_2=0$), defined through the ratio of probabilities for negative (1-p) and positive (p) values of τ_{ij} : $e^{-K_1}/e^{K_1}=(1-p)/p$. Equation (16) suggests that the probability of a single bond to be positive (ferromagnetic) increases as we introduce small K_2 . This implies (but does not prove) that the low-temperature ferromagnetic phase becomes more stable by the introduction of $K_2>0$. This is intuitively reasonable because K_2 suppresses frustration, increasing stability of the ferromagnetic phase. Since the existence of a paramagnetic phase for $K=K_1\ll 1$ is trivial, we may reasonably expect that there is a phase transition as a function of $K=K_1$ for $K_2>0$ in spite of the absence of singularity in the average energy.

The range of K_2 where the analyticity of average energy holds depends on the spatial dimension and the lattice structure. In the case of the two-dimensional square lattice, the Z_2 gauge Higgs model of Eq. (12) is equivalent to the conventional ferromagnetic Ising model in a finite field on the square lattice according to the duality relation [48]. Since the ferromagnetic Ising model in a field has no singularity, the energy of Eq. (12) is analytic for any positive value of K_2 [49]. In three dimensions, Monte Carlo simulation shows the range of analyticity to be $0 \le K_2 < 0.6893$ on the cubic lattice [47].

3. Identities for the average energy in three dimensions

It is generally difficult to derive an explicit compact expression of the average energy for $K = K_1$ except for the case of $K_2 = 0$ as written in Eq. (13). Nevertheless, identities involving the average energy can be obtained in the case of the three-dimensional cubic lattice using the self duality of the model [48].

Let us define the average energy per spin as

$$e(K_1, K_2) = \frac{1}{N} E(K_1, K_1, K_2) = -\frac{1}{N Z_{\tau}(K_1, K_2)} \sum_{\{\tau_{ij} = \pm 1\}} \left(\sum_{\langle ij \rangle} \tau_{ij} \right) e^{K_1 \sum_{\tau_{ij} + K_2 \sum_{\tau_{\square}}} .$$
 (17)

and the average plaquette energy per spin by

$$e_p(K_1, K_2) = -\frac{1}{N Z_{\tau}(K_1, K_2)} \sum_{\{\tau_{ij} = \pm 1\}} \left(\sum_{\square} \tau_{\square} \right) e^{K_1 \sum_{\tau_{ij}} + K_2 \sum_{\tau_{\square}}}.$$
 (18)

For these quantities, we can derive the following relation for a sufficiently large system on the cubic lattice where the boundary effects are sufficiently small,

$$e(K_1, K_2) + 3\tanh K_1 = -\frac{1}{\sinh 2K_1} \left(e_p(K_1^*, K_2^*) + 3 \right), \tag{19}$$

where the dual couplings K_1^* and K_2^* are defined as

$$\tanh K_1^* = e^{-2K_2}, \ \tanh K_2^* = e^{-2K_1}.$$
 (20)

The proof is based on the duality relation of Wegner [48] for the \mathbb{Z}_2 gauge Higgs model in three dimensions,

$$\frac{Z_{\tau}(K_1, K_2)}{2^{3N}(\cosh K_1 \cosh K_2)^{3N}} = \frac{Z_{\tau}(K_1^*, K_2^*)}{e^{3N(K_1^* + K_2^*)}}.$$
(21)

Self duality manifests itself as the same function Z_{τ} appearing on both sides. Taking the logarithmic derivative of both sides with respect to K_1 and dividing both sides by N, we obtain Eq. (19).

On the self-dual line $K_1 = K_1^*$ (equivalently $K_2 = K_2^*$) in the K_1 - K_2 plane, the above relation (19) becomes

$$\sinh 2K_1 \cdot e(K_1, K_2) + e_p(K_1, K_2) = -3 \tanh K_1 \sinh K_1 - 3. \tag{22}$$

This is a weighted sum of the averages of τ_{ij} and τ_{\square} . A further simplification is realized when we impose another condition $K_1 = K_2$ in addition to the self duality, which, together with $K_1 = K_1^*$ and $K_2 = K_2^*$, yields $\sinh 2K_1 = \sinh 2K_2 = 1$, resulting in

$$e(K_c, K_c) + e_p(K_c, K_c) = -3\sqrt{2},$$
 (23)

where $K_c = \frac{1}{2} \ln(\sqrt{2} + 1)$ is the solution to $\sinh 2K_c = 1$. This is the thermal average of the effective Hamiltonian $K_c \sum \tau_{ij} + K_c \sum \tau_{\Box}$ of the Z_2 gauge Higgs model under the probability weight P_{τ} at a special point (K_c, K_c) on the K_1 - K_2 phase diagram.

4. Specific heat and correlation functions

Identities and inequalities can be proven for the specific heat and correlation functions in the subspace $K = K_1$, generalizing the results known in the case of uncorrelated disorder [1, 6].

The specific heat in the subspace $K = K_1$, $C(K, K, K_2)$, is bounded from above as follows.

$$T^{2}C(K, K, K_{2}) = -\frac{\partial E(K, K_{1}, K_{2})}{\partial K} \bigg|_{K_{1}=K}$$

$$= \frac{1}{2^{N}Z_{\tau}(K_{1}, K_{2})} \sum_{\{\tau_{ij}=\pm 1\}} \sum_{\{\sigma_{i}=\pm 1\}} e^{K_{1}\sum \tau_{ij}\sigma_{i}\sigma_{j} + K_{2}\sum \tau_{\square}}$$

$$\times \left\{ \frac{\partial_{K}^{2}\sum_{\{S_{i}\}} e^{K\sum \tau_{ij}S_{i}S_{j}}}{\sum_{\{S_{i}\}} e^{K\sum \tau_{ij}S_{i}S_{j}}} - \left(\frac{\partial_{K}\sum_{\{S_{i}\}} e^{K\sum \tau_{ij}S_{i}S_{j}}}{\sum_{\{S_{i}\}} e^{K\sum \tau_{ij}S_{i}S_{j}}} \right)^{2} \right\} \bigg|_{K_{1}=K}$$

$$\leq \frac{1}{2^{N}Z_{\tau}(K_{1}, K_{2})} \sum_{\{\tau_{ij}=\pm 1\}} e^{K_{2}\sum \tau_{\square}} \frac{\partial^{2}}{\partial K^{2}} \sum_{\{S_{i}=\pm 1\}} e^{K\sum \tau_{ij}S_{i}S_{j}}$$

$$- \left(\frac{1}{Z_{\tau}(K_{1}, K_{2})} \frac{\partial}{\partial K} \sum_{\{\tau_{ij}=\pm 1\}} e^{K\sum \tau_{ij}+K_{2}\sum \tau_{\square}} \right)^{2}$$

$$= -\frac{\partial}{\partial K} E(K, K, K_{2}) < \infty. \tag{24}$$

In deriving the third line from the second, we have replaced the average of the squared quantity $(\cdots)^2$ by the square of the average and have applied the condition $K_1 = K$. The last inequality is based on the analyticity of the average energy for $K(=K_1)$. This inequality means that the specific heat does not diverge in the subspace $K = K_1$ for small $K_2 > 0$, generalizing the result known for uncorrelated disorder [1, 6].

Similarly, we can verify the following relations on correlation functions using the method of gauge transformation as above, generalizing known relations [1, 6],

$$[\langle S_i \rangle_K]_{K_1, K_2} = [\langle S_i \rangle_{K_1} \langle S_i \rangle_K]_{K_1, K_2} \tag{25}$$

$$[\langle S_i \rangle_K]_{K_1, K_2} \le [|\langle \sigma_i \rangle_{K_1}||\langle S_i \rangle_K|]_{K_1, K_2} \le [|\langle \sigma_i \rangle_{K_1}|]_{K_1, K_2} \tag{26}$$

$$\left[\frac{1}{\langle S_i \rangle_{K_1}}\right]_{K_1, K_2} = 1. \tag{27}$$

Here the square brackets $[\cdot\cdot\cdot]_{K_1,K_2}$ denote the average over $\{\tau_{ij}\}$ by the probability $P(\tau,K_1,K_2)$. Notice that Eqs. (25) and (26) hold for any values of K_1,K_2 , and K, not just in the subspace $K=K_1$. The first identity means that there is no spin glass phase, which has $[\langle S_i \rangle_K]_{K_1,K_2} = 0$ and $[\langle S_i \rangle_K^2]_{K_1,K_2} > 0$, when $K=K_1$. It is assumed that boundary spins are fixed to +1 such that the thermal average $\langle S_i \rangle_K$ is finite in the ferromagnetic phase.

5. More general correlations

Analyticity of the average energy holds for a more general class of correlation distributions as long as the invariance property under gauge transformation is satisfied. A simple example is an additional term of the product of two neighboring plaquette variables,

$$P(\tau, K_1, K_2, K_3) = \frac{1}{Z_{\tau}(K_1, K_2, K_3)} \exp\left(K_1 \sum_{\langle ij \rangle} \tau_{ij} + K_2 \sum_{\Box} \tau_{\Box} + K_3 \sum_{\Box_1 \Box_2} \tau_{\Box_1} \tau_{\Box_2}\right), \tag{28}$$

where \Box_1 and \Box_2 are two neighboring plaquettes sharing a bond. It is straightforward to confirm that the proof of Theorem 1 in Appendix A remains valid with minor changes after this modification with sufficiently small K_2 and K_3 .

It is unclear whether or not Theorem 1 holds in the case of infinitely many (in the thermodynamic limit) terms in the exponent of the probability distribution of bond variables, not just a finite number of terms as in Eq. (28). Nevertheless, the following example suggests that similar analyticity may be valid for a large class of probability distributions. Let us consider the probability distribution of correlated disorder,

$$P(\tau, K_1, K_0) = \frac{1}{Z_{\tau}(K_1, K_0)} e^{K_1 \sum \tau_{ij}} Z_{I}(K_0, \{\tau_{ij}\}) = \frac{1}{Z_{\tau}(K_1, K_0)} e^{K_1 \sum \tau_{ij} - F(K_0, \{\tau_{ij}\})}, \tag{29}$$

where $Z_{\rm I}(K_0, \{\tau_{ij}\})$ is the partition function of the Ising model on the same lattice with coupling K_0 and $F(K_0, \{\tau_{ij}\})$ is the corresponding free energy,

$$Z_{I}(K_{0}, \{\tau_{ij}\}) = \sum_{\{\xi_{i} = \pm 1\}} e^{K_{0} \sum_{\langle ij \rangle} \tau_{ij} \xi_{i} \xi_{j}} = e^{-F(K_{0}, \{\tau_{ij}\})}.$$
(30)

The denominator $Z_{\tau}(K_1, K_0)$ is for normalization. The free energy can be expressed in terms of a cluster expansion (high-temperature expansion) for small K_0 , generalizing Eq. (28) to infinitely many (in the thermodynamic limit) gauge invariant terms,

$$-F(K_0, \{\tau_{ij}\}) = K_1(K_0) \sum_{\tau_{\square}} \tau_{\square} + K_2(K_0) \sum_{\text{connected}} \tau_{\square} \tau_{\square} + K_3(K_0) \sum_{\text{connected}} \tau_{\square} \tau_{\square} \tau_{\square} + \cdots,$$
(31)

where the summations run over connected clusters with positive coefficients K_1, K_2, K_3, \cdots [50].

The following result holds for this distribution of correlated disorder.

Theorem 2. For sufficiently small K_0 , the average energy is analytic in K under the distribution function of Eq. (29) of correlated disorder.

Proof. Similarly to Eq. (11), the average energy after gauge transformation is

$$-E(K, K_{1}, K_{0}) = \frac{1}{2^{N} Z_{\tau}(K_{1}, K_{0})} \sum_{\{\tau_{ij}\}} \sum_{\{\sigma_{i}\}} e^{K_{1} \sum \tau_{ij} \sigma_{i} \sigma_{j}} \sum_{\{\xi_{i}\}} e^{K_{0} \sum \tau_{ij} \xi_{i} \xi_{j}} \frac{\partial_{K} \sum_{\{S_{i}\}} e^{K \sum \tau_{ij} S_{i} S_{j}}}{\sum_{\{S_{i}\}} e^{K \sum \tau_{ij} S_{i} S_{j}}}.$$
(32)

When $K = K_1$, the summation over $\{\sigma_i\}$ cancels out with the partition function (summation over $\{S_i\}$) in the denominator,

$$-E(K, K, K_0) = \frac{1}{2^N Z_{\tau}(K, K_0)} \sum_{\{\tau_{ij}\}} \sum_{\{\xi_i\}} e^{K_0 \sum \tau_{ij} \xi_i \xi_j} \frac{\partial}{\partial K} \sum_{\{S_i\}} e^{K \sum \tau_{ij} S_i S_j}.$$
(33)

We can carry out the summation over $\{\tau_{ij}\}$ at each $\langle ij\rangle$ independently,

$$\sum_{\{\tau_{ij}\}} \prod_{\langle ij \rangle} e^{\tau_{ij}(K_0 \xi_i \xi_j + KS_i S_i)} \propto e^{\tilde{K} \sum_{\langle ij \rangle} \xi_i \xi_j S_i S_j}, \tag{34}$$

where $\tanh \tilde{K} = \tanh K_0 \tanh K$. This relation can be verified by inserting all possible values of $\xi_i \xi_j = \pm 1$ and $S_i S_j = \pm 1$. By the gauge transformation $\xi_i S_i \to S_i$ for fixed ξ_i , we find that $E(K,K,K_0)$ is the energy of the ferromagnetic Ising model with uniform coupling \tilde{K} without disorder. Since the effective coupling \tilde{K} runs from 0 to K_0 as K changes from 0 to ∞ , the effective coupling \tilde{K} stays below the critical point of the ferromagnetic Ising model for sufficiently small K_0 . Therefore the energy is analytic in K.

III. CONCLUSION

We have introduced correlations in disorder for the Ising spin glass problem in such a way that the degree of frustration is controlled. We have analyzed the properties of the system in a subspace of the phase diagram in which the average energy is known to be analytic even across a transition point in the case of uncorrelated disorder. We have proved that the analyticity of the average energy is preserved under the introduction of correlations in disorder. This result remains valid for a few more complex types of correlations as long as correlations are gauge invariant. Bounds on the specific heat and identities and inequalities have been derived, generalizing the results for uncorrelated disorder.

Our derivation of these results relies on gauge invariance of correlations. It is difficult to analyze the case without gauge invariance such as a simple product of two neighboring bond variables $\tau_{ij}\tau_{jk}$. It would be safe not to expect similar analyticity to hold in such a case because our result has been derived through a delicate balance of the term in the probability distribution and the partition function, which breaks down for gauge non-invariant probabilities.

For uncorrelated disorder, outstanding properties of the system in the special subspace have been found useful in applications in many fields including statistical inference [37, 51–81], quantum error correction [39, 82–98], quantum Hall effect [99], and localization [100, 101]. The present result may stimulate further developments in these fields in addition to the spin glass theory itself.

Appendix A: Proof of Theorem 1

We prove by a cluster expansion that the following expression in Eq. (12)

$$C_{01}(K_1, K_2) = \frac{\sum_{\{\tau_{ij}\}} \tau_{01} e^{K_1 \sum \tau_{ij} + K_2 \sum (\tau_{\square} + 1)}}{\sum_{\{\tau_{ij}\}} e^{K_1 \sum \tau_{ij} + K_2 \sum (\tau_{\square} + 1)}}$$
(A1)

is analytic in K_1 for sufficiently small $K_2(>0)$.

Notice that the exponent above has an additional term $K_2 \sum 1$ both in the numerator and the denominator compared to Eq. (12) but they cancel out to give the same quantity as in Eq. (12). This term is useful in the proof. The proof below is basically an adaptation of the theory in Ref. [102] to the present context of the Z_2 lattice gauge model. See also Ref. [42], where an abridged description of the theory is provided.

If we define $P(\tau, K_1)$ as $P(\tau, K_1, 0)$, the above Eq. (A1) is expressed as

$$C_{01}(K_1, K_2) = \frac{\sum_{\{\tau_{ij}\}} \tau_{01} P(\tau, K_1) e^{K_2 \sum (\tau_{\square} + 1)}}{\sum_{\{\tau_{ij}\}} P(\tau, K_1) e^{K_2 \sum (\tau_{\square} + 1)}}.$$
(A2)

To show that the small- K_2 expansion of $C_{01}(K_1, K_2)$ converges absolutely and uniformly and thus $C_{01}(K_1, K_2)$ is analytic in K_1 , we introduce ρ_{\square} as

$$e^{K_2(\tau_{\Box}+1)} = 1 + \rho_{\Box}.$$
 (A3)

This ρ_{\square} is positive semi-definite and small for small K_2 . The exponential factor in Eq. (A2) is expanded as

$$e^{K_2 \sum_{\square} (\tau_{\square} + 1)} = \prod_{\square} (1 + \rho_{\square}) = \sum_{Q} \prod_{\square \in Q} \rho_{\square}, \tag{A4}$$

where Q is the set of products of plaquettes. Let us divide Q into Q_1 and Q_2 , where Q_1 is the set of connected products of plaquettes involving the bond (01) and Q_2 is $Q \setminus Q_1$. Then

$$C_{01}(K_1, K_2) = \frac{1}{Z_{\tau}^{+}(K_1, K_2)} \sum_{Q_1} \sum_{Q_2} \sum_{\{\tau_{i,i}\}} \tau_{01} P(\tau, K_1) \prod_{\square \in Q_1} \rho_{\square} \prod_{\square \in Q_2} \rho_{\square}, \tag{A5}$$

where

$$Z_{\tau}^{+}(K_1, K_2) = \sum_{\{\tau_{ij}\}} P(\tau, K_1) e^{K_2 \sum (\tau_{\square} + 1)} = Z_{\tau}(K_1, K_2) e^{K_2 \sum 1}.$$
(A6)

Since Q_1 and Q_2 are disjoint,

$$C_{01}(K_1, K_2) = \frac{1}{Z_{\tau}^+(K_1, K_2)} \sum_{Q_1} \sum_{\tau \in Q_1} \tau_{01} P(\tau \in Q_1, K_1) \Big(\prod_{\square \in Q_1} \rho_{\square} \Big) \cdot Z_{Q_2}(K_1, K_2), \tag{A7}$$

where $Z_{Q_2}(K_1, K_2)$ is the partial partition function,

$$Z_{Q_2}(K_1, K_2) = \sum_{Q_2} \sum_{\tau \in Q_2} P(\tau \in Q_2, K_1) \prod_{\Box \in Q_2} \rho_{\Box}.$$
(A8)

This partial partition function is bounded from above by $Z_{\tau}^+(K_1, K_2)$ because the coupling K_2 for Q_1 is set to 0 in Z_{Q_2} in comparison with the full Z_{τ} and thus the summand is smaller. See Remark 1 of Lemma 3.2 of Ref. [102]. We therefore replace $Z_{Q_2}(K_1, K_2)/Z_{\tau}^+(K_1, K_2)$ by 1 to have a bound

$$C_{01}(K_1, K_2) \leq \sum_{Q_1} \sum_{\tau \in Q_1} P(\tau \in Q_1, K_1) \prod_{\Box \in Q_1} \rho_{\Box}$$

$$\leq \sum_{k} \sum_{|Q_1| = k} \sum_{\tau \in Q_1 : |Q_1| = k} P(\tau \in Q_1, K_1) \prod_{\Box \in Q_1} \rho_{\Box}.$$
(A9)

We have classified Q_1 by the number of elements in it. Using the trivial inequality

$$\rho_{\square} \le e^{2K_2} - 1,\tag{A10}$$

we can replace the average of $\prod_{\square \in Q_1} \rho_{\square}$ with respect to the weight $P(\tau \in Q_1, K_1)$ by the upper bound,

$$\sum_{\tau \in Q_1: |Q_1| = k} P(\tau \in Q_1, K_1) \prod_{\square \in Q_1} \rho_{\square} \le (e^{2K_2} - 1)^k.$$
(A11)

Equation (A9) is then simplified as

$$C_{01}(K_1, K_2) \le \sum_{k} \sum_{|Q_1|=k} (e^{2K_2} - 1)^k.$$
 (A12)

As stated in Lemma 1 below, the number of possible connected graphs involving the bond (01) is bounded as

$$N(k) \le c_1 \cdot c_2^{\ k},\tag{A13}$$

where c_1 and c_2 are positive constants. We therefore obtain

$$C_{01}(K_1, K_2) \le c_1 \sum_k c_2^k (e^{2K_2} - 1)^k.$$
 (A14)

The upper bound of k runs to infinity in the thermodynamic limit, but the summation clearly converges for sufficiently small K_2 . Uniform convergence follows this absolute convergence of an upper bound, proving analyticity of the left-hand side $C_{01}(K_1, K_2)$ as a function of K_1 if we regard K_1 as a complex number in the neighborhood of the origin. \square Remark. The proof remains valid when τ_{01} is replaced by the product of a finite number of bond variables.

Lemma 1. The number of graphs in Q_1 is bounded as in Eq. (A13).

This is Lemma 3.3 of Ref. [102] and is a plaquette version of the well-established bound for the number of connected graphs with variables on the bonds (edges), not on plaquettes, as described, for example, in Exercise 5.3 of Ref. [50]. It is straightforward to replace bonds with plaquettes to derive the above Lemma.

Appendix B: Evaluating p_+ to first order in $\tanh K_2$

According to the definition of Eq. (15), p_+ is written as

$$p_{+} = \frac{\sum_{\{\tau_{ij}\}} \delta_{\tau_{ij},1} e^{K_{1} \sum \tau_{ij}} \prod_{\square} (1 + \tau_{\square} \tanh K_{2})}{\sum_{\{\tau_{ij}\}} e^{K_{1} \sum \tau_{ij}} \prod_{\square} (1 + \tau_{\square} \tanh K_{2})}.$$
 (B1)

The denominator is expanded to first order in $\tanh K_2$ as

$$(\cosh K_1)^{N_B} + N_p \tanh K_2 (\cosh K_1)^{N_B - 4} (\sinh K_1)^4, \tag{B2}$$

where N_B is the number of bonds and N_p is for the number of plaquettes. Similarly, the numerator is

$$(\cosh K_{1})^{N_{B}-1} \sum_{\tau_{ij}} \delta_{\tau_{ij},1} e^{K_{1}\tau_{ij}}$$

$$+ (\cosh K_{1})^{N_{B}-5} \tanh K_{2} (\sinh K_{1})^{4} \sum_{\tau_{ij}} \delta_{\tau_{ij},1} e^{K_{1}\tau_{ij}} (N_{B} - a)$$

$$+ (\cosh K_{1})^{N_{B}-4} \tanh K_{2} (\sinh K_{1})^{3} \sum_{\tau_{ij}} \delta_{\tau_{ij},1} \tau_{ij} e^{K_{1}\tau_{ij}} \cdot a,$$
(B3)

where a the number of plaquettes to which a given single bond belongs. Inserting these expansions to Eq. (B1) and leaving the leading order, we have

$$p_{+} = p \Big(1 - a(2p - 1)^{4} \tanh K_{2} + a(2p - 1)^{3} \tanh K_{2} \Big),$$
(B4)

where we used $(1 - p)/p = e^{-2K_1}$. This is Eq. (16).

^[1] H. Nishimori, Statistical Physics of Spin Glasses and Information Processing: An Introduction (Oxford Univ. Press, 2001).

^[2] M. Mézard, G. Parisi, and M. A. Virasoro, Spin Glass Theory and Beyond (World Scientific, 1987).

^[3] D. Sherrington and S. Kirkpatrick, Solvable model of a spin-glass, Phys. Rev. Lett. 35, 1792 (1975).

^[4] G. Parisi, Infinite number of order parameters for spin-glasses, Phys. Rev. Lett. 43, 1754 (1979).

^[5] M. Talagrand, Spin glasses: a challenge for mathematicians: cavity and mean field models, Vol. 46 (Springer Science & Business Media, 2003).

^[6] H. Nishimori, Internal energy, specific heat and correlation function of the bond-random Ising model, Prog. Theor. Phys. 66, 1169 (1981).

- [7] H. Nishimori, Exact results and critical properties of the Ising model with competing interactions, J, Phys. C 13, 4071 (1980).
- [8] T. Horiguchi and T. Morita, Existence of the ferromagnetic phase in a random-bond Ising model on the square lattice, J. Phys. A 15, L75 (1982).
- [9] C. Garban and T. Spencer, Continuous symmetry breaking along the Nishimori line, arXiv:2109.01617 (2021).
- [10] S. F. Edwards and P. W. Anderson, Theory of spin glasses, J. Phys. F 5, 965 (1975).
- [11] J. Mydosh, Spin Glasses: An Experimental Introduction (Taylor & Francis, 1993).
- [12] J. A. Hoyos, N. Laflorencie, A. P. Vieira, and T. Vojta, Protecting clean critical points by local disorder correlations, Epl 93, 30004 (2011).
- [13] V. Bonzom, R. Gurau, and M. Smerlak, Universality in p-spin glasses with correlated disorder, J. Stat. Mech. 2013, L02003 (2013).
- [14] A. G. Cavaliere and A. Pelissetto, Disordered ising model with correlated frustration, J. Phys. A 52, 174002 (2019).
- [15] L. Münster, C. Norrenbrock, A. K. Hartmann, and A. P. Young, Ordering behavior of the two-dimensional Ising spin glass with long-range correlated disorder, Phys. Rev. E 103, 042117 (2021).
- [16] J. B. Kogut, An introduction to lattice gauge theory and spin systems, Rev. Mod. Phys. 51, 659 (1979).
- [17] D. C. Mattis, Solvable spin systems with random interactions, Phys. Lett. A 56, 421 (1976).
- [18] J. J. Hopfield, Neural networks and physical systems with emergent collective computational abilities, Proc, Nat. Acad. Sci. USA 79, 2554 (1982).
- [19] D. J. Amit, H. Gutfreund, and H. Sompolinsky, Storing infinite numbers of patterns in a spin-glass model of neural networks, Phys. Rev. Lett. 55, 1530 (1985).
- [20] D. J. Amit, H. Gutfreund, and H. Sompolinsky, Spin-glass models of neural networks, Phys. Rev. A 32, 1007 (1985).
- [21] D. J. Amit, H. Gutfreund, and H. Sompolinsky, Statistical mechanics of neural networks near saturation, Ann. Phys. 67, 30 (1987).
- [22] We can consider other possibilities. For example, on the two-dimensional triangular lattice, the product will run over three bond variables around a unit triangle, $\tau_{ij}\tau_{ji}\tau_{ki}$. We work on the hypercubic lattice in the present paper for simplicity.
- [23] A. Georges, D. Hansel, P. L. Doussal, and J. P. Bouchaud, Exact properties of spin glasses. II. Nishimori's line: new results and physical implications, J. Phys. France 46, 1827 (1985).
- [24] P. Le Doussal and A. B. Harris, Location of the Ising spin-glass multicritical point on Nishimori's line, Phys. Rev. Lett. **61**, 625 (1988).
- [25] P. Le Doussal and A. B. Harris, ϵ expansion for the Nishimori multicritical point of spin glasses, Phys. Rev. B **40**, 9249 (1989).
- [26] Y. Iba, The Nishimori line and Bayesian statistics, J. Phys. A 32, 3875 (1998).
- [27] I. A. Gruzberg, N. Read, and A. W. W. Ludwig, Random-bond Ising model in two dimensions: The Nishimori line and supersymmetry, Phys. Rev. B 63, 104422 (2001).
- [28] A. Honecker, M. Picco, and P. Pujol, Universality class of the Nishimori point in the 2d $\pm J$ random-bond Ising model, Phys. Rev. Lett. 87, 047201 (2001).
- [29] F. D. Nobre, Phase diagram of the two-dimensional $\pm J$ Ising spin glass, Phys. Rev. E 64, 046108 (2001).
- [30] F. Merz and J. T. Chalker, Two-dimensional random-bond Ising model, free fermions, and the network model, Phys. Rev. B 65, 054425 (2002).
- [31] M. Hasenbusch, F. P. Toldin, A. Pelissetto, and E. Vicari, Multicritical Nishimori point in the phase diagram of the $\pm J$ Ising model on a square lattice, Phys. Rev. E 77, 051115 (2008).
- [32] H. Kitatani, Griffiths inequalities for ising spin glasses on the Nishimori line, J. Phys. Soc. Jpn. 78, 044714 (2009).
- [33] C. Yamaguchi, Percolation thresholds of the Fortuin-Kasteleyn cluster for the Edwards-Anderson Ising model on complex networks: — Analytical results on the Nishimori line —, Prog. Theor. Phys. 124, 399 (2010).
- [34] F. Krzakala and L. Zdeborová, On melting dynamics and the glass transition. II. Glassy dynamics as a melting process., J. Chem. Phys. 134, 034513 (2011).
- [35] M. Ohzeki, Fluctuation theorems on the Nishimori line, Phys. Rev. E 86, 061110 (2012).
- [36] Y. Sasagawa, H. Ueda, J. Genzor, A. Gendiar, and T. Nishino, Entanglement entropy on the boundary of the square-lattice ±J Ising model, J. Phys. Soc. Jpn. 89, 114005 (2020).
- [37] D. Alberici, F. Camilli, P. Contucci, and E. Mingione, The multi-species mean-field spin-glass on the Nishimori line, J. Stat. Phys. 182, 2 (2021).
- [38] D. Alberici, F. Camilli, P. Contucci, and E. Mingione, The solution of the deep Boltzmann machine on the Nishimori line, Commun. Math. Phys. 387, 1191 (2021).
- [39] E. Dennis, A. Kitaev, A. Landahl, and J. Preskill, Topological quantum memory, J. Math. Phys. 43, 4452 (2002).
- [40] Y. Ozeki and H. Nishimori, Phase diagram and critical exponents of the $\pm J$ Ising model in finite dimensions by Monte Carlo renormalization group, J. Phys. Soc. Jpn. **56**, 1568 (1987).
- [41] Notice that the general expression of the energy of Eq. (7) is also regarded as the expectation value of a function of $\{\tau_{ij}\}$, $\langle H \rangle_K(\{\tau_{ij}\})$, with respect to the gauge Higgs probability weight P_{τ} . However, this $\langle H \rangle_K(\{\tau_{ij}\})$ is a very complicated function of the bond variables $\{\tau_{ij}\}$ involving all types of products and summations of $\{\tau_{ij}\}$, which makes it hard to analyze its properties.
- [42] E. Fradkin and S. H. Shenker, Phase diagrams of lattice gauge theories with Higgs fields, Phys. Rev. D 19, 3682 (1979).
- [43] M. Creutz, Phase diagrams for coupled spin-gauge systems, Phys. Rev. D 21, 1006 (1980).
- [44] G. A. Jongeward, J. D. Stack, and C. Jayaprakash, Monte carlo calculations on Z₂ gauge-higgs theories, Phys. Rev. D 21, 3360 (1980).

- [45] M. Creutz, L. Jacobs, and C. Rebbi, Monte carlo computations in lattice gauge theories, Phys. Rep. 95, 201 (1983).
- [46] L. Genovese, F. Gliozzi, A. Rago, and C. Torrero, The phase diagram of the three-dimensional z_2 gauge higgs system at zero and finite temperature, Nucl. Phys. B Proc. Suppl. 119, 894 (2003), Proceedings of the XXth International Symposium on Lattice Field Theory.
- [47] I. S. Tupitsyn, A. Kitaev, N. V. Prokof'ev, and P. C. E. Stamp, Topological multicritical point in the phase diagram of the toric code model and three-dimensional lattice gauge Higgs model, Phys. Rev. B 82, 085114 (2010).
- [48] F. J. Wegner, Duality in generalized Ising models and phase transitions without local order parameters, J. Math. Phys. 12, 2259 (1971).
- [49] This duality equivalence to the two-dimensional ferromagnetic Ising model in a field also implies the impossibility of the exact closed-form formula of the average energy.
- [50] S. Friedli and Y. Velenik, Statistical Mechanics of Lattice Systems (Cambridge Univ. Press, 2017).
- [51] P. Ruján, Finite temperature error-correcting codes, Phys. Rev. Lett. 70, 2968 (1993).
- [52] N. Sourlas, Spin glasses, error-correcting codes and finite-temperature decoding, EPL 25, 159 (1994).
- [53] T. Murayama, Y. Kabashima, D. Saad, and R. Vicente, Statistical physics of regular low-density parity-check error-correcting codes, Phys. Rev. E 62, 1577 (2000).
- [54] Y. Kabashima, T. Murayama, and D. Saad, Typical performance of Gallager-type error-correcting codes, Phys. Rev. Lett. 84, 1355 (2000).
- [55] Y. Kabashima, T. Murayama, and D. Saad, Cryptographical properties of Ising spin systems, Phys. Rev. Lett. 84, 2030 (2000).
- [56] A. Montanari, The glassy phase of Gallager codes, Eur. Phys. J. B 23, 121 (2001).
- [57] Y. Kabashima, N. Sazuka, K. Nakamura, and D. Saad, Tighter decoding reliability bound for Gallager's error-correcting code, Phys. Rev. E 64, 046113 (2001).
- [58] K. Tanaka, Statistical-mechanical approach to image processing, J. Phys. A 35, R81 (2002).
- [59] Franz, M. Leone, A. Montanari, and F. Ricci-Tersenghi, Dynamic phase transition for decoding algorithms, Phys. Rev. E 66, 046120 (2002).
- [60] Y. Kabashima, A CDMA multiuser detection algorithm on the basis of belief propagation, J. Phys. A 36, 11111 (2003).
- [61] N. Macris, On the relation between map and bp gexit functions of low density parity check codes, in 2006 IEEE Information Theory Workshop - ITW '06 Punta del Este (2006) pp. 312–316.
- [62] N. Macris, A useful tool in the theory of error correcting codes, IEEE Trans. Inf. Theor. 53, 664 (2007).
- [63] A. Decelle, F. Krzakala, C. Moore, and L. Zdeborová, Asymptotic analysis of the stochastic block model for modular networks and its algorithmic applications, Phys. Rev. E 84, 066106 (2011).
- [64] A. Manoel and R. Vicente, Statistical mechanics of reputation systems in autonomous networks, J. Stat. Mech. 2013, P08002 (2013).
- [65] F. Caltagirone, L. Zdeborová, and F. Krzakala, On convergence of approximate message passing, in 2014 IEEE International Symposium on Information Theory (2014) pp. 1812–1816.
- [66] Y. Xu, Y. Kabashima, and L. Zdeborová, Bayesian signal reconstruction for 1-bit compressed sensing, J, Stat. Mech. 2014, P11015 (2014).
- [67] T. Lesieur, F. Krzakala, and L. Zdeborová, Phase transitions in sparse PCA, in 2015 IEEE International Symposium on Information Theory (ISIT) (2015) pp. 1635–1639.
- [68] H. Huang and T. Toyoizumi, Unsupervised feature learning from finite data by message passing: Discontinuous versus continuous phase transition, Phys. Rev. E **94**, 062310 (2016).
- [69] L. Zdeborová and F. Krzakala, Statistical physics of inference: thresholds and algorithms, Adv. Phys. 65, 453 (2016), 1511.02476.
- [70] H. Huang, Statistical mechanics of unsupervised feature learning in a restricted Boltzmann machine with binary synapses, J. Stat. Mech. 2017, 053302 (2017).
- [71] T. Lesieur, F. Krzakala, and L. Zdeborová, Constrained low-rank matrix estimation: phase transitions, approximate message passing and applications, J. Stat. Mech. 2017, 073403 (2017).
- [72] T. Kawamoto, Algorithmic detectability threshold of the stochastic block model, Phys. Rev. E 97, 032301 (2018).
- [73] B. Aubin, A. Maillard, J. Barbier, F. Krzakala, N. Macris, and L. Zdeborová, The committee machine: computational to statistical gaps in learning a two-layers neural network, J. Stat. Mech. **2019**, 124023 (2019).
- [74] F. Antenucci, F. Krzakala, P. Urbani, and L. Zdeborová, Approximate survey propagation for statistical inference, J. Stat. Mech. **2019**, 023401 (2019).
- [75] J. Kadmon and S. Ganguli, Statistical mechanics of low-rank tensor decomposition, J. Stat. Mech. 2019, 124016 (2019).
- [76] T. Murayama, A. Saito, and P. Davis, Rate distortion theorem and the multicritical point of a spin glass, Phys. Rev. E 102, 042122 (2020).
- [77] U. Vasiliy and E. Sergey, Hyper neural network as the diffeomorphic domain for short code soft decision beyound belief propagation decoding problem, in 2020 IEEE East-West Design Test Symposium (EWDTS) (2020) pp. 1–6.
- [78] T. Hou and H. Huang, Statistical physics of unsupervised learning with prior knowledge in neural networks, Phys. Rev. Lett. 124, 248302 (2020).
- [79] L. Dall'Amico, R. Couillet, and N. Tremblay, Nishimori meets Bethe: a spectral method for node classification in sparse weighted graphs, J. Stat. Mech. 2021, 093405 (2021).
- [80] S. Kawaguchi, Spread-spectrum watermarking model using a parity-check code for simultaneous restoration of message and image, J. Phys. Soc. Jpn. 90, 104003 (2021).

- [81] S. Arai, M. Ohzeki, and K. Tanaka, Mean field analysis of reverse annealing for code-division multiple-access multiuser detection, Phys. Rev. Research 3, 033006 (2021).
- [82] C. Wang, J. Harrington, and J. Preskill, Confinement-Higgs transition in a disordered gauge theory and the accuracy threshold for quantum memory, Ann. Phys. 303, 31 (2002).
- [83] H. G. Katzgraber, H. Bombin, and M. A. Martin-Delgado, Error threshold for color codes and random three-body Ising models, Phys. Rev. Lett. 103, 090501 (2009).
- [84] H. G. Katzgraber, H. Bombin, R. S. Andrist, and M. A. Martin-Delgado, Topological color codes on union jack lattices: a stable implementation of the whole Clifford group, Phys. Rev. A 81, 012319 (2010).
- [85] T. M. Stace and S. D. Barrett, Error correction and degeneracy in surface codes suffering loss, Phys. Rev. A 81, 022317 (2010).
- [86] R. S. Andrist, H. G. Katzgraber, H. Bombin, and M. a. Martin-Delgado, Tricolored lattice gauge theory with randomness: fault tolerance in topological color codes, New J. Phys. 13, 083006 (2011).
- [87] H. Bombin, R. S. Andrist, M. Ohzeki, H. G. Katzgraber, and M. A. Martin-Delgado, Strong resilience of topological codes to depolarization, Phys. Rev. X 2, 021004 (2012).
- [88] K. Fujii, Y. Nakata, M. Ohzeki, and M. Murao, Measurement-based quantum computation on symmetry breaking thermal states, Phys. Rev. Lett. 110, 120502 (2013).
- [89] K. Fujii, M. Negoro, N. Imoto, and M. Kitagawa, Measurement-free topological protection using dissipative feedback, Phys. Rev. X 4, 041039 (2014).
- [90] R. S. Andrist, J. R. Wootton, and H. G. Katzgraber, Error thresholds for Abelian quantum double models: Increasing the bit-flip stability of topological quantum memory, Phys. Rev. A 91, 042331 (2015).
- [91] P. Iyer and D. Poulin, Hardness of decoding quantum stabilizer codes, IEEE Transactions on Information Theory 61, 5209 (2015).
- [92] A. Kubica, M. E. Beverland, F. Brandão, J. Preskill, and K. M. Svore, Three-dimensional color code thresholds via statistical-mechanical mapping, Phys. Rev. Lett. 120, 180501 (2018).
- [93] A. A. Kovalev, S. Prabhakar, I. Dumer, and L. P. Pryadko, Numerical and analytical bounds on threshold error rates for hypergraph-product codes, Phys. Rev. A 97, 062320 (2018).
- [94] M. Li, D. Miller, M. Newman, Y. Wu, and K. R. Brown, 2d compass codes, Phys. Rev. X 9, 021041 (2019).
- [95] C. Vuillot, H. Asasi, Y. Wang, L. P. Pryadko, and B. M. Terhal, Quantum error correction with the toric Gottesman-Kitaev-Preskill code, Phys. Rev. A 99, 032344 (2019).
- [96] M. H. Zarei and A. Ramezanpour, Noisy toric code and random-bond Ising model: The error threshold in a dual picture, Phys. Rev. A 100, 062313 (2019).
- [97] O. Viyuela, S. Vijay, and L. Fu, Scalable fermionic error correction in Majorana surface codes, Phys. Rev. B 99, 205114 (2019).
- [98] C. T. Chubb and S. Flammia, Statistical mechanical models for quantum codes with correlated noise, Ann. Inst. Henri Poincare 8, 629 (2021).
- [99] N. Read and A. W. W. Ludwig, Absence of a metallic phase in random-bond Ising models in two dimensions: Applications to disordered superconductors and paired quantum Hall states, Phys. Rev. B 63, 024404 (2000).
- [100] T. Senthil and M. P. A. Fisher, Quasiparticle localization in superconductors with spin-orbit scattering, Phys. Rev. B 61, 9690 (2000).
- [101] D. Vodola, M. Rispler, S. Kim, and M. Müller, Fundamental thresholds of realistic quantum error correction circuits from classical spin models, arXiv:2104.04847 (2021).
- [102] K. Osterwalder and E. Seiler, Gauge field theories on a lattice, Ann. Phys. 110, 440 (1978).