

CONSTRUCTION OF p -ENERGY AND ASSOCIATED ENERGY MEASURES ON SIERPIŃSKI CARPETS

RYOSUKE SHIMIZU

ABSTRACT. We establish the existence of a scaling limit \mathcal{E}_p of discrete p -energies on the graphs approximating a generalized Sierpiński carpet for $p > d_{\text{ARC}}$, where d_{ARC} is the Ahlfors regular conformal dimension of the underlying generalized Sierpiński carpet. Furthermore, the function space \mathcal{F}_p defined as the collection of functions with finite p -energies is shown to be a reflexive and separable Banach space that is dense in the set of continuous functions with respect to the supremum norm. In particular, $(\mathcal{E}_2, \mathcal{F}_2)$ recovers the canonical regular Dirichlet form constructed by Barlow and Bass [5] or Kusuoka and Zhou [54]. We also provide \mathcal{E}_p -energy measures associated with the constructed p -energy and investigate its basic properties like self-similarity and chain rule.

1. INTRODUCTION

On Euclidean spaces, the *nonlinear potential theory* is built on the theory of the $(1, p)$ -Sobolev spaces $W^{1,p}$ and the p -energy $\int |\nabla f|^p dx$. The main aim of this paper is to construct and study p -energies on Sierpiński carpets as a prototype of nonlinear potential theory on complicated metric spaces like “fractals” (see also [41, Problem 7.6]). There has been significant progress on “analysis and probability” on complicated spaces beyond Euclidean spaces over the last several decades. The earlier works are the constructions of diffusion processes, which is called the *Brownian motions*, on self-similar sets in 1980s and 1990s. (For details and precise history of “analysis on fractals”, see the ICM survey of Kumagai [52] for example.) In particular, a class of self-similar sets called *generalized Sierpiński carpets* (see Figure 1), is one of the successful examples. In this introduction, we restrict to the case of *the standard Sierpiński carpet* (the left in Figure 1), SC for short, for simplicity. The first Brownian motion on the SC was given by Barlow and Bass in [5], where they obtained the Brownian motion as a scaling limit of Brownian motions on Euclidean regions approximating the SC. From an analytic viewpoint, the result of Barlow and Bass gives 2-energy \mathcal{E}_2 and the associated $(1, 2)$ -“Sobolev” space \mathcal{F}_2 , namely *regular Dirichlet form* on the SC. Recall that a tuple of 2-energy $\int |\nabla f|^2 dx$ (on $L^2(\mathbb{R}^N, dx)$) and $(1, 2)$ -Sobolev space $W^{1,2}$ is a typical example of regular Dirichlet forms, which corresponds to the classical Brownian motion on \mathbb{R}^N . Although it is difficult to define the gradient ∇f on the SC, we can say that a suitable 2-energy “ $\int |\nabla f|^2 dx$ ” exists on the SC. Later, Kusuoka and Zhou [54]

Date: March 26, 2023.

2020 Mathematics Subject Classification. Primary 28A80, 30L99, 31E99; Secondary 46E36, 31C45.

Key words and phrases. Sierpiński carpet, p -energy, p -energy measure, nonlinear potential theory.

The author was supported in part by JSPS KAKENHI Grant Number JP20J20207.




FIGURE 1. Sierpiński carpet (left), two other generalized Sierpiński carpets and Menger sponge (right)

gave an alternative construction of a regular Dirichlet form as a scaling limit of discrete 2-energies on a series of graphs approximating the SC as shown in Figure 2. Our work gives a “canonical” construction of p -energy \mathcal{E}_p and the associated $(1, p)$ -“Sobolev” space \mathcal{F}_p on the SC, which play the same roles as the pair of $\int |\nabla f|^p dx$ and the Sobolev space $W^{1,p}$, by extending and simplifying the method of Kusuoka and Zhou.

Let us describe briefly our strategy to construct $(\mathcal{E}_p, \mathcal{F}_p)$ on the SC. We write (K, d, μ) to denote the SC as a metric measure space, that is, K is the Sierpiński carpet, d is the Euclidean metric of \mathbb{R}^2 and μ is the $\dim_{\mathbb{H}}(K, d)$ -dimensional Hausdorff measure on (K, d) , where $\dim_{\mathbb{H}}(K, d) = \log 8 / \log 3$ is the Hausdorff dimension of (K, d) . Let $\{G_n\}_{n \geq 1}$ be a series of finite graphs approximating the SC whose edge set is denoted by E_n (see Figure 2 and Definition 2.9). Then discrete p -energy $\mathcal{E}_p^{G_n}$ on G_n is

$$\mathcal{E}_p^{G_n}(f) = \frac{1}{2} \sum_{(x,y) \in E_n} |M_n f(x) - M_n f(y)|^p,$$

where M_n is a discretization operator from $L^p(K, \mu)$ to \mathbb{R}^{G_n} (see Section 2 for the precise definition). To obtain an appropriate non-trivial limit of discrete p -energies, some renormalization is necessary (see [4] for example). We will see that the behavior of $\mathcal{R}_p^{(n)}$ defined as

$$\mathcal{R}_p^{(n)} := \left(\inf \left\{ \mathcal{E}_p^{G_n}(M_n f) \mid \begin{array}{l} f \in L^p(K, \mu) \text{ with } M_n f \equiv 0 \text{ on the left side of } G_n \\ \text{and } M_n f \equiv 1 \text{ on the right side} \end{array} \right\} \right)^{-1}$$

gives us the proper renormalization constant of discrete p -energies $\mathcal{E}_p^{G_n}$. In fact, for $p = 2$, Barlow and Bass [6] have proved that there exist constants $\rho_2 > 0$ (the so-called *resistance scaling factor*) and $C \geq 1$ such that

$$(1.1) \quad C^{-1} \rho_2^n \leq \mathcal{R}_2^{(n)} \leq C \rho_2^n, \quad n \in \mathbb{N}.$$

What Kusuoka and Zhou have shown is that, roughly speaking, the Dirichlet form $(\mathcal{E}_2, \mathcal{F}_2)$ on the SC is obtained as

$$\mathcal{F}_2 = \left\{ f \in L^2(K, \mu) \mid \sup_{n \geq 1} \rho_2^n \mathcal{E}_2^{G_n}(M_n f) < \infty \right\}$$

and $\mathcal{E}_2(f) = \lim_{k \rightarrow \infty} \rho_2^{n_k} \mathcal{E}_2^{G_{n_k}}(M_{n_k} f)$ for some subsequence $\{n_k\}_{k \geq 1}$.

By using p -combinatorial modulus, which is one of fundamental tools in “quasi-conformal geometry”, Bourdon and Kleiner [15] have generalized (1.1), i.e. they




FIGURE 2. Graphical approximation $\{G_n\}_{n \geq 1}$ of the SC (This figure draws G_1 and G_2 in blue)

have ensured the existence of a constant $\rho_p > 0$ such that

$$(1.2) \quad C^{-1} \rho_p^n \leq \mathcal{R}_p^{(n)} \leq C \rho_p^n, \quad n \in \mathbb{N}.$$

Then our $(1, p)$ -“Sobolev” space \mathcal{F}_p equipped with the norm $\|\cdot\|_{\mathcal{F}_p}$ is defined by

$$\mathcal{F}_p = \left\{ f \in L^p(K, \mu) \mid \sup_{n \geq 1} \rho_p^n \mathcal{E}_p^{G_n}(M_n f) < \infty \right\},$$

and

$$\|f\|_{\mathcal{F}_p} = \|f\|_{L^p} + \left(\sup_{n \geq 1} \rho_p^n \mathcal{E}_p^{G_n}(M_n f) \right)^{1/p}.$$

Under the following assumption (see Assumption 4.7):

$$(1.3) \quad 1 < p < \infty \text{ and } \rho_p > 1,$$

we will prove that \mathcal{F}_p is continuously embedded in the Hölder space:

$$\mathcal{C}^{0, \theta_p} = \left\{ f: K \rightarrow \mathbb{R} \mid \sup_{x \neq y \in K} \frac{|f(x) - f(y)|}{d(x, y)^{\theta_p}} < \infty \right\},$$

where $\theta_p := \log \rho_p / p \log 3$ (Theorem 5.1). This embedding result is very powerful. Indeed, we will deduce the *closedness*, i.e. $(\mathcal{F}_p, \|\cdot\|_{\mathcal{F}_p})$ is a Banach space, and the *regularity*, i.e. \mathcal{F}_p is dense in $\mathcal{C}(K) = \{f: K \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$ with the sup norm, from this embedding (see Theorems 5.2 and 5.5).

Moreover, the *separability* of $(\mathcal{F}_p, \|\cdot\|_{\mathcal{F}_p})$ will be deduced from the *reflexivity* of \mathcal{F}_p (Theorems 5.9 and 5.10). Thanks to the separability, one easily sees that, by the diagonal procedure, a subsequential limit $\lim_{k \rightarrow \infty} \rho_p^{n_k} \mathcal{E}_p^{G_{n_k}}(M_{n_k} f)$ exists for all $f \in \mathcal{F}_p$. Our final object \mathcal{E}_p called the p -energy on the SC will be constructed through these subsequential limits¹.

The assumption (1.3) is essential for the continuous embedding of \mathcal{F}_p in the Hölder space $\mathcal{C}^{0, \theta_p}$ and has a close connection with the *Ahlfors regular conformal dimension* $\dim_{\text{ARC}}(K, d)$ which is defined by

$$(1.4) \quad \dim_{\text{ARC}}(K, d) = \inf \left\{ \alpha \mid \begin{array}{l} \text{there exists a metric } \rho \text{ on } K \text{ which is} \\ \text{quasisymmetric to } d \text{ and } \alpha\text{-Ahlfors regular} \end{array} \right\}.$$

¹To construct “canonical” p -energy \mathcal{E}_p on the SC, we need to follow some additional procedures as shown in the work of Kusuoka and Zhou. In this paper, we will introduce new graphs $\{G_n\}_{n \geq 1}$ and consider discrete p -energies on them to get a “good” p -energy. These procedures are described in Section 6. See Theorem 2.23 for the meaning of canonical p -energies.

(For the precise definitions of *Ahlfors regularity* and being *quasisymmetric*, see (2.2) and Definition 4.7.) Indeed, by results of Carrasco Piaggio [20] and Kigami [47], the condition (1.3) is equivalent to

$$(1.5) \quad p > \dim_{\text{ARC}}(K, d).$$

We expect that this condition (1.5) represents a “low-dimensional” phase. More precisely, we regard the Hölder embedding $\mathcal{F}_p \hookrightarrow \mathcal{C}^{0, \theta_p}$ as a generalization of the classical Sobolev embedding (a consequence of Morrey’s inequality). For this reason, we naturally arrive at the following conjecture.

Conjecture 1.1. $\dim_{\text{ARC}}(K, d) = \inf\{p \mid \mathcal{F}_p \text{ is embedded in a subset of } \mathcal{C}(K)\}$.

To show this conjecture, what we need is the regularity of \mathcal{F}_p , i.e. the density of $\mathcal{F}_p \cap \mathcal{C}(K)$ in $\mathcal{C}(K)$ with the sup norm, for $p \leq \dim_{\text{ARC}}(K, d)$ (see [7] for $p = 2$). This is a big open problem for future work.

Besides our “Sobolev spaces” \mathcal{F}_p , there has already been an established theory of “Sobolev spaces on metric measure spaces” based on the notion of *upper gradients*, which is a counter part of $|\nabla f|$ introduced by Heinonen and Koskela in [32]. We refer to [31, 33] for details. From the viewpoint of this theory, our $(1, p)$ -“Sobolev” space \mathcal{F}_p can be seen as a *fractional Korevaar–Schoen Sobolev space*. Indeed, we will give the following representation of \mathcal{F}_p (Theorem 2.22):

$$(1.6) \quad \mathcal{F}_p = \left\{ f \in L^p(K, \mu) \mid \overline{\lim}_{r \downarrow 0} \int_K \int_{B_d(x, r)} \frac{|f(x) - f(y)|^p}{r^{\beta_p}} d\mu(y) d\mu(x) < \infty \right\},$$

where $\beta_p = \log(8\rho_p)/\log 3$. When $p = 2$, this result is well-known (see [28, 51, 53] for example) and the parameter β_2 is called the *walk dimension*. For detailed expositions of β_2 , see [28, 52, 53] for example. If $\beta_p = p$, then the expression (1.6) coincides with (a slight modification of) the Korevaar–Schoen $(1, p)$ -Sobolev space [49, 50]. However, it is well-known that a strict inequality $\beta_2 > 2$ holds on the SC (see [7, Proposition 5.1] or [39]). This phenomenon suggests that the existing theory of “Sobolev spaces on metric measure spaces” do not give any non-trivial $(1, p)$ -Sobolev spaces on the SC². This is one of the reasons why we try to provide an alternative theory of $(1, p)$ -“Sobolev” space and p -energy on the SC.

Another major objective of this paper is the \mathcal{E}_p -energy measures associated with p -energy \mathcal{E}_p . In terms of a Dirichlet form $(\mathcal{E}_2, \mathcal{F}_2)$, \mathcal{E}_2 -energy measure of a function $f \in \mathcal{F}_2$ is defined as the unique Borel measure $\mu_{\langle f \rangle}^2$ on K such that

$$(1.7) \quad \int_K g d\mu_{\langle f \rangle}^2 = \mathcal{E}_2(f, fg) - \frac{1}{2}\mathcal{E}_2(f^2, g), \quad g \in \mathcal{F}_2.$$

(Note that we can define the form $\mathcal{E}_2(f, g)$ by the polarization: $\mathcal{E}_2(f, g) := \frac{1}{4}(\mathcal{E}_2(f + g) - \mathcal{E}_2(f - g))$.) This measure plays the role of $|\nabla f(x)|^2 dx$ if the underlying space is Euclidean. On the other hand, for any $f \in \mathcal{F}_2$ with $\mathcal{E}_2(f) \neq 0$, the \mathcal{E}_2 -energy measure $\mu_{\langle f \rangle}^2$ and the $\log_3 8$ -dimensional Hausdorff measure μ on the SC are mutually singular due to the fact that $\beta_2 > 2$ by a result of Hino [35]. See [40] for an extension of this fact to general metric measure Dirichlet spaces. This

²It is also well-known that the *Newtonian $(1, p)$ -Sobolev space* on the SC becomes $L^p(K, \mu)$ due to the lack of plenty rectifiable curves in the SC. See [55, Proposition 4.3.3] and [33, Proposition 7.1.33] for example.

phenomenon is also far different from “smooth” settings and motivates the study of \mathcal{E}_2 -energy measures on fractals.

For general p , due to the lack of a counterpart of the expression in the right-hand side of (1.7), we will choose to generalize Hino’s alternative method of the construction of \mathcal{E}_2 -energy measure. Namely, for any $f \in \mathcal{F}_p$, we first construct a measure $\mathfrak{m}_{\langle f \rangle}^p$ on the shift space $\{1, \dots, 8\}^{\mathbb{N} \cup \{0\}}$ associated with the SC and define our \mathcal{E}_p -energy measure $\mu_{\langle f \rangle}^p$ as the pushforward measure of $\mathfrak{m}_{\langle f \rangle}^p$ under the natural quotient map $\pi: \{1, \dots, 8\}^{\mathbb{N} \cup \{0\}} \rightarrow K$ (see Proposition 2.6 for a description of π), i.e. $\mu_{\langle f \rangle}^p(A) = \mathfrak{m}_{\langle f \rangle}^p(\pi^{-1}(A))$ for any Borel set A of K . Then our \mathcal{E}_p -energy measure $\mu_{\langle f \rangle}^p$ is associated with \mathcal{E}_p in the sense that $\mu_{\langle f \rangle}^p(K) = \mathcal{E}_p(f)$ (for more details on relations between $\mu_{\langle f \rangle}^p$ and \mathcal{E}_p , see Theorem 2.25-(c)).

Furthermore, we will show the *chain rule*: for any $\Phi \in \mathcal{C}^1(\mathbb{R})$,

$$(1.8) \quad d\mu_{\langle \Phi \circ f \rangle}^p = |\Phi' \circ f|^p d\mu_{\langle f \rangle}^p.$$

When $p = 2$, the chain rule (1.8) is proved by using integral expressions of \mathcal{E}_2 (see [25, (3.2.12)] for example), but such representations take full advantage of the fact that $p = 2$. Alternatively, we prove (1.8) by introducing a new series of graphs $\{\mathbb{G}_n\}_{n \geq 1}$ (see the beginning of subsection 6.1), which is embedded in the SC, and analyzing discrete p -energies $\{\mathcal{E}_p^{\mathbb{G}_n}(\Phi \circ f)\}_{n \geq 1}$. This approach is actually valid since our p -energies are based on subsequential limits of $\{\rho_p^n \mathcal{E}_p^{\mathbb{G}_n}\}_{n \geq 1}$.

The first result on the existence of suitable p -energy on fractals is due to [34], where the Sierpiński gaskets are considered. (Added in revision: for *p.c.f. self-similar sets*, there are also recent studies [9, 18, 26].) In the very recent paper [44], Kigami has established a theory of $(1, p)$ -Sobolev space and p -energy on *p-conductively homogeneous* compact metric spaces (see [44] for details). His paper [44] includes new construction results even if $p = 2$ (see [44, Sections 12 and 13] for a gallery). Also, a class of highly symmetric p.c.f. self-similar sets called *nested fractals* is also treated in [44, Section 14]. However, the construction of \mathcal{E}_p -energy measures associated with the p -energy \mathcal{E}_p is not treated in earlier works.

Outline. This paper is organized as follows. In Section 2, we prepare basic frameworks in this paper and state the main results. In particular, we give the definition of generalized Sierpiński carpets. Sections 3 and 4 are devoted to extending results of Kusuoka and Zhou to fit our purpose. Section 3 is a collection of basic estimates of (p, p) -Poincaré constants and $\mathcal{R}_p^{(n)}$. In Section 4, we prove powerful results concerning (p, p) -Poincaré constants (uniform Hölder estimates and a condition called p -Knight Move (KM_p) for example) under Assumption 4.7 and finish all preparations to construct p -energy \mathcal{E}_p and $(1, p)$ -“Sobolev” space \mathcal{F}_p . Section 5 is devoted to investigating detailed properties of \mathcal{F}_p . Then, in Section 6, we introduce another graphical approximation $\{\mathbb{G}_n\}_{n \geq 1}$ and construct a canonical p -energy \mathcal{E}_p (see Theorem 2.21 for the precise meaning of ‘canonical’). Section 7 is devoted to discussions on \mathcal{E}_p -energy measures. Finally, in Section 8, we prove $\beta_p > p$ (Theorem 2.27) under the assumption that the underlying generalized Sierpiński carpet is embedded in \mathbb{R}^2 . The appendix contains proofs of some elementary lemmas.

Notation. In this paper, we use the following notation and conventions.

- (1) $\mathbb{N} := \{n \in \mathbb{Z} \mid n > 0\}$ and $\mathbb{Z}_{\geq 0} := \mathbb{N} \cup \{0\}$.
- (2) We set $a \vee b := \max\{a, b\}$, $a \wedge b := \min\{a, b\}$ for $a, b \in [-\infty, \infty]$.

- (3) For any countable set V , we define $\mathbb{R}^V := \{f \mid f: V \rightarrow \mathbb{R}\}$.
- (4) For $f: \mathbb{R} \rightarrow \mathbb{R}$, define $\text{Lip}(f) := \sup_{x \neq y \in \mathbb{R}} \frac{|f(x) - f(y)|}{|x - y|}$.
- (5) Let X be a compact topological space. We set $\mathcal{C}(X) := \{f: X \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$ and write its sup norm by $\|f\|_{\mathcal{C}(X)} := \sup_{x \in X} |f(x)|$.
- (6) Let X be a topological space and let A be a subset of X . The topological boundary of A is denoted by ∂A , that is $\partial A := \overline{A}^X \setminus \text{int}_X A$.
- (7) Let (X, d) be a metric space. The open ball with center $x \in X$ and radius $r > 0$ is denoted by $B_d(x, r)$, that is,

$$B_d(x, r) := \{y \in X \mid d(x, y) < r\}.$$

If the metric d is clear in context, then we write $B(x, r)$ for short.

- (8) Let K be a compact metrizable space and let $\mathcal{B}(K)$ denote the Borel σ -algebra of K . Let μ be a Borel (regular) measure on K . For any $A \in \mathcal{B}(K)$ with $\mu(A) > 0$ and $f \in L^1(K, \mu)$, we define

$$\int_A f d\mu := \frac{1}{\mu(A)} \int_A f d\mu.$$

- (9) We use \sqcup to denote disjoint unions.
- (10) Let $D \in \mathbb{N}$. Set $\mathbf{0} = \mathbf{0}^D := (0)_{k=1}^D \in \mathbb{R}^D$ and $\mathbf{e}_j = \mathbf{e}_j^D := (\delta_{k,j})_{k=1}^D \in \mathbb{R}^D$ for each $j \in \{1, \dots, D\}$, where $\delta_{j,k}$ is the Dirac delta. For $x = (x_k)_{k=1}^D, y = (y_k)_{k=1}^D \in \mathbb{R}^D$, we write $|x - y|_{\mathbb{R}^D} = \left(\sum_{k=1}^D |x_k - y_k|^2\right)^{1/2}$.

Acknowledgments This paper is mainly originated from the author's doctoral thesis at Kyoto University. The author would like to express his deepest gratitude toward Professor Jun Kigami for guidance during his graduate course, reading drafts, and stimulating conversations. He would also like to thank Professor Naotaka Kajino for giving him several valuable comments. In particular, Prof. Kajino has suggested using the approach in [39] to prove Theorem 2.27 and given him detailed comments about the results in [8]. The author would also like to thank Professors Mathav Murugan and Toshiyuki Tanaka for their careful readings of an earlier version of the manuscript. In particular, Prof. Murugan has pointed out a gap in the previous proof of Theorem 5.1. Finally, he would like to thank Professor Fabrice Baudoin for a comment regarding the critical Besov exponent in [3] (Remark 8.8) and anonymous referees for helpful comments and suggestions.

2. PRELIMINARY AND RESULTS

2.1. Generalized Sierpiński carpets and graphical approximations. We start with the definition of *generalized Sierpiński carpets* and related notations. The reader is referred to [46] for further background and more general framework, namely, self-similar structure.

Let $D, a \in \mathbb{N}$ with $D \geq 2, a \geq 3$ and set $Q_0 := [-1, 1]^D$. Let $S \subsetneq \{0, 1, \dots, a-1\}^D$ be non-empty, define $f_i: \mathbb{R}^D \rightarrow \mathbb{R}^D$ by $f_i(x) := a^{-1}x + 2a^{-1} \cdot i - (a-1)a^{-1} \sum_{j=1}^D \mathbf{e}_j$ for each $i \in \{0, 1, \dots, a-1\}^D$ and set $Q_1 := \bigcup_{i \in S} f_i(Q_0)$, so that $Q_1 \subsetneq Q_0$. Let K be the *self-similar set* associated with $\{f_i\}_{i \in S}$, i.e., the unique non-empty compact subset of \mathbb{R}^D such that $K = \bigcup_{i \in S} f_i(K)$, which exists and satisfies $K \subsetneq Q_0$ thanks to $Q_1 \subsetneq Q_0$ by [46, Theorem 1.1.4]. Define $F_i := f_i|_K$ for each $i \in S$ and $\text{GSC}(D, a, S) := (K, S, \{F_i\}_{i \in S})$.

By following [39], we will introduce the notion of *generalized Sierpiński carpets*. The following definition is essentially due to Barlow and Bass [7, Section 2]. The non-diagonality condition in [7, Hypotheses 2.1] has been modified later in [8]. See [8, Remark 2.10-1.] for details of this correction.

Definition 2.1 (Generalized Sierpiński carpet, [8, Subsection 2.2]). $\text{GSC}(D, a, S)$ is called a *generalized Sierpiński carpet* if and only if the following four conditions are satisfied:

- (GSC1) (Symmetry) $f(Q_1) = Q_1$ for any isometry f of \mathbb{R}^D with $f(Q_0) = Q_0$.
- (GSC2) (Connectedness) Q_1 is connected.
- (GSC3) (Non-diagonality) $\text{int}_{\mathbb{R}^D}(Q_1 \cap \prod_{k=1}^D [2(i_k - \varepsilon_k)a^{-1}, 2(i_k + 1)a^{-1}])$ is either empty or connected for any $(i_k)_{k=1}^D \in \mathbb{Z}^D$ and any $(\varepsilon_k)_{k=1}^D \in \{0, 1\}^D$.
- (GSC4) (Borders included) $[-1, 1] \times \{1\}^{D-1} \subset Q_1$.

Remark 2.2. In [8, 39], generalized Sierpiński carpets are defined as subspaces of D -dimensional unit cube $[0, 1]^D$. In this paper, we consider GSC as subspaces of $[-1, 1]^D$ instead of $[0, 1]^D$ to follow [44, Section 11].

As special cases of Definition 2.1, the *standard Sierpiński carpet* (left in Figure 1) and *Menger sponge* (right in Figure 1), are given by $\text{GSC}(2, 3, \{0, 1, 2\}^2 \setminus \{(1, 1)\})$ and $\text{GSC}(3, 3, \{(i_1, i_2, i_3) \in \{0, 1, 2\}^3 \mid \sum_{k=1}^3 \mathbf{1}_1(i_k) \leq 1\})$ respectively.

In this paper, we suppose that $\text{GSC}(D, a, S) = (K, S, \{F_i\}_{i \in S})$ is a generalized Sierpiński carpet and that $d: K \times K \rightarrow [0, +\infty)$ is the normalized Euclidean metric on K , i.e. $d(x, y) = \frac{1}{2\sqrt{D}} |x - y|_{\mathbb{R}^D}$.

Next, by following [39, 44, 46], we introduce useful notations to express the symmetries of $(K, S, \{F_i\}_{i \in S})$ and to describe the topological structure as a self-similar set of $(K, S, \{F_i\}_{i \in S})$.

Definition 2.3. Define

$$B_{j, \sigma} = \{(x_1, \dots, x_D) \in Q_0 \mid x_j = \sigma\}$$

for $j \in \{1, \dots, D\}$ and $\sigma \in \{-1, 0, +1\}$. We also define the hyperplane

$$\mathcal{H}_{j_1, j_2}^+ = \{(x_1, \dots, x_D) \in \mathbb{R}^D \mid x_{j_1} = x_{j_2}\},$$

and

$$\mathcal{H}_{j_1, j_2}^- = \{(x_1, \dots, x_D) \in \mathbb{R}^D \mid x_{j_1} = -x_{j_2}\},$$

for $j_1, j_2 \in \{1, \dots, D\}$ with $j_1 \neq j_2$. Moreover, define

$$\mathcal{H}_{j_1, j_2}^{+, \leq} = \{(x_1, \dots, x_D) \in \mathbb{R}^D \mid x_{j_1} \leq x_{j_2}\}$$

and

$$\mathcal{H}_{j_1, j_2}^{+, \geq} = \{(x_1, \dots, x_D) \in \mathbb{R}^D \mid x_{j_1} \geq x_{j_2}\}.$$

We also define $\mathcal{H}_{j_1, j_2}^{-, \leq}$ and $\mathcal{H}_{j_1, j_2}^{-, \geq}$ in similar ways.

Definition 2.4. We define

$$\mathbb{B}_D = \{T \mid T: \mathbb{R}^D \rightarrow \mathbb{R}^D \text{ is an isometry such that } T(Q_0) = Q_0\},$$

and

$$(2.1) \quad \mathcal{G}_0 = \{f|_K \mid f \in \mathbb{B}_D\}.$$

Then \mathcal{G}_0 is a finite subgroup of the set of homeomorphism of K by virtue of (GSC1). Furthermore, define $R_j \in \mathbb{B}_D$ as the reflection in the hyperplane $B_{j,0}$ for each

$j \in \{1, \dots, D\}$ and define $R_{j_1, j_2}^\pm \in \mathbb{B}_D$ as the (restriction of the) reflection in the hyperplane $\mathcal{H}_{j_1, j_2}^\pm$ for each $j_1, j_2 \in \{1, \dots, D\}$. We also use the same symbols R_j and R_{j_1, j_2}^\pm to denote these restrictions to K , which are elements of \mathcal{G}_0 .

Definition 2.5. (1) We set $W_m := S^m = \{w_1 \cdots w_m \mid w_i \in S \text{ for } i \in \{1, \dots, m\}\}$ for $m \in \mathbb{N}$ and $W_\# := \bigcup_{m=1}^\infty W_m$. For $w = w_1 \cdots w_m \in W_\#$, the unique $m \in \mathbb{N}$ with $w \in W_m$ is denoted by $|w|$ and set $f_w := f_{w_1} \circ \cdots \circ f_{w_m}$, $F_w := F_{w_1} \circ \cdots \circ F_{w_m}$, $K_w := F_w(K)$, $O_w := K_w \setminus \bigcup_{v \in W_m; v \neq w} (K_w \cap K_v)$, and $[w]_n := w_1 \cdots w_n$ for $n \in \{1, \dots, m\}$. We define $W_0 := \{\emptyset\}$ and $[w]_0 := \emptyset$, where \emptyset is an empty word. Set $W_* = W_0 \cup W_\#$ and $F_\emptyset := \text{id}_{\mathbb{R}^D}|_K$. We also set $i^m := i \cdots i \in W_m$ for each $i \in S$. For $n, m \geq 0$, $v = v_1 \cdots v_n \in W_n$ and $w = w_1 \cdots w_m \in W_m$, define $v \cdot w \in W_{n+m}$ by $v \cdot w = v_1 \cdots v_n w_1 \cdots w_m$. We also write vw for $v \cdot w$ if there is no confusion. For $n \in \mathbb{N}$ and non-empty subset A of W_n , we define $A \cdot W_m$ by setting

$$A \cdot W_m = \{vw \mid v \in A, w \in W_m\}.$$

When $A = \{v\}$ for some $v \in W_n$, we write $v \cdot W_m$ to denote $\{v\} \cdot W_m$ for simplicity.

(2) The collection of one-sided infinite sequences of symbols S is denoted by Σ , that is,

$$\Sigma = \{\omega = \omega_1 \omega_2 \omega_3 \cdots \mid \omega_i \in S \text{ for any } i \in \mathbb{N}\},$$

which is called the *one-sided shift space* of symbols S . We define the *shift map* $\sigma : \Sigma \rightarrow \Sigma$ by $\sigma(\omega_1 \omega_2 \cdots) = \omega_2 \omega_3 \cdots$ for each $\omega_1 \omega_2 \cdots \in \Sigma$. The branches of σ are denoted by σ_i ($i \in S$), namely $\sigma_i : \Sigma \rightarrow \Sigma$ is defined as $\sigma_i(\omega_1 \omega_2 \cdots) = i \omega_1 \omega_2 \cdots$ for each $i \in S$ and $\omega_1 \omega_2 \cdots \in \Sigma$. For $w = w_1 \cdots w_m \in W_\#$, we write $\sigma_w = \sigma_{w_1} \circ \cdots \circ \sigma_{w_m}$ and $\Sigma_w := \sigma_w(\Sigma)$. For $\omega = \omega_1 \omega_2 \cdots \in \Sigma$ and $m \in \mathbb{Z}_{\geq 0}$, we define $[\omega]_m = \omega_1 \cdots \omega_m \in W_m$.

(3) For any $w = w_1 \cdots w_m \in W_m$, we define $\xi(w) = w_1 \cdots w_{m-1}$. For a subset $A \subseteq W_m$, we write $\xi(A)$ for $\{\xi(w) \mid w \in A\}$.

(4) For $A \subseteq \mathbb{R}^D$ and $n \in \mathbb{Z}_{\geq 0}$, define

$$W_n[A] = \{w \in W_n \mid A \cap K_w \neq \emptyset\}.$$

(5) For $T \in \mathcal{G}_0$ and $n \in \mathbb{Z}_{\geq 0}$, $\tau[T] : W_n \rightarrow W_n$ is the bijection such that $T(K_w) = K_{\tau[T](w)}$ for any $w \in W_n$.

We consider Σ as a topological space equipped with the product topology of $S^\mathbb{N}$. Then the following fact is elemental (see [46, Theorem 1.2.3]).

Proposition 2.6. *For any $\omega = \omega_1 \omega_2 \cdots \in \Sigma$, the set $\bigcap_{m \geq 1} K_{[\omega]_m}$ contains only one point. If we define $\pi : \Sigma \rightarrow K$ by $\{\pi(\omega)\} = \bigcap_{m \geq 1} K_{[\omega]_m}$, then π is a continuous surjective map. Furthermore, it holds that $\pi \circ \sigma_i = F_i \circ \pi$ for each $i \in S$.*

Set $N_* := \#S$ and $\alpha := \log N_* / \log a$. Note that $\alpha < D$ by $S \subsetneq \{0, 1, \dots, a-1\}^D$. Let μ be the self-similar probability measure on K with weight $(1/N_*, \dots, 1/N_*)$, namely μ is the unique Borel probability measure on K such that $\mu = N_*(\mu \circ F_i)$ for any $i \in S$. It is known that α is the Hausdorff dimension of (K, d) and that μ is a constant multiple of the α -dimensional Hausdorff measure on (K, d) ; see [46, Proposition 1.5.8 and Theorem 1.5.7] for example. In particular, d is α -Ahlfors regular, that is, there exists a constant $C_{\text{AR}} \geq 1$ such that

$$(2.2) \quad C_{\text{AR}}^{-1} r^\alpha \leq \mu(B(x, r)) \leq C_{\text{AR}} r^\alpha,$$

for any $x \in K$ and $r \in (0, 1)$. The following lemma on the self-similar measure μ is standard (see [39, Lemma 3.3] for example).

Lemma 2.7. *Let $w \in W_{\#}$ and let $f: K \rightarrow [-\infty, \infty]$ be Borel measurable. Then*

$$\int_K |f \circ F_w| d\mu = N_*^{|w|} \int_{K_w} |f| d\mu \quad \text{and} \quad \int_{K_w} |f \circ F_w^{-1}| d\mu = N_*^{-|w|} \int_K |f| d\mu.$$

Now, we define some operators that are frequently used in this paper.

Definition 2.8. Let $p \in [1, \infty)$. For $w \in W_{\#}$, we define F_w^* , $(F_w)_*$: $L^p(K, \mu) \rightarrow L^p(K, \mu)$ by setting

$$F_w^* f := f \circ F_w, \quad (F_w)_* f := \begin{cases} f \circ F_w^{-1} & \text{on } K_w, \\ 0 & \text{on } K \setminus K_w, \end{cases}$$

for each $f \in L^p(K, \mu)$. For $n \in \mathbb{N}$, define $M_n: L^p(K, \mu) \rightarrow \mathbb{R}^{W_n}$ by setting

$$M_n f(w) := \int_{K_w} f d\mu = N_*^n \int_{K_w} f d\mu, \quad w \in W_n,$$

for each $f \in L^p(K, \mu)$.

Note that, from Lemma 2.7, $M_n f(w) = \int_K F_w^* f d\mu$ for any $f \in L^p(K, \mu)$, which implies that, for $m \in \mathbb{N}$ and $v \in W_m$,

$$(2.3) \quad M_n(F_v^* f)(w) = \int_K F_w^*(F_v^* f) d\mu = \int_K F_{vw}^* f d\mu = M_{n+m} f(vw).$$

We introduce graphical approximations of K and related notations by following [47] and [44, section 2].

Definition 2.9. We define $G_n := (W_n, E_n)$ by setting

$$E_n := \{(v, w) \mid v, w \in W_n \text{ with } v \neq w \text{ and } K_v \cap K_w \neq \emptyset\}.$$

(This series of graphs G_n is called the horizontal networks in [47].) We also define $\tilde{G}_n = (W_n, \tilde{E}_n)$ by

$$\tilde{E}_n := \{(v, w) \in E_n \mid F_v(Q_0) \cap F_w(Q_0) \text{ is a } (D-1)\text{-dimensional hypercube}\}.$$

We use d_{G_n} to denote the graph distance of G_n . By (GSC2), the graph G_n is connected. Furthermore, by virtue of (GSC3), (W_n, \tilde{E}_n) is also connected (see [37, Proposition 2.5]). Moreover, by [37, Theorem 2.6], the following result holds.

Proposition 2.10. *Let $m \in \mathbb{N}$ and let $v, w \in W_m$ satisfy $K_v \cap K_w \neq \emptyset$. Then it holds that $d_{\tilde{G}_m}(v, w) \leq D$.*

For $n, m, k \in \mathbb{N}$ and $w \in W_m$, we define a subset $\mathcal{B}_n(w, k)$ of W_{n+m} by setting

$$\mathcal{B}_n(w, k) := \bigcup_{v \in W_m; d_{G_m}(v, w) \leq k} v \cdot W_n.$$

For each $x \in K$ and $s > 0$, we also define a subset $U_1(x, s)$ of K by setting

$$(2.4) \quad U_1(x, s) := \bigcup_{w \in \Lambda_{s,1}(x)} K_w,$$

where

$$\Lambda_{s,1}(x) = \left\{ w \mid \begin{array}{l} w \in W_n \text{ with } x \in K_w \text{ or } (v, w) \in E_n \text{ for some } v \in W_n \\ \text{s.t. } x \in K_v, \text{ where } n \in \mathbb{Z}_{\geq 0} \text{ with } a^{-n} \leq s < a^{-n+1}. \end{array} \right\}$$

(See [47, Definition 2.3.6].) Then the following proposition says that generalized Sierpiński carpets equipped with the (normalized) Euclidean metrics satisfy the *basic framework* of [47]. See also [44, Assumption 2.15 and Proposition 11.4].

Proposition 2.11. *Let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet. Then the following properties hold:*

(0) (minimal, strongly finite) for any $m \geq 0$ and $w \in W_m$,

$$O_w := K_w \setminus \left(\bigcup_{v \in W_m \setminus \{w\}} K_v \right) \neq \emptyset.$$

Furthermore, if we define

$$(2.5) \quad L_* := \sup_{w \in W_\#} \#\{v \in W_{|w|} \mid (v, w) \in E_{|w|}\},$$

then $L_* \leq 3^D - 1$;

(1) for any $w \in W_*$, K_w is connected;

(2A) (1-adapted) there exists a constant C_{AD} (depending only on a, D) such that for any $x \in K$ and $s \in (0, 1]$,

$$(2.6) \quad U_1(x, C_{\text{AD}}^{-1}s) \subseteq B_d(x, s) \subseteq U_1(x, C_{\text{AD}}s);$$

(2B) for any $m \geq 0$ and $w \in W_m$, $\text{diam}(K_w, d) = a^{-m}$;

(2C) (thick) for any $n \geq 1$ and $w \in W_n$, there exists $x \in K$ such that

$$K_w \supseteq B_d\left(x, \frac{1}{\sqrt{D}}a^{-n}\right);$$

(3) for $m \geq 0$ and $v, w \in W_m$ with $v \neq w$,

$$\mu(K_v \cap K_w) = 0 \quad \text{and} \quad \mu(K_v) = N_*^{-m} = a^{-\alpha m};$$

(4) for any $m \in \mathbb{N}$ and $w \in W_m$, it holds that

$$\xi\left(\{v \in W_m \mid d_{G_m}(v, w) \leq 1\}\right) \subseteq \{z \in W_{m-1} \mid d_{G_{m-1}}(z, \xi(w)) \leq 1\}.$$

Proof. As mentioned in [44, Proposition 11.4], all statements can be easily verified. Indeed, (0), (1), (2B), (2C) and (4) are immediate from the definition of generalized Sierpiński carpets. A proof of (3) can be found in [39, Lemma 3.2] for example. Finally, the condition (2A) follows by noting that

$$(2.7) \quad \inf_{x \in K_v, y \in K_w} d(x, y) \geq \frac{1}{\sqrt{D}}a^{-m} \quad \text{if } m \geq 1, v, w \in W_m \text{ satisfy } K_v \cap K_w = \emptyset.$$

□

An intrinsic boundary $\partial_* G_n$ of the graph G_n is the set of words that the associated n -cells intersect with the topological boundary of $[-1, 1]^D$, that is,

$$\partial_* G_n := W_n \left[\partial[-1, 1]^D \right] = \left\{ w \in W_n \mid K_w \cap \partial[-1, 1]^D \neq \emptyset \right\}.$$

We have the following proposition (see [44, Assumption 2.10 and Proposition 2.16]).

Proposition 2.12. *Let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet. Then $W_k \setminus \partial_* G_k \neq \emptyset$ for any $k \geq 2$.*

Proof. By (GSC1) and (GSC4), we have

$$i_* := (0)_{k=1}^D \in S \quad \text{and} \quad \widehat{i}_* := (a-1)_{k=1}^D \in S.$$

Then we easily see that $i_* \widehat{i}_* \in W_2 \setminus \partial_* G_2$.

□

We conclude this subsection by giving another aspect of (2.6). Define

$$n(x, y) := \max \left\{ m \in \mathbb{Z}_{\geq 0} \mid \begin{array}{l} \text{there exist } v, w \in W_m \text{ such that} \\ x \in K_v, y \in K_w \text{ and } K_v \cap K_w \neq \emptyset \end{array} \right\},$$

where we set $K_\emptyset := K$.

Lemma 2.13. *Let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet. Then*

$$\frac{1}{a\sqrt{D}} a^{-n(x,y)} \leq d(x, y) \leq 2a^{-n(x,y)}, \quad x, y \in K.$$

Proof. Let $x, y \in K$ and let $n = n(x, y)$. The definition of $n(x, y)$ immediately implies that there exist $v, w \in W_n$ such that $x \in K_v, y \in K_w$ and $K_v \cap K_w \neq \emptyset$. Hence

$$d(x, y) \leq \text{diam}(K_v \cup K_w, d) \leq 2a^{-n}.$$

We next prove the converse inequality. Let $v, w \in W_{n+1}$ such that $x \in K_v$ and $y \in K_w$. Then, by the definition of $n(x, y)$, we have $K_v \cap K_w = \emptyset$. Therefore,

$$d(x, y) \geq \inf_{x' \in K_v, y' \in K_w} d(x', y') \geq \frac{1}{\sqrt{D}} a^{-(n+1)},$$

where we used (2.7). This completes the proof. \square

2.2. p -energies and Poincaré constants on finite graphs. In this subsection, we review some basic results and definitions in discrete nonlinear potential theory and introduce (p, p) -Poincaré constants that will play essential roles in this paper.

Let $G = (V, E)$ be a directed, connected, simple finite graph, and let $p > 0$. We always suppose that $(x, y) \in E$ if and only if $(y, x) \in E$.

Definition 2.14. For $f: V \rightarrow \mathbb{R}$, we define its p -energy $\mathcal{E}_p^G(f)$ by setting

$$\mathcal{E}_p^G(f) := \frac{1}{2} \sum_{(x,y) \in E} |f(x) - f(y)|^p.$$

Definition 2.15. For disjoint subsets A, B of V , we define their p -conductance $\mathcal{C}_p^G(A, B)$ by setting

$$\mathcal{C}_p^G(A, B) := \inf \{ \mathcal{E}_p^G(f) \mid f|_A \equiv 1, f|_B \equiv 0 \}.$$

For a given subset A of V , define

$$E^A := \{(x, y) \in E \mid x, y \in A\},$$

and

$$\mathcal{E}_{p,A}(f) := \frac{1}{2} \sum_{(x,y) \in E^A} |f(x) - f(y)|^p.$$

To clarify the underlying graph, we also write $\mathcal{E}_{p,A}^G(f)$ for $\mathcal{E}_{p,A}(f)$. We also set

$$\bar{A} := \{x \in V \mid x \in A \text{ or } (x, y) \in E \text{ for some } y \in A\},$$

and $\partial A := \bar{A} \setminus A$.

Then the following monotonicity of p -conductance is immediate (see [57, Proposition 3.7-(2)] for example).

Proposition 2.16. *Let $A, B, A', B' \subseteq V$ with $A \subseteq A'$ and $B \subseteq B'$. Then $\mathcal{C}_p^G(A, B) \leq \mathcal{C}_p^G(A', B')$.*

The following property states the Markov property of discrete p -energy. (This naming is borrowed from the case $p = 2$.) This is also immediate from the definition.

Proposition 2.17. *Let $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ with $\text{Lip}(\varphi) \leq 1$. Then $\mathcal{E}_p^G(\varphi \circ f) \leq \mathcal{E}_p^G(f)$ for any $f: V \rightarrow \mathbb{R}$. In particular, if we define $f^\# := (f \vee 0) \wedge 1$, then $\mathcal{E}_p^G(f^\#) \leq \mathcal{E}_p^G(f)$.*

Next we define some types of (p, p) -Poincaré constants. Let ν be a non-negative measure on V , and let $\partial_* G \subsetneq V$ be a given non-empty subset.

Definition 2.18. For a non-empty subset A of V and $f: A \rightarrow \mathbb{R}$, define its mean $\langle f \rangle_{A, \nu}$ by setting

$$\langle f \rangle_{A, \nu} := \frac{1}{\sum_{x \in A} \nu(x)} \sum_{x \in A} f(x) \nu(x).$$

We define $\lambda_p^{(G, \nu)}$ on (G, ν) by setting

$$\lambda_p^{(G, \nu)} := \sup \left\{ \frac{\sum_{x \in V} |f(x) - \langle f \rangle_{V, \nu}|^p \nu(x)}{\mathcal{E}_p^G(f)} \mid f \in \mathbb{R}^V, \mathcal{E}_p^G(f) \neq 0 \right\}.$$

We consider its Dirichlet boundary conditioned version $\lambda_{p, \text{Dir}}^{(G, \nu)}(\partial_* G)$ defined as

$$\lambda_{p, \text{Dir}}^{(G, \nu)}(\partial_* G) := \sup \left\{ \frac{|\langle f \rangle_{V, \nu}|^p}{\mathcal{E}_p^G(f)} \mid f \in \mathbb{R}^V, \mathcal{E}_p^G(f) \neq 0 \text{ and } f|_{\partial_* G} \equiv 0 \right\}.$$

For disjoint subsets A, B of V , we also define

$$\sigma_p^{(G, \nu)}(A, B) := \sup \left\{ \frac{|\langle f \rangle_{A, \nu} - \langle f \rangle_{B, \nu}|^p}{\mathcal{E}_{p, A \cup B}(f)} \mid f \in \mathbb{R}^{A \cup B}, \mathcal{E}_{p, A \cup B}(f) \neq 0 \right\}.$$

By standard arguments in calculus of variations (see [57, proof of Lemma 3.3] for example), one can easily prove the following proposition.

Proposition 2.19. *Suppose that $p > 1$ and that G is connected. Let A, B be non-empty disjoint subsets of V , and let $\partial_* G \subsetneq V$ be non-empty.*

- (1) *There exists a unique $f \in \mathbb{R}^V$ such that $f|_A \equiv 1$, $f|_B \equiv 0$ and $\mathcal{E}_p^G(f) = \mathcal{C}_p^G(A, B)$.*
- (2) *There exists a unique $f \in \mathbb{R}^V$ such that $f|_{\partial_* G} \equiv 0$, $\langle f \rangle_{V, \nu} = 1$ and $\mathcal{E}_p^G(f)^{-1} = \lambda_{p, \text{Dir}}^{(G, \nu)}(\partial_* G)$.*
- (3) *If both A and B are connected, then there exists $f \in \mathbb{R}^{A \cup B}$ such that*

$$\left| \langle f \rangle_{A, \nu} - \langle f \rangle_{B, \nu} \right| = 1 \quad \text{and} \quad \mathcal{E}_{p, A \cup B}(f)^{-1} = \sigma_p^{(G, \nu)}(A, B).$$

Moreover, such f is unique up to an additive constant and to the multiplication by -1 .

We conclude this subsection by introducing notations of these quantities in specific settings. We mainly consider p -conductance and (p, p) -Poincaré constants on approximating graphs G_m introduced in subsection 2.1. Note that, by the self-similarity of $\{K_w\}_w$, each subgraph $(w \cdot W_m, E^{w \cdot W_m})$ is a copy of G_m for any $w \in W_\#$ and $m \in \mathbb{N}$. Recall that μ denotes the self-similar probability measure on K with weight $(1/N_*, \dots, 1/N_*)$. We consider that μ is also a measure on W_n by

setting $\mu(w) := \mu(K_w) = N_*^{-n}$ for each $w \in W_n$. Then, for any subset A of W_n and $f: A \rightarrow \mathbb{R}$,

$$\langle f \rangle_{A,\mu} = \frac{1}{\#A} \sum_{w \in A} f(w),$$

and thus we write $\langle f \rangle_A$ to denote $\langle f \rangle_{A,\mu}$ for simplicity. For $w \in W_\#$ and $n \in \mathbb{N}$, we define

$$\mathcal{C}_p^{(n)} := \sup_{w \in W_\#} \mathcal{C}_p^{G_{n+|w|}}(w \cdot W_n, W_{n+|w|} \setminus \mathcal{B}_n(w, 1)),$$

and $\mathcal{R}_p^{(n)} := (\mathcal{C}_p^{(n)})^{-1}$. We also set $\lambda_p^{(n)} := \lambda_p^{(G_n, \mu)}$ and $\lambda_{p,\text{Dir}}^{(n)} := \lambda_{p,\text{Dir}}^{(G_n, \mu)}(\partial_* G_n)$. Finally, for $v, w \in W_\#$ with $|v| = |w|$, define

$$\sigma_p^{(n)}(v, w) := \sigma_p^{(G_{|v|+n}, \mu)}(v \cdot W_n, w \cdot W_n),$$

and

$$\sigma_p^{(n)} := \sup_{m \geq 1} \max_{(v,w) \in \bar{E}_m} \sigma_p^{(n)}(v, w).$$

Remark 2.20. Our definitions of Poincaré constants are slightly changed from the original definitions adopted in [54]. Indeed, $N_*^{-n} \lambda_2^{(n)}$ in our notation is the same as λ_n in [54]. The situations are the same for other Poincaré constants $\sigma_2^{(n)}, \lambda_{2,\text{Dir}}^{(n)}$.

2.3. Main results. Now, we are ready to state the main results of this paper. Let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet. Then, for $p > 0$, it is well-known that there exists $\rho_p > 0$ such that $\lim_{n \rightarrow \infty} (\mathcal{C}_p^{(n)})^{1/n} = \rho_p^{-1}$ (see Theorem 3.4).

The following two theorems state detailed properties of our $(1, p)$ -“Sobolev” space \mathcal{F}_p on (K, d, μ) .

Theorem 2.21. *Assume that $p > \dim_{\text{ARC}}(K, d)$. Then a function space \mathcal{F}_p defined as*

$$\mathcal{F}_p := \left\{ f \in L^p(K, \mu) \mid \sup_{n \geq 1} \rho_p^n \mathcal{E}_p^{G_n}(M_n f) < \infty \right\}$$

is a reflexive and separable Banach space equipped with a norm $\|\cdot\|_{\mathcal{F}_p}$ defined by

$$\|f\|_{\mathcal{F}_p} := \|f\|_{L^p} + \left(\sup_{n \geq 1} \rho_p^n \mathcal{E}_p^{G_n}(M_n f) \right)^{1/p}.$$

Moreover, \mathcal{F}_p is continuously embedded in a Hölder space $\mathcal{C}^{0,(\beta_p - \alpha)/p}$ on K , where $\beta_p := \log(N_ \rho_p) / \log a$ and*

$$\mathcal{C}^{0,(\beta_p - \alpha)/p} := \left\{ f \in \mathcal{C}(K) \mid \sup_{x \neq y \in K} \frac{|f(x) - f(y)|}{d(x, y)^{(\beta_p - \alpha)/p}} < \infty \right\}.$$

Furthermore, \mathcal{F}_p is dense in $\mathcal{C}(K)$ with respect to the supremum norm.

Theorem 2.22 (Theorem 5.15). *Assume that $p > \dim_{\text{ARC}}(K, d)$. Let β_p be the same constant as in Theorem 2.21. Then \mathcal{F}_p has the following expression:*

$$(2.8) \quad \mathcal{F}_p = \left\{ f \in L^p(K, \mu) \mid \overline{\lim}_{r \downarrow 0} \int_K \int_{B_d(x, r)} \frac{|f(x) - f(y)|^p}{r^{\beta_p}} d\mu(y) d\mu(x) < \infty \right\}.$$

Note that we will show that $\beta_p \geq p$ for any $p > 0$ in Proposition 3.5.

In Section 6, we construct a “canonical” p -energy \mathcal{E}_p on (K, d, μ) , which satisfies the following properties. For the definition of Clarkson’s inequality, see Definition 5.6.

Theorem 2.23. *Assume that $p > \dim_{\text{ARC}}(K, d)$. Then there exists a functional $\mathcal{E}_p: \mathcal{F}_p \rightarrow [0, \infty)$ such that $\mathcal{E}_p(\cdot)^{1/p}$ is a semi-norm satisfying Clarkson’s inequality and the associated norm $\|\cdot\|_{\mathcal{E}_p} := \|\cdot\|_{L^p} + \mathcal{E}_p(\cdot)^{1/p}$ is equivalent to $\|\cdot\|_{\mathcal{F}_p}$. Furthermore, $(\mathcal{E}_p, \mathcal{F}_p)$ satisfies the following conditions:*

- (1) $\mathbf{1}_K \in \mathcal{F}_p$, and, for $f \in \mathcal{F}_p$, $\mathcal{E}_p(f) = 0$ if and only if f is constant. Furthermore, $\mathcal{E}_p(f + a\mathbf{1}_K) = \mathcal{E}_p(f)$ for any $f \in \mathcal{F}_p$ and $a \in \mathbb{R}$;
- (2) (Regularity) \mathcal{F}_p is dense in $\mathcal{C}(K)$ with respect to the sup norm;
- (3) (Markov property) if $f \in \mathcal{F}_p$ and $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ with $\text{Lip}(\varphi) \leq 1$, then $\varphi \circ f \in \mathcal{F}_p$ and $\mathcal{E}_p(\varphi \circ f) \leq \mathcal{E}_p(f)$;
- (4) (Symmetry) if $f \in \mathcal{F}_p$ and $T \in \mathcal{G}_0$, then $f \circ T \in \mathcal{F}_p$ and $\mathcal{E}_p(f \circ T) = \mathcal{E}_p(f)$;
- (5) (Self-similarity) it holds that

$$(2.9) \quad \mathcal{F}_p = \{f \in \mathcal{C}(K) \mid f \circ F_i \in \mathcal{F}_p \text{ for all } i \in S\}$$

and, for every $f \in \mathcal{F}_p$,

$$(2.10) \quad \mathcal{E}_p(f) = \rho_p \sum_{i \in S} \mathcal{E}_p(f \circ F_i);$$

- (6) (Strong locality) if $f, g \in \mathcal{F}_p$ satisfy $\text{supp}[f] \cap \text{supp}[g - a\mathbf{1}_K] = \emptyset$ for some $a \in \mathbb{R}$, then $\mathcal{E}_p(f + g) = \mathcal{E}_p(f) + \mathcal{E}_p(g)$.

Remark 2.24. When $p = 2$, there exists the unique Dirichlet form (up to constant multiples) satisfying all conditions (1)-(5) by [8, Theorem 1.2], [36, Proposition 5.1] and [38, Proposition 5.9]³. This is the reason why we say that a p -energy \mathcal{E}_p satisfying these conditions (1)-(5) is canonical. (We will see that the condition (6) is automatically deduced from a combination of (1) and (5).) However, we do not know whether or not such uniqueness also holds for p -energy.

We next introduce \mathcal{E}_p -energy measure $\mu_{\langle f \rangle}^p$ for $f \in \mathcal{F}_p$ and establish a few properties of it in Section 7 (Theorems 7.4, 7.5 and 7.7).

Theorem 2.25. *Assume that $p > \dim_{\text{ARC}}(K, d)$. For any $f \in \mathcal{F}_p$, there exists a Borel finite measure $\mu_{\langle f \rangle}^p$ on K with $\mu_{\langle f \rangle}^p(K) = \mathcal{E}_p(f)$ satisfying the following conditions:*

- (1) if $f, g \in \mathcal{F}_p$ and $A \in \mathcal{B}(K)$ satisfy $(f - g)|_A \equiv \text{const.}$, then $\mu_{\langle f \rangle}^p(A) = \mu_{\langle g \rangle}^p(A)$;
- (2) (Chain rule) for any $\Phi \in \mathcal{C}^1(\mathbb{R})$, it holds that $d\mu_{\langle \Phi \circ f \rangle}^p = |\Phi'| \circ f|^p d\mu_{\langle f \rangle}^p$;
- (3) (Self-similarity) for any $n \in \mathbb{N}$, it holds that

$$\mu_{\langle f \rangle}^p(dx) = \rho_p^n \sum_{w \in W_n} (F_w)_* \mu_{\langle f \circ F_w \rangle}^p(dx),$$

where $(F_w)_* \mu_{\langle f \circ F_w \rangle}^p(A) := \mu_{\langle f \circ F_w \rangle}^p(F_w^{-1}(A))$ for any $A \in \mathcal{B}(K)$.

³To be precise, the uniqueness was proved in [8] in an alternative formulation of (1)-(5). In particular, there is no proof of the self-similarity condition (5) in [8]. The identity (2.9) was proved in [36, Proposition 5.1] and an explicit proof of (2.10) was given in [38, Proposition 5.9].

As mentioned in the introduction, this measure $\mu_{(f)}^p$ plays the role of $|\nabla f(x)|^p dx$ in the case of Euclidean spaces. To treat \mathcal{E}_2 -energy measures, there are established frameworks in terms of Dirichlet forms. For further development of \mathcal{E}_p -energy measures, the lack of p -energy form “ $\mathcal{E}_p(f; g)$ ” (formally written as $(f, g) \mapsto \int \langle |\nabla f|^{p-2} \nabla f, \nabla g \rangle dx$) is a big obstacle. This paper contains no results in this direction.

Remark 2.26. For the Sierpiński gasket, Herman, Peirone and Strichartz [34] have constructed p -energy $\mathcal{E}_p^{\text{HPS}}(f)$, and Strichartz and Wong [58] have suggested an approach to interpret $\mathcal{E}_p^{\text{HPS}}(f; g)$ as subderivatives of $t \mapsto \frac{1}{p} \mathcal{E}_p^{\text{HPS}}(f + tg)$ at $t = 0$. The notion of p -harmonicity and p -Laplacian based on this form $\mathcal{E}_p^{\text{HPS}}(f; g)$ are also considered in [58].

Lastly, we prove $\beta_p > p$ for *planar* generalized Sierpiński carpets.

Theorem 2.27. *Suppose that $D = 2$. Then $\rho_p > N_*^{-1}a^p$. In particular, $\beta_p > p$ for any $p > 0$.*

Remark 2.28. Our proof of Theorem 2.27 is inspired by [39], where $\beta_2 > 2$ is proved for all generalized Sierpiński carpets. However, our argument is limited to the planar case due to the lack of suitable p -energy form “ $\mathcal{E}_p(f; g)$ ” (see also Remark 2.26). In a forthcoming paper [42], the required p -energy forms will be constructed. Using these p -energy forms and following the arguments in [39], we can show $\beta_p > p$ for all generalized Sierpiński carpets without assuming $D = 2$. The details will be provided in [42].

3. ESTIMATES OF POINCARÉ CONSTANTS AND CONDUCTANCES

In this section and Section 4, we investigate relations among (p, p) -Poincaré constants $\lambda_p^{(n)}$, $\lambda_{p, \text{Dir}}^{(n)}$, $\sigma_p^{(n)}$ and p -conductances $\mathcal{C}_p^{(n)}$ (and its reciprocal $\mathcal{R}_p^{(n)}$). Almost all parts of this section are p -energy analogs of [54, Section 2]. The ultimate goal is to show that $\lambda_p^{(n)}$, $\sigma_p^{(n)}$ and $\mathcal{R}_p^{(n)}$ are comparable without depending on the level n . In particular, the estimate $\sigma_p^{(n)} \vee \lambda_p^{(n)} \leq C \mathcal{R}_p^{(n)}$ will be needed in later sections (especially Theorem 4.15 and Corollary 4.16). However, we need some hard preparations to this end. In the case $p = 2$, this was done in [54, Theorem 7.16] under two assumptions: [54, (B-1) and (B-2)]. The following conditions are generalizations of these assumptions to fit our p -energy context.

(B_p) There exist $k_* \in \mathbb{Z}_{\geq 0}$ and a positive constant C_* (that depends only on p and N_*) such that $\sigma_p^{(n)} \leq C_* \lambda_{p, \text{Dir}}^{(n+k_*)}$ for every $n \in \mathbb{N}$.

(KM_p) There exists $C_{\text{KM}} > 0$ such that $\lambda_p^{(n)} \leq C_{\text{KM}} \mathcal{R}_p^{(n)}$ for every $n \in \mathbb{N}$.

A proof of (B₂) for the Sierpiński carpet is given in [54, Proposition 8.1], and we also prove (B_p) for all p by a similar method to theirs in Section 4 (see Proposition 4.5). The condition (KM_p) is essential for our goals. We prove (KM_p) and show that $\lambda_p^{(n)}$, $\sigma_p^{(n)}$ and $\mathcal{R}_p^{(n)}$ are comparable in the next section (see Theorem 4.14). This section is devoted to a part of preparations toward Theorem 4.14.

Remark 3.1. Kusuoka and Zhou have proved (KM₂) using the result of Barlow and Bass [5] that is called the *Knight Move argument* (see [54, Theorem 7.16]). The original Knight Move condition [54, condition (KM)] is a uniform estimate for

discrete harmonic functions with some boundary conditions. We can check that a p -harmonic analog of [54, condition (KM)] is equivalent to (KM_p) under Assumption 4.7, which will be introduced later, and so we call the condition (KM_p) p -Knight Move instead. A recent study by Kigami reveals new important aspects of (KM_p) , and he introduced an important condition that is called p -conductive homogeneity, which plays a similar role as (KM_p) (see [44, Theorem 1.1 and 1.3]).

In this section, let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet and let $p > 0$.

3.1. Basic estimates without (\mathbf{B}_p) and (KM_p) . Let us start by preparing some basic facts. The following proposition is easily derived from the definition of (p, p) -Poincaré constants (for $p = 2$, see [54, Proposition 1.5]).

Proposition 3.2. *Let $n \geq 1$, $m \geq 0$, $w \in W_m$ and $f \in \mathbb{R}^{W_{n+m}}$.*

(1) *It holds that*

$$\sum_{v \in w \cdot W_n} |f(v) - \langle f \rangle_{w \cdot W_n}|^p \leq N_*^n \lambda_p^{(n)} \mathcal{E}_{p, w \cdot W_n}^{G_{n+m}}(f).$$

In particular,

$$(3.1) \quad \sum_{v \in W_n} |f(v) - \langle f \rangle_{W_n}|^p \leq N_*^n \lambda_p^{(n)} \mathcal{E}_p^{G_n}(f).$$

(2) *It holds that*

$$(3.2) \quad \left| \langle f \rangle_{w \cdot W_n} - \langle f \rangle_{W_{n+m}} \right|^p \leq N_*^m \left(1 \vee N_*^{-(p-1)n} \right) \lambda_p^{(n+m)} \mathcal{E}_p^{G_{n+m}}(f).$$

Moreover, for $w \in W_n$, $k \in \{1, \dots, n\}$ and $f \in \mathbb{R}^{W_n}$,

$$(3.3) \quad \left| \langle f \rangle_{[w]_{n-k} \cdot W_k} - \langle f \rangle_{[w]_{n-k+1} \cdot W_{k-1}} \right|^p \leq N_* \lambda_p^{(k)} \mathcal{E}_{p, [w]_{n-k} \cdot W_k}^{G_{n+m}}(f).$$

(3) *For any $(v, w) \in \tilde{E}_m$,*

$$(3.4) \quad \left| \langle f \rangle_{v \cdot W_n} - \langle f \rangle_{w \cdot W_n} \right|^p \leq \sigma_p^{(n)} \mathcal{E}_{p, \{v, w\} \cdot W_n}^{G_{n+m}}(f).$$

Proof. (1) This is immediate from the definition.

(2) Note that a simple computation yields that

$$\langle f \rangle_{w \cdot W_n} - \langle f \rangle_{W_{n+m}} = N_*^{-n} \sum_{v \in w \cdot W_n} (f(v) - \langle f \rangle_{W_{n+m}}).$$

Applying Hölder's inequality, we have that

$$\begin{aligned} \left| \langle f \rangle_{w \cdot W_n} - \langle f \rangle_{W_{n+m}} \right|^p &\leq N_*^{-pn} \left(N_*^{(p-1)n} \vee 1 \right) \sum_{v \in w \cdot W_n} \left| f(v) - \langle f \rangle_{W_{n+m}} \right|^p \\ &\leq N_*^m \left(N_*^{-(n+m)} \vee N_*^{-(pn+m)} \right) \sum_{v \in W_{n+m}} \left| f(v) - \langle f \rangle_{W_{n+m}} \right|^p \\ &\leq N_*^m \left(1 \vee N_*^{-(p-1)n} \right) \lambda_p^{(n+m)} \mathcal{E}_p^{G_{n+m}}(f), \end{aligned}$$

which proves (3.2). Lastly, by viewing $[w]_{n-k} \cdot W_k$ as a copy of W_k , we see that the estimate (3.2) becomes (3.3).

(3) It is obvious from the definition. \square

For $n \geq 0$ and $m \geq 1$, we define $P_{n+m,n}: \mathbb{R}^{W_{n+m}} \rightarrow \mathbb{R}^{W_n}$ by setting

$$P_{n+m,n}f(w) := \langle f \rangle_{w \cdot W_m}, \quad w \in W_n.$$

When m is clear in the context, we abbreviate P_n to denote $P_{n+m,n}$. Note that $\langle P_{n+m,n}f \rangle_{W_n} = \langle f \rangle_{W_{n+m}}$ by a simple calculation.

While the following lemma is also immediate from the definition of $\sigma_p^{(n)}$ (for $p = 2$, see [54, Lemma 2.12]), this lemma will derive some important properties later. In particular, the weak monotonicity (Corollary 4.16) comes from this lemma.

Lemma 3.3. *For all $n \geq 0$, $m \geq 1$, $f \in \mathbb{R}^{W_{n+m}}$ and a subset A of W_n ,*

$$(3.5) \quad \mathcal{E}_{p,A}^{G_n}(P_{n+m,n}f) \leq \left(D^{(p-1)} \vee 1\right) L_* \sigma_p^{(m)} \mathcal{E}_{p,A \cdot W_m}^{G_{n+m}}(f).$$

Proof. If f is a constant function on W_{n+m} , then we have nothing to be proved. Let $f: W_{n+m} \rightarrow \mathbb{R}$ be a function that is not constant. For each $v, w \in W_n$, define a function $\tilde{f}[v, w]$ on W_{n+m} by setting $\tilde{f}[v, w] := \mathcal{E}_{p,\{v,w\} \cdot W_m}^{G_{n+m}}(f)^{-1/p} \cdot f$. Now, it is a simple computation that

$$\begin{aligned} \mathcal{E}_{p,A}^{G_n}(P_n f) &= \sum_{(v,w) \in E_n^A} \left| \langle f \rangle_{v \cdot W_m} - \langle f \rangle_{w \cdot W_m} \right|^p \\ &= \sum_{(v,w) \in E_n^A} \left| \left\langle \tilde{f}[v, w] \right\rangle_{v \cdot W_m} - \left\langle \tilde{f}[v, w] \right\rangle_{w \cdot W_m} \right|^p \mathcal{E}_{p,\{v,w\} \cdot W_m}^{G_{n+m}}(f) \\ &\leq \left(D^{p-1} \vee 1\right) \sigma_p^{(m)} \sum_{(v,w) \in \tilde{E}_n^A} \mathcal{E}_{p,\{v,w\} \cdot W_m}^{G_{n+m}}(f) \\ &\leq \left(D^{(p-1)} \vee 1\right) L_* \sigma_p^{(m)} \mathcal{E}_{p,A \cdot W_m}^{G_{n+m}}(f), \end{aligned}$$

where we used Proposition 2.10 in the third line. \square

The following theorem states the *submultiplicative inequality* of $\mathcal{C}_p^{(n)}$, whose proof can be found in many literatures (e.g. [15, Proposition 3.6], [20, Lemma 3.7], [47, Lemma 4.9.3] or [44, Theorem 4.3]).

Theorem 3.4. *There exists $C_{3.4} > 0$ (depending only on p, L_*) such that*

$$(3.6) \quad \mathcal{C}_p^{(n+m)} \leq C_{3.4} \mathcal{C}_p^{(n)} \mathcal{C}_p^{(m)} \quad \text{for any } n, m \in \mathbb{N}.$$

In particular, the limit $\lim_{n \rightarrow \infty} (\mathcal{C}_p^{(n)})^{1/n} =: \rho_p^{-1} > 0$ exists and

$$\mathcal{C}_p^{(n)} \geq C_{3.4}^{-1} \rho_p^{-n} \quad \text{for any } n \in \mathbb{N}.$$

The constant ρ_p in the above theorem will play indispensable roles in this paper. The following proposition is an extension of [54, Proposition 2.7] and gives an estimate of ρ_p .

Proposition 3.5. *There exists a positive constant $C_{3.5}$ depending only on p, a, D, L_* such that*

$$\mathcal{C}_p^{(n)} \leq C_{3.5} (N_* a^{-p})^n \quad \text{for all } n \in \mathbb{N}.$$

In particular, it holds that $\rho_p \geq N_^{-1} a^p$.*

Proof. Let $z \in W_m$ and set $A := z \cdot W_n$, $B := W_{n+m} \setminus \mathcal{B}_n(z, 1)$, $K_A := \bigcup_{w \in A} K_w$, and $K_B := \bigcup_{w \in B} K_w$. Then, by Lemma 2.13, we have that $\text{dist}(K_A, K_B) := \inf\{d(x, y) \mid x \in K_A, y \in K_B\} \geq ca^{-m}$, where c is a positive constant depending only on a, D . Define a continuous function $f: K \rightarrow \mathbb{R}$ by setting

$$f(x) := \frac{\text{dist}(x, K_A)}{\text{dist}(K_A, K_B)} \wedge 1$$

for each $x \in K$, where $\text{dist}(x, F) := \inf_{y \in F} d(x, y)$ for any subset F of K . Then it is immediate that $f|_{K_A} \equiv 0$ and $f|_{K_B} \equiv 1$, and thus $M_{n+m}f|_A \equiv 0$ and $M_{n+m}f|_B \equiv 1$. This yields that $\mathcal{C}_p^{G_{n+m}}(A, B) \leq \mathcal{E}_p^{G_{n+m}}(M_{n+m}f)$.

Next, we will estimate the p -energy of $M_{n+m}f$ by estimating distances. For $(v, w) \in E_{n+m}$, by the triangle inequality, we have

$$|\text{dist}(F_v(x), K_A) - \text{dist}(F_w(x), K_A)| \leq d(F_v(x), F_w(x)) \leq 2a^{-(n+m)}.$$

By Lemma 2.7,

$$\begin{aligned} & |M_{n+m}f(v) - M_{n+m}f(w)| \\ &= \left| \int_K F_v^* f \, d\mu - \int_K F_w^* f \, d\mu \right| \\ &\leq \frac{1}{\text{dist}(K_A, K_B)} \int_K |\text{dist}(F_v(x), K_A) - \text{dist}(F_w(x), K_A)| \, d\mu(x) \\ &\leq c^{-1}a^{-n}. \end{aligned}$$

Consequently, we conclude that

$$\begin{aligned} \mathcal{C}_p^{G_{n+m}}(A, B) &\leq \mathcal{E}_p^{G_{n+m}}(M_{n+m}f) \\ &\leq \sum_{w \in \mathcal{B}_n(z, 1)} \sum_{v \in W_{n+m}} |M_{n+m}f(v) - M_{n+m}f(w)|^p \mathbf{1}_{E_{n+m}}((v, w)) \\ &\leq c^{-p} L_* (\#\mathcal{B}_n(z, 1)) a^{-pn} \leq c^{-p} L_* (L_* + 1) N_*^n a^{-pn}. \quad \square \end{aligned}$$

Since ‘‘gluing’’ maximizers of $\lambda_{p, \text{Dir}}^{(n)}$ does not increase energies, the next proposition follows (for $p = 2$, see [54, Proposition 2.11]).

Proposition 3.6. *For all $n \geq 1$,*

$$\lambda_{p, \text{Dir}}^{(n)} \leq N_* \left(1 \vee N_*^{-(p-1)n}\right) \lambda_p^{(n+1)}.$$

Proof. Let $f: W_n \rightarrow \mathbb{R}$ satisfy $f|_{\partial_* G_n} \equiv 0$, $\mathcal{E}_p^{G_n}(f) = 1$, and $\langle f \rangle_{W_n} = (\lambda_{p, \text{Dir}}^{(n)})^{1/p}$ (see Proposition 2.19-(2)). Fix $i \neq j \in S$ and define $f_*: W_{n+1} \rightarrow \mathbb{R}$ by

$$f_*(z) := \begin{cases} f(w) & \text{if } z = iw \text{ for some } w \in W_n, \\ -f(w) & \text{if } z = jw \text{ for some } w \in W_n, \\ 0 & \text{otherwise.} \end{cases}$$

Since $f|_{\partial_* G_n} = 0$, we easily see that $\langle f_* \rangle_{W_{n+1}} = 0$ and $\mathcal{E}_p^{G_{n+1}}(f_*) = 2$. By Hölder's inequality,

$$\begin{aligned} & \left(N_*^{-n} \vee N_*^{-pn} \right) \sum_{z \in W_{n+1}} \left| f_*(z) - \langle f_* \rangle_{W_{n+1}} \right|^p \\ &= 2 \left((\#W_n)^{-1} \vee (\#W_n)^{-p} \right) \sum_{w \in W_n} |f(w)|^p \\ &\geq 2 (\#W_n)^{-p} \left| \sum_{w \in W_n} f(w) \right|^p = 2 |\langle f \rangle_{W_n}|^p = 2 \lambda_{p, \text{Dir}}^{(n)}. \end{aligned}$$

Combining with Proposition 3.2-(1), we finish the proof. \square

Next, we see relations between two (p, p) -Poincaré constants $\lambda_p^{(n)}$ and $\sigma_p^{(n)}$. The following proposition states that the submultiplicative inequality of $\sigma_p^{(n)}$ holds (for $p = 2$, see [54, a part of Proposition 2.13]).

Proposition 3.7. (1) For any $n, m \in \mathbb{N}$,

$$\lambda_p^{(n+m)} \leq (2^{p-1} \vee 1) \left\{ \lambda_p^{(m)} N_*^{-n} + L_*(D^{p-1} \vee 1) \lambda_p^{(n)} \sigma_p^{(m)} \right\}.$$

(2) For any $n, m \in \mathbb{N}$,

$$\sigma_p^{(n+m)} \leq (D^{p-1} \vee 1) L_* \sigma_p^{(n)} \sigma_p^{(m)}.$$

Proof. (1) Let $f: W_{n+m} \rightarrow \mathbb{R}$ with $\mathcal{E}_p^{G_{n+m}}(f) = 1$. Then we see from Proposition 3.2-(1) and Lemma 3.3 that

$$\begin{aligned} & N_*^{-(n+m)} \sum_{v \in W_n} \sum_{w \in v \cdot W_m} \left| f(w) - \langle f \rangle_{W_{n+m}} \right|^p \\ &\leq (2^{p-1} \vee 1) N_*^{-(n+m)} \sum_{\substack{v \in W_n, \\ w \in v \cdot W_m}} \left(|f(w) - \langle f \rangle_{v \cdot W_m}|^p + \left| \langle f \rangle_{v \cdot W_m} - \langle P_{n+m, n} f \rangle_{W_n} \right|^p \right) \\ &\leq (2^{p-1} \vee 1) \lambda_p^{(m)} N_*^{-n} + (2^{p-1} \vee 1) N_*^{-n} \sum_{v \in W_n} \left| P_{n+m, n} f(v) - \langle P_{n+m, n} f \rangle_{W_n} \right|^q \\ &\leq (2^{p-1} \vee 1) \left\{ \lambda_p^{(m)} N_*^{-n} + L_*(D^{p-1} \vee 1) \lambda_p^{(n)} \sigma_p^{(m)} \right\}. \end{aligned}$$

Since f with $\mathcal{E}_p^{G_{n+m}}(f) = 1$ is arbitrary, we obtain the desired estimate.

(2) Let $k \in \mathbb{N}$, let $(v, w) \in \tilde{E}_k$ and let $f \in \mathbb{R}^{W_{n+m+k}}$ satisfy $\mathcal{E}_{p, \{v, w\} \cdot W_{n+m}}(f) = 1$. Note that $\langle f \rangle_{v \cdot W_{n+m}} = \langle P_{n+k} f \rangle_{v \cdot W_n}$, where $P_{n+k} = P_{n+m+k, n+k}$. Indeed,

$$\begin{aligned} \langle f \rangle_{v \cdot W_{n+m}} &= N_*^{-(n+m)} \sum_{v' \in W_n} \sum_{z \in vv' \cdot W_m} f(z) = N_*^{-n} \sum_{v' \in W_n} \langle f \rangle_{vv' \cdot W_m} \\ &= N_*^{-n} \sum_{v' \in W_n} P_{n+k} f(vv') = \langle P_{n+k} f \rangle_{v \cdot W_n}. \end{aligned}$$

Similar computation yields that $\langle f \rangle_{v \cdot W_{n+m}} = \langle P_{m+k} f \rangle_{v \cdot W_m}$. From Lemma 3.3,

$$\begin{aligned} \left| \langle f \rangle_{v \cdot W_{n+m}} - \langle f \rangle_{w \cdot W_{n+m}} \right|^p &= \left| \langle P_{n+k} f \rangle_{v \cdot W_n} - \langle P_{n+k} f \rangle_{w \cdot W_n} \right|^p \\ &\leq \sigma_p^{(n)} \mathcal{E}_{p, \{v, w\} \cdot W_n}^{G_{n+k}}(P_{n+k} f) \\ &\leq (D^{p-1} \vee 1) L_* \sigma_p^{(n)} \sigma_p^{(m)}. \end{aligned}$$

The desired result is immediate from this estimate. \square

In the rest of this subsection, we prove the following relation with (p, p) -Poincaré constant $\lambda_p^{(n)}$ and p -conductance $\mathcal{C}_p^{(n)}$ (see [54, Proposition 2.10] for $p = 2$).

Proposition 3.8. *Let $p \geq 1$. For every $n, m, k \in \mathbb{N}$ with $W_k \setminus \partial_* G_k \neq \emptyset$,*

$$\frac{\#(W_k \setminus \partial_* G_k)}{N_*^k \#(\partial_* G_k)} \mathcal{R}_p^{(m)} \lambda_p^{(n)} \leq C_{3.8} \lambda_p^{(n+m+k)},$$

where $C_{3.8}$ is a positive constant depending only on p and L_* .

Similar ideas of its proof appear in many contexts (see [6, proof of Theorem 3.3], [15, proof of Proposition 3.6] for example). In the following two lemmas, we prepare estimates for “partition of unity” (for $p = 2$, see [54, Lemmas 2.8 and 2.9]).

Lemma 3.9. *Let $n, m \in \mathbb{N}$, and let $\{\varphi_w^{(m)}\}_{w \in W_n}$ be a family of $[0, 1]$ -valued functions on W_{n+m} such that $\sum_{w \in W_n} \varphi_w^{(m)} \equiv 1$ and $\varphi_w^{(m)}|_{W_{n+m} \setminus \mathcal{B}_m(w, 1)} \equiv 0$ for each $w \in W_n$. If $f \in \mathbb{R}^{W_n}$, then*

$$\mathcal{E}_p^{G_{n+m}}(f_*) \leq C_{3.9} \mathcal{E}_p^{G_n}(f) \max_{w \in W_n} \mathcal{E}_p^{G_{n+m}}(\varphi_w^{(m)})$$

where $C_{3.9} > 0$ is constant depending only on p, L_* , and $f_* \in \mathbb{R}^{W_{n+m}}$ is defined as

$$f_*(z) := \sum_{w \in W_n} f(w) \varphi_w^{(m)}(z), \quad z \in W_{n+m}.$$

Proof. For each $z, z' \in W_{n+m}$, we set

$$A(z, z') := \left\{ w \in W_n \mid \varphi_w^{(m)}(z) \vee \varphi_w^{(m)}(z') > 0 \right\}.$$

Since $\text{supp}[\varphi_w^{(m)}] \subseteq \mathcal{B}_m(w, 1)$, we can verify that there exists $M \in \mathbb{N}$ depending only on L_* such that $\#A(z, z') \leq M$ for any $n, m \in \mathbb{N}$ and $z, z' \in W_{n+m}$. Furthermore, we see that

$$f_*(z) - f_*(z') = \sum_{w \in A(z, z')} f(w) \left(\varphi_w^{(m)}(z) - \varphi_w^{(m)}(z') \right)$$

and $\sum_{w \in A(z, z')} (\varphi_w^{(m)}(z) - \varphi_w^{(m)}(z')) = 0$. From these identities, we have that

$$\begin{aligned} (3.7) \quad \mathcal{E}_p^{G_{n+m}}(f_*) &= \frac{1}{2} \sum_{w \in W_n} \sum_{z \in w \cdot W_m} \sum_{\substack{z' \in W_{n+m}; \\ (z, z') \in E_{n+m}}} |f_*(z) - f_*(z')|^p \\ &= \frac{1}{2} \sum_{\substack{w \in W_n, \\ z \in w \cdot W_m}} \sum_{\substack{z' \in W_{n+m}; \\ (z, z') \in E_{n+m}}} \left| \sum_{v \in A(z, z')} (f(v) - f(w)) \left(\varphi_v^{(m)}(z) - \varphi_v^{(m)}(z') \right) \right|^p. \end{aligned}$$

We first consider the case $p > 1$. By Hölder's inequality we obtain

$$(3.8) \quad \left| \sum_{v \in A(z, z')} (f(v) - f(w)) (\varphi_v^{(m)}(z) - \varphi_v^{(m)}(z')) \right|^p \\ \leq \left(\sum_{v \in A(z, z')} |f(v) - f(w)|^p \right) \left(\sum_{v \in A(z, z')} \left| \varphi_v^{(m)}(z) - \varphi_v^{(m)}(z') \right|^{p/(p-1)} \right)^{p-1}.$$

To bound the term $\sum_{v \in A(z, z')} |f(v) - f(w)|^p$, for each $v \in A(z, z')$, $w \in W_n$ with $z \in w \cdot W_m$, and $(z, z') \in E_{n+m}$, we find a path $[w^1, \dots, w^l]$ in G_n from v to w with $l \leq 3$, that is, $(w^i, w^{i+1}) \in E_n$ or $w^i = w^{i+1}$ for each $i = 1, \dots, l-1$, and $w^1 = v, w^l = w$. Define

$$\Gamma_{\leq 3}^{(n)}(w) := \left\{ [w^1, \dots, w^l] \mid \begin{array}{l} l \leq 3, w^i \in W_n, w^l = w, \text{ and} \\ (w^i, w^{i+1}) \in E_n \text{ for each } i = 1, \dots, l-1 \end{array} \right\} \neq \emptyset.$$

Then, for any $(z, z') \in E_{n+m}$, we see that

$$(3.9) \quad \sum_{v \in A(z, z')} |f(v) - f(w)|^p \\ \leq C_1 \sum_{v \in A(z, z')} \sum_{i=1}^{l-1} |f(w^i) - f(w^{i+1})|^p \\ \leq C_1 \sum_{[w^1, \dots, w^l] \in \Gamma_{\leq 3}^{(n)}(w)} \sum_{i=1}^{l-1} |f(w^i) - f(w^{i+1})|^p =: S_f(w),$$

where C_1 is a constant depending only on p, l . Note that the number $\#\Gamma_{\leq 3}^{(n)}(w)$ is bounded above by a constant depending only on L_* . Thus we conclude that there exists a constant C_2 depending only on C_1 and L_* such that $\sum_{w \in W_n} S_f(w) \leq C_2 \mathcal{E}_p^{G_n}(f)$ for any $n \in \mathbb{N}$ and $f: W_n \rightarrow \mathbb{R}$. Combining these estimates (3.7), (3.8) and (3.9), we obtain

$$\mathcal{E}_p^{G_{n+m}}(f_*) \\ \leq \frac{1}{2} \sum_{w \in W_n} S_f(w) \sum_{z \in w \cdot W_m} \sum_{\substack{z' \in W_{n+m}; \\ (z, z') \in E_{n+m}}} \left(\sum_{v \in A(z, z')} \left| \varphi_v^{(m)}(z) - \varphi_v^{(m)}(z') \right|^{p/(p-1)} \right)^{p-1} \\ \leq M^{p-1} \sum_{w \in W_n} S_f(w) \sum_{z \in w \cdot W_m} \sum_{\substack{z' \in W_{n+m}; \\ (z, z') \in E_{n+m}}} \max_{v \in W_n} \left| \varphi_v^{(m)}(z) - \varphi_v^{(m)}(z') \right|^p \\ \leq M^{p-1} \sum_{w \in W_n} S_f(w) \max_{v \in W_n} \mathcal{E}_{p, w \cdot W_m}^{G_{n+m}}(\varphi_v^{(m)}) \\ \leq C_2 M^{p-1} \mathcal{E}_p^{G_n}(f) \max_{v \in W_n} \mathcal{E}_p^{G_{n+m}}(\varphi_v^{(m)}),$$

which finishes the proof when $p > 1$.

If $p \in (0, 1]$, then we can use

$$\begin{aligned} & \left| \sum_{v \in A(z, z')} (f(v) - f(w)) (\varphi_v^{(m)}(z) - \varphi_v^{(m)}(z')) \right|^p \\ & \leq \left(\sum_{v \in A(z, z')} |f(v) - f(w)|^p \right) \left(\sum_{v \in A(z, z')} |\varphi_v^{(m)}(z) - \varphi_v^{(m)}(z')|^p \right) \end{aligned}$$

instead of (3.8) and we obtain the desired estimate in a similar way. \square

Lemma 3.10. *Let $n, m, k \in \mathbb{N}$ with $W_k \setminus \partial_* G_k \neq \emptyset$. If $f \in \mathbb{R}^{W_n}$, then there exists a function $f_* \in \mathbb{R}^{W_{n+m+k}}$ satisfying*

$$(3.10) \quad f_*(v) = f(w) \quad \text{if } w \in W_n \text{ and } v \in ww' \cdot W_m \text{ for some } w' \in W_k \setminus \partial_* G_k,$$

and

$$(3.11) \quad \mathcal{E}_p^{G_{n+m+k}}(f_*) \leq C_{3.10} \#(\partial_* G_k) \mathcal{C}_p^{(m)} \mathcal{E}_p^{G_n}(f),$$

where $C_{3.10}$ is a positive constant depending only on p and L_* .

Proof. For each $w \in W_{n+k}$, let $h_w^{(m)}: W_{n+m+k} \rightarrow [0, 1]$ satisfy $h_w^{(m)}|_{w \cdot W_m} \equiv 1$, $h_w^{(m)}|_{W_{n+m+k} \setminus \mathcal{B}_m(w, 1)} \equiv 0$, and

$$\mathcal{E}_p^{G_{n+m+k}}(h_w^{(m)}) \leq 2\mathcal{C}_p^{G_{n+m+k}}(w \cdot W_m, W_{n+m+k} \setminus \mathcal{B}_m(w, 1)).$$

Define $\Psi := \sum_{w \in W_{n+k}} h_w^{(m)}$. Then it is obvious that $\Psi \geq 1$, and so a family $\{\varphi_w^{(m)}\}_{w \in W_{n+k}}$ given by $\varphi_w^{(m)} := \Psi^{-1} h_w^{(m)}$ satisfies the conditions in Lemma 3.9. For each $f: W_n \rightarrow \mathbb{R}$, define a function $f_*: W_{n+m+k} \rightarrow \mathbb{R}$ by setting

$$f_*(v) := \sum_{z \in W_{n+k}} f([z]_n) \varphi_z^{(m)}(v), \quad v \in W_{n+m+k}.$$

We will prove that f_* is the required function.

First, we check (3.10). Define $f_{n+k}: W_{n+k} \rightarrow \mathbb{R}$ by

$$f_{n+k}(w) := f([w]_n), \quad w \in W_{n+k}.$$

Since $\text{supp}[\varphi_w^{(m)}] \subseteq \mathcal{B}_m(w, 1)$, we can write

$$f_*(v) = \sum_{\substack{z \in W_{n+k}: \\ v \in \mathcal{B}_m(z, 1)}} f_{n+k}(z) \varphi_z^{(m)}(v), \quad v \in W_{n+m+k}.$$

Let $v \in W_{n+m+k}$ and $w \in W_n$ such that $v \in ww' \cdot W_m$ for some $w' \in W_k \setminus \partial_* G_k$. From $w' \notin \partial_* G_k$, it follows that $\mathcal{B}_m(ww', 1) \cap (w \cdot W_{m+k})^c = \emptyset$, and thus, for any $z \in W_{n+k}$ with $v \in \mathcal{B}_m(z, 1)$, we obtain $[z]_n = w$. From this observation, it holds that $f_{n+k}(z) = f(w)$, and thus we obtain

$$f_*(v) = \sum_{\substack{z \in W_{n+k}; \\ v \in \mathcal{B}_m(z, 1)}} f([z]_n) \varphi_z^{(m)}(v) = \sum_{\substack{z \in W_{n+k}; \\ v \in \mathcal{B}_m(z, 1)}} f(w) \varphi_z^{(m)}(v) = f(w),$$

which proves (3.10).

To prove (3.11), it will suffice to show the bound

$$(3.12) \quad \max_{w \in W_{n+k}} \mathcal{E}_p^{G_{n+m+k}}(\varphi_w^{(m)}) \leq c_1 \mathcal{C}_p^{(m)},$$

where c_1 is a positive constant depending only on p and L_* . Indeed, by Lemma 3.9, we have

$$\begin{aligned} \mathcal{E}_p^{G_{n+m+k}}(f_*) &\leq C_{3.9} \mathcal{E}_p^{G_{n+k}}(f_{n+k}) \max_{w \in W_{n+k}} \mathcal{E}_p^{G_{n+m+k}}(\varphi_w^{(m)}) \\ &\leq 2C_{3.9} \#(\partial_* G_k) \mathcal{E}_p^{G_n}(f) \max_{w \in W_{n+k}} \mathcal{E}_p^{G_{n+m+k}}(\varphi_w^{(m)}). \end{aligned}$$

A combination of this estimate and (3.12) yields (3.11). Towards proving (3.12), we start by observing that $\#A_m(w) \leq M$ for some constant M depending only on L_* , where

$$A_m(w) := \left\{ z \in W_{n+k} \mid \mathcal{B}_m(w, 1) \cap \overline{\mathcal{B}_m(z, 1)} \neq \emptyset \right\},$$

for each $w \in W_{n+k}$. Indeed, we have $z \in A_m(w)$ if $z \in W_{n+k}$ satisfies $h_w^{(m)}(v) \wedge h_z^{(m)}(v) \neq 0$ for some $v \in W_{n+m+k}$, and thus we obtain (similarly to the bound of $A(z, z')$ in the proof of Lemma 3.9) that $\#A_m(w) \leq M$ for any $n, m, k \in \mathbb{N}$ and $w \in W_{n+k}$. Now, it is a simple computation that, for any $v, v' \in W_{n+m+k}$,

$$\varphi_w^{(m)}(v) - \varphi_w^{(m)}(v') = \frac{1}{\Psi(v)\Psi(v')} \left(\Psi(v) \left(h_w^{(m)}(v) - h_w^{(m)}(v') \right) - h_w^{(m)}(v) (\Psi(v) - \Psi(v')) \right).$$

If we put $h_w(v, v') = h_w^{(m)}(v) - h_w^{(m)}(v')$ for each $w \in W_{n+k}$ and $(v, v') \in E_{n+m+k}$, then we have from the above identity that

$$\begin{aligned} &\mathcal{E}_p^{G_{n+m+k}}(\varphi_w^{(m)}) \\ &\leq (2^{p-1} \vee 1) \sum_{(v, v') \in E_{n+m+k}} \left(\frac{1}{|\Psi(v')|^p} |h_w(v, v')|^p + \frac{|h_w^{(m)}(v)|^p}{|\Psi(v)\Psi(v')|^p} |\Psi(v) - \Psi(v')|^p \right) \\ &\leq (2^{p-1} \vee 1) \left(\mathcal{E}_p^{G_{n+m+k}}(h_w^{(m)}) + \sum_{(v, v') \in E_{n+m+k}} \left| \sum_{w' \in A_m(w)} h_{w'}(v, v') \right|^p \right) \\ &\stackrel{(*)}{\leq} ((2M)^{p-1} \vee 1) \left(\mathcal{E}_p^{G_{n+m+k}}(h_w^{(m)}) + \sum_{w' \in A_m(w)} \mathcal{E}_p^{G_{n+m+k}}(h_{w'}^{(m)}) \right) \\ &\leq ((2M)^{p-1} \vee 1)(M+1) \max_{w \in W_{n+k}} \mathcal{E}_p^{G_{n+m+k}}(h_w^{(m)}) \leq 2((2M)^{p-1} \vee 1)(M+1) \mathcal{C}_p^{(m)}, \end{aligned}$$

where we used Hölder's inequality in (*). This shows (3.12). \square

With these preparations in place, we are ready to prove Proposition 3.8.

Proof of Proposition 3.8. Let $f: W_n \rightarrow \mathbb{R}$ with $\mathcal{E}_p^{G_n}(f) = 1$, and let $f_* \in \mathbb{R}^{W_{n+m+k}}$ be a function obtained by applying Lemma 3.10 to f . From Lemma 3.10 and $\mathcal{E}_p^{G_n}(f) = 1$, we have $\mathcal{E}_p^{G_{n+m+k}}(f_*) \leq C_{3.10} \#(\partial_* G_k) \mathcal{C}_p^{(m)}$. On the one hand, Proposition 3.2-(1) yields that

$$N_*^{-(n+m+k)} \sum_{w \in W_{n+m+k}} \left| f_*(w) - \langle f_* \rangle_{W_{n+m+k}} \right|^p \leq C_{3.10} \lambda_p^{(n+m+k)} \#(\partial_* G_k) \mathcal{C}_p^{(m)}.$$

On the other hand, from the property of f_* in (3.10), we have that

$$\begin{aligned}
& N_*^{-(n+m+k)} \sum_{w \in W_{n+m+k}} \left| f_*(w) - \langle f_* \rangle_{W_{n+m+k}} \right|^p \\
& \geq N_*^{-(n+m+k)} \sum_{w \in W_n} \sum_{w' \in W_k \setminus \partial_* G_k} \sum_{z \in ww' \cdot W_m} \left| f(w) - \langle f_* \rangle_{W_{n+m+k}} \right|^p \\
& \geq \#(W_k \setminus \partial_* G_k) N_*^{-(n+k)} \sum_{w \in W_n} \left| f(w) - \langle f_* \rangle_{W_{n+m+k}} \right|^p \\
& \geq 2^{-p} \frac{\#(W_k \setminus \partial_* G_k)}{\#W_k} N_*^{-n} \sum_{w \in W_n} |f(w) - \langle f \rangle_{W_n}|^p,
\end{aligned}$$

where we used $\sum_{w \in W_n} |f(w) - c|^p \geq 2^{-p} \sum_{w \in W_n} |f(w) - \langle f \rangle_{W_n}|^p$ for any $c \in \mathbb{R}$ (see [11, Lemma 4.17] for example) in the last line, which requires $p \geq 1$. Since f with $\mathcal{E}_p^{G_n}(f) = 1$ is arbitrary, we obtain the desired estimate. \square

We conclude this subsection by giving a submultiplicative-like inequality of Poincaré constants.

Lemma 3.11. *Let $p \geq 1$. There exists a positive constant $C_{3.11}$ depending only on p, a, D, L_*, N_* such that for all $n, m \in \mathbb{N}$,*

$$\lambda_p^{(n)} \leq C_{3.11} N_*^m \lambda_p^{(m)} \sigma_p^{(n)}.$$

In particular,

$$(3.13) \quad \lambda_p^{(n+m)} \leq (2D)^{p-1} L_* C_{3.11} \lambda_p^{(m)} \sigma_p^{(n)}.$$

Proof. Recall that, by Proposition 2.12, $W_k \setminus \partial_* G_k \neq \emptyset$ if $k \geq 2$. By Proposition 3.8, we have that $\mathcal{R}_p^{(m-2)} \lambda_p^{(n)} \leq c_1 \lambda_p^{(n+m)}$ for all $n \in \mathbb{N}$ and $m \geq 3$, where c_1 depends only on a, D, N_* and $C_{3.8}$. Combining this estimate with Proposition 3.7-(1), we obtain

$$\mathcal{R}_p^{(m-2)} \lambda_p^{(n)} \leq c_2 \left(N_*^{-m} \lambda_p^{(n)} + \lambda_p^{(m)} \sigma_p^{(n)} \right),$$

where $c_2 := c_1 L_* (2D)^{p-1} > 0$. Since $N_*^m \mathcal{R}_p^{(m-2)} \rightarrow \infty$ as $m \rightarrow \infty$ by Proposition 3.5, there exists $M_0 \in \mathbb{N}$ such that

$$\inf_{m \geq M_0} N_*^m \mathcal{R}_p^{(m-2)} \geq c_2 + 1.$$

From these estimates, we have that $(c_2 + 1) N_*^{-m} \lambda_p^{(n)} \leq c_2 (N_*^{-m} \lambda_p^{(n)} + \lambda_p^{(m)} \sigma_p^{(n)})$ for all $n \in \mathbb{N}$ and $m \geq M_0$. Hence, we conclude that

$$\lambda_p^{(n)} \leq c_2 N_*^m \lambda_p^{(m)} \sigma_p^{(n)}.$$

This together with Proposition 3.7-(1) implies (3.13). \square

3.2. Comparability under (B_p) and (KM_p) . Now we will prove submultiplicative inequality of $\lambda_p^{(\cdot)}$ under assuming (B_p) (see [54, Theorem 2.1] for $p = 2$). The following theorem gives a nonlinear analog of [54, Propositions 2.13 and 4.1].

Theorem 3.12. *Let $p \geq 1$ and assume that (B_p) holds. Then there exists a positive constant $C_{3.12}$ depending only on p, a, D, L_*, N_* and the constants associated with (B_p) such that the following statements hold:*

(1) for every $n \in \mathbb{N}$,

$$(3.14) \quad C_{3.12}^{-1} \sigma_p^{(n)} \leq \lambda_p^{(n)} \leq C_{3.12} \sigma_p^{(n)};$$

(2) for every $n, m \in \mathbb{N}$,

$$(3.15) \quad \lambda_p^{(n+m)} \leq C_{3.12} \lambda_p^{(n)} \lambda_p^{(m)};$$

(3) for every $n \in \mathbb{N}$,

$$(3.16) \quad C_{3.12}^{-1} \lambda_p^{(n)} \leq \lambda_{p, \text{Dir}}^{(n)} \leq C_{3.12} \lambda_p^{(n)};$$

(4) for every $n \in \mathbb{N}$,

$$(3.17) \quad \mathcal{R}_p^{(n)} \leq C_{3.12} \lambda_p^{(n)}.$$

Proof. (1) By (B_p) , Proposition 3.6 and (3.13) in Lemma 3.11, for $n \geq 1$,

$$\sigma_p^{(n)} \leq (2D)^{p-1} C_* N_* L_* C_{3.11} \sigma_p^{(k_*+1)} \lambda_p^{(n)}.$$

This implies that $\sigma_p^{(n)} \leq c_1 \lambda_p^{(n)}$ for any $n \in \mathbb{N}$, where c_1 is a positive constant depending only on p, a, D, N_*, L_* and the constants associated with (B_p) . Lemma 3.11 also implies that $\lambda_p^{(n)} \leq C_{3.11} N_* \lambda_p^{(1)} \sigma_p^{(n)}$ for any $n \in \mathbb{N}$. Hence we get (3.14).

(2) The submultiplicative inequality (3.15) is immediate from Proposition 3.7-(2) and (3.14).

(3) Applying Proposition 3.7-(2) and (B_p) , we have that $\sigma_p^{(n)} \leq C_* L_* \sigma_p^{(k_*)} \lambda_{p, \text{Dir}}^{(n)}$ for all $n \geq k_* + 1$, which together with (3.14) implies that

$$(3.18) \quad \lambda_p^{(n)} \leq c_2 \lambda_{p, \text{Dir}}^{(n)} \quad \text{for any } n \in \mathbb{N},$$

where c_2 is a positive constant depending only on p, a, D, N_*, L_* and the constants associated with (B_p) . The converse inequality of (3.18) is immediate from Proposition 3.6 and (3.15).

(4) We immediately get (3.17) from Proposition 3.8 and (3.15). \square

Next we derive supermultiplicative inequalities of (p, p) -Poincaré constants under assuming both (B_p) and (KM_p) .

Theorem 3.13. *Let $p \geq 1$ and assume that both (B_p) and (KM_p) hold. Then there exists a positive constant $C_{3.13}$ (depending only on p, a, D, L_*, N_* and the constants associated with (B_p) and (KM_p)) such that for all $n, m \in \mathbb{N}$*

$$(3.19) \quad C_{3.13}^{-1} \lambda_p^{(n)} \lambda_p^{(m)} \leq \lambda_p^{(n+m)} \leq C_{3.13} \lambda_p^{(n)} \lambda_p^{(m)},$$

and

$$(3.20) \quad C_{3.13}^{-1} \mathcal{R}_p^{(n)} \mathcal{R}_p^{(m)} \leq \mathcal{R}_p^{(n+m)} \leq C_{3.13} \mathcal{R}_p^{(n)} \mathcal{R}_p^{(m)}.$$

Proof. Thanks to Theorem 3.12-(2), in order to show (3.19) it will suffice to prove the supermultiplicative inequality: $\lambda_p^{(n)} \lambda_p^{(m)} \lesssim \lambda_p^{(n+m)}$. From Proposition 3.8 and Theorem 3.12-(2), we have that $\lambda_p^{(n)} \mathcal{R}_p^{(m)} \leq c_1 \lambda_p^{(n+m)}$ for any $n, m \in \mathbb{N}$, where c_1 depends only on p, a, D, L_*, N_* and the constants associated with (B_p) . By (KM_p) , we deduce that

$$C_{\text{KM}}^{-1} \lambda_p^{(n)} \lambda_p^{(m)} \leq \lambda_p^{(n)} \mathcal{R}_p^{(m)} \leq c_1 \lambda_p^{(n+m)},$$

and hence we have $\lambda_p^{(n)} \lambda_p^{(m)} \leq c_1 C_{\text{KM}} \lambda_p^{(n+m)}$.

Since $\lambda_p^{(n)}$ and $\mathcal{R}_p^{(n)}$ are comparable by Theorem 3.12-(4) and (KM_p) , the multiplicative inequality (3.20) follows from (3.19). \square

Note that $\sigma_p^{(n)}$ and $\lambda_p^{(n)}$ are also comparable under the situation of Theorem 3.13. From these multiplicative inequalities, we deduce the following behaviors of $\mathcal{R}_p^{(n)}$, $\lambda_p^{(n)}$ and $\sigma_p^{(n)}$: for all $n \in \mathbb{N}$,

$$(3.21) \quad c_*^{-1} \rho_p^n \leq \mathcal{R}_p^{(n)} \leq c_* \rho_p^n, \quad c_*^{-1} \rho_p^n \leq \lambda_p^{(n)} \leq c_* \rho_p^n \quad \text{and} \quad c_*^{-1} \rho_p^n \leq \sigma_p^{(n)} \leq c_* \rho_p^n,$$

where $\rho_p = \lim_{n \rightarrow \infty} (\mathcal{R}_p^{(n)})^{1/n}$ (see Theorem 3.4) and c_* depends only on p, a, D, N_*, L_* and the constants associated with (B_p) and (KM_p) .

4. CHECKING (B_p) AND (KM_p) FOR GSCs

This section gives p -energy analogs of [54, Lemma 3.9, Proposition 3.10, Theorem 7.2, (B-1) and (B-2)]. The condition (B_p) can be proved in a combinatorial way. In order to prove (KM_p) , we will assume that $p > \dim_{\text{ARC}}(K, d)$. This assumption is needed to derive (uniform) Hölder estimates in Theorem 4.10.

In this section, let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet and let $p > 1$.

4.1. Proof of (B_p) . The condition (B_p) plays the converse role of the Proposition 3.6. Let us start by providing preparations from *asymptotic geometry* in order to simplify several arguments. The following definition extends the notion of *rough isometry among graphs* to that among sequences of graphs.

Definition 4.1. For each $i = 1, 2$, let $\{G_n^i = (V_n^i, E_n^i)\}_{n \geq 1}$ be a series of finite graphs with

$$(4.1) \quad L_*^i := \sup_{n \in \mathbb{N}} \max_{x \in V_n^i} \#\{y \in V_n^i \mid (x, y) \in E_n^i\} < \infty.$$

A family of maps $\{\varphi_n\}_{n \geq 1}$, where $\varphi_n: V_n^1 \rightarrow V_n^2$, is said to be a *uniform rough isometry* from $\{G_n^1\}_{n \geq 1}$ to $\{G_n^2\}_{n \geq 1}$ if:

- (1) there exist constants C_1, C_2 such that, for every $n \in \mathbb{N}$ and $x, y \in V_n^1$,

$$C_1^{-1} d_{G_n^1}(x, y) - C_2 \leq d_{G_n^2}(\varphi_n(x), \varphi_n(y)) \leq C_1 d_{G_n^1}(x, y) + C_2;$$

- (2) there exists a constant C_3 such that, for every $n \in \mathbb{N}$,

$$\bigcup_{x \in V_n^1} B_{d_{G_n^2}}(\varphi_n(x), C_3) = V_n^2;$$

- (3) there exists a constant C_4 such that, for every $n \in \mathbb{N}$ and $x \in V_n^1$,

$$C_4^{-1} \leq \frac{\#\{y' \in V_n^2 \mid (\varphi_n(x), y') \in E_n^2\}}{\#\{y \in V_n^1 \mid (x, y) \in E_n^1\}} \leq C_4.$$

Remark 4.2. Since each φ_n is a rough isometry from G_n^1 to G_n^2 , there exists a rough isometry $\tilde{\varphi}_n$ from G_n^2 to G_n^1 . Moreover, we can choose $\tilde{\varphi}_n$ so that $\{\tilde{\varphi}_n\}_{n \geq 1}$ is a uniform rough isometry from $\{G_n^2\}_{n \geq 1}$ to $\{G_n^1\}_{n \geq 1}$. Consequently, being uniform rough isometry gives an equivalence relation among series of finite graphs satisfying (4.1).

Then the following stability result holds. Its proof is a straightforward modification of [57, proof of Lemma 8.5], and so we omit it here (see Appendix A.1 for a proof).

Lemma 4.3. *Let $\{G_n^i = (V_n^i, E_n^i)\}_{n \geq 1}$ be a series of finite graphs with*

$$L_*^i = \sup_{n \in \mathbb{N}} \max_{x \in V_n^i} \#\{y \in V_n^i \mid (x, y) \in E_n^i\} < \infty,$$

for each $i = 1, 2$, and let $\varphi_n: V_n^1 \rightarrow V_n^2$ be a uniform rough isometry from $\{G_n^1\}_{n \geq 1}$ to $\{G_n^2\}_{n \geq 1}$. Then there exists a positive constant C_{URI} (depending only on C_1, C_2 in Definition 4.1, L_*^1 and p) such that

$$(4.2) \quad \mathcal{E}_p^{G_n^1}(f \circ \varphi_n) \leq C_{\text{URI}} \mathcal{E}_p^{G_n^2}(f),$$

for every $n \in \mathbb{N}$ and $f: V_n^2 \rightarrow \mathbb{R}$. In particular,

$$(4.3) \quad C_p^{G_n^1}(\varphi_n^{-1}(A_n), \varphi_n^{-1}(B_n)) \leq C_{\text{URI}} C_p^{G_n^2}(A_n, B_n)$$

for every $n \in \mathbb{N}$ and disjoint subsets A_n, B_n of V_n^2 .

Next we introduce variants of Poincaré constants $\tilde{\lambda}_{p, \text{Dir}}^{(\cdot)}$ and $\tilde{\sigma}_p^{(\cdot)}$ by setting

$$\tilde{\lambda}_{p, \text{Dir}}^{(n)} := \lambda_{p, \text{Dir}}^{(\tilde{G}_n, \mu)}(\partial_* G_n), \quad \text{and} \quad \tilde{\sigma}_p^{(n)} := \sup_{m \geq 1} \max_{(v, w) \in \tilde{E}_m} \sigma_p^{(\tilde{G}_{m+n}, \mu)}(v \cdot W_n, w \cdot W_n).$$

We can easily verify that there exists a uniform rough isometry between $\{G_n\}_{n \geq 0}$ and $\{\tilde{G}_n\}_{n \geq 0}$ (with C_1, C_2, C_3, C_4 depend only on D, L_*). Our new Poincaré constants have the following variational expressions:

$$\tilde{\lambda}_{p, \text{Dir}}^{(n)} = \inf \left\{ \mathcal{E}_p^{\tilde{G}_n}(f) \mid f \in \mathbb{R}^{W_n}, f|_{\partial_* G_n} \equiv 0, \langle f \rangle_{G_n} = 1 \right\}^{-1}$$

and, for $(v, w) \in \tilde{E}_m$,

$$\begin{aligned} & \sigma_p^{(\tilde{G}_{m+n}, \mu)}(v \cdot W_n, w \cdot W_n) \\ &= \inf \left\{ \mathcal{E}_{p, \{v, w\} \cdot W_n}^{\tilde{G}_{n+m}} \mid f \in \mathbb{R}^{\{v, w\} \cdot W_n}, \langle f \rangle_{v \cdot W_n} - \langle f \rangle_{w \cdot W_n} = 1 \right\}^{-1}. \end{aligned}$$

By noting that average $\langle f \rangle_{A, \mu}$ does not depend on edge sets E_n and \tilde{E}_n , we have the following lemma as a consequence of Lemma 4.3 (see also [44, Section 18]).

Lemma 4.4. *There exists a constant $C_{4.4}$ depending only on p, D, L_* such that*

$$C_{4.4}^{-1} \lambda_{p, \text{Dir}}^{(n)} \leq \tilde{\lambda}_{p, \text{Dir}}^{(n)} \leq C_{4.4} \lambda_{p, \text{Dir}}^{(n)} \quad \text{for all } n \in \mathbb{N},$$

and

$$C_{4.4}^{-1} \sigma_p^{(n)} \leq \tilde{\sigma}_p^{(n)} \leq C_{4.4} \sigma_p^{(n)} \quad \text{for all } n \in \mathbb{N}.$$

Now, we prove (B_p) . We heavily use the symmetries of the underlying (generalized) Sierpiński carpet, namely the symmetries of $[-1, 1]^D$, to prove (B_p) . Recall notations in Definitions 2.3, 2.4 and 2.5.

Proposition 4.5. *There exists a positive constant C_* depending only on p, D, N_*, L_* such that $\sigma_p^{(n)} \leq C_* \lambda_{p, \text{Dir}}^{(n+2)}$ for any $n \in \mathbb{N}$. In particular, (B_p) holds (with $k_* = 2$).*

Proof. The proof is a straightforward (but complicated) generalization of [54, Proposition 8.1], where only the standard Sierpiński carpet is considered. Thanks to Lemma 4.4, it is enough to show that $\tilde{\sigma}_p^{(n)} \lesssim \tilde{\lambda}_{p, \text{Dir}}^{(n+2)}$.

Let $n \in \mathbb{N}$. Using (GSC1), (GSC4) and the self-similarity of K , we easily see that $\tilde{\sigma}_p^{(n)} = \tilde{\sigma}_p^{(\tilde{G}_{n+1}, \mu)}(\mathbf{0} \cdot W_n, \mathbf{e}_D \cdot W_n)$. By Proposition 2.19-(3), there exists $f: \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n \rightarrow \mathbb{R}$ with $\mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}_{n+1}}(f) = 1$ such that $|\langle f \rangle_{\mathbf{0} \cdot W_n} - \langle f \rangle_{\mathbf{e}_D \cdot W_n}| = (\tilde{\sigma}_p^{(n)})^{1/p}$. Adding a constant if necessary, we may assume that $\langle f \rangle_{\mathbf{0} \cdot W_n} = -\langle f \rangle_{\mathbf{e}_D \cdot W_n}$. Then, by the self-similarity and the uniqueness (Proposition 2.19-(3)), we can show that

$$(4.4) \quad f(\mathbf{e}_D \cdot w) = -f(\mathbf{0} \cdot \tau[R_D](w)), \quad w \in W_n.$$

Indeed, a function $f_*: \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n \rightarrow \mathbb{R}$ given by

$$f_*(iw) = \begin{cases} f(\mathbf{e}_D \cdot \tau[R_D](w)) & \text{if } i = \mathbf{0} \text{ and } w \in W_n, \\ f(\mathbf{0} \cdot \tau[R_D](w)) & \text{if } i = \mathbf{e}_D \text{ and } w \in W_n, \end{cases}$$

satisfies $\mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}_{n+1}}(f_*) = 1$, $|\langle f_* \rangle_{\mathbf{0} \cdot W_n} - \langle f_* \rangle_{\mathbf{e}_D \cdot W_n}| = (\tilde{\sigma}_p^{(n)})^{1/p}$ and $\langle f_* \rangle_{\mathbf{0} \cdot W_n} = -\langle f_* \rangle_{\mathbf{e}_D \cdot W_n}$. Moreover, it is immediate that $\langle f \rangle_{\mathbf{0} \cdot W_n} = -\langle f_* \rangle_{\mathbf{0} \cdot W_n}$. The uniqueness in Proposition 2.19-(3) implies that $(\sigma_p^{(n)})^{-1/p} f = \pm (\sigma_p^{(n)})^{-1/p} f_*$. If $f = f_*$, then we have $\langle f_* \rangle_{\mathbf{0} \cdot W_n} = -\langle f_* \rangle_{\mathbf{0} \cdot W_n}$, which leads to a contradiction since $\tilde{\sigma}_p^{(n)} = |\langle f_* \rangle_{\mathbf{0} \cdot W_n} - \langle f_* \rangle_{\mathbf{e}_D \cdot W_n}|^p = 0$. Hence the case $f = f_*$ does not happen and thus $f = -f_*$. This proves (4.4).

Next we will show that $f|_{\mathbf{0} \cdot W_n} \geq 0$ or $f_*|_{\mathbf{0} \cdot W_n} \geq 0$ holds. Suppose that $\langle f \rangle_{\mathbf{0} \cdot W_n} \geq 0$. Obviously, a function $f_*: \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n \rightarrow \mathbb{R}$ given by

$$f_*(iw) = \begin{cases} |f(\mathbf{0} \cdot w)| & \text{if } i = \mathbf{0} \text{ and } w \in W_n, \\ -|f(\mathbf{e}_D \cdot w)| & \text{if } i = \mathbf{e}_D \text{ and } w \in W_n, \end{cases}$$

satisfies

$$|\langle f_* \rangle_{\mathbf{0} \cdot W_n} - \langle f_* \rangle_{\mathbf{e}_D \cdot W_n}| \geq |\langle f \rangle_{\mathbf{0} \cdot W_n} - \langle f \rangle_{\mathbf{e}_D \cdot W_n}|.$$

Furthermore, by noting that $f = -f_*$ implies $|f_*(x) - f_*(y)| = |f(x) - f(y)|$ for $(x, y) \in \tilde{E}_{n+1}$ with $x \in \mathbf{0} \cdot W_n$ and $y \in \mathbf{e}_D \cdot W_n$, we have

$$\mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}_{n+1}}(f_*) \leq \mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}_{n+1}}(f) = 1.$$

Since f is an optimizer of $\tilde{\sigma}_p^{(\tilde{G}_{n+1}, \mu)}(\mathbf{0} \cdot W_n, \mathbf{e}_D \cdot W_n)$, we deduce from Proposition 2.19-(3) that $f = \pm f_*$. The condition $\langle f \rangle_{\mathbf{0} \cdot W_n} \geq 0$ implies $f|_{\mathbf{0} \cdot W_n} \geq 0$. The proof in the case $-\langle f_* \rangle_{\mathbf{0} \cdot W_n} = \langle f \rangle_{\mathbf{0} \cdot W_n} \leq 0$ is similar and we have $f_*|_{\mathbf{0} \cdot W_n} \geq 0$ in this case.

By considering f_* instead of f if necessary, we can assume that f satisfies $f|_{\mathbf{0} \cdot W_n} \geq 0$. Then we have $\langle f \rangle_{\mathbf{0} \cdot W_n} = -\langle f \rangle_{\mathbf{e}_D \cdot W_n} \geq 0$ and hence

$$(4.5) \quad \langle f \rangle_{\mathbf{0} \cdot W_n} = \frac{1}{2} \left(\tilde{\sigma}_p^{(n)} \right)^{1/p}.$$

Furthermore, by the Markov property of \mathcal{E}_p (Proposition 2.17),

$$(4.6) \quad \mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}_{n+1}}(f \vee 0) \leq \mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}_{n+1}}(f) = 1.$$

Note that $(f \vee 0)|_{\mathbf{0} \cdot W_n} \equiv f|_{\mathbf{0} \cdot W_n}$ and $(f \vee 0)|_{\mathbf{e}_D \cdot W_n} \equiv 0$. Roughly speaking, the estimate (4.6) tells us that values of a function $f_0: W_n \rightarrow [0, +\infty)$ given by

$$f_0(w) = f(\mathbf{0} \cdot w), \quad w \in W_n,$$

on $W_n[B_{D,+1}]$ are small so that its p -energy arising from ‘‘boundaries’’ can be controlled. Clearly, $\langle f_0 \rangle_{W_n} = \langle f \rangle_{\mathbf{0} \cdot W_n} = \frac{1}{2}(\tilde{\sigma}_p^{(n)})^{1/p}$. By the uniqueness, we have that $f_0 \circ \tau[R_j] = f_0$ for any $j = 1, \dots, D-1$ and that $f_0 \circ \tau[R_{j,k}^\pm] = f_0$ for any $j, k = 1, \dots, D-1$. Next, for $j = 1, \dots, D-1$, we inductively define $f_j: W_n \rightarrow [0, +\infty)$ by

$$f_j(w) = \begin{cases} f_{j-1}(w) & \text{if } w \in W_n[\mathcal{H}_{j,D}^{+,\leq}], \\ f_{j-1}(\tau[R_{j,D}^+](w)) & \text{if } w \in W_n[\mathcal{H}_{j,D}^{+,\geq}]. \end{cases}$$

This construction yields $f_j \circ \tau[R_k] = f_j$ if $0 \leq j < k < D$. Moreover, $f_j \circ \tau[R_{k,l}^+] = f_j$ for $k, l = 1, \dots, j$ with $k < l$. It is also immediate that $\mathcal{E}_p^{\tilde{G}_n}(f_0) \leq \mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}_{n+1}}(f) = 1$ and $\mathcal{E}_p^{\tilde{G}_n}(f_j) \leq 2\mathcal{E}_p^{\tilde{G}_n}(f_0) \leq 2$ for each $j = 0, \dots, D-1$ (we used both (4.5) and (4.6)).

Finally, we construct a function $h: W_{n+2} \rightarrow [0, +\infty)$, whose estimates of energies and averages will deduce (B_p) . To this end, let us introduce ‘fundamental region’ $\Delta^{a,D} = \Delta$ of $[-3a^{-1}, 3a^{-1}]^{D-1} \times [1-2a^{-1}, 1] =: \mathcal{R}_{a,D}$ that is given by

$$\Delta = \{(x_1, \dots, x_D) \mid x_k \in [0, 3a^{-1}] (1 \leq k \leq D-1), x_D \in [1-2a^{-1}, 1]\} \cap \bigcap_{i=1}^{D-2} \mathcal{H}_{i,i+1}^{+,\geq}.$$

Here, we regard $\bigcap_{i=1}^{D-2} \mathcal{H}_{i,i+1}^{+,\geq}$ as Q_0 when $D = 2$. For each $k \in \{0, \dots, D-1\}$, define $u_k \in \mathbb{R}^D$ by $u_k := \sum_{j=1}^k \mathbf{e}_j$, where we set $u_0 := \mathbf{0}^D$. Then we have

$$\Delta \subseteq \bigcup_{k=0}^{D-1} f_{u_k}(Q_0).$$

Next we inductively define $\{\Delta_k\}_{k=0}^{D-2}$ as follows. Define $\Delta_0 := \Delta$. For given $\Delta_0, \dots, \Delta_{k-1}$, define Δ_k as

$$\Delta_k := \Delta_{k-1} \cup R_{k,k+1}^+(\Delta_{k-1}).$$

Note that this construction yields $\Delta_{D-2} = [0, 3a^{-1}]^{D-1} \times [1-2a^{-1}, 1]$ for all $D \geq 2$. Also, we define subsets $\{\square_k\}_{k=0}^{D-1}$ of $\mathcal{R}_{a,D}$ as follows. Define $\square_0 := \Delta_{D-2}$. For given $\square_0, \dots, \square_{k-1}$, define \square_k as

$$\square_k := \square_{k-1} \cup R_k(\square_{k-1}).$$

This construction yields $\square_{D-1} = \mathcal{R}_{a,D}$.

By $S \subsetneq \{0, 1, \dots, a-1\}^D$, $a \geq 3$, (GSC1) and (GSC4), we can find $v = v_1 v_2 \in W_2$ such that $v_1 - \mathbf{e}_D \notin S$, $v \in W_2[B_{D,-1}]$ and $\mathcal{N}(v) \subseteq v_1 \cdot W_1$, where $\mathcal{N}(v) := \{z \in W_2 \mid d_{G_2}(z, v) \leq 1\}$. Set

$$A_v := \{v_1 \cdot (v_2 + u_k) \mid k \in \{1, \dots, D-1\} \text{ with } v_2 + u_k \in S\}$$

and define $\tilde{h}: A_v \cdot W_n \rightarrow [0, +\infty)$ by

$$\tilde{h}(v_1 \cdot (v_2 + u_k) \cdot z) = f_k(z) \quad \text{for } z \in W_n.$$

Note that $A_v \cdot W_n \supseteq W_{n+2}[f_{v_1}(\Delta_0)]$. The desired function $h: W_{n+2} \rightarrow [0, +\infty)$ will be constructed by ‘unfolding’ \tilde{h} in a suitable way as described below. Define $\tilde{h}_0 := \tilde{h}$. Inductively, for $z \in W_{n+1}$ with $v_1 z \in W_{n+2}[f_{v_1}(\Delta_k)]$, define $\tilde{h}_k: W_{n+2}[f_{v_1}(\Delta_k)] \rightarrow [0, +\infty)$ by

$$\tilde{h}_k(v_1 z) = \begin{cases} \tilde{h}_{k-1}(v_1 z) & \text{if } v_1 z \in W_{n+2}[f_{v_1}(\Delta_{k-1})], \\ \tilde{h}_{k-1}(v_1 \cdot \tau[R_{k,k+1}^+](z)) & \text{otherwise.} \end{cases}$$

Also, define $h_0 := \tilde{h}_{D-2}: W_{n+2}[f_{v_1}(\square_0)] \rightarrow [0, +\infty)$. Inductively, for $z \in W_{n+1}$ with $v_1 z \in W_{n+2}[f_{v_1}(\square_k)]$, define $h_k: W_{n+2}[f_{v_1}(\square_k)] \rightarrow [0, +\infty)$ by

$$h_k(v_1 z) = \begin{cases} h_{k-1}(v_1 z) & \text{if } v_1 z \in W_{n+2}[f_{v_1}(\square_{k-1})], \\ h_{k-1}(v_1 \cdot \tau[R_k](z)) & \text{otherwise.} \end{cases}$$

Lastly, we define $h: W_{n+2} \rightarrow [0, +\infty)$ by

$$h(z) = \begin{cases} h_{D-1}(z) & \text{if } z \in W_{n+2}[f_{v_1}(\mathcal{R}_{a,D})], \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, we have $h|_{W_{n+2} \setminus \mathcal{N}(v) \cdot W_n} \equiv 0$ and $h|_{\partial_* G_{n+2}} \equiv 0$ from the construction. Furthermore, we have from the symmetries of $\{f_j\}_{j=0}^{D-1}$ and the definition of h that

$$\begin{aligned} \mathcal{E}_p^{\tilde{G}^{n+2}}(h) &\leq \sum_{j=0}^{D-1} 2^j \binom{D-1}{j} \left\{ \mathcal{E}_p^{\tilde{G}^n}(f_j) + (j+1) \mathcal{E}_{p, \{\mathbf{0}, \mathbf{e}_D\} \cdot W_n}^{\tilde{G}^{n+1}}(f \vee 0) \right\} \\ &\leq \sum_{j=0}^{D-1} 2^j (j+3) \binom{D-1}{j} \\ &= (D+2) \sum_{j=0}^{D-1} 2^j \binom{D-1}{j} = 3^{D-1} (D+2), \end{aligned}$$

and that

$$\langle h \rangle_{W_{n+2}} \geq N_*^{-(n+2)} \sum_{w \in W_n} f_0(w) = N_*^{-2} \langle f_0 \rangle_{W_n} = \frac{1}{2} N_*^{-2} \left(\tilde{\sigma}_p^{(n)} \right)^{1/p}.$$

Hence, by putting $h_* := \mathcal{E}_p^{\tilde{G}^{n+2}}(h)^{-1/p} \cdot h$, we conclude that

$$\tilde{\lambda}_{p, \text{Dir}}^{(n+2)} \geq \left| \langle h_* \rangle_{W_{n+2}} \right|^p = \mathcal{E}_p^{\tilde{G}^{n+2}}(h)^{-1} \left| \langle h \rangle_{W_{n+2}} \right|^p \geq C_*^{-1} \tilde{\sigma}_p^{(n)}$$

which shows (B_p), where $k_* = 2$ and $C_* = 3^{D-1} (D+2) \cdot (2N_*^2)^p C_{4,4}^2$. \square

4.2. Uniform Hölder estimate: $p > \dim_{\text{ARC}}(K, d)$. Next we will prove useful Hölder type estimates. In order to obtain these estimates, a ‘‘low-dimensional’’ condition: Assumption 4.7, which is described in terms of the *Ahlfors regular conformal dimension*, will be essential. The notion of Ahlfors regular conformal dimension was implicitly introduced by Bourdon and Pajot [16]. The exact definition of this dimension is as follows.

Definition 4.6. Let X be a metrizable space (without isolated points) and let d_i ($i = 1, 2$) be compatible metrics on X . We say that d_1 and d_2 are *quasisymmetric* to each other if there exists a homeomorphism $\eta: [0, \infty) \rightarrow [0, \infty)$ such that

$$\frac{d_2(x, a)}{d_2(x, b)} \leq \eta \left(\frac{d_1(x, a)}{d_1(x, b)} \right),$$

for every triple $x, a, b \in X$ with $x \neq b$. (It is easy to show that being quasisymmetric gives an equivalence relation among metrics.) The *Ahlfors regular conformal gauge* $\mathcal{J}_{\text{AR}}(X, d_1)$ of (X, d_1) is defined as

$$\mathcal{J}_{\text{AR}}(X, d_1) := \left\{ d_2 \mid \begin{array}{l} d_2 \text{ is a metric on } X, d_2 \text{ is quasisymmetric to } d_1, \\ \text{and } d_2 \text{ is } \alpha'\text{-Ahlfors regular for some } \alpha' > 0. \end{array} \right\}.$$

(For the definition of Ahlfors regularity, recall (2.2). Note that $\dim_{\text{H}}(X, d_2) = \alpha'$ if d_2 is α' -Ahlfors regular.) Then the *Ahlfors regular conformal dimension* (ARC-dimension for short) of (X, d_1) is

$$(4.7) \quad \dim_{\text{ARC}}(X, d_1) := \inf_{d_2 \in \mathcal{J}_{\text{AR}}(X, d_1)} \dim_{\text{H}}(X, d_2).$$

The notion of quasisymmetric and the exact definition of ARC-dimension are not essential in this paper. We refer the reader to a monograph [55] and surveys [13, 48] for details of the ARC-dimension and related subjects.

The following assumption describes our “low-dimensional” setting (see [54, the condition (R)] in the context of probability theory).

Assumption 4.7. A positive real number p satisfies $p > \dim_{\text{ARC}}(K, d)$.

Remark 4.8. The following bound concerning the Ahlfors regular conformal dimension of the standard Sierpiński carpet is known:

$$1 + \frac{\log 2}{\log 3} \leq \dim_{\text{ARC}}(K, d) < \alpha = \frac{\log N_*}{\log a} = \frac{\log 8}{\log 3}.$$

The lower bound follows from a general result due to Tyson [59]. The strict inequality in the upper bound is proved by Keith and Laakso [43].

To promote understanding Assumption 4.7, we recall characterization results by Carrasco Piaggio [20] and Kigami [47] in our setting. Recall $\lim_{n \rightarrow \infty} (\mathcal{R}_p^{(n)})^{1/n} = \rho_p$ (Theorem 3.4).

Theorem 4.9 ([20, Theorem 1.3], [47, Theorems 4.6.9 and 4.7.6]). *Assumption 4.7 is equivalent to $\rho_p > 1$. In particular, if Assumption 4.7 holds, then there exist $C_{4.9} > 0$ and $\theta > 0$ (depending only on a, ρ_p) such that*

$$\mathcal{C}_p^{(n)} \leq C_{4.9} a^{-\theta n} \quad \text{for all } n \in \mathbb{N}.$$

Under Assumption 4.7, we can show the following powerful Hölder continuity.

Theorem 4.10. *Suppose Assumption 4.7 holds. Then there exist constants $\tilde{C}_{\text{UH}} > 0$ (depending only on $p, a, D, N_*, L_*, \rho_p$) and $\theta > 0$ (depending only on a, ρ_p) such that, for any $n, m \in \mathbb{N}$, $z \in W_m$, $v, w \in \mathcal{B}_n(z, 1)$ and $f \in \mathbb{R}^{W_{n+m}}$,*

$$(4.8) \quad |f(v) - f(w)|^p \leq \tilde{C}_{\text{UH}} \lambda_p^{(n+m)} \mathcal{E}_p^{G_{n+m}}(f) a^{-m\theta}.$$

This theorem is proved by iterating Proposition 3.2-(2). Kusuoka and Zhou [54] prepared a general estimate using signed measures ([54, Lemma 3.9]) to show Hölder type estimates, but we need only the case of Dirac measures for our purpose. Here we give a simplified extension of [54, Lemma 3.9 and Proposition 3.10].

Lemma 4.11. *Let $n, m \in \mathbb{N}$, let $v \in W_m$ and let $f \in \mathbb{R}^{W_{n+m}}$. Then, for any $w \in v \cdot W_n$,*

$$(4.9) \quad |f(w) - \langle f \rangle_{v \cdot W_n}| \leq N_*^{1/p} \mathcal{E}_{p, v \cdot W_n}^{G_{n+m}}(f)^{1/p} \sum_{k=1}^n (\lambda_p^{(k)})^{1/p}.$$

Proof. Let $w \in v \cdot W_n$ and set $w_l := [w]_l$ for each $l = m, \dots, n+m$. Note that $w_m = v$ and $w_{n+m} = w$. From Proposition 3.2-(2), we see that

$$\begin{aligned} |f(w) - \langle f \rangle_{v \cdot W_n}| &\leq \sum_{l=m}^{n+m-1} \left| \langle f \rangle_{w_l \cdot W_{n+m-l}} - \langle f \rangle_{w_{l+1} \cdot W_{n+m-l-1}} \right| \\ &\leq N_*^{1/p} \mathcal{E}_{p, v \cdot W_n}^{G_{n+m}}(f)^{1/p} \sum_{l=m}^{n+m-1} (\lambda_p^{(n+m-l)})^{1/p} \\ &= N_*^{1/p} \mathcal{E}_{p, v \cdot W_n}^{G_{n+m}}(f)^{1/p} \sum_{k=1}^n (\lambda_p^{(k)})^{1/p}. \quad \square \end{aligned}$$

Proof of Theorem 4.10. By Assumption 4.7 and Theorem 4.9, there exists $\theta > 0$ such that

$$(4.10) \quad \mathcal{C}_p^{(n)} \leq C_{4.9} a^{-n\theta} \quad \text{for every } n \in \mathbb{N}.$$

From (4.10), Proposition 3.8 and Theorem 3.12-(2), we have $\lambda_p^{(n)} \leq c_1 \lambda_p^{(n+m)} a^{-m\theta}$ for every $n, m \in \mathbb{N}$, where $c_1 > 0$ depends only on $p, a, D, L_*, N_*, \rho_p$. In particular,

$$(4.11) \quad \lambda_p^{(k)} \leq c_1 \lambda_p^{(n+m)} a^{-(n+m-k)\theta} \quad \text{for every } n, m, k \in \mathbb{N} \text{ with } k \leq n.$$

By Lemma 4.11, for any $z \in W_m$, $v \in z \cdot W_n$ and $w \in \mathcal{B}_n(z, 1)$,

$$\begin{aligned} &|f(v) - f(w)| \\ &\leq |f(v) - \langle f \rangle_{z \cdot W_n}| + \left| \langle f \rangle_{z \cdot W_n} - \langle f \rangle_{[w]_m \cdot W_n} \right| + |f(w) - \langle f \rangle_{[w]_m \cdot W_n}| \\ &\leq \sum_{i=1}^D \left| \langle f \rangle_{z^{(i-1)} \cdot W_n} - \langle f \rangle_{z^{(i)} \cdot W_n} \right| + 2N_*^{1/p} \mathcal{E}_p^{G_{n+m}}(f)^{1/p} \sum_{k=1}^n (\lambda_p^{(k)})^{1/p}, \end{aligned}$$

where $z^{(i)} \in W_m$ ($i = 0, \dots, D$) with $z^{(0)} = z$, $z^{(D)} = [w]_m$ satisfy $(z^{(i-1)}, z^{(i)}) \in \tilde{E}_m$ or $z^{i-1} = z^i$ for $i = 1, \dots, D$. (Such $\{z^{(i)}\}_i$ exists due to Proposition 2.10.) From Proposition 3.2-(3), Theorem 3.12-(1) and (4.11), we see that

$$\begin{aligned} |f(v) - f(w)| &\leq 2D^{(p-1)/p} \mathcal{E}_p^{G_{n+m}}(f)^{1/p} \left((\sigma_p^{(n)})^{1/p} + N_*^{1/p} \sum_{k=1}^n (\lambda_p^{(k)})^{1/p} \right) \\ &\leq c_2 \mathcal{E}_p^{G_{n+m}}(f)^{1/p} \sum_{k=1}^n (\lambda_p^{(k)})^{1/p} \\ &\leq c_1^{1/p} c_2 \left(\lambda_p^{(n+m)} \mathcal{E}_p^{G_{n+m}}(f) \right)^{1/p} \sum_{k=1}^n a^{-(n+m-k)\theta/p}, \end{aligned}$$

where c_2 is a positive constant depending only on p, a, D, N_*, L_* . Since $\theta > 0$ and

$$\begin{aligned} \sum_{k=1}^n a^{-(n+m-k)\theta/p} &= a^{-m\theta/p} \sum_{k=0}^{n-1} a^{-k\theta/p} \\ &= a^{-m\theta/p} \cdot \frac{1 - a^{-n\theta/p}}{1 - a^{-\theta/p}} \leq \frac{1}{1 - a^{-\theta/p}} \cdot a^{-m\theta/p}, \end{aligned}$$

we have the desired estimate for $v \in z \cdot W_n$ and $w \in \mathcal{B}_n(z, 1)$. A combination of this estimate and the triangle inequality proves the desired estimate. \square

4.3. Proof of (KM_p) : $p > \dim_{\text{ARC}}(K, d)$. The aim of this subsection is to prove (KM_p) under Assumption 4.7. Our strategy for proving (KM_p) comes from a recent study by Cao and Qiu [19], where they give an “analytic” proof of (KM_2) using estimates of Poincaré constants in [54]. Although our proof of (KM_p) is similar to the argument in [19, Section 4], we give a complete proof of (KM_p) for the reader’s convenience. Our argument will depend heavily on the uniform Hölder estimate (Theorem 4.10) and on behaviors of “chain” type p -conductance $\mathcal{C}_p^{(n, M)}$ (its definition will be given later).

Let us start by introducing a new graph $G_{n, M}$, which is a “horizontal chain” consisting of M copies of G_n . The exact definition of $G_{n, M}$ is as follows. Let $n, M \in \mathbb{N}$ with $M \geq 2$ and pick $m \in \mathbb{N}$ such that $a^m \geq M$. Then, by (GSC4), we can find a simple path $[w(1), \dots, w(M)]$ in (V_m, \tilde{E}_m) such that

$$F_{w(i)}(\mathcal{K}^+) = F_{w(i+1)}(\mathcal{K}^-) \quad \text{for each } i = 1, \dots, M-1,$$

where

$$(4.12) \quad \mathcal{K}^- := K \cap B_{1, -1}, \quad \mathcal{K}^+ := K \cap B_{1, +1}.$$

(\mathcal{K}^\pm denotes a couple of opposite faces of K .) We define $G_{n, M} = (V_{n, M}, E_{n, M})$ as a subgraph of G_{n+m} given by

$$V_{n, M} := \bigcup_{i=1}^M w(i) \cdot W_n \quad \text{and} \quad E_{n, M} := \{(x, y) \in E_{n+m} \mid x, y \in V_{n, M}\}.$$

We also consider a horizontal network $\tilde{E}_{n, M}$ defined by

$$\tilde{E}_{n, M} := \{(x, y) \in \tilde{E}_{n+m} \mid x, y \in V_{n, M}\}.$$

Now, we set

$$V_{n, M}^- := w(1) \cdot W_n \quad \text{and} \quad V_{n, M}^+ := w(M) \cdot W_n.$$

and define $\mathcal{C}_p^{(n, M)}$ (see Figure 3) by setting

$$\mathcal{C}_p^{(n, M)} := \mathcal{C}_p^{G_{n, M}}(V_{n, M}^-, V_{n, M}^+).$$

We easily see that these definitions do not depend on choices of large $m \in \mathbb{N}$ and horizontal chain $[w(1), \dots, w(M)]$.

The following lemma describes a key behavior of $\mathcal{C}_p^{(n, M)}$.

Lemma 4.12. *For every $M \geq 3$ there exists a constant $C(M) \geq 1$ depending only on p, L_*, N_*, M such that*

$$(4.13) \quad C(M)^{-1} \mathcal{C}_p^{(n)} \leq \mathcal{C}_p^{(n, M)} \leq C(M) \mathcal{C}_p^{(n)} \quad \text{for any } n \in \mathbb{N}.$$




FIGURE 3. The conductance $\mathcal{C}_p^{(n,M)}$ (the planar case)

Its proof will be a straightforward modification of [19, Lemma 4.7]. In order to prove Lemma 4.12, we need to show that $\mathcal{C}_p^{(n)}$ behaves similarly to the conductance that appeared in the work of Barlow and Bass (see the quantity R_n^{-1} in [6]). To state rigorously, we define

$$W_n^- := W_n[B_{1,-1}], \quad W_n^+ := W_n[B_{1,+1}] \quad \text{and} \quad \mathcal{C}_{p,\text{face}}^{(n)} := \mathcal{C}_p^{G_n}(W_n^-, W_n^+).$$

The next lemma is proved in [15, Lemma 4.4] for some special cases of Sierpiński carpets by using p -combinatorial modulus instead of p -conductance. We give a simple proof without using p -modulus.

Lemma 4.13. *There exists a constant $C_{4.13} \geq 1$ depending only on p, D, L_* such that*

$$C_{4.13}^{-1} \mathcal{C}_p^{(n)} \leq \mathcal{C}_{p,\text{face}}^{(n)} \leq C_{4.13} \mathcal{C}_p^{(n)} \quad \text{for any } n \in \mathbb{N}.$$

Proof. By the self-similarity of K and (GSC1), there exist $m \in \mathbb{N}$ and $w \in W_m$ such that $\mathcal{C}_p^{(n)} = \mathcal{C}_p^{G_{n+m}}(w \cdot W_n, W_{n+m} \setminus \mathcal{B}_n(w, 1))$. Define $A_n := \overline{w \cdot W_n}$ and $B_n := \overline{W_{n+m} \setminus \mathcal{B}_n(w, 1)}$. It is immediate from Proposition 2.16 that $\mathcal{C}_p^{G_{n+m}}(A_n, B_n) \geq \mathcal{C}_p^{(n)}$. For $n \in \mathbb{N}$ and $v \in \partial(w \cdot W_n)$ (resp. $v \in \partial \mathcal{B}_n(w, 1)^c$), we fix $d(v) \in w \cdot W_n$ (resp. $d(v) \in \mathcal{B}_n(w, 1)^c$) such that $(v, d(v)) \in E_{n+m}$. Define $\varphi_n: W_{n+m} \rightarrow W_{n+m}$ by

$$\varphi_n(v) = \begin{cases} v & \text{if } v \notin \partial(w \cdot W_n) \cup \partial \mathcal{B}_n(w, 1)^c, \\ d(v) & \text{if } v \in \partial(w \cdot W_n) \cup \partial \mathcal{B}_n(w, 1)^c. \end{cases}$$

Then we easily see that φ_n is a rough isometry with $C_1 = 1$, $C_2 = 2$, $C_3 = 2$ and $C_4 = L_*/2$ in Definition 4.1. Note that $\mathcal{C}_p^{G_{n+m}}(\partial(w \cdot W_n), \partial \mathcal{B}_n(w, 1)^c) = \mathcal{C}_p^{G_{n+m}}(A_n, B_n)$. Applying (4.3) in Lemma 4.3 and Proposition 2.16, we deduce that there exists $c_1 > 0$ (depending only on p, L_*) such that, for any $n \in \mathbb{N}$,

$$\mathcal{C}_p^{G_{n+m}}(A_n, B_n) \leq c_1 \mathcal{C}_p^{(n)}.$$

Let $f: W_{n+m} \rightarrow \mathbb{R}$ satisfy $f|_{A_n} \equiv 1$, $f|_{B_n} \equiv 0$ and $\mathcal{E}_p^{G_{n+m}}(f) = \mathcal{C}_p^{G_{n+m}}(A_n, B_n)$. If $v \in W_m$ satisfies $(v, w) \in \tilde{E}_m$, then we have that

$$\mathcal{C}_p^{G_{n+m}}(A_n, B_n) \geq \mathcal{E}_{p,v \cdot W_n}^{G_{n+m}}(f) \geq \mathcal{C}_{p,\text{face}}^{(n)},$$

and thus we conclude that $\mathcal{C}_{p,\text{face}}^{(n)} \leq c_1 \mathcal{C}_p^{(n)}$.

The converse can be shown in a very similar way as Proposition 4.5. Define ‘fundamental region’ $\tilde{\Delta}^{a,D} = \tilde{\Delta}$ of $[-3a^{-1}, 3a^{-1}]^D =: \mathcal{Q}_{a,D}$ by

$$\tilde{\Delta} = [0, 3a^{-1}]^D \cap \bigcap_{i=1}^{D-1} \mathcal{H}_{i,i+1}^{+,\geq}.$$

For each $k \in \{0, \dots, D\}$, define $u_k \in \mathbb{R}^D$ by $u_k := \sum_{j=1}^k \mathbf{e}_j$, where we set $u_0 := \mathbf{0}^D$. Then we have

$$\tilde{\Delta} \subseteq \bigcup_{k=0}^D f_{u_k}(Q_0).$$

Next we inductively define $\{\tilde{\Delta}_k\}_{k=0}^{D-1}$ as follows. Define $\tilde{\Delta}_0 := \tilde{\Delta}$. For given $\tilde{\Delta}_0, \dots, \tilde{\Delta}_{k-1}$, define $\tilde{\Delta}_k$ as

$$\tilde{\Delta}_k := \tilde{\Delta}_{k-1} \cup R_{k,k+1}^+(\tilde{\Delta}_{k-1}).$$

Note that this construction yields $\tilde{\Delta}_{D-1} = [0, 3a^{-1}]^D$ for all $D \geq 2$. Also, we define subsets $\{\tilde{\square}_k\}_{k=0}^D$ of $\mathcal{Q}_{a,D}$ as follows. Define $\tilde{\square}_0 := \tilde{\Delta}_{D-1}$. For given $\tilde{\square}_0, \dots, \tilde{\square}_{k-1}$, define $\tilde{\square}_k$ as

$$\tilde{\square}_k := \tilde{\square}_{k-1} \cup R_k(\tilde{\square}_{k-1}).$$

This construction yields $\tilde{\square}_D = \mathcal{Q}_{a,D}$.

Next we will introduce a new graph Γ_n ($n \geq 1$) as follows. For each $n \geq 1$, define $\tilde{W}_n \subseteq (\{0, \dots, a-1\}^D)^n$ by

$$\tilde{W}_n := \left\{ (\sigma_k)_{k=1}^D \cdot w \mid \sigma_k \in \{-1, 0, 1\} \text{ for each } k \in \{1, \dots, D\}, w \in W_{n-1} \right\},$$

and $E(\tilde{W}_n) \subseteq \tilde{W}_n \times \tilde{W}_n$ by

$$E(\tilde{W}_n) :=$$

$$\{(v, w) \mid v, w \in \tilde{W}_n, f_v(Q_0) \cap f_w(Q_0) \text{ is a } (D-1)\text{-dimensional hypercube}\}.$$

Then $\Gamma_n := (\tilde{W}_n, E(\tilde{W}_n))$. Define subsets I_n, O_n of \tilde{W}_n by

$$I_n := \{w \in \tilde{W}_n \mid f_w(Q_0) \cap f_{\mathbf{0}}(Q_0) \neq \emptyset\} = \{w \in \tilde{W}_n \mid f_w(Q_0) \cap [-a^{-1}, a^{-1}]^D \neq \emptyset\},$$

and

$$O_n := \{w \in \tilde{W}_n \mid f_w(Q_0) \cap \partial \mathcal{Q}_{a,D} \neq \emptyset\}.$$

By the cutting law of p -conductances (see [57, Proposition 3.18] for example),

$$\mathcal{C}_p^{\Gamma_{n+1}}(I_{n+1}, O_{n+1}) \geq \mathcal{C}_p^{\tilde{G}^{n+m}}(A_n, B_n) \geq c \cdot \mathcal{C}_p^{G^{n+m}}(A_n, B_n) \geq c \cdot \mathcal{C}_p^{(n)},$$

where $c > 0$ depends only on p, D, L_* (we used Lemma 4.3). Let $f: W_n \rightarrow [0, 1]$ satisfy

$$f|_{W_n^-} \equiv 1, \quad f|_{W_n^+} \equiv 0 \quad \text{and} \quad \mathcal{E}_p^{G^n}(f) = \mathcal{C}_{p,\text{face}}^{(n)}.$$

From the uniqueness of the optimizer of $\mathcal{C}_{p,\text{face}}^{(n)}$, we have $f \circ \tau[R_j] = f$ and $f \circ \tau[R_{j,k}^\pm] = f$ for any $j, k = 2, \dots, D$. Define $h_1: W_n \rightarrow [0, 1]$ as $h_1 := f$. Inductively, define $h_k: W_n \rightarrow [0, 1]$ by

$$h_k(w) = \begin{cases} h_{k-1}(w) & \text{if } w \in W_n[\mathcal{H}_{1,k}^{+,\geq}], \\ h_{k-1}(\tau[R_{1,j}^+](w)) & \text{if } w \in W_n[\mathcal{H}_{1,k}^{+,\leq}]. \end{cases}$$

Then we have $\mathcal{E}_p^{G_n}(h_k) \leq 2\mathcal{E}_p^{G_n}(h_{k-1})$ and hence

$$\max_{k \in \{1, \dots, D\}} \mathcal{E}_p^{G_n}(h_k) \leq 2^{D-1} \mathcal{E}_p^{G_n}(h_0) = 2^{D-1} \mathcal{C}_{p, \text{face}}^{(n)}.$$

Let $\tilde{h}: \{u_k \cdot w \mid k \in \{1, \dots, D\}, w \in W_n\} \rightarrow [0, 1]$ such that

$$\tilde{h}(u_k \cdot w) = h_k(w) \quad \text{for all } k \in \{1, \dots, D\} \text{ and } w \in W_n.$$

Note that $\{u_k \cdot w \mid k \in \{1, \dots, D\}, w \in W_n\}$ contains $\{w \in \widetilde{W}_{n+1} \mid f_w(Q_0) \cap \widetilde{\Delta} \neq \emptyset\}$ and that

$$\mathcal{E}_p(\tilde{h}) = \sum_{k=1}^D \mathcal{E}_p^{\tilde{G}_n}(h_k),$$

where $\mathcal{E}_p(\tilde{h})$ denotes the p -energy of \tilde{h} on the induced subgraph of Γ_{n+1} whose vertex set is given by $\{u_k \cdot w \mid k \in \{1, \dots, D\}, w \in W_n\}$. Similar construction by ‘unfolding’ \tilde{h} as in the proof of Proposition 4.5 yields a function $h_*: \widetilde{W}_{n+1} \rightarrow [0, 1]$ satisfying

$$h_*|_{I_{n+1}} \equiv 1, \quad h_*|_{O_{n+1}} \equiv 0$$

and

$$\mathcal{E}_p^{\Gamma_{n+1}}(h_*) = \sum_{k=1}^D 2^k \binom{D}{k} \mathcal{E}_p^{\tilde{G}_n}(h_k) \leq 2^{D-1} \mathcal{C}_{p, \text{face}}^{(n)} \sum_{k=1}^D 2^k \binom{D}{k} \leq 2^{D-1} 3^D \mathcal{C}_{p, \text{face}}^{(n)}.$$

Hence we conclude that

$$c \cdot \mathcal{C}_p^{(n)} \leq \mathcal{C}_p^{\Gamma_{n+1}}(I_{n+1}, O_{n+1}) \leq \mathcal{E}_p^{\Gamma_{n+1}}(h_*) \leq 2^{D-1} 3^D \mathcal{C}_{p, \text{face}}^{(n)},$$

which completes the proof. \square

Now we are ready to prove Lemma 4.12.

Proof of Lemma 4.12. Thanks to Lemma 4.13, it will suffice to compare $\mathcal{C}_{p, \text{face}}^{(n)}$ and $\mathcal{C}_p^{(n, M)}$. Suppose that $G_{n, M}$ is realized as a subgraph of G_{n+M} using a horizontal chain $[w(1), \dots, w(M)]$ in (V_m, \tilde{E}_m) , that is, $V_{n, M} = \bigcup_{i=1}^M w(i) \cdot W_n$. First, we consider the case $M = 3$. By the monotonicity of p -conductance (Proposition 2.16), we immediately have that $\mathcal{C}_p^{(n, 3)} \leq \mathcal{C}_{p, \text{face}}^{(n)}$. We will prove the converse by using Lemma 4.3. Let us define a subgraph $\widehat{G}_{n, 3} = (\widehat{V}_{n, 3}, \widehat{E}_{n, 3})$ of $G_{n, 3}$ by

$$\widehat{V}_{n, 3} := \{w(1)v \mid v \in W_n^+\} \cup w(2) \cdot W_n \cup \{w(3)v \mid v \in W_n^-\},$$

and

$$\widehat{E}_{n, 3} := \{(v, w) \in E_{n, 3} \mid v, w \in \widehat{V}_{n, 3}\}.$$

Then we easily see that

$$\mathcal{C}_p^{(n, 3)} = \mathcal{C}_p^{\widehat{G}_{n, 3}}(\{\{w(1)v \mid v \in W_n^+\}, \{w(3)v \mid v \in W_n^-\}\}).$$

Define $\varphi_n: W_n \rightarrow \widehat{V}_{n, 3}$ by

$$\varphi_n(w) := \begin{cases} w^1 \tau[R_1](w) & \text{if } w \in W_n^-, \\ w^2 w & \text{if } w \notin W_n^- \cup W_n^+, \\ w^3 \tau[R_1](w) & \text{if } w \in W_n^+. \end{cases}$$

Then $\{\varphi_n\}_{n \geq 1}$ is a uniform rough isometry between $\{G_n\}_{n \geq 1}$ and $\{\widehat{G}_{n,3}\}_{n \geq 1}$ (with $C_1 = 1, C_2 = 2, C_3 = 2$ and $C_4 = L_*/2$ in Definition 4.1). Applying Lemma 4.3, we get $\mathcal{C}_{p,\text{face}}^{(n)} \leq c_1 \mathcal{C}_p^{(n,3)}$, where $c_1 > 0$ depends only on p and L_* .

Next, let us consider the case $M_k := 2^k + 2$ for $k \in \mathbb{N}$. Define $\widetilde{\mathcal{C}}_p^{(n,M)}$ by

$$\widetilde{\mathcal{C}}_p^{(n,M)} = \mathcal{C}_p^{\widetilde{G}_{n,M}}(V_{n,M}^-, V_{n,M}^+),$$

where $\widetilde{G}_{n,M} = (V_{n,M}, \widetilde{E}_{n,M})$. We easily see that $G_{n,M}$ and $(V_{n,M}, \widetilde{E}_{n,M})$ are uniformly rough isometric and thus Lemma 4.3 implies that there exists a constant $c_2 > 0$ (depending only on p, L_*, D) such that

$$c_2^{-1} \mathcal{C}_p^{(n,M)} \leq \widetilde{\mathcal{C}}_p^{(n,M)} \leq c_2 \mathcal{C}_p^{(n,M)}$$

Let $f_{n,k}: V_{n,M_k} \rightarrow \mathbb{R}$ satisfy

$$f_{n,k}|_{V_{n,M_k}^-} \equiv 1, \quad f_{n,k}|_{V_{n,M_k}^+} \equiv 0, \quad \text{and} \quad \mathcal{E}_p^{G_{n,M_k}}(f_{n,k}) = \widetilde{\mathcal{C}}_p^{(n,M_k)}.$$

We shall show that

$$(4.14) \quad \min\{f_{n,k}(v) \mid v \in w(i) \cdot W_n, 1 \leq i \leq 2^{k-1} + 1\} \geq \frac{1}{2}.$$

(See Figure 4.) To this end, let us consider a function $\widehat{f}_{n,k}$ given by

$$\widehat{f}_{n,k}(v) := \begin{cases} f_{n,k}(v) \vee (1 - f_{n,k}(v)) & \text{if } v \in \bigcup_{i=1}^{2^{k-1}+1} w(i) \cdot W_n, \\ f_{n,k}(v) \wedge (1 - f_{n,k}(v)) & \text{if } v \in \bigcup_{i=2^k}^{2^k+2} w(i) \cdot W_n, \end{cases}$$

which obviously satisfies $\widehat{f}_{n,k}|_{V_{n,M_k}^-} \equiv 1$ and $\widehat{f}_{n,k}|_{V_{n,M_k}^+} \equiv 0$. From the uniqueness of $f_{n,k}$, for $i = 1, \dots, M_k$ and $w \in W_n$,

$$f_{n,k}(w(i)w) = 1 - f_{n,k}(w(M_k - i + 1)\tau[R_1](w)).$$

This yields that

$$|f_{n,k}(x) - f_{n,k}(y)| = \left| \widehat{f}_{n,k}(x) - \widehat{f}_{n,k}(y) \right|, \quad (x, y) \in E_n(w(2^{k-1} + 1), w(2^{k-1} + 2)),$$

where $E_n(w(2^{k-1} + 1), w(2^{k-1} + 2))$ denotes

$$\left\{ (x_1, x_2) \in E_{n+m} \mid x_i \in w(2^{k-1} + i) \cdot W_n \text{ for } i = 1, 2 \right\}.$$

Hence we have $\mathcal{E}_p(\widehat{f}_{n,k}) \leq \mathcal{E}_p(f_{n,k})$. By the uniqueness of $f_{n,k}$, we have $\widehat{f}_{n,k} = f_{n,k}$, which deduces (4.14). Similarly, we also have

$$(4.15) \quad \max\{f_{n,k}(v) \mid v \in w(i) \cdot W_n, 2^k \leq i \leq 2^k + 2\} \leq \frac{1}{2}.$$


By (4.14), (4.15) and the Markov property of p -energies on graphs (Proposition 2.17),

$$\begin{aligned} \widetilde{\mathcal{C}}_p^{(n,M_k)} &= \mathcal{E}_p^{\widetilde{G}_{n,M_k}}(f_{n,k}) \\ &\geq \frac{1}{2} \left[\mathcal{E}_p^{\widetilde{G}_{n,M_k}} \left(f_{n,k} \vee \frac{1}{2} \right) + \mathcal{E}_p^{\widetilde{G}_{n,M_k}} \left(f_{n,k} \wedge \frac{1}{2} \right) \right] = 2^{-p} \widetilde{\mathcal{C}}_p^{(n,M_{k-1})}. \end{aligned}$$

Iterating this estimate, we conclude that, for any $k \in \mathbb{N}$,

$$\mathcal{C}_p^{(n,M_k)} \geq c_2^{-1} \widetilde{\mathcal{C}}_p^{(n,M_k)} \geq c_2^{-1} 2^{-pk} \widetilde{\mathcal{C}}_p^{(n,3)} \geq c_1^{-1} c_2^{-2} 2^{-pk} \mathcal{C}_{p,\text{face}}^{(n)}$$

Since $\mathcal{C}_p^{(n,M)} \geq \mathcal{C}_p^{(n,M')}$ for $M \leq M'$, we obtain the desired estimate for all M . \square

FIGURE 4. Values of $f_{n,k}$

Finally, we prove (KM_p) . We mainly follow the method in [19, Lemma 4.8].

Theorem 4.14. *Suppose Assumption 4.7 holds. Then (KM_p) holds.*

Proof. Let $f_n: W_n \rightarrow \mathbb{R}$ satisfy

$$\mathcal{E}_p^{G_n}(f_n) = 1, \quad f_n|_{\partial_* G_n} \equiv 0, \quad \text{and} \quad \langle f_n \rangle_{W_n} = \left(\lambda_{p, \text{Dir}}^{(n)} \right)^{1/p}.$$

Note that f_n is non-negative. Pick $w^* \in W_n$ such that $f_n(w^*) = \max_{w \in W_n} f_n(w)$. Then we easily see that $f_n(w^*) \geq \left(\lambda_{p, \text{Dir}}^{(n)} \right)^{1/p}$. Let $\theta, \tilde{C}_{\text{UH}} > 0$ be constants in Theorem 4.10 and choose $l_* \in \mathbb{N}$ such that

$$(4.16) \quad \left(\tilde{C}_{\text{UH}} a^{-l_* \theta} \right)^{1/p} \leq \frac{1}{8DN_*}.$$

Then, by Theorem 4.10, for any $n \geq l_*$,

$$(4.17) \quad \max\{|f_n(v) - f_n(w)| \mid k \leq n - l_*, z \in W_{l_*+k}, v, w \in z \cdot W_{n-l_*-k}\} \leq \frac{1}{8DN_*} \left(\lambda_{p, \text{Dir}}^{(n)} \right)^{1/p}.$$

Now, we will choose a chain of $(n-2)$ -cells from $\partial_* G_n$ to w^* . Define

$$S_{\text{corner}} := \left\{ \sum_{k=1}^D \sigma_k \mathbf{e}_k \mid \sigma_k \in \{-1, +1\} \right\}$$

that satisfies $\#S_{\text{corner}} = 2^D$. Fix a path $[i(1), \dots, i(L)]$ in G_1 , i.e. $i(k) \in S$ and $(i(k), i(k+1)) \in E_1$ for any l , such that $i(1) \in \partial_* G_1$ and $w^* \in i(L) \cdot W_{n-1}$. Note that $L \leq \text{diam } G_1 < N_*$. We can find ‘‘corners’’ $z(1), z(2), \dots, z(2L-1) \in \{i^{n-1} \mid i \in S_{\text{corner}}\}$ satisfying $v^* := i(1)z(1) \in \partial_* G_n$ and

$$(i(k)z(2k), i(k+1)z(2k+1)) \in E_n \quad \text{for } k = 1, \dots, L-1.$$

Since $f_n(v^*) = 0$ and $\mathcal{E}_p^{G_n}(f_n) = 1$,

$$\begin{aligned} \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{1/p} &\leq f_n(w^*) = f_n(w^*) - f_n(v^*) \\ &\leq |f_n(v^*) - f_n(w^*)| \\ &\leq \sum_{k=1}^{L-1} \left(|f_n(i(k)z(2k-1)) - f_n(i(k)z(2k))| \right. \\ &\quad \left. + |f_n(i(k)z(2k)) - f_n(i(k+1)z(2k+1))| \right) \\ &\leq \left\{ \sum_{k=1}^{L-1} |f_n(i(k)z(2k-1)) - f_n(i(k)z(2k))| \right\} + L - 1. \end{aligned}$$

Now, we have $\lambda_{p,\text{Dir}}^{(n)} \rightarrow \infty$ as $n \rightarrow \infty$ by Assumption 4.7 and Theorem 3.12. Hence we may assume that

$$\sum_{k=1}^{L-1} |f_n(i(k)z(2k-1)) - f_n(i(k)z(2k))| \geq \frac{1}{2} \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{1/p},$$

for all large n . Then there exists $l \in \{1, \dots, L-1\}$ such that

$$|f_n(i(l)z(2l-1)) - f_n(i(l)z(2l))| \geq \frac{1}{2(L-1)} \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{1/p}.$$

Moreover, by the triangle inequality, there exist $i^*(0), i^*(1) \in S_{\text{corner}}$ and $j \in \{1, \dots, D\}$ with $i^*(0) - i^*(1) \in \{-\mathbf{e}_j, \mathbf{e}_j\}$ such that

$$(4.18) \quad |f_n(i(l)i^*(0)^{n-1}) - f_n(i(l)i^*(1)^{n-1})| \geq \frac{1}{2D(L-1)} \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{1/p}.$$

By using the symmetries of f_n if necessary, we may assume that

$$f_n(i(l)i^*(1)^{n-1}) \geq f_n(i(l)i^*(0)^{n-1}).$$

Then we see from (4.17) and (4.18) that

$$\begin{aligned} \delta_n &:= \min_{i(l)i^*(1)^{l_*} \cdot W_{n-l_*-1}} f_n - \max_{i(l)i^*(0)^{l_*} \cdot W_{n-l_*-1}} f_n \\ &\geq \left[\frac{1}{2D(L-1)} - 2 \cdot \frac{1}{8DN_*} \right] \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{1/p} \geq \frac{1}{4DN_*} \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{1/p}, \end{aligned}$$

where we used a bound $L-1 \leq N_*$ in the last inequality. Let us consider the ‘‘horizontal chain of $(n-l_*-1)$ -cell along j -axis’’ from $i(l)i^*(0)^{l_*} \cdot W_{n-l_*-1}$ to $i(l)i^*(1)^{l_*} \cdot W_{n-l_*-1}$. Then, by putting $g_n := ((\delta_n^{-1} f_n + c_n) \vee 0) \wedge 1$, where c_n is a constant such that $g_n|_{i(l)i^*(\varepsilon)^{l_*} \cdot W_{n-l_*-1}} \equiv \varepsilon$ for $\varepsilon = 0, 1$, we have that

$$\mathcal{C}_p^{(n-l_*-1, a^{l_*})} \leq \mathcal{E}_p^{G_n}(g_n) \leq \delta_n^{-p} \mathcal{E}_p^{G_n}(f_n) \leq (4DN_*)^p \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{-1}.$$

From Lemma 4.12, Theorems 3.4 and 3.12-(3), we conclude that

$$\begin{aligned} \mathcal{C}_p^{(n)} &\leq C_{3.4} \mathcal{C}_p^{(l_*+1)} \mathcal{C}_p^{(n-l_*-1)} \\ &\leq C_{3.4} C(a^{l_*}) \mathcal{C}_p^{(l_*+1)} \mathcal{C}_p^{(n-l_*-1, a^{l_*})} \\ &\leq C_{3.4} C(a^{l_*}) \mathcal{C}_p^{(l_*+1)} (4DN_*)^p \left(\lambda_{p,\text{Dir}}^{(n)}\right)^{-1} \leq C_{\text{KM}} \left(\lambda_p^{(n)}\right)^{-1}, \end{aligned}$$

where $C_{\text{KM}} = C(a^{l_*})\mathcal{C}_p^{(l_*+1)}(4DN_*)^p C_{3.12}^{-1}$, which depends only on $p, a, D, L_*, N_*, \rho_p$. This proves (KM_p) . \square

We conclude this section by giving some useful estimates to construct p -energies. Under Assumption 4.7, we have multiplicative inequality of $\mathcal{R}_p^{(\cdot)}$ by virtue of Proposition 4.5, Theorems 4.14 and 3.13. In particular, $\mathcal{R}_p^{(n)} \asymp \rho_p^n$ for all $n \in \mathbb{N}$. Then Theorem 4.9) implies that

$$(4.19) \quad \beta_p := \frac{\log N_* \rho_p}{\log a}$$

satisfies $\beta_p - \alpha > 0$, where $\alpha = \log N_*/\log a$ is the Hausdorff dimension of (K, d) .

Now, we define the rescaled discrete p -energy $\tilde{\mathcal{E}}_p^{G_n}$ by setting

$$\tilde{\mathcal{E}}_p^{G_n}(f) := \rho_p^n \mathcal{E}_p^{G_n}(f), \quad f \in \mathbb{R}^{W_n}.$$

Similarly to Theorem 4.10, we can show the following Hölder estimate.

Theorem 4.15. *For every $n, m \in \mathbb{N}$, $z \in W_m$, $v, w \in \mathcal{B}_n(z, 1)$ and $f \in \mathbb{R}^{W_{n+m}}$,*

$$(4.20) \quad |f(v) - f(w)|^p \leq C_{\text{UH}} \tilde{\mathcal{E}}_p^{G_{n+m}}(f) a^{-(\beta_p - \alpha)m},$$

where $C_{\text{UH}} > 0$ is a constant depending only on $p, a, D, L_*, N_*, \rho_p$.

Proof. The required uniform Hölder estimate (4.20) can be proved by using $\mathcal{C}_p^{(n)} \lesssim a^{-(\beta_p - \alpha)n}$, which is a consequence of supermultiplicative inequality of $\mathcal{C}_p^{(\cdot)}$, instead of (4.10) in the proof of Theorem 4.10. \square

Lastly, we observe a monotonicity result (the so-called *weak monotonicity* in [29]). This is proved in [54, Proposition 5.2] for $p = 2$.

Corollary 4.16. *For every $n, m \in \mathbb{N}$ and $f \in L^p(K, \mu)$,*

$$(4.21) \quad \tilde{\mathcal{E}}_p^{G_n}(M_n f) \leq C_{\text{WM}} \tilde{\mathcal{E}}_p^{G_{n+m}}(M_{n+m} f),$$

where $C_{\text{WM}} > 0$ is a constant depending only on $p, a, D, L_*, N_*, \rho_p$. In particular,

$$(4.22) \quad \sup_{n \in \mathbb{N}} \tilde{\mathcal{E}}_p^{G_n}(M_n f) \leq C_{\text{WM}} \varliminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_p^{G_n}(M_n f).$$

Proof. By Proposition 2.11-(3), for any $n, m \in \mathbb{N}$, $w \in W_n$ and $f \in L^p(K, \mu)$,

$$P_{n+m,n}(M_{n+m} f)(w) = N_*^n \sum_{v \in w \cdot W_m} \int_{K_v} f d\mu = M_n f(w).$$

Hence $P_{n+m,n}(M_{n+m} f) = M_n f$. We get (4.21) from Lemma 3.3 and (3.21). \square

Remark 4.17. One can derive a uniform Harnack type estimate for discrete p -harmonic functions as an application of (KM_p) . For $p = 2$, this was done by [5, Theorem 3.1] or [54, Lemma 7.8]. We expect that such type estimate will be important for future work, but we omit this since its proof does not fit the purpose of this paper.

5. THE DOMAIN OF p -ENERGY

This section aims to prove a part of our main results: Theorems 2.21 and 2.22.

Let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet. Thanks to Corollary 4.16, we know that the following quantity:

$$(5.1) \quad |f|_{\mathcal{F}_p} := \sup_{n \in \mathbb{N}} \widetilde{\mathcal{E}}_p^{G_n}(M_n f)^{1/p}.$$

describes the limit behavior of rescaled p -energy $\widetilde{\mathcal{E}}_p^{G_n}(M_n f)$. Then we easily see that $|\cdot|_{\mathcal{F}_p}$ defines a ($[0, \infty]$ -valued) semi-norm. We also define a function space \mathcal{F}_p and its norm $\|\cdot\|_{\mathcal{F}_p}$ by setting

$$\mathcal{F}_p := \{f \in L^p(K, \mu) \mid |f|_{\mathcal{F}_p} < \infty\} \quad \text{and} \quad \|f\|_{\mathcal{F}_p} := \|f\|_{L^p} + |f|_{\mathcal{F}_p}.$$

Ideally, \mathcal{F}_p plays the same role as the Sobolev space $W^{1,p}$ in smooth settings like Euclidean spaces. As stated in [41, Section 7], this $(1, p)$ -“Sobolev” space \mathcal{F}_p should be *closable* and have *regularity*, that is,

- any Cauchy sequence $\{f_n\}_{n \geq 1}$ in $|\cdot|_{\mathcal{F}_p}$ with $f_n \rightarrow 0$ in L^p converges to 0 in \mathcal{F}_p ;
- $\mathcal{F}_p \cap \mathcal{C}(K)$ is dense in $\mathcal{C}(K)$ with respect to the supremum norm.

We prove these properties in subsection 5.1. In addition, the *separability* of \mathcal{F}_p is proved in subsection 5.2. The separability will be essential to follow our construction of p -energy in section 6. We also see in subsection 5.3 that \mathcal{F}_p has a Besov-like representation, which is an extension of results for \mathcal{F}_2 in [28, 51].

Throughout this section, we suppose Assumption 4.7 holds.

5.1. Closability and regularity. First, we derive the following Hölder estimate from the uniform Hölder estimates on graphical approximations (Theorem 4.15) in the same way as [44, Lemmas 6.10 and 6.13].

Theorem 5.1. *There exists a positive constant $C_{\text{Hö}}l$ (depending only on $p, a, D, L_*, N_*, \rho_p$) such that every $f \in \mathcal{F}_p$ has a continuous modification $f_* \in \mathcal{C}(K)$ with*

$$|f_*(x) - f_*(y)|^p \leq C_{\text{Hö}}l |f|_{\mathcal{F}_p}^p d(x, y)^{\beta_p - \alpha},$$

for every $x, y \in K$. Moreover, the inclusion map $\mathcal{F}_p \ni f \mapsto f_* \in \mathcal{C}(K)$ is injective. In particular, \mathcal{F}_p is continuously embedded in the Hölder space $\mathcal{C}^{0, (\beta_p - \alpha)/p}$.

Proof. Let $f \in \mathcal{F}_p$. By Proposition 2.11-(3) and Lemma 2.7, for each $n \geq 1$, we have that $\int_K f d\mu = N_*^{-n} \sum_{w \in W_n} M_n f(w)$. From this identity, there exists $w(n) \in W_n$ for each $n \in \mathbb{N}$ such that $M_n |f|(w(n)) \leq \int_K |f| d\mu$. Then, by Theorem 4.15, for any $n \in \mathbb{N}$ and $v \in W_n$,

$$\begin{aligned} |M_n f(v)|^p &\leq 2^{p-1} |M_n f(v) - M_n f(w(n))|^p + 2^{p-1} |M_n f(w(n))|^p \\ &\leq 2^{p-1} C_{\text{UH}} |f|_{\mathcal{F}_p}^p + 2^{p-1} \left(\int_K |f|^p d\mu \right)^p \\ &\leq 2^{p-1} C_{\text{UH}} |f|_{\mathcal{F}_p}^p + 2^{p-1} \|f\|_{L^p}^p, \end{aligned}$$

and hence we obtain the following uniform bound of $f \in \mathcal{F}_p$:

$$(5.2) \quad \sup_{n \geq 1} \max_{v \in W_n} |M_n f(v)| \leq c_1 \|f\|_{\mathcal{F}_p} < \infty,$$

where $c_1 > 0$ depends only on p and C_{UH} .

For each $n \in \mathbb{N}$, enumerate the elements W_n as $W_n = \{w_n(1), \dots, w_n(N_*^n)\}$ and inductively define $\{\widehat{K}_{w_n(i)}\}_{i=1}^{N_*^n}$ as follows: $\widehat{K}_{w_n(1)} := K_{w_n(1)}$ and

$$\widehat{K}_{w_n(i+1)} := K_{w_n(i+1)} \setminus \bigcup_{j \leq i} \widehat{K}_{w_n(j)}.$$

Note that each $\widehat{K}_{w_n(i)}$ is a Borel set of K , $\widehat{K}_{w_n(i)}$ ($i = 1, \dots, N_*^n$) are disjoint, and $K = \bigcup_{i=1}^{N_*^n} \widehat{K}_{w_n(i)}$. Also, by Proposition 2.11-(3), we have $\mu(K_w \setminus \widehat{K}_w) = 0$ for any $w \in W_n$. Next, define a Borel measurable function $f_n: K \rightarrow \mathbb{R}$ by setting

$$f_n := \sum_{w \in W_n} M_n f(w) \mathbb{1}_{\widehat{K}_w}.$$

Then (5.2) yields that

$$(5.3) \quad \sup_{n \geq 1} \sup_{x \in K} |f_n(x)| \leq c_1 \|f\|_{\mathcal{F}_p}.$$

Let $n \in \mathbb{N}$, let $x \neq y \in K$ and set $n_* := n(x, y) \in \mathbb{Z}_{\geq 0}$. If $n > n_*$, then there exist $v, w \in W_{n_*}$ such that $x \in K_v$, $y \in K_w$ and $K_v \cap K_w \neq \emptyset$. We can find $v', w' \in W_n$ such that $x \in \widehat{K}_{v'}$ and $y \in \widehat{K}_{w'}$. Then $v' \in \mathcal{B}_{n-n_*}(v, 1)$, $w' \in \mathcal{B}_{n-n_*}(w, 1)$ and $\mathcal{B}_{n-n_*}(v, 1) \cap \mathcal{B}_{n-n_*}(w, 1) \neq \emptyset$. Fix $z' \in \mathcal{B}_{n-n_*}(v, 1) \cap \mathcal{B}_{n-n_*}(w, 1)$. Applying Theorem 4.15, we have that

$$\begin{aligned} |f_n(x) - f_n(y)|^p &\leq 2^{p-1} (|M_n f(v') - M_n f(z')|^p + |M_n f(z') - M_n f(w')|^p) \\ &\leq 2^p C_{\text{UH}} \widetilde{\mathcal{E}}_p^{G_n}(M_n f) a^{-(\beta_p - \alpha)n_*} \\ &\leq 2^{p+1} a \sqrt{D} C_{\text{UH}} |f|_{\mathcal{F}_p}^p d(x, y)^{\beta_p - \alpha}, \end{aligned}$$

where we used Lemma 2.13 in the last line. If $n \leq n_*$, then there exist $v, w \in W_n$ such that $x \in K_v$, $y \in K_w$ and $K_v \cap K_w \neq \emptyset$. Let $v', w' \in W_n$ with $x \in \widehat{K}_{v'}$ and $y \in \widehat{K}_{w'}$. Then $[v', v, w, w']$ is a path in G_n , and hence we have that

$$\begin{aligned} |f_n(x) - f_n(y)|^p &\leq 3^{p-1} (|M_n f(v') - M_n f(v)|^p + |M_n f(v) - M_n f(w)|^p + |M_n f(w) - M_n f(w')|^p) \\ &\leq 3^p \mathcal{E}_p^{G_n}(M_n f) \leq 3^p |f|_{\mathcal{F}_p}^p \rho_p^{-n}. \end{aligned}$$

As a result of this observation, we conclude that

$$(5.4) \quad |f_n(x) - f_n(y)|^p \leq c_2 |f|_{\mathcal{F}_p}^p (d(x, y)^{\beta_p - \alpha} + \rho_p^{-n}), \quad f \in \mathcal{F}_p, n \in \mathbb{N}, x, y \in K,$$

where c_2 is a positive constant depending only on p, a, D, C_{UH} .

Thanks to (5.3) and (5.4), we can apply an Arzelà–Ascoli type argument for $\{f_n\}_{n \geq 1}$ (see [44, Lemma D.1]). For reader's convenience, we provide a complete proof. Set $A_n := \{F_w(\sum_{k=1}^D \sigma_k \mathbf{e}_k)\}_{\sigma_k \in \{-1, +1\}, w \in W_n}$ and $A := \bigcup_{n \geq 1} A_n$. Then A is a countable dense subset of K . Since $\{f_n(x)\}_{n \geq 1}$ is bounded for each $x \in A$ by (5.3), by a diagonal argument, we obtain a subsequence $\{n_k\}_{k \geq 1}$ such that $\{f_{n_k}(x)\}_{k \geq 1}$ converges as $k \rightarrow \infty$ for any $x \in A$. Define $f_*(x) := \lim_{k \rightarrow \infty} f_{n_k}(x)$ for any $x \in A$. From (5.4) and Assumption 4.7, we see that

$$|f_*(x) - f_*(y)|^p \leq c_2 |f|_{\mathcal{F}_p}^p d(x, y)^{\beta_p - \alpha} \quad \text{for any } x, y \in A.$$

Since A is dense in K , f_* is extended to a continuous function on K , which is again denoted by $f_* \in \mathcal{C}(K)$, and it follows that

$$|f_*(x) - f_*(y)|^p \leq c_2 |f|_{\mathcal{F}_p}^p d(x, y)^{\beta_p - \alpha} \quad \text{for any } x, y \in K.$$

(We can also show that $\sup_{x \in K} |f_*(x) - f_{n_k}(x)| \rightarrow 0$ as $k \rightarrow \infty$. For a proof, see [44, Lemma D.1].) Then, for any $m \in \mathbb{N}$ and $w \in W_m$, we have $\int_{K_w} f_{n_k} d\mu \rightarrow \int_{K_w} f_* d\mu$ as $k \rightarrow \infty$. By Proposition 2.11-(3) and $\mu(K_w \setminus \widehat{K}_w) = 0$,

$$\begin{aligned} \int_{K_w} f_{n_k} d\mu &= \int_{K_w} \sum_{z \in W_{n_k}} M_{n_k} f(z) \mathbf{1}_{\widehat{K}_z} d\mu = \int_{K_w} \sum_{z \in w \cdot W_{n_k-m}} M_{n_k} f(z) \mathbf{1}_{K_z} d\mu \\ &= \sum_{z \in w \cdot W_{n_k-m}} \frac{1}{\mu(K_z)} \left(\int_{K_z} f d\mu \right) \int_{K_w} \mathbf{1}_{K_z} d\mu = \int_{K_w} f d\mu, \end{aligned}$$

whenever $w \in W_m$ and $n_k > m$. Letting $k \rightarrow \infty$, we obtain $\int_{K_w} f_* d\mu = \int_{K_w} f d\mu$ for all $w \in W_\#$. By Proposition 2.11 and Dynkin's π - λ theorem, we conclude that f_* is a continuous modification of f . The injectivity of $f \mapsto f_*$ is obvious. We complete the proof. \square

Next, we prove the closability by proving that $(\mathcal{F}_p, \|\cdot\|_{\mathcal{F}_p})$ is complete. See also [44, Lemmas 6.15 and 6.16].

Theorem 5.2. *$(\mathcal{F}_p, \|\cdot\|_{\mathcal{F}_p})$ is a Banach space.*

Proof. Let $\{f_n\}_{n \geq 1}$ be a Cauchy sequence in $(\mathcal{F}_p, \|\cdot\|_{\mathcal{F}_p})$. Then $\{f_n\}_{n \geq 1}$ converges to some $f \in L^p(K, \mu)$ in L^p . Fix $x_0 \in K$ and set $g_n := f_n - f_n(x_0)$. Then, by the Hölder estimate in Theorem 5.1, for all $n, m \geq 1$ and $x \in K$,

$$|g_n(x) - g_m(x)|^p \leq C_{\text{Hö}} |f_n - f_m|_{\mathcal{F}_p}^p d(x, x_0)^{\beta_p - \alpha} \leq C_{\text{Hö}} |f_n - f_m|_{\mathcal{F}_p}^p,$$

which implies that $\{g_n\}_{n \geq 1}$ is a Cauchy sequence in $\mathcal{C}(K)$. Since $\mathcal{C}(K)$ is complete, $\{g_n\}_{n \geq 1}$ converges to some $g \in \mathcal{C}(K)$ in the supremum norm.

It is immediate that $\{f_n - g_n\}_{n \geq 1}$ converges to $f - g$ in L^p , and thus we can pick a subsequence $\{n_k\}_{k \geq 1}$ so that $f_{n_k} - g_{n_k} \rightarrow f - g$ for μ -a.e. as $k \rightarrow \infty$. On the other hand, the definition of g_n implies that $f_n - g_n \equiv f_n(x_0)$. Hence the limit $\lim_{k \rightarrow \infty} f_{n_k}(x_0) =: b$ exists and $f - g = b$ for μ -a.e. In particular, f admits a continuous modification. We again write f to denote this continuous version. Then f is the limit of f_{n_k} . Indeed, we have

$$\|f - f_{n_k}\|_{\mathcal{C}(K)} \leq \|g - g_{n_k}\|_{\mathcal{C}(K)} + |f_{n_k}(x_0) - b| \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Since $\{f_n\}_{n \geq 1}$ is a Cauchy sequence in \mathcal{F}_p , for any $\varepsilon > 0$ there exists $N(\varepsilon) \geq 1$ such that

$$\sup_{i \wedge j \geq N(\varepsilon)} \sup_{k \geq 1} \widetilde{\mathcal{E}}_p^{G_k}(M_k f_{n_i} - M_k f_{n_j}) \leq \varepsilon,$$

which implies that

$$(5.5) \quad \sup_{i \geq N(\varepsilon)} \sup_{k \geq 1} \widetilde{\mathcal{E}}_p^{G_k}(M_k f_{n_i} - M_k f) \leq \varepsilon.$$

Therefore, we have that, for large $i \geq 1$ with $n_i \geq N(\varepsilon)$,

$$\widetilde{\mathcal{E}}_p^{G_k}(M_k f)^{1/p} \leq \widetilde{\mathcal{E}}_p^{G_k}(M_k f_{n_i} - M_k f)^{1/p} + \widetilde{\mathcal{E}}_p^{G_k}(M_k f_{n_i})^{1/p} \leq \varepsilon + \sup_{n \geq 1} |f_n|_{\mathcal{F}_p},$$

which implies that $f \in \mathcal{F}_p$. In addition, (5.5) yields that $\|f - f_{n_i}\|_{\mathcal{F}_p} \rightarrow 0$ as $i \rightarrow \infty$.

The convergence $\|f - f_n\|_{\mathcal{F}_p} \rightarrow 0$ is easily derived by applying the above arguments for any subsequence of $\{f_n\}_{n \geq 1}$. We complete the proof. \square

Moreover, we can show that \mathcal{F}_p is compactly embedded in $L^p(K, \mu)$.

Proposition 5.3. *The inclusion map from \mathcal{F}_p to $L^p(K, \mu)$ is a compact operator.*

Proof. Let $\{f_n\}_{n \geq 1}$ be a bounded sequence in \mathcal{F}_p . Since the embedding of \mathcal{F}_p in $\mathcal{C}^{0,(\beta_p - \alpha)/p}$ is continuous, we obtain a subsequence $\{f_{n_k}\}_{k \geq 1}$ and $f \in \mathcal{C}(K)$ such that f_{n_k} converges to f in the supremum norm by applying the Arzelá–Ascoli theorem. This proves our assertion. \square

Towards the regularity of \mathcal{F}_p , the following lemma gives a “partition of unity” in \mathcal{F}_p . See also [44, Lemma 6.18].

Lemma 5.4. *There exists a family $\{\varphi_w\}_{w \in W_\#}$ in \mathcal{F}_p such that*

- (1) for any $w \in W_\#$, $0 \leq \varphi_w \leq 1$;
- (2) for any $n \in \mathbb{N}$, $\sum_{w \in W_n} \varphi_w \equiv 1$;
- (3) for any $n \in \mathbb{N}$ and $w \in W_n$, $\text{supp}[\varphi_w] \subseteq U_1^{(n)}(w)$, where $U_1^{(n)}(w)$ is defined as

$$(5.6) \quad U_1^{(n)}(w) := \bigcup_{v \in W_n; d_{G_n}(v, w) \leq 1} K_v;$$

- (4) there exists a constant $C_{5.4} > 0$ (depending only on $p, a, D, L_*, N_*, \rho_p$) such that

$$|\varphi_w|_{\mathcal{F}_p}^p \leq C_{5.4} \rho_p^n \quad \text{for any } n \in \mathbb{N} \text{ and } w \in W_n.$$

Proof. For $n, m \in \mathbb{N}$ and $w \in W_m$, let $\psi_w^{(n)} : W_{n+m} \rightarrow [0, 1]$ satisfy $\psi_w^{(n)}|_{w \cdot W_n} \equiv 1$, $\psi_w^{(n)}|_{W_{n+m} \setminus \mathcal{B}_n(w, 1)} \equiv 0$ and $\mathcal{E}_p^{G_{n+m}}(\psi_w^{(n)}) = \mathcal{C}_p^{G_{n+m}}(w \cdot W_n, W_{n+m} \setminus \mathcal{B}_n(w, 1))$. Define $\Psi_m^{(n)} := \left(\sum_{v \in W_m} \psi_v^{(n)} \right)^{-1}$ and $\varphi_w^{(n)} := \Psi_m^{(n)} \psi_w^{(n)}$. Note that $\varphi_w^{(n)}$ coincides with the function denoted by the same symbol in the proof of Lemma 3.10. We also set $\tilde{\varphi}_w^{(n)} : K \rightarrow \mathbb{R}$ by setting

$$\tilde{\varphi}_w^{(n)} := \sum_{z \in W_{n+m}} \varphi_w^{(n)}(z) \mathbf{1}_{\hat{K}_z},$$

where $\{\hat{K}_z\}_{z \in W_\#}$ is the same as in the proof of Theorem 5.1. Then $M_{n+m} \tilde{\varphi}_w^{(n)} = \varphi_w^{(n)}$ and, from (3.12) in the proof of Lemma 3.10 and (3.21), we have $\tilde{\mathcal{E}}_p^{G_{n+m}}(\varphi_w^{(n)}) \leq c_1 \rho_p^m$ for all $w \in W_m$ and $n \in \mathbb{N}$, where $c_1 > 0$ depends only on p, a, D, N_*, L_* . In particular, by Theorem 5.1, we obtain

$$\left| \tilde{\varphi}_w^{(n)}(x) - \tilde{\varphi}_w^{(n)}(y) \right|^p \leq c_1 C_{\text{Hö}} \rho_p^m d(x, y)^{\beta_p - \alpha},$$

for $x, y \in K$ with $n(x, y) < n + m$. Similarly to the Arzelá–Ascoli type argument in the proof of Theorem 5.1, we can find a subsequence $\{n_k\}_{k \geq 1}$ and a continuous function $\varphi_w \in \mathcal{C}(K)$ such that $\lim_{k \rightarrow \infty} \tilde{\varphi}_w^{(n_k)}(x) = \varphi_w(x)$ for any $x \in K$ and

$$|\varphi_w(x) - \varphi_w(y)|^p \leq c_1 C_{\text{Hö}} \rho_p^{|w|} d(x, y)^{\beta_p - \alpha} \quad \text{for any } x, y \in K.$$

Then the properties (1), (2) and (3) are immediate from this convergence and the associated properties of $\tilde{\varphi}_w^{(n)}$, so it will suffice to show (4). By the weak monotonicity (Corollary 4.16),

$$\begin{aligned} \tilde{\mathcal{E}}_p^{G_l} \left(M_l \tilde{\varphi}_w^{(n_k)} \right) &\leq C_{\text{WM}} \tilde{\mathcal{E}}_p^{G_{n_k+m}} \left(M_{n_k+m} \tilde{\varphi}_w^{(n_k)} \right) \\ &\leq C_{\text{WM}} \sup_{n \geq 1} \tilde{\mathcal{E}}_p^{G_{n+m}} \left(\varphi_w^{(n)} \right) \\ &\leq c_1 C_{\text{WM}} \rho_p^m, \end{aligned}$$

whenever $l \leq n_k + m$. Passing to the limit $k \rightarrow \infty$ and supremum over $l \in \mathbb{N}$ in this estimate, we conclude that $|\varphi_w|_{\mathcal{F}_p}^p \leq c_1 C_{\text{WM}} \rho_p^m$ for all $m \geq 1$ and $w \in W_m$ and complete the proof. \square

Now, define a subspace \mathcal{H}_p^* of \mathcal{F}_p by setting

$$(5.7) \quad \mathcal{H}_p^* := \left\{ \sum_{w \in A} a_w \varphi_w \mid A \text{ is a finite subset of } W_\#, a_w \in \mathbb{R} \text{ for each } w \in A \right\},$$

where $\{\varphi_w\}_{w \in W_\#}$ is a family of functions in \mathcal{F}_p appeared in Lemma 5.4. Then we achieve the regularity of \mathcal{F}_p (see also [44, Lemma 6.19]).

Theorem 5.5. *The space \mathcal{H}_p^* is dense in $\mathcal{C}(K)$ with respect to the sup norm. In particular, \mathcal{F}_p is dense in $\mathcal{C}(K)$.*

Proof. Let $f \in \mathcal{C}(K)$ and define f_n by setting $f_n := \sum_{w \in W_n} M_n f(w) \varphi_w \in \mathcal{H}_p^*$. Then

$$\begin{aligned} |f(x) - f_n(x)| &\leq L_*^2 \max_{w \in W_n, x \in \text{supp}[\varphi_w]} |f(x) - M_n f(w)| \\ &\leq L_*^2 \max_{w \in W_n} \sup_{x \in U_1^{(n)}(w), y \in K_w} |f(x) - f(y)|. \end{aligned}$$

Since $\max_{w \in W_n} \text{diam}(U_1^{(n)}(w), d) \leq 2a^{-n} \rightarrow 0$ as $n \rightarrow \infty$ and f is uniformly continuous, we have $\|f - f_n\|_{\mathcal{C}(K)} \rightarrow 0$. \square

5.2. Separability. In this subsection, we prove that \mathcal{F}_p is separable with respect to $\|\cdot\|_{\mathcal{F}_p}$. In the case $p = 2$, this is done by applying easy functional analytic arguments since the polarization formula of \mathcal{E}_2 yields a non-negative definite closed quadratic form. For example, by Proposition 5.3, the inclusion map from \mathcal{F}_2 to $L^2(K, \mu)$ is a compact operator, and thus there exists a countable complete orthonormal system of \mathcal{F}_2 (see [23, Exercise 4.2 and Corollary 4.2.3] for example). One can also give a short proof of the separability of \mathcal{F}_2 using resolvents (see [25, proof of Theorem 1.4.2-(iii)] for example). However, it is hopeless to execute similar arguments in our setting.

To overcome this difficulty, we directly show that the space \mathcal{H}_p^* defined in (5.7) is dense in \mathcal{F}_p and hence \mathbb{Q} -hull of $\{\varphi_w\}_{w \in W_\#}$ is also dense. Our strategy is standard in calculus of variations, namely, we extract a strong convergent approximation from \mathcal{H}_p^* by using Mazur's lemma. To this end, it will be a key ingredient to ensure the reflexivity of \mathcal{F}_p , which is deduced from a combination of *Clarkson's inequality* and the Milman–Pettis theorem. We will derive Clarkson's inequality by using Γ -convergence to find a norm of \mathcal{F}_p having the required properties.

We start by recalling Clarkson's inequality.

Definition 5.6 (Clarkson’s inequality). Let \mathcal{X} be a vector space (over \mathbb{R}) and let $\|\cdot\|$ be a semi-norm on \mathcal{X} . The semi-norm $\|\cdot\|$ satisfies *Clarkson’s inequality* if and only if one of the following holds:

- (1) There exists $p \in (1, 2]$ such that for every $x, y \in \mathcal{X}$,

$$\|x + y\|^{\frac{p}{p-1}} + \|x - y\|^{\frac{p}{p-1}} \leq 2(\|x\|^p + \|y\|^p)^{\frac{1}{p-1}};$$

- (2) There exists $p \in [2, \infty)$ such that for every $x, y \in \mathcal{X}$,

$$\|x + y\|^p + \|x - y\|^p \leq 2^{p-1}(\|x\|^p + \|y\|^p).$$

It is well-known that L^p -norm on a measurable space satisfies Clarkson’s inequality, and that a normed space satisfying Clarkson’s inequality is *uniformly convex*. Recall that Milman–Pettis theorem says that any Banach spaces that possess a uniformly convex norm is reflexive.

Next, let us recall the definition of Γ -convergence and its basic properties. The reader is referred to [22] for details on Γ -convergence.

Definition 5.7 (Γ -convergence). Let $\{\Phi_n\}_{n \geq 1}$ be a sequence of $[-\infty, \infty]$ -valued functional on $L^p(K, \mu)$. We say that a functional $\Phi: L^p(K, \mu) \rightarrow [-\infty, \infty]$ is a Γ -limit of $\{\Phi_n\}_{n \geq 1}$ as $n \rightarrow \infty$ if the following two inequalities hold;

- (1) (liminf inequality) If $f_n \rightarrow f$ in L^p , then $\Phi(f) \leq \underline{\lim}_{n \rightarrow \infty} \Phi_n(f_n)$.
(2) (limsup inequality) For any $f \in L^p(K, \mu)$, there exists a sequence $\{f_n\}_{n \geq 1}$ such that

$$(5.8) \quad f_n \rightarrow f \text{ in } L^p \text{ and } \overline{\lim}_{n \rightarrow \infty} \Phi_n(f_n) \leq \Phi(f).$$

A sequence $\{f_n\}_{n \geq 1}$ satisfying (5.8) is called a *recovery sequence* of f .

Since (K, d) is a separable metric space, the following fact holds.

Theorem 5.8 ([22, Theorem 8.5]). *Let $\{\Phi_n\}_{n \geq 1}$ be a sequence of functionals on $L^p(K, \mu)$. Then there exists a subsequence $\{n_k\}_{k \geq 1}$ and a functional Φ on $L^p(K, \mu)$ such that Φ is a Γ -limit of $\{\Phi_{n_k}\}_{k \geq 1}$.*

Now, we regard $\tilde{\mathcal{E}}_p^{G_n}(\cdot)$ as a $[0, \infty]$ -valued functional on $L^p(K, \mu)$ defined by $f \mapsto \tilde{\mathcal{E}}_p^{G_n}(M_n f)$. Then, by Theorem 5.8, there exists a Γ -convergent subsequence $\{\tilde{\mathcal{E}}_p^{G_{n_k}}(\cdot)\}_{k \geq 1}$ and we write $E_p(\cdot)$ to denote its Γ -limit. We define $\|\!\| \cdot \|\!\|_{\mathcal{F}_p} := (\|\cdot\|_{L^p}^p + E_p(\cdot))^{1/p}$. This new “norm” $\|\!\| \cdot \|\!\|_{\mathcal{F}_p}$ establishes the reflexivity. (We also need to show that $\|\!\| \cdot \|\!\|_{\mathcal{F}_p}$ is a norm.)

Theorem 5.9. *The norm $\|\!\| \cdot \|\!\|_{\mathcal{F}_p}$ is equivalent to $\|\cdot\|_{\mathcal{F}_p}$ and satisfies Clarkson’s inequality. In particular, the Banach space \mathcal{F}_p is reflexive.*

Proof. Let $f, g \in L^p(K, \mu)$ and let $\{f_n\}_{n \geq 1}, \{g_n\}_{n \geq 1}$ be their recovery sequences throughout the proof. To verify the triangle inequality of $\|\!\| \cdot \|\!\|_{\mathcal{F}_p}$, define

$$\|f\|_{p,n} := \left(\|f\|_{L^p}^p + \tilde{\mathcal{E}}_p^{G_n}(M_n f) \right)^{1/p}.$$

Note that the Γ -limit of $\{\|\cdot\|_{p,n_k}\}_{k \geq 1}$ coincides with $\|\!\| \cdot \|\!\|_{\mathcal{F}_p}$ and that the norm $\|\cdot\|_{p,n}$ can be regarded as a suitable L^p -norm on $K \sqcup E_n$, where \sqcup denotes the

disjoint union. Using the triangle inequality of $\|\cdot\|_{p,n}$, we see that

$$\begin{aligned} \|f + g\|_{\mathcal{F}_p} &\leq \liminf_{k \rightarrow \infty} \|f_{n_k} + g_{n_k}\|_{p,n_k} \leq \overline{\lim}_{k \rightarrow \infty} \|f_{n_k}\|_{p,n_k} + \overline{\lim}_{k \rightarrow \infty} \|g_{n_k}\|_{p,n_k} \\ &\leq \|f\|_{\mathcal{F}_p} + \|g\|_{\mathcal{F}_p}, \end{aligned}$$

and thus $\|\cdot\|_{\mathcal{F}_p}$ is an extended norm on $L^p(K, \mu)$ (we admit $\|f\|_{\mathcal{F}_p} = \infty$).

Next, we prove $C_{\text{WM}}^{-1} |f|_{\mathcal{F}_p}^p \leq \mathbf{E}_p(f) \leq |f|_{\mathcal{F}_p}^p$ for every $f \in L^p(K, \mu)$ to conclude that $\|\cdot\|_{\mathcal{F}_p}$ and $\|\cdot\|_{p,n}$ are equivalent. From the liminf inequality, we immediately have that $\mathbf{E}_p(f) \leq |f|_{\mathcal{F}_p}^p$ for $f \in L^p(K, \mu)$. To prove the converse, note that $M_n f_{n_k}(w) \rightarrow M_n f(w)$ for any $w \in W_n$ as $k \rightarrow \infty$ by the dominated convergence theorem. By the weak monotonicity (Corollary 4.16), we obtain

$$\tilde{\mathcal{E}}_p^{G_n}(M_n f) \leq C_{\text{WM}} \liminf_{k \rightarrow \infty} \tilde{\mathcal{E}}_p^{G_{n_k}}(M_{n_k} f_{n_k}) \leq C_{\text{WM}} \mathbf{E}_p(f)$$

for all $n \geq 1$. We thus conclude that $|f|_{\mathcal{F}_p}^p \leq C_{\text{WM}} \mathbf{E}_p(f)$.

The rest of the proof is mainly devoted to Clarkson's inequalities. Let $p \leq 2$. Recall that $\|\cdot\|_{p,n}$ can be regarded a suitable L^p -norm on $K \sqcup E_n$. By Clarkson's inequality for $\|\cdot\|_{p,n}$, we have

$$\|f + g\|_{p,n}^{\frac{p}{p-1}} + \|f - g\|_{p,n}^{\frac{p}{p-1}} \leq 2(\|f\|_{p,n}^p + \|g\|_{p,n}^p)^{\frac{1}{p-1}}.$$

Thus, we see that

$$\begin{aligned} \|f + g\|_{\mathcal{F}_p}^{\frac{p}{p-1}} + \|f - g\|_{\mathcal{F}_p}^{\frac{p}{p-1}} &\leq \liminf_{k \rightarrow \infty} \|f_{n_k} + g_{n_k}\|_{p,n_k}^{\frac{p}{p-1}} + \liminf_{k \rightarrow \infty} \|f_{n_k} - g_{n_k}\|_{p,n_k}^{\frac{p}{p-1}} \\ &\leq \overline{\lim}_{k \rightarrow \infty} (\|f_{n_k} + g_{n_k}\|_{p,n_k}^{\frac{p}{p-1}} + \|f_{n_k} - g_{n_k}\|_{p,n_k}^{\frac{p}{p-1}}) \\ &\leq 2 \overline{\lim}_{k \rightarrow \infty} (\|f_{n_k}\|_{p,n_k}^p + \|g_{n_k}\|_{p,n_k}^p)^{\frac{1}{p-1}} \\ &\leq 2 \left(\overline{\lim}_{k \rightarrow \infty} \|f_{n_k}\|_{p,n_k}^p + \overline{\lim}_{k \rightarrow \infty} \|g_{n_k}\|_{p,n_k}^p \right)^{\frac{1}{p-1}} \\ &\leq 2(\|f\|_{\mathcal{F}_p}^p + \|g\|_{\mathcal{F}_p}^p)^{\frac{1}{p-1}}, \end{aligned}$$

which is Clarkson's inequality of $\|\cdot\|_{\mathcal{F}_p}$ when $p \leq 2$. Similarly, we get Clarkson's inequality for $p \geq 2$.

Consequently, we get a new norm $\|\cdot\|_{\mathcal{F}_p}$ of \mathcal{F}_p satisfying Clarkson's inequality. Thus, we see that the Banach space $(\mathcal{F}_p, \|\cdot\|_{\mathcal{F}_p})$ is uniformly convex (see [17, proof of Theorem 4.10 and its remark] for example). Therefore, we finish the proof by the Milman–Pettis theorem (see [17, Theorem 3.31] for example). \square

Theorem 5.10. *The space \mathcal{H}_p^* defined in (5.7) is dense in \mathcal{F}_p . Furthermore, \mathcal{F}_p is separable.*

Proof. Recall the definition of $\tilde{\varphi}_w^{(n)}$ in the construction of φ_w (see the proof of Lemma 5.4). By the diagonal procedure, we can pick a subsequence $\{n_k\}_{k \geq 1}$ such that $\{\tilde{\varphi}_w^{(n_k)}\}_{k \geq 1}$ converges to φ_w with respect to the supremum norm for all $w \in W_{\#}$. Next, for $f \in \mathcal{F}_p$, we define f_n and $f_n^{(k)}$ by setting

$$f_n := \sum_{w \in W_n} M_n f(w) \varphi_w, \quad f_n^{(k)} := \sum_{w \in W_n} M_n f(w) \tilde{\varphi}_w^{(n_k)}.$$

Similarly to the proof of Theorem 5.5, we see that $\{f_n\}_{n \geq 1}$ converges to f with respect to the supremum norm. Also, by Lemma 3.9, we obtain $\tilde{\mathcal{E}}_p^{G_{n+n_k}}(M_{n+n_k}f_n^{(k)}) \leq c_1 |f|_{\mathcal{F}_p}^p$, where $c_1 > 0$ depends only on p, a, D, N_*, L_* . Thus, by the weak monotonicity (Corollary 4.16), it holds that $\tilde{\mathcal{E}}_p^{G_l}(M_l f_n^{(k)}) \leq c_1 C_{\text{WM}} |f|_{\mathcal{F}_p}^p$ whenever $l \leq n + n_k$. Letting $k \rightarrow \infty$, we see that $\{f_n\}_{n \geq 1}$ is bounded in \mathcal{F}_p . Since \mathcal{F}_p is reflexive, we may assume that a subsequence $\{f_{n_k}\}_{k \geq 1}$ converges to f with respect to the weak topology of \mathcal{F}_p . By Mazur's lemma (see [17, Corollary 3.8] for example), we can find a sequence $\{g_l\}_{l \geq 1}$ such that each g_l is a convex combination of $\{f_{n_k}\}_{k \geq 1}$ and $g_l \rightarrow f$ in \mathcal{F}_p as $l \rightarrow \infty$. In particular, we obtain $\overline{\mathcal{H}_p^*}^{\|\cdot\|_{\mathcal{F}_p}} = \mathcal{F}_p$. Clearly, the \mathbb{Q} -hull of $\{\varphi_w\}_{w \in W_\#}$ also gives this approximation, that is,

$$\overline{\left\{ \sum_{w \in A} a_w \varphi_w \mid A \text{ is a finite subset of } W_\# \text{ and } a_w \in \mathbb{Q} (w \in A) \right\}}^{\|\cdot\|_{\mathcal{F}_p}} = \overline{\mathcal{H}_p^*}^{\|\cdot\|_{\mathcal{F}_p}}.$$

Therefore, \mathcal{F}_p is separable. \square

5.3. Lipschitz–Besov type expression. This subsection is devoted to proving Theorem 2.22. Let us start by introducing the definition of Lipschitz–Besov spaces on the underlying generalized Sierpiński carpet (K, d, μ) in some specific cases (see [12] for example).

Definition 5.11. For $s \in (0, \infty)$ and $p \in [1, \infty)$, the *Lipschitz–Besov space* $\Lambda_{p,\infty}^s$ is defined as

$$\Lambda_{p,\infty}^s := \{f \in L^p(K, \mu) \mid |f|_{\Lambda_{p,\infty}^s} < \infty\},$$

where

$$|f|_{\Lambda_{p,\infty}^s} := \sup_{n \in \mathbb{N}} \left(\int_K \int_{B_d(x, a^{-n})} \frac{|f(x) - f(y)|^p}{a^{-nsp}} d\mu(y) d\mu(x) \right)^{1/p}.$$

We also define its norm $\|\cdot\|_{\Lambda_{p,\infty}^s}$ by setting $\|f\|_{\Lambda_{p,\infty}^s} := \|f\|_{L^p} + |f|_{\Lambda_{p,\infty}^s}$.

Then $(\Lambda_{p,\infty}^s, \|\cdot\|_{\Lambda_{p,\infty}^s})$ is a Banach space. Furthermore, for any $c \in [1, \infty)$ there exists a positive constant $C_{\text{LB}}(c)$, which depends only on c, a, N_*, C_{AR} , such that, for any $f \in L^p(K, \mu)$,

$$(5.9) \quad |f|_{\Lambda_{p,\infty}^s}^p \leq C_{\text{LB}}(c) \sup_{n \in \mathbb{N}} \int_K \int_{B_d(x, ca^{-n})} \frac{|f(x) - f(y)|^p}{a^{-nsp}} d\mu(y) d\mu(x),$$

and

$$(5.10) \quad \begin{aligned} C_{\text{LB}}(c)^{-1} \overline{\lim}_{n \rightarrow \infty} \int_K \int_{B_d(x, ca^{-n})} \frac{|f(x) - f(y)|^p}{a^{-nsp}} d\mu(y) d\mu(x) \\ \leq \overline{\lim}_{r \downarrow 0} \int_K \int_{B_d(x, r)} \frac{|f(x) - f(y)|^p}{r^{sp}} d\mu(y) d\mu(x) \\ \leq C_{\text{LB}}(c) \overline{\lim}_{n \rightarrow \infty} \int_K \int_{B_d(x, ca^{-n})} \frac{|f(x) - f(y)|^p}{a^{-nsp}} d\mu(y) d\mu(x). \end{aligned}$$

First, we prove a (p, p) -Poincaré inequality in the sense of Kumagai and Sturm (see [53, pp. 315]). Recall that $\alpha = \log N_*/\log a$ denotes the Hausdorff dimension of (K, d) , and that $\beta_p = \log(N_* \rho_p)/\log a$, where ρ_p is the resistance scaling factor (see Theorem 3.4 and (3.21)).

Lemma 5.12. *There exists a positive constant $C_{\text{PI-KS}}$ (depending only on $p, a, D, L_*, N_*, \rho_p, C_{\text{AD}}$) such that*

$$(5.11) \quad a^{\beta p n} \sum_{w \in A} \int_{K_w} |f(x) - M_n f(w)|^p d\mu(x) \leq C_{\text{PI-KS}} \varliminf_{l \rightarrow \infty} \tilde{\mathcal{E}}_{p, A, W_l}^{G_{l+n}}(M_{l+n} f),$$

for every $n \in \mathbb{N}$, $f \in \mathcal{F}_p$ and subset $A \subseteq W_n$. In particular,

$$a^{\beta p n} \sum_{w \in W_n} \int_{K_w} |f(x) - M_n f(w)|^p d\mu(x) \leq C_{\text{PI-KS}} |f|_{\mathcal{F}_p}^p.$$

Proof. Let $f \in \mathcal{F}_p$ be the continuous version. Then, by the mean value theorem, for any $n \in \mathbb{N}$ and $w \in W_n$ there exists $x_w \in K_w$ such that $f(x_w) = M_n f(w)$. From the Hölder estimate (Theorem 5.1), we have that, for any $x \in K_w$,

$$\begin{aligned} |f(x) - M_n f(w)|^p &= |F_w^* f(F_w^{-1}(x)) - F_w^* f(F_w^{-1}(x_w))|^p \\ &\leq C_{\text{HöI}} |F_w^* f|_{\mathcal{F}_p}^p d(F_w^{-1}(x), F_w^{-1}(x_w))^{\beta p - \alpha} \\ &\leq C_{\text{HöI}} \text{diam}(K, d)^{\beta p - \alpha} |F_w^* f|_{\mathcal{F}_p}^p. \end{aligned}$$

Consequently, we obtain

$$\int_{K_w} |f(x) - M_n f(w)|^p d\mu(x) \leq C_{\text{HöI}} a^{-\alpha n} |F_w^* f|_{\mathcal{F}_p}^p.$$

Summing over $w \in W_n$, we conclude that

$$\begin{aligned} \sum_{w \in A} \int_{K_w} |f(x) - M_n f(w)|^p d\mu(x) &\leq C_{\text{HöI}} a^{-\alpha n} \sum_{w \in A} |F_w^* f|_{\mathcal{F}_p}^p \\ &\leq C_{\text{HöI}} C_{\text{WM}} a^{-\alpha n} \sum_{w \in A} \varliminf_{l \rightarrow \infty} \tilde{\mathcal{E}}_p^{G_l}(M_l(F_w^* f)) \\ &\leq C_{\text{HöI}} C_{\text{WM}} a^{-\alpha n} \varliminf_{l \rightarrow \infty} \sum_{w \in A} \tilde{\mathcal{E}}_p^{G_l}(M_l(F_w^* f)), \end{aligned}$$

where we used the weak monotonicity (Corollary 4.16) in the second line. From (2.3), we see that $\sum_{w \in A} \tilde{\mathcal{E}}_p^{G_l}(M_l(F_w^* f)) \leq \rho_p^{-n} \tilde{\mathcal{E}}_{p, A, W_l}^{G_{l+n}}(M_{l+n} f)$. In particular,

$$a^{-\alpha n} \varliminf_{l \rightarrow \infty} \sum_{w \in A} \tilde{\mathcal{E}}_p^{G_l}(M_l(F_w^* f)) \leq a^{-\beta p n} \varliminf_{l \rightarrow \infty} \tilde{\mathcal{E}}_{p, A, W_l}^{G_{l+n}}(M_{l+n} f),$$

which proves (5.11). \square

Next, we give an extension of [28, Theorem 4.11-(iii)]. This is essentially proved in [2, Theorem 5.1], so its proof is omitted here. Since the function space $\mathbf{B}^{p, \alpha}$ in [2, Theorem 5.1] is defined using heat kernels, we give a direct proof of the lemma in Appendix A.2 for reader's convenience.

Lemma 5.13. *Let $\beta > \alpha$ and $p > 1$. Then there exists a positive constant $C_{5.13}$ (depending only on $p, \beta, a, N_*, C_{\text{AR}}$) such that*

$$\begin{aligned} &|f(x) - f(y)|^p \\ &\leq C_{5.13} d(x, y)^{\beta - \alpha} \sup_{r \in (0, 3d(x, y)]} r^{-\beta} \int_K \int_{B_d(z, r)} |f(z) - f(z')|^p d\mu(z') d\mu(z), \end{aligned}$$

for every $f \in \Lambda_{p,\infty}^{\beta/p}$ and μ -a.e. $x, y \in K$. In particular, for any $g \in \mathcal{C}(K)$ and $x, y \in K$,

$$|g(x) - g(y)|^p \leq C_{5.13} |g|_{\Lambda_{p,\infty}^{\beta/p}}^p d(x, y)^{\beta-\alpha}.$$

An important consequence of the above lemma is the following type “ (p, p) -Poincaré inequality”.

Lemma 5.14. *Let $\beta > \alpha$ and $p > 1$. Then there exists a positive constant $C_{\text{PI-LB}}$ (depending only on $p, \beta, a, D, N_*, C_{\text{AR}}$) such that*

$$(5.12) \quad a^{\beta n} \sum_{w \in W_n} \int_{K_w} |f(x) - M_n f(w)|^p d\mu(x) \\ \leq C_{\text{PI-LB}} \sup_{r \in (0, 3a^{-n}]} r^{-\beta} \int_K \int_{B_d(x, r)} |f(x) - f(y)|^p d\mu(y) d\mu(x),$$

for every $n \in \mathbb{N}$ and $f \in \mathcal{C}(K)$.

Proof. We adopt a method in [29, proof of Theorem 3.5] and generalize it to fit our context. Let $f \in \Lambda_{p,\infty}^{\beta/p}$, let $w = w_1 \cdots w_n \in W_n$ and fix $k \in \mathbb{N}$ that we choose later. Then, by the mean value theorem, there exists $x_w \in K_w$ such that $M_n f(w) = f(x_w)$. Let $\omega \in \pi^{-1}(\{x_w\})$ such that $[\omega]_n = w$. For each $m \in \mathbb{Z}_{\geq 0}$, we define $w(m) := [\omega]_{n+km} \in W_{n+km}$. Then, for $z_m \in K_{w(m)}$ ($m = 0, \dots, n$),

$$(5.13) \quad |f(x_w) - f(z_0)|^p \\ \leq 2^{p-1} |f(x_w) - f(z_n)|^p + 2^{p-1} \sum_{i=0}^{n-1} 2^{i(p-1)} |f(z_i) - f(z_{i+1})|^p.$$

Integrating (5.13), we obtain

$$(5.14) \quad \int_{K_w} |f(z) - M_n f(w)|^p d\mu(z) \\ \leq 2^{p-1} \int_{K_{w(n)}} |f(x_w) - f(z_n)|^p d\mu(z_n) \\ + 2^{2(p-1)} \sum_{i=0}^{n-1} 2^{i(p-1)} \int_{K_{w(i)}} \int_{K_{w(i+1)}} |f(z_i) - f(z_{i+1})|^p d\mu(z_{i+1}) d\mu(z_i).$$

Set $c := 3 \text{diam}(K, d) = 3$ and define

$$(5.15) \quad S_{p,\beta}^{(n)}(f) := \sup_{r \in (0, ca^{-n}]} r^{-\beta} \int_K \int_{B_d(z, r)} |f(z) - f(z')|^p d\mu(z') d\mu(z).$$

By Lemma 5.13, the first term of the right-hand side of (5.14) has a bound:

$$(5.16) \quad \int_{K_{w(n)}} |f(x_w) - f(z_n)|^p d\mu(z_n) \\ \leq C_{5.13} S_{p,\beta}^{(n)}(f) \int_{K_{w(n)}} d(x_w, z_n)^{\beta-\alpha} d\mu(z_n) \\ \leq (C_{5.13} \text{diam}(K, d)^{\beta-\alpha}) a^{-(n+kn)(\beta-\alpha)} S_{p,\beta}^{(n)}(f).$$

For the second term of (5.14), we see that

$$(5.17) \quad \begin{aligned} & 2^{i(p-1)} \int_{K_{w(i)}} \int_{K_{w(i+1)}} |f(z_i) - f(z_{i+1})|^p d\mu(z_{i+1})d\mu(z_i) \\ & \leq c_1 2^{i(p-1)} a^{\alpha k} a^{2\alpha(n+ki)} I_i(f), \end{aligned}$$

where

$$I_i(f) := \int_{K_{w(i)}} \int_{B_d(z_i, ca^{-(n+ki)})} |f(z_i) - f(z_{i+1})|^p d\mu(z_{i+1})d\mu(z_i)$$

and $c_1 > 0$ depends only on c and C_{AR} .

Now, we consider $k \in \mathbb{N}$ large enough so that

$$(5.18) \quad k(\beta - \alpha) \geq \alpha \quad \text{and} \quad N_* a^{-(\beta-\alpha)k} \vee 2^{p-1} a^{-(\beta-\alpha)k} < 1.$$

Then, by summing (5.16) and (5.17) over $w \in W_n$, we have from (5.18) that

$$\begin{aligned} \sum_{w \in W_n} \int_{K_{w(n)}} |f(x_w) - f(z_n)|^p d\mu(z_n) & \leq C_{5.13} S_{p,\beta}^{(n)}(f) N_*^n a^{-(n+kn)(\beta-\alpha)} \\ & \leq C_{5.13} S_{p,\beta}^{(n)}(f) a^{-(\beta-\alpha)n}, \end{aligned}$$

and from Proposition 2.11-(3) and (5.18) that

$$\begin{aligned} & \sum_{i=0}^{n-1} \sum_{w \in W_n} 2^{i(p-1)} \int_{K_{w(i)}} \int_{K_{w(i+1)}} |f(z_i) - f(z_{i+1})|^p d\mu(z_{i+1})d\mu(z_i) \\ & \leq c_1 a^{\alpha k} \sum_{i=0}^{n-1} 2^{i(p-1)} a^{2\alpha(n+ki)} \int_K \int_{B(x, ca^{-(n+ki)})} |f(x) - f(y)|^p d\mu(y)d\mu(x) \\ & \leq c_2 a^{\alpha k} S_{p,\beta}^{(n)}(f) \left(\sum_{i=0}^{n-1} 2^{i(p-1)} a^{-(\beta-\alpha)(n+ki)} \right) \\ & \leq c_2 a^{\alpha k} S_{p,\beta}^{(n)}(f) \left(\sum_{i=0}^{\infty} \left(2^{p-1} a^{-(\beta-\alpha)k} \right)^i \right) a^{-(\beta-\alpha)n} =: c_3 S_{p,\beta}^{(n)}(f) a^{-(\beta-\alpha)n}, \end{aligned}$$

where $c_2, c_3 > 0$ depend only on $C_{AR}, c, \beta, a, N_*, p$. From these estimates and (5.14), we finish the proof. \square

Now we are ready to finish the proof of Theorem 2.22.

Theorem 5.15.

$$\mathcal{F}_p = \Lambda_{p,\infty}^{\beta_p/p} = \left\{ f \in L^p(K, \mu) \left| \overline{\lim}_{r \downarrow 0} \int_K \int_{B_d(x,r)} \frac{|f(x) - f(y)|^p}{r^{\beta_p}} d\mu(y)d\mu(x) < \infty \right. \right\}.$$

Proof. Let $c > 0$ such that $\max_{(v,w) \in E_n} \sup_{x \in K_v, y \in K_w} d(x,y) < ca^{-n}$. We can choose such c depending only on C_{AD} by Lemma 2.13 and we can assume that $c \geq 3 \text{diam}(K, d)$ without loss of generality. Then $y \in B(x, ca^{-n})$ whenever $(v, w) \in E_n$ and $x \in K_v, y \in K_w$. Let $\beta > \alpha$ and set

$$A_{p,\beta}^{(n)}(f) := a^{\beta n} \int_K \int_{B_d(x, ca^{-n})} |f(x) - f(y)|^p d\mu(y)d\mu(x),$$

and define $S_{p,\beta}^{(n)}(f)$ as in (5.15) for each $n \in \mathbb{N}$ and $f \in L^p(K, \mu)$. Then, thanks to Lemma 5.12, it will suffice to show the following two estimates:

$$(5.19) \quad A_{p,\beta_p}^{(n)}(f) \leq c_{5.19} \left(\tilde{\mathcal{E}}_p^{G_n}(M_n f) + a^{\beta_p n} \sum_{w \in W_n} \int_{K_w} |f(x) - M_n f(w)|^p d\mu(x) \right), \quad f \in L^p(K, \mu),$$

$$(5.20) \quad a^{(\beta-\alpha)n} \cdot \mathcal{E}_p^{G_n}(M_n f) \leq c_{5.20} S_{p,\beta}^{(n)}(f), \quad f \in \mathcal{C}(K),$$

for some positive constants $c_{5.19}, c_{5.20}$ (without depending on f and n). Indeed, by (5.9), (5.19) and Lemma 5.12, we immediately see that

$$(5.21) \quad C_{\text{LB}}(c)^{-1} |f|_{\Lambda_{p,\infty}^{\beta_p/p}}^p \leq \sup_{n \in \mathbb{N}} A_{p,\beta_p}^{(n)}(f) \leq c_{5.19}(1 + C_{\text{PLKS}}) |f|_{\mathcal{F}_p}^p.$$

Additionally, by the weak monotonicity (Corollary 4.16), (5.20) and (5.10), we have

$$(5.22) \quad |f|_{\mathcal{F}_p}^p \leq c_{5.20} C_{\text{WM}} \overline{\lim}_{r \downarrow 0} r^{-\beta_p} \int_K \int_{B_d(x,r)} |f(x) - f(y)|^p d\mu(y) d\mu(x) \leq c'_{5.20} \overline{\lim}_{n \rightarrow \infty} a^{\beta_p n} \int_K \int_{B_d(x,a^{-n})} |f(x) - f(y)|^p d\mu(y) d\mu(x),$$

where $c'_{5.20} = c_{5.20} C_{\text{WM}} C_{\text{LB}}(c) C_{\text{LB}}(1)$. Our assertion follows from (5.21) and (5.22).

The rest of the proof is devoted to proving (5.19) and (5.20). First, we will prove (5.19). Let $x, y \in K$ with $d(x, y) < ca^{-n}$. Then, by the metric doubling property of (K, d) (see [31, pp. 81] for example), there exists a constant $L \geq 2$ depending only on C_{AR} such that, for any $w \in W_n$ with $x \in K_w$, we can choose $v \in W_n$ satisfying $d_{G_n}(v, w) \leq L$ and $y \in K_v$. From this observation, we have that

$$(5.23) \quad A_{p,\beta_p}^{(n)}(f) \leq a^{\beta_p n} \sum_{\substack{w \in W_n, v \in W_n; \\ d_{G_n}(v, w) \leq L}} \int_{K_w} \frac{1}{\mu(B_d(x, ca^{-n}))} \int_{K_v} |f(x) - f(y)|^p d\mu(y) d\mu(x).$$

To estimate the integral in (5.23), let $v, w \in W_n$ with $d_{G_n}(v, w) \leq L$. Then we can pick a path $[w(0), w(1), \dots, w(L)]$ in G_n from w to v , that is, $w(i)$ ($i = 0, \dots, L$) satisfy $w(0) = w$, $w(L) = v$ and

$$w(i-1) = w(i) \text{ or } (w(i-1), w(i)) \in E_n \text{ for each } i = 1, \dots, L.$$

Let $x_i \in K_{w(i)}$ for each $i = 0, \dots, L$. Then Hölder's inequality implies that

$$|f(x_0) - f(x_L)|^p \leq L^{p-1} \sum_{i=1}^L |f(x_{i-1}) - f(x_i)|^p.$$

Now, by integrating this, we deduce that

$$\begin{aligned} & \left(\prod_{i=1}^{L-1} \mu(K_{w(i)}) \right) \int_{K_w} \int_{K_v} |f(x) - f(y)|^p d\mu(x) d\mu(y) \\ & \leq L^{p-1} \sum_{i=1}^L \frac{\prod_{j=0}^L \mu(K_{w(j)})}{\mu(K_{w(i-1)}) \mu(K_{w(i)})} \int_{K_{w(i-1)}} \int_{K_{w(i)}} |f(x_i) - f(x_{i-1})|^p d\mu(x_i) d\mu(x_{i-1}). \end{aligned}$$

Since μ is the self-similar measure with weights $(N_*^{-1}, \dots, N_*^{-1})$, it is a simple computation that

$$\frac{\prod_{j=0}^L \mu(K_{w(j)})}{\mu(K_{w(i-1)})\mu(K_{w(i)})} \frac{1}{\prod_{i=1}^{L-1} \mu(K_{w(i)})} = \frac{\mu(K_v)\mu(K_w)}{\mu(K_{w(i-1)})\mu(K_{w(i)})} = 1.$$

Furthermore, the Ahlfors regularity of μ (more precisely, the volume doubling property of μ) implies that there exists a constant $c_1 > 0$ depending only on C_{AR}, a, N_* , c such that $\mu(K_z) \leq c_1 \mu(B_d(x, ca^{-n}))$ for any $n \in \mathbb{N}$, $z \in W_n$ and $x \in K$. Thus, it follows from (5.23) that

$$\begin{aligned} & A_{p, \beta_p}^{(n)}(f) \\ & \leq (c_1 L^{p-1} L_*^L) a^{\beta_p n} \sum_{(v, w) \in E_n} \int_{K_w} \int_{K_v} |f(x) - f(y)|^p d\mu(y) d\mu(x) \\ & \leq c_{5.19} \left(a^{\beta_p n} \sum_{v \in W_n} \int_{K_v} |f(x) - M_n f(v)|^p d\mu(x) \right. \\ & \quad \left. + \rho_p^n \sum_{(v, w) \in E_n} |M_n f(v) - M_n f(w)|^p \right), \end{aligned}$$

where $c_{5.19} := c_1 (2L)^{p-1} L_*^{L+1}$. This proves (5.19).

Next let us prove (5.20). Let $\beta > \alpha$, let $p > 1$ and let $f \in \mathcal{C}(K)$. For $n \in \mathbb{N}$, $(v, w) \in E_n$, $x \in K_v$ and $y \in K_w$, we see that

$$\begin{aligned} & |M_n f(v) - M_n f(w)|^p \\ & \leq 3^{p-1} (|M_n f(v) - f(x)|^p + |f(x) - f(y)|^p + |M_n f(w) - f(y)|^p). \end{aligned}$$

Integrating this over K_v and K_w , we obtain

$$\begin{aligned} & a^{(\beta-\alpha)n} \cdot |M_n f(v) - M_n f(w)|^p \\ & \leq 3^{p-1} \left(a^{\beta n} \int_{K_v} |M_n f(v) - f(x)|^p d\mu(x) \right. \\ & \quad \left. + a^{\beta n} \int_{K_w} \int_{K_v} |f(x) - f(y)|^p d\mu(x) d\mu(y) + a^{\beta n} \int_{K_w} |M_n f(w) - f(y)|^p d\mu(y) \right). \end{aligned}$$

Summing over $(v, w) \in E_n$, we obtain

$$\begin{aligned} & a^{(\beta-\alpha)n} \cdot \mathcal{E}_p^{G_n}(M_n f) \\ & \leq 2 \cdot 3^{p-1} L_* \left(a^{\beta n} \sum_{v \in W_n} \int_{K_v} |M_n f(v) - f(x)|^p d\mu(x) \right. \\ & \quad \left. + a^{\beta n} \sum_{(v, w) \in E_n} \int_{K_w} \int_{K_v} |f(x) - f(y)|^p d\mu(x) d\mu(y) \right). \end{aligned}$$

A bound of the first term in the right-hand side is obtained in Lemma 5.14. Noting that $K_v \subseteq B_d(y, ca^{-n})$ for $(v, w) \in E_n$ and $y \in K_w$, we can estimate the second

term as follows:

$$\begin{aligned} & a^{\beta n} \sum_{(v,w) \in E_n} \int_{K_w} \int_{K_v} |f(x) - f(y)|^p d\mu(x) d\mu(y) \\ & \leq c_2 a^{\beta n} \sum_{(v,w) \in E_n} \int_{K_w} \int_{B_d(y, ca^{-n})} |f(x) - f(y)|^p d\mu(x) d\mu(y) \leq c_2 c^{-\beta} L_* S_{p,\beta}^{(n)}(f), \end{aligned}$$

where $c_2 > 0$ depends only on C_{AR}, c . This proves (5.20) and finishes the proof. \square

As an immediate consequence of Theorem 5.15, we have a characterization of β_p as *critical Besov exponents*. For details on critical Besov exponents, see [3, 28] for example. This result is well-known when $p = 2$ (see [28, Theorem 4.6]).

Corollary 5.16. *It holds that $\beta_p = p \cdot \sup\{s > 0 \mid \Lambda_{p,\infty}^s \neq \{\text{constant}\}\}$.*

Proof. Note that $\Lambda_{p,\infty}^{\beta'/p} \subseteq \Lambda_{p,\infty}^{\beta/p}$ for any $\beta \leq \beta'$. It is immediate that

$$\beta_p \leq p \cdot \sup\{s > 0 \mid \Lambda_{p,\infty}^s \neq \{\text{constant}\}\}.$$

To prove the converse, let $\beta > \beta_p$. If $f \in \mathcal{C}(K)$ is not constant, then there exists $N \in \mathbb{N}$ such that $\tilde{\mathcal{E}}_p^{G_N}(M_N f) > 0$. By Corollary 4.16, for any $n \geq N$,

$$a^{(\beta-\alpha)n} \cdot \mathcal{E}_p^{G_n}(M_n f) = a^{(\beta-\alpha)n} \rho_p^{-n} \cdot \tilde{\mathcal{E}}_p^{G_n}(M_n f) \geq C_{WM}^{-1} a^{(\beta-\alpha)n} \rho_p^{-n} \cdot \tilde{\mathcal{E}}_p^{G_N}(M_N f).$$

Letting $n \rightarrow \infty$ in this inequality, we obtain $\overline{\lim}_{n \rightarrow \infty} a^{(\beta-\alpha)n} \cdot \mathcal{E}_p^{G_n}(M_n f) = \infty$ since $\rho_p^{-1} a^{\beta-\alpha} > 1$. By (5.20), we conclude that $|f|_{\Lambda_{p,\infty}^{\beta/p}} = \infty$ whenever $\beta > \beta_p$ and $f \in \mathcal{C}(K)$ is not constant. This proves our assertion. \square

6. CONSTRUCTION OF A CANONICAL SCALING LIMIT OF p -ENERGIES

To construct a canonical Dirichlet form on fractals, there is already an established way as appeared in [54, proof of Theorem 6.9]. However, in the original argument of [54], the Markov property of their ‘‘Dirichlet form’’ was not clarified. In [45], Kigami has pointed out this gap and filled it.

After Kigami’s work, another very simple way to check the Markov property is given by Barlow, Bass, Kumagai, and Teplyaev [8, proof of Theorem 2.1]. This method deduces that the Dirichlet forms of Kusuoka and Zhou in [54] have the Markov property, but it very heavily relies on being Dirichlet forms, that is, the use of bilinearity (and locality) is essential to follow [8, proof of Theorem 2.1].

Regrettably, these ways are insufficient to follow the arguments in Section 7, where the main results about \mathcal{E}_p -energy measures will be proved. Indeed, our strategy to prove the chain rule (Theorem 2.25) will require some expression of constructed p -energy \mathcal{E}_p (see (6.22)) due to the lack of representation formula (see [25, (3.2.10)] for example) that is very useful in Dirichlet form theory. Also, as shown in [19, 29, 44], the usage of Γ -convergence is very useful to construct energies. However, we need to adopt alternative approach because our argument in Theorem 2.25 will heavily use the compactness of $f(K)$ for a fixed function $f \in \mathcal{F}_p$. If we construct \mathcal{E}_p by using Γ -convergence, then we lose this compactness since we have to consider $\bigcup_{n \geq 1} f_n(K)$ instead of $f(K)$, where $\{f_n\}_{n \geq 1}$ is a sequence converging to f in L^p .

In order to overcome these difficulties, we will introduce a new series of graphs \mathbb{G}_n approximating the underlying Sierpiński carpet in subsection 6.1. Then in subsection 6.2 we directly construct p -energy \mathcal{E}_p as a subsequential scaling limit of




FIGURE 5. Modified Sierpiński carpet graph $\{\mathbb{G}_n\}_{n \geq 1}$ (This figure draws \mathbb{G}_3 in the SC case)

discrete p -energies on this new series of graphs. Since we already prove (B_p) and (KM_p) , Kigami's result [44, Theorem 9.3] gives p -energies on GSCs satisfying all properties in Theorem 2.23 (except Clarkson's inequality). We emphasize that a main aim of this section is to construct p -energies that will be useful to prove the chain rule in Section 7 (Theorem 2.25-(2)).

Throughout this section, let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet and suppose that Assumption 4.7 holds.

6.1. Behavior of p -energies on modified Sierpiński carpet graphs. For $n \in \mathbb{Z}_{\geq 0}$, define subsets $\mathbb{V}_n \subseteq K$ by

$$\mathbb{V}_n = \left\{ F_w \left(\sum_{k=1}^D \sigma_k \mathbf{e}_k \right) \mid w \in W_n, \sigma_k \in \{-1, +1\} \text{ for } k = 1, \dots, D \right\}.$$

(The conditions (GSC1) and (GSC4) ensure that $\sum_{k=1}^D \sigma_k \mathbf{e}_k \in K$.) Note that $\#\mathbb{V}_0 = 2^D$. Next, we inductively define edge sets \mathbb{E}_n by

$$\mathbb{E}_0 := \{(x, y) \mid x \neq y, x, y \in \mathbb{V}_0\},$$

and

$$\mathbb{E}_n := \{(F_i(x), F_i(y)) \mid i \in S, (x, y) \in \mathbb{E}_{n-1}\}.$$

Define a new finite graph by $\mathbb{G}_n := (\mathbb{V}_n, \mathbb{E}_n)$ (see Figure 5). Note that \mathbb{G}_0 coincides with the complete graph \mathbb{K}_{2^D} having 2^D vertices. We will write $d_{\mathbb{G}_n}$ for the graph distance of \mathbb{G}_n . By (GSC2), we see that \mathbb{G}_n is connected. Furthermore, we easily see that $\{\mathbb{G}_n\}_{n \geq 0}$ is an increasing sequence. It is also immediate that

$$L_{*, \text{modif}} := \sup_{n \in \mathbb{Z}_{\geq 0}} \max_{x \in \mathbb{V}_n} \#\{y \in \mathbb{V}_n \mid (x, y) \in \mathbb{E}_n\} \leq L_* 2^D.$$

For any $n, m \in \mathbb{Z}_{\geq 0}$ and $w \in W_n$, we define a subset \mathbb{V}_m^w of \mathbb{V}_{n+m} by setting $\mathbb{V}_m^w := \{F_w(x) \mid x \in \mathbb{V}_m\}$, and define a subgraph $\mathbb{G}_m^w := (\mathbb{V}_m^w, \mathbb{E}_m^w)$, where

$$\mathbb{E}_m^w = \{(x, y) \in \mathbb{E}_{n+m} \mid x, y \in \mathbb{V}_m^w\}.$$

Note that, for $v, w \in W_*$ with $|v| = |w|$, $\mathbb{V}_m^w \cap \mathbb{V}_m^v \neq \emptyset$ if and only if $K_v \cap K_w \neq \emptyset$.

For simplicity, we write $\mathcal{R}_p^{\mathbb{G}_n}(x, y)$ to denote $\mathcal{C}_p^{\mathbb{G}_n}(\{x\}, \{y\})^{-1}$, that is,

$$\mathcal{R}_p^{\mathbb{G}_n}(x, y) := \sup \left\{ \frac{|f(x) - f(y)|^p}{\mathcal{E}_p^{\mathbb{G}_n}(f)} \mid f: \mathbb{V}_n \rightarrow \mathbb{R} \text{ is not constant} \right\},$$

for each $x, y \in \mathbb{V}_n$. Then one of the key ingredients is the next proposition; $\mathcal{R}_p^{\mathbb{G}_n}(x, y)$ behaves like $\mathcal{R}_p^{(n)}$.

Proposition 6.1. *There exists a positive constant $C_{6.1}$ (depending only on $p, a, D, N_*, L_*, \rho_p$) such that, for any $n \in \mathbb{N}$ and $x, y \in \mathbb{V}_0$,*

$$(6.1) \quad \mathcal{R}_p^{\mathbb{G}_n}(x, y) \leq C_{6.1} \rho_p^n.$$

Remark 6.2. For $p = 2$ in the SC case, such point-to-point estimates on a series of Sierpiński carpet graphs are proved in [56, Appendix], where a uniform Hanarck inequality by Barlow and Bass (see also Remark 4.17) is used. In [44, Lemma 8.5], Kigami also shows similar estimates for all $p > \dim_{\text{ARC}}$ assuming p -conductive homogeneity, where he also uses some uniform Hölder estimate. Our proof also relies on the uniform Hölder estimate: Theorem 4.15.

To prove this proposition, we need an estimate of p -conductance between points on the original graph $\{G_n\}_{n \geq 1}$. For $n \in \mathbb{Z}_{\geq 0}$ and x , we fix $w^x(n) \in W_n$ such that $x \in K_{w^x(n)}$. If $x = \sum_{k=1}^D \sigma_k \mathbf{e}_k \in \mathbb{V}_0$, then such $w^x(n)$ is uniquely given by

$$w^x(n) = (\sigma_1, \dots, \sigma_D)^n \in W_n.$$

We can show the following lemma in a similar way to the ‘chain argument’ in the proof of Theorem 4.14.

Lemma 6.3. *There exists a constant $C_{6.3} \geq 1$ (depending only on $p, a, D, N_*, L_*, \rho_p$) such that, for any $n \in \mathbb{N}$ and $x, y \in \mathbb{V}_0$ with $x \neq y$,*

$$(6.2) \quad C_{6.3}^{-1} \rho_p^{-n} \leq \mathcal{C}_p^{\mathbb{G}_n}(\{w^x(n)\}, \{w^y(n)\}) \leq C_{6.3} \rho_p^{-n}.$$

Proof. An upper bound is easy. Indeed, by Proposition 2.16, Theorems 3.4 and 3.13,

$$(6.3) \quad \mathcal{C}_p^{\mathbb{G}_n}(\{w^x(n)\}, \{w^y(n)\}) \leq \mathcal{C}_p^{(n-1)} \leq c_* \rho_p \cdot \rho_p^{-n},$$

where c_* is the constant in (3.21).

In order to prove the converse, we first consider the case $|x - y|_{\mathbb{R}^D} = 2$. Let $C_{\text{UH}} > 0$ be the constant (depending only on $p, a, D, L_*, N_*, \rho_p$) in Theorem 4.15 and choose $l_* \in \mathbb{N}$ such that

$$(6.4) \quad C_{\text{UH}} a^{-(\beta_p - \alpha)l_*} \leq \frac{1}{4}.$$

We also set $v(x) := [w^x(n)]_{l_*} \in W_{l_*}$ for each $x \in \mathbb{V}_0$. Since $|x - y|_{\mathbb{R}^D} = 2$, there exists a horizontal chain $[z(1), \dots, z(L)]$ in \tilde{G}_{l_*} such that $z(1) = v(x)$, $z(L) = v(y)$, $z(k) \in \partial_* G_{l_*}$ for any $k = 1, \dots, L$ and $(z(k), z(k+1)) \in \tilde{E}_{l_*}$ for each $k = 1, \dots, L-1$, where $L = a^{l_*}$.

Let $f_n: W_n \rightarrow \mathbb{R}$ satisfy $f_n(w^x(n)) = 0$, $f_n(w^y(n)) = 1$ and

$$\mathcal{E}_p^{\mathbb{G}_n}(f_n) = \mathcal{C}_p^{\mathbb{G}_n}(\{w^x(n)\}, \{w^y(n)\}).$$

Note that f_n is $[0, 1]$ -valued. From Theorem 4.15 and (6.4),

$$\max_{v(x) \cdot W_{n-l_*}} f_n \leq \frac{1}{4} \quad \text{and} \quad \min_{v(y) \cdot W_{n-l_*}} f_n \geq \frac{3}{4}.$$

Now we define g_n by setting $g_n := 2((f_n \vee 1/4) \wedge 3/4)$. Then we have that

$$\mathcal{C}_p^{(n-l_*, L)} \leq \mathcal{E}_p^{\mathbb{G}_n}(g_n) \leq 2^p \mathcal{E}_p^{\mathbb{G}_n}(f_n).$$

(Recall the definition of $\mathcal{C}_p^{(n, L)}$ in subsection 4.3.) By Lemma 4.12 and Theorem 3.4, there exists a constant $C(L) > 0$ depending only on p, a, L_*, N_* such that

$$\mathcal{C}_p^{(n-l_*, L)} \geq C(L)^{-1} \mathcal{C}_p^{(n-l_*)} \geq (C(L) C_{3.4} \mathcal{C}_p^{(l_*)})^{-1} \mathcal{C}_p^{(n)}.$$

Hence, by using (3.21),

$$\mathcal{C}_p^{G_n}(\{w^x(n)\}, \{w^y(n)\}) \geq (2^p C(L) C_{3.4} \mathcal{C}_p^{(l_*)})^{-1} \mathcal{C}_p^{(n)} \geq c_* (2^p C(L) C_{3.4} \mathcal{C}_p^{(l_*)})^{-1} \rho_p^{-n},$$

which proves the case $|x - y|_{\mathbb{R}^D} = 2$.

Finally, note that $(v, w) \mapsto \mathcal{C}_p^G(\{v\}, \{w\})^{-1/p}$ is a metric on a graph G . Indeed, this fact immediately follows from the representation:

$$\mathcal{C}_p^G(\{v\}, \{w\})^{-1/p} = \max \left\{ \frac{|f(v) - f(w)|}{\mathcal{E}_p^G(f)^{1/p}} \mid f: V \rightarrow \mathbb{R} \text{ with } \mathcal{E}_p^G(f) > 0 \right\}.$$

Applying the triangle inequality of this metric, we obtain

$$\max_{x \neq y \in \mathbb{V}_0} \mathcal{C}_p^{G_n}(\{w^x(n)\}, \{w^y(n)\})^{-1/p} \leq DC_1 \rho_p^{n/p},$$

where $C_1 = (2^p c_*^{-1} C(L) C_{3.4} \mathcal{C}_p^{(l_*)})^{1/p}$. \square

Remark 6.4. It is known that $(x, y) \mapsto \mathcal{C}_p^G(\{x\}, \{y\})^{-1/(p-1)}$ also becomes a metric, and thus $\mathcal{C}_p^G(\cdot, \cdot)^{-1/(p-1)}$ gives a generalization of the *resistance metric*. This fact is proved in [1, Theorem 8] when G is a finite graph. One can check the case of infinite graphs in [57, Theorem 4.3].

Now we are ready to prove Proposition 6.1 by using Lemma 4.3.

Proof of Proposition 6.1. For each $n \in \mathbb{N}$, let $\varphi_n: W_n \rightarrow \mathbb{V}_n$ be a function such that $\varphi_n(w) \in \mathbb{V}_0^w$ for $w \in W_n$ and

$$\varphi_n((\sigma_1, \dots, \sigma_D)^n) = \sum_{k=1}^D \sigma_k \mathbf{e}_k \quad \text{if } (\sigma_k)_{k=1}^D \in \{-1, +1\}^D.$$

Then we easily see that φ_n is a rough isometry from G_n to \mathbb{G}_n with $C_1 = 1, C_2 = 2, C_3 = 2, C_4 = ((2^D - 1)/L_*) \vee (L_{*, \text{modif}}/2)$ (recall Definition 4.1). Hence $\{\varphi_n\}_{n \geq 1}$ is a uniform rough isometry from $\{G_n\}_{n \geq 1}$ to $\{\mathbb{G}_n\}_{n \geq 1}$. Note that for $(\sigma_k)_{k=1}^D \in \{-1, +1\}^D$,

$$\varphi_n(w) = \sum_{k=1}^D \sigma_k \mathbf{e}_k \quad \text{if and only if } w = (\sigma_1, \dots, \sigma_D)^n,$$

i.e. $\varphi_n^{-1}(\{\sum_{k=1}^D \sigma_k \mathbf{e}_k\}) = \{(\sigma_1, \dots, \sigma_D)^n\}$. Thus, for $x \neq y \in \mathbb{V}_0$, Lemma 4.3 yields that

$$\mathcal{C}_p^{G_n}(\{w^x(n)\}, \{w^y(n)\}) \leq C_{\text{URI}} \mathcal{R}_p^{\mathbb{G}_n}(x, y)^{-1},$$

which together with Lemma 6.3 implies (6.1). \square

Next, we define rescaled p -energy $\tilde{\mathcal{E}}_p^{\mathbb{G}_n}: \mathcal{C}(K) \rightarrow \mathbb{R}$ on \mathbb{G}_n by setting

$$\tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f) := \frac{\rho_p^n}{2} \sum_{(x, y) \in \mathbb{E}_n} |f(x) - f(y)|^p,$$

for each $f \in \mathcal{C}(K)$. Then the following lemma, especially statement (4) below, is a collection of benefits of the new graphical approximation $\{\mathbb{G}_n\}_{n \geq 0}$.

Lemma 6.5. *Let $f \in \mathcal{C}(K)$ and let $n \in \mathbb{Z}_{\geq 0}$. The following statements hold.*

- (1) *It holds that $\sup_{n \geq 0} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f) = 0$ if and only if f is constant.*
- (2) *If $T \in \mathcal{G}_0$, then $\tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f \circ T) = \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f)$.*

(3) For $m \in \mathbb{Z}_{\geq 0}$,

$$(6.5) \quad L_*^{-1} \rho_p^m \sum_{w \in W_m} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f \circ F_w) \leq \tilde{\mathcal{E}}_p^{\mathbb{G}_{n+m}}(f) \leq \rho_p^m \sum_{w \in W_m} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f \circ F_w).$$

(4) If $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ with $\text{Lip}(\varphi) \leq 1$, then $\tilde{\mathcal{E}}_p^{\mathbb{G}_n}(\varphi \circ f) \leq \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f)$.

Proof. (1) It is immediate that $\sup_{n \in \mathbb{N}} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f) = 0$ if $f \in \mathcal{C}(K)$ is constant. We easily see that $\overline{\bigcup_{n \in \mathbb{N}} \mathbb{V}_n^K} = K$. Thus, if $f \in \mathcal{C}(K)$ satisfies $\mathcal{E}_p^{\mathbb{G}_n}(f) = 0$ for any $n \in \mathbb{Z}_{\geq 0}$, then f is constant.

(2) This immediately follows from the symmetries of \mathbb{G}_n .

(3) Let $f \in \mathcal{C}(K)$ and $n \in \mathbb{Z}_{\geq 0}$. Since $\bigcup_{w \in W_m} \mathbb{V}_n^w = \mathbb{V}_{n+m}$, we have

$$\begin{aligned} \tilde{\mathcal{E}}_p^{\mathbb{G}_{n+m}}(f) &\leq \rho_p^m \sum_{w \in W_m} \left(\frac{\rho_p^n}{2} \sum_{\substack{(x,y) \in \mathbb{E}_{n+m}; \\ x,y \in K_w}} |f(F_w(F_w^{-1}(x))) - f(F_w(F_w^{-1}(y)))|^p \right) \\ &= \rho_p^m \sum_{w \in W_m} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f \circ F_w). \end{aligned}$$

Next, by noting that

$$\sup_{(x,y) \in \mathbb{E}_{n+m}} \#\{w \in W_m \mid x, y \in \mathbb{V}_n^w\} \leq L_*,$$

we see that

$$\begin{aligned} \rho_p^m \sum_{w \in W_m} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f \circ F_w) &= \frac{\rho_p^{n+m}}{2} \sum_{w \in W_m} \sum_{\substack{(x,y) \in \mathbb{E}_{n+m}; \\ x,y \in \mathbb{V}_n^w}} |f(x) - f(y)|^p \\ &\leq L_* \frac{\rho_p^{n+m}}{2} \sum_{(x,y) \in \mathbb{E}_{n+m}} |f(x) - f(y)|^p = L_* \tilde{\mathcal{E}}_p^{\mathbb{G}_{n+m}}(f), \end{aligned}$$

which proves (6.5).

(4) The required estimate immediately follows from the fact that $|\varphi(a) - \varphi(b)| \leq |a - b|$ for any $a, b \in \mathbb{R}$ whenever $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ satisfies $\text{Lip}(\varphi) \leq 1$. \square

We also have the weak monotonicity of $\tilde{\mathcal{E}}_p^{\mathbb{G}_n}$ as follows. Its proof is similar to [29, Theorem 7.1], where 2-energies are considered.

Lemma 6.6. *There exists a constant $C_{\text{WM,modif}} > 0$ (depending only on $p, a, D, N_*, L_*, \rho_p$) such that*

$$(6.6) \quad \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f) \leq C_{\text{WM,modif}} \tilde{\mathcal{E}}_p^{\mathbb{G}_{n+m}}(f),$$

for every $n, m \in \mathbb{Z}_{\geq 0}$ and $f \in \mathcal{C}(K)$. In particular, for any $f \in \mathcal{C}(K)$,

$$\sup_{n \in \mathbb{N}} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f) \leq C_{\text{WM,modif}} \liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f).$$

Proof. Let $(x, y) \in \mathbb{E}_n$. Then there exists $w \in W_{n-1}$ such that $x, y \in K_w$. Furthermore, there exist $x_0, y_0 \in \mathbb{V}_0$ such that $x = F_w(x_0)$ and $y = F_w(y_0)$. Now we define

$$\mathcal{R}_p^{\mathbb{G}_m^w}(z_1, z_2) := \sup \left\{ \frac{|f(z_1) - f(z_2)|^p}{\mathcal{E}_p^{\mathbb{G}_m^w}(f)} \mid f \in \mathcal{C}(K) \text{ with } f|_{\mathbb{V}_m^w} \text{ is not constant} \right\}.$$

From Proposition 6.1 and the cutting law ([57, Proposition 3.18]),

$$\begin{aligned} |f(x) - f(y)|^p &\leq \mathcal{R}_p^{\mathbb{G}^m}(F_w(x_0), F_w(y_0)) \mathcal{E}_p^{\mathbb{G}^m}(f) \\ &\leq \mathcal{R}_p^{\mathbb{G}^m}(x_0, y_0) \mathcal{E}_p^{\mathbb{G}^m}(f) \\ &\leq C_{6.1} \rho_p^m \mathcal{E}_p^{\mathbb{G}^{n+m}}(f). \end{aligned}$$

Summing over $(x, y) \in \mathbb{E}_n$, we obtain $\mathcal{E}_p^{\mathbb{G}^n}(f) \leq 2^{-1} C_{6.1} \rho_p^m \mathcal{E}_p^{\mathbb{G}^{n+m}}(f)$, which deduces our assertion with $C_{\text{WM,modif}} = 2^{-1} C_{6.1}$. \square

The rest of this subsection is devoted to proving the following lemma. Recall the definition of $|\cdot|_{\mathcal{F}_p}$ that is a semi-norm of \mathcal{F}_p (see (5.1)).

Lemma 6.7. *There exists a constant $C_{6.7} \geq 1$ depending only on $p, a, D, N_*, L_*, C_{\text{AR}}, \rho_p$ such that*

$$(6.7) \quad C_{6.7}^{-1} |f|_{\mathcal{F}_p}^p \leq \sup_{n \in \mathbb{N}} \tilde{\mathcal{E}}_p^{\mathbb{G}^n}(f) \leq C_{6.7} |f|_{\mathcal{F}_p}^p \quad \text{for any } f \in L^p(K, \mu).$$

Remark 6.8. Such discrete characterizations of Lipschitz–Besov space are treated in [12], but we need some modification as stated in [30, Remark 1 in Section 3]. To be self-contained, we give complete proofs in a similar way as in [29, Theorems 3.5 and 3.6 and Proposition 11.1], where they give discrete characterizations of \mathcal{F}_2 on the SC.

By virtue of Theorem 2.22, it will suffice to show that $\sup_{n \in \mathbb{N}} \tilde{\mathcal{E}}_p^{\mathbb{G}^n}(f)$ and $|f|_{\Lambda_{p,\infty}^{\beta_{p/p}}}$ are comparable. Similarly to [12], we will apply an argument using discrete approximations of measure μ with respect to the weak convergence of probability measures (see Lemma 6.10-(2)). The following lemma is elementary (see also [12, Lemma 3.12], [30, Remark 1 in Section 3]). Recall the definition of $U_1(x, s)$ in (2.4).

Lemma 6.9. *Let $\{\mu_n\}_{n \geq 1}$ be a sequence of probability measures on K given by*

$$\mu_n := \frac{1}{\#\mathbb{V}_n} \sum_{x \in \mathbb{V}_n} \delta_x,$$

where δ_x denotes the Dirac measure with support $\{x\}$ for each $x \in K$. Then there exist a subsequence $\{n_k\}_{k \geq 1}$ and a constant $c_{6.9} > 0$ (depending only on $a, D, N_*, L_*, C_{\text{AR}}$) such that for any $m \in \mathbb{Z}_{\geq 0}$, $p > 1$ and $f \in \mathcal{C}(K)$,

$$(6.8) \quad \begin{aligned} &\lim_{k \rightarrow \infty} \int_K \int_{U_1(x, a^{-m})} |f(x) - f(y)|^p d\mu_{n_k}(y) d\mu_{n_k}(x) \\ &\geq c_{6.9} \int_K \int_{U_1(x, a^{-m})} |f(x) - f(y)|^p d\mu(y) d\mu(x). \end{aligned}$$

Proof. Since K is compact, by Prokhorov’s theorem (see [10, Theorem 5.1] for example), there exist a subsequence $\{n_k\}_{k \geq 1}$ and a Borel probability measure $\tilde{\mu}$ on K such that μ_{n_k} converges weakly to $\tilde{\mu}$ as $k \rightarrow \infty$. From the definition of \mathbb{G}_n , by noting that

$$\sup_{n \geq 0, x \in \mathbb{V}_n} \#\{w \in W_n \mid x \in \mathbb{V}_0^w\} \leq L_*,$$

we have

$$(6.9) \quad L_*^{-1} 2^D N_*^n \leq \#\mathbb{V}_n \leq 2^D N_*^n, \quad n \in \mathbb{Z}_{\geq 0}.$$

Next we will see that $\tilde{\mu}$ is “ α -Ahlfors regular” and $\tilde{\mu}(\partial K_w) = 0$ for any $w \in W_*$. (Recall that $\alpha = \log N_*/\log a$ and thus $a^\alpha = N_*$.) For $r \in (0, 1]$, let $n(r) \in \mathbb{Z}_{\geq 0}$ denote the unique non-negative integer such that $a^{-n(r)} \leq r < a^{-n(r)+1}$. Let $C_{\text{AD}} \geq 1$ be the constant in Proposition 2.11-(2A) and let $n_* \in \mathbb{Z}_{\geq 0}$ such that $a^{-n_*} \leq C_{\text{AD}}^{-1} < a^{-n_*+1}$. Note that n_* depends only on C_{AD} and a . In addition, note that we can choose $r_0 \in (0, 1)$ (depending only on C_{AD} and a) so that $n(r) \geq n_*$ for all $r \in (0, r_0]$. Hereafter, we fix such a small $r_0 \in (0, 1)$. Let $x \in K$ and $r \in (0, r_0]$. By Proposition 2.11-(2A), we know that $U_1(x, C_{\text{AD}}^{-1}r) \subseteq B_d(x, r) \subseteq U_1(x, C_{\text{AD}}r)$. From $U_1(x, C_{\text{AD}}^{-1}r) \subseteq B_d(x, r)$, we can find $w \in W_{n(r)+n_*}$ such that $K_w \subseteq B_d(x, r)$. The portmanteau theorem (see [10, Theorem 2.1] for example) and (6.9) yield that

$$\tilde{\mu}(\overline{B_d(x, r)}) \geq \overline{\lim}_{k \rightarrow \infty} \mu_{n_k}(\overline{B_d(x, r)}) \geq \overline{\lim}_{k \rightarrow \infty} \mu_{n_k}(K_w) \geq L_*^{-1} a^{-(\alpha n_* + 1)} r^\alpha.$$

Similarly, by $B_d(x, r) \subseteq U_1(x, C_{\text{AD}}r)$, portmanteau theorem and (6.9),

$$\tilde{\mu}(B_d(x, r)) \leq \underline{\lim}_{k \rightarrow \infty} \mu_{n_k}(B_d(x, r)) \leq \underline{\lim}_{k \rightarrow \infty} \mu_{n_k}(U_1(x, C_{\text{AD}}r)) \leq L_*^2 a^{\alpha n_*} r^\alpha.$$

Consequently, for any $x \in K$ and $r \in (0, \text{diam}(K, d)]$, we obtain

$$(aN_*L_*C_{\text{AD}}^\alpha)^{-1}r^\alpha \leq \tilde{\mu}(\overline{B_d(x, r)}), \quad \tilde{\mu}(B_d(x, r)) \leq L_*^2N_*C_{\text{AD}}^\alpha r^\alpha.$$

By [31, Exercise 8.11] (or by following the argument in [27, Lemma 2.13]), there exists a constant $C_1 \geq 1$ depending only on $a, N_*, L_*, C_{\text{AD}}$ such that

$$C_1^{-1}\mathcal{H}^\alpha(A) \leq \tilde{\mu}(A) \leq C_1\mathcal{H}^\alpha(A) \quad \text{for any } A \in \mathcal{B}(K),$$

where \mathcal{H}^α is the normalized α -dimensional Hausdorff measure on (K, d) . Since μ is α -Ahlfors regular, the measure μ is also comparable to \mathcal{H}^α . Hence Proposition 2.11-(3) yields that $\tilde{\mu}(\partial K_w) = 0$ for all $w \in W_*$.

Finally, we will prove (6.8). Let $f \in \mathcal{C}(K)$ be not constant. Then there exists $N \geq 1$ such that $f|_{\mathbb{V}_N}$ is not constant. For a Borel measure ν on K and $n \geq N$, define

$$I(f, \nu) := \int_K \int_K |f(x) - f(y)|^p d\nu(y)d\nu(x).$$

Since $f|_{\mathbb{V}_N}$ is not constant and f is bounded, we have $I(f, \tilde{\mu}), I(f, \mu_n) \in (0, \infty)$. The weak convergence of $\mu_{n_k} \times \mu_{n_k}$ to $\tilde{\mu} \times \tilde{\mu}$ (see [10, Theorem 2.8] for example) implies that $I(f, \mu_{n_k}) \rightarrow I(f, \tilde{\mu})$ as $n \rightarrow \infty$. Next, we define $\nu_n(dx \otimes dy) := I(f, \mu_n)^{-1} |f(x) - f(y)|^p d\mu_n(x)d\mu_n(y)$. Since f is continuous, we easily see that ν_{n_k} converges weakly to the probability measure ν on $K \times K$ given by

$$\nu(dx \otimes dy) := I(f, \tilde{\mu})^{-1} |f(x) - f(y)|^p d\tilde{\mu}(x)d\tilde{\mu}(y).$$

Thanks to $\tilde{\mu}(\partial K_w) = 0$, it is immediate that

$$\nu(\partial(K \times U_1(x, a^{-m}))) \leq \nu(\partial K \times K) + \nu(K \times \partial U_1(x, a^{-m})) = 0.$$

(Note that $\partial U_1(x, a^{-m}) \subseteq \bigcup_{w \in W_m} \partial K_w$.) The portmanteau theorem and the fact that $\mu \asymp \tilde{\mu}$ deduce (6.8). \square

We next prove some (p, p) -Poincaré type inequalities (in the sense of Kumagai and Sturm), which plays the same role as Lemma 5.14.

Lemma 6.10. *Suppose that $\beta > \alpha$ and $p > 1$. Then there exists a positive constant $C_{6.10}$ (depending only on $p, a, D, N_*, \beta, C_{AR}$ and ρ_p) such that for every $n \in \mathbb{Z}_{\geq 0}$ and $f \in L^p(K, \mu)$,*

$$(6.10) \quad a^{\beta n} \sum_{w \in W_n} \sum_{x \in \mathbb{V}_0^w} \int_{K_w} |f(x) - f(z)|^p d\mu(z) \leq C_{6.10} |f|_{\Lambda_{p,\infty}^{\beta/p}}^p.$$

Proof. The proof is essentially the same as that of Lemma 5.14, so we omit the proof. (Consider $x \in \mathbb{V}_0^w$ instead of x_w in Lemma 5.14 and note that $\max_{w \in W_n} \#\mathbb{V}_0^w = 2^D$.) \square

Now, we are ready to prove Lemma 6.7.

Proof of Lemma 6.7. Let $\beta > \alpha$. We will prove the following two bounds:

$$\begin{aligned} \text{Upper bound: } \sup_{n \in \mathbb{N}} a^{(\beta-\alpha)n} \cdot \mathcal{E}_p^{\mathbb{G}_n}(f) &\leq C_{\text{upper}} |f|_{\Lambda_{p,\infty}^{\beta/p}}^p \quad \text{for every } f \in L^p(K, \mu), \\ \text{Lower bound: } \sup_{n \in \mathbb{N}} a^{(\beta_p-\alpha)n} \cdot \mathcal{E}_p^{\mathbb{G}_n}(f) &\geq C_{\text{lower}} |f|_{\Lambda_{p,\infty}^{\beta/p}}^p \quad \text{for every } f \in \mathcal{C}(K), \end{aligned}$$

for some positive constants $C_{\text{upper}}, C_{\text{lower}}$ (without depending on f). Note that, from Theorem 2.22, the case $\beta = \beta_p$ in these bounds includes our assertion.

Upper bound. For $f \in L^p(K, \mu)$, we easily see that

$$a^{(\beta-\alpha)n} \cdot \mathcal{E}_p^{\mathbb{G}_n}(f) \leq 2^{p-2} a^{\beta n} L_{*,\text{modif}} \sum_{w \in W_n} \sum_{x \in \mathbb{V}_0^w} \int_{K_w} |f(x) - f(z)|^p d\mu(z).$$

Applying Lemma 6.10, we get the desired bound with $C_{\text{upper}} = 2^{p-2} L_{*,\text{modif}} C_{6.10}$ that depends only on $p, a, D, N_*, L_*, \beta, C_{AR}, \rho_p$.

Lower bound. The desired bound can be obtained by applying [12, Theorem 3.18], but we give a complete proof for reader's convenience. Our proof is inspired by the arguments in those of [12, Lemma 4.2] and [29, Theorem 3.6].

Let $f \in \mathcal{C}(K)$ and let $n, m \in \mathbb{Z}_{\geq 0}$. Then we see that

$$(6.11) \quad \begin{aligned} &\int_K \int_{U_1(x, a^{-n})} |f(x) - f(y)|^p d\mu_{n+m}(y) d\mu_{n+m}(x) \\ &\leq L_* \sum_{w \in W_n} \int_{K_w} \int_{U_1(x, a^{-n})} |f(x) - f(y)|^p d\mu_{n+m}(y) d\mu_{n+m}(x) \\ &\leq L_* \sum_{\substack{v, w \in W_n; \\ d_{G_n}(v, w) \leq 2}} \int_{K_w} \int_{K_v} |f(x) - f(y)|^p d\mu_{n+m}(y) d\mu_{n+m}(x). \end{aligned}$$

Now, for $v, w \in W_n$, we define

$$\begin{aligned} I_m(f; v, w) &:= \int_{K_w} \int_{K_v} |f(x) - f(y)|^p d\mu_{n+m}(y) d\mu_{n+m}(x) \\ &= (\#\mathbb{V}_{n+m})^{-2} \sum_{x \in \mathbb{V}_n^w} \sum_{y \in \mathbb{V}_n^v} |f(x) - f(y)|^p. \end{aligned}$$

For $v, w \in W_n$ with $d_{G_n}(v, w) \leq 2$, we fix $\mathfrak{t}[v, w] \in W_n$ such that

$$\max_{z \in \{v, w\}} d_{G_n}(z, \mathfrak{t}[v, w]) \leq 1,$$

i.e. $K_v \cap K_{\mathfrak{t}[v,w]} \neq \emptyset$ and $K_w \cap K_{\mathfrak{t}[v,w]} \neq \emptyset$. We also fix $x_0 \in \mathbb{V}_0^w \cap \mathbb{V}_0^{\mathfrak{t}[v,w]}$ and $x_1 \in \mathbb{V}_0^v \cap \mathbb{V}_0^{\mathfrak{t}[v,w]}$. Then, by Hölder's inequality,

$$\begin{aligned} I_m(f; v, w) &\leq 3^{p-1} (\#\mathbb{V}_{n+m})^{-2} \sum_{x \in \mathbb{V}_m^w} \sum_{y \in \mathbb{V}_m^v} \{ |f(x) - f(x_0)|^p + |f(x_0) - f(x_1)|^p \\ &\quad + |f(y) - f(x_1)|^p \}. \end{aligned}$$

Summing over suitable $v, w \in W_n$ and using (6.9), we get

$$\begin{aligned} (6.12) \quad &\sum_{\substack{v, w \in W_n; \\ d_{G_n}(v, w) \leq 2}} I_m(f; v, w) \\ &\leq 3^{p-1} L_*^2 (\#\mathbb{V}_{n+m})^{-2} (\#\mathbb{V}_m) \sum_{w \in W_n} \max_{y \in \mathbb{V}_0^w} \sum_{x \in \mathbb{V}_m^w} |f(x) - f(y)|^p \\ &\leq 2^{-D} 3^{p-1} L_*^4 a^{-\alpha(2n+m)} \sum_{w \in W_n} \max_{y \in \mathbb{V}_0^w} \sum_{x \in \mathbb{V}_m^w} |f(x) - f(y)|^p. \end{aligned}$$

To obtain estimates of the sums in (6.12), we will find “good sequences in \mathbb{V}_{n+m} ”. Let $z \in w \cdot W_m$ and define $z(k) := [z]_k \in W_k$ for $k = n, \dots, n+m$. Let $q_k \in \mathbb{V}_0^{z(k)}$ for $k = n, \dots, n+m$. Then, by using Hölder's inequality repeatedly,

$$\begin{aligned} (6.13) \quad &|f(q_n) - f(q_{n+m})|^p \\ &\leq 2^{p-1} (|f(q_n) - f(q_{n+1})|^p + |f(q_{n+1}) - f(q_{n+m})|^p) \\ &\leq 2^{p-1} |f(q_n) - f(q_{n+1})|^p \\ &\quad + 2^{2(p-1)} (|f(q_{n+1}) - f(q_{n+2})|^p + |f(q_{n+2}) - f(q_{n+m})|^p) \\ &\leq \dots \leq 2^{p-1} \sum_{k=0}^{m-1} 2^{k(p-1)} |f(q_{n+k}) - f(q_{n+k+1})|^p. \end{aligned}$$

Since $q_k, q_{k+1} \in \mathbb{V}_1^{z(k)}$ for any k , there exists $[x_k(0), \dots, x_k(L)]$ such that $x_k(j) \in \mathbb{V}_1^{z(k)}$, $x_k(0) = q_k$, $x_k(L) = q_{k+1}$ and $d_{G_{k+1}}(x_k(j), x_k(j+1)) \leq 1$, where $L := \text{diam}(\mathbb{G}_1, d_{G_1}) \leq \#W_1 = N_*$. Again by Hölder's inequality, for $k = n, \dots, n+m$,

$$|f(q_{n+k}) - f(q_{n+k+1})|^p \leq L^{p-1} \sum_{j=0}^{L-1} |f(x_{n+k}(j)) - f(x_{n+k}(j+1))|^p.$$

We can assume that $[x_k(0), \dots, x_k(L)]$ is a simple path in the sense that

$$\max_{\substack{k=n, \dots, n+m; \\ (q, q') \in \mathbb{E}_1^{z(k)}}} \#\{j \mid (q, q') = (x_k(j), x_k(j+1))\} = 1.$$

Then we have

$$|f(q_{n+k}) - f(q_{n+k+1})|^p \leq 2L^{p-1} \mathcal{E}_p^{\mathbb{G}_1^{z(n+k)}}(f) \leq 2N_*^{p-1} \mathcal{E}_p^{\mathbb{G}_1^{z(n+k)}}(f).$$

Combining with (6.13), we get

$$(6.14) \quad |f(q_n) - f(q_{n+m})|^p \leq 2^p N_*^{p-1} \sum_{k=0}^{m-1} 2^{k(p-1)} \mathcal{E}_p^{\mathbb{G}_1^{z(n+k)}}(f).$$

Note that choices of $w \in W_n$, q_n and q_{n+m} are arbitrary. Hence, by noting that

$$\max_{v \in W_{n+k}} \#\{z \in w \cdot W_m \mid [z]_{n+k} = v\} = N_*^{m-k} = a^{\alpha(m-k)} \quad (k = 0, \dots, m)$$

and using (6.14), we obtain the following estimate of the sum in (6.12):

$$\begin{aligned} & \sum_{w \in W_n} \max_{y \in \mathbb{V}_0^w} \sum_{x \in \mathbb{V}_m^w} |f(x) - f(y)|^p \\ & \leq \sum_{w \in W_n} \max_{y \in \mathbb{V}_0^w} \sum_{z \in w \cdot W_m} \sum_{x \in \mathbb{V}_0^z} |f(x) - f(y)|^p \\ & \leq 2^{p+D} N_*^{p-1} \sum_{w \in W_n} \sum_{z \in w \cdot W_m} \sum_{k=0}^{m-1} 2^{k(p-1)} \mathcal{E}_p^{\mathbb{G}_1^{z(n+k)}}(f) \\ & \leq 2^{p+D} N_*^{p-1} \sum_{k=0}^{m-1} 2^{k(p-1)} a^{\alpha(m-k)} \sum_{w \in W_n} \sum_{z' \in w \cdot W_k} \mathcal{E}_p^{\mathbb{G}_1^{z'}}(f) \\ & \leq 2^{p+D} N_*^{p-1} L_* a^{\alpha m} \sum_{k=0}^{m-1} 2^{k(p-1)} a^{-\alpha k} \mathcal{E}_p^{\mathbb{G}_{n+k+1}}(f). \end{aligned}$$

Hence (6.12) becomes

$$(6.15) \quad \sum_{\substack{v, w \in W_n; \\ d_{G_n}(v, w) \leq 2}} I_m(f; v, w) \leq C_1 a^{-2\alpha n} \sum_{k=1}^m 2^{k(p-1)} a^{-\alpha k} \mathcal{E}_p^{\mathbb{G}_{n+k}}(f).$$

where $C_1 = 2a(3N_*)^{p-1} L_*^5$.

From (6.11), (6.15) and (6.9), we see that

$$(6.16) \quad \begin{aligned} & a^{(\alpha+\beta_p)n} \int_K \int_{U_1(x, a^{-n})} |f(x) - f(y)|^p d\mu_{n+m}(y) d\mu_{n+m}(x) \\ & \leq C_1 L_* a^{(\beta_p - \alpha)n} \sum_{k=1}^m 2^{k(p-1)} a^{-\alpha k} \mathcal{E}_p^{\mathbb{G}_{n+k}}(f) \\ & \leq C_1 L_* \left(\sup_{n \geq 0} a^{(\beta_p - \alpha)n} \mathcal{E}_p^{\mathbb{G}_n}(f) \right) \sum_{k=0}^{m-1} 2^{k(p-1)} a^{-\beta_p k} \leq C_2 \sup_{n \geq 0} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f). \end{aligned}$$

where $C_2 = C_1 L_* \sum_{k=0}^{\infty} a^{(p-1-\beta_p)k} < +\infty$. (Note that $p - \beta_p \leq 0$ by Proposition 3.5.)

Letting $m \rightarrow \infty$ in (6.16), we see from Lemma 6.9 that

$$a^{(\alpha+\beta_p)n} \int_K \int_{U_1(x, a^{-n})} |f(x) - f(y)|^p d\mu(y) d\mu(x) \leq c_{6.9}^{-1} C_2 \sup_{n \geq 0} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f),$$

for any $n \in \mathbb{Z}_{\geq 0}$. Since $U_1(x, a^{-n}) \supseteq B_d(x, C_{\text{AD}}^{-1} a^{-n})$ by Lemma 2.13, we conclude that $|f|_{\Lambda_{p, \infty}^{\beta_p/p}} \leq C_{\text{lower}}^{-1} \sup_{n \in \mathbb{N}} a^{(\beta_p - \alpha)n} \cdot \mathcal{E}_p^{\mathbb{G}_n}(f)$ for all $f \in \mathcal{C}(K)$, where C_{lower} depends only on $p, a, D, N_*, L_*, \rho_p, C_{\text{AR}}$. \square

6.2. Proof of Theorem 2.23. Now, we construct the desired p -energy \mathcal{E}_p on K . *Proof of Theorem 2.23.* From (6.5) and Lemma 6.7, we immediately conclude that

$$(6.17) \quad \mathcal{F}_p = \{f \in \mathcal{C}(K) \mid f \circ F_i \in \mathcal{F}_p \text{ for any } i \in S\}.$$

Since \mathcal{F}_p is separable by Theorem 5.10, there exists a countable dense subset $\mathcal{F}_p^0 = \{f_j\}_{j \geq 1}$ of \mathcal{F}_p . By virtue of Lemma 6.7, we know that

$$(6.18) \quad C_{6.7}^{-1} |f|_{\mathcal{F}_p}^p \leq \sup_{n \geq 0} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f) \leq C_{6.7} |f|_{\mathcal{F}_p}^p.$$

In particular, for each $f \in \mathcal{F}_p$, a sequence $\{\tilde{\mathcal{E}}_p^{\mathbb{G}_n}(f)\}_{n \geq 0}$ is bounded. By the diagonal argument, we can take a subsequence $\{n_k\}_{k \geq 1}$ such that

$$\left\{ \tilde{\mathcal{E}}_p^{\mathbb{G}_{n_k}}(f_j) \right\}_{k \geq 1} \text{ converges for each } j \geq 1 \text{ as } k \rightarrow \infty.$$

For simplicity, we write $\mathcal{E}_{p,n}$ to denote $\tilde{\mathcal{E}}_p^{\mathbb{G}_n}$. Note that $\mathcal{E}_{p,n}(\cdot)^{1/p}$ is a semi-norm on $\mathbb{R}^{\mathbb{V}^n}$. Let $f \in \mathcal{F}_p$, let $\varepsilon > 0$ and let $f_* \in \mathcal{F}_p^0$ such that $\|f - f_*\|_{\mathcal{F}_p} < \varepsilon$. For $k, l \geq 1$, by using the triangle inequality of $\mathcal{E}_{p,n}(\cdot)^{1/p}$,

$$\begin{aligned} & \left| \mathcal{E}_{p,n_k}(f)^{1/p} - \mathcal{E}_{p,n_l}(f)^{1/p} \right| \\ & \leq \left| \mathcal{E}_{p,n_k}(f)^{1/p} - \mathcal{E}_{p,n_k}(f_*)^{1/p} \right| \\ & \quad + \left| \mathcal{E}_{p,n_k}(f_*)^{1/p} - \mathcal{E}_{p,n_l}(f_*)^{1/p} \right| + \left| \mathcal{E}_{p,n_l}(f)^{1/p} - \mathcal{E}_{p,n_l}(f_*)^{1/p} \right| \\ & \leq 2C_{6.7} |f - f_*|_{\mathcal{F}_p} + \left| \mathcal{E}_{p,n_k}(f_*)^{1/p} - \mathcal{E}_{p,n_l}(f_*)^{1/p} \right|, \end{aligned}$$

and hence we obtain $\overline{\lim}_{k \wedge l \rightarrow \infty} \left| \mathcal{E}_{p,n_k}(f)^{1/p} - \mathcal{E}_{p,n_l}(f)^{1/p} \right| \leq 2C_{6.7}\varepsilon$. This shows that a sequence $\{\mathcal{E}_{p,n_k}(f)\}_{k \geq 1}$ is Cauchy for any $f \in \mathcal{F}_p$. We conclude that $\lim_{k \rightarrow \infty} \mathcal{E}_{p,n_k}(f)$ exists for all $f \in \mathcal{F}_p$. Now, define

$$\mathcal{E}(f) := \lim_{k \rightarrow \infty} \mathcal{E}_{p,n_k}(f), \quad f \in \mathcal{F}_p.$$

Next, for $f \in \mathcal{C}(K)$ and $m \in \mathbb{N}$, we define

$$\bar{\mathcal{E}}_{p,m}(f) := \rho_p^m \sum_{w \in W_m} \mathcal{E}(f \circ F_w),$$

and

$$\hat{\mathcal{E}}_{p,m}(f) := \frac{1}{m} \sum_{l=0}^{m-1} \bar{\mathcal{E}}_{p,l}(f),$$

where we set $\bar{\mathcal{E}}_{p,0}(f) := \mathcal{E}(f)$. Note that

$$(6.19) \quad \begin{aligned} \bar{\mathcal{E}}_{p,m}(f) &= \lim_{k \rightarrow \infty} \rho_p^m \sum_{w \in W_m} \tilde{\mathcal{E}}_p^{\mathbb{G}_{n_k}}(f \circ F_w) \\ &= \lim_{k \rightarrow \infty} \frac{\rho_p^{m+n_k}}{2} \sum_{w \in W_m} \sum_{(x,y) \in \mathbb{E}_{n_k}^w} |f(x) - f(y)|^p, \end{aligned}$$

and that

$$(6.20) \quad \hat{\mathcal{E}}_{p,m}(f) = \lim_{k \rightarrow \infty} \frac{1}{2m} \sum_{l=0}^{m-1} \rho_p^{l+n_k} \sum_{w \in W_l} \sum_{(x,y) \in \mathbb{E}_{n_k}^w} |f(x) - f(y)|^p.$$

From (6.19), Lemmas 6.7, 6.6 and 6.5-(3), for any $m \in \mathbb{Z}_{\geq 0}$, we have

$$(C_{6.7} C_{\text{WM,modif}})^{-1} |f|_{\mathcal{F}_p}^p \leq \bar{\mathcal{E}}_{p,m}(f) \leq L_* C_{6.7} |f|_{\mathcal{F}_p}^p, \quad f \in \mathcal{F}_p.$$

From this comparability and the definition of $\widehat{\mathcal{E}}_{p,m}$, it is immediate that

$$(6.21) \quad (C_{6.7}C_{\text{WM,modif}})^{-1} |f|_{\mathcal{F}_p}^p \leq \widehat{\mathcal{E}}_{p,m}(f) \leq L_* C_{6.7} |f|_{\mathcal{F}_p}^p, \quad f \in \mathcal{F}_p.$$

Hence $\{\widehat{\mathcal{E}}_{p,m}(f)\}_{m \geq 1}$ is bounded for each $f \in \mathcal{F}_p$. We also know that $\widehat{\mathcal{E}}_{p,m}(\cdot)^{1/p}$ satisfies the triangle inequality from the expression (6.20). From the argument used to define \mathcal{E} , we conclude that there exists a subsequence $\{m_j\}_{j \geq 1}$ such that

$$\left\{ \widehat{\mathcal{E}}_{p,m_j}(f) \right\}_{j \geq 1} \text{ converges for any } f \in \mathcal{F}_p.$$

Define

$$(6.22) \quad \mathcal{E}_p(f) := \lim_{j \rightarrow \infty} \widehat{\mathcal{E}}_{p,m_j}(f) \\ = \lim_{j \rightarrow \infty} \lim_{k \rightarrow \infty} \frac{1}{2m_j} \sum_{l=0}^{m_j-1} \rho_p^{l+n_k} \sum_{w \in W_l(x,y) \in \mathbb{E}_{n_k}^w} |f(x) - f(y)|^p, \quad f \in \mathcal{F}_p.$$

We shall show that \mathcal{E}_p have the desired properties.

First, we will see that $\mathcal{E}_p(\cdot)^{1/p}$ is a semi-norm, which is comparable to $|\cdot|_{\mathcal{F}_p}$, and satisfies Clarkson's inequality (see Definition 5.6). Obviously, we have from (6.21) that

$$(6.23) \quad (C_{6.7}C_{\text{WM,modif}})^{-1} |f|_{\mathcal{F}_p}^p \leq \mathcal{E}_p(f) \leq L_* C_{6.7} |f|_{\mathcal{F}_p}^p, \quad f \in \mathcal{F}_p.$$

By virtue of the expression (6.20), we can regard $\widehat{\mathcal{E}}_{p,m}(\cdot)^{1/p}$ as a limit of ℓ^p -norms on suitable finite sets, i.e.

$$I_{m,k}(f) := \left(\frac{1}{2m} \sum_{l=0}^{m-1} \rho_p^{l+n_k} \sum_{w \in W_l(x,y) \in \mathbb{E}_{n_k}^w} |f(x) - f(y)|^p \right)^{1/p}, \quad f \in \mathcal{F}_p,$$

coincides with suitable ℓ^p -norm on $\bigsqcup_{l=0}^{m-1} \bigsqcup_{w \in W_l} \mathbb{E}_{n_k}^w$ and

$$\widehat{\mathcal{E}}_{p,m}(f)^{1/p} = \lim_{k \rightarrow \infty} I_{m,k}(f).$$

By Clarkson's inequality for ℓ^p spaces, for $m, k \in \mathbb{N}$ and $f, g \in \mathcal{F}_p$,

- if $p \leq 2$, then

$$(6.24) \quad I_{m,k}(f+g)^{\frac{p}{p-1}} + I_{m,k}(f-g)^{\frac{p}{p-1}} \leq 2(I_{m,k}(f)^p + I_{m,k}(g)^p)^{\frac{1}{p-1}};$$

- if $p \geq 2$, then

$$(6.25) \quad I_{m,k}(f+g)^p + I_{m,k}(f-g)^p \leq 2^{p-1}(I_{m,k}(f)^p + I_{m,k}(g)^p).$$

Passing limits in (6.24) and (6.25), we get Clarkson's inequalities of $\mathcal{E}_p(\cdot)^{1/p}$.

Next, we will show that \mathcal{E}_p satisfies the properties (1)-(6).

- (1) Let $f \in \mathcal{C}(K)$ be constant. Then we easily see that $\widehat{\mathcal{E}}_p^{\mathbb{G}_n}(f \circ F_w) = 0$ for any $n \in \mathbb{Z}_{\geq 0}$ and $w \in W_*$. From (6.19) and (6.20), we have $\bar{\mathcal{E}}_{p,m}(f) = \widehat{\mathcal{E}}_{p,m}(f) = 0$ and thus $\mathcal{E}_p(f) = 0$. Conversely, if $\mathcal{E}_p(f) = 0$, then, by (6.21) and Lemma 6.7, $\widehat{\mathcal{E}}_p^{\mathbb{G}_n}(f) = 0$ for any $n \in \mathbb{Z}_{\geq 0}$. Lemma 6.5-(1) yields that $f|_{\bigcup_{n \geq 0} \mathbb{V}_n}$ is constant. Since $\bigcup_{n \geq 0} \mathbb{V}_n$ is dense in K , we conclude that f is constant. Next, let $f \in \mathcal{F}_p$ and let $a \in \mathbb{R}$. Then it is immediate that $\widehat{\mathcal{E}}_p^{\mathbb{G}_n}(f) = \widehat{\mathcal{E}}_p^{\mathbb{G}_n}(f + a\mathbb{1}_K)$ for any $n \in \mathbb{Z}_{\geq 0}$. Hence $\mathcal{E}_p(f) = \mathcal{E}_p(f + a\mathbb{1}_K)$.
- (2) This is proved in Theorem 5.5.

(3) Note that $\|\varphi \circ f\|_{L^p}^p \leq 2^{p-1}(\|\varphi(0)\|^p + \|f\|_{L^p}^p)$ whenever $\varphi: \mathbb{R} \rightarrow \mathbb{R}$ satisfies $\text{Lip}(\varphi) \leq 1$. Since $\widehat{\mathcal{E}}_p^{\mathbb{G}^n}$ has the Markov property (Lemma 6.5-(4)), we see that $\mathcal{E}, \bar{\mathcal{E}}_{p,m}$ and $\widehat{\mathcal{E}}_{p,m}$ also have the same property. This immediately deduces the Markov property of \mathcal{E}_p .

(4) For the same reason as (3), \mathcal{E}_p has the required symmetries (see Lemma 6.5-(2)).

(5) For any $m \in \mathbb{Z}_{\geq 0}$, the definition of $\bar{\mathcal{E}}_{p,m}$ implies that

$$\begin{aligned} \bar{\mathcal{E}}_{p,m+1}(f) &= \rho_p^{m+1} \sum_{w \in W_{m+1}} \mathcal{E}(f \circ F_w) \\ &= \rho_p^{m+1} \sum_{i \in S} \sum_{w \in W_m} \mathcal{E}(f \circ F_i \circ F_w) = \rho_p \sum_{i \in S} \bar{\mathcal{E}}_{p,m}(f \circ F_i). \end{aligned}$$

Hence, we see that

$$\begin{aligned} \rho_p \sum_{i \in S} \mathcal{E}_p(f \circ F_i) &= \rho_p \sum_{i \in S} \lim_{j \rightarrow \infty} \widehat{\mathcal{E}}_{p,m_j}(f \circ F_i) \\ &= \lim_{j \rightarrow \infty} \frac{1}{m_j} \sum_{l=0}^{m_j-1} \rho_p \sum_{i \in S} \bar{\mathcal{E}}_{p,l}(f \circ F_i) \\ &= \lim_{j \rightarrow \infty} \frac{1}{m_j} \sum_{l=0}^{m_j-1} \bar{\mathcal{E}}_{p,l+1}(f) \\ &= \lim_{j \rightarrow \infty} \frac{1}{m_j} \left(\sum_{l=0}^{m_j-1} \bar{\mathcal{E}}_{p,l}(f) + \bar{\mathcal{E}}_{p,m_j}(f) - \bar{\mathcal{E}}_{p,0}(f) \right) = \mathcal{E}_p(f). \end{aligned}$$

(6) Let $A_1 := \text{supp}[f]$ and let $A_2 := \text{supp}[g - a\mathbb{1}_K]$. Since $\text{dist}(A_1, A_2) > 0$, there exists $N \in \mathbb{N}$ such that $\sup_{n \geq N} \max_{w \in W_n} \text{diam}(K_w, d) < \text{dist}(A_1, A_2)$. Then $f \circ F_w$ or $(g - a\mathbb{1}_K) \circ F_w$ is equal to 0 for any $w \in W_n$ and $n \geq N$. From the self-similarity, we deduce that, for $n \geq N$,

$$\begin{aligned} \mathcal{E}_p(f + g) &= \mathcal{E}_p(f + g - a\mathbb{1}_K) \\ &= \rho_p^n \sum_{w \in W_n} \mathcal{E}_p(f \circ F_w + (g - a\mathbb{1}_K) \circ F_w) \\ &= \rho_p^n \sum_{w \in W_n[A_1]} \mathcal{E}_p(f \circ F_w) + \rho_p^n \sum_{w \in W_n[A_2]} \mathcal{E}_p((g - a\mathbb{1}_K) \circ F_w) \\ &= \mathcal{E}_p(f) + \mathcal{E}_p(g). \end{aligned}$$

We complete the proof. \square

Remark 6.11. (1) Since $\mathcal{E}_p(\cdot)^{1/p}$ satisfies Clarkson's inequality, $\mathcal{E}_p(\cdot)^{1/p}$ is strictly convex, that is, if $\lambda \in (0, 1)$ $f, g \in \mathcal{F}_p$ with $f - g$ is not constant, then

$$\mathcal{E}_p(\lambda f + (1 - \lambda)g)^{1/p} < \lambda \mathcal{E}_p(f)^{1/p} + (1 - \lambda) \mathcal{E}_p(g)^{1/p}.$$

The convexity of $\mathcal{E}_p(\cdot)$ is also immediate from the construction. Moreover, from the convexity of $x \mapsto |x|^p$, we can show that $\mathcal{E}_p(\cdot)$ is strictly convex.

(2) The framework in [54] includes not only the standard planar Sierpiński carpet but also Sierpiński gaskets and other self-similar sets (nested fractals for example). A recent paper by Kigami [44] gives a more general framework

to construct canonical p -energy on p -conductively homogeneous compact metric spaces, which includes new results even when $p = 2$.

For future work, it is useful to provide the following estimate concerning products of functions in \mathcal{F}_p . When $p = 2$, this result is standard (see [25, Theorem 1.4.2-(ii)] for example). See also [44, Lemma 6.17-(2)].

Proposition 6.12. *For any $f, g \in \mathcal{F}_p$,*

$$\mathcal{E}_p(f \cdot g) \leq 2^{p-1} (\|f\|_{\mathcal{C}(K)}^p \mathcal{E}_p(g) + \|g\|_{\mathcal{C}(K)}^p \mathcal{E}_p(f)).$$

In particular, $f \cdot g \in \mathcal{F}_p$.

Proof. For any $n \in \mathbb{N}$, we have

$$\begin{aligned} \mathcal{E}_p^{\mathbb{G}_n}(f \cdot g) &\leq 2^{p-1} \frac{1}{2} \sum_{(x,y) \in \mathbb{E}_n} (|g(x)|^p |f(x) - f(y)|^p + |f(y)|^p |g(x) - g(y)|^p) \\ &\leq 2^{p-1} (\|g\|_{\mathcal{C}(K)}^p \mathcal{E}_p^{\mathbb{G}_n}(f) + \|f\|_{\mathcal{C}(K)}^p \mathcal{E}_p^{\mathbb{G}_n}(g)). \end{aligned}$$

In view of the proof of Theorem 2.23, this immediately implies our assertion. \square

7. CONSTRUCTION AND BASIC PROPERTIES OF \mathcal{E}_p -ENERGY MEASURES

In this section, we construct \mathcal{E}_p -energy measures in the same way as Hino's work [35, Lemma 4.1]. We also investigate some properties, especially the chain rule of \mathcal{E}_p -energy measures. Let $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ be a generalized Sierpiński carpet. We always suppose that Assumption 4.7 holds in this section.

Let $(\mathcal{E}_p, \mathcal{F}_p)$ be the p -energy in subsection 6.2, and let $f \in \mathcal{F}_p$. For each $n \geq 0$, we define a measure $\mathbf{m}_{\langle f \rangle}^{p,n}$ on W_n (equipped with the σ -algebra 2^{W_n}) by setting

$$\mathbf{m}_{\langle f \rangle}^{p,n}(A) := \rho_p^n \sum_{w \in A} \mathcal{E}_p(f \circ F_w), \quad A \subseteq W_n.$$

Then we easily see that the total mass of $\mathbf{m}_{\langle f \rangle}^{p,n}$ is equal to $\mathcal{E}_p(f) < \infty$. Furthermore, it follows from the self-similarity of \mathcal{E}_p that, for any $A \subseteq W_n$,

$$\mathbf{m}_{\langle f \rangle}^{p,n+1}(A \cdot W_1) = \rho_p^n \sum_{w \in A} \rho_p \sum_{i \in S} \mathcal{E}_p(f \circ F_{wi}) = \rho_p^n \sum_{w \in A} \mathcal{E}_p(f \circ F_w) = \mathbf{m}_{\langle f \rangle}^{p,n}(A).$$

Therefore, $\{\mathbf{m}_{\langle f \rangle}^{p,n}\}_{n \geq 0}$ satisfies the consistency condition, and hence Kolmogorov's extension theorem (see [24, Theorem 12.1.2] for example) yields a unique Borel finite measure $\mathbf{m}_{\langle f \rangle}^p$ on Σ such that $\mathbf{m}_{\langle f \rangle}^p(\Sigma_w) = \mathbf{m}_{\langle f \rangle}^{p,|w|}(\{w\})$ for every $w \in W_*$. Then we define $\mu_{\langle f \rangle}^p := \pi_* \mathbf{m}_{\langle f \rangle}^p$, where π is the natural projection (recall Proposition 2.6). Note that $\mu_{\langle f \rangle}^p$ is Borel regular (see [24, Theorem 7.1.3] for example).

Remark 7.1. We used only the self-similar property of \mathcal{E}_p to verify the consistency condition of Kolmogorov's extension theorem. Therefore, our definition of $\mu_{\langle \cdot \rangle}^p$ works if we have a self-similar p -energy on a self-similar set. In other words, specific structures of generalized Sierpiński carpets except their self-similarities are irrelevant for the above approach.

Before proving Theorem 2.25, we observe two fundamental properties of $\mu_{\langle f \rangle}^p$.

Proposition 7.2. *Let $f \in \mathcal{F}_p$. Then $\mu_{\langle f \rangle}^p \equiv 0$ if and only if f is constant.*

Proof. It is immediate from $\mu_{\langle f \rangle}^p(K) = \mathcal{E}_p(f)$ and Theorem 2.23-(1). \square

Proposition 7.3. *For every $f, g \in \mathcal{F}_p$ and $A \in \mathcal{B}(K)$, it holds that*

$$(7.1) \quad \left| \mu_{\langle f \rangle}^p(A)^{1/p} - \mu_{\langle g \rangle}^p(A)^{1/p} \right| \leq \mu_{\langle f-g \rangle}^p(A)^{1/p}.$$

In particular, if $f_n \in \mathcal{F}_p$ converges to f in \mathcal{F}_p , then $\mu_{\langle f_n \rangle}^p(A) \rightarrow \mu_{\langle f \rangle}^p(A)$ for every $A \in \mathcal{B}(K)$.

Proof. Since $\mu_{\langle f \rangle}^p$ is Borel regular, it will suffice to prove (7.1) when A is a closed set. Let A be a closed set of K and define $C_l := \{w \in W_l \mid \Sigma_w \cap \pi^{-1}(A) \neq \emptyset\}$ for each $l \in \mathbb{N}$. Then, as proved in [35, proof of Lemma 4.1], one can show that $\{\Sigma_{C_l}\}_{l \geq 1}$ is a decreasing sequence and $\bigcap_{l \in \mathbb{N}} \Sigma_{C_l} = \pi^{-1}(A)$, where $\Sigma_{C_l} := \{\omega \in \Sigma \mid [\omega]_l \in C_l\}$.

Recall that \mathcal{E}_p is obtained as a subsequential limit of $\{\widehat{\mathcal{E}}_{p,n}\}_{n \geq 1}$, where $\widehat{\mathcal{E}}_{p,n}$ is given in (6.20). We may assume that $\mathcal{E}_p(f) = \lim_{n \rightarrow \infty} \widehat{\mathcal{E}}_{p,n}(f)$ for every $f \in \mathcal{F}_p$. For each $l, n \in \mathbb{N}$, we can regard $\left(\sum_{w \in C_l} \widehat{\mathcal{E}}_{p,n}(f \circ F_w)\right)^{1/p}$ as a limit of suitable l^p -norms on finite sets. Consequently, we have that

$$\begin{aligned} & \left| \left(\rho_p^l \sum_{w \in C_l} \widehat{\mathcal{E}}_{p,n}(f \circ F_w) \right)^{1/p} - \left(\rho_p^l \sum_{w \in C_l} \widehat{\mathcal{E}}_{p,n}(g \circ F_w) \right)^{1/p} \right| \\ & \leq \left(\rho_p^l \sum_{w \in C_l} \widehat{\mathcal{E}}_{p,n}((f-g) \circ F_w) \right)^{1/p}. \end{aligned}$$

Letting $n \rightarrow \infty$ in this inequality, we conclude that

$$\left| \mathbf{m}_{\langle f \rangle}^p(\Sigma_{C_l})^{1/p} - \mathbf{m}_{\langle g \rangle}^p(\Sigma_{C_l})^{1/p} \right| \leq \mathbf{m}_{\langle f-g \rangle}^p(\Sigma_{C_l})^{1/p},$$

for any $l \in \mathbb{N}$. Letting $l \rightarrow \infty$, we obtain (7.1) for any closed set A . \square

First, we prove Theorem 2.25-(2) and (3)

Theorem 7.4 (Theorem 2.25-(2)). *For any $\Phi \in \mathcal{C}^1(\mathbb{R})$ and $f \in \mathcal{F}_p$,*

$$(7.2) \quad d\mu_{\langle \Phi \circ f \rangle}^p = |\Phi' \circ f|^p d\mu_{\langle f \rangle}^p.$$

Proof. Note that $\Phi \circ f \in \mathcal{F}_p$ for any $f \in \mathcal{F}_p$ by the Markov property of \mathcal{E}_p (Theorem 2.23-(c)) and the compactness of $f(K)$.

First, we prove (7.2) when Φ is a polynomial. Let Φ be a polynomial and let $f \in \mathcal{F}_p$. Since $f(K)$ is compact, for any $\varepsilon > 0$ there exists $N(\varepsilon) \in \mathbb{N}$ such that

$$\left| \left| \frac{\Phi(f(y)) - \Phi(f(y'))}{f(y) - f(y')} \right|^p - |\Phi'(f(x))|^p \right| < \varepsilon$$

whenever $x, y, y' \in K_w$ for some $w \in W_n$ with $n \geq N(\varepsilon)$. Thus, for any $m \in \mathbb{N}$, $n \geq N(\varepsilon)$, $(y, y') \in \mathbb{E}_m$, $w \in W_n$ and $x \in K_w$,

$$\begin{aligned} & \left| |\Phi(f(F_w(y))) - \Phi(f(F_w(y')))|^p - |\Phi'(x)|^p |f(F_w(y)) - f(F_w(y'))|^p \right| \\ & \leq \varepsilon |f(F_w(y)) - f(F_w(y'))|^p, \end{aligned}$$

and we conclude that

$$\left| \widetilde{\mathcal{E}}_p^{\mathbb{G}_m}((\Phi \circ f) \circ F_w) - |\Phi'(f(x))|^p \widetilde{\mathcal{E}}_p^{\mathbb{G}_m}(f \circ F_w) \right| \leq \varepsilon \widetilde{\mathcal{E}}_p^{\mathbb{G}_m}(f \circ F_w).$$

Taking an appropriate limit, we have

$$\left| \mathcal{E}_p((\Phi \circ f) \circ F_w) - |\Phi'(f(x))|^p \mathcal{E}_p(f \circ F_w) \right| \leq \varepsilon \mathcal{E}_p(f \circ F_w),$$

whenever $w \in W_n$, $n \geq N(\varepsilon)$ and $f \in \mathcal{F}_p$, where \mathcal{E} is the same as in the proof of Theorem 2.23. Combining with the definition of \mathcal{E}_p , we obtain

$$(7.3) \quad |\mathcal{E}_p((\Phi \circ f) \circ F_w) - |\Phi'(f(x))|^p \mathcal{E}_p(f \circ F_w)| \leq \varepsilon \mathcal{E}_p(f \circ F_w),$$

For any $m \in \mathbb{N}$, $w \in W_m$ and $n \geq N(\varepsilon)$, we see from the self-similarity of \mathcal{E}_p that

$$\begin{aligned} & \left| \mathbf{m}_{\langle \Phi(f) \rangle}^p(\Sigma_w) - \int_{\Sigma_w} |\Phi'(f(\pi(\omega)))|^p d\mathbf{m}_{\langle f \rangle}^p(\omega) \right| \\ & \leq \sum_{v \in W_n} \int_{\Sigma_{wv}} \left| \frac{\mathcal{E}_p(F_{wv}^*(\Phi(f)))}{\mathcal{E}_p(F_{wv}^*f)} - |\Phi'(f(\pi(\omega)))|^p \right| d\mathbf{m}_{\langle f \rangle}^p(\omega) \\ & \leq \varepsilon \mathbf{m}_{\langle f \rangle}^p(\Sigma_w). \end{aligned}$$

Hence, for any $w \in W_\#$,

$$\mathbf{m}_{\langle \Phi(f) \rangle}^p(\Sigma_w) = \int_{\Sigma_w} |\Phi'(f(\pi(\omega)))|^p d\mathbf{m}_{\langle f \rangle}^p(\omega).$$

By Dynkin's π - λ theorem, we get $\mathbf{m}_{\langle \Phi(f) \rangle}^p(d\omega) = |\Phi'(f(\pi(\omega)))| \mathbf{m}_{\langle f \rangle}^p(d\omega)$. By the change of variable formula (see [24, Theorem 4.1.11] for example), we have (7.2) in this case.

Next, let $\Phi \in \mathcal{C}^1(\mathbb{R})$. Then, by applying Weierstrass' approximation theorem for Φ' , we can obtain a sequence of polynomials $\{\Phi_k\}_{k \geq 1}$ with $\Phi_k(0) = \Phi(0)$ such that $\Phi_k \rightarrow \Phi$ and $\Phi'_k \rightarrow \Phi'$ uniformly on $f(K)$. By the argument in the last paragraph, we know that

$$(7.4) \quad \mu_{\langle \Phi_k(f) \rangle}^p(dx) = |\Phi'_k(f(x))|^p \mu_{\langle f \rangle}^p(dx),$$

for every $k \in \mathbb{N}$. For any $\tilde{\Phi} \in \mathcal{C}^1(\mathbb{R})$, it is immediate that

$$\left| \tilde{\Phi}(f(F_w(y))) - \tilde{\Phi}(f(F_w(y'))) \right| \leq \sup_{s \in f(K_w)} \left| \tilde{\Phi}'(s) \right| |f(F_w(y)) - f(F_w(y'))|,$$

and hence,

$$\tilde{\mathcal{E}}_p^{\mathbb{G}_m}(F_w^*(\tilde{\Phi} \circ f)) \leq \sup_{s \in f(K_w)} \left| \tilde{\Phi}'(s) \right|^p \tilde{\mathcal{E}}_p^{\mathbb{G}_m}(f \circ F_w).$$

From the construction in subsection 6.2 and the self-similarity of \mathcal{E}_p , we get

$$(7.5) \quad \mathcal{E}_p(\tilde{\Phi} \circ f) \leq \rho_p^n \sum_{w \in W_n} \sup_{s \in f(K_w)} \left| \tilde{\Phi}'(s) \right|^p \tilde{\mathcal{E}}_p^{\mathbb{G}_m}(f \circ F_w),$$

for every $n \in \mathbb{N}$. From (7.5) and the self-similarity of \mathcal{E}_p , since the convergence $\Phi'_k \rightarrow \Phi'$ is uniform, we obtain $\lim_{k \rightarrow \infty} \mathcal{E}_p(\tilde{\Phi} \circ f - \Phi_k \circ f) = 0$. We deduce our assertion by letting $k \rightarrow \infty$ in (7.4) and applying Proposition 7.3. \square

Theorem 7.5 (Theorem 2.25-(3)). *For any $n \in \mathbb{N}$ and $f \in \mathcal{F}_p$,*

$$(7.6) \quad \mu_{\langle f \rangle}^p(dx) = \rho_p^n \sum_{w \in W_n} (F_w)_* \mu_{\langle F_w^* f \rangle}^p(dx).$$

Proof. Let $n, m \in \mathbb{N}$, let $w \in W_m$ and let $f \in \mathcal{F}_p$. If $m \leq n$, then we see that

$$\begin{aligned} \rho_p^n \sum_{v \in W_n} (\sigma_v)_* \mathbf{m}_{\langle F_v^* f \rangle}^p(\Sigma_w) &= \rho_p^n \sum_{v \in w \cdot W_{n-m}} (\sigma_v)_* \mathbf{m}_{\langle F_v^* f \rangle}^p(\Sigma_w) \\ &= \rho_p^n \sum_{v \in w \cdot W_{n-m}} \mathbf{m}_{\langle F_v^* f \rangle}^p(\Sigma) \\ &= \rho_p^{n+m} \sum_{v \in w \cdot W_{n-m}} \mathcal{E}_p(F_v^* f) = \mathbf{m}_{\langle f \rangle}^p(\Sigma_w). \end{aligned}$$

If $m \geq n$, then we have that

$$\begin{aligned} \rho_p^n \sum_{v \in W_n} (\sigma_v)_* \mathbf{m}_{\langle F_v^* f \rangle}^p(\Sigma_w) &= \rho_p^n (\sigma_{[w]_n})_* \mathbf{m}_{\langle F_{[w]_n}^* f \rangle}^p(\Sigma_w) \\ &= \rho_p^m \mathcal{E}_p(F_w^* f) = \mathbf{m}_{\langle f \rangle}^p(\Sigma_w). \end{aligned}$$

Therefore, by Dynkin's π - λ theorem, we deduce that

$$\mathbf{m}_{\langle f \rangle}^p(d\omega) = \rho_p^n \sum_{w \in W_n} (\sigma_w)_* \mathbf{m}_{\langle F_w^* f \rangle}^p(d\omega),$$

for every $n \in \mathbb{N}$. By Proposition 2.6, we have the desired result. \square

As an immediate consequence of Theorems 7.4 and 7.5, we can prove the following *energy image density property* (we borrow this naming from [14, Theorem I.7.1.1]).

Proposition 7.6. *For any $f \in \mathcal{F}_p$, it holds that the image measure of $\mu_{\langle f \rangle}^p$ by f is absolutely continuous with respect to the one-dimensional Lebesgue measure \mathcal{L}^1 on \mathbb{R} . In particular, $\mu_{\langle f \rangle}^p(\{x\}) = 0$ for any $x \in K$.*

Proof. We follow [21, Theorem 4.3.8]. It will suffice to show that $f_* \mu_{\langle f \rangle}^p(F) = 0$ whenever $f \in \mathcal{F}_p$ and F is a compact subset of \mathbb{R} with $\mathcal{L}^1(F) = 0$. We can choose a sequence $\{\varphi_n\}_{n \geq 1}$ from continuous functions on \mathbb{R} with compact supports such that $|\varphi_n| \leq 1$, $\lim_{n \rightarrow \infty} \varphi_n(x) = \mathbf{1}_F(x)$ for each $x \in \mathbb{R}$, and

$$\int_0^\infty \varphi_n(t) dt = \int_{-\infty}^0 \varphi_n(t) dt = 0,$$

for each $n \in \mathbb{N}$. Define $\Phi_n(x) := \int_0^x \varphi_n(t) dt$ for each $x \in \mathbb{R}$ and $n \in \mathbb{N}$. Then we easily see that $\Phi_n \in C^1(\mathbb{R})$ with compact support, $\Phi_n(0) = 0$, and $|\Phi_n'| \leq 1$ for each $n \in \mathbb{N}$. By the dominated convergence theorem, it is immediate that $\lim_{n \rightarrow \infty} \Phi_n(x) = 0$ for each $x \in \mathbb{R}$ and $\Phi_n \circ f$ converges to 0 in $L^p(K, \mu)$. Since $\mathcal{E}_p(\Phi_n \circ f) \leq \mathcal{E}_p(f)$ by the Markov property of \mathcal{E}_p , we deduce that $\{\Phi_n\}_{n \geq 1}$ is \mathcal{F}_p -bounded. Therefore, there exists a subsequence $\{n_k\}_{k \geq 1}$ such that $\{\Phi_{n_k} \circ f\}_{k \geq 1}$ converges to 0 weakly in \mathcal{F}_p . By Mazur's lemma, there exist $N(l) \in \mathbb{N}$ and $\{a(l)_k\}_{k=l}^{N(l)}$ with $a(l)_k \geq 0$ and $\sum_{k=l}^{N(l)} a(l)_k = 1$ such that $\Psi_l \circ f := \sum_{k=l}^{N(l)} a(l)_k \Phi_{n_k} \circ f$ converges to 0 in \mathcal{F}_p as $l \rightarrow \infty$. Then, by Fatou's lemma and the change of variable formula,

we conclude that

$$\begin{aligned}
f_*\mu_{\langle f \rangle}^p(F) &= \int_{\mathbb{R}} \lim_{l \rightarrow \infty} \left| \sum_{k=l}^{N(l)} a(l)_k \Phi'_{n_k}(t) \right|^p f_*\mu_{\langle f \rangle}^p(dt) \\
&\leq \lim_{l \rightarrow \infty} \int_K |\Psi'_l(f(x))|^p \mu_{\langle f \rangle}^p(dx) \\
&= \lim_{l \rightarrow \infty} \mu_{\langle \Psi_l \circ f \rangle}^p(K) = \lim_{l \rightarrow \infty} \mathcal{E}_p(\Psi_l \circ f) = 0. \quad \square
\end{aligned}$$

Finally, we prove Theorem 2.23-(1).

Theorem 7.7 (Theorem 2.25-(1)). *Let $f, g \in \mathcal{F}_p$. If $(f - g)|_A$ is constant for some Borel subset A of K , then $\mu_{\langle f \rangle}^p(A) = \mu_{\langle g \rangle}^p(A)$.*

Proof. Let $f \in \mathcal{F}_p$ and let $A \in \mathcal{B}(K)$. Suppose that $f|_A = c$ for some $c \in \mathbb{R}$. Then, by Proposition 7.6, we have $\mu_{\langle f \rangle}^p(f^{-1}(\{c\})) = 0$, which implies that $\mu_{\langle f \rangle}^p(A) = 0$. Combining this result and Proposition 7.3, we finish the proof. \square

We conclude this section by showing a consequence of the symmetries of \mathcal{E}_p that will be used to prove Theorem 2.27 in the next section.

Proposition 7.8. *For any $f \in \mathcal{F}_p$ and $T \in \mathcal{G}_0$, it holds that $T_*\mu_{\langle f \rangle}^p = \mu_{\langle T^*f \rangle}^p$.*

Proof. Let $A \in \mathcal{B}(K)$ be a closed set and let $T \in \mathcal{G}_0$. For each $l \in \mathbb{N}$, define

$$C_l := \{w \in W_l \mid \Sigma_w \cap \pi^{-1}(A) \neq \emptyset\} \quad \text{and} \quad C_l^T := \{w \in W_l \mid \Sigma_w \cap \pi^{-1}(T^{-1}(A)) \neq \emptyset\}.$$

Then we easily see that $\tau[T]|_{W_l}$ gives a bijection between C_l and C_l^T . Hence, for any $n \in \mathbb{N}$,

$$\sum_{w \in C_l} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(F_w^*(T^*f)) = \sum_{w \in C_l} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}((T \circ F_w)^*f) = \sum_{w \in C_l^T} \tilde{\mathcal{E}}_p^{\mathbb{G}_n}(F_w^*f).$$

From (6.20), we get $\sum_{w \in C_l} \hat{\mathcal{E}}_{p,n}(F_w^*(T^*f)) = \sum_{w \in C_l^T} \hat{\mathcal{E}}_{p,n}(F_w^*f)$, and thus

$$\begin{aligned}
\mathbf{m}_{\langle T^*f \rangle}^p(\Sigma_{C_l}) &= \rho_p^l \sum_{w \in C_l} \mathcal{E}_p(T^*f \circ F_w) \\
&= \rho_p^l \lim_{k \rightarrow \infty} \sum_{w \in C_l} \hat{\mathcal{E}}_{p,n_k}(F_w^*(T^*f)) \\
&= \rho_p^l \lim_{k \rightarrow \infty} \sum_{w \in C_l^T} \hat{\mathcal{E}}_{p,n}(F_w^*f) \\
&= \rho_p^l \sum_{w \in C_l^T} \mathcal{E}_p(f \circ F_w) = \mathbf{m}_{\langle f \rangle}^p(\Sigma_{C_l^T}).
\end{aligned}$$

Letting $l \rightarrow \infty$, we obtain $\mu_{\langle T^*f \rangle}^p(A) = T_*\mu_{\langle f \rangle}^p(A)$ because $\bigcap_{l \in \mathbb{N}} \Sigma_{C_l} = \pi^{-1}(A)$ and $\bigcap_{l \in \mathbb{N}} \Sigma_{C_l^T} = \pi^{-1}(T^{-1}(A))$ as seen in the proof of Proposition 7.3. Since both of these measures $\mu_{\langle T^*f \rangle}^p$ and $T_*\mu_{\langle f \rangle}^p$ are Borel regular, we complete the proof. \square

8. PROOF OF THEOREM 2.27

We conclude this paper by proving Theorem 2.27: a strict inequality $\beta_p > p$. Our argument is similar to [39, Section 3]. A key to prove Theorem 2.27 is the notion of \mathcal{E}_p -harmonicity (see Definition 8.3). In this section, we suppose that Assumption 4.7 holds (except Theorem 2.27) and we write $(\mathcal{E}_p, \mathcal{F}_p)$ for the p -energy on a generalized Sierpiński carpet $(K, S, \{F_i\}_{i \in S}) = \text{GSC}(D, a, S)$ in Theorem 2.23. Recall that $\mathcal{F}_p \subseteq \mathcal{C}(K)$ in this case.

Definition 8.1. Let U be a non-empty open subset of K . We define

$$(8.1) \quad \mathcal{C}^U := \{f \in \mathcal{F}_p \mid \text{supp}[f] \subseteq U\}, \quad \text{and} \quad \mathcal{F}_p^U := \overline{\mathcal{C}^U}^{\|\cdot\|_{\mathcal{F}_p}}.$$

Proposition 8.2. *It holds that*

$$(8.2) \quad \mathcal{F}_p^U = \{f \in \mathcal{F}_p \mid f(x) = 0 \text{ for any } x \in K \setminus U\}.$$

Proof. It is easy to show that $\mathcal{F}_p^U \subseteq \{f \in \mathcal{F}_p \mid f(x) = 0 \text{ for any } x \in K \setminus U\} =: \tilde{\mathcal{F}}_p^U$, so we will prove the converse. Let $f \in \tilde{\mathcal{F}}_p^U$ be non-negative. For $n \geq 1$, since f is uniformly continuous on K , we can choose $r_n > 0$ such that

$$f(x) < n^{-1} \text{ for all } x \in O_n := \bigcup_{x \in K \setminus U} B(x, r_n).$$

We can assume that $\{r_n\}_{n \geq 1}$ is non-increasing. Also, we define

$$f_n := (f - n^{-1}) \vee 0 \in \mathcal{F}_p.$$

Then it is clear that $\text{supp}[f_n] \subseteq K \setminus U$ and $f_n \rightarrow f$ in $\mathcal{C}(K)$ as $n \rightarrow \infty$. In particular, $f_n \in \mathcal{C}^U$ for all $n \geq 1$. By the Markov property of \mathcal{E}_p , we have

$$\mathcal{E}_p(f_n) \leq \mathcal{E}_p(f - n^{-1}) = \mathcal{E}_p(f),$$

and hence $\{f_n\}_{n \geq 1}$ is bounded in \mathcal{F}_p . Since \mathcal{F}_p is reflexive, there exists a subsequence $\{n_k\}_{k \geq 1}$ such that f_{n_k} converges weakly to f . By Mazur's lemma, we can find a sequence $\{g_m\}_{m \geq 1}$ from

$$\left\{ \sum_{k=1}^N a_k f_{n_k} \mid N \in \mathbb{N}, a_k \geq 0 \text{ and } \sum_{k=1}^N a_k = 1 \right\} \subseteq \mathcal{C}^U$$

such that $g_m \rightarrow f$ in \mathcal{F}_p , which completes the proof. \square

Definition 8.3. Let U be a non-empty open subset of K . For $f \in \mathcal{F}_p$, we set

$$f + \mathcal{F}_p^U := \{f + g \mid g \in \mathcal{F}_p^U\}.$$

A function $h \in \mathcal{F}_p$ is \mathcal{E}_p -harmonic on U if

$$(8.3) \quad \mathcal{E}_p(h) = \inf\{\mathcal{E}_p(f) \mid f \in h + \mathcal{F}_p^U\}.$$

Proposition 8.4. *Let U be a non-empty open subset of K with $U \neq K$ and let $g \in \mathcal{F}_p$. Then there exists a unique function $h \in \mathcal{F}_p$ that is \mathcal{E}_p -harmonic on U and $h|_{K \setminus U} \equiv g|_{K \setminus U}$.*

Proof. If $g|_{K \setminus U} \equiv a$ for some $a \in \mathbb{R}$, then $h := a$ is the required function. Suppose that $g \in \mathcal{F}_p$ is not constant on $K \setminus U$. Since g is bounded and $\mathcal{E}_p(f + a\mathbf{1}_K) = \mathcal{E}_p(f)$ for any $f \in \mathcal{F}_p$ and $a \in \mathbb{R}$, we may assume that $0 \leq g \leq 1$. Clearly, $\{f \in \mathcal{F}_p \mid f|_{K \setminus U} \equiv g|_{K \setminus U}\}$ is non-empty. For each $\lambda \geq 0$, define

$$c_\lambda := \inf\{\mathcal{E}_p(f) + \lambda \|f\|_{L^p}^p \mid f \in \mathcal{F}_p \text{ with } f|_{K \setminus U} \equiv g|_{K \setminus U}\}.$$

Note that $c_\lambda < \infty$. Let $f \in \mathcal{F}_p$ satisfy $f|_{K \setminus U} \equiv g|_{K \setminus U}$. Set $f^\# := (f \vee 0) \wedge 1 \in \mathcal{F}_p$. Then, it follows from $0 \leq g \leq 1$ that $f^\#|_{K \setminus U} = g|_{K \setminus U}$. Thus,

$$\mathcal{E}_p(f) \geq \mathcal{E}_p(f^\#) + \lambda \|f^\#\|_{L^p}^p - \lambda \geq c_\lambda - \lambda,$$

which implies that $c_0 \geq c_\lambda - \lambda$ for any $\lambda \geq 0$. For each $n \in \mathbb{N}$, we can choose $f_n \in \mathcal{F}_p$ with $f_n|_{K \setminus U} \equiv g|_{K \setminus U}$ such that

$$\mathcal{E}_p(f_n) + n^{-1} \|f_n\|_{L^p}^p < c_{n^{-1}} + n^{-1}.$$

Then $\mathcal{E}_p(f_n^\#) \leq c_0 + 2n^{-1}$ for each $n \in \mathbb{N}$, where $f_n^\# := (f_n \vee 0) \wedge 1 \in \mathcal{F}_p$. Since $\|f_n^\#\|_{L^p}^p \leq 1$ for any $n \in \mathbb{N}$, there exist $h \in L^p(K, \mu)$ and a subsequence $\{n_k\}_{k \geq 1}$ such that $\{f_{n_k}^\#\}_{k \geq 1}$ converges weakly to h in L^p . Applying Mazur's lemma, we find convex combinations $u_k = \sum_{j=k}^{N_k} a_{k,j} f_{n_j}^\#$ (i.e. $N_k \in \mathbb{N}$, $a_{k,j} \geq 0$ and $\sum_{j=k}^{N_k} a_{k,j} = 1$) such that u_k converges to h in L^p as $k \rightarrow \infty$. Note that $f_{n_j}^\#|_{K \setminus U} \equiv g|_{K \setminus U}$ and thus $u_k|_{K \setminus U} \equiv g|_{K \setminus U}$. Also, we obtain $h|_{K \setminus U} = g|_{K \setminus U}$ μ -a.e. since $u_k \rightarrow h$ in L^p as $k \rightarrow \infty$. By the triangle inequality of $\mathcal{E}_p(\cdot)^{1/p}$, we have that $\mathcal{E}_p(u_k) \in [c_0, c_0 + 2n_k^{-1}]$, which together with Clarkson's inequality implies that $\lim_{k \wedge l \rightarrow \infty} \mathcal{E}_p(u_k - u_l) = 0$. Indeed, when $p < 2$,

$$\begin{aligned} \mathcal{E}_p(u_k - u_l)^{\frac{1}{p-1}} &\leq 2(\mathcal{E}_p(u_k) + \mathcal{E}_p(u_l))^{\frac{1}{p-1}} - \mathcal{E}_p(u_k + u_l)^{\frac{1}{p-1}} \\ &\leq 2(2c_0 + 2n_k^{-1} + 2n_l^{-1})^{\frac{1}{p-1}} - 2^{\frac{p}{p-1}} c_0^{\frac{1}{p-1}} \\ &\xrightarrow{k \wedge l \rightarrow \infty} 2(2c_0)^{\frac{1}{p-1}} - 2^{\frac{p}{p-1}} c_0^{\frac{1}{p-1}} = 0. \end{aligned}$$

The case $p \geq 2$ is similar to the above. Therefore, $\{u_k\}_{k \geq 1}$ is a Cauchy sequence in \mathcal{F}_p . Since \mathcal{F}_p is a Banach space, we see that $h \in \mathcal{F}_p$ and u_k converges to h in \mathcal{F}_p . Moreover, by $\lim_{k \rightarrow \infty} \|u_k\|_{L^p} = \|h\|_{L^p}$, we conclude that $\mathcal{E}_p(h) = \lim_{k \rightarrow \infty} \mathcal{E}_p(u_k) = c_0$, that is, h is a minimizer of $\inf\{\mathcal{E}_p(f) \mid f \in \mathcal{F}_p \text{ with } f|_{K \setminus U} \equiv g|_{K \setminus U}\}$.

Lastly, we prove the uniqueness. Let $h_i \in \mathcal{F}_p$ ($i = 1, 2$) be \mathcal{E}_p -harmonic on U with $h_i|_{K \setminus U} \equiv g|_{K \setminus U}$. When $p < 2$, by Clarkson's inequality of \mathcal{E}_p ,

$$\begin{aligned} \mathcal{E}_p(h_1 - h_2)^{\frac{1}{p-1}} &\leq 2(\mathcal{E}_p(h_1) + \mathcal{E}_p(h_2))^{\frac{1}{p-1}} - \mathcal{E}_p(h_1 + h_2)^{\frac{1}{p-1}} \\ &\leq 2^{1+\frac{1}{p-1}} c_0^{\frac{1}{p-1}} - 2^{\frac{p}{p-1}} c_0^{\frac{1}{p-1}} = 0. \end{aligned}$$

Thus $h_1 - h_2$ is constant by Theorem 2.23-(1). Since $K \setminus U$ is not empty, we have that $h_1 = h_2$. The case $p \leq 2$ is similar. \square

Recall the definitions of \mathcal{K}^- , \mathcal{K}^+ (see (4.12)). It is immediate from Theorems 2.21 and 2.23-(3) that $\{f \in \mathcal{F}_p \mid f|_{\mathcal{K}^-} \equiv 0, f|_{\mathcal{K}^+} \equiv 1\} \neq \emptyset$. Thus we have the following lemma by applying Proposition 8.4 and using the symmetries of Sierpiński carpets.

Lemma 8.5. *There exists a function $h_0 \in \mathcal{F}_p$ such that $h_0|_{\mathcal{K}^-} \equiv 0$, $h_0|_{\mathcal{K}^+} \equiv 1$ and h_0 is \mathcal{E}_p -harmonic on $K \setminus (\mathcal{K}^- \cup \mathcal{K}^+)$. Moreover, it holds that $h_0 \circ R_j = h_0$ for all $j = 2, \dots, D$.*

Let h_0 be the \mathcal{E}_p -harmonic function given in Lemma 8.5. Since $\mathcal{E}_p(f) = 0$ if and only if f is constant, we immediately have that $\mathcal{E}_p(h_0) > 0$. Inductively, we define

$$(8.4) \quad h_n := \sum_{i=(i_k)_{k=1}^D \in S} (F_i)_* (a^{-1}(h_{n-1} + i_1 \mathbf{1}_K)).$$

The following proposition is clear by its definition and the self-similarity of \mathcal{E}_p .

Lemma 8.6. *For any $n \in \mathbb{Z}_{\geq 0}$, it holds that $h_n \in \mathcal{F}_p$, $h_n|_{\mathcal{K}^-} \equiv 0$, and $h_n|_{\mathcal{K}^+} \equiv 1$.*

Hereafter we suppose that $D = 2$. The following lemma is a key to prove Theorem 2.27.

Lemma 8.7. *Suppose that $D = 2$. Then the function h_2 is not \mathcal{E}_p -harmonic on $K \setminus (\mathcal{K}^- \cup \mathcal{K}^+)$.*

Proof. Set

$$\mathcal{K}_*^- := K \cap B_{2,-1}, \quad \mathcal{K}_*^+ := K \cap B_{2,+1}.$$

(Recall notations in Definition 2.3.) Suppose to the contrary that h_2 were \mathcal{E}_p -harmonic on $K \setminus (\mathcal{K}^- \cup \mathcal{K}^+)$. We claim that then a contradiction that $h_0|_{\mathcal{K}_*^+} \equiv 0$ would be implied. Let $\varphi \in \mathcal{F}_p^{K \setminus (\mathcal{K}^- \cup \mathcal{K}_*^-)}$, i.e. $\varphi \in \mathcal{F}_p$ satisfies $\varphi(x) = 0$ for any $x \in \mathcal{K}^- \cup \mathcal{K}_*^-$.

From $S \subsetneq \{0, \dots, a-1\}^2$, $a \geq 3$ and (GSC4), there exist $v_1, v_2 \in S \subseteq \mathbb{R}^2$ such that $v_1 + \mathbf{e}_1 \notin S$, $v_2 \in W_1[\mathcal{K}^+]$ and $v_2 + \mathbf{e}_2 \in S$. Now, we define $\varphi_* \in \mathcal{C}(K)$ by setting

$$\varphi_*(x) := \begin{cases} a^{-2}(F_{v_1 v_2})_* \varphi(x) & \text{if } x \in K_{v_1 v_2}, \\ a^{-2}(F_{v_1(v_2+\mathbf{e}_2)})_*(\varphi \circ R_2)(x) & \text{if } x \in K_{v_1(v_2+\mathbf{e}_2)}, \\ 0 & \text{otherwise.} \end{cases}$$

Note that φ_* is well-defined since

$$(F_{v_1 v_2})_* \varphi(x)|_{F_{v_1 v_2}(\mathcal{K}^- \cup \mathcal{K}_*^+)} = 0, \quad (F_{v_1(v_2+\mathbf{e}_2)})_* \varphi(x)|_{F_{v_1(v_2+\mathbf{e}_2)}(\mathcal{K}^- \cup \mathcal{K}_*^-)} = 0,$$

and

$$\partial(K_{v_1 v_2} \cup K_{v_1(v_2+\mathbf{e}_2)}) = F_{v_1 v_2}(\mathcal{K}^- \cup \mathcal{K}_*^+) \cup F_{v_1(v_2+\mathbf{e}_2)}(\mathcal{K}^- \cup \mathcal{K}_*^-) \cup \bigcup_{\mathbf{s} \in \{\mathbf{0}, \mathbf{e}_2\}} F_{v_1(v_2+\mathbf{s})}(\mathcal{K}^+).$$

Moreover, it follows that $\varphi_* \in \mathcal{F}_p^{K \setminus (\mathcal{K}^- \cup \mathcal{K}^+)}$. Since h_2 is \mathcal{E}_p -harmonic on $K \setminus (\mathcal{K}^- \cup \mathcal{K}^+)$, we have $\mathcal{E}_p(h_2 + \varphi_*) > \mathcal{E}_p(h_2)$ unless $\varphi \equiv 0$. Using Theorem 2.23-(4), (5) and $\varphi \circ R_2 = \varphi$, we see that

$$\begin{aligned} & \mathcal{E}_p(h_2 + \varphi_*) - \mathcal{E}_p(h_2) \\ &= \rho_p^2 \sum_{w \in W_2} (\mathcal{E}_p(F_w^* h_2 + F_w^* \varphi_*) - \mathcal{E}_p(F_w^* h_2)) \\ &= \rho_p^2 \sum_{\mathbf{s} \in \{\mathbf{0}, \mathbf{e}_2\}} \left(\mathcal{E}_p(F_{v_1(v_2+\mathbf{s})}^* h_2 + F_{v_1(v_2+\mathbf{s})}^* \varphi_*) - \mathcal{E}_p(F_{v_1(v_2+\mathbf{s})}^* h_2) \right) \\ &= 2\rho_p^2 a^{-2p} (\mathcal{E}_p(h_0 + \varphi) - \mathcal{E}_p(h_0)). \end{aligned}$$

Hence, we conclude that $\mathcal{E}_p(h_0 + \varphi) > \mathcal{E}_p(h_0)$ for any $\varphi \in \mathcal{F}_p^{K \setminus (\mathcal{K}^- \cup \mathcal{K}_*^-)} \setminus \{0\}$. This implies that h_0 is the minimizer of $\inf\{\mathcal{E}_p(f) \mid f \in \mathcal{F}_p \text{ with } f|_{\mathcal{K}^- \cup \mathcal{K}_*^-} \equiv h_0|_{\mathcal{K}^- \cup \mathcal{K}_*^-}\}$.

Next, we define $\tilde{h}_0 \in \mathcal{C}(K)$ by

$$(8.5) \quad \tilde{h}_0(x) := \begin{cases} h_0(x) & \text{if } x \in \mathcal{H}_{1,2}^{-;\leq}, \\ (h_0 \circ R_{1,2}^-)(x) & \text{if } x \in \mathcal{H}_{1,2}^{-;\geq}. \end{cases}$$

Then it is clear that $\tilde{h}_0 \in \mathcal{F}_p$ and $\tilde{h}_0|_{\mathcal{K}^- \cup \mathcal{K}_*^-} \equiv h_0|_{\mathcal{K}^- \cup \mathcal{K}_*^-}$. Moreover, we can show that $\mathcal{E}_p(\tilde{h}_0) = \mathcal{E}_p(h_0)$. To prove this, let

$$A_1 := \mathcal{H}_{1,2}^{-;\leq} \setminus \mathcal{H}_{1,2}^-, \quad A_2 := \mathcal{H}_{1,2}^- \quad \text{and} \quad A_3 := \mathcal{H}_{1,2}^{-;\geq} \setminus \mathcal{H}_{1,2}^-.$$

Since $h_0|_{A_1 \sqcup A_2} = \tilde{h}_0|_{A_1 \sqcup A_2}$, Theorem 7.7 yields $\mu_{\langle h_0 \rangle}^p(A_1 \sqcup A_2) = \mu_{\langle \tilde{h}_0 \rangle}^p(A_1 \sqcup A_2)$. Moreover, since $R_{1,2}^+(A_3) = A_3$ and $(\tilde{h}_0 \circ R_{1,2}^+)|_{A_3} = (1 - h_0)|_{A_3}$, Proposition 7.8 and Theorem 7.7 imply

$$\mu_{\langle \tilde{h}_0 \rangle}^p(A_3) = \mu_{\langle (R_{1,2}^+)^* \tilde{h}_0 \rangle}^p(A_3) = \mu_{\langle 1 - h_0 \rangle}^p(A_3) = \mu_{\langle h_0 \rangle}^p(A_3).$$

Therefore, we obtain $\mu_{\langle h_0 \rangle}^p(K) = \mu_{\langle \tilde{h}_0 \rangle}^p(K)$, which implies that $\mathcal{E}_p(\tilde{h}_0) = \mathcal{E}_p(h_0)$.

By Proposition 8.4, we have $h_0 = \tilde{h}_0$. Hence $h_0|_{\mathcal{K}^+} \equiv 0$, which contradicts the fact that $h_0((1, 1)) = 1$. We complete the proof. \square

Now we are ready to prove Theorem 2.27.

Proof of Theorem 2.27. By Lemmas 8.6 and 8.7, we obtain $\mathcal{E}_p(h_2) > \mathcal{E}_p(h_0)$. Since

$$\mathcal{E}_p(h_2) = \rho_p^2 \sum_{w \in W_2} \mathcal{E}_p(F_w^* h_2) = \rho_p^2 a^{-2p} \sum_{w \in W_2} \mathcal{E}_p(h_0),$$

we conclude that $\rho_p^2 a^{-2p} N_*^2 > 1$, which proves our assertion for $p > \dim_{\text{ARC}}(K, d)$. We know that β_p/p is monotonically non-increasing by [47, Lemma 4.7.4], and thus we obtain the desired result. \square

Remark 8.8. A recent study of L^p Besov critical exponent in [3] implies a partial result of Theorem 2.27. (Note that sub-Gaussian type heat kernels estimates on GSCs are obtained by Barlow and Bass in [7, Theorem 1.3].) Indeed, β_p is characterized as the L^p Besov critical exponent in Corollary 5.16. Thus a critical exponent $\alpha_p^\#$ in [3, equation (7)] coincides with $\beta_p/(p\beta_2)$. Therefore, [3, Theorem 3.11] gives a lower bound of β_p :

- $\beta_p \geq \frac{p\beta_2}{2}$ for $p \in (\dim_{\text{ARC}}(K, d), 2)$;
- $\beta_p \geq (p - 2)(\beta_2 - \alpha) + \beta_2$ for $p \geq 2$.

This bound implies $\beta_p > p$ for $p \in (\dim_{\text{ARC}}(K, d), (2\alpha - \beta_2)/(\alpha - \beta_2 + 1))$.

APPENDIX A. MISCELLANEOUS FACTS

A.1. Proof of Lemma 4.3. This lemma is obtained by observing that the estimates in [57, Lemma 8.4] depend only on the constants controlling rough isometries. For the reader's convenience, we give a complete proof.

Lemma A.1. *For each $i = 1, 2$, let $\{G_n^i = (V_n^i, E_n^i)\}_{n \geq 1}$ be a series of finite graphs with*

$$L_*^i := \sup_{n \in \mathbb{N}} \max_{x \in V_n^i} \#\{y \in V_n^i \mid (x, y) \in E_n^i\} < \infty,$$

and let $\varphi_n: V_n^1 \rightarrow V_n^2$ be a uniform rough isometry from $\{G_n^1\}_{n \geq 1}$ to $\{G_n^2\}_{n \geq 1}$. Then there exists a positive constant C_{URI} (depending only on C_1, C_2 in Definition 4.1, L_^1 and p) such that*

$$\mathcal{E}_p^{G_n^1}(f \circ \varphi_n) \leq C_{\text{URI}} \mathcal{E}_p^{G_n^2}(f),$$

for every $n \in \mathbb{N}$ and $f: V_n^2 \rightarrow \mathbb{R}$. In particular,

$$C_p^{G_n^1}(\varphi_n^{-1}(A_n), \varphi_n^{-1}(B_n)) \leq C_{\text{URI}} C_p^{G_n^2}(A_n, B_n)$$

for every $n \in \mathbb{N}$, where A_n, B_n are disjoint subsets of V_n^2 .

Proof. Let $n \in \mathbb{N}$, let $f: V_n^2 \rightarrow \mathbb{R}$ and let $(x, y) \in E_n^1$. We set $x' = \varphi_n(x)$ and $y' = \varphi_n(y)$. Let C_i ($i = 1, \dots, 4$) be constants in the definition of uniform rough isometry. Then we get

$$0 \leq d_{G_n^2}(x', y') \leq C_1 + C_2.$$

We set $L \in \mathbb{N}$ such that $L - 1 < C_1 + C_2 \leq L$. Since $d_{G_n^2}(x', y') \leq L$, there exist $l \leq L$ and a path $[z_0, z_1, \dots, z_l]$ in G_n^2 from x' to y' , that is $z_0 = x'$, $z_l = y'$ and $(z_{i-1}, z_i) \in E_n^2$ for each $i = 1, \dots, l$. Now, by Hölder's inequality, we have that

$$\begin{aligned} |f \circ \varphi_n(y) - f \circ \varphi_n(x)| &= |f(x') - f(y')| \leq \sum_{i=1}^l |f(z_{i-1}) - f(z_i)| \\ &\leq L^{(p-1)/p} \left(\sum_{i=1}^l |f(z_{i-1}) - f(z_i)|^p \right)^{1/p}. \end{aligned}$$

In particular, it follows that

$$(A.1) \quad |f \circ \varphi_n(y) - f \circ \varphi_n(x)|^p \leq L^{p-1} \sum_{i=1}^l |f(z_{i-1}) - f(z_i)|^p.$$

For each $(x, y) \in E_n^1$, fix a path $\gamma'_{xy} := [z'_0, \dots, z'_l]$ in G_n^2 from $\varphi_n(x)$ to $\varphi_n(y)$ with $l \leq L$. For each $(v, w) \in E_n^2$, we set

$$M_{(v,w)} := \#\{(x, y) \in E_n^1 \mid \text{path } \gamma'_{xy} \text{ contains } (v, w)\}.$$

We also define $\mathcal{A}_v^{(n)} := \{x \in V_n^1 \mid \varphi_n(x) \in B_{d_{G_n^2}}(v, L)\}$ for each $v \in V_n^2$. Then, for any $a, b \in \mathcal{A}_v^{(n)}$,

$$C_1^{-1} d_{G_n^1}(a, b) - C_2 \leq d_{G_n^2}(\varphi_n(a), \varphi_n(b)) \leq d_{G_n^2}(\varphi_n(a), v) + d_{G_n^2}(\varphi_n(b), v) \leq 2L.$$

Therefore, we have $\text{diam}(\mathcal{A}_v^{(n)}, d_{G_n^1}) \leq C_1(2L + C_2) =: C_*$, which implies that $\#\mathcal{A}_v^{(n)} \leq C_* L_*^1$ for any $n \in \mathbb{N}$ and $v \in V_n^2$. Now, since the length of $\gamma_{\varphi_n(x)\varphi_n(y)}$ is less than L , it follows that

$$\{(x, y) \in E_n^1 \mid \text{a path } \gamma'_{xy} \text{ contains a edge } (v, w) \in E_n^2\} \subseteq \mathcal{A}_v^{(n)} \times \mathcal{A}_v^{(n)}.$$

This yields that $\#M_{(v,w)} \leq (C_* L_*^1)^2$ for any $n \in \mathbb{N}$ and $(v, w) \in E_n^2$. Using this bound and summing (A.1) over $(x, y) \in E_n^1$, we conclude that

$$\mathcal{E}_p^{G_n^1}(f \circ \varphi_n) \leq L^{p-1} (C_* D_*^1)^2 \mathcal{E}_p^{G_n^2}(f). \quad \square$$

A.2. Proof of Lemma 5.13. We prove Lemma 5.13 in a metric measure space setting by extending [28, Theorem 4.11-(iii)]. Let (\mathcal{X}, d, μ) be α -Ahlfors regular, that is, (\mathcal{X}, d) is a non-empty metric space, μ is a Borel regular measure on (\mathcal{X}, d) without point mass, and there exist $\alpha > 0$ and $C_{\text{AR}} \geq 1$ such that

$$C_{\text{AR}}^{-1} r^\alpha \leq \mu(B_d(x, r)) \leq C_{\text{AR}} r^\alpha$$

for every $x \in X$ and $r \in (0, \text{diam}(\mathcal{X}, d))$. Note that $\dim_{\text{H}}(\mathcal{X}, d) = \alpha$.

Lemma A.2. *Let $\beta > \alpha$ and $p > 1$. Then there exists a positive constant $C_{5.13}$ (depending only on $p, \beta, \alpha, C_{\text{AR}}$) such that*

$$\begin{aligned} & |f(x) - f(y)|^p \\ & \leq C_{5.13} d(x, y)^{\beta-\alpha} \sup_{r \in (0, 3d(x, y)]} r^{-\beta} \int_{\mathcal{X}} \int_{B_d(z, r)} |f(z) - f(z')|^p d\mu(z') d\mu(z), \end{aligned}$$

for every $f \in \Lambda_{p, \infty}^{\beta/p}$ and μ -a.e. $x, y \in \mathcal{X}$.

Proof. For $f \in L_{\text{loc}}^1(\mathcal{X}, \mu)$, $x \in \mathcal{X}$ and $r > 0$, we set $f_{B_d(x, r)} := \int_{B_d(x, r)} f(z) d\mu(z)$ and

$$A_r(f) := \sup_{\rho \in (0, 3r]} \rho^{-\beta} \int_{\mathcal{X}} \int_{B_d(z, \rho)} |f(z) - f(z')|^p d\mu(z') d\mu(z).$$

Let $f \in \Lambda_{p, \infty}^{\beta/p}$, let $x \neq y \in \mathcal{X}$ and $r > 0$ such that $d(x, y) \leq r$. By Fubini's theorem,

$$f_{B_d(x, r)} = \frac{1}{\mu(B_d(x, r))\mu(B_d(y, r))} \int_{B_d(x, r)} \int_{B_d(y, r)} f(z) d\mu(z') d\mu(z).$$

Also, we have

$$f_{B_d(y, r)} = \frac{1}{\mu(B_d(x, r))\mu(B_d(y, r))} \int_{B_d(x, r)} \int_{B_d(y, r)} f(z') d\mu(z') d\mu(z).$$

From these identities, we have that

$$\begin{aligned} \text{(A.2)} \quad & |f_{B_d(x, r)} - f_{B_d(y, r)}|^p \\ & = \left| \frac{1}{\mu(B_d(x, r))\mu(B_d(y, r))} \int_{B_d(x, r)} \int_{B_d(y, r)} (f(z) - f(z')) d\mu(z') d\mu(z) \right|^p \\ & \leq \frac{1}{\mu(B_d(x, r))\mu(B_d(y, r))} \int_{B_d(x, r)} \int_{B_d(y, r)} |f(z) - f(z')|^p d\mu(z') d\mu(z) \\ & \leq \frac{1}{\mu(B_d(x, r))\mu(B_d(y, r))} \int_{\mathcal{X}} \int_{B_d(z, 3r)} |f(z) - f(z')|^p d\mu(z') d\mu(z) \\ \text{(A.3)} \quad & \leq c_1 r^{-\alpha} \int_{\mathcal{X}} \int_{B_d(z, 3r)} |f(z) - f(z')|^p d\mu(z') d\mu(z) \leq c_1 r^{-\alpha+\beta} A_r(f), \end{aligned}$$

where we used Hölder's inequality in the third line and the Ahlfors regularity in the fifth line. (c_1 is a positive constant depending only on C_{AR} .) Similarly, we obtain

$$\text{(A.4)} \quad |f_{B_d(x, 2r)} - f_{B_d(x, r)}|^p \leq c_1 r^{-\alpha+\beta} A_r(f).$$

Next, let \mathcal{X}_* be the set of Lebesgue points with respect to f and let $r > 0$. By Lebesgue's differential theorem on Ahlfors regular metric measure space (see [31, Theorem 1.8] for example), it holds that $\mu(\mathcal{X} \setminus \mathcal{X}_*) = 0$. Set $r_k := 2^{-k}r$ for any $k \in \mathbb{Z}_{\geq 0}$. Then, for any $x \in \mathcal{X}_*$ and any $\varepsilon > 0$ there exists $K \in \mathbb{N}$ such that $|f(x) - f_{B_d(x, r_k)}| < \varepsilon$ for all $k \geq K$. Now we have that

$$\begin{aligned} & |f(x) - f_{B_d(x, r)}| \\ & \leq |f(x) - f_{B_d(x, r_k)}| + |f_{B_d(x, r_k)} - f_{B_d(x, r_0)}| \leq \varepsilon + \sum_{k=0}^{\infty} |f_{B_d(x, r_k)} - f_{B_d(x, r_{k+1})}|. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, we get

$$|f(x) - f_{B_d(x,r)}| \leq \sum_{k=0}^{\infty} |f_{B_d(x,r_k)} - f_{B_d(x,r_{k+1})}|,$$

for any $x \in \mathcal{X}_*$. From this inequality and (A.4), we see that

$$(A.5) \quad |f(x) - f_{B_d(x,r)}| \leq \sum_{k=0}^{\infty} |f_{B_d(x,r_k)} - f_{B_d(x,2r_k)}| \leq c_2 r^{(\beta-\alpha)/p} A_r(f)^{1/p},$$

where $c_2 := c_1^{1/p} \sum_{k=0}^{\infty} 2^{-k(\beta-\alpha)/p}$.

Since μ has no point mass, it holds that $\mathcal{X}_* \setminus \{x\} \neq \emptyset$ for any $x \in \mathcal{X}_*$. Let $y \in \mathcal{X}_* \setminus \{x\}$ and set $r := d(x, y) > 0$. From (A.2) and (A.5), we conclude that

$$\begin{aligned} |f(x) - f(y)| &\leq |f(x) - f_{B_d(x,r)}| + |f_{B_d(x,r)} - f_{B_d(y,r)}| + |f(y) - f_{B_d(y,r)}| \\ &\leq c_3 r^{(\beta-\alpha)/p} A_r(f)^{1/p}, \end{aligned}$$

where $c_3 := c_1^{1/p} + 2c_2$. This proves our assertion. \square

REFERENCES

- [1] Nathan Albin, Jason Clemens, Nethali Fernando, and Pietro Poggi-Corradini, *Blocking duality for p -modulus on networks and applications*, Ann. Mat. Pura Appl. (4) **198** (2019), no. 3, 973–999. MR3954401
- [2] Patricia Alonso Ruiz and Fabrice Baudoin, *Gagliardo-Nirenberg, Trudinger-Moser and Morrey inequalities on Dirichlet spaces*, J. Math. Anal. Appl. **497** (2021), no. 2, Paper No. 124899, 26. MR4196573
- [3] Patricia Alonso-Ruiz, Fabrice Baudoin, Li Chen, Luke Rogers, Nageswari Shanmugalingam, and Alexander Teplyaev, *Besov class via heat semigroup on Dirichlet spaces III: BV functions and sub-Gaussian heat kernel estimates*, Calc. Var. Partial Differential Equations **60** (2021), no. 5, Paper No. 170, 38. MR4290383
- [4] Martin T. Barlow, *Analysis on the Sierpiński carpet*, Analysis and geometry of metric measure spaces, 2013, pp. 27–53. MR3060498
- [5] Martin T. Barlow and Richard F. Bass, *The construction of Brownian motion on the Sierpiński carpet*, Ann. Inst. H. Poincaré Probab. Statist. **25** (1989), no. 3, 225–257. MR1023950
- [6] ———, *On the resistance of the Sierpiński carpet*, Proc. Roy. Soc. London Ser. A **431** (1990), no. 1882, 345–360. MR1080496
- [7] ———, *Brownian motion and harmonic analysis on Sierpiński carpets*, Canad. J. Math. **51** (1999), no. 4, 673–744. MR1701339
- [8] Martin T. Barlow, Richard F. Bass, Takashi Kumagai, and Alexander Teplyaev, *Uniqueness of Brownian motion on Sierpiński carpets*, J. Eur. Math. Soc. (JEMS) **12** (2010), no. 3, 655–701. MR2639315
- [9] Fabrice Baudoin and Li Chen, *Sobolev spaces and Poincaré inequalities on the Vicsek fractal*, Ann. Fenn. Math. **48** (2023), no. 1, 3–26. MR4494039
- [10] Patrick Billingsley, *Convergence of probability measures*, Second, Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons, Inc., New York, 1999. A Wiley-Interscience Publication. MR1700749
- [11] Anders Björn and Jana Björn, *Nonlinear potential theory on metric spaces*, EMS Tracts in Mathematics, vol. 17, European Mathematical Society (EMS), Zürich, 2011. MR2867756
- [12] Mats Bodin, *Discrete characterisations of Lipschitz spaces on fractals*, Math. Nachr. **282** (2009), no. 1, 26–43. MR2473129
- [13] Mario Bonk, *Quasiconformal geometry of fractals*, International Congress of Mathematicians. Vol. II, 2006, pp. 1349–1373. MR2275649
- [14] Nicolas Bouleau and Francis Hirsch, *Dirichlet forms and analysis on Wiener space*, De Gruyter Studies in Mathematics, vol. 14, Walter de Gruyter & Co., Berlin, 1991. MR1133391

- [15] Marc Bourdon and Bruce Kleiner, *Combinatorial modulus, the combinatorial Loewner property, and Coxeter groups*, Groups Geom. Dyn. **7** (2013), no. 1, 39–107. MR3019076
- [16] Marc Bourdon and Hervé Pajot, *Cohomologie l_p et espaces de Besov*, J. Reine Angew. Math. **558** (2003), 85–108. MR1979183
- [17] Haim Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Universitext, Springer, New York, 2011. MR2759829
- [18] Shiping Cao, Qingsong Gu, and Hua Qiu, *p -energies on $p.c.f.$ self-similar sets*, Adv. Math. **405** (2022), Paper No. 108517, 58. MR4437617
- [19] Shiping Cao and Hua Qiu, *Dirichlet forms on unconstrained sierpinski carpets*. preprint, arXiv:2104.01529.
- [20] Matias Carrasco Piaggio, *On the conformal gauge of a compact metric space*, Ann. Sci. Éc. Norm. Supér. (4) **46** (2013), no. 3, 495–548 (2013). MR3099984
- [21] Zhen-Qing Chen and Masatoshi Fukushima, *Symmetric Markov processes, time change, and boundary theory*, London Mathematical Society Monographs Series, vol. 35, Princeton University Press, Princeton, NJ, 2012. MR2849840
- [22] Gianni Dal Maso, *An introduction to Γ -convergence*, Progress in Nonlinear Differential Equations and their Applications, vol. 8, Birkhäuser Boston, Inc., Boston, MA, 1993. MR1201152
- [23] E. B. Davies, *Spectral theory and differential operators*, Cambridge Studies in Advanced Mathematics, vol. 42, Cambridge University Press, Cambridge, 1995. MR1349825
- [24] R. M. Dudley, *Real analysis and probability*, Cambridge Studies in Advanced Mathematics, vol. 74, Cambridge University Press, Cambridge, 2002. Revised reprint of the 1989 original. MR1932358
- [25] Masatoshi Fukushima, Yoichi Oshima, and Masayoshi Takeda, *Dirichlet forms and symmetric Markov processes*, extended, De Gruyter Studies in Mathematics, vol. 19, Walter de Gruyter & Co., Berlin, 2011. MR2778606
- [26] Jun Gao, Zhenyu Yu, and Junda Zhang, *Convergence of p -energy forms on homogeneous $p.c.f.$ self-similar sets*, Potential Anal. (2022).
- [27] Alexander Grigor'yan, *Heat kernels on metric measure spaces with regular volume growth*, Handbook of geometric analysis, No. 2, 2010, pp. 1–60. MR2743439
- [28] Alexander Grigor'yan, Jiaxin Hu, and Ka-Sing Lau, *Heat kernels on metric measure spaces and an application to semilinear elliptic equations*, Trans. Amer. Math. Soc. **355** (2003), no. 5, 2065–2095. MR1953538
- [29] Alexander Grigor'yan and Meng Yang, *Local and non-local Dirichlet forms on the Sierpiński carpet*, Trans. Amer. Math. Soc. **372** (2019), no. 6, 3985–4030. MR4009425
- [30] Qingsong Gu and Ka-Sing Lau, *Dirichlet forms and convergence of Besov norms on self-similar sets*, Ann. Acad. Sci. Fenn. Math. **45** (2020), no. 2, 625–646. MR4112250
- [31] Juha Heinonen, *Lectures on analysis on metric spaces*, Universitext, Springer-Verlag, New York, 2001. MR1800917
- [32] Juha Heinonen and Pekka Koskela, *Quasiconformal maps in metric spaces with controlled geometry*, Acta Math. **181** (1998), no. 1, 1–61. MR1654771
- [33] Juha Heinonen, Pekka Koskela, Nageswari Shanmugalingam, and Jeremy T. Tyson, *Sobolev spaces on metric measure spaces*, New Mathematical Monographs, vol. 27, Cambridge University Press, Cambridge, 2015. An approach based on upper gradients. MR3363168
- [34] P. Edward Herman, Roberto Peirone, and Robert S. Strichartz, *p -energy and p -harmonic functions on Sierpinski gasket type fractals*, Potential Anal. **20** (2004), no. 2, 125–148. MR2032945
- [35] Masanori Hino, *On singularity of energy measures on self-similar sets*, Probab. Theory Related Fields **132** (2005), no. 2, 265–290. MR2199293
- [36] ———, *Upper estimate of martingale dimension for self-similar fractals*, Probab. Theory Related Fields **156** (2013), no. 3-4, 739–793. MR3078285
- [37] Naotaka Kajino, *Remarks on non-diagonality conditions for Sierpinski carpets*, Probabilistic approach to geometry, 2010, pp. 231–241. MR2648262
- [38] ———, *Non-regularly varying and non-periodic oscillation of the on-diagonal heat kernels on self-similar fractals*, Fractal geometry and dynamical systems in pure and applied mathematics. II. Fractals in applied mathematics, 2013, pp. 165–194. MR3203831
- [39] ———, *An elementary proof that walk dimension is greater than two for Brownian motion on Sierpiński carpets*, Bull. Lond. Math. Soc. **55** (2023), no. 1, 508–521.

- [40] Naotaka Kajino and Mathav Murugan, *On singularity of energy measures for symmetric diffusions with full off-diagonal heat kernel estimates*, Ann. Probab. **48** (2020), no. 6, 2920–2951. MR4164457
- [41] ———, *On the conformal walk dimension: quasisymmetric uniformization for symmetric diffusions*, Invent. Math. **231** (2023), no. 1, 263–405. MR4526824
- [42] Naotaka Kajino and Ryosuke Shimizu, in preparation.
- [43] S. Keith and T. Laakso, *Conformal Assouad dimension and modulus*, Geom. Funct. Anal. **14** (2004), no. 6, 1278–1321. MR2135168
- [44] Jun Kigami, *Conductive homogeneity of compact metric spaces and construction of p -energy*, preprint (to appear in *Memoirs of the European Mathematical Society*), arXiv:2109.08335.
- [45] ———, *Markov property of Kusuoka-Zhou’s Dirichlet forms on self-similar sets*, J. Math. Sci. Univ. Tokyo **7** (2000), no. 1, 27–33. MR1749979
- [46] ———, *Analysis on fractals*, Cambridge Tracts in Mathematics, vol. 143, Cambridge University Press, Cambridge, 2001. MR1840042
- [47] ———, *Geometry and analysis of metric spaces via weighted partitions*, Lecture Notes in Mathematics, vol. 2265, Springer, Cham, [2020] ©2020. MR4175733
- [48] Bruce Kleiner, *The asymptotic geometry of negatively curved spaces: uniformization, geometrization and rigidity*, International Congress of Mathematicians. Vol. II, 2006, pp. 743–768. MR2275621
- [49] Nicholas J. Korevaar and Richard M. Schoen, *Sobolev spaces and harmonic maps for metric space targets*, Comm. Anal. Geom. **1** (1993), no. 3-4, 561–659. MR1266480
- [50] Pekka Koskela and Paul MacManus, *Quasiconformal mappings and Sobolev spaces*, Studia Math. **131** (1998), no. 1, 1–17. MR1628655
- [51] Takashi Kumagai, *Brownian motion penetrating fractals: an application of the trace theorem of Besov spaces*, J. Funct. Anal. **170** (2000), no. 1, 69–92. MR1736196
- [52] ———, *Anomalous random walks and diffusions: from fractals to random media*, Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. IV, 2014, pp. 75–94. MR3727603
- [53] Takashi Kumagai and Karl-Theodor Sturm, *Construction of diffusion processes on fractals, d -sets, and general metric measure spaces*, J. Math. Kyoto Univ. **45** (2005), no. 2, 307–327. MR2161694
- [54] Shigeo Kusuoka and Zhou Xian Yin, *Dirichlet forms on fractals: Poincaré constant and resistance*, Probab. Theory Related Fields **93** (1992), no. 2, 169–196. MR1176724
- [55] John M. Mackay and Jeremy T. Tyson, *Conformal dimension*, University Lecture Series, vol. 54, American Mathematical Society, Providence, RI, 2010. Theory and application. MR2662522
- [56] Kôhei Sasaya, *Some relation between spectral dimension and ahlfors regular conformal dimension on infinite graphs*, preprint, arXiv:2109.00851.
- [57] Ryosuke Shimizu, *Parabolic index of an infinite graph and ahlfors regular conformal dimension of a self-similar set* (Alexander Grigor’yan and Yuhua Sun, eds.), De Gruyter, 2021.
- [58] Robert S. Strichartz and Carto Wong, *The p -Laplacian on the Sierpinski gasket*, Nonlinearity **17** (2004), no. 2, 595–616. MR2039061
- [59] Jeremy T. Tyson, *Sets of minimal Hausdorff dimension for quasiconformal maps*, Proc. Amer. Math. Soc. **128** (2000), no. 11, 3361–3367. MR1676353

GRADUATE SCHOOL OF INFORMATICS, KYOTO UNIVERSITY, YOSHIDA-HONCHO, SAKYO-KU, KYOTO 606-8501, JAPAN

Email address: r.shimizu@acs.i.kyoto-u.ac.jp