Dynamics and oscillations of a predator–prey model with modified Leslie–Gower Holling-type II schemes time-dependent delays

Haifa Ben Fredj*

Farouk Chérif[†]

November 2, 2021

Abstract

A predator–prey system is investigated in this research, which is based on a modified version of the Leslie–Gower scheme and a Holling-type II scheme with time-dependent delays. Using Schauder's fixed-point theorem, we studied the existence of pseudo almost periodic solution for the suggested model. Based on the suitable Lyapunov functional, sufficient conditions are established for the globally attractive pseudo almost periodic solution. At the end, two numerical examples are presented to demonstrate the effectiveness of our results.

keywords: Prey-predator model, Pseudo almost periodic solution, continuous delays, Global exponential stability. **Mathematics Subject Classification (2020)** 34C27, 34D23, 93C43, 47H10.

1 Introduction

One of the most attractive and significant issues in mathematical ecology is the connection between prey and predator. The predator–prey interaction is heavily influenced by functional responses. As a result, both mathematicians and ecologists have looked at predator–prey systems with a wide variety of functional responses. In particulier, Aziz-Alaoui and Daher [3] designed and investigated the following predator–prey model with modified Leslie–Gower and Holling-type II:

$$\begin{cases} \frac{du}{dt} = \left(a_1 - bu - \frac{c_1 v}{k_1 + u}\right) u, \\ \frac{dv}{dt} = \left(a_2 - \frac{c_2 v}{k_2 + u}\right) v, \end{cases}$$

$$\tag{1}$$

with initial value $u(0) = u_0 > 0$, $v(0) = v_0 > 0$, where u(t) and v(t) stand for the prey population size and the predator population size respectively, and a_1 , a_2 , b, c_1 , c_2 , k_1 and k_2 are all positives with the ecology meaning as follows

 a_1 the growth rate of prey

 a_2 the growth rate of predator

b measures the strength of competition among individuals of species u

 c_1 is the maximum value which per capita reduction rate of u can attain

 c_2 is the maximum value which per capita reduction rate of v can attain

 k_1 measures the extent to which environment provides protection to prey u

 k_2 measures the extent to which environment provides protection to predator v

This model can treat the interaction prey-predator which based on the following assumptions

- 1. the prey growth following the logistic equation (i.e $u'(t) = (a_1 bu(t))u(t)$) in the absence of his predator.
- 2. the link between the attack rate and predator size describe following the Holling type II $\left(\frac{c_1 v}{k_1 + u}\right)$ which a predator's rate of prey consumption grows as prey density grows, but finally reaches a plateau (or asymptote) where the rate of consumption remains constant regardless of prey density increases.
- 3. the Leslie–Gower formula is built on the premise that a predator population's decrease is proportional to the availability of its favorite food per capita. It is $\frac{dv}{dt} = a_2 v \left(1 \frac{v}{\alpha u}\right)$, in which the growth of the predator population is of logistic form i.e. $\frac{dv}{dt} = a_2 v \left(1 \frac{v}{C}\right)$. Here, "C" measures the carry capacity set by the environmental resources

^{*}MaPSFA, ESSTHS, University of Sousse-Tunisia. Email: haifabenfredjh@gmail.com

[†]ISSATs and MaPSFA (LR11ES35), ESSTHS, University of Sousse-Tunisia. Email: farouk.y.cherif@gmail.com

and is proportional to prey abundance, $C = \alpha u$, where is the conversion factor of prey into predator [14, 13, 20]. In the case of severe scarcity, v can switch over to other populations but its growth will be limited by the fact that its most favorite food u is not available in abundance. This case can be taken care of by adding a positive constant k to the denominator, see [2, 3].

Many natural and man-made processes in biology, medicine, and other fields now incorporate time-delays, according to current research. Time delays occur so often in nearly every circumstance that ignoring them is ignoring reality. Kuang [12] mentioned that animals must take time to digest their food before further activities and responses take place, and hence any model of species dynamics without delays is an approximation at best. It is now beyond question that the influence of time-delay owing to the time necessary to transition from egg to adult stage, gestation duration, and other factors must be considered in a better study. The famous works of Macdonald [17], Gopalsamy [11], and Kuang [12] include detailed arguments for the relevance and use of time-delays in realistic models. As a result, the "Ordinary Differential Equation", which is at the heart of Mathematical Biology, should be replaced by the "Delay Differential Equation".

Furthermore, the occurrence of almost periodic solutions is one of the most fascinating subjects in qualitative differential equations since they may be used to dynamic of prey-predator system [15, 18, 21, 22]. An extension of the almost periodic function is the pseudo almost periodic function. It was defined in [23]. It is worth noting that due to their potential applicability in a wide range of fields, almost periodic and pseudo almost periodic solutions have received a lot of attention in the last decade [1, 4, 6].

Roughly speaking, we shall consider the following differential system of predator-prey model which incorporates the Holling type II and a modified Leslie-Gower functional response:

$$\begin{cases}
 u'(t) = \left(a_1(t) - b(t)u(t) - \frac{c_1(t)v(t - \tau_1(t))}{u(t - \sigma_1(t)) + k_1(t)}\right) u(t); \\
 v'(t) = \left(a_2(t) - \frac{c_2(t)v(t - \tau_2(t))}{u(t - \sigma_2(t)) + k_2(t)}\right) v(t),
\end{cases}$$
(2)

where $a_i, b, c_i, k_i, \tau_i, \sigma_i : [0, +\infty[\rightarrow]0, +\infty[$, i = 1, 2, are continuous functions. The term $\frac{c_2(t)v(t - \tau_2(t))}{u(t - \sigma_2(t)) + k_2(t)}$ is of this

equation is called the Leslie-Gower term and the term $\frac{c_1(t)u(t)}{u(t-\sigma_1(t))+k_1(t)}$ is the Holling II functional response. Pose

$$r = \sup_{t \in \mathbb{R}} (\tau_i(t), \sigma_i(t); i = 1, 2).$$

Denote by $BC([-r,0],\mathbb{R}^2_+)$ the set of bounded continuous functions from [-r,0] to \mathbb{R}^2_+ . If z(.) is defined on $[-r+t_0,\rho[$ with $t_0,\rho\in\mathbb{R}$, then we define $z_t\in C([-r,0],\mathbb{R}^2)$ where $z_t(\theta)=z(t+\theta)$ for all $\theta\in[-r,0]$. Notice that we restrict our selves to \mathbb{R}^2_+ -valued functions since only non-negative solutions of (2) are biologically meaningful. So, let us consider the following initial condition

$$z_{t_0} = \phi, \quad \phi = (\phi_1, \phi_2) \in BC([-r, 0], \mathbb{R}^2_+) \text{ and } \phi_1(0), \phi_2(0) > 0.$$
 (3)

We write $z_t(t_0, \phi)$ for a solution of the admissible initial value problem (2) and (3). Also, let $[t_0, \eta(\phi)]$ be the maximal right-interval of existence of $z_t(t_0, \phi)$.

2 Preliminaries and definitions

Throughout this paper, for all functions $f \in BC(\mathbb{R}, \mathbb{R})$, we note :

$$f^s = \sup_{x \in \mathbb{R}} |f| \text{ and } \quad f^i = \inf_{x \in \mathbb{R}} |f|.$$

Definition 2.1. [7]

Let $f \in BC(\mathbb{R}, \mathbb{X})$. f is said almost periodic (a,p) if for any $\epsilon > 0$, there exists $l_{\epsilon} > 0$, such that

$$\exists \tau \in [a, a + l_{\epsilon}], \forall a \in \mathbb{R}, \quad ||f(x + \tau) - f(x)||_{\mathbb{X}} < \epsilon,$$

As well know τ is called ϵ – period of f. We denote by $AP(\mathbb{R}, \mathbb{X})$ the set of such functions.

It is well known that the set $AP(\mathbb{R}, \mathbb{X})$ is a Banach space with the supremum norm:

$$||f||_{\infty} = \sup_{x \in \mathbb{R}} ||f(x)||_{\mathbb{X}}.$$

In the early 1990's, the concept of pseudo almost periodicity (p.a.p) was introduced by Zhang (see [7]). It is a generalization of the almost periodicity. Define the class of functions $PAP_0(\mathbb{R}, \mathbb{X})$) as follows:

$$PAP_0(\mathbb{R}, \mathbb{X})) = \left\{ f \in BC(\mathbb{R}, \mathbb{X}); \lim_{T \to +\infty} \int_{-T}^{T} ||f(t)||_{\mathbb{X}} dt = 0 \right\}.$$

Definition 2.2. [7]

A function $f \in BC\mathbb{R}, \mathbb{X}$) is called pseudo almost-periodic if it can be expressed as

$$f = f_1 + f_2$$
, where $f_1, f_2 \in AP(\mathbb{R}, \mathbb{X}) \times PAP_0(\mathbb{R}, \mathbb{X})$.

Proposition 2.3. [7] 1. $(PAP(\mathbb{R}, \mathbb{X})), ||.||_{\infty})$ is Banach space and

$$AP(\mathbb{R}, \mathbb{X})) \subsetneq PAP(\mathbb{R}, \mathbb{X})) \subsetneq BC(\mathbb{R}, \mathbb{X}).$$

2. For $f \in PAP(\mathbb{R}, \mathbb{X})$) and $g \in PAP(\mathbb{R}, \mathbb{R})$ with $\inf_{t \in \mathbb{R}} |g(t)| > 0$, then $\frac{f}{g} \in PAP(\mathbb{R}, \mathbb{X})$).

Definition 2.4. (*Definition 2.12*, [7])

Let $\Omega \subseteq Y$. An continuous function $f: \mathbb{R} \times \Omega \longrightarrow \mathbb{X}$ is called pseudo almost periodic (p.a.p). in t uniformly with respect $x \in \Omega$ if the two following conditions are satisfied:

- i) $\forall x \in \Omega, f(.,x) \in PAP(\mathbb{R}, \mathbb{X}),$
- ii) for all compact K of Ω , $\forall \epsilon > 0, \exists \delta > 0, \forall t \in \mathbb{R}, \forall x_1, x_2 \in K$,

$$||x_1 - x_2||_{\mathbb{Y}} \le \delta \Rightarrow ||f(t, x_1) - f(t, x_2)||_{\mathbb{X}} \le \epsilon.$$

Denote by $PAP_U(\mathbb{R} \times \Omega; \mathbb{X})$ the set of all such functions.

3 Positivity and Bounded of the solution

Theorem 3.1. Let $(u, v) \in \mathbb{R}^2$ solution of system (2). If the initial condition (3) is satisfied, then the solution (u, v) is strictly positive.

Proof. By integration from t_0 into t of the system (2), we have

$$u(t) = \phi_1(0)exp\left(\int_{t_0}^t a_1(s) - b(s)u(s) - \frac{c_1(s)v(s - \tau_1(s))}{u(s - \sigma_1(s)) + k_1(s)}ds\right),$$

$$v(t) = \phi_2(0)exp\left(\int_{t_0}^t a_2(s) - \frac{c_2(s)v(s - \tau_2(s))}{u(s - \sigma_2(s)) + k_2(s)}ds\right).$$

then it is clear that the solution (u, v) has the same sign as the initial condition (3). Hence, the solution is strictly positive.

Definition 3.2. [16]

We will say that the solution (u, v) of (2) is:

1. permanent if there exists $M_1, M_2 > 0$ such that

$$0 < \liminf_{t \to +\infty} u(t) \leq \limsup_{t \to +\infty} u(t) \leq M_1 \quad \& \quad 0 < \liminf_{t \to +\infty} v(t) \leq \limsup_{t \to +\infty} v(t) \leq M_2$$

2. is uniformly permanent if there exists $M_1 > m_1 > 0$ and $M_2 > m_2 > 0$ such that

$$m_1 \geq \liminf_{t \to +\infty} u(t) \leq \limsup_{t \to +\infty} u(t) \leq M_1 \quad \& \quad m_2 < \liminf_{t \to +\infty} v(t) \leq \limsup_{t \to +\infty} v(t) \leq M_2$$

Lemme 3.3. [5] Let a > 0, b > 0.

- 1. If $dx(t)/dt \ge x(b-ax)$, then $\lim \inf_{t\to+\infty} x(t) \ge b/a$ for $t\ge 0$ and x(0)>0;
- 2. If $dx(t)/dt \le x(b-ax)$, then $\lim \sup_{t\to +\infty} x(t) \le b/a$ for $t\ge 0$ and x(0)>0.

Theorem 3.4. [Uniform permanent]

$$(C0) a_1^i k_1^i - M_2 c_1^s > 0$$

holds, then any positive solution $(u(t), v(t))^T$ of differential system (2) satisfies

$$m_1 \leq \lim \inf_{t \to +\infty} u(t) \leq \lim \sup_{t \to +\infty} u(t) \leq M_1;$$

 $m_2 \leq \lim \inf_{t \to +\infty} v(t) \leq \lim \sup_{t \to +\infty} v(t) \leq M_2.$

where

$$\begin{cases} M_1 := \frac{a_1^s}{b^i}, & m_1 := \frac{a_1^i k_1^i - M_2 c_1^s}{b^s k_1^i} \\ M_2 := \frac{a_2^s (M_1 + k_2^s) exp(a_2^s \tau_2^s)}{c_2^i}, & m_2 := \frac{a_2^i (m_1 + k_2^i)}{c_2^s exp\left(\frac{c_2^s M_2 \tau_2^s}{k_2^i + m_1}\right)} \end{cases}$$

Proof. Let $(u(t), v(t))^T$ be any positive solution of system (2). It follows from the first equation system (2) that

$$\frac{du(t)}{dt} \le u(t) \left[a_1^s - b^i u(t) \right]; \tag{4}$$

From Lemma 3.3, we get

$$\limsup_{t \to +\infty} u(t) \le \frac{a_1^s}{b^i} := M_1. \tag{5}$$

From the second equation in system (2), we have

$$\frac{dv(t)}{dt} \le a_2^s v(t). \tag{6}$$

By integrating (6) from $t - \tau_2(t)$ to t, we have

$$v(t) \le v(t - \tau_2(t))e^{a_2^s \tau_2(t)} \le v(t - \tau_2(t))e^{a_2^s \tau_2^s}. \tag{7}$$

Then,

$$v(t - \tau_2(t)) \ge v(t)exp(-a_2^s \tau_2^s). \tag{8}$$

By (8) and the second equation of system (2)

$$\frac{dv(t)}{dt} \le v(t) \left[a_2^s - \frac{c_2^i exp(-a_2^s \tau_2^s)}{M_1 + k_2^s} v(t) \right]. \tag{9}$$

From lemma 3.3, we get

$$v(t) \le \frac{a_2^s(M_1 + k_2^s)}{c_2^i e^{-a_2^s \tau_2^s}} := M_2.$$
(10)

By positivity of the solution (u, v) and from (10), we have

$$\frac{du(t)}{dt} \ge u(t) \left[a_1^i - \frac{M_2 c_1^s}{k_1^i} - b_1^s u(t) \right]. \tag{11}$$

From lemma 3.3, we obtain

$$\liminf_{t \to +\infty} u(t) \geq \frac{a_1^i k_1^i - M_2 c_1^s}{b_1^s k_1^i} := m_1.$$

By integrating second inequality of system (2) from t to $t - \tau_2(t)$, we have

$$v(t) \ge v(t - \tau_2(t))exp\left(-\frac{c_2^s M_2}{k_2^s + m_1}\tau_2(t)\right). \tag{12}$$

Then,

$$v(t - \tau_2(t)) \le v(t) exp\left(\frac{c_2^s M_2 \tau_2(t)}{k_2^i + m_1}\right) \le v(t) exp\left(\frac{c_2^s M_2 \tau_2^s}{k_2^i + m_1}\right). \tag{13}$$

On the other hand, from the 2^{sd} equation of (2), one has

$$\frac{dv(t)}{dt} \ge v(t) \left[a_2^i - \frac{c_2^s exp\left(\frac{c_2^s M_2 \tau_2^s}{k_2^i + m_1}\right)}{m_1 + k_2^i} v(t) \right]. \tag{14}$$

which yields

$$\liminf_{t \to +\infty} v(t) \geq \frac{a_2^i(m_1 + k_2^i)}{c_2^s exp\bigg(\frac{c_2^s M_2 \tau_2^s}{k_2^i + m_1}\bigg)} := m_2.$$

Remark: If the condition (C0) is not satisfied, every solution (u, v) of system (2) is permanent. So, at this case we consider $m_1 = 0$.

4 Existence the p.a.p solution

We will offer here adequate criteria that assure the existence of the pseudo almost periodic solution of (2), as stated in the introduction. The following lemmas will be stated in order to show this conclusion.

Lemme 4.1. (Theorem 2.17,[8]). If $f \in PAP_U(\mathbb{R} \times \mathbb{X}; \mathbb{X})$ and for each bounded subset B of \mathbb{X} , f is bounded on $\mathbb{R} \times B$, then the Nymetskii operator

$$N_f: PAP(\mathbb{R}, \mathbb{X}) \to PAP(\mathbb{R}, \mathbb{X})$$
 with $N_f(u) = f(.; u(.))$

is well defined.

Lemme 4.2. Let $\psi, \tau \in PAP(\mathbb{R}, \mathbb{R})$). Then $\psi(. - \tau(.)) \in PAP(\mathbb{R}, \mathbb{R})$).

Proof. let $h(t,z) = \psi(t-z)$, since the numerical application ψ is continuous function and the space $PAP(\mathbb{R},\mathbb{R})$ is a translation invariant then

- i. for any $z \in \mathbb{R}$, the function $h(.,z) \in PAP(\mathbb{R}?\mathbb{R})$.
- ii. for all compact K of \mathbb{R} , $\forall \epsilon > 0, \exists \delta > 0, \forall t \in \mathbb{R}, \forall z_1, z_2 \in K$,

$$|z_1 - z_2| \le \delta \Rightarrow |h(t, z_1) - h(t, z_2)| \le \epsilon.$$

Furthermore ψ is bounded $(PAP(\mathbb{R},\mathbb{R}) \subset BC(\mathbb{R},\mathbb{R}))$, then h is bounded on $\mathbb{R} \times B$ where B is bounded interval. By the lemma 4.1, the Nymetskii operator

$$N_f: PAP(\mathbb{R}, \mathbb{R}) \longrightarrow PAP(\mathbb{R}, \mathbb{R})$$
 $\tau_i \longmapsto h(., \tau(.))$

is well defined for $\tau \in PAP(\mathbb{R},\mathbb{R})$. Consequently, $\left[t \longmapsto \psi(t-\tau(t))\right] \in PAP(\mathbb{R},\mathbb{R})$.

Remark 4.3. In addition, many authors suppose that ϕ is bounded or uniformly continuous or $\inf_{t \in \mathbb{R}} (1 - \tau'(t)) > 0$ for given the result. So this is the first one give the proof for any $\phi, \tau \in PAP(\mathbb{R}, \mathbb{R})$, we have $\phi(t - \tau(t)) \in PAP(\mathbb{R}, \mathbb{R})$.

Lemme 4.4.

If $a, b: t \in \mathbb{R} \to \mathbb{R}$ continuous functions, then

$$e^{-\int_s^t a(u+\alpha)du} - e^{-\int_s^t b(u)du} = \int_s^t e^{-\int_r^t a(u+\alpha)du} \cdot e^{-\int_s^r b(u)du} (b(r) - a(r+\alpha))dr, \ \forall \alpha \in \mathbb{R}.$$
 (15)

Proof. Let us denote by h the function defined by $h(t,s)=g_a(t,s)-g_b(t,s)$, where $g_a(t,s)=e^{-\int_s^t a(u+\alpha)du}$ and $g_b(t,s)=e^{-\int_s^t b(u)du}$. Then, by partial differentiation of h with respect to the first variable, we get

$$d_t h(t,s) = -a(t+\alpha)g_a(t,s) + b(t)g_a(t,s),$$

= -a(t+\alpha)h(t,s) + g_b(t,s)(b(t) - a(t+\alpha)).

Now, by multiplying by $g_a(t,r)$ and simplifying, we deduce that

$$g_b(r,s)(b(r) - a(r+\alpha))g_a(t,r) = d_r h(r,s)g_a(t,r) + a(r+\alpha)g_a(t,r)h(r,s),$$

= $d_r h(r,s)g_a(t,r) + d_r g_a(t,r)h(r,s),$
= $d_r (h(r,s)g_a(t,r)).$

Thus, by integration over the interval [s, t], we have that

$$h(t,s)g_a(t,t) - g_a(t,s)h(s,s) = \int_s^t g_b(r,s)g_a(t,r)(b(r) - a(r+\alpha))dr,$$

which implies the result by noticing that $g_a(t,t) = 1$ and h(s,s) = 0.

Corollary 4.5. If $a, b : t \in \mathbb{R} \to \mathbb{R}$ continuous functions, then

$$e^{\int_{s}^{t} a(u)du} - e^{\int_{s}^{t} b(u)du} = \int_{s}^{t} e^{\int_{r}^{t} a(u)du} e^{\int_{s}^{r} b(u)du} (a(r) - b(r)) dr.$$
 (16)

Proof.

We just replace a (b resp.) with $-a_1$ ($-b_1$ resp.) in lemma 4.4.

Lemme 4.6. If $a, f \in AP(\mathbb{R}, \mathbb{R})$, then the following function

$$F_{\infty}(t) = \int_{-\infty}^{t} e^{-\int_{s}^{t} a(u)du} f(s) \, ds, \, F^{\infty}(t) = \int_{t}^{+\infty} e^{-\int_{s}^{t} a(u)du} f(s) \, ds \in AP(\mathbb{R}), \quad \forall t \in \mathbb{R}$$

Proof. For $a \in AP(\mathbb{R}^+)$ and $f \in AP(\mathbb{R},\mathbb{R})$, for $\epsilon > 0$; $\exists l_{\epsilon} > 0$; $\exists \tau \in [l_{\epsilon} - n, l_{\epsilon} + n]$ where $n \in \mathbb{R}$, then

$$|f(t+\tau)-f(t)| \le \epsilon$$
 and $|a(t+\tau)-a(t)| \le \epsilon$, $\forall t \in \mathbb{R}$.

We pose $\epsilon = \epsilon' \frac{2\underline{a}^2}{\|f\|_{\infty} + \underline{a}}$ where $\epsilon' > 0$. Therefore, applying the lemma 4.4, we get

$$|F_{\infty}(t+\tau) - F_{\infty}(t)| = \left| \int_{-\infty}^{t} e^{-\int_{s}^{t} a(u-\tau)du} f(s-\tau) ds - \int_{-\infty}^{t} e^{-\int_{s}^{t} a(u)du} f(s) ds \right|$$

$$= \left| \int_{-\infty}^{t} e^{-\int_{s}^{t} a(u-\tau)du} f(s-\tau) ds - \int_{-\infty}^{t} e^{-\int_{s}^{t} a(u)du} f(s) ds \right|$$

$$+ \int_{-\infty}^{t} e^{-\int_{s}^{t} a(u)du} f(s-\tau) ds - \int_{-\infty}^{t} e^{-\int_{s}^{t} a(u)du} f(s-\tau) ds \right|$$

$$\leq \int_{-\infty}^{t} \left| e^{-\int_{s}^{t} a(u-\tau)du} - e^{-\int_{s}^{t} a(u)du} \right| ds ||f||_{\infty} + \int_{-\infty}^{t} e^{-2(t-s)\underline{a}} |f(s-\tau) - f(s)| ds$$

$$\leq ||f||_{\infty} \int_{-\infty}^{t} \int_{s}^{t} e^{-\int_{r}^{t} a(u-\tau)du} e^{-\int_{s}^{r} a(u)du} |a(r-\tau) - a(r)| dr ds + \frac{\epsilon}{2\underline{a}}$$

$$\leq \epsilon ||f||_{\infty} \int_{-\infty}^{t} e^{-2\underline{a}(t-s)} (t-s) ds + \frac{\epsilon}{2\underline{a}},$$

$$\leq ||f||_{\infty} \frac{\epsilon}{2a^{2}} + \frac{\epsilon}{2a} = \epsilon'.$$

and, here, we pose $\epsilon = \epsilon'' \frac{2\overline{a}^2}{\|f\|_{\infty} + \overline{a}}$ where $\epsilon'' > 0$. Therefore, applying the corollary 4.5, we get

$$|F^{\infty}(t+\tau) - F^{\infty}(t)| \le ||f||_{\infty} \frac{\epsilon}{2\overline{a}^2} + \frac{\epsilon}{2\overline{a}} = \epsilon''.$$

Define the non-linear operator Υ as follows, for each $(\phi, \psi) \in PAP(\mathbb{R}, \mathbb{R}) \times PAP(\mathbb{R}, \mathbb{R})$, $\Upsilon(\phi, \psi) = (\Upsilon_1(\phi, \psi), \Upsilon_2(\phi, \psi))$ where

$$\begin{pmatrix} \Upsilon_1(\phi, \psi)(t) \\ \Upsilon_2(\phi, \psi)(t) \end{pmatrix} = \begin{pmatrix} \int_t^{+\infty} e^{\int_s^t a_1(r)dr} \left[b_1(s)\phi^2(s) + \frac{c_1(s)\psi(s - \tau_1(s))\phi(s)}{\phi(s - \sigma_1(s)) + k_1(s)} \right] ds \\ \int_t^{+\infty} e^{\int_s^t a_2(r)dr} \frac{c_2(s)\psi(s - \tau_2(s))\psi(s)}{\phi(s - \sigma_2(s)) + k_2(s)} ds \end{pmatrix}$$

and Pose

$$\begin{pmatrix} f_1(\phi(t), \phi(t - \sigma_1(t)), \psi(t - \tau_1(t))) \\ f_2(\phi(t - \sigma_2(t)), \psi(t), \psi(t - \tau_2(t))) \end{pmatrix} = \begin{pmatrix} b_1(t)\phi^2(t) + \frac{c_1(t)\psi(t - \tau_1(t))\phi(t)}{\phi(t - \sigma_1(t)) + k_1(t)} \\ \frac{c_2(t)\psi(t - \tau_2(t))\psi(t)}{\phi(t - \sigma_2(t)) + k_2(t)} \end{pmatrix}$$

Lemme 4.7.

If all the functions $a_{1,2}, b, c_{1,2}, k_{1,2}, \tau_{1,2}$ and $\sigma_{1,2}$ are p.a.p and positives, Then Υ maps $PAP^2(\mathbb{R}, \mathbb{R}_+)$ into itself.

Proof. Fix $\phi, \psi \in PAP(\mathbb{R}, \mathbb{R}_+)$. It follows from Lemma 4.2 that $\phi(. - \sigma_1(.)), \psi(. - \tau_1(.)) \in PAP(\mathbb{R}, \mathbb{R}_+)$. From $inf_{s \in \mathbb{R}}\phi(s - \sigma_1(s)) + k_1(s) > 0$ and from proprieties of space $PAP(\mathbb{R}, \mathbb{R})$, we infer that

$$f_1(\phi(s), \phi(s - \sigma_1(s)), \psi(s - \tau_1(s))) = b_1(s)\phi^2(s) + \frac{c_1(s)\psi(s - \tau_1(s))\phi(s)}{\phi(s - \sigma_1(s)) + k_1(s)} \in PAP(\mathbb{R}_+).$$

Similarly stages,

$$f_2(\phi(s - \sigma_2(s)), \psi(s), \psi(s - \tau_2(s))) = \frac{c_2(s)\psi(s - \tau_2(s))\psi(s)}{\phi(s - \sigma_2(s)) + k_2(s)} \in PAP(\mathbb{R}, \mathbb{R}_+).$$

Consequently, for all $1 \le j \le 2$, f_j can be expressed as

$$f_j = f_{j1} + f_{j2}$$
.

where $f_{j1} \in AP(\mathbb{R}, \mathbb{R}_+)$ and $f_{j2} \in PAP_0(\mathbb{R}, \mathbb{R}, \mathbb{R}_+)$. So,

$$\Upsilon_{j}(\phi, \psi)(t) = \int_{t}^{+\infty} e^{\int_{s}^{t} a_{j}(r)dr} f_{j}(\phi(s), \phi(s - \sigma_{j}(s)), \psi(s - \tau_{j}(s))) ds,$$

$$= \int_{t}^{+\infty} e^{\int_{s}^{t} a_{j}(r)dr} \left(f_{j1}(\phi(s), \phi(s - \sigma_{j}(s)), \psi(s - \tau_{j}(s))) + f_{j2}(\phi(s), \phi(s - \sigma_{j}(s)), \psi(s - \tau_{j}(s))) \right) ds$$

$$= I_{j}(t) + II_{j}(t).$$

By lemma 4.6, $I_j(t) \in AP(\mathbb{R}, \mathbb{R}^+)$, for j = 1, 2. In the other hand, we prove that $II_j(t) \in PAP_0(\mathbb{R}, \mathbb{R}_+)$, for j = 1, 2. We have

$$\int_{-T}^{T} |II_{j}(t)| dt = \int_{-T}^{T} |\int_{t}^{+\infty} e^{\int_{s}^{t} a_{j}(r)dr} f_{j2}(\phi(s), \phi(s - \sigma_{j}(s)), \psi(s - \tau_{j}(s))) ds | dt.$$

Pose $f_{j2}(s) = f_{j2}(\phi(s), \phi(s - \sigma_j(s)), \psi(s - \tau_j(s)))$ and $\vartheta = t - s$. Then by Fubini Tonnelli's Theorem one has

$$\frac{1}{2T} \int_{-T}^{T} |II_{j}(t)| dt = \frac{1}{2T} \int_{-T}^{T} \int_{t}^{+\infty} e^{\int_{s}^{t} a_{j}(r)dr} f_{j2}(s) ds dt,
\leq \frac{1}{2T} \int_{-\infty}^{0} \int_{-T}^{T} e^{-a_{j}^{i}\vartheta} f_{j2}(t-\vartheta) dt d\vartheta,$$

Pose $s = t - \vartheta$, then

$$\frac{1}{2T}\int_{-T}^{T} \big|II_{j}(t)\big|dt \leq \int_{-\infty}^{0} \frac{1}{2T}\int_{-(T+\vartheta)}^{T+\vartheta} e^{-a_{j}^{i}\vartheta}f_{j2}(s)dsd\vartheta,$$

Since the function $f_{j2} \in PAP_0(\mathbb{R}, \mathbb{R}^+)$, then the following function Ψ_T

$$\Psi_T(\vartheta) = \frac{T+\vartheta}{T} \frac{1}{2(T+\vartheta)} \int_{-(T+\vartheta)}^{T+\vartheta} f_{j2}(s) ds$$

is bounded and $\lim_{T\to +\infty} \Psi_T(\vartheta)=0$. Hence from dominated convergence Theorem, we obtain that $II_j(t)\in PAP_0(\mathbb{R},\mathbb{R}_+)$, for j=1,2. So for all $j=1,2, \Upsilon_j(\phi,\psi)$ belongs to $PAP(\mathbb{R},\mathbb{R}_+)$ and consequently Υ belongs to $PAP^2(\mathbb{R},\mathbb{R}_+)$.

Definition 4.8. [9]Let $X \in \mathbb{R}^n$ and let A(t) be a $n \times n$ continuous matrix defined on \mathbb{R} . The linear system

$$X'(t) = A(t)X(t) \tag{17}$$

is said to admit an exponential dichotomy on \mathbb{R} if there exist positive constants k; h_1 ; h_2 a projection P (i.e $P^2 = I$), and the fundamental solution matrix Y(t) of (17) satisfying

$$||Y(t)PY^{-1}(s)|| \le ke^{-h_1(t-s)}, t \ge s$$

 $||Y(t)(I-P)Y^{-1}(s)|| \le ke^{-h_2(s-t)}, s \ge t$

where I is the identity matrix.

Lemme 4.9. [9] If the linear system (17) admits an exponential dichotomy and $f \in BC(\mathbb{R}, \mathbb{R}^n)$, then the system

$$X'(t) = A(t)X(t) + f(t)$$
(18)

has a bounded solution X(t), and

$$\widetilde{X}(t) = \int_{-\infty}^{t} Y(t)PY^{-1}(s)f(s)ds - \int_{t}^{+\infty} Y(t)(I-P)Y^{-1}(s)f(s)ds;$$
(19)

where Y(t) is the fundamental solution matrix of (17).

Theorem 4.10. (Schauder Theorem)[19]:

Let M be a non-empty convex subset of a normed space B. Let T be a continuous mapping of M into a compact set $H \subset M$. Then T has at least fixed point.

Let define the following set

$$\mathcal{M} = \{\phi, \psi \in PAP(\mathbb{R}); m_1 \leq \phi \leq M_1 \text{ and } m_2 \leq \psi \leq M_2\}.$$

Theorem 4.11. The differential system (2) has at least pap solution in M.

Proof. Fix $\phi, \psi \in \mathcal{M}$. Let us consider the following differential system

$$\begin{cases} u'(t) = a_1(t)u(t) - b(t)\phi^2(t) - \frac{c_1(t)\phi(t)\psi(t - \tau_1(t))}{k_1(t) + \phi(t - \sigma_1(t))}, \\ v'(t) = a_2(t)v(t) - \frac{c_2(t)\psi(t)\psi(t - \tau_2(t))}{k_2(t) + \phi(t - \sigma_2(t))}. \end{cases}$$
(20)

It is clearly that

$$b(t)\phi^{2}(t) + \frac{c_{1}(t)\phi(t)\psi(t-\tau_{1}(t))}{k_{1}(t) + \phi(t-\sigma_{1}(t))}, \frac{c_{2}(t)\psi(t)\psi(t-\tau_{2}(t))}{k_{2}(t) + \phi(t-\sigma_{2}(t))} \in BC(\mathbb{R}, \mathbb{R}).$$

By Lemma 4.9, the system (20) has a unique bounded solution given by

$$\widetilde{X}(t) = \begin{pmatrix} \int_{t}^{+\infty} e^{\int_{s}^{t} a_{1}(u)du} f_{1}(\phi(s), \phi(s - \sigma_{1}(s)), \psi(s - \tau_{1}(s))) ds \\ \int_{t}^{+\infty} e^{\int_{s}^{t} a_{2}(u)du} f_{2}(\phi(s), \phi(s - \sigma_{2}(s)), \psi(s - \tau_{2}(s))) ds \end{pmatrix}$$

where

$$\begin{pmatrix} f_1(\phi(t), \phi(t - \sigma_1(t)), \psi(t - \tau_1(t))) \\ f_2(\phi(t - \sigma_2(t)), \psi(t), \psi(t - \tau_2(t))) \end{pmatrix} = \begin{pmatrix} b_1(t)\phi^2(t) + \frac{c_1(t)\psi(t - \tau_1(t))\phi(t)}{\phi(t - \sigma_1(t)) + k_1(t)} \\ \frac{c_2(t)\psi(t - \tau_2(t))\psi(t)}{\phi(t - \sigma_2(t)) + k_2(t)} \end{pmatrix}.$$

Then, we consider the operator Υ . By lemma 4.7, we get Υ maps $PAP(\mathbb{R}, \mathbb{R}^+) \times PAP(\mathbb{R}, \mathbb{R}^+)$ into itself. By the way, ones has the following inequality

$$u'(t) \le a_1^s u(t) - b^i \phi^2(t).$$

By the same lemma 4.9 and by definition of M_1 , the equation $z(t) = a_1^s z(t) - b^i \phi^2(t)$ has a unique bounded solution given by

$$\widetilde{z}(t) = \int_{t}^{+\infty} e^{a_1^s(t-s)} b^i \phi^2(s) ds \text{ and } \widetilde{z}(t) \leq M_1.$$

Using the comparison theorem, we get

$$\Upsilon_1(\phi,\psi)(t) \leq \widetilde{z}(t) \leq M_1.$$

We consider the inequality (9)

$$v'(t) \le a_2^s v(t) - \frac{c_2^i exp(-a_2^s \tau_2^s)}{M_1 + k_2^s} \psi^2(t).$$

By the same lemma 4.9 and by definition of M_2 , the equation $z'(t) = a_2^s z(t) - \frac{c_2^i exp(-a_2^s \tau_2^s)}{M_1 + k_2^s} \psi^2(t)$ has a unique bounded solution given by

$$\overset{\leadsto}{z}(t) = \int_t^{+\infty} e^{a_2^s(t-s)} \frac{c_2^i exp(-a_2^s \tau_2^s)}{M_1 + k_2^s} \psi^2(s) ds \text{ and } \overset{\leadsto}{z}(t) \leq M_2.$$

Using the comparison theorem, we get

$$\Upsilon_2(\phi,\psi)(t) \leq \tilde{z}(t) \leq M_2.$$

From the inequality (11), ones has

$$\frac{du(t)}{dt} \ge a_1^i u(t) - \frac{M_2 c_1^s}{k_1^i} \phi(t) - b_1^s \phi^2(t).$$

By the same lemma 4.9 and by definition of m_1 , the equation $z'(t) = a_1^i z(t) - \frac{M_2 c_1^s}{k_1^i} \phi(t) - b_1^s \phi^2(t)$ has a unique bounded solution given by

$$\widetilde{z}(t) = \int_t^{+\infty} e^{a_1^i(t-s)} \left(\frac{M_2 c_1^s}{k_1^i} \phi(t) + b_1^s \phi^2(s) \right) ds \text{ and } \widetilde{z}(t) \ge m_1.$$

Using the comparison theorem, we get

$$\Upsilon_1(\phi,\psi)(t) \geq \widetilde{z}(t) \geq m_1.$$

From the inequality (14), ones has

$$\frac{dv(t)}{dt} \ge a_2^i v(t) - \frac{c_2^s exp\left(\frac{c_2^s M_2 \tau_2^s}{k_2^i + m_1}\right)}{m_1 + k_2^i} \psi^2(t).$$

Using the lemma 4.9 and by definition of m_2 , the equation $z(t) = a_2^i z(t) - \frac{c_2^s exp\left(\frac{c_2^s M_2 \tau_2^s}{k_2^i + m_1}\right)}{m_1 + k_2^i} \psi^2(t)$ has a unique bounded solution given by

$$\widetilde{z}(t) = \int_t^{+\infty} e^{a_2^i(t-s)} \frac{c_2^s exp\left(\frac{c_2^s M_2 \tau_2^s}{k_2^i + m_1}\right)}{m_1 + k_2^i} \psi^2(s) ds \text{ and } \widetilde{z}(t) \ge m_2.$$

Using the comparison theorem, we get

$$\Upsilon_2(\phi, \psi)(t) \ge \tilde{z}(t) \ge m_2.$$

Therefore, $\Upsilon(\mathcal{M}) \subseteq \mathcal{M}$. Next step, we prove that Υ is continuous. For $(\phi_1, \psi_1), (\phi_2, \psi_2) \in \mathcal{M}$, such that $\|\phi_1 - \phi_2\|_{\infty} \leq \frac{\min(a_1^i, a_2^i)}{6\alpha} \epsilon$ and $\|\psi_1 - \psi_2\|_{\infty} \leq \frac{\min(a_1^i, a_2^i)}{2\beta} \epsilon$, where

$$\alpha = \max\left(2.M_1b^s, \frac{c_1^sM_1M_2}{(k_1^i)^2}, \frac{c_2^sM_2^2}{(k_2^i)^2}, \frac{c_1^sM_2}{k_1^i}\right) \text{ and } \beta = \max\left(\frac{c_1^sM_1}{k_1^i}, \frac{2c_2^sM_1}{k_2^i}\right),$$

one has

$$|\Upsilon_{1}(\phi_{1},\psi_{1})(t) - \Upsilon_{1}(\phi_{2},\psi_{2})(t)| = \left| \int_{t}^{+\infty} e^{\int_{s}^{t} a_{1}(r)dr} f_{1}\left(\phi_{1}(s),\phi_{1}(s-\sigma_{1}(s)),\psi_{1}(s-\tau_{1}(s))\right)\right) ds$$

$$- \int_{t}^{+\infty} e^{\int_{s}^{t} a_{1}(r)dr} f_{1}\left(\phi_{2}(s),\phi_{2}(s-\sigma_{1}(s)),\psi_{2}(s-\tau_{1}(s))\right)) ds$$

$$= \left| \int_{t}^{+\infty} e^{\int_{s}^{t} a_{1}(r)dr} \left[b(s)\phi_{1}^{2}(s) + \frac{c_{1}(s)\psi_{1}(s-\tau_{1}(s))\phi_{1}(s)}{\phi_{1}(s-\sigma_{1}(s)) + k_{1}(s)} \right] ds$$

$$- \int_{t}^{+\infty} e^{\int_{s}^{t} a_{1}(r)dr} \left[b(s)\phi_{2}^{2}(s) + \frac{c_{1}(s)\psi_{2}(s-\tau_{1}(s))\phi_{2}(s)}{\phi_{2}(s-\sigma_{1}(s)) + k_{1}(s)} \right] ds$$

$$\leq \int_{t}^{+\infty} e^{\int_{s}^{t} a_{1}(r)dr} b(s) \left| \phi_{1}^{2}(s) - \phi_{2}^{2}(s) \right| ds$$

$$+ \int_{t}^{+\infty} e^{\int_{s}^{t} a_{1}(r)dr} c_{1}(s) \left| \frac{\psi_{1}(s-\tau_{1}(s))\phi_{1}(s)}{\phi_{1}(s-\sigma_{1}(s)) + k_{1}(s)} - \frac{\psi_{2}(s-\tau_{1}(s))\phi_{2}(s)}{\phi_{2}(s-\sigma_{1}(s)) + k_{1}(s)} \right| ds$$

For simplicity, pose that

$$\psi_{j_s}(\tau_1(s)) = \psi_j(s - \tau_1(s), \phi_{j_s}(\sigma_1(s)) = \phi_j(s - \sigma_1(s)), \text{ for } j = 1, 2.$$

We have

$$\begin{split} \left| \frac{\psi_{1_s}(\tau_1(s))\phi_1(s)}{\phi_{1_s}(\sigma_1(s)) + k_1(s)} - \frac{\psi_{2_s}(\tau_1(s))\phi_2(s)}{\phi_{2_s}(\sigma_1(s)) + k_1(s)} \right| &\leq \frac{\psi_{1_s}(\tau_1(s))}{\phi_{1_s}(\sigma_1(s)) + k_1(s)} \|\phi_1 - \phi_2\|_{\infty} + \frac{\phi_2(s)}{\phi_{1_s}(\sigma_1(s)) + k_1(s)} \|\psi_1 - \psi_2)\|_{\infty} \\ &+ \left| \frac{\psi_{2_s}(\tau_1(s))\phi_2(s)}{\phi_{1_s}(\sigma_1(s)) + k_1(s)} - \frac{\psi_{2_s}(\tau_1(s))\phi_2(s)}{\phi_{2_s}(\sigma_1(s)) + k_2(s)} \right| \\ &\leq \frac{M_2}{k_1^i} \|\phi_1 - \phi_2\|_{\infty} + \frac{M_1}{k_1^i} \|\psi_1 - \psi_2\|_{\infty} + \frac{M_1M_2}{(k_1^i)^2} \|\phi_1 - \phi_2\|_{\infty}. \end{split}$$

And we have

$$|\phi_1^2(s) - \phi_2^2(s)| = |(\phi_1(s) + \phi_2(s))(\phi_1(s) - \phi_2(s))|$$

$$\leq 2M_1 ||\phi_1 - \phi_2||_{\infty}.$$

Therefore, we obtain

$$\begin{split} |\Upsilon_1(\phi_1,\psi_1)(t) - \Upsilon_1(\phi_2,\psi_2)(t)| &\leq \int_t^{+\infty} e^{\int_s^t a_1(r)dr} \bigg\{ 2b^s N 1 \|\phi_1 - \phi_2\|_{\infty} \\ &\quad + c_1(s) \bigg(\frac{M_2}{k_1^i} + \frac{M_1 M_2}{(k_1^i)^2} \bigg) \|\phi_1 - \phi_2\|_{\infty} + \frac{M_1}{k_1^i} \|\psi_1 - \psi_2\|_{\infty} \bigg\} \, ds \\ &\leq \int_t^{+\infty} e^{a_1^i(t-s)} \bigg\{ \left(2b^s M_1 + \frac{c_1^s M_2}{k_1^i} + \frac{c_1^s M_1 M_2}{(k_1^i)^2} \right) \|\phi_1 - \phi_2\|_{\infty} \\ &\quad + \frac{c_1^s M_1}{k_1^i} \|\psi_1 - \psi_2\|_{\infty} \bigg\} \, ds \\ &\leq \int_t^{+\infty} e^{a_1^i(t-s)} \left\{ 3\alpha \|\phi_1 - \phi_2\|_{\infty} + \beta \|\psi_1 - \psi_2\|_{\infty} \right\} \, ds \\ &= 3\frac{\alpha}{a_1^i} \|\phi_1 - \phi_2\|_{\infty} + \frac{\beta}{a_1^i} \|\psi_1 - \psi_2\|_{\infty} \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{split}$$

By conclusion, Υ_1 is a continuous operator.

Now, we should prove that Υ_2 is a continuous operator.

$$\begin{split} |\Upsilon_2(\phi_1,\psi_1)(t) - \Upsilon_2(\phi_2,\psi_2)(t)| &= \left| \int_t^{+\infty} e^{\int_s^t a_2(r)dr} f_2\left(\phi_1(s-\sigma_2(s)),\psi_1(s),\psi_1(s-\tau_2(s))\right)\right) \ ds \\ &- \int_t^{+\infty} e^{\int_s^t a_2(r)dr} f_2\left(\phi_2(-\sigma_2(s)s),\psi_2(s),\psi_2(s-\tau_2(s))\right)\right) \ ds \right| \\ &= \left| \int_t^{+\infty} e^{\int_s^t a_2(r)dr} \frac{c_2(s)\psi_1(s-\tau_2(s))\psi_1(s)}{\phi_1(s-\sigma_2(s))+k_2(s)} \ ds \right| \\ &- \int_t^{+\infty} e^{\int_s^t a_2(r)dr} \frac{c_2(s)\psi_2(s-\tau_2(s))\psi_2(s)}{\phi_2(s-\sigma_2(s))+k_2(s)} \ ds \right| \\ &\leq \int_t^{+\infty} e^{\int_s^t a_2(r)dr} c_2(s) \left| \frac{\psi_{1_s}(\tau_2(s))\psi_1(s)}{\phi_{1_s}(\sigma_2(s))+k_2(s)} - \frac{\psi_{2_s}(\tau_2(s))\psi_2(s)}{\phi_{2_s}(\sigma_2(s))+k_2(s)} \right| \ ds. \end{split}$$

For j = 1 or 2, pose that

$$\psi_{j_s}(\tau_2(s)) = \psi_j(s - \tau_2(s)), \phi_{j_s}(\sigma_2(s)) = \phi_j(s - \sigma_2(s)).$$

We get

$$\begin{split} \left| \frac{\psi_{1_s}(\tau_2(s))\psi_1(s)}{\phi_{1_s}(\sigma_2(s)) + k_2(s)} - \frac{\psi_{2_s}(\tau_2(s))\psi_2(s)}{\phi_{2_s}(\sigma_2(s)) + k_2(s)} \right| \leq & \frac{\psi_{1_s}(\tau_2(s))}{\phi_{1_s}(\sigma_2(s)) + k_2(s)} \|\psi_1 - \psi_2\|_{\infty} + \frac{\psi_2(s)}{\phi_{1_s}(\sigma_2(s)) + k_2(s)} \|\psi_1 - \psi_2\|_{\infty} \\ & + \frac{\psi_{1_s}(\tau_2(s))\psi_1(s)}{(\phi_{1_s}(\sigma_2(s)) + k_2(s))(\phi_{2_s}(\sigma_2(s)) + k_2(s))} \|\phi_1 - \phi_2\|_{\infty} \\ \leq & \frac{2M_2}{k_2^i} \|\psi_1 - \psi_2\|_{\infty} + \frac{(M_2)^2}{(k_2^i)^2} \|\phi_1 - \phi_2\|_{\infty}. \end{split}$$

Thus, we get

$$\begin{split} |\Upsilon_2(\phi_1, \psi_1)(t) - \Upsilon_2(\phi_2, \psi_2)(t)| &\leq \int_t^{+\infty} e^{\int_s^t a_2(r)dr} c_2(s) \left\{ \frac{2M_2}{k_2^i} \|\psi_1 - \psi_2\|_{\infty} + \frac{M_2^2}{(k_2^i)^2} \|\phi_1 - \phi_2\|_{\infty} \right\} ds \\ &\leq \int_t^{+\infty} e^{a_2^i(t-s)} \left\{ \frac{2c_2^s M_2}{k_2^i} \|\psi_1 - \psi_2\|_{\infty} + \frac{c_2^s M_2^2}{(k_2^i)^2} \|\phi_1 - \phi_2\|_{\infty} \right\} ds \\ &= \frac{\beta}{a_2^i} \|\psi_1 - \psi_2\|_{\infty} + \frac{\alpha}{a_2^i} \|\phi_1 - \phi_2\|_{\infty} \\ &\leq \epsilon. \end{split}$$

By conclusion, Υ_2 is a continuous operator.

Next, we need to prove that Υ is a compact operator. In fact, we show that the following two statements are true:

- 1) $\{\Upsilon(\phi,\psi): (\phi,\psi) \in M\} \subset BC(\mathbb{R},\mathbb{R}^2)$ is equi-continuous.
- 2) $\{\Upsilon(\phi,\psi)(t); (\phi,\psi) \in M\}$ is relatively compact subset of \mathbb{R}^2 for each $t \in \mathbb{R}$.

To prove 1), given $u, v \in \mathbb{R}$, such that u < v and $|u - v| < \epsilon$, we have

$$\begin{split} \left| \Upsilon_{1}(\phi,\psi)(u) - \Upsilon_{1}(\phi,\psi)(v) \right| &= \left| \int_{u}^{+\infty} e^{\int_{s}^{u} a_{1}(r)dr} \left[b(s)\phi^{2}(s) + \frac{c_{1}(s)\psi(s - \tau_{1}(s))\psi(s)}{\phi(s - \sigma_{1}(s)) + k_{1}(s)} \right] ds \right. \\ &- \int_{v}^{+\infty} e^{\int_{s}^{v} a_{1}(r)dr} \left[b(s)\phi^{2}(s) + \frac{c_{1}(s)\psi(s - \tau_{1}(s))\psi(s)}{\phi(s - \sigma_{1}(s)) + k_{1}(s)} \right] ds \\ &= \left| \int_{u}^{+\infty} e^{\int_{s}^{u} a_{1}(r)dr} \left[b(s)\phi^{2}(s) + \frac{c_{1}(s)\psi(s - \tau_{1}(s))\psi(s)}{\phi(s - \sigma_{1}(s)) + k_{1}(s)} \right] ds \right. \\ &- \int_{v}^{u} e^{\int_{s}^{v} a_{1}(r)dr} \left[b(s)\phi^{2}(s) + \frac{c_{1}(s)\psi(s - \tau_{1}(s))\psi(s)}{\phi(s - \sigma_{1}(s)) + k_{1}(s)} \right] ds \\ &- e^{\int_{u}^{u} a_{1}(r)dr} \int_{u}^{+\infty} e^{\int_{s}^{u} a_{1}(r)dr} \left[b(s)\phi^{2}(s) + \frac{c_{1}(s)\psi(s - \tau_{1}(s))\psi(s)}{\phi(s - \sigma_{1}(s)) + k_{1}(s)} \right] ds \\ &\leq \left| 1 - e^{\int_{u}^{v} a_{1}(r)dr} \left| \int_{u}^{+\infty} e^{\int_{s}^{u} a_{1}(r)dr} \left[b(s)\phi^{2}(s) + \frac{c_{1}(s)\psi(s - \tau_{1}(s))\psi(s)}{\phi(s - \sigma_{1}(s)) + k_{1}(s)} \right] ds \right. \\ &+ \int_{v}^{u} e^{\int_{s}^{v} a_{1}(r)dr} \left[b(s)\phi^{2}(s) + \frac{c_{1}(s)\psi(s - \tau_{1}(s))\psi(s)}{\phi(s - \sigma_{1}(s)) + k_{1}(s)} \right] ds \end{split}$$

Or $\int_u^v a_1(r)dr > 0 \Rightarrow e^{\int_u^v a_1(r)dr} > 1$. Therefore

$$\begin{split} \left| \Upsilon_1(\phi,\psi)(u) - \Upsilon_1(\phi,\psi)(v) \right| &\leq \left(e^{\int_u^v a_1(r)dr} - 1 \right) \int_u^{+\infty} e^{\int_s^u a_1(r)dr} \left[b(s)\phi^2(s) + \frac{c_1(s)\psi(s - \tau_1(s))\psi(s)}{\phi(s - \sigma_1(s)) + k_1(s)} \right] \, ds \\ &+ \int_u^v e^{\int_s^v a_1(r)dr} \left[b(s)\phi^2(s) + \frac{c_1(s)\psi(s - \tau_1(s))\psi(s)}{\phi(s - \sigma_1(s)) + k_1(s)} \right] \, ds \\ &\leq \left(e^{a_1^s(v - u)} - 1 \right) \int_u^{+\infty} e^{\int_s^u a_1(r)dr} \left[b(s)\phi^2(s) + \frac{c_1(s)\psi(s - \tau_1(s))\psi(s)}{\phi(s - \sigma_1(s)) + k_1(s)} \right] \, ds \\ &+ \int_u^v e^{\int_s^v a_1(r)dr} \left[b(s)\phi^2(s) + \frac{c_1(s)\psi(s - \tau_1(s))\psi(s)}{\phi(s - \sigma_1(s)) + k_1(s)} \right] \, ds \\ &\leq \frac{e^{a_1^s(v - u)} - 1}{a_1^i} \left[b^s M_1^2 + \frac{c_1^s M_1 M_2}{k_1^s} \right] + \frac{e^{a_1^s(v - u)} - 1}{a_1^i} \left[b^s M_1^2 + \frac{c_1^s M_1 M_2}{k_1^s} \right]. \end{split}$$

Or

$$e^{a_1^s(v-u)} - 1 \underset{u \to v}{\to} 0.$$
 (21)

Then

$$\left| \Upsilon_1(\phi, \psi)(u) - \Upsilon_1(\phi, \psi)(v) \right| \underset{u \to v}{\longrightarrow} 0.$$

And

$$\begin{split} |\Upsilon_2(\phi,\psi)(u) - \Upsilon_2(\phi,\psi)(v)| &= \left| \int_u^{+\infty} e^{\int_s^u a_2(r) dr} \frac{c_2(s) \psi(s - \tau_2(s)) \psi(s)}{\phi(s - \sigma_2(s)) + k_2(s)} ds \right. \\ &- \int_v^{+\infty} e^{\int_s^v a_2(r) dr} \frac{c_2(s) \psi(s - \tau_2(s)) \psi(s)}{\phi(s - \sigma_2(s)) + k_2(s)} ds \right| \\ &\leq \left| 1 - e^{\int_u^v a_2(r) dr} \right| \int_u^{+\infty} e^{\int_s^u a_2(r) dr} \frac{c_2(s) \psi(s - \tau_1(s)) \psi(s)}{\phi(s - \sigma_1(s)) + k_2(s)} ds \\ &+ \int_u^v e^{\int_s^v a_2(r) dr} \frac{c_2(s) \psi(s - \tau_1(s)) \psi(s)}{\phi(s - \sigma_1(s)) + k_2(s)} ds \\ &\leq 2 \left(e^{a_2^s(v - u)} - 1 \right) \frac{c_2^s M_1 M_2}{a_2^i k_2^s}. \end{split}$$

Then from 21

$$|\Upsilon_2(\phi,\psi)(u) - \Upsilon_2(\phi,\psi)(v)| \underset{u \to v}{\to} 0.$$

which shows that 1) holds.

For the 2) statement, given any $t \in \mathbb{R}$ and for any $(\phi, \psi) \in M$, we get

$$\Upsilon_1(\phi,\psi)(t) \leq M_1 \text{ And } \Upsilon_2(\phi,\psi)(t) \leq M_2.$$

Pose now $\rho = max(M_1, M_2)$. Therefore $\Upsilon(\phi, \psi)(t) \in B(0, \rho)$ which proves the result.

Denote the closed convex hull of ΥM by $\overline{co} \Upsilon M$. Since $\Upsilon M \subseteq M$ and M is closed convex, $\overline{co} \Upsilon M \subseteq M$. Thus $\Upsilon(\overline{co} \Upsilon M) \subseteq \Upsilon M \subseteq \overline{co} \Upsilon M$. It is easy to verify that $\overline{co} \Upsilon M$ has the properties 1) and 2). More explicitly, $\{\Upsilon(\phi,\psi)(t); (\phi,\psi) \in \overline{co} \Upsilon M\}$ is relatively compact \mathbb{R}^2 for each $t \in \mathbb{R}$, and $\overline{co} \Upsilon M \subseteq BC(\mathbb{R}^2)$ is uniformly bounded and equi-continuous. By the Arzela-Ascoli theorem [10], the restriction of $\overline{co} \Upsilon M$ to every bounded interval I of \mathbb{R} , namely $\{\Upsilon(\phi,\psi)(t); (\phi,\psi) \in \overline{co} \Upsilon M\}_{t \in I}$, is relatively compact in $C(I,\mathbb{R}^2)$. Thus, $\Upsilon: \overline{co} \Upsilon M \to \overline{co} \Upsilon M$ is a compact operator. It follows from Schauder's fixed point theorem that Υ has a fixed point (u^*,v^*) in M. The proof is complete.

5 Stability of the p.a.p solution

Before the stability theorem, we need the following lemma

Lemme 5.1. [11]Let f be a non-negative function defined on $[0; +\infty[$ such that f is integrable on $[0; +\infty[$ and is uniformly continuous on $[0; +\infty[$. Then

$$\lim_{t \to +\infty} f(t) = 0.$$

Definition 5.2. If (u^*, v^*) is a pseudo almost periodic solution of system (2), and (u, v) is an any solution of (2) satisfying $\lim_{t\to +\infty} |u(t)-u^*(t)| = \lim_{t\to +\infty} |v(t)-v^*(t)| = 0$, then we call this solution (u^*, v^*) is globally attractive.

Theorem 5.3. Assume that

$$\liminf \alpha(t), \liminf \beta(t) > 0.$$

where $\alpha(t)$ and $\beta(t)$ are defined in 28 and 29. The pseudo almost periodic solution (u^*, v^*) is globally attractive.

Proof.

Suppose (u, v) is an any solution of the system (2). Define the Lyapunov functional as follows

$$W_1(t) = \left| \ln u(t) - \ln u^*(t) \right| + \left| \ln v(t) - \ln v^*(t) \right|.$$

Pose that $w_1 = u - u^*$, $w_2 = v - v^*$. And define ζ_i^{-1} , ζ_i^{-1} are the inverse functions of $\zeta_i = t - \sigma_i(t)$, $\zeta_i = t - \tau_i(t)$, respectively i = 1, 2.

By calculating Dini derivative of V along system (2), we get

$$D^{+}W_{1}(t) = sgn(w_{1}(t)) \left[\frac{u'(t)}{u(t)} - \frac{u^{*'}(t)}{u^{*}(t)} \right] + sgn(w_{2}(t)) \left[\frac{v'(t)}{v(t)} - \frac{v^{*'}(t)}{v^{*}(t)} \right]$$

$$= sgn(w_{1}(t)) \left[-b_{1}(t)w_{1}(t) - c_{1}(t) \left(\frac{v(\zeta_{1}(t))}{u(\varsigma_{1}(t)) + k_{1}(t)} - \frac{v^{*}(\zeta_{1}(t))}{u^{*}(\varsigma_{1}(t) + k_{1}(t)} \right) \right]$$

$$- sgn(w_{2}(t))c_{2}(t) \left[\frac{v(\zeta_{1}(t))}{u(\varsigma_{2}(t) + k_{2}(t)} - \frac{v^{*}(\zeta_{2}(t))}{u^{*}(\varsigma_{2}(t)) + k_{2}(t)} \right]$$

$$= -b_{1}(t) |w_{1}(t)| - \frac{c_{1}(t)sgn(w_{1}(t))w_{2}(\zeta_{1}(t))}{u(\varsigma_{1}(t)) + k_{1}(t)} + \frac{c_{1}(t)sgn(w_{1}(t))v^{*}(\zeta_{1}(t))w_{1}(\varsigma_{1}(t))}{(u(\varsigma_{1}(t)) + k_{1}(t))(u^{*}(\varsigma_{1}(t)) + k_{1}(t))}$$

$$- \frac{c_{2}(t)sgn(w_{2}(t))w_{2}(\zeta_{2}(t))}{u(\varsigma_{2}(t)) + k_{2}(t)} + \frac{c_{2}(t)sgn(w_{2}(t))v^{*}(\zeta_{2}(t))w_{1}(\varsigma_{2}(t))}{(u(\varsigma_{1}(t)) + k_{1}(t))(u^{*}(\varsigma_{2}(t)) + k_{2}(t))}$$

$$= -b_{1}(t) |w_{1}(t)| - c_{1}(t) \frac{sgn(w_{1}(t))w_{2}(t)}{u(\varsigma_{1}(t)) + k_{1}(t)} + c_{1}(t) \frac{sgn(w_{1}(t))}{u(\varsigma_{1}(t)) + k_{1}(t)} \int_{\zeta_{1}(t)}^{t} w'_{2}(s)ds$$

$$+ \frac{c_{1}(t)v^{*}(\zeta_{1}(t))sgn(w_{1}(t))}{(u(\varsigma_{1}(t)) + k_{1}(t))(u^{*}(\varsigma_{1}(t)) + k_{1}(t))} \left[w_{1}(t) - \int_{\zeta_{1}(t)}^{t} w'_{1}(s)ds \right]$$

$$- c_{2}(t) \frac{sgn(w_{2}(t))}{u(\varsigma_{2}(t)) + k_{2}(t)} \left[w_{2}(t) - \int_{\zeta_{2}(t)}^{t} w'_{2}(s)ds \right]$$

$$+ \frac{c_{2}(t)v^{*}(\zeta_{2}(t))sgn(w_{2}(t))}{(u(\varsigma_{2}(t)) + k_{2}(t))(u^{*}(\varsigma_{2}(t)) + k_{2}(t))} \left[w_{1}(t) - \int_{\zeta_{2}(t)}^{t} w'_{1}(s)ds \right]$$

Integrating both sides of $w'_1(t)$ on the interval $[\varsigma_i(t), t]$ where j=1 or =2, we have

$$\int_{\zeta_{j}(t)}^{t} w_{1}'(s)ds = \int_{\zeta_{j}(t)}^{t} u(s) \left[a_{1}(s) - b_{1}(s)u(s) - \frac{c_{1}(s)v(\zeta_{1}(s))u(s)}{u(\zeta_{1}(s)) + k_{1}(s)} \right]
- u^{*}(s) \left[a_{1}(s) - b_{1}(s)u^{*}(s) - \frac{c_{1}(s)v^{*}(\zeta_{1}(s))u^{*}(s)}{u^{*}(\zeta_{1}(s)) + k_{1}(s)} \right] ds
= \int_{\zeta_{j}(t)}^{t} a_{1}(s)w_{1}(s) - b_{1}(s)(u(s) + u^{*}(s)w_{1}(s) - \frac{c_{1}(t)u(s)}{u(\zeta_{1}(s)) + k_{1}(s)} w_{2}(\zeta_{1}(s))
- \frac{c_{1}(s)v^{*}(\zeta_{1}(s))}{u(\zeta_{1}(s)) + k_{1}(s)} w_{1}(s) + \frac{c_{1}(s)v^{*}(\zeta_{1}(s))u^{*}(s)}{(u(\zeta_{1}(s)) + k_{1}(s))(u^{*}(\zeta_{1}(s)) + k_{1}(s))} w_{1}(\zeta_{1}(s)) ds$$
(23)

and Integrating both sides of $w_2'(t)$ on the interval $[\zeta_i(t), t]$ where j=1 or =2, we have

$$\int_{\zeta_{j}(t)}^{t} w_{2}'(s)ds = \int_{\zeta_{j}(t)}^{t} v(s) \left[a_{2}(s) - \frac{c_{2}(s)v(\zeta_{2}(s))v(s)}{u(\varsigma_{2}(s)) + k_{2}(s)} \right] - v^{*}(s) \left[a_{2}(s) - \frac{c_{2}(s)v^{*}(\zeta_{2}(s))v^{*}(s)}{u^{*}(\varsigma_{2}(s)) + k_{2}(s)} \right] ds$$

$$= \int_{\zeta_{j}(t)}^{t} a_{2}(s)w_{2}(s) - \frac{c_{2}(t)v(s)}{u(\varsigma_{2}(s)) + k_{2}(s)} w_{2}(\zeta_{1}(s))$$

$$- \frac{c_{2}(s)v^{*}(\zeta_{2}(s))}{u(\varsigma_{2}(s)) + k_{2}(s)} w_{2}(s) + \frac{c_{2}(s)v^{*}(\zeta_{2}(s))v^{*}(s)}{(u(\varsigma_{2}(s)) + k_{2}(s))(u^{*}(\varsigma_{2}(s)) + k_{2}(s))} w_{1}(\varsigma_{2}(s)) ds$$
(24)

From theorem 3.4, we have for all solution (u, v) of system (2) $\exists M_1 > m_1 \geq 0, M_2 > m_2 > 0$ such that

$$m_1 \le u(t) \le M_1; \quad m_2 \le v(t) \le M_2 \qquad \forall t \in \mathbb{R}.$$

Therefore, By substituting (23)–(24) into (22), we get

$$D^{+}W_{1}(t) \leq -\left(b_{1}^{i} - \frac{c_{1}^{s}M_{2}}{(m_{1} + k_{1}^{i})^{2}} - \frac{c_{2}^{s}M_{2}}{(m_{1} + k_{2}^{i})^{2}}\right) |w_{1}(t)| - \left(\frac{c_{2}^{i}}{M_{1} + k_{2}^{s}} - \frac{c_{1}^{s}}{m_{1} + k_{1}^{i}}\right) |w_{2}(t)|$$

$$+ \frac{c_{1}^{s}}{m_{1} + k_{1}^{i}} \left[a_{2}^{s} + \frac{c_{2}^{s}M_{2}}{m_{1} + k_{2}^{i}}\right] \int_{\zeta_{1}(t)}^{t} |w_{2}(s)| ds + \frac{c_{1}^{s}c_{2}^{s}M_{2}}{(m_{1} + k_{1}^{i})(m_{1} + k_{2}^{i})} \int_{\zeta_{1}(t)}^{t} |w_{2}(\zeta_{1}(s))| ds$$

$$+ \frac{c_{1}^{s}c_{2}^{s}M_{2}^{2}}{(m_{1} + k_{1}^{i})(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{1}(t)}^{t} |w_{1}(\zeta_{2}(s))| ds + \frac{(c_{1}^{s})^{2}M_{1}M_{2}^{2}}{(m_{1} + k_{1}^{i})^{4}} \int_{\zeta_{1}(t)}^{t} |w_{1}(\zeta_{1}(s))| ds$$

$$+ \frac{(c_{1}^{s})^{2}M_{1}M_{2}}{(m_{1} + k_{1}^{i})^{3}} \int_{\zeta_{1}(t)}^{t} |w_{2}(\zeta_{1}(s))| ds + \left[\frac{c_{1}^{s}a_{1}^{s}M_{2}}{(m_{1} + k_{1}^{i})^{2}} + \frac{2c_{1}^{s}b_{1}^{s}M_{1}M_{2}}{(m_{1} + k_{1}^{i})^{2}} + \frac{(c_{1}^{s})^{2}M_{2}^{2}}{(m_{1} + k_{1}^{i})^{3}}\right] \int_{\zeta_{1}(t)}^{t} |w_{1}(s)| ds$$

$$+ \left[\frac{c_{2}^{s}a_{2}^{s}}{m_{1} + k_{2}^{s}} + \frac{(c_{2}^{s})^{2}M_{2}}{(m_{1} + k_{2}^{i})^{2}}\right] \int_{\zeta_{2}(t)}^{t} |w_{2}(s)| ds + \frac{(c_{2}^{s})^{2}M_{2}}{(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{2}(t)}^{t} |w_{1}(s)| ds$$

$$+ \left[\frac{2c_{2}^{s}b_{1}^{s}M_{1}M_{2}}{(m_{1} + k_{2}^{i})^{2}} + \frac{c_{2}^{s}c_{1}^{s}M_{2}^{2}}{(m_{1} + k_{2}^{i})^{2}} + \frac{c_{2}^{s}a_{1}^{s}M_{2}}{(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{2}(t)}^{t} |w_{1}(s_{1}(s))| ds$$

$$+ \frac{c_{2}^{s}M_{1}M_{2}}{(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{2}(t)}^{t} |w_{2}(\zeta_{1}(s))| ds + \frac{c_{2}^{s}c_{1}^{s}M_{1}M_{2}^{2}}{(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{2}(t)}^{t} |w_{1}(\varsigma_{1}(s))| ds$$

$$+ \frac{c_{2}^{s}M_{1}M_{2}}{(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{2}(t)}^{t} |w_{1}(\varsigma_{2}(s)| ds$$

$$+ \frac{c_{2}^{s}M_{1}M_{2}}{(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{2}(t)}^{t} |w_{1}(\varsigma_{2}(s)| ds$$

$$+ \frac{c_{2}^{s}M_{1}M_{2}}{(m_{1} + k_{2}^{i})^{2}} \int_{\zeta_{2}(t)}^{t} |w_{1}(\varsigma_{2}(s)| ds$$

Now, Let

$$\begin{split} W_2(t) = & \frac{c_1^s}{m_1 + k_1^i} \left[a_2^s + \frac{c_2^s M_2}{m_1 + k_2^i} \right] \int_t^{\zeta_1^{-1}(t)} \int_{\zeta_1(u)}^t \left| w_2(s) \middle| ds du + \frac{c_2^s a_1^s M_2}{(m_1 + k_2^i)^2} \right] \int_t^{\zeta_2^{-1}(t)} \int_{\zeta_2(u)}^t \left| w_1(s) \middle| ds du \right. \\ & + \frac{c_1^s c_2^s M_2^2}{(m_1 + k_1^i)(m_1 + k_2^i)^2} \int_t^{\zeta_1^{-1}(t)} \int_{\zeta_1(u)}^t \left| w_1(\varsigma_2(s)) \middle| ds du + \frac{(c_1^s)^2 M_1 M_2}{(m_1 + k_1^i)^3} \int_t^{\varsigma_1^{-1}(t)} \int_{\zeta_1(u)}^t \left| w_2(\zeta_1(s)) \middle| ds du \right. \\ & + \left[\frac{c_1^s a_1^s M_2}{(m_1 + k_1^i)^2} + \frac{2c_1^s b_1^s M_1 M_2}{(m_1 + k_1^i)^2} + \frac{(c_1^s)^2 M_2^2}{(m_1 + k_1^i)^3} \right] \int_t^{\varsigma_1^{-1}(t)} \int_{\zeta_1(u)}^t \left| w_1(s) \middle| ds du \right. \\ & + \left. \frac{(c_1^s)^2 M_1 M_2^2}{(m_1 + k_1^i)^4} \int_t^{\varsigma_1^{-1}(t)} \int_{\zeta_1(u)}^t \left| w_1(\zeta_1(s)) \middle| ds du + \frac{c_2^s a_2^s}{m_1 + k_2^i} \int_t^{\zeta_2^{-1}(t)} \int_{\zeta_2(u)}^t \left| w_2(s) \middle| ds du \right. \\ & + \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^2} \int_t^{\zeta_2^{-1}(t)} \int_{\zeta_2(u)}^t \left| w_2(\zeta_2(s)) \middle| ds du + \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^2} \int_t^{\zeta_2^{-1}(t)} \int_{\zeta_2(u)}^t \left| w_2(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{(c_2^s)^2 M_2^2}{(m_1 + k_2^i)^3} \int_t^{\zeta_2^{-1}(t)} \int_{\zeta_2(u)}^t \left| w_1(\zeta_2(s)) \middle| ds du + \frac{c_2^s M_1 M_2}{(m_1 + k_2^i)^2(m_1 + k_1^i)} \int_t^{\zeta_2^{-1}(t)} \int_{\zeta_2(u)}^t \left| w_2(\zeta_1(s)) \middle| ds du \right. \\ & + \left. \left[\frac{2c_2^s b_1^s M_1 M_2}{(m_1 + k_2^i)^2} + \frac{c_2^s c_1^s M_2^2}{(m_1 + k_1^i)^2} + \frac{c_1^s c_2^s M_2}{(m_1 + k_1^i)(m_1 + k_2^i)} \int_t^{\zeta_1^{-1}(t)} \int_{\zeta_1(u)}^t \left| w_2(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2(m_1 + k_1^i)^2} \int_t^{\zeta_2(u)} \left| w_1(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2(m_1 + k_1^i)^2} \int_t^{\zeta_2(u)} \left| w_1(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2(m_1 + k_1^i)^2} \int_t^{\zeta_2(u)} \left| w_1(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2(m_1 + k_1^i)^2} \int_t^{\zeta_2(u)} \left| w_1(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2(m_1 + k_1^i)^2} \int_t^{\zeta_2(u)} \left| w_1(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2(m_1 + k_1^i)^2} \int_t^{\zeta_2(u)} \left| w_1(\zeta_1(s)) \middle| ds du \right. \\ & + \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2(m_$$

then its derivative is as follows:

$$\begin{split} W_2'(t) &= \frac{c_1^s}{m_1 + k_1^i} \left[a_2^s + \frac{c_2^s M_2}{m_1 + k_2^i} \right] \left[|w_2(t)| (\zeta_1^{-1}(t) - t) - \int_{\zeta_1(t)}^t |w_2(s)| ds \right] \\ &+ \frac{c_1^s c_2^s M_2}{(m_1 + k_1^i)(m_1 + k_2^i)} \left[|w_2(\zeta_1(t))| (\zeta_1^{-1}(t) - t) - \int_{\zeta_1(t)}^t |w_2(\zeta_1(s))| ds \right] \\ &+ \frac{c_1^s c_2^s M_2^2}{(m_1 + k_1^i)(m_1 + k_2^i)^2} \left[|w_1(\varsigma_2(t))| (\zeta_1^{-1}(t) - t) - \int_{\zeta_1(t)}^t |w_1(\varsigma_2(s))| ds \right] \\ &+ \frac{(c_1^s)^2 M_1 M_2}{(m_1 + k_1^i)^3} \left[|w_2(\zeta_1(t))| (\zeta_1^{-1}(t) - t) - \int_{\varsigma_1(t)}^t |w_2(\zeta_1(s))| ds \right] \\ &+ \left[\frac{c_1^s a_1^s M_2}{(m_1 + k_1^i)^2} + \frac{2c_1^s b_1^s M_1 M_2}{(m_1 + k_1^i)^2} + \frac{(c_1^s)^2 M_2^2}{(m_1 + k_1^i)^3} \right] \left[|w_1(t)| (\varsigma_1^{-1}(t) - t) - \int_{\varsigma_1(t)}^t |w_1(s)| ds \right] \\ &+ \frac{(c_1^s)^2 M_1 M_2^2}{(m_1 + k_1^i)^4} \left[|w_1(\varsigma_1(t))| (\varsigma_1^{-1}(t) - t) - \int_{\varsigma_1(t)}^t |w_1(\varsigma_1(s))| ds \right] \\ &+ \left[\frac{c_2 a_2^s}{m_1 + k_2^i} + \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^2} \right] \left[|w_2(t)| (\zeta_2^{-1}(t) - t) - \int_{\zeta_2(t)}^t |w_2(s)| ds \right] \\ &+ \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^3} \left[|w_1(\varsigma_2(t))| (\zeta_2^{-1}(t) - t) - \int_{\zeta_2(t)}^t |w_1(\varsigma_2(s))| ds \right] \\ &+ \frac{c_2^s M_1 M_2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \left[|w_2(\zeta_1(t))| (\varsigma_2^{-1}(t) - t) - \int_{\varsigma_2(t)}^t |w_2(\zeta_1(s))| ds \right] \\ &+ \left[\frac{2c_2^s b_1^s M_1 M_2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \left[|w_1(\varsigma_1(t))| (\varsigma_2^{-1}(t) - t) - \int_{\varsigma_2(t)}^t |w_1(\varsigma_1(s))| ds \right] \\ &+ \frac{c_2^s c_1^s M_1 M_2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \left[|w_1(\varsigma_1(t))| (\varsigma_2^{-1}(t) - t) - \int_{\varsigma_2(t)}^t |w_1(\varsigma_1(s))| ds \right] \\ &+ \frac{c_2^s c_1^s M_1 M_2^2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \left[|w_1(\varsigma_1(t))| (\varsigma_2^{-1}(t) - t) - \int_{\varsigma_2(t)}^t |w_1(\varsigma_1(s))| ds \right] \\ &+ \frac{c_2^s c_1^s M_1 M_2^2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \left[|w_1(\varsigma_1(t))| (\varsigma_2^{-1}(t) - t) - \int_{\varsigma_2(t)}^t |w_1(\varsigma_1(s))| ds \right] \\ &+ \frac{c_2^s c_1^s M_1 M_2^2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \left[|w_1(\varsigma_1(t))| (\varsigma_2^{-1}(t) - t) - \int_{\varsigma_2(t)}^t |w_1(\varsigma_1(s))| ds \right] \\ \end{aligned}$$

Define the Lyapunov functional by $W(t) = W_1(t) + W_2(t)$, then

$$D^{+}W(t) = D^{+}W_{1}(t) + W_{2}'(t). (27)$$

Substitution of (25)–(26) into (27) gives:

$$\begin{split} D^+W(t) & \leq -\left[b_1^i - \frac{c_1^s M_2}{(m_1 + k_1^i)^2} - \frac{c_2^s M_2}{(m_1 + k_2^i)^2} - \frac{c_1^s c_2^s M_2^2}{(m_1 + k_1^i)(m_1 + k_2^i)^2} (\zeta_1^{-1}(t) - t) \right. \\ & - \left(\frac{c_1^s a_1^s M_2}{(m_1 + k_1^i)^2} + \frac{2c_1^s b_1^s M_1 M_2}{(m_1 + k_1^i)^2} + \frac{(c_1^s)^2 M_2^2}{(m_1 + k_1^i)^3} + \frac{(c_1^s)^2 M_1 M_2^2}{(m_1 + k_1^i)^4} \right) (\varsigma_1^{-1}(t) - t) \\ & - \left(\frac{(c_2^s)^2 M_2^2}{(m_1 + k_2^i)^3} + \frac{2c_2^s b_1^s M_1 M_2}{(m_1 + k_2^i)^2} + \frac{c_2^s c_1^s M_2^2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} + \frac{c_2^s a_1^s M_2}{(m_1 + k_2^i)^2} \right) (\varsigma_2^{-1}(t) - t) \right] |w_1(t)| \\ & - \left[\frac{c_2^i}{M_1 + k_2^s} - \frac{c_1^s}{m_1 + k_1^i} - \frac{c_1^s}{m_1 + k_1^i} \left(a_2^s + \frac{c_2^s M_2}{m_1 + k_2^i} + \frac{c_1^s c_2^s M_2}{(m_1 + k_1^i)(u^i + k_2^i)} \right) (\zeta_1^{-1}(t) - t) \right. \\ & - \frac{(c_1^s)^2 M_1 M_2}{(m_1 + k_1^i)^3} (\varsigma_1^{-1}(t) - t) - \left(\frac{c_2^s a_2^s}{m_1 + k_2^i} + \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^2} + \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^2} \right) (\zeta_2^{-1}(t) - t) \\ & - \left(\frac{c_2^s M_1 M_2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)} + \frac{c_2^s c_1^s M_1 M_2^2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \right) (\varsigma_2^{-1}(t) - t) \right] |w_2(t)| \end{split}$$

Let's denote

$$\alpha(t) = b_1^i - \frac{c_1^s M_2}{(m_1 + k_1^i)^2} - \frac{c_2^s M_2}{(m_1 + k_2^i)^2} - \frac{c_1^s c_2^s M_2^2}{(m_1 + k_1^i)(m_1 + k_2^i)^2} (\zeta_1^{-1}(t) - t)$$

$$- \left(\frac{c_1^s a_1^s M_2}{(m_1 + k_1^i)^2} + \frac{2c_1^s b_1^s M_1 M_2}{(m_1 + k_1^i)^2} + \frac{(c_1^s)^2 M_2^2}{(m_1 + k_1^i)^3} + \frac{(c_1^s)^2 M_1 M_2^2}{(m_1 + k_1^i)^4} \right) (\zeta_1^{-1}(t) - t)$$

$$- \left(\frac{(c_2^s)^2 M_2^2}{(m_1 + k_2^i)^3} + \frac{2c_2^s b_1^s M_1 M_2}{(m_1 + k_2^i)^2} + \frac{c_2^s c_1^s M_2^2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} + \frac{c_2^s a_1^s M_2}{(m_1 + k_2^i)^2} \right) (\zeta_2^{-1}(t) - t)$$

$$(28)$$

And

$$\beta(t) = \frac{c_2^i}{M_2 + k_2^s} - \frac{c_1^s}{m_1 + k_1^i} - \frac{c_1^s}{m_1 + k_1^i} \left(a_2^s + \frac{c_2^s M_2}{m_1 + k_2^i} + \frac{c_1^s c_2^s M_2}{(m_1 + k_1^i)(m_1 + k_2^i)} \right) (\zeta_1^{-1}(t) - t)$$

$$- \frac{(c_1^s)^2 M_1 M_2}{(m_1 + k_1^i)^3} (\zeta_1^{-1}(t) - t) - \left(\frac{c_2^s a_2^s}{m_1 + k_2^i} + \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^2} + \frac{(c_2^s)^2 M_2}{(m_1 + k_2^i)^2} \right) (\zeta_2^{-1}(t) - t)$$

$$- \left(\frac{c_2^s M_1 M_2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)} + \frac{c_2^s c_1^s M_1 M_2^2}{(m_1 + k_2^i)^2 (m_1 + k_1^i)^2} \right) (\zeta_2^{-1}(t) - t)$$

$$(29)$$

From hypothesis of the theorem, ones have $\alpha^i = \liminf \alpha(t)$ and $\beta^i = \liminf \beta(t)$ verified for sufficiently large T, we obtained

$$D^+W(t) \le -\alpha^i |w_1(t)| - \beta^i |w_2(t)| < 0.$$

which implies W(t) is non-increasing on $[T, +\infty[$. An integration of above inequality from T to t yields

$$W(t) + \alpha^{i} \int_{t}^{T} |u(s) - u^{*}(s)| ds + \beta^{i} \int_{t}^{T} |v(s) - v^{*}(s)| ds \le W(T) < +\infty, \quad \forall t > T.$$

Then

$$\limsup_{t\to +\infty} \int_t^T |u(s)-u^*(s)| ds \leq \frac{W(T)}{\alpha^i} < +\infty \text{ and } \limsup_{t\to +\infty} \int_t^T |v(s)-v^*(s)| ds \leq \frac{W(T)}{\beta^i} < +\infty.$$

Thus we have

$$\lim_{t \to +\infty} \left| u(t) - u^*(t) \right| = \lim_{t \to +\infty} \left| v(t) - v^*(t) \right| = 0.$$

6 Example and Stimulation

In order to illustrate some feature of our main results, in this section, we will apply our main results to some special prey-predator systems and demonstrate the efficiencies of our criteria.

6.1 Example 1:

In this example, we consider a system without the condition (C0). Then, $m_1 = 0$. The system is considered


$$\begin{cases} u'(t) = \left(0.04 + 0.125|\cos(\sqrt{2}t)| + 0.125exp(-t) - (2.6 + 0.5\cos(t))u(t) - \frac{3.2v(t - 0.75)}{u(t - 0.75) + 17}\right)u(t); \\ v'(t) = \left(0.01 + 0.25|\sin(\sqrt{7}t)| - \frac{3.5v(t - 0.75)}{u(t - 0.75) + 3.4}\right)v(t), \end{cases}$$
(30)


By a direct calculation, ones have the following table

Table 1: The ecological parameters of u and v.

j	a_j^i	a_j^s	b^i	b^s	c^i_j	c_j^s	k_j^i	k_j^s	σ_j^s	$ au_j^s$
1	0.04	0.29	2.6	3.1	3.2	3.2	17	17	0.75	0.75
2	0.01	0.26	_	_	3.5	3.5	3.4	3.4	0.75	0.75

And $\alpha^i \simeq 2.4$; $\beta^i \simeq 0.012$; $M_1 = 0.7085$; $m_1 = 0$; $M_2 = 0.6506$; and $m_2 = 0.0829$. The theorem (4.11) is verified and the conditions of theorem (5.3) are satisfied. Therefore, there exist at least a pseudo almost periodic which is globally attractive.





6.2 Example 2:

In this example, the condition (C0) holds. Let us consider


$$\begin{cases} u'(t) = \left(4.8 + 0.125(|\cos(\sqrt{2}t)| + |\cos(\sqrt{2}t)|\right) - \left(0.25|\cos(t)| + \frac{33.72 + 32.72t^2}{4 + 4t^2}\right)u(t) - \frac{0.32v(t - 0.92)}{u(t - 0.92) + 16.7}\right)u(t); \\ v'(t) = \left(0.03 + 0.125(|\sin(\sqrt{2}t)| + |\cos(\sqrt{5}t)|) - \frac{3.6v(t - 0.92)}{u(t - 0.92) + 5.7}\right)v(t), \end{cases}$$
(31)


By a direct calculation, ones have the following table

Table 2: The ecological parameters of u and v.

j	a_j^i	a_j^s	b^i	b^s	c_j^i	c_j^s	k_j^i	k_j^s	σ_j^s	$ au_j^s$
1	4.8	5.05	8.1	8.6	0.32	0.32	16.7	16.7	0.92	0.92
2	0.03	0.28	_	_	3.6	3.6	5.7	5.7	0.92	0.92

And $\alpha^i \simeq 7.25$; $\beta^i \simeq 0.009$; $M_1 = 0.6226$; $m_1 = 0.5567$; $M_2 = 0.6403$; and $m_2 = 0.0408$. The theorem (4.11) is verified and the conditions of theorem (5.3) are satisfied. Therefore, there exist at least a pseudo almost periodic which is globally attractive.





7 Conclusion

The aim of this paper is to prove the existence of positive almost periodic solution in a Leslie-Gower predator-prey model with continuous delays. Based on new conditions, the global attractivity of the above model is obtained by building a suitable Lyapunov functional. Moreover, some numerical examples show that the our theoretical results are effective

References

- [1] M. Amdouni and F. Chérif. The pseudo almost periodic solutions of the new class of lotka–volterra recurrent neural networks with mixed delays. Chaos, Solitons & Fractals, 113:79–88, 2018.
- [2] M. Aziz-Alaoui. Study of a leslie–gower-type tritrophic population model. Chaos, Solitons & Fractals, 14(8):1275–1293, 2002.
- [3] M. Aziz-Alaoui and M. D. Okiye. Boundedness and global stability for a predator-prey model with modified leslie-gower and holling-type II schemes. Applied Mathematics Letters, 16(7):1069–1075, 2003.
- [4] D. Békollè, K. Ezzinbi, S. Fatajou, D. E. H. Danga, and F. M. Béssémè. Attractiveness of pseudo almost periodic solutions for delayed cellular neural networks in the context of measure theory. Neurocomputing, 435:253–263, 2021.
- [5] F. Chen, Z. Li, and Y. Huang. Note on the permanence of a competitive system with infinite delay and feedback controls. Nonlinear Analysis: Real World Applications, 8(2):680–687, 2007.
- [6] X. Chen and H. Ding. Positive pseudo almost periodic solutions for a hematopoiesis model. J Nonlinear Evol Equ Appl ISSN, 2016(2):25–36, 2016.
- [7] Z. Chuanyi. Almost periodic type functions and ergodicity. Springer Science & Business Media, 2003.
- [8] P. Cieutat, S. Fatajou, and G. M. N'Guérékata. Composition of pseudo almost periodic and pseudo almost automorphic functions and applications to evolution equations. Applicable Analysis, 89(1):11–27, 2010.
- [9] W. A. Coppel. Dichotomies in stability theory, volume 629. Springer, 2006.
- [10] T. Diagana. Almost automorphic type and almost periodic type functions in abstract spaces. Springer, 2013.
- [11] K. Gopalsamy. Stability and osilation in delay differential equation in delay differential equations of population dynamics, 1992.
- [12] Y. Kuang and H. Smith. Global stability for infinite delay lotka-volterra type systems. Journal of differential equations, 103(2):221–246, 1993.
- [13] P. Leslie and J. Gower. The properties of a stochastic model for the predator-prey type of interaction between two species. Biometrika, 47(3/4):219–234, 1960.
- [14] P. H. Leslie. Some further notes on the use of matrices in population mathematics. Biometrika, 35(3/4):213–245, 1948.
- [15] Q. Liao. Almost periodic solution for a lotka-volterra predator-prey system with feedback controls on time scales. In 2018 6th International Conference on Machinery, Materials and Computing Technology (ICMMCT 2018), pages 46–51. Atlantis Press, 2018.
- [16] H. Lu and G. Yu. Permanence of a gilpin-ayala predator-prey system with time-dependent delay. Advances in Difference Equations, 2015(1):1–15, 2015.
- [17] N. MacDonald and N. MacDonald. Biological delay systems: linear stability theory. Cambridge University Press, 2008.
- [18] M. Menouer, A. Moussaoui, and E. A. Dads. Existence and global asymptotic stability of positive almost periodic solution for a predator-prey system in an artificial lake. Chaos, Solitons & Fractals, 103:271–278, 2017.
- [19] D. Smart. Fixed point theorems, cambridge uni. Press., Cambridge, 1980.
- [20] K. Steinmuller. Pielou, ec: Mathematical ecology. john wiley & sons, new york-london-sydneytoronto 1977. x, 385 s.,£13.00; 22.45, 1978.

- [21] J. P. Tripathi. Almost periodic solution and global attractivity for a density dependent predatorprey system with mutual interference and crowley–martin response function. Differential Equations and Dynamical Systems, 28(1):19–37, 2020.
- [22] C. Xu, P. Li, and Y. Guo. Global asymptotical stability of almost periodic solutions for a nonautonomous competing model with time-varying delays and feedback controls. Journal of biological dynamics, 13(1):407–421, 2019.
- [23] C. Y. Zhang. Pseudo almost periodic solutions of some differential equations. Journal of Mathematical Analysis and Applications, 181(1):62–76, 1994.