

Proving dualities for q MZVs with connected sums

Benjamin Brindle *

November 2, 2021

ABSTRACT. This paper gives a new application of so-called connected sums, introduced recently by Seki and Yamamoto [SY]. Special about our approach is that it proves a duality for the Schlesinger–Zudilin and the Bradley–Zhao model of q MZVs simultaneously. The latter implies the duality for MZVs and the former can be used to prove the shuffle product formula for MZVs. Furthermore, the q -Ohno relations, a generalization of Bradley–Zhao duality, is also obtained.

2010 *Mathematics Subject Classification*. 11M32, 05A30.

Key words: Multiple zeta values; q -multiple zeta values; duality; connected sums.

1 Notation and definitions

For an *admissible index* $\mathbf{k} = (k_1, \dots, k_r) \in \mathbb{N}^r$, i.e., $r \geq 0$ and $k_1 \geq 2$, its *multiple zeta value* (MZV) is

$$\zeta(\mathbf{k}) := \sum_{m_1 > \dots > m_r > 0} \frac{1}{m_1^{k_1}} \cdots \frac{1}{m_r^{k_r}}.$$

To understand the algebraic structure of MZVs better on the one hand and to get connections to holomorphic functions, in particular, modular forms (see [GKZ], [Bac]), on the other hand, it is useful to introduce q -analogs of MZVs. There are several models of q -analogs. We focus in this paper on the one by Schlesinger, and Zudilin and Bradley and Zhao. For further details on these and other models, we refer to [Zha], [Bri]. In this note q will be a formal variable or a real number with $0 < q < 1$.

The Bradley–Zhao model is defined as follows: Set $\zeta_q^{\text{BZ}}(\emptyset) := 1$ and for $\mathbf{k} = (k_1, \dots, k_r)$ an admissible index define

$$\zeta_q^{\text{BZ}}(\mathbf{k}) := \sum_{m_1 > \dots > m_r > 0} \frac{q^{m_1(k_1-1)}}{(1-q^{m_1})^{k_1}} \cdots \frac{q^{m_r(k_r-1)}}{(1-q^{m_r})^{k_r}}.$$

Similar, but different, we define Schlesinger–Zudilin q MZVs via $\zeta_q^{\text{SZ}}(\emptyset) := 1$ and for every *SZ-admissible index* \mathbf{k} , i.e., $\mathbf{k} \in \mathbb{N}_0^r$ for some $r \geq 0$ with $k_1 \geq 1$, we set

$$\zeta_q^{\text{SZ}}(\mathbf{k}) := \sum_{m_1 > \dots > m_r > 0} \frac{q^{m_1 k_1}}{(1-q^{m_1})^{k_1}} \cdots \frac{q^{m_r k_r}}{(1-q^{m_r})^{k_r}}.$$

*The author has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 101001179).

2 Dualities

Write an admissible index \mathbf{k} in the shape $\mathbf{k} = (k_1 + 1, \{1\}^{d_1-1}, \dots, k_r + 1, \{1\}^{d_r-1})$ with $k_j, d_j \geq 1$ unique ($\{1\}^d$ means that 1 is repeated d -times). For the next two theorems, we need the *dual index*,

$$\mathbf{k}^\vee := \left(d_r + 1, \{1\}^{k_r-1}, \dots, d_1 + 1, \{1\}^{k_1-1} \right).$$

Theorem 1 (MZV-Duality, [Zag, §9]). *For every admissible index \mathbf{k} , we have $\zeta(\mathbf{k}) = \zeta(\mathbf{k}^\vee)$.*

The next theorem can be seen as a q -analogue of MZV-duality since MZV-duality follows immediately from it (cf. the proof):

Theorem 2 (BZ-Duality, [Bra, Thm. 5]). *For every admissible index \mathbf{k} , we have $\zeta_q^{\text{BZ}}(\mathbf{k}) = \zeta_q^{\text{BZ}}(\mathbf{k}^\vee)$.*

A generalization of BZ-duality are the so-called q -Ohno relations, of which BZ-duality is the special case $c = 0$:

Theorem 3 (q -Ohno relation, [OT, Thm. 1]). *For any admissible index $\mathbf{k} = (k_1, \dots, k_r)$ and any $c \in \mathbb{N}_0$ we have*

$$\sum_{|\mathbf{c}|=c} \zeta_q^{\text{BZ}}(\mathbf{k} + \mathbf{c}) = \sum_{|\mathbf{c}|=c} \zeta_q^{\text{BZ}}(\mathbf{k}^\vee + \mathbf{c}),$$

where we sum over all $\mathbf{c} = (c_1, \dots, c_r) \in \mathbb{N}_0^r$ with $|\mathbf{c}| := c_1 + \dots + c_r = c$.

For the SZ-model, we write an SZ-admissible index in the shape $\mathbf{k} = (k_1 + 1, \{0\}^{d_1}, \dots, k_r + 1, \{0\}^{d_r})$ with $k_j, d_j \geq 0$ unique and define the *SZ-dual index*,

$$\mathbf{k}^\dagger := \left(d_r + 1, \{0\}^{k_r}, \dots, d_1 + 1, \{0\}^{k_1} \right).$$

Theorem 4 (SZ-Duality, [Zha, Thm. 8.3]). *For \mathbf{k} SZ-admissible, we have $\zeta_q^{\text{SZ}}(\mathbf{k}) = \zeta_q^{\text{SZ}}(\mathbf{k}^\dagger)$.*

Note that BZ- and SZ-duality on algebraic level look the same, both can be obtained by the same anti-automorphism on the non-commutative free algebra in two variables (see, e.g., [Bri, Thm. 3.5, Thm. 3.16]). But they imply different things. BZ-duality gives direct duality for MZVs, while SZ-duality does not. However, SZ-duality implies another important result in the theory of MZVs, namely the shuffle product formula (cf. [EMS], [Sin], for details [Bri, Thm. 3.46]).

For some calculations in the next section we need the connection between admissible and SZ-admissible index: An index \mathbf{k} is admissible if and only if $\mathbf{k} - \mathbf{1}$ is SZ-admissible ($\mathbf{k} + \mathbf{1}$ is the index, which is \mathbf{k} with every entry increased by 1; similar for $\mathbf{k} - \mathbf{1}$). Furthermore, we have for \mathbf{k} admissible

$$(\mathbf{k} - \mathbf{1})^\dagger = \mathbf{k}^\vee - \mathbf{1}. \tag{2.1}$$

3 Connected sums & proof of dualities

As a new tool for proving identities among (q) -multiple zeta values, Seki and Yamamoto introduced the concept of so-called connected sums (this notion is independent of connected sums in topology). With connected sums, they have proven, e.g., the duality of MZVs, Hoffman's identity, and the q -analogue of Ohno's relation, cf. [Sek] or [SY].

Using connected sums, we give a new proof of the duality of Schlesinger–Zudilin q MZVs, the duality of Bradley–Zhao q MZVs and the usual duality of MZVs. It turns out that the connected sum

defined below has the power to prove all three statements at once. As a by-product, we also get a proof for the q -Ohno relation. The proof is inspired by the one of Seki and Yamamoto ([SY]), where the authors proved q -Ohno's relation for non-modified Bradley–Zhao q MZVs. We work with modified q MZVs, which will be here the reason that we can prove also Schlesinger–Zudilin duality at the same time.

Definition 5 (Connected sum). Let be $r, s \geq 0$, $\mathbf{k} = (k_1, \dots, k_r) \in \mathbb{N}_0^r$, $\boldsymbol{\ell} = (\ell_1, \dots, \ell_s) \in \mathbb{N}_0^s$ and x real with $|x| < 1$. Define the connected sum as

$$Z_q(\mathbf{k}; \boldsymbol{\ell}; x) := \sum_{\substack{m_1 > \dots > m_r > m_{r+1} = 0 \\ n_1 > \dots > n_s > n_{s+1} = 0}} \prod_{i=1}^r \frac{q^{m_i k_i}}{(1 - q^{m_i x})(1 - q^{m_i})^{k_i}} \prod_{j=1}^s \frac{q^{n_j \ell_j}}{(1 - q^{n_j x})(1 - q^{n_j})^{\ell_j}} \\ \times \frac{q^{m_1 n_1} f_q(m_1; x) f_q(n_1; x)}{f_q(m_1 + n_1; x)},$$

where $f_q(m; x) := \prod_{h=1}^m (1 - q^h x)$.

Remark 6.

- (i) The connected sum Z_q is symmetric in \mathbf{k} and $\boldsymbol{\ell}$ by definition.
- (ii) Notice that the connected sum is well-defined in the sense that it is a series over positive real numbers and hence either a positive real number (if convergent) or $+\infty$ (if not convergent).
- (iii) If $k_1 \geq 1$, then $Z_q(\mathbf{k}; \emptyset; 0) = \zeta_q^{\text{SZ}}(\mathbf{k})$.
- (iv) If $k_1 \geq 1$, then $\lim_{x \rightarrow 1} Z_q(\mathbf{k}; \emptyset; x) = \zeta_q^{\text{BZ}}(\mathbf{k} + \mathbf{1})$.

Proposition 7 (Boundary conditions). *If $k_1 \geq 1$, $0 < q < 1$ and $x \in \mathbb{R}$ with $|x| < 1$, then $Z_q(\mathbf{k}; \emptyset; x)$ is a well-defined real number.*

Proof. One has

$$Z_q(\mathbf{k}; \emptyset; x) = \sum_{m_1 > \dots > m_{r+1} := 0} \prod_{i=1}^r \frac{q^{m_i k_i}}{(1 - q^{m_i x})(1 - q^{m_i})^{k_i}} \\ \leq \frac{1}{(1 - qx)^r} \sum_{m_1 > \dots > m_{r+1} := 0} \prod_{i=1}^r \frac{q^{m_i k_i}}{(1 - q^{m_i})^{k_i}} = \frac{1}{(1 - qx)^r} \zeta_q^{\text{SZ}}(k_1, \dots, k_r),$$

which is well-defined since $k_1 \geq 1$, i.e., (k_1, \dots, k_r) SZ-admissible. \square

After we have checked well-definedness of Z_q , we state and prove now distinguished relations among our connected sums.

Theorem 8 (Transport relations). *Let be $r, s \geq 0$ and $k_1, \dots, k_r, \ell_1, \dots, \ell_s \geq 0$. If $s > 0$,*

$$Z_q((0, k_1, \dots, k_r); (\ell_1, \dots, \ell_s); x) = Z_q((k_1, \dots, k_r); (\ell_1 + 1, \ell_2, \dots, \ell_s); x) \quad (3.1)$$

and if $r > 0$,

$$Z_q((k_1 + 1, k_2, \dots, k_r); (\ell_1, \dots, \ell_s); x) = Z_q((k_1, \dots, k_r); (0, \ell_1, \ell_2, \dots, \ell_s); x). \quad (3.2)$$

Proof. The second equality follows from the first by symmetry and the first one is obtained from

$$\begin{aligned} & \sum_{a>m} \frac{1}{1-q^a x} \frac{q^{an} f_q(a; x) f_q(n; x)}{f_q(a+n; x)} \\ &= \frac{q^n}{1-q^n} \sum_{a>m} \left(\frac{q^{(a-1)n} f_q(a-1; x) f_q(n; x)}{f_q(a+n-1; x)} - \frac{q^{an} f_q(a; x) f_q(n; x)}{f_q(a+n; x)} \right) \\ &= \frac{q^n}{1-q^n} \frac{q^{mn} f_q(m; x) f_q(n; x)}{f_q(m+n; x)} \end{aligned}$$

and setting $m = m_1$, $n = n_1$, $a = m_0$. \square

This theorem is the key of proving Theorems 1–4. Especially, the following corollary will be needed, together with the connection of Z_q with ζ_q^{BZ} resp. ζ_q^{SZ} (Remark 6).

Corollary 9. *For every SZ-admissible index \mathbf{k} and real x with $|x| < 1$ we have*

$$Z_q(\mathbf{k}; \emptyset; x) = Z_q(\emptyset; \mathbf{k}^\dagger; x).$$

Proof. For all indices \mathbf{k} and ℓ and $k \geq 1$, $d \geq 0$ we obtain (by $(k, \{0\}^d, \mathbf{k})$ we mean the concatenation of the indices $(k, \{0\}^d)$ and \mathbf{k}) by applying (3.2) first and then $(d+1)$ -times (3.1)

$$Z_q\left(\left(k, \{0\}^d, \mathbf{k}\right); \ell; x\right) = Z_q\left(\left(\{0\}^{d+1}, \mathbf{k}\right); \left(\{0\}^k, \ell\right); x\right) = Z_q\left(\mathbf{k}, \left(d+1, \{0\}^{k-1}; \ell\right); x\right). \quad (3.3)$$

Now, set $\ell = \emptyset$ and write an SZ-admissible index \mathbf{k} in the form

$$\mathbf{k} = \left(k_1, \{0\}^{d_1}, \dots, k_r, \{0\}^{d_r}\right).$$

Then we obtain the corollary by induction on r and using (3.3) in the induction step. \square

With the connection of Z_q and ζ_q^{SZ} (Rem. 6 (iii)), SZ-duality follows directly:

Proof of Theorem 4. Take some SZ-admissible index \mathbf{k} . Using the symmetry of Z_q and setting $x = 0$, the claim follows by Corollary 9:

$$\zeta_q^{\text{SZ}}(\mathbf{k}) = Z_q(\mathbf{k}; \emptyset; 0) = Z_q(\emptyset; \mathbf{k}^\dagger; 0) = Z_q(\mathbf{k}^\dagger; \emptyset; 0) = \zeta_q^{\text{SZ}}(\mathbf{k}^\dagger). \quad \square$$

Analogously, we are able to prove BZ-duality:

Proof of Theorem 2. For an admissible index \mathbf{k} we have, using Remark 6 and Corollary 9,

$$\begin{aligned} \zeta_q^{\text{BZ}}(\mathbf{k}) &= \lim_{x \rightarrow 1} Z_q(\mathbf{k} - \mathbf{1}; \emptyset; x) = \lim_{x \rightarrow 1} Z_q(\emptyset; (\mathbf{k} - \mathbf{1})^\dagger; x) \\ &= \lim_{x \rightarrow 1} Z_q((\mathbf{k} - \mathbf{1})^\dagger; \emptyset; x) = \zeta_q^{\text{BZ}}((\mathbf{k} - \mathbf{1})^\dagger + \mathbf{1}) = \zeta_q^{\text{BZ}}(\mathbf{k}^\vee). \quad \square \end{aligned}$$

Example 10. We give a concrete example of applying transport relations step by step to make clear what happens:

$$\begin{aligned} Z_q((1, 0); \emptyset; x) &= Z_q((0, 0); (0); x) \\ &= Z_q((0); (1); x) = Z_q(\emptyset; (2); x) \end{aligned}$$

By Remark 6 (iii) resp. (iv), we obtain $\zeta_q^{\text{SZ}}(1, 0) = \zeta_q^{\text{SZ}}(2)$ resp. $\zeta_q^{\text{BZ}}(2, 1) = \zeta_q^{\text{BZ}}(3)$. We have $(1, 0)^\dagger = (2)$ and $(2, 1)^\vee = (3)$, why these results indeed correspond to SZ- resp. BZ-duality.

We derive in the following the proof of MZV-duality, Theorem 1, from BZ-duality:

Proof of Theorem 1. Let \mathbf{k} be any admissible index. Denote by $\text{wt}(\mathbf{k}) := k_1 + \dots + k_r$ the sum of all entries, the *weight* of \mathbf{k} . Obviously, one has $\text{wt}(\mathbf{k}) = \text{wt}(\mathbf{k}^\vee)$. We have

$$\zeta(\mathbf{k}) = \lim_{q \rightarrow 1} (1-q)^{\text{wt}(\mathbf{k})} \zeta_q^{\text{BZ}}(\mathbf{k}) = \lim_{q \rightarrow 1} (1-q)^{\text{wt}(\mathbf{k}^\vee)} \zeta_q^{\text{BZ}}(\mathbf{k}^\vee) = \zeta(\mathbf{k}^\vee). \quad \square$$

Consider for the proof of Theorem 3 connected sums of the form $Z_q(\mathbf{k}; \emptyset; x)$ and the related one of the form $Z_q(\emptyset; \ell; x)$ using transport relations. In both, we will develop all occurring terms as a Taylor series at $x = 1$, mainly we use that for all $m \in \mathbb{N}$, we have

$$\frac{1}{1-q^m x} = \frac{1}{1-q^m} \frac{1}{1 - \frac{q^m}{1-q^m}(x-1)} = \frac{1}{1-q^m} \sum_{c \geq 0} \left(\frac{q^m}{1-q^m} \right)^c (x-1)^c = \sum_{c \geq 0} \frac{q^{mc}}{(1-q^m)^{c+1}} (x-1)^c.$$

Proof of Theorem 3. Let $\mathbf{k} = (k_1, \dots, k_r)$ be an admissible index. Then we have

$$\begin{aligned} Z_q(\mathbf{k} - \mathbf{1}; \emptyset; x) &= \sum_{m_1 > \dots > m_r > 0} \prod_{j=1}^r \frac{1}{1-q^{m_j} x} \frac{q^{m_j(k_j-1)}}{(1-q^{m_j})^{k_j-1}} \\ &= \sum_{m_1 > \dots > m_r > 0} \prod_{j=1}^r \left(\sum_{c_j \geq 0} \frac{q^{m_j c_j}}{(1-q^{m_j})^{c_j+1}} (x-1)^{c_j} \frac{q^{m_j(k_j-1)}}{(1-q^{m_j})^{k_j-1}} \right) \\ &= \sum_{c_1, \dots, c_r \geq 0} \sum_{m_1 > \dots > m_r > 0} \left(\prod_{j=1}^r \frac{q^{m_j(k_j+c_j-1)}}{(1-q^{m_j})^{k_j+c_j}} \right) (x-1)^{c_1+\dots+c_r} \\ &= \sum_{c_1, \dots, c_r \geq 0} \zeta_q^{\text{BZ}}(\mathbf{k} + \mathbf{c}) (x-1)^{|\mathbf{c}|}. \end{aligned}$$

Since \mathbf{k} was an arbitrary admissible index and \mathbf{k}^\vee is admissible too, we get

$$Z_q(\emptyset; \mathbf{k}^\vee - \mathbf{1}; x) = \sum_{c_1, \dots, c_r \geq 0} \zeta_q^{\text{BZ}}(\mathbf{k}^\vee + \mathbf{c}) (x-1)^{|\mathbf{c}|}.$$

Now, since $Z_q(\mathbf{k} - \mathbf{1}; \emptyset; x) = Z_q(\emptyset; \mathbf{k}^\vee - \mathbf{1}; x)$ for every admissible index by using the transport relations, the result follows by comparing the coefficient of $(x-1)^c$ on both sides. \square

In the same way, we can consider $Z_q(\mathbf{k} - \mathbf{1}; \emptyset; x)$ when developing $\frac{1}{1-q^m x}$ around some $a \in \mathbb{R}$, i.e.,

$$\frac{1}{1-q^m x} = \frac{1}{1-aq^m - q^m(x-a)} = \frac{1}{1-aq^m} \frac{1}{1 - \frac{q^m}{1-aq^m}(x-a)} = \sum_{c \geq 0} \frac{q^{mc}}{(1-aq^m)^{c+1}} (x-a)^c.$$

Then it is

$$\begin{aligned} Z_q(\mathbf{k}; \emptyset; x) &= \sum_{m_1 > \dots > m_r > 0} \prod_{j=1}^r \frac{1}{1-q^{m_j} x} \frac{q^{m_j k_j}}{(1-q^{m_j})^{k_j}} \\ &= \sum_{m_1 > \dots > m_r > 0} \prod_{j=1}^r \sum_{c_j \geq 0} \frac{q^{m_j c_j}}{(1-aq^{m_j})^{c_j+1}} \frac{q^{m_j k_j}}{(1-q^{m_j})^{k_j}} (x-a)^{c_j}. \end{aligned}$$

Remark 11. The series

$$\sum_{m_1 > \dots > m_r > 0} \prod_{j=1}^r \frac{q^{m_j c_j}}{(1 - aq^{m_j})^{c_j+1}} \frac{q^{m_j k_j}}{(1 - q^{m_j})^{k_j}}$$

for $c_1, \dots, c_r \geq 0$, $k_1 \geq 2$, $k_2, \dots, k_r \geq 1$ and $a \in [0, 1]$ can be seen as q -analogue of MZVs: For $a = 1$ we have seen already by proving the q -Ohno relation, how this works. For arbitrary a , it is not clear so far, whether we can prove more identities among q MZVs with this shape of the connected sum. This could be interesting for the future.

Acknowledgements

I would like to thank Kathrin Bringmann for her helpful comments on the paper. Furthermore, I thank Henrik Bachmann and Ulf Kühn for fruitful discussions and lots of comments while supervising my master thesis, of which this paper is part of.

References

- [Bac] H. Bachmann: *Multiple Eisenstein series and q -analogues of multiple zeta values*, PhD Thesis (2015).
- [Bra] D. Bradley: *Multiple q -zeta values*, J. of Algebra **283** (2005), 752–798.
- [Bri] B. Brindle: *Dualities of q -analogues of multiple zeta values*, Master thesis, University of Hamburg, <https://sites.google.com/view/benjamin-brindle/start>, 2021.
- [EMS] K. Ebrahimi-Fard, D. Manchon, J. Singer: *Duality and (q -)Multiple Zeta Values*, Adv. Math. **298** (2016), 254–285.
- [GKZ] H. Gangl, M. Kaneko, D. Zagier: *Double zeta values and modular forms*, Automorphic forms and zeta functions, Tokyo, Japan, World Scientific (2006).
- [OT] J. Okuda, Y. Takeyama: *On relations for the multiple q -zeta values*, Ramanujan J. **14**(3) (2007), 379–387.
- [Sek] S. Seki: *Connectors*, RIMS Kôkyûroku 2160 (2020), 15–27.
- [Sin] J. Singer: *q -Analogues of Multiple Zeta Values and their application in renormalization*, Dissertation, Erlangen-Nürnberg University (2017).
- [SY] S. Seki, S. Yamamoto: *A new proof of the duality of Multiple Zeta Values and its generalizations*, Int. J. of Number Theory, **15**(6) (2019), 1261–1265.
- [Zag] D. Zagier: *Values of Zeta Functions and Their Applications*, First European Congress of Mathematics Paris (1994).
- [Zha] J. Zhao: *Uniform approach to double shuffle and duality relations of various q -Analogues of Multiple Zeta Values via Rota-Baxter algebras*, In J. I. Burgos Gil, K. Ebrahimi-Fard, H. Gangl (Eds.), *Periods in quantum field theory and arithmetic*. Cham: Springer (2014), 259–292.

UNIVERSITY OF COLOGNE, DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES
Weyertal 86-90, 50931 Cologne, Germany
E-mail address: bbrindle@uni-koeln.de