RECURSIVE FORMULAS FOR THE KRONECKER QUANTUM CLUSTER ALGEBRA WITH PRINCIPAL COEFFICIENTS

MING DING, FAN XU AND XUEQING CHEN

Abstract. We use the quantum version of Chebyshev polynomials to explicitly construct the recursive formulas for the Kronecker quantum cluster algebra with principal coefficients. As a byproduct, we obtain two bar-invariant positive \mathbb{Z}^p bases with one being the atomic basis.

1. INTRODUCTION

Cluster algebras were invented by Fomin and Zelevinsky [\[16\]](#page-20-0), and their quantum analogues (called quantum cluster algebras) were introduced by Berenstein and Zelevinsky [\[3\]](#page-19-0). A (quantum) cluster algebra is generated by distinguished generators called (quantum) cluster variables which are grouped into overlapping subsets called clusters. The first important result is the Laurent phenomenon which asserts that any (quantum) cluster algebra element is a Laurent polynomial in the cluster variables in any cluster. A (quantum) cluster algebra element is said to be positive if the Laurent polynomial has non-negative coefficients.

In order to obtain more structural results in cluster theory, one needs to construct cluster multiplication formulas. For the classical cluster algebras of acyclic quivers, Sherman and Zelevinsky [\[23\]](#page-20-1) firstly provided the cluster multiplication formulas in rank 2 cluster algebras of finite and affine types. This result was generalized to rank 3 cluster algebra of affine type $A_2^{(1)}$ by Cerulli [\[7\]](#page-20-2). Caldero and Keller [\[4\]](#page-19-1) constructed the cluster multiplication formulas between two generalized cluster variables for simply laced Dynkin quivers, which were generalized to affine types in [\[17\]](#page-20-3) and to acyclic types in [\[24,](#page-20-4) [25\]](#page-20-5). In the quantum setting, Ding and Xu [\[10\]](#page-20-6) firstly gave the cluster multiplication formulas of the quantum cluster algebra of type $A_1^{(1)}$ without coefficients, which were generalized to type $A_2^{(2)}$ by Bai, Chen, Ding and Xu in [\[1\]](#page-19-2). Recently, Chen, Ding and Zhang [\[9\]](#page-20-7) constructed the cluster multiplication formulas in the acyclic quantum cluster algebras with arbitrary coefficients through some quotients of derived Hall algebras of acyclic valued quivers. Note that, for general cases, although the structure constants deduced from cluster multiplication formulas have the homological interpretations, it is still difficult to explicitly compute them in these formulas.

²⁰²⁰ Mathematics Subject Classification. 13F60, 16G20, 18E30.

Key words and phrases. quantum cluster algebra, cluster variable, positive basis.

2 MING DING, FAN XU AND XUEQING CHEN

Cluster multiplication formulas play a very important role in constructing "good" bases of cluster algebras. A basis of a cluster algebra is said to be positive if its structure constants are positive, and to be atomic (or canonical) if positive elements are exactly linear combinations of basis elements with non-negative coefficients. Note that the atomic basis must be a positive basis. It attracts a lot of attention to find "good" bases such as the atomic basis or canonical basis of a cluster algebra. Sherman and Zelevinsky [\[23\]](#page-20-1) introduced and constructed atomic bases in rank 2 cluster algebras of finite and affine types, which were originally called canonical bases. Cerulli constructed the atomic basis in the rank 3 cluster algebra of affine type $A_2^{(1)}$ 2 in [\[7\]](#page-20-2). He also proved the coincidence of the atomic basis and the set of cluster monomials for cluster algebras of finite type [\[8\]](#page-20-8). For non-finite type cluster algebras, the set of cluster monomials does not form a basis even it is linearly independent. Dupont and Thomas [\[14\]](#page-20-9) constructed the atomic bases of cluster algebras of affine type A . They also provided a new proof of Cerulli's result for cluster algebras of type A. In the quantum setting, the atomic bases of the quantum cluster algebras of type $A_1^{(1)}$ $\stackrel{(1)}{1}$ (or $\widetilde{A}_{1,1}$), $A_2^{(2)}$ $\sum_{i=1}^{2}$ and $A_{2n-1,1}$ without coefficients were constructed in [\[10,](#page-20-6) [1,](#page-19-2) [12\]](#page-20-10) by using the explicit cluster multiplication formulas. However, most of quantum cluster algebras must have coefficients involved. It is natural to ask whether there exists an explicit treatment of the atomic bases for the acyclic quantum cluster algebras with coefficients by using representation theory of quivers. Sherman and Zelevinsky [\[23\]](#page-20-1) initiated the use of Chebyshev polynomials to construct the atomic bases of rank 2 cluster algebras of affine type. In order to obtain atomic bases of cluster algebras with principle coefficients, Dupont [\[15\]](#page-20-11) introduced the deformed Chebyshev polynomials and conjectured that the deformed Chebyshev polynomials have interactions with atomic bases or canonically positive bases in affine cluster algebras. He proved this conjecture for cluster algebra of affine type $A_1^{(1)}$ $_1^{(1)}$. Note that for the $A_2^{(1)}$ $2^{(1)}$ case, Cerulli [\[7\]](#page-20-2) already used the deformed Chebyshev polynomials to prove the existence of the atomic basis.

As the Chebyshev polynomials are used to construct the positive bases in classical cluster algebras, it is important to study them in quantized cases. In this paper, we modify the definition of Chebyshev polynomials which can be viewed as the quantum analogue of those in [\[7,](#page-20-2) [15\]](#page-20-11). Then we explicitly construct the recursive formulas for the Kronecker quantum cluster algebra with principal coefficients based on the representation theory of the Kronecker quiver. As a direct corollary, we obtain two bar-invariant positive ZP-bases with one being the atomic basis. We expect to apply results and methods used in this paper in the future studies for all affine types.

The paper is organized as follows: in Section 2, we collect basic concepts of quantum cluster algebras and quantum cluster characters and recall some multiplication formulas; Section 3 provides detailed calculations of recursive formulas in the Kronecker quantum cluster algebra with principle coefficients; in Section 4, we obtain an explicit treatment of two bar-invariant positive ZP-bases in terms of quiver representations.

2. Quantum cluster algebra and quantum cluster character

We recall some basic features of quantum cluster algebras and quantum cluster characters.

2.1. Quantum cluster algebra. Let $\Lambda : \mathbb{Z}^m \times \mathbb{Z}^m \to \mathbb{Z}$ be a skew-symmetric bilinear form on \mathbb{Z}^m . The natural basis of \mathbb{Z}^m is denoted by $\{\mathbf{e}_1, \cdots, \mathbf{e}_m\}$. Denote by $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ be the ring of integer Laurent polynomials of an indeterminate q. The quantum torus \mathcal{T}_{q} associated to Λ as the $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ -algebra is freely generated by the set $\{X^{\mathbf{e}} : \mathbf{e} \in \mathbb{Z}^m\}$ with the following twisted product:

$$
X^{\mathbf{e}} \cdot X^{\mathbf{f}} = \mathfrak{q}^{\Lambda(\mathbf{e}, \mathbf{f})/2} X^{\mathbf{e} + \mathbf{f}} \quad \text{ for any } \mathbf{e}, \mathbf{f} \in \mathbb{Z}^{\mathbf{m}}.
$$

In what follows, we omit the twisted product \cdot for the multiplication in algebras. The skew-field of fractions for \mathcal{T}_{q} is denoted by \mathcal{F}_{q} . It will cause no confusion if we use the same letter Λ to designate the $m \times m$ skew-symmetric integer matrix associated to the bilinear form Λ.

Let $\widetilde{B}=(b_{ij})$ be an $m \times n$ integer matrix with $n \leq m$ and \widetilde{B}^{tr} the transpose of \widetilde{B} . The *principle* part of \widetilde{B} is the upper $n \times n$ submatrix of \widetilde{B} denoted by B. We call the pair (Λ, \widetilde{B}) compatible if $\widetilde{B}^{tr} \Lambda = (D|0)$ for some diagonal matrix D with positive entries. An *initial quantum seed* $(\Lambda, \widetilde{B}, X)$ of $\mathcal{F}_{\mathfrak{q}}$ consists of a compatible pair (Λ, \widetilde{B}) and the set $X = \{X_1, \dots, X_m\}$ with $X_i := X^{\mathbf{e_i}}$ for $1 \leq i \leq m$. For any $1 \leq k \leq n$, the mutation μ_k of the quantum seed $(\Lambda, \widetilde{B}, X)$ in direction k is the new quantum seed $\mu_k(\Lambda, \tilde{B}, X) := (\Lambda', \tilde{B}', X')$ given as follows:

(1) $\Lambda' = E^{tr} \Lambda E$, where the $m \times m$ matrix $E = (e_{ij})$ is given by

$$
e_{ij} = \begin{cases} \delta_{ij} & \text{if } j \neq k; \\ -1 & \text{if } i = j = k; \\ [-b_{ik}]_+ & \text{if } i \neq j = k. \end{cases}
$$

(2) $B' = (b'_{ij})$ with

$$
b'_{ij} = \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k; \\ b_{ij} + [b_{ik}]_+ b_{kj} + b_{ik}[-b_{kj}]_+ & \text{otherwise.} \end{cases}
$$

(3) $X' = \{X'_1, \cdots, X'_m\}$ with

$$
X'_{i} = \begin{cases} X_{i} & \text{if } i \neq k; \\ X^{\sum_{1 \leq j \leq m} [b_{jk}]_{+} \mathbf{e}_{j} - \mathbf{e}_{k}} + X^{\sum_{1 \leq j \leq m} [-b_{jk}]_{+} \mathbf{e}_{j} - \mathbf{e}_{k}} & \text{if } i = k, \end{cases}
$$

where $[a]_+ := \max\{0, a\}$ for $a \in \mathbb{Z}$.

Note that the mutation is an involution, i.e., $\mu_k^2(\Lambda, \tilde{B}, X) = (\Lambda, \tilde{B}, X)$. Thus we have an equivalence relation: the quantum seed $(\Lambda, \widetilde{B}, X)$ is mutation-equivalent to $(\Lambda', \tilde{B}', X')$, denoted by $(\Lambda, \tilde{B}, X) \sim (\Lambda', \tilde{B}', X')$, if they can be obtained from each other by finite sequence of mutations. In the context of a quantum seed $(\Lambda', \tilde{B}', X')$,

we call the set $\{X'_i \mid 1 \leq i \leq n\}$ the *cluster* of the seed and call the elements X'_i the cluster variables. The cluster monomials are the elements $X^{\prime e}$ with $e \in (\mathbb{Z}_{\geq 0})^m$ and $e_i = 0$ for $n + 1 \le i \le m$. The elements in the set $\mathbb{P} := \{X_i \mid n + 1 \le i \le m\}$ are called the *coefficients*. Let $\mathbb{Z}P$ be the ring of Laurent polynomials in the elements of P with coefficients in $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}]$. The quantum cluster algebra $\mathcal{A}_{\mathfrak{q}}(\Lambda, \widetilde{B})$ is defined as the $\mathbb{Z}\mathbb{P}$ -subalgebra of \mathcal{F}_{q} generated by all cluster variables. On \mathcal{T}_{q} , we can define the Z-linear bar-involution:

$$
\overline{\mathfrak{q}^{\frac{r}{2}}X^{\mathbf{e}}} = \mathfrak{q}^{-\frac{r}{2}}X^{\mathbf{e}}, \quad \text{for any } r \in \mathbb{Z} \text{ and } \mathbf{e} \in \mathbb{Z}^{m}.
$$

It is straightforward to show that $\overline{XY} = \overline{Y} \ \overline{X}$ for any $X, Y \in \mathcal{T}_{\mathfrak{q}}$ and that quantum cluster monomials are bar-invariant. The bar-involution on $\mathcal{A}_{q}(\Lambda, B)$ can be naturally induced.

The following celebrated quantum Laurent phenomenon proved by Berenstein and Zelevinsky is one of the most important structural results on quantum cluster algebras, which asserts that every cluster variable can be written as a Laurent polynomial of cluster variables in any other cluster.

Theorem 2.1. [\[3,](#page-19-0) Theorem 5.1] The quantum cluster algebra $\mathcal{A}_{\mathfrak{a}}(\Lambda, \widetilde{B})$ is a subalgebra of the ring of Laurent polynomials in the cluster variables in any cluster over ZP.

2.2. Quantum cluster character. Let $k = \mathbb{F}_q$ be a finite field of order q. Fix two positive integers $n \leq m$. Let \tilde{Q} be an acyclic valued quiver with vertex set $\{1, 2, \ldots, m\}$. The full subquiver Q on the vertices $\{1, \ldots, n\}$ is called the principal part of \tilde{Q} . The elements in the subset $\{n+1,\ldots,m\}$ are called the frozen vertices.

Let $\mathfrak{S} = kQ$ and $\mathfrak{S} = kQ$ be the path algebras of Q and Q, respectively. Let S_i , P_i and I_i be the simple, projective and injective $\tilde{\mathfrak{S}}$ -module associated to the vertex i for any $1 \leq i \leq m$, respectively. The $m \times n$ matrix $\widetilde{B} = (b_{ij})$ associated to \widetilde{Q} is defined by

$$
b_{ij} = \text{dim}_{\text{End}_{\widetilde{\mathfrak{S}}}(S_i)^{op}} \text{Ext}^1_{\widetilde{\mathfrak{S}}}(S_i, S_j) - \text{dim}_{\text{End}_{\widetilde{\mathfrak{S}}}(S_i)} \text{Ext}^1_{\widetilde{\mathfrak{S}}}(S_j, S_i)
$$

for $1 \leq i \leq m, 1 \leq j \leq n$. Note that the upper $n \times n$ part of \widetilde{B} is B. The left $m \times n$ submatrix of the identity matrix of size $m \times m$ is denoted by I.

Assume that there exists some skew-symmetric integer matrix Λ of size $m \times m$ such that

$$
\Lambda(-\widetilde{B})=\left(\begin{array}{c}D_n\\0\end{array}\right),\,
$$

where $D_n = \text{diag}(d_1, \dots, d_n)$ with $d_i \in \mathbb{N}$ for $1 \leq i \leq n$. Note that if we enlarge the quiver Q to \tilde{Q} , where we attach principal frozen vertices to the valued quiver Q , such a matrix Λ always exists [\[26\]](#page-20-12).

The quantum cluster algebra associated to this obtained compatible pair (Λ, \widetilde{B}) is simply denoted by $\mathcal{A}_{\mathfrak{q}}(Q)$.

Let $R = R_{\widetilde{Q}} = (\widetilde{r}_{ij})$ be the $m \times n$ matrix with

$$
\widetilde{r}_{ij} := \mathrm{dim}_{\mathrm{End}_{\widetilde{\mathfrak{S}}}}(S_i) \mathrm{Ext}^1_{\widetilde{\mathfrak{S}}}(S_j, S_i)
$$

for $1 \leq i \leq m$ and $1 \leq j \leq n$. And define $\widetilde{R}' := \widetilde{R}_{\widetilde{Q}^{op}}$. Note that the principal part of \widetilde{R} is $R, \widetilde{B} = \widetilde{R}' - \widetilde{R}$ and $B = R' - R$.

Let $\mathcal{C}_{\tilde{O}}$ be the cluster category of the valued quiver \tilde{Q} with the shift functor [1] and the Aulander-Reiten translation functor τ . For more details about cluster category, the readers can refer to [\[2,](#page-19-3) [22\]](#page-20-13). Note that each object M of $\mathcal{C}_{\tilde{Q}}$ can be uniquely decomposed up to isomorphism as follows:

$$
M = M_0 \oplus P_M[1]
$$

where M_0 is an \mathfrak{S} -module and P_M is a projective \mathfrak{S} -module. Let $P_M = \bigoplus_{1 \leq i \leq m} m_i P_i$. The definition of dimension vector \dim on \mathfrak{S} -modules can be extended on objects of $\mathcal{C}_{\tilde{\mathcal{O}}}$ by defining

$$
\underline{\dim} M = \underline{\dim} M_0 - (m_i)_{1 \le i \le m}.
$$

In the following, we will denote the dimension vector of an $\mathfrak{S}\text{-module }X$ by the corresponding underlined lower case letter \underline{x} viewed as a column vector in \mathbb{Z}^n . For simplifying notations, we write $Ext^1(M, N) := Ext^1_{\mathfrak{S}}(M, N)$ and $Hom(M, N) :=$ $\text{Hom}_{\widetilde{\mathfrak{S}}}(M, N)$. The Euler form on $\widetilde{\mathfrak{S}}$ -modules M and N is given by

$$
\langle M, N \rangle = \dim_k \text{Hom}(M, N) - \dim_k \text{Ext}^1(M, N).
$$

Let $\mathcal{A}_q(Q)$ be the specialization of the quantum cluster algebra $\mathcal{A}_q(Q)$ by evaluating $q = q$. For an G-module N, we denote its socle and radical by Soc N and rad N. Let M be an $\mathfrak{S}\text{-module}$ and I an injective $\mathfrak{S}\text{-module}$, then the quantum cluster character of $\mathcal{A}_q(Q)$ is defined as:

$$
X_{M\oplus I[-1]} = \sum_{\underline{e}} |\mathrm{Gr}_{\underline{e}}M| q^{-\frac{1}{2}\langle \underline{e}, \underline{m}-\underline{e}-\underline{i}\rangle} X^{-\widetilde{B}\underline{e}-(\widetilde{I}-\widetilde{R}')\underline{m}+\underline{\dim}} \mathrm{Soc} I,
$$

where $\underline{\dim} I = \underline{i}$, $\underline{\dim} M = \underline{m}$ and $\operatorname{Gr}_{e} M$ is the Grassmannian of sub-representations of M with dimension vector \underline{e} . Note that for any projective $\mathfrak{S}\text{-module }P$ and injective $\mathfrak{S}\text{-module } I$ with $\text{Soc } I = P/\text{rad } P$, we have that

$$
X_{P[1]} = X_{\tau P} = X^{\dim P/\text{rad }P} = X^{\dim \text{Soc }I} = X_{I[-1]} = X_{\tau^{-1}I}.
$$

The main results in [\[20,](#page-20-14) [22\]](#page-20-13) show that $\mathcal{A}_q(Q)$ as the ZP-subalgebra of \mathcal{F}_q is generated by cluster variables

 ${X_M|M}$ is an indecomposable rigid G-module} $\sqcup {X_{I_i[-1]}}|1 \leq i \leq n$.

Let M and N be two G-modules and $\theta : N \to \tau M$ be a morphism, we can deduce an exact sequence

$$
0 \longrightarrow D \longrightarrow N \stackrel{\theta}{\longrightarrow} \tau M \longrightarrow \tau A \oplus I \longrightarrow 0
$$

where $D = \ker \theta$, $\tau A \oplus I = \operatorname{coker} \theta$, I is an injective $\widetilde{\mathfrak{S}}$ -module, A and M have the same maximal projective summand.

The following theorems are the generalizations of those in [\[11,](#page-20-15) [20\]](#page-20-14) to acyclic valued quivers.

Theorem 2.2. [\[22,](#page-20-13) Theorem 4.5] Assume M and N are $\mathfrak{S}\text{-modules with a unique (up)}$ to scalar) nontrivial extension $E \in Ext^1(M, N)$, in particular $dim_{End(M)} Ext^1(M, N)$ 1. Let $\theta \in Hom(N, \tau M)$ be the equivalent morphism with A, D, I described as above. Furthermore assume that $Hom(A \oplus D, I) = 0 = Ext^1(A, D)$. Then we have that

$$
\begin{split} &X_M X_N \\ =& q^{\frac{1}{2}\Lambda((\widetilde{I}-\widetilde{R}')\underline{m}, (\widetilde{I}-\widetilde{R}')\underline{n})}X_E+q^{\frac{1}{2}\Lambda((\widetilde{I}-\widetilde{R}')\underline{m}, (\widetilde{I}-\widetilde{R}')\underline{n})+\frac{1}{2}\langle M, N\rangle-\frac{1}{2}\langle A, D\rangle}X_{D\oplus A\oplus I[-1]}.\end{split}
$$

Let M be an \mathfrak{S} -module and I an injective $\widetilde{\mathfrak{S}}$ -module. Let ν be the Nakayama functor on \mathfrak{S} -modules and write $P = \nu^{-1}(I)$. From morphisms $f : M \to I$ and $g: P \to M$, we can deduce two exact sequences

$$
0\longrightarrow G\longrightarrow M\stackrel{f}{\longrightarrow} I\longrightarrow I'\longrightarrow 0
$$

and

$$
0 \longrightarrow P' \longrightarrow P \stackrel{g}{\longrightarrow} M \longrightarrow F \longrightarrow 0
$$

where $G = \text{Ker } f$, $I' = \text{coker } f$ is an injective $\tilde{\mathfrak{S}}$ -module, $P' = \text{Ker } g$ is a projective $\mathfrak{S}\text{-module and } F = \text{coker } g.$

Theorem 2.3. [\[22,](#page-20-13) Theorem 4.8] Let M be an $\mathfrak{S}\text{-modules}$ I and P be $\widetilde{\mathfrak{S}}\text{-modules}$ defined as above. Assume that there exist unique (up to scalar) morphisms $f \in$ $Hom(M, I)$ and $g \in Hom(P, M)$, in particular

 $dim_{End(I)}Hom(M, I) = dim_{End(P)}Hom(P, M) = 1.$

Define F, G, I', P' as above and assume further that $Hom(P', F) = Hom(G, I') = 0$. Then we have

$$
X_M X_{I[-1]}=q^{-\frac{1}{2}\Lambda((\widetilde{I}-\widetilde{R}')\underline{m},\underline{\dim}SocI)}X_{G\oplus I'[-1]}+q^{-\frac{1}{2}\Lambda((\widetilde{I}-\widetilde{R}')\underline{m},\underline{\dim}SocI)-\frac{1}{2}\dim_k\operatorname{End}(I)}X_{F\oplus P'[1]}.
$$

3. Recursive formulas for the Kronecker quantum cluster algebra WITH PRINCIPLE COEFFICIENTS

Consider the Kronecker quiver Q , i.e., the quiver of type $\tilde{A}_{1,1}$:

$$
1\bullet \mathrel{\overbrace{\text{---}}} \mathbin{\mathbin{\raisebox{.3pt}{:}}\mathbin{\raisebox{.3pt}{:}}}
$$

It is well-known [\[13\]](#page-20-16) that indecomposable kQ-modules are grouped into (up to isomorphism) three families: the indecomposable preprojective modules with dimension vector $(l, l + 1)$ for $l \in \mathbb{Z}_{\geq 0}$ denoted by $V(-l)$, the indecomposable regular modules (in particular, denote by $R(l)$ the indecomposable regular module with dimension vector (l, l) for $l \in \mathbb{Z}_{\geq 1}$ from the same homogeneous tube whose mouth has dimension vector $(1, 1)$, and the indecomposable preinjective modules with dimension vector $(l + 1, l)$ for $l \in \mathbb{Z}_{\geq 0}$ denoted by $V(l + 3)$.

The preprojective component looks like the following:

The preinjective components looks like the following:

We consider the following ice quiver \widetilde{Q} with frozen vertices 3 and 4:

Thus we have

$$
\widetilde{R}' = \begin{pmatrix} 0 & 2 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}, \ \widetilde{R} = \begin{pmatrix} 0 & 0 \\ 2 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \ \widetilde{I} - \widetilde{R}' = \begin{pmatrix} 1 & -2 \\ 0 & 1 \\ -1 & 0 \\ 0 & -1 \end{pmatrix}, \ \widetilde{B} = \begin{pmatrix} 0 & 2 \\ -2 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}.
$$

An easy calculation shows that the following skew-symmetric 4×4 integer matrix

$$
\Lambda = \left(\begin{array}{rrrr} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & -2 \\ 0 & 1 & 2 & 0 \end{array}\right)
$$

satisfying

$$
\Lambda(-\widetilde{B})=\left(\begin{array}{c}D_2\\0\end{array}\right),\,
$$

where $D_2 =$ $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. The quantum cluster algebra $\mathcal{A}_q(Q)$ associated to this pair (Λ, B) is called the Kronecker quantum cluster algebra with principle coefficients. We denote the coefficients by $y_1 = X_{I_3[-1]}$, $y_2 = X_{I_4[-1]}$, and all cluster variables by

 $x_l = X_{V(l)}$ for $l \in \mathbb{Z}$ where $x_1 = X_{V(1)} = X_{I_1[-1]}$ and $x_2 = X_{V(2)} = X_{I_2[-1]}$ are initial cluster variables.

By the definition of quantum cluster characters, it follows that

$$
X_{R(1)} = X^{(1,-1,1,1)} + X^{(-1,-1,1,0)} + X^{(-1,1,0,0)}.
$$

Therefore the expression of $X_{R(1)}$ is independent of the choice of the indecomposable regular module with dimension vector $(1, 1)$ and then we denote $X_{R(1)}$ by X_{δ} . Note that the element X_{δ} coincides with the quantum loop element s_1 defined in [\[6\]](#page-20-17) in terms of perfect matchings associated to the Kronecker quantum cluster algebra with principle coefficients. A direct calculation shows that

$$
X_{\delta} = x_0 x_3 - q^{\frac{3}{2}} x_1 x_2 y_2,
$$

and thus X_{δ} belongs to $\mathcal{A}_{q}(Q)$ (cf. [\[6\]](#page-20-17)).

The *n*-th Chebyshev polynomials of the second kind $S_n(x)$ appeared in [\[5\]](#page-20-18) is related to the dual semi-canonical basis of the Kronecker cluster algebra. We modify the definition of $S_n(x)$ for $n \in \mathbb{Z}_{\geq 0}$ on X_δ as follows

$$
S_0(X_\delta) = 1, S_1(X_\delta) = X_\delta, S_2(X_\delta) = S_1(X_\delta)S_1(X_\delta) - X^{(0,0,1,1)},
$$

\n
$$
S_{n+1}(X_\delta) = S_n(X_\delta)S_1(X_\delta) - X^{(0,0,1,1)}S_{n-1}(X_\delta), \text{ for } n \ge 2.
$$

In the following, we denote $S_n(X_\delta)$ by S_n for simplicity. According to the definition of S_n and the fact that $S_1 \in \mathcal{A}_q(Q)$, it is easy to see $S_n \in \mathcal{A}_q(Q)$ for any $n \geq 1$.

We start with some technical lemmas:

Lemma 3.1. For any $n \geq 0$ and $m \geq 0$, we have that

(1) $y_1 S_n = q^{-n} S_n y_1;$ (2) $y_2S_n = q^nS_n y_2;$ (3) $X^{(0,0,1,1)}S_n = S_n X^{(0,0,1,1)};$ (4) $S_m S_n = S_n S_m$.

Proof. (1) When $n = 0$, it is trivial. When $n = 1$, we have

$$
y_1 S_1 = X^{(0,0,1,0)}(X^{(1,-1,1,1)} + X^{(-1,-1,1,0)} + X^{(-1,1,0,0)})
$$

= $q^{-1}(X^{(1,-1,1,1)} + X^{(-1,-1,1,0)} + X^{(-1,1,0,0)})X^{(0,0,1,0)}$
= $q^{-1}S_1y_1$.

By induction, we have

$$
y_1 S_{n+1} = y_1 (S_n S_1 - X^{(0,0,1,1)} S_{n-1})
$$

= $q^{-(n+1)} S_n S_1 y_1 - q^{-2} X^{(0,0,1,1)} y_1 S_{n-1}$
= $q^{-(n+1)} S_n S_1 y_1 - q^{-2} q^{-(n-1)} X^{(0,0,1,1)} S_{n-1} y_1$
= $q^{-(n+1)} S_{n+1} y_1$.

- (2) The proof is similar to (1).
- (3) Note that $X^{(0,0,1,1)} = qy_1y_2$, the proof follows from (1) and (2).
- (4) According to (3) and by induction, the statement follows immediately. \Box

Lemma 3.2. For $n \geq 0$, we have that

(1)
$$
\overline{S_n} = S_n;
$$

(2) $\overline{X^{(0,0,1,1)}S_n} = X^{(0,0,1,1)}S_n.$

Proof. (1) When $n = 0$, it is trivial. When $n = 1$, by the definition of X_{δ} , we see that $\overrightarrow{X_{\delta}} = X_{\delta}$, i.e, $\overline{S_1} = S_1$. Then by induction and Lemma [3.1,](#page-7-0) we have

$$
\overline{S_{n+1}} = \overline{S_n S_1 - X^{(0,0,1,1)} S_{n-1}}
$$

\n
$$
= \overline{S_1} \overline{S_n} - \overline{S_{n-1}} \overline{X^{(0,0,1,1)}}
$$

\n
$$
= S_1 S_n - S_{n-1} X^{(0,0,1,1)}
$$

\n
$$
= S_n S_1 - X^{(0,0,1,1)} S_{n-1}
$$

\n
$$
= S_{n+1}
$$

(2) By (1) and Lemma [3.1,](#page-7-0) we have

$$
\overline{X^{(0,0,1,1)}S_n} = \overline{S_n} \overline{X^{(0,0,1,1)}} = S_n X^{(0,0,1,1)} = X^{(0,0,1,1)} S_n.
$$

Hence, the proof is finished. \Box

The following proposition gives multiplication formulas for the modified Chebyshev polynomials of the second kind on X_{δ} .

Proposition 3.3. For $n \geq m \geq 1$, we have that

$$
S_m S_n = S_{n+m} + X^{(0,0,1,1)} S_{n+m-2} + \dots + X^{(0,0,m-1,m-1)} S_{n-m+2} + X^{(0,0,m,m)} S_{n-m}.
$$

Proof. When $m = 1$, the statement is just the definition. Assume that we have the equality for all positive integers m less than or equal to k. Then by Lemma [3.1,](#page-7-0) we have

$$
S_{k+1}S_n = (S_1S_k - X^{(0,0,1,1)}S_{k-1})S_n
$$

= $S_1(S_{n+k} + X^{(0,0,1,1)}S_{n+k-2} + \cdots X^{(0,0,k-1,k-1)}S_{n-k+2} + X^{(0,0,k,k)}S_{n-k}) -$
 $X^{(0,0,1,1)}(S_{n+k-1} + X^{(0,0,1,1)}S_{n+k-3} + \cdots X^{(0,0,k-2,k-2)}S_{n-k+3} + X^{(0,0,k-1,k-1)}S_{n-k+1})$
= $(S_{n+k+1} + X^{(0,0,1,1)}S_{n+k-1}) + X^{(0,0,1,1)}(S_{n+k-1} + X^{(0,0,1,1)}S_{n+k-3}) + \cdots +$
 $X^{(0,0,k-1,k-1)}(S_{n-k+3} + X^{(0,0,1,1)}S_{n-k+1}) + X^{(0,0,k,k)}(S_{n-k+1} + X^{(0,0,1,1)}S_{n-k-1})$
 $- X^{(0,0,1,1)}(S_{n+k-1} + X^{(0,0,1,1)}S_{n+k-3} + \cdots$
 $+ X^{(0,0,k-2,k-2)}S_{n-k+3} + X^{(0,0,k-1,k-1)}S_{n-k+1})$
= $S_{n+k+1} + X^{(0,0,1,1)}S_{n+k-1} + \cdots + X^{(0,0,k,k)}S_{n-k+1} + X^{(0,0,k+1,k+1)}S_{n-k-1}.$

Hence, the proof is finished.

Now, we give a representation-theoretic interpretation of the element S_n for any $n \geq 1$.

Proposition 3.4. For $n \geq 1$, we have that $S_n = X_{R(n)}$.

Proof. The case when $n = 1$ is trivial. Note that we have

$$
0 \longrightarrow R(1) \longrightarrow R(n+1) \longrightarrow R(n) \longrightarrow 0
$$

and since dim $Ext^1(R(n), R(1)) = dim Hom(R(1), \tau R(n)) = 1$, we have that

$$
0 \longrightarrow R(1) \longrightarrow \tau R(n) \longrightarrow \tau R(n-1) \oplus I_3 \oplus I_4 \longrightarrow 0.
$$

By Theorem [2.2,](#page-5-0) we then have that

$$
X_{R(n)}X_{R(1)} = X_{R(n+1)} + X_{R(n-1)\oplus (I_3\oplus I_4)[-1]}.
$$

A direct calculation shows that

$$
X^{(0,0,1,1)}X^{-\widetilde{B}\underline{e}-(\widetilde{I}-\widetilde{R}')(n-1,n-1)^{tr}}=X^{-\widetilde{B}\underline{e}-(\widetilde{I}-\widetilde{R}')(n-1,n-1)^{tr}+(0,0,1,1)},
$$

which implies $X_{R(n)}X_{R(1)} = X_{R(n+1)} + X^{(0,0,1,1)}X_{R(n-1)}$.

Note that $S_nS_1 = S_{n+1} + X^{(0,0,1,1)}S_{n-1}$. Therefore, by induction, we can obtain that $S_{n+1} = X_{R(n+1)}$.

Lemma 3.5. In $\mathcal{A}_q(Q)$, we have that

(1) $S_1x_0 = x_{-1} + q^{-\frac{1}{2}}X^{(1,0,0,1)}$; (2) $S_1x_{-m} = x_{-(m+1)} + X^{(0,0,1,1)}x_{-(m-1)}$ for $m \ge 1$.

Proof. (1) Note that we have

$$
0 \longrightarrow V(0) \longrightarrow V(-1) \longrightarrow R(1) \longrightarrow 0
$$

and since dim $\text{Ext}^1(R(1), V(0)) = \dim \text{Hom}(V(0), \tau R(1)) = 1$, we have

$$
0 \longrightarrow V(0) \longrightarrow \tau R(1) \longrightarrow I_1 \oplus I_4 \longrightarrow 0.
$$

By Theorem [2.2,](#page-5-0) we can then obtain that $S_1x_0 = X_{R(1)}X_{V(0)} = x_{-1} + q^{-\frac{1}{2}}X^{(1,0,0,1)}$;

(2) Note that dim $\text{Ext}^1(R(1), V(-m)) = \dim \text{Hom}(V(-m), \tau R(1)) = 1$, we have that

$$
0 \longrightarrow V(-m) \longrightarrow V(-m-1) \longrightarrow R(1) \longrightarrow 0
$$

and

$$
0 \longrightarrow V(-m+1) \longrightarrow V(-m) \longrightarrow \tau R(1) \longrightarrow I_3 \oplus I_4 \longrightarrow 0.
$$

By Theorem [2.2,](#page-5-0) we have that $S_1 x_{-m} = x_{-(m+1)} + q^{-\frac{1}{2}} X_{V(-m+1) \oplus (I_3 \oplus I_4)[-1]}$. Note that

$$
X^{(0,0,1,1)}X^{-\widetilde{B}\underline{e}-(\widetilde{I}-\widetilde{R}')(m-1,m)^{tr}}=q^{-\frac{1}{2}}X^{-\widetilde{B}\underline{e}-(\widetilde{I}-\widetilde{R}')(m-1,m)^{tr}+(0,0,1,1)},
$$

which implies $S_1x_{-m} = x_{-(m+1)} + X^{(0,0,1,1)}x_{-(m-1)}$ for $m \ge 1$.

We are now able to prove the following recursive formulas between modified Chebyshev polynomials of the second kind on X_{δ} and cluster variables.

Theorem 3.6. In $\mathcal{A}_q(Q)$, we have that

(1) $S_n x_0 = x_{-n} + q^{-\frac{1}{2}} S_{n-1} X^{(1,0,0,1)}$, for $n \ge 1$; (2) $S_n x_{-m} = x_{-(m+n)} + X^{(0,0,1,1)} S_{n-1} x_{-(m-1)}$, for $n \ge 1$ and $m \ge 1$; (3) $x_1S_n = x_{n+1} + q^{-\frac{1}{2}}x_0y_1S_{n-1}$, for $n \ge 1$; (4) $x_m S_n = x_{n+m} + x_{m-1} X^{(0,0,1,1)} S_{n-1}$, for $n \ge 1$ and $m \ge 2$.

Proof. We only prove (1) and (2). The proofs of (3) and (4) are similar.

(1) When $n = 1$, by Lemma [3.5](#page-9-0) (1), the statement is true. Assume that we have the equality for all positive integers less than or equal to k . Then

$$
S_{k+1}x_0 = (S_1S_k - X^{(0,0,1,1)}S_{k-1})x_0
$$

= $S_1(x_{-k} + q^{-\frac{1}{2}}S_{k-1}X^{(1,0,0,1)}) - X^{(0,0,1,1)}(x_{-(k-1)} + q^{-\frac{1}{2}}S_{k-2}X^{(1,0,0,1)})$
= $x_{-(k-1)} + X^{(0,0,1,1)}x_{-(k-1)} + q^{-\frac{1}{2}}(S_k + X^{(0,0,1,1)}S_{k-2})X^{(1,0,0,1)}$
 $- X^{(0,0,1,1)}(x_{-(k-1)} + q^{-\frac{1}{2}}S_{k-2}X^{(1,0,0,1)})$
= $x_{-(k+1)} + q^{-\frac{1}{2}}S_kX^{(1,0,0,1)};$

(2) When $n = 1$, by Lemma [3.5](#page-9-0) (2), the statement is true. Assume that we have the equality for all positive integers less than or equal to k . Then

$$
S_{k+1}x_{-m} = (S_1S_k - X^{(0,0,1,1)}S_{k-1})x_{-m}
$$

= $S_1(x_{-(m+k)} + X^{(0,0,1,1)}S_{k-1}x_{-(m-1)}) - X^{(0,0,1,1)}(x_{-(m+k-1)} + X^{(0,0,1,1)}S_{k-2}x_{-(m-1)})$
= $x_{-(m+k+1)} + X^{(0,0,1,1)}x_{-(m+k-1)} + X^{(0,0,1,1)}(S_k + X^{(0,0,1,1)}S_{k-2})x_{-(m-1)}$
 $- X^{(0,0,1,1)}(x_{-(m+k-1)} + X^{(0,0,1,1)}S_{k-2}x_{-(m-1)})$
= $x_{-(m+k+1)} + X^{(0,0,1,1)}S_kx_{-(m-1)}$.

Hence, the proof is finished.

Remark 3.7. (1) In Theorem [3.6](#page-9-1) (4), let $n = 1$ and $m \geq 2$, then we obtain

$$
x_m S_1 = x_{m+1} + x_{m-1} X^{(0,0,1,1)},
$$

which is exactly the equality in [\[6,](#page-20-17) Lemma 4.10].

(2) In Theorem [3.6](#page-9-1) (4), let $m = 2$ and $n \ge 1$, then we obtain

$$
x_2 S_n = x_{n+2} + x_1 X^{(0,0,1,1)} S_{n-1}.
$$

Compare this equality with [\[6,](#page-20-17) Lemma 4.16 (b)], it follows that $S_n = s_n$. Thus, by Proposition [3.4,](#page-8-0) the elements $X_{R(n)}$ give a representation-theoretic interpretation of those elements s_n which are expressed in terms of perfect matchings in [\[6\]](#page-20-17).

We now prove the following recursive formulas between cluster variables:

Theorem 3.8. In $\mathcal{A}_q(Q)$, we have that

(1)
$$
x_{-(m+n+2)}x_{-m} = q^{-\frac{1}{2}}S_nX^{(0,0,m,m+1)} + x_{-(n+m+1)}x_{-(m+1)}
$$
, for $m \ge 0$ and $n \ge 0$;

- (2) $x_{m+n+2}x_m = q^{\frac{1}{2}}S_n X^{(0,0,m,m-1)} + x_{n+m+1}x_{m+1}$, for $m \ge 1$ and $n \ge 0$;
- (3) $x_0x_{n+2} = q^{\frac{2n+1}{2}}x_1x_{n+1}y_2 + S_n$, for $n \ge 0$;
- (4) $x_{-m}x_1 = S_{m-1} + q^{\frac{1}{2}}x_{-(m-1)}x_0y_1$, for $m \ge 1$;
- (5) $x_{-m}x_n = qx_{-(m-1)}x_{n-1}X^{(0,0,1,1)} + S_{m+n-2}$, for $m \ge 1$ and $n \ge 2$.

Proof. (1) Note that dim $Ext^1(V(-m-2), V(-m)) = \dim Hom(V(-m), \tau V(-m 2)$) = 1. We have

$$
0 \longrightarrow V(-m) \longrightarrow 2V(-m-1) \longrightarrow V(-m-2) \longrightarrow 0
$$

and

$$
0 \longrightarrow V(-m) \longrightarrow \tau V(-m-2) \longrightarrow mI_3 \oplus (m+1)I_4 \longrightarrow 0.
$$

By Theorem [2.2,](#page-5-0) we have that $X_{V(-m-2)}X_{V(-m)} = X_{V(-m-1)}^2 + q^{-\frac{1}{2}}X_{mI_3[-1]\oplus(m+1)I_4[-1]},$ which is exactly the equality $x_{-(m+2)}x_{-m} = q^{-\frac{1}{2}} X^{(0,0,m,m+1)} + x_{-(m+1)}^2$. Therefore the statement is true for the case when $n = 0$.

When $n = 1$, we have

$$
x_{-(m+3)}x_{-m} = (S_1x_{-(m+2)} - X^{(0,0,1,1)}x_{-(m+1)})x_{-m}
$$

= $S_1(x_{-(m+1)}^2 + q^{-\frac{1}{2}}X^{(0,0,m,m+1)}) - X^{(0,0,1,1)}x_{-(m+1)}x_{-m}$
= $(x_{-(m+2)} + X^{(0,0,1,1)}x_{-m})x_{-(m+1)} + q^{-\frac{1}{2}}S_1X^{(0,0,m,m+1)} - X^{(0,0,1,1)}x_{-(m+1)}x_{-m}$
= $x_{-(m+2)}x_{-(m+1)} + q^{-\frac{1}{2}}S_1X^{(0,0,m,m+1)}$.

By induction, we then have

$$
q^{-\frac{1}{2}}S_{n+1}X^{(0,0,m,m+1)} = q^{-\frac{1}{2}}S_1S_nX^{(0,0,m,m+1)} - q^{-\frac{1}{2}}X^{(0,0,1,1)}S_{n-1}X^{(0,0,m,m+1)}
$$

= $S_1(x_{-(m+n+2)}x_{-m} - x_{-(m+n+1)}x_{-(m+1)})$
 $- X^{(0,0,1,1)}(x_{-(m+n+1)}x_{-m} - x_{-(m+n)}x_{-(m+1)})$
= $(x_{-(m+n+3)} + X^{(0,0,1,1)}x_{-(m+n+1)})x_{-m} - (x_{-(m+n+2)} + X^{(0,0,1,1)}x_{-(m+1)})x_{-(m+1)}$
 $- X^{(0,0,1,1)}(x_{-(m+n+1)}x_{-m} - x_{-(m+n)}x_{-(m+1)})$
= $x_{-(m+n+3)}x_{-m} - x_{-(m+n+2)}x_{-(m+1)}$.

(2) When $n = 0$, it follows from Theorems [2.2](#page-5-0) and [2.3.](#page-5-1) When $n = 1$, we have

$$
x_{m+3}x_m = (S_1x_{m+2} - X^{(0,0,1,1)}x_{m+1})x_m
$$

= $S_1(x_{m+1}^2 + q^{\frac{1}{2}}X^{(0,0,m,m-1)}) - X^{(0,0,1,1)}x_{m+1}x_m$
= $(x_{m+2} + X^{(0,0,1,1)}x_m)x_{m+1} + q^{\frac{1}{2}}S_1X^{(0,0,m,m-1)} - X^{(0,0,1,1)}x_{m+1}x_m$
= $x_{m+2}x_{m+1} + q^{\frac{1}{2}}S_1X^{(0,0,m,m-1)}$.

By induction, we then have

$$
q^{\frac{1}{2}}S_{n+1}X^{(0,0,m,m-1)} = q^{\frac{1}{2}}S_1S_nX^{(0,0,m,m-1)} - q^{\frac{1}{2}}X^{(0,0,1,1)}S_{n-1}X^{(0,0,m,m-1)}
$$

= $S_1(x_{n+m+2}x_m - x_{n+m+1}x_{m+1})$
 $- X^{(0,0,1,1)}(x_{n+m+1}x_m - x_{n+m}x_{m+1})$
= $(x_{n+m+3} + X^{(0,0,1,1)}x_{n+m+1})x_m - (x_{n+m+2} + X^{(0,0,1,1)}x_{n+m})x_{m+1}$
 $- X^{(0,0,1,1)}(x_{n+m+1}x_m - x_{n+m}x_{m+1})$
= $x_{n+m+3}x_m - x_{n+m+2}x_{m+1}$.

(3) When $n = 0$, it follows from Theorem [2.3.](#page-5-1) When $n = 1$, the equality follows from the definition of S_1 . According to Lemma [3.1](#page-7-0) and by induction, we have

$$
S_{n+1} = S_n S_1 - X^{(0,0,1,1)} S_{n-1}
$$

= $(x_0 x_{n+2} - q^{\frac{2n+1}{2}} x_1 x_{n+1} y_2) S_1 - (x_0 x_{n+1} - q^{\frac{2n-1}{2}} x_1 x_n y_2) X^{(0,0,1,1)}$
= $x_0 (x_{n+3} + qx_{n+1} y_1 y_2) - q^{\frac{2n+1}{2}} qx_1 x_{n+1} S_1 y_2 - (x_0 x_{n+1} - q^{\frac{2n-1}{2}} x_1 x_n y_2) q y_1 y_2$
= $x_0 x_{n+3} + qx_0 x_{n+1} y_1 y_2 - q^{\frac{2n+3}{2}} x_1 (x_{n+2} + qx_n y_1 y_2) y_2 - (x_0 x_{n+1} - q^{\frac{2n-1}{2}} x_1 x_n y_2) q y_1 y_2$
= $x_0 x_{n+3} - q^{\frac{2n+3}{2}} x_1 x_{n+2} - q^{\frac{2n+5}{2}} x_1 x_n y_1 y_2^2 + q^{\frac{2n+1}{2}} x_1 x_n y_2 y_1 y_2$
= $x_0 x_{n+3} - q^{\frac{2n+3}{2}} x_1 x_{n+2}$.

(4) When $m = 1$, it follows from Theorem [2.3.](#page-5-1) When $m = 2$, we have that

$$
x_{-2}x_1 = (S_1x_{-1} - X^{(0,0,1,1)}x_0)x_1
$$

= $S_1(1 + q^{\frac{1}{2}}x_0^2y_1) - X^{(0,0,1,1)}x_0x_1$
= $S_1 + q^{\frac{1}{2}}(x_{-1} + q^{-\frac{1}{2}}X^{(1,0,0,1)})x_0y_1 - X^{(0,0,1,1)}x_0x_1$
= $S_1 + q^{\frac{1}{2}}x_{-1}x_0y_1 + x_1y_2x_0y_1 - X^{(0,0,1,1)}x_0x_1$
= $S_1 + q^{\frac{1}{2}}x_{-1}x_0y_1$.

By induction, we have that

$$
x_{-(m+1)}x_1 = (S_1x_{-m} - X^{(0,0,1,1)}x_{-(m-1)})x_1
$$

= $S_1(S_{m-1} + q^{\frac{1}{2}}x_{-(m-1)}x_0y_1) - X^{(0,0,1,1)}(S_{m-2} + q^{\frac{1}{2}}x_{-(m-2)}x_0y_1)$
= $S_m + X^{(0,0,1,1)}S_{m-2} + q^{\frac{1}{2}}(x_{-m} + X^{(0,0,1,1)}x_{-(m-2)})x_0y_1$
 $- X^{(0,0,1,1)}(S_{m-2} + q^{\frac{1}{2}}x_{-(m-2)}x_0y_1)$
= $S_m + q^{\frac{1}{2}}x_{-m}x_0y_1$.

(5) When $m = 1$ and $n \ge 2$, we need to prove

$$
x_{-1}x_n = qx_0x_{n-1}X^{(0,0,1,1)} + S_{n-1}.
$$

When $n = 2$, by Lemma [3.1,](#page-7-0) we have that

$$
x_{-1}x_2 = x_{-1}(x_1S_1 - q^{-\frac{1}{2}}x_0y_1)
$$

\n
$$
= (q^{\frac{1}{2}}x_0^2y_1 + 1)S_1 - q^{-\frac{1}{2}}x_{-1}x_0y_1
$$

\n
$$
= q^{-\frac{1}{2}}x_0^2S_1y_1 + S_1 - q^{-\frac{1}{2}}x_{-1}x_0y_1
$$

\n
$$
= q^{-\frac{1}{2}}x_0(x_{-1} + q^{\frac{1}{2}}X^{(1,0,0,1)})y_1 + S_1 - q^{-\frac{1}{2}}x_{-1}x_0y_1
$$

\n
$$
= x_0X^{(1,0,0,1)}X^{(0,0,1,0)} + S_1 = qx_0x_1X^{(0,0,1,1)} + S_1.
$$

By induction, we have that

$$
x_{-1}x_{n+1} = x_{-1}(x_nS_1 - x_{n-1}X^{(0,0,1,1)})
$$

\n
$$
= (qx_0x_{n-1}X^{(0,0,1,1)} + S_{n-1})S_1 - (qx_0x_{n-2}X^{(0,0,1,1)} + S_{n-2})X^{(0,0,1,1)}
$$

\n
$$
= qx_0(x_{n-1}S_1)X^{(0,0,1,1)} + S_n + S_{n-2}X^{(0,0,1,1)} - (qx_0x_{n-2}X^{(0,0,1,1)} + S_{n-2})X^{(0,0,1,1)}
$$

\n
$$
= qx_0(x_n + x_{n-2}X^{(0,0,1,1)})X^{(0,0,1,1)} + S_n + S_{n-2}X^{(0,0,1,1)}
$$

\n
$$
- (qx_0x_{n-2}X^{(0,0,1,1)} + S_{n-2})X^{(0,0,1,1)}
$$

\n
$$
= qx_0x_nX^{(0,0,1,1)} + S_n.
$$

Now we proceed the induction on m , we have

$$
x_{-(m+1)}x_n = (S_1x_{-m} - X^{(0,0,1,1)}x_{-(m-1)})x_n
$$

= $S_1(qx_{-(m-1)}x_{n-1}X^{(0,0,1,1)} + S_{m+n-2}) - X^{(0,0,1,1)}(qx_{-(m-2)}x_{n-1}X^{(0,0,1,1)} + S_{m+n-3})$
= $q(x_{-m} + X^{(0,0,1,1)}x_{-(m-2)})x_{n-1}X^{(0,0,1,1)} + S_{m+n-1} + X^{(0,0,1,1)}S_{m+n-3}$
 $- X^{(0,0,1,1)}(qx_{-(m-2)}x_{n-1}X^{(0,0,1,1)} + S_{m+n-3})$
= $qx_{-m}x_{n-1}X^{(0,0,1,1)} + S_{m+n-1}$.

Thus, the proof is completed.

Remark 3.9. According to [\[6,](#page-20-17) Definition 4.15, Remark 4.18, Theorem 4.20], it follows that the equality in [\[6,](#page-20-17) Lemma 4.16 (a)] is exactly $x_{n+3}x_1 = q^{\frac{1}{2}}S_ny_1 + x_{n+2}x_2$ which is a special case in Theorem [3.8](#page-10-0) (2) for $m = 1$.

In order to study atomic bases in the rank two cluster algebras of affine types, Sherman and Zelevinsky $[23]$ introduced the *n*-th Chebyshev polynomials of the first kind $F_n(x)$. We modify the definition of $F_n(x)$ for $n \in \mathbb{Z}_{\geq 0}$ on X_δ as follows

$$
F_0(X_\delta) = 1, F_1(X_\delta) = X_\delta, F_2(X_\delta) = F_1(X_\delta)F_1(X_\delta) - 2X^{(0,0,1,1)},
$$

$$
F_{n+1}(X_\delta) = F_n(X_\delta)F_1(X_\delta) - X^{(0,0,1,1)}F_{n-1}(X_\delta) \text{ for } n \ge 2.
$$

In the following, we denote $F_n(X_\delta)$ by F_n for simplicity. According to the definition of F_n and the fact that $F_1 \in \mathcal{A}_q(Q)$, it is easy to see $F_n \in \mathcal{A}_q(Q)$ for any $n \geq 1$.

Lemma 3.10. In $\mathcal{A}_q(Q)$, we have that

(1)
$$
F_m x_0 = x_{-m} + q^{-\frac{2m-1}{2}} y_2 x_m
$$
, for $m \ge 1$;
\n(2) $F_m x_{-1} = x_{-(m+1)} + q^{-\frac{2m-1}{2}} X^{(0,0,1,2)} x_{m-1}$, for $m \ge 2$;
\n(3) $F_2 x_{-2} = x_{-4} + X^{(0,0,2,2)} x_0$, and $F_m x_{-2} = x_{-(m+2)} + q^{-\frac{2m-1}{2}} X^{(0,0,2,3)} x_{m-2}$, for $m \ge 3$.

Proof. (1) The case when $m = 1$ follows from Lemma [3.5](#page-9-0) (1). When $m = 2$, we have

$$
F_2x_0 = X_\delta^2 x_0 - 2X^{(0,0,1,1)}x_0
$$

= $X_\delta(x_{-1} + q^{-\frac{1}{2}}y_2x_1) - 2X^{(0,0,1,1)}x_0$
= $x_{-2} + X^{(0,0,1,1)}x_0 + q^{-\frac{3}{2}}y_2(x_2 + q^{\frac{1}{2}}y_1x_0) - 2X^{(0,0,1,1)}x_0$
= $x_{-2} + q^{-\frac{3}{2}}y_2x_2$.

When $m = 3$, we have

$$
F_3x_0 = (X_\delta F_2 - X^{(0,0,1,1)}X_\delta)x_0
$$

= $X_\delta(x_{-2} + q^{-\frac{3}{2}}y_2x_2) - X^{(0,0,1,1)}(x_{-1} + q^{-\frac{1}{2}}y_2x_1)$
= $x_{-3} + X^{(0,0,1,1)}x_{-1} + q^{-\frac{5}{2}}y_2(x_3 + X^{(0,0,1,1)}x_1) - X^{(0,0,1,1)}(x_{-1} + q^{-\frac{1}{2}}y_2x_1)$
= $x_{-3} + q^{-\frac{5}{2}}y_2x_3$.

Suppose that $F_m x_0 = x_{-m} + q^{-\frac{2m-1}{2}} y_2 x_m$. Then we have that

$$
F_{m+1}x_0 = (X_{\delta}F_m - X^{(0,0,1,1)}F_{m-1})x_0
$$

= $X_{\delta}(x_{-m} + q^{-\frac{2m-1}{2}}y_2x_m) - X^{(0,0,1,1)}(x_{-(m-1)} + q^{-\frac{2m-3}{2}}y_2x_{m-1})$
= $x_{-(m+1)} + X^{(0,0,1,1)}x_{-(m-1)} + q^{-\frac{2m+1}{2}}y_2(x_{m+1} + X^{(0,0,1,1)}x_{m-1})$
 $- X^{(0,0,1,1)}(x_{-(m-1)} + q^{-\frac{2m-3}{2}}y_2x_{m-1})$
= $x_{-(m+1)} + q^{-\frac{2m+1}{2}}y_2x_{m+1}.$

(2) When $m = 2$, we have

$$
F_2x_{-1} = X_{\delta}^2 x_{-1} - 2X^{(0,0,1,1)}x_{-1}
$$

= $X_{\delta}(x_{-2} + X^{(0,0,1,1)}x_0) - 2X^{(0,0,1,1)}x_{-1}$
= $x_{-3} + X^{(0,0,1,1)}x_{-1} + X^{(0,0,1,1)}(x_{-1} + q^{-\frac{1}{2}}y_2x_1) - 2X^{(0,0,1,1)}x_{-1}$
= $x_{-3} + q^{-\frac{3}{2}}X^{(0,0,1,2)}x_1$.

When $m = 3$, we have

$$
F_3x_{-1} = X_{\delta}F_2x_{-1} - X^{(0,0,1,1)}X_{\delta}x_{-1}
$$

= $X_{\delta}(x_{-3} + q^{-\frac{3}{2}}X^{(0,0,1,2)}x_1) - X^{(0,0,1,1)}(x_{-2} + X^{(0,0,1,1)}x_0)$
= $x_{-4} + X^{(0,0,1,1)}x_{-2} + q^{-\frac{5}{2}}X^{(0,0,1,2)}(x_2 + q^{\frac{1}{2}}y_1x_0) - X^{(0,0,1,1)}(x_{-2} + X^{(0,0,1,1)}x_0)$
= $x_{-4} + q^{-\frac{5}{2}}X^{(0,0,1,2)}x_2$.

Suppose that $F_m x_{-1} = x_{-(m+1)} + q^{-\frac{2m-1}{2}} X^{(0,0,1,2)} x_{m-1}$. Then we have that

$$
F_{m+1}x_{-1} = X_{\delta}F_m x_{-1} - X^{(0,0,1,1)}F_{m-1}x_{-1}
$$

= $X_{\delta}(x_{-(m+1)} + q^{-\frac{2m-1}{2}} X^{(0,0,1,2)} x_{m-1}) - X^{(0,0,1,1)}(x_{-m} + q^{-\frac{2m-3}{2}} X^{(0,0,1,2)} x_{m-2})$
= $x_{-(m+2)} + X^{(0,0,1,1)}x_{-m} + q^{-\frac{2m+1}{2}} X^{(0,0,1,2)}(x_m + -X^{(0,0,1,1)} x_{m-2})$
 $- X^{(0,0,1,1)}(x_{-m} + q^{-\frac{2m-3}{2}} X^{(0,0,1,2)} x_{m-2})$
= $x_{-(m+2)} + q^{-\frac{2m+1}{2}} X^{(0,0,1,2)} x_m.$

(3) The first equality can be calculated as follows

$$
F_2x_{-2} = X_{\delta}^2x_{-2} - 2X^{(0,0,1,1)}x_{-2}
$$

= $X_{\delta}(x_{-3} + X^{(0,0,1,1)}x_{-1}) - 2X^{(0,0,1,1)}x_{-2}$
= $x_{-4} + X^{(0,0,1,1)}x_{-2} + X^{(0,0,1,1)}(x_{-2} + X^{(0,0,1,1)}x_0) - 2X^{(0,0,1,1)}x_{-2}$
= $x_{-4} + X^{(0,0,2,2)}x_0$.

When $m = 3$, we have that

$$
F_3x_{-2} = X_{\delta}F_2x_{-2} - X^{(0,0,1,1)}X_{\delta}x_{-2}
$$

= $X_{\delta}(x_{-4} + X^{(0,0,2,2)}x_0) - X^{(0,0,1,1)}(x_{-3} + X^{(0,0,1,1)}x_{-1})$
= $x_{-5} + X^{(0,0,1,1)}x_{-3} + X^{(0,0,2,2)}(x_{-1} + q^{-\frac{1}{2}}y_2x_1) - X^{(0,0,1,1)}(x_{-3} + X^{(0,0,1,1)}x_{-1})$
= $x_{-5} + q^{-\frac{5}{2}}X^{(0,0,2,3)}x_1$.

When $m = 4$, we have that

$$
F_4x_{-2} = X_{\delta}F_3x_{-2} - X^{(0,0,1,1)}F_2x_{-2}
$$

= $X_{\delta}(x_{-5} + q^{-\frac{5}{2}}X^{(0,0,2,3)}x_1) - X^{(0,0,1,1)}(x_{-4} + X^{(0,0,2,2)}x_0)$
= $x_{-6} + X^{(0,0,1,1)}x_{-4} + q^{-\frac{7}{2}}X^{(0,0,2,3)}(x_2 + q^{\frac{1}{2}}y_1x_0) - X^{(0,0,1,1)}(x_{-4} + X^{(0,0,2,2)}x_0)$
= $x_{-6} + q^{-\frac{7}{2}}X^{(0,0,2,3)}x_2$.

Suppose that $F_m x_{-2} = x_{-(m+2)} + q^{-\frac{2m-1}{2}} X^{(0,0,2,3)} x_{m-2}$. Then we have that

$$
F_{m+1}x_{-2} = X_{\delta}F_m x_{-2} - X^{(0,0,1,1)}F_{m-1}x_{-2}
$$

= $X_{\delta}(x_{-(m+2)} + q^{-\frac{2m-1}{2}} X^{(0,0,2,3)} x_{m-2}) - X^{(0,0,1,1)}(x_{-(m+1)} + q^{-\frac{2m-3}{2}} X^{(0,0,2,3)} x_{m-3})$
= $x_{-(m+3)} + X^{(0,0,1,1)} x_{-(m+1)} + q^{-\frac{2m+1}{2}} X^{(0,0,2,3)}(x_{m-1} + X^{(0,0,1,1)} x_{m-3})$
 $- X^{(0,0,1,1)}(x_{-(m+1)} + q^{-\frac{2m-3}{2}} X^{(0,0,2,3)} x_{m-3})$
= $x_{-(m+3)} + q^{-\frac{2m+1}{2}} X^{(0,0,2,3)} x_{m-1}.$

Hence, the proof is finished. \Box

Similarly, we have the following results.

Lemma 3.11. In $A_q(Q)$, we have that

(1) $x_1 F_m = x_{m+1} + q^{-\frac{2m-1}{2}} x_{-(m-1)} y_1$, for $m \ge 1$; (2) $x_2 F_m = x_{m+2} + q^{-\frac{2m-1}{2}} x_{-(m-2)} X^{(0,0,2,1)}$, for $m \ge 2$.

We can now prove multiplication formulas between the modified Chebyshev polynomials of the first kind on X_{δ} and cluster variables.

Theorem 3.12. In $A_q(Q)$, we have that

(1)

$$
F_mx_{-n} = \begin{cases} x_{-(n+m)} + X^{(0,0,m,m)}x_{-(n-m)} & \text{for } 1 \le m \le n; \\ x_{-(n+m)} + q^{-\frac{2m-1}{2}} X^{(0,0,n,n+1)} x_{m-n} & \text{for } n \ge 0, n+1 \le m. \end{cases}
$$

$$
(2)
$$

$$
x_n F_m = \begin{cases} x_{n+m} + x_{n-m} X^{(0,0,m,m)} & \text{for } 1 \le m < n; \\ x_{n+m} + q^{-\frac{2m-1}{2}} x_{-(m-n)} X^{(0,0,n,n-1)} & \text{for } 1 \le n \le m. \end{cases}
$$

Proof. (1) The cases when $n = 0, 1, 2$ follow from Lemmas [3.5](#page-9-0) and [3.10.](#page-13-0) Assume that $n \geq 2$, then we have $X_{\delta}x_{-(n+1)} = x_{-(n+2)} + X^{(0,0,1,1)}x_{-n}$, and

$$
F_2x_{-(n+1)} = X_{\delta}^2 x_{-(n+1)} - 2X^{(0,0,1,1)}x_{-(n+1)}
$$

= $X_{\delta}(x_{-(n+2)} + X^{(0,0,1,1)}x_{-n}) - 2X^{(0,0,1,1)}x_{-(n+1)}$
= $x_{-(n+3)} + X^{(0,0,1,1)}x_{-(n+1)} + X^{(0,0,1,1)}(x_{-(n+1)} + X^{(0,0,1,1)}x_{-(n-1)}) - 2X^{(0,0,1,1)}x_{-(n+1)}$
= $x_{-(n+3)} + X^{(0,0,2,2)}x_{-(n-1)}$.

Suppose that $F_k x_{-(n+1)} = x_{-(n+1+k)} + X^{(0,0,k,k)} x_{-(n+1-k)}$, for $k \leq n$, then we have that

$$
F_{k+1}x_{-(n+1)} = (X_{\delta}F_k - X^{(0,0,1,1)}F_{k-1})x_{-(n+1)}
$$

= $X_{\delta}(x_{-(n+1+k)} + X^{(0,0,k,k)}x_{-(n+1-k)}) - X^{(0,0,1,1)}(x_{-(n+k)} + X^{(0,0,k-1,k-1)}x_{-(n+2-k)})$
= $x_{-(n+2+k)} + X^{(0,0,1,1)}x_{-(n+k)} + X^{(0,0,k,k)}(x_{-(n+2-k)} + X^{(0,0,1,1)}x_{-(n-k)})$

$$
- X^{(0,0,1,1)}(x_{-(n+k)} + X^{(0,0,k-1,k-1)}x_{-(n+2-k)})
$$

= $x_{-(n+2+k)} + X^{(0,0,k+1,k+1)}x_{-(n-k)};$

Thus,

$$
F_{n+2}x_{-(n+1)} = (X_{\delta}F_{n+1} - X^{(0,0,1,1)}F_n)x_{-(n+1)}
$$

= $X_{\delta}(x_{-(2n+2)} + X^{(0,0,n+1,n+1)}x_0) - X^{(0,0,1,1)}(x_{-(2n+1)} + X^{(0,0,n,n)}x_{-1})$
= $x_{-(2n+3)} + X^{(0,0,1,1)}x_{-(2n+1)} + X^{(0,0,n+1,n+1)}(x_{-1} + q^{-\frac{1}{2}}y_2x_1)$
 $- X^{(0,0,1,1)}(x_{-(2n+1)} + X^{(0,0,n,n)}x_{-1})$
= $x_{-(2n+3)} + q^{-\frac{2n+3}{2}}X^{(0,0,n+1,n+2)}x_1;$

Similarly, we have that

$$
F_{n+3}x_{-(n+1)} = x_{-(2n+4)} + q^{-\frac{2n+5}{2}} X^{(0,0,n+1,n+2)} x_2;
$$

$$
F_{n+4}x_{-(n+1)} = x_{-(2n+5)} + q^{-\frac{2n+7}{2}} X^{(0,0,n+1,n+2)} x_3.
$$

Now, for $m \geq n+2$, suppose that

$$
F_m x_{-(n+1)} = x_{-(m+n+1)} + q^{-\frac{2m-1}{2}} X^{(0,0,n+1,n+2)} x_{m-n-1},
$$

then we have that

$$
F_{m+1}x_{-(n+1)} = (X_{\delta}F_m - X^{(0,0,1,1)}F_{m-1})x_{-(n+1)}
$$

= $X_{\delta}(x_{-(m+n+1)} + q^{-\frac{2m-1}{2}}X^{(0,0,n+1,n+2)}x_{m-n-1})$
 $- X^{(0,0,1,1)}(x_{-(m+n)} + q^{-\frac{2m-3}{2}}X^{(0,0,n+1,n+2)}x_{m-n-2})$
= $x_{-(m+n+2)} + X^{(0,0,1,1)}x_{-(m+n)} + q^{-\frac{2m+1}{2}}X^{(0,0,n+1,n+2)}(x_{m-n} + X^{(0,0,1,1)}x_{m-n-2})$
 $- X^{(0,0,1,1)}(x_{-(m+n)} + q^{-\frac{2m-3}{2}}X^{(0,0,n+1,n+2)}x_{m-n-2})$
= $x_{-(m+n+2)} + q^{-\frac{2m+1}{2}}X^{(0,0,n+1,n+2)}x_{m-n}.$

(2) The proof is similar to (1), where we need to use Lemma [3.11](#page-16-0) and Theorem $3.6.$

Remark 3.13. Since all coefficients appear in multiplication formulas are Laurent polynomials of q which are independent of the choice of the finite field \mathbb{F}_q , generically all multiplication formulas also hold in $\mathcal{A}_{q}(Q)$ if we replace q by the indeterminate q.

4. BAR-INVARIANT POSITIVE $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm},y_2^{\pm}]$ -BASES OF $\mathcal{A}_{\mathfrak{q}}(Q)$

In this section, we use multiplication formulas established in Section 3 to construct two positive $\mathbb{Z}[q^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -bases.

Definition 4.1. An element in $\mathcal{A}_{q}(Q)$ is called positive if the coefficients of its Laurent expansion associated to any cluster belong to $\mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}].$

Definition 4.2. A basis of $\mathcal{A}_{\mathfrak{q}}(Q)$ is called a positive $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -basis if its structure constants belong to $\mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}].$

Denote the following

 $S = \{\text{cluster monomials}\}\sqcup \{S_n|n \geq 1\}.$

Theorem 4.3. The set S is a bar-invariant positive $\mathbb{Z}[q^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -basis of $\mathcal{A}_q(Q)$.

Proof. According to Proposition [3.3,](#page-8-1) Theorem [3.6,](#page-9-1) Theorem [3.8,](#page-10-0) Lemma [3.1](#page-7-0) and Lemma [3.2,](#page-7-1) we can deduce that the $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -combinations of the elements in S is $\mathcal{A}_{\mathfrak{q}}(Q)$.

In order to prove the $\mathbb{Z}[q^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -independency of these elements, we define a partial order \leq on \mathbb{Z}^2 as follows: $(r_1, r_2) \leq (s_1, s_2)$ if $r_i \leq s_i$ for $1 \leq i \leq 2$. Moreover if $r_i < s_i$ for some i, we will write $(r_1, r_2) < (s_1, s_2)$. According to Proposition [3.3,](#page-8-1) every element s_n has a minimal non-zero term $a_n X^{-\underline{dim}R(n)}$ where $a_n \in \mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$. Ac-cording to Theorem [3.6,](#page-9-1) every element x_m has a minimal non-zero term $b_m X^{-\underline{dim}V(m)}$ where $b_m \in \mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$. It follows from [\[23,](#page-20-1) Proposition 3.1], there exists a bijection between the set of all minimal non-zero terms in the elements in S and \mathbb{Z}^2 , which implies that the elements in S are $\mathbb{Z}[{\bf q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -independent.

By Lemma [3.2](#page-7-1) and the fact that cluster monomials are bar-invariant, we know that any element in S is bar-invariant. Again from Proposition [3.3,](#page-8-1) Theorem [3.6,](#page-9-1) Theorem [3.8,](#page-10-0) Lemma [3.1](#page-7-0) and Lemma [3.2,](#page-7-1) we can deduce that the structure constants belong to $\mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$].

Remark 4.4. Using the same arguments as [\[19,](#page-20-19) Corollary 3.3.10] [\[21,](#page-20-20) Corollary 8.3.3, it follows that all elements in S are positive.

Definition 4.5. A basis of $\mathcal{A}_{\mathfrak{q}}(Q)$ is called the atomic $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -basis if positive elements in $\mathcal{A}_{\mathfrak{q}}(Q)$ are exactly $\mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -combinations of these basis elements.

It follows from the definition that the atomic basis, if exists, must be a positive basis.

Denote the following

 $\mathcal{B} = \{\text{cluster monomials}\}\sqcup \{F_n|n \geq 1\}.$

Theorem 4.6. The set B is the bar-invariant atomic $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -basis of $\mathcal{A}_{\mathfrak{q}}(Q)$.

Proof. By Theorem [4.3](#page-18-0) and the relation between S_n and F_n , we have that $\mathcal B$ is a $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -basis. By applying similar discussions as in Theorem [4.3,](#page-18-0)we can show that the elements in β are bar-invariant. Note that cluster variables are positive (see also Remark [4.4\)](#page-18-1), then by Theorem [3.12,](#page-16-1) we have that F_n is a positive element for $n \geq 1$. Thus, all elements in β are positive.

Let Y be a positive element in $\mathcal{A}_{\mathfrak{q}}(Q)$, then $Y = \sum_{b \in \mathcal{B}} \lambda_b b$ for some $\lambda_b \in$ $\mathbb{Z}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}].$ We need to prove $\lambda_b \in \mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}].$ Firstly, we consider the case when the element b is a cluster monomial. Without loss of generality, suppose that b is a cluster monomial in the cluster $\{x_m, x_{m+1}\}$. If b appears in the $\mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$ -expansion of some other basis element b' of $\mathcal B$ associated to the cluster ${x_m, x_{m+1}}$, then $b|_{q=1,y_1=1,y_2=1}$ appears in the $\mathbb{Z}_{\geq 0}$ -expansion of some other basis element $b'|_{q=1,y_1=1,y_2=1}$ associated to the cluster $\{x_m, x_{m+1}\}\vert_{q=1,y_1=1,y_2=1}$ due to the positivity of the elements in \mathcal{B} . However this can't happen in the corresponding classical cluster algebra, for more details, refer to [\[14,](#page-20-9) Section 4, Proof of (B3) in Theorem 1.2 or [\[23,](#page-20-1) Section 5, Proof of (5.10)]. Thus by the assumption that Y is positive associated to the cluster $\{x_m, x_{m+1}\}\$, we have $\lambda_b \in \mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$. Now, we consider the case when $b = F_l$ for some $l \geq 1$. By using the same discussions and references as above, we can find a Laurent monomial Y_l in certain cluster expansion of F_l with coefficients $\mathfrak{q}^{\frac{l_0}{2}}y_1^{l_1}y_2^{l_2}$ for some $l_0, l_1, l_2 \in \mathbb{Z}$, but Y_l doesn't appear in this cluster expansion of any other basis element in the above sum terms. Thus we have $\lambda_b \mathfrak{q}^{\frac{l_0}{2}} y_1^{l_1} y_2^{l_2} \in \mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}] [y_1^{\pm}, y_2^{\pm}]$ by using the assumption that Y is positive associated to this cluster. It follows that $\lambda_b \in \mathbb{Z}_{\geq 0}[\mathfrak{q}^{\pm \frac{1}{2}}][y_1^{\pm}, y_2^{\pm}]$. The proof is completed. \Box

Remark 4.7. By specializing $q = 1$ in Theorems [4.3](#page-18-0) and [4.6,](#page-18-2) we obtain a positive integral basis and the atomic basis of the classical cluster algebra associated to the Kronecker quiver with principal coefficients. Note that Dupont constructed these bases [\[15,](#page-20-11) Theorem 7.3, Theorem 7.4] in the classical Kronecker cluster algebra with opposite principal coefficients.

ACKNOWLEDGMENTS

Ming Ding was supported by NSF of China (No. 11771217) and Fan Xu was supported by NSF of China (No. 12031007). We thank the referees for their very helpful comments and suggestions.

REFERENCES

- [1] L. Bai, X. Chen, M. Ding and F. Xu, Cluster multiplication theorem in the quantum cluster algebra of type $A_2^{(2)}$ and the triangular basis, J. Algebra 533 (2019), 106-141.
- [2] A. Buan, R. Marsh, M. Reineke, I. Reiten and G. Todorov, Tilting theory and cluster combinatorics, Adv. Math. **204** (2006), 572-618.
- [3] A. Berenstein and A. Zelevinsky, *Quantum cluster algebras*, Adv. Math. **195** (2005), 405–455.
- [4] P. Caldero and B. Keller, From triangulated categories to cluster algebras, Invent. Math. 172 (2008) 169-211.
- [5] P. Caldero and A. Zelevinsky, Laurent expansions in cluster algebras via quiver representations, Mosc. Math. J. 6 (2006), 411–429.
- [6] I. Canaki and P. Lampe, An expansion formula for type A and Kronecker quantum cluster algebras, Journal of Combinatorial Theory, Series A 171 (2020) 105132.
- [7] G. Cerulli Irelli, *Cluster algebras of type* $A_2^{(1)}$, Algebr. Represent. Theory 15 (2012), no. 5, 977–1021.
- [8] G. Cerulli Irelli, Positivity in skew-symmetric cluster algebras of finite type, Preprint, arXiv:1102.3050.
- [9] X. Chen, M. Ding and H.Zhang, Acyclic quantum cluster algebras via derived Hall algebras, Preprint, arXiv:2108.03558 [math.RT].
- [10] M. Ding and F. Xu, Bases of the quantum cluster algebra of the Kronecker quiver, Acta Math. Sin. (Engl. Ser.) 28 (2012), no. 6, 1169–1178.
- [11] M. Ding and F. Xu, A quantum analogue of generic bases for affine cluster algebras, Sci. China Math. 55 (2012), no. 10, 2045–2066.
- [12] M. Ding, F. Xu and X. Chen, Atomic bases of quantum cluster algebras of type $\tilde{A}_{2n-1,1}$, Journal of Algebra 590 (2022) 1-25.
- [13] V. Dlab and C. M. Ringel, *Indecomposable representations of graphs and algebras*, Mem. Amer. Math. Soc., vol. 173, Amer. Math. Soc., Providence, RI, 1976.
- [14] G. Dupont and H. Thomas, Atomic bases of cluster algebras of types A and A , Proc. Lond. Math. Soc. **107** (2013), no. 4, 825–850.
- [15] G. Dupont, Quantized Chebyshev polynomials and cluster characters with coefficients, J. Algebraic Combin. 31 (2010), no. 4, 501-532.
- [16] S. Fomin and A. Zelevinsky, Cluster algebras. I. Foundations, J. Amer. Math. Soc. 15 (2002), no. 2, 497–529.
- [17] A. Hubery, Acyclic cluster algebras via Ringel-Hall algebras, preprint (2005).
- [18] A. Hubery, *Hall polynomials for affine quivers*, Represent. Theory 14 (2010), 355-378.
- [19] Y. Kimura and F. Qin, Graded quiver varieties, quantum cluster algebras and dual canonical *basis*, Adv. Math. **262** (2014), 261-312.
- [20] F. Qin, Quantum cluster variables via Serre polynomials. With an appendix by Bernhard Keller, J. Reine Angew. Math. 668 (2012), 149–190.
- [21] F. Qin, t-analog of q-characters, bases of quantum cluster algebras, and a correction technique, Int. Math. Res. Not. IMRN 2014, no. 22, 6175-6232.
- [22] D. Rupel, *Quantum cluster characters for valued quivers*, Trans. Amer. Math. Soc. 367(10), (2015), 7061–7102
- [23] P. Sherman and A. Zelevinsky, Positivity and canonical bases in rank 2 cluster algebras of finite and affine types, Mosc. Math. J. 4 (2004), no. 4, 947–974, 982.
- [24] J. Xiao and F. Xu, Green's formula with C^{*}-action and Caldero-Keller's formula for cluster algebras, in: Representation Theory of Algebraic Groups and Quantum Groups, in: Progr. Math., vol. 284, Birkhauser/Springer, New York, 2010: 313-348.
- [25] F. Xu, On the cluster multiplication theorem for acyclic cluster algebras, Trans. Amer. Math. Soc. 362(2) (2010), 753-776.
- [26] A. Zelevinsky, Quantum Cluster Algebras: Oberwolfach talk, February 2005, math.QA/0502260.

School of Mathematics and Information Science, Guangzhou University, Guangzhou, P. R. China

 $Email \ address: m\text{-ding}04@mails.tsinghua.edu.cn (M.Ding)$

Department of Mathematical Sciences, Tsinghua University, Beijing 100084, P. R. China

Email address: fanxu@mail.tsinghua.edu.cn (F.Xu)

Department of Mathematics, University of Wisconsin-Whitewater, 800 W. Main Street, Whitewater, WI.53190. USA

Email address: chenx@uww.edu (X.Chen)