

GLOBALLY CONFORMALLY KÄHLER EINSTEIN METRICS ON CERTAIN HOLOMORPHIC BUNDLES

ZHIMING FENG

ABSTRACT. The subject of this paper is the explicit momentum construction of complete Einstein metrics by ODE methods. Using the Calabi ansatz, further generalized by Hwang-Singer, we show that there are non-trivial complete conformally Kähler Einstein metrics on certain Hermitian holomorphic vector bundles and their subbundles over complete Kähler-Einstein manifolds. In special cases, we give the explicit expressions of these metrics. These examples show that there is a compact Kähler manifold M and its subvariety N whose codimension is greater than 1 such that there is a complete conformally Kähler Einstein metric on $M - N$.

1. INTRODUCTION AND MAIN RESULTS

Let M be a complex m -dimensional Kähler manifold endowed with a Kähler metric g with respect to an integrable almost complex structure J , and f be a positive smooth function f on M . A Hermitian metric $\tilde{g} = \frac{1}{f^2}g$ on (M, J) is called a globally conformally Kähler metric.

If there is a constant μ such that the Ricci tensor $\text{Ric}_{\tilde{g}}$ of the metric \tilde{g} satisfies $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$, then the Hermitian metric \tilde{g} on (M, J) is called a conformally Kähler Einstein metric; if f also satisfies $df \wedge d\Delta f = 0$, then the metric \tilde{g} is called a strongly conformally Kähler Einstein metric; this is automatic if $m \geq 3$.

If $\text{Ric}_{\tilde{g}}(J\cdot, J\cdot) = \text{Ric}_{\tilde{g}}(\cdot, \cdot)$ and the scalar curvature $s_{\tilde{g}}$ of \tilde{g} is constant, then the metric \tilde{g} on (M, J) is called a conformally Kähler, Einstein-Maxwell metric (cKEM metric for short), see [27, 28]. Note that $\text{Ric}_{\tilde{g}}(J\cdot, J\cdot) = \text{Ric}_{\tilde{g}}(\cdot, \cdot)$ if and only if $\bar{\partial}(\text{grad}^{1,0} f) = \bar{\partial}(g^{j\bar{q}} f_{\bar{q}} \partial_j) = 0$. Obviously, a conformally Kähler Einstein metric is a cKEM metric. When f is constant on M , a cKEM metric is a constant scalar curvature Kähler (cscK for short) metric, so a cKEM metric with nonconstant f is referred to as non-trivial.

For the case of compact, so far, not many examples of non-trivial cKEM metrics are known. The conformally Kähler Einstein metrics constructed by Page [29] on the one-point-blow-up of $\mathbb{C}\mathbb{P}^2$, by Chen-LeBrun-Weber [9] on the two-point-blow-up of $\mathbb{C}\mathbb{P}^2$, by Apostolov-Calderbank-Gauduchon [3, 4] on 4-orbifolds and by Bérard-Bergery [7] on \mathbb{P}^1 -bundles over Fano Kähler-Einstein manifolds. Non-Einstein cKEM examples are given by LeBrun [27, 28] on $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ and the one-point-blow-up of $\mathbb{C}\mathbb{P}^2$, by Koca-Tønnesen-Friedman [23] on ruled surfaces of higher genus, and by Futaki-Ono [17] on $\mathbb{C}\mathbb{P}^1 \times M$ where M is a compact constant scalar curvature Kähler manifold of arbitrary dimension. For more research on cKEM metrics, please refer to Apostolov-Maschler [5], Apostolov-Maschler-Tønnesen-Friedman [6], Futaki-Ono [18] and Lahdili [24, 25, 26].

In [11], Derdzinski-Maschler gave the local classification of strongly conformally Kähler Einstein metrics. In [2], Apostolov-Calderbank-Gauduchon also gave the local classification of strongly conformally Kähler Einstein metrics by using the more general Hamiltonian 2-forms method. That

2010 *Mathematics Subject Classification.* 32Q15, 53C55.

Key words and phrases. Einstein metrics, Conformally Kähler Einstein metrics, CKEM metrics, Holomorphic bundles.

is, if $\tilde{g} = \frac{1}{f^2}g$ is a strongly conformally Kähler Einstein metric, then the metric g can be expressed as

$$\begin{aligned} g &= zg_0 + \frac{z^{m-1}}{F(z)}dz^2 + \frac{F(z)}{z^{m-1}}\theta^2, \\ \omega &= dd^c \int^z \frac{u^m}{F(u)}du = z\omega_0 + dz \wedge \theta, \quad d\theta = \omega_0, \quad Jdz = \frac{F(z)}{z^{m-1}}\theta, \quad J\theta = -\frac{z^{m-1}}{F(z)}dz, \\ \rho_0 &= c\omega_0, \end{aligned}$$

where c is a constant, $f = qz + p$ and

$$(1.1) \quad F''(z) - \left(\frac{2(m-1)q}{qz+p} + \frac{m}{z} \right) F'(z) + \frac{2m(m-1)q}{z(qz+p)} F(z) + 2cz^{m-2} = 0.$$

In [12, 13], Derdzinski-Maschler gave the classification compact conformally Einstein Kähler manifolds M , namely M are the total spaces of holomorphic $\mathbb{C}\mathbb{P}^1$ bundles over compact Kähler-Einstein manifolds.

For the case of non-compact, in [15], we study the existence of cKEM metrics on rotationally invariant domains in \mathbb{C}^d . For non-compact examples of conformally Kähler Einstein metrics, see [10, 1].

Let N be the generic norm of an irreducible bounded symmetric domain Ω , for $\lambda > 0$ and $r \in \mathbb{N}_+ = \mathbb{N} - \{0\}$, the Cartan-Hartogs domain defined by

$$M = \left\{ (z, w) \in \Omega \times \mathbb{C}^r : e^{\lambda\phi(z)} \|w\|^2 < 1 \right\}, \quad \phi(z) = -\log N(z, z),$$

its holomorphic invariant Kähler metrics can be expressed as

$$\omega = \sqrt{-1}\partial\bar{\partial}(\nu\phi(z) + F(t)), \quad t = \lambda\phi(z) + \log \|w\|^2, \quad \nu > 0.$$

So holomorphic invariant conformally Kähler metrics \tilde{g} associated with Kähler forms $\tilde{\omega}$ on the Cartan-Hartogs domain M can be expressed as

$$\tilde{\omega} = \frac{\sqrt{-1}}{f^2(t)}\partial\bar{\partial}(\nu\phi(z) + F(t)).$$

In this paper, using the Calabi ansatz and the momentum construction [22], we construct non-trivial complete conformally Kähler Einstein metrics \tilde{g} associated with Kähler forms $\tilde{\omega}$ on certain holomorphic bundles, which are locally expressed as

$$M = \left\{ (z, w) \in \Omega \times \mathbb{C}^r : e^{\lambda\phi(z)} \|w\|^2 \in \mathcal{I} \right\} \quad (\mathcal{I} \text{ are intervals on } \mathbb{R})$$

and

$$\tilde{\omega} = \frac{\sqrt{-1}}{f^2(t)}\partial\bar{\partial}(\nu\phi(z) + F(t)), \quad t = \lambda\phi(z) + \log \|w\|^2, \quad \nu \geq 0.$$

The following Theorem 1.1 is the main result of this paper.

Theorem 1.1. *Let (L_0, h_0, Π_0) be a Hermitian holomorphic line bundle over a complete Kähler manifold (M_0, g_0) of complex dimension d such that $2\pi c_1(L_0, h_0) = -\lambda\omega_0$, where ω_0 denotes the Kähler form associated with the Kähler metric g_0 , $c_1(L_0, h_0)$ denotes the curvature of the Chern connection on the Hermitian holomorphic bundle (L_0, h_0) , and $\Pi_0 : L_0 \rightarrow M_0$ is the bundle projection. The curvature $c_1(L_0, h_0)$ is given by*

$$c_1(L_0, h_0) = -\frac{\sqrt{-1}}{2\pi}\partial\bar{\partial}\log h_0(\sigma(z), \sigma(z))$$

for a trivializing holomorphic section $\sigma : U \rightarrow L_0 \setminus \{0\}$.

Let E be the direct sum of r copies of L_0 , i.e. $E = L_0^{\oplus r}$ with an associated hermitian metric denoted by h . The projective bundle $\mathbb{P}(E \oplus 1)$ can be viewed as a compactification of E , where 1 stands for the trivial line bundle $M_0 \times \mathbb{C}$. The projection of the vector bundle E to M_0 denoted by $\Pi : E \rightarrow M_0$.

Let M be one of the following sets:

$$E = L_0^{\oplus r}, E^* = \{u \in E : 0 < h(u, u)\}, E - \{u \in E : h(u, u) \leq 1\},$$

$$\mathbb{B}_{L_0}^r = \{u \in E : h(u, u) < 1\}, \mathbb{B}_{L_0}^{r*} = \{u \in E : 0 < h(u, u) < 1\},$$

$$\mathbb{P}(E \oplus 1) - M_0 \text{ and } \mathbb{P}(E \oplus 1) - \{u \in E : h(u, u) \leq 1\}.$$

Let g_F be a Kähler metric on M associated with the Kähler form

$$\omega_F(u) = \nu (\Pi^* \omega_0)(u) + \sqrt{-1} \partial \bar{\partial} F(\log h(u, u)) \quad (u \in E, \nu = 0 \text{ or } 1).$$

(ω_F can be extended across $M - E$ if $M = \mathbb{P}(E \oplus 1)$, the extension of ω_F on M is still expressed as ω_F).

Set

$$t = \log h(u, u), \quad x = F'(t), \quad \varphi(x) = F''(t), \quad n = d + r,$$

$$x_0 = \inf\{F'(t)\}, \quad x_1 = \sup\{F'(t)\}, \quad f = aF'(t) + b = ax + b > 0, \quad a\lambda \neq 0.$$

Let $\tilde{g} = \frac{1}{f^2} g_F$ be a conformally Kähler metric on M . The following conclusions hold.

(I) For $\nu = 1$ and $b = 1$, \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) M if and only if

$$(1.2) \quad \begin{cases} \text{Ric}_{g_0} = (\lambda + \mu - 2a) g_0, \\ \varphi(x) = \frac{(1+ax)^{2d+1}(a\lambda x + 2a - \lambda)}{(1+\lambda x)^d} \int_0^x \frac{(1+\lambda u)^d (\mu(1+\lambda u) - (\lambda + \mu - 2a)(1+au)^2)}{(1+au)^{2d+2}(a\lambda u + 2a - \lambda)^2} du, \quad x_0 \leq x < x_1, \\ \mu \leq 0, a \leq \lambda, a < 0, r = 1, \end{cases}$$

here $x_0 = 0$, $x_1 = -\frac{1}{a}$, $M = \mathbb{B}_{L_0}^r$ for $\mu < 0$, and $M = L_0^{\oplus r}$ for $\mu = 0$.

(II)

(II - 1) \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $\mathbb{B}_{L_0}^{r*}$ if and only if

$$(1.3) \quad \begin{cases} \text{Ric}_{g_0} = \mu \frac{\nu + \lambda x_0}{(b + ax_0)^2} g_0, \\ \varphi(x) = \mu \frac{(b+ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_0}^x \frac{(\nu + \lambda u)^d \left((\nu + \lambda u) - \frac{\nu + \lambda x_0}{(b + ax_0)^2} (b + au)^2 \right)}{(b + au)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ \mu < 0, a < 0, r = 1, \end{cases}$$

where

$$\begin{cases} -\infty \leq x_0 < x_1 = -\frac{b}{a} \text{ for } \lambda < 0 \text{ and } -\frac{b}{a} \leq -\frac{\nu}{\lambda}, \\ -\frac{\nu}{\lambda} \leq x_0 < x_1 = -\frac{b}{a} \text{ for } \lambda > 0 \text{ and } -\frac{\nu}{\lambda} < -\frac{b}{a}. \end{cases}$$

(II – 2) \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $M = E - \{u \in E : h(u, u) \leq 1\}$ if and only if

$$(1.4) \quad \begin{cases} \text{Ric}_{g_0} = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} g_0, \\ \varphi(x) = \mu \frac{(b + ax)^{2d+1} (a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d \left((\nu + \lambda u) - \frac{\nu + \lambda x_1}{(b + ax_1)^2} (b + au)^2 \right)}{(b + au)^{2d+2} (a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ \mu < 0, a > 0, r = 1, \end{cases}$$

where

$$\begin{cases} -\frac{b}{a} = x_0 < x_1 \leq +\infty \text{ for } \lambda > 0 \text{ and } -\frac{b}{a} \geq -\frac{\nu}{\lambda}, \\ -\frac{b}{a} = x_0 < x_1 \leq -\frac{\nu}{\lambda} \text{ for } \lambda < 0 \text{ and } -\frac{\nu}{\lambda} > -\frac{b}{a}. \end{cases}$$

(III) Let $r = 1$.

(III – 1) \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $M = \mathbb{P}(E \oplus 1) - M_0$ if and only if

$$(1.5) \quad \begin{cases} \text{Ric}_{g_0} = \gamma g_0, \\ \gamma = \frac{a\lambda x_1 + 2a\nu - b\lambda}{ax_1 + b}, \\ \varphi(x) = -\gamma \frac{(a\lambda x + 2a\nu - b\lambda)(ax + b)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d}{(au + b)^{2d} (a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ a > 0, \mu = 0, \end{cases}$$

where

$$\begin{cases} -\frac{b}{a} = x_0 < x_1 < +\infty \text{ for } \lambda > 0 \text{ and } -\frac{b}{a} \geq -\frac{\nu}{\lambda}, \\ -\frac{b}{a} = x_0 < x_1 \leq -\frac{\nu}{\lambda} \text{ for } \lambda < 0 \text{ and } -\frac{\nu}{\lambda} > -\frac{b}{a}. \end{cases}$$

(III – 2) \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $M = \mathbb{P}(E \oplus 1) - \{u \in E : h(u, u) \leq 1\}$ if and only if

$$(1.6) \quad \begin{cases} \text{Ric}_{g_0} = \gamma g_0, \\ \gamma = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{ax_1 + b}, \\ \varphi(x) = \frac{(a\lambda x + 2a\nu - b\lambda)(ax + b)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(au + b)^{2d} (a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ a > 0, \mu < 0, \end{cases}$$

where

$$\begin{cases} -\frac{b}{a} = x_0 < x_1 < +\infty \text{ for } \lambda > 0 \text{ and } -\frac{b}{a} \geq -\frac{\nu}{\lambda}, \\ -\frac{b}{a} = x_0 < x_1 \leq -\frac{\nu}{\lambda} \text{ for } \lambda < 0 \text{ and } -\frac{\nu}{\lambda} > -\frac{b}{a}. \end{cases}$$

(IV) Let $r > 1$.

(IV - 1) \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $M = \mathbb{P}(E \oplus 1) - M_0$ if and only if

$$(1.7) \quad \begin{cases} \text{Ric}_{g_0} = r\lambda g_0, \\ \varphi(x) = -r \frac{(ax+b)^{2n-1}(ax-b)}{x^{n-1}} \int_{x_1}^x \frac{u^{n-1}}{(au+b)^{2n-2}(au-b)^2} du, \quad -\frac{b}{a} = x_0 < x < x_1 = -\frac{(r+1)b}{(r-1)a}, \\ a > 0, b < 0, \lambda > 0, \mu = \nu = 0. \end{cases}$$

(IV - 2) \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $M = \mathbb{P}(E \oplus 1) - \{u \in E : h(u, u) \leq 1\}$ if and only if

$$(1.8) \quad \begin{cases} \text{Ric}_{g_0} = r\lambda g_0, \\ \mu = \frac{(ax_1+b)(r(ax_1+b)-(ax_1-b))}{x_1} < 0, \\ \varphi(x) = \frac{(ax+b)^{2n-1}(ax-b)}{x^{n-1}} \int_{x_1}^x \frac{u^{n-1}(\mu u - r(b+au)^2)}{(b+au)^{2n}(au-b)^2} du, \quad -\frac{b}{a} = x_0 < x < x_1 < -\frac{(r+1)b}{(r-1)a}, \\ a > 0, b < 0, \lambda > 0, \nu = 0. \end{cases}$$

According to Theorem 1.1, we have Corollary 1.1, Corollary 1.2, Corollary 1.3, Corollary 1.4 and Corollary 1.5 for specific parameters a, b, λ, μ and ν .

Corollary 1.1. *Under the situation of Theorem 1.1, let $a = \lambda = -1$ and $b = \nu = 1$, then \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on M if and only if*

$$(1.9) \quad \begin{cases} \text{Ric}_{g_0} = (\mu + 1) g_0, \\ \varphi(x) = \frac{\mu+1}{d+1} (1-x - (1-x)^{d+2}) - \frac{\mu}{d+2} (1 - (1-x)^{d+2}), \quad 0 \leq x < 1, \\ \mu \leq 0, r = 1, \end{cases}$$

here $M = \mathbb{B}_{L_0}^r$ for $\mu < 0$, and $M = L_0^{\oplus r}$ for $\mu = 0$.

In particular, for $\mu = 0$, (1.9) is simplified as

$$(1.10) \quad \begin{cases} \text{Ric}_{g_0} = g_0, \\ \varphi(x) = \frac{1}{d+1} (1-x - (1-x)^{d+2}), \quad 0 \leq x < 1, \\ r = 1. \end{cases}$$

For $\mu = -1$, (1.9) is simplified as

$$(1.11) \quad \begin{cases} \text{Ric}_{g_0} = 0, \\ \varphi(x) = \frac{1}{d+2} (1 - (1-x)^{d+2}), \quad 0 \leq x < 1, \\ r = 1. \end{cases}$$

For $\mu = -(d+2)$, (1.9) is simplified as

$$(1.12) \quad \begin{cases} \text{Ric}_{g_0} = -(d+1) g_0, \\ \varphi(x) = x, 0 \leq x < 1, \\ F(t) = e^t, -\infty \leq t < 0, \\ r = 1. \end{cases}$$

Remark 1.1. We denote the dimension d , the genus p , and the generic norm $N(z, \bar{w})$ for a Cartan domain Ω (an irreducible bounded symmetric domain Ω in \mathbb{C}^d). For a given positive integer r and a positive real number μ , let

$$\phi(z) = -\mu \log N(z, \bar{z}), \quad h = e^{\lambda \phi(z)} \|w\|^2, \quad M = \left\{ (z, w) \in \Omega \times \mathbb{C}^r : \|w\|^2 < N(z, \bar{z})^{\lambda \mu} \right\}.$$

When $r = 1$ and $\mu = \frac{p}{d+1}$, by (1.12), $\frac{\sqrt{-1}}{1-h} \partial \bar{\partial}(\phi + h)$ is the Kähler form of a complete conformally Kähler Einstein metric on M for $\lambda = -1$. Note that $\sqrt{-1} \partial \bar{\partial}(\phi - \log(1-h))$ is the Kähler form of a complete Kähler-Einstein metric on M when $\mu = \frac{p}{d+1}$ and $\lambda = 1$ [30].

Corollary 1.2. Under the situation of Theorem 1.1, let $r = 1$, $a = \lambda = -1$ and $b = \nu = 0$, then \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $\mathbb{B}_{L_0}^{r*}$ if and only if

$$(1.13) \quad \begin{cases} \text{Ric}_{g_0} = -\frac{\mu}{x_0} g_0, \\ \varphi(x) = \mu \left(\frac{1}{d+2} \left(\left(\frac{x}{x_0} \right)^{d+2} - 1 \right) - \frac{\mu}{d+1} \frac{x}{x_0} \left(\left(\frac{x}{x_0} \right)^{d+1} - 1 \right) \right), -\infty \leq x_0 < x < 0, \\ \mu < 0. \end{cases}$$

In particular, for $\mu = -(d+2)$ and $x_0 = -\infty$, (1.13) is simplified as

$$(1.14) \quad \begin{cases} \text{Ric}_{g_0} = 0, \\ \varphi(x) = 1, -\infty < x < 0, \\ F(t) = \frac{1}{2} t^2, -\infty < t < 0. \end{cases}$$

Corollary 1.3. Under the situation of Theorem 1.1, let $a < 0$, $\lambda > 0$, $r = 1$, $b = 1$ and $\nu = 0$, then \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $\mathbb{B}_{L_0}^{r*}$ if and only if

$$(1.15) \quad \begin{cases} \text{Ric}_{g_0} = \frac{\lambda \mu x_0}{(1+ax_0)^2} g_0, \\ \varphi(x) = -\frac{\mu}{(1+ax_0)^2} \frac{(1-ax)(1+ax)^{2n-1}}{x^{n-1}} \int_{x_0}^x \frac{(u-x_0)u^{n-1}(1-a^2x_0u)}{(1-au)^2(1+au)^{2n}} du, 0 \leq x_0 < x < -\frac{1}{a}, \\ \mu < 0. \end{cases}$$

In particular, for $x_0 = 0$, (1.15) is simplified as

$$(1.16) \quad \begin{cases} \text{Ric}_{g_0} = 0, \\ \varphi(x) = -\frac{\mu}{n+1}x^2 - \frac{\mu}{a^2} \sum_{k=3}^{n+1} \left(\frac{k-1}{n+k-1} \prod_{j=1}^{k-2} \frac{n-j}{n+j} \right) (ax)^k, \quad -1 < ax < 0, \\ \mu < 0. \end{cases}$$

As $a \rightarrow 0^-$, the limit of (1.16) is

$$(1.17) \quad \begin{cases} \text{Ric}_{g_0} = 0, \\ \omega_F = \frac{n+1}{\mu} \sqrt{-1} \partial \bar{\partial} \log |\lambda \phi + \log |w|^2|, \\ \mu < 0. \end{cases}$$

Remark 1.2. Let $\phi = \|z\|^2$, $h = e^{\lambda \|z\|^2} |w|^2$, and $M = \{(z, w) \in \mathbb{C}^d \times \mathbb{C} : 0 < |w|^2 < e^{-\lambda \|z\|^2}\}$. Then from (1.14) and (1.17), $\frac{\sqrt{-1}}{2 \log h} \partial \bar{\partial} (\log h)^2$ is the Kähler form of a complete conformally Kähler Einstein metric on M for $\lambda = -1$, and $-\sqrt{-1} \partial \bar{\partial} \log(-\log h)$ is the Kähler form of a complete Kähler Einstein metric on M for $\lambda = 1$.

Corollary 1.4. Under the situation of Theorem 1.1, let $r = a = 1$, $\lambda > 0$ and $b = \nu = 0$, then \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $M = \mathbb{P}(E \oplus 1) - M_0$ if

$$(1.18) \quad \begin{cases} \text{Ric}_{g_0} = \lambda g_0, \\ \varphi(x) = \frac{1}{d+1} x \left(1 - \left(\frac{x}{x_1} \right)^{d+1} \right), \quad 0 < x < x_1 < +\infty, \\ \mu = 0. \end{cases}$$

Corollary 1.5. Let M_0 be an irreducible Hermitian symmetric space of compact type, i.e. M_0 is the compact dual of an irreducibly bounded symmetric domain Ω in \mathbb{C}^d , $\Omega \subset \mathbb{C}^d \subset M$ and \mathbb{C}^d is dense in M_0 . There is a Kähler-Einstein metric ω_0 on M_0 , the restriction of ω_0 on \mathbb{C}^d can be expressed as

$$\omega_0 = \sqrt{-1} \partial \bar{\partial} \log N(z, -\bar{z}),$$

where $N(z, \bar{z})$ is the generic norm of the Cartan domain Ω .

Set $\mathcal{O}(1)$ is the positive generator of the Picard group $\text{Pic}(M_0) = \mathbb{Z}$. For positive integers $r > 1$ and λ satisfying a condition $r\lambda = p$, there exists a metric h_0 on a holomorphic line bundle $L_0 = \mathcal{O}(-\lambda)$ such that $2\pi c_1(L_0, h_0) = -\lambda \omega_0$, here p is the genus of the Cartan domain Ω .

Under the situation of Theorem 1.1, let $a = 1$, $\nu = 0$ and $b < 0$, then \tilde{g} is a complete Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) on $M = \mathbb{P}(L_0^{\oplus r} \oplus 1) - M_0$ if

$$(1.19) \quad \begin{cases} \varphi(x) = -r \frac{(x+b)^{2n-1} (x-b)}{x^{n-1}} \int_{x_1}^x \frac{u^{n-1}}{(u+b)^{2n-2} (u-b)^2} du, \quad -b < x < -\frac{(r+1)b}{(r-1)}, \\ \mu = 0. \end{cases}$$

Remark 1.3. By Gao-Yau-Zhou [19], if M is a compact Kähler manifold and N is a subvariety with codimension greater than or equal to 2, then there are no complete Kähler-Einstein metrics on $M - N$. Corollary 1.5 shows that this conclusion does not hold for the conformally Kähler Einstein metric.

The paper is organized as follows. In Section 2, by the momentum profiles $\varphi(x)$ (refer to Hwang-Singer [22]) of unitary-invariant Kähler metrics g_F , we derive ordinary differential equations for conformally Kähler, Einstein metrics. In Section 3, we discuss the completeness of conformally Kähler metrics. In Section 4, by using the conclusions of Section 2 and Section 3, we obtain the proof for Theorem 1.1.

2. EQUATIONS OF CONFORMALLY KÄHLER, EINSTEIN METRICS

For convenience, we need Lemma 2.1 and Remark 2.1 of [14], namely:

Lemma 2.1. *Let ϕ be globally defined real Kähler potentials on a domain $\Omega \subset \mathbb{C}^d$, F be a smooth real-valued function on some interval $\mathcal{I} \subset \mathbb{R}$ and*

$$\Phi_F(z, w) = \nu\phi(z) + F(t),$$

where $z \in \mathbb{C}^d$, $t = \lambda\phi(z) + \log \rho^2$, $\rho = \|w\|$, $w \in \mathbb{C}^r$, and $\lambda, \nu \in \mathbb{R}$.

Set

$$Z = (z, w), \quad T \equiv (T_{ij}) := \frac{\partial^2 \Phi_F}{\partial Z^t \partial \bar{Z}} \equiv \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix} \equiv \begin{pmatrix} \frac{\partial^2 \Phi_F}{\partial z^t \partial \bar{z}} & \frac{\partial^2 \Phi_F}{\partial z^t \partial \bar{w}} \\ \frac{\partial^2 \Phi_F}{\partial w^t \partial \bar{z}} & \frac{\partial^2 \Phi_F}{\partial w^t \partial \bar{w}} \end{pmatrix}$$

and

$$T^{-1} := \begin{pmatrix} (T^{-1})_{11} & (T^{-1})_{12} \\ (T^{-1})_{21} & (T^{-1})_{22} \end{pmatrix}.$$

Then

$$T_{11} = (\nu + \lambda F') \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} + \lambda^2 F'' \frac{\partial \phi}{\partial z^t} \frac{\partial \phi}{\partial \bar{z}}, \quad T_{12} = \lambda \frac{F''}{\rho^2} \frac{\partial \phi}{\partial z^t} w,$$

$$T_{21} = \lambda \frac{F''}{\rho^2} \bar{w}^t \frac{\partial \phi}{\partial \bar{z}}, \quad T_{22} = \frac{F'}{\rho^2} I_r + \frac{F'' - F'}{\rho^4} \bar{w}^t w,$$

$$\det T = \frac{1}{\rho^{2r}} (F')^{r-1} F'' (\nu + \lambda F')^d \det \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right).$$

$$(T^{-1})_{11} = \frac{1}{\nu + \lambda F'} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1},$$

$$(T^{-1})_{12} = -\frac{\lambda}{\nu + \lambda F'} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \frac{\partial \phi}{\partial z^t} w,$$

$$(T^{-1})_{21} = -\frac{\lambda}{\nu + \lambda F'} \bar{w}^t \frac{\partial \phi}{\partial \bar{z}} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1}$$

and

$$(T^{-1})_{22} = \frac{\rho^2}{F'} I_r + \left(\frac{1}{F''} - \frac{1}{F'} \right) \bar{w}^t w + \frac{\lambda^2}{\nu + \lambda F'} \frac{\partial \phi}{\partial \bar{z}} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \frac{\partial \phi}{\partial z^t} \bar{w}^t w.$$

Where Z^t and \bar{Z} denote the transpose and the conjugate of the row vector $Z = (z, w)$, respectively, I_r denotes the identity matrix of order r , and symbols

$$\frac{\partial}{\partial z^t} = \left(\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_d} \right)^t, \quad \frac{\partial}{\partial \bar{z}} = \left(\frac{\partial}{\partial \bar{z}_1}, \dots, \frac{\partial}{\partial \bar{z}_d} \right), \quad \frac{\partial^2}{\partial z^t \partial \bar{z}} = \left(\frac{\partial^2}{\partial z_i \partial \bar{z}_j} \right).$$

Remark 2.1. Under assumptions of Lemma 2.1, Φ_F is a Kähler potential on a domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : t \in \mathcal{I}\}$ if and only if

- (i) $\nu + \lambda F'(t) > 0$, $F''(t) > 0$ for $r = 1$;
- (ii) $\nu + \lambda F'(t) > 0$, $F'(t) > 0$, $F''(t) > 0$ for $r > 1$.

Lemma 2.2. Under the situation of Lemma 2.1, if a Hermitian matrix T is positive definite, f is a function in t and $\bar{\partial} \left(\frac{\partial f}{\partial \bar{Z}} T^{-1} \right) = 0$, then there exist constants a, b such that $f = aF'(t) + b$.

Proof. Let $x = F'(t)$ and $\varphi(x) = F''(t)$. By Lemma 2.1, we have

$$\frac{\partial f}{\partial \bar{Z}} = \frac{df}{dx} \varphi(x) \left(\lambda \frac{\partial \phi}{\partial \bar{z}}, \frac{w}{\rho^2} \right)$$

and

$$\frac{\partial f}{\partial \bar{Z}} T^{-1} = \left(0, \frac{df}{dx} w \right).$$

Thus

$$\bar{\partial} \left(\frac{\partial f}{\partial \bar{Z}} T^{-1} \right) = \left(0, \bar{\partial} \left(\frac{df}{dx} w \right) \right) = 0.$$

That is $\bar{\partial} \left(\frac{df}{dx} \right) = 0$.

Since $\bar{\partial} \left(\frac{df}{dx} \right) = \frac{d^2 f}{dx^2} \varphi(x) \bar{\partial} t$ and $\varphi(x) > 0$, so $\frac{d^2 f}{dx^2} = 0$. Then there exist constants a, b such that $f = ax + b$. □

By Lemma 2.2, if g_F is a Kähler metric associated with the Kähler form $\omega_F = \sqrt{-1} \partial \bar{\partial} \Phi_F$, and $\tilde{g} = \frac{1}{f^2} g_F$ is a cKEM metric, then f must be forms $f = aF'(t) + b$.

Theorem 2.1. Let g_ϕ be a Kähler metric on a domain $\Omega \subset \mathbb{C}^d$ associated with the Kähler form $\omega_\phi = \sqrt{-1} \partial \bar{\partial} \phi$, namely $g_\phi(X, Y) = \omega_\phi(X, JY)$, where ϕ is globally defined the Kähler potential on Ω , and J is the canonical complex structure on the complex Euclidean space. Suppose that F is a smooth real-valued function on some interval $\mathcal{I} \subset \mathbb{R}$ such that

$$\Phi_F(z, w) := \nu \phi(z) + F(t) \quad (t = \lambda \phi(z) + \log \|w\|^2, \nu = 0, 1, \lambda \neq 0)$$

is a Kähler potential on a domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : t \in \mathcal{I}\}$. Set g_F is a Kähler metric on the domain M associated with the Kähler form $\omega_F = \sqrt{-1} \partial \bar{\partial} \Phi_F$.

Let $\tilde{g} = \frac{1}{f^2} g_F$, $n = d + r$, $x = F'(t)$, $\varphi(x) = F''(t)$ and $f = ax + b \neq 0$ on M .

(I) If the scalar curvature $s_{\tilde{g}}$ is constant on M for the metric $\tilde{g} = \frac{1}{f^2} g_F$, then the scalar curvature s_{g_ϕ} is constant on Ω for the metric g_ϕ and

$$(2.1) \quad \left(\frac{(\nu + \lambda x)^d x^{r-1}}{f^{2n-1}} \varphi \right)'' = \frac{(\nu + \lambda x)^d x^{r-1}}{f^{2n-1}} \left(\frac{1}{\nu + \lambda x} k_{g_\phi} + \frac{r(r-1)}{x} - \frac{k_{\tilde{g}}}{f^2} \right),$$

where $k_{g_\phi} = \frac{1}{2} s_{g_\phi}$ and $k_{\tilde{g}} = \frac{1}{2} s_{\tilde{g}}$

(II) $\tilde{g} = \frac{1}{f^2} g_F$ is an Einstein metric on M (that is $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) if and only if

$$(2.2) \quad \begin{cases} \text{Ric}_{g_\phi} = \gamma g_\phi, \\ \varphi'(x) + \left(\frac{d\lambda}{\lambda x + \nu} - \frac{(2d+1)a}{ax+b} - \frac{\lambda a}{a\lambda x + 2a\nu - b\lambda} \right) \varphi(x) - \frac{\mu(\nu + \lambda x) - \gamma(ax+b)^2}{(ax+b)(a\lambda x + 2a\nu - b\lambda)} = 0, \\ r = 1 \end{cases}$$

for $\nu a \neq 0$, and

$$(2.3) \quad \begin{cases} \text{Ric}_{g_\phi} = \gamma g_{g_\phi}, \\ \varphi'(x) + \left(\frac{d\lambda}{\lambda x + \nu} + \frac{r-1}{x} - \frac{(2n-1)a}{ax+b} - \frac{\lambda a}{a\lambda x + 2a\nu - b\lambda} \right) \varphi(x) - \frac{\mu(\nu + \lambda x) - \gamma(ax+b)^2}{(ax+b)(a\lambda x + 2a\nu - b\lambda)} = 0 \end{cases}$$

for $\nu a = 0$. Where γ is a constant, and $\gamma = \frac{\mu\nu}{f^2} + r\lambda = \frac{\mu\nu}{b^2} + r\lambda$ for $r > 1$ and $\nu a = 0$.

Proof. (I)

Set

$$\rho = \|w\|, \quad Z = (z, w), \quad T \equiv (T_{ij}) := \frac{\partial^2 \Phi_F}{\partial Z^t \partial \bar{Z}} \equiv \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix} \equiv \begin{pmatrix} \frac{\partial^2 \Phi_F}{\partial z^t \partial \bar{z}} & \frac{\partial^2 \Phi_F}{\partial z^t \partial \bar{w}} \\ \frac{\partial^2 \Phi_F}{\partial w^t \partial \bar{z}} & \frac{\partial^2 \Phi_F}{\partial w^t \partial \bar{w}} \end{pmatrix}$$

and

$$T^{-1} := \begin{pmatrix} (T^{-1})_{11} & (T^{-1})_{12} \\ (T^{-1})_{21} & (T^{-1})_{22} \end{pmatrix}.$$

Let $k_{\tilde{g}} = \frac{1}{2}s_{\tilde{g}}$ and $k_{g_F} = \frac{1}{2}s_{g_F}$, then

$$k_{\tilde{g}} = f^2 k_{g_F} + 2(2n-1)f \Delta_{g_F} f - n(2n-1)|df|_{g_F}^2 = f^2 k_{g_F} - \frac{2(2n-1)}{n-1} f^{n+1} \Delta_{g_F} f^{-n+1},$$

where

$$\Delta_{g_F} = \text{Tr} \left(T^{-1} \frac{\partial^2}{\partial Z^t \partial \bar{Z}} \right), \quad k_{g_F} = -\text{Tr} \left(T^{-1} \frac{\partial^2 \log \det T}{\partial Z^t \partial \bar{Z}} \right),$$

for the expression of T , see Lemma 2.1.

Since

$$\frac{\partial t}{\partial Z} = \left(\lambda \frac{\partial \phi}{\partial z}, \frac{\bar{w}}{\rho^2} \right), \quad \frac{\partial t}{\partial \bar{Z}} = \left(\lambda \frac{\partial \phi}{\partial \bar{z}}, \frac{w}{\rho^2} \right), \quad \frac{\partial^2 t}{\partial Z^t \partial \bar{Z}} = \begin{pmatrix} \lambda \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} & 0 \\ 0 & \frac{1}{\rho^2} I_r - \frac{\bar{w}^t w}{\rho^4} \end{pmatrix}$$

and

$$\frac{\partial^2 f^\alpha}{\partial Z^t \partial \bar{Z}} = ((f^\alpha)' \varphi)' \varphi \frac{\partial t}{\partial Z^t} \frac{\partial t}{\partial \bar{Z}} + (f^\alpha)' \varphi \frac{\partial^2 t}{\partial Z^t \partial \bar{Z}},$$

by Lemma 2.1, it follows that

$$\begin{aligned} & \text{Tr} \left(T^{-1} \frac{\partial t}{\partial Z^t} \frac{\partial t}{\partial \bar{Z}} \right) = \frac{\partial t}{\partial \bar{Z}} T^{-1} \frac{\partial t}{\partial Z^t} \\ &= \left(\lambda \frac{\partial \phi}{\partial \bar{z}}, \frac{w}{\rho^2} \right) \begin{pmatrix} (T^{-1})_{11} & (T^{-1})_{12} \\ (T^{-1})_{21} & (T^{-1})_{22} \end{pmatrix} \begin{pmatrix} \lambda \frac{\partial \phi}{\partial z^t} \\ \frac{\bar{w}^t}{\rho^2} \end{pmatrix} \\ &= \lambda^2 \frac{\partial \phi}{\partial \bar{z}} (T^{-1})_{11} \frac{\partial \phi}{\partial z^t} + \lambda \frac{\partial \phi}{\partial \bar{z}} (T^{-1})_{12} \frac{\bar{w}^t}{\rho^2} + \lambda \frac{w}{\rho^2} (T^{-1})_{21} \frac{\partial \phi}{\partial z^t} + \frac{1}{\rho^4} w (T^{-1})_{22} \bar{w}^t \\ &= \frac{\lambda^2}{\nu + \lambda F'} (1 - 1 - 1 + 1) \frac{\partial \phi}{\partial \bar{z}} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \frac{\partial \phi}{\partial z^t} + \frac{1}{F'} + \left(\frac{1}{F''} - \frac{1}{F'} \right) \\ &= \frac{1}{F''} = \frac{1}{\varphi}, \end{aligned}$$

$$\begin{aligned}
 & \operatorname{Tr} \left(T^{-1} \frac{\partial^2 t}{\partial Z^t \partial \bar{Z}} \right) \\
 &= \lambda \operatorname{Tr} \left((T^{-1})_{11} \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right) + \operatorname{Tr} \left((T^{-1})_{22} \left(\frac{1}{\rho^2} I_r - \frac{\bar{w}^t w}{\rho^4} \right) \right) \\
 &= \frac{\lambda d}{\nu + \lambda F'} + \frac{1}{F'} \operatorname{Tr} \left(I_r - \frac{\bar{w}^t w}{\rho^2} \right) \\
 &= \frac{\lambda d}{\nu + \lambda F'} + \frac{r-1}{F'} = \frac{\lambda d}{\nu + \lambda x} + \frac{r-1}{x},
 \end{aligned}$$

$$\begin{aligned}
 \Delta_{g_F} f^\alpha &= \operatorname{Tr} \left(T^{-1} \frac{\partial^2 f^\alpha}{\partial Z^t \partial \bar{Z}} \right) \\
 &= ((f^\alpha)' \varphi)' \varphi \operatorname{Tr} \left(T^{-1} \frac{\partial t}{\partial Z^t} \frac{\partial t}{\partial \bar{Z}} \right) + (f^\alpha)' \varphi \operatorname{Tr} \left(T^{-1} \frac{\partial^2 t}{\partial Z^t \partial \bar{Z}} \right) \\
 &= \frac{((f^\alpha)' \varphi)' \varphi}{\varphi} + (f^\alpha)' \varphi \left(\frac{\lambda d}{\nu + \lambda x} + \frac{r-1}{x} \right) \\
 &= ((f^\alpha)' \varphi)' + \left(\frac{\lambda d}{\nu + \lambda x} + \frac{r-1}{x} \right) (f^\alpha)' \varphi
 \end{aligned}$$

and

$$f^{-\alpha} \Delta_{g_F} f^\alpha = \frac{(f^\alpha)'}{f^\alpha} \varphi' + \left(\frac{(f^\alpha)''}{f^\alpha} + \frac{(f^\alpha)' p'}{f^\alpha p} \right) \varphi,$$

where $p := (\nu + \lambda x)^d x^{r-1}$.

So by (2.24) of [14], we have

$$\begin{aligned}
 \frac{k_{\tilde{g}}}{f^2} &= k_{g_F} + \frac{2(2n-1)}{\alpha} f^{-\alpha} \Delta_{g_F} f^\alpha \\
 &= \left(\frac{1}{\nu + \lambda x} k_{g_\phi} + \frac{r(r-1)}{x} - \frac{(p\varphi)''}{p} \right) + \frac{2(2n-1)}{\alpha} \left(\frac{(f^\alpha)'}{f^\alpha} \varphi' + \left(\frac{(f^\alpha)''}{f^\alpha} + \frac{(f^\alpha)' p'}{f^\alpha p} \right) \varphi \right) \\
 &= \frac{1}{\nu + \lambda x} k_{g_\phi} + \frac{r(r-1)}{x} - \left(\varphi'' + 2 \left(\frac{p'}{p} - \frac{(2n-1)}{\alpha} \frac{(f^\alpha)'}{f^\alpha} \right) \varphi' \right. \\
 &\quad \left. + \left(\frac{p''}{p} - \frac{2(2n-1)}{\alpha} \left(\frac{(f^\alpha)''}{f^\alpha} + \frac{(f^\alpha)' p'}{f^\alpha p} \right) \right) \varphi \right),
 \end{aligned}$$

here $\alpha = 1 - n$. Thus, we get

$$\begin{aligned}
 & \varphi'' + 2 \left(\frac{p'}{p} - \frac{(2n-1)}{\alpha} \frac{(f^\alpha)'}{f^\alpha} \right) \varphi' + \left(\frac{p''}{p} - \frac{2(2n-1)}{\alpha} \left(\frac{(f^\alpha)''}{f^\alpha} + \frac{(f^\alpha)' p'}{f^\alpha p} \right) \right) \varphi \\
 &= \frac{1}{\nu + \lambda x} k_{g_\phi} + \frac{r(r-1)}{x} - \frac{k_{\tilde{g}}}{f^2}.
 \end{aligned}$$

Using

$$\frac{(f^\alpha)''}{f^\alpha} + \left(\frac{2n-1}{\alpha} + 1 \right) \left(\frac{(f^\alpha)'}{f^\alpha} \right)^2 = 0,$$

we have (2.1).

(II) We complete the proof in four steps.

(II-1) In this part, we prove the following conclusions:

$\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric on M (that is $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) if and only if

$$(2.4) \quad \text{Ric}_{g_\phi} = \gamma g_{g_\phi},$$

$$(2.5) \quad \varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f} + \frac{a\lambda}{\lambda f - 2a(\nu + \lambda x)} \right) \varphi + \frac{\mu(\nu + \lambda x) - \gamma f^2}{f(\lambda f - 2a(\nu + \lambda x))} = 0$$

and

$$(2.6) \quad \varphi''(x) + \left(\frac{p'}{p} - \frac{2an}{f} \right) \varphi' + \left(\frac{2(2n-1)a^2}{f^2} - \frac{2ap'}{pf} + \left(\frac{p'}{p} \right)' \right) \varphi + \frac{\mu}{f^2} = 0,$$

where $p = (\nu + \lambda x)^d x^{r-1}$, and γ is a constant. For $r > 1$, $\varphi(x)$ is also satisfied

$$(2.7) \quad \left(1 - \frac{2ax}{f} \right) \varphi' + \left(\frac{p'}{p} - \frac{2a(n-1)}{f} - \frac{2axp'}{pf} + \frac{2(2n-1)a^2x}{f^2} \right) \varphi + \frac{\mu x}{f^2} - r = 0.$$

Using (see e.g. [8], 1.161)

$$\text{Ric}_{\tilde{g}} = \text{Ric}_{g_F} + \frac{2(n-1)}{f} D_{g_F}^2 f - \frac{1}{f^2} (-2f \Delta_{g_F} f + (2n-1) |df|_{g_F}^2) g_F,$$

here D_{g_F} denotes the Levi-Civita connection of g_F , we get

$$\text{Ric}_{g_F} + \frac{2(n-1)}{f} D_{g_F}^2 f = \frac{1}{f^2} (\mu - 2f \Delta_{g_F} f + (2n-1) |df|_{g_F}^2) g_F,$$

namely

$$\frac{2(n-1)}{f} \sqrt{-1} \partial \bar{\partial} f = \frac{1}{f^2} (\mu - 2f \Delta_{g_F} f + (2n-1) |df|_{g_F}^2) \sqrt{-1} \partial \bar{\partial} \Phi_F + \sqrt{-1} \partial \bar{\partial} \log \det(\partial \bar{\partial} \Phi_F)$$

or

$$(2.8) \quad \frac{2(n-1)}{f} \frac{\partial^2 f}{\partial Z^t \partial \bar{Z}} = \frac{1}{f^2} (\mu - 2f \Delta_{g_F} f + (2n-1) |df|_{g_F}^2) \frac{\partial^2 \Phi_F}{\partial Z^t \partial \bar{Z}} + \frac{\partial^2 \log \det(\partial \bar{\partial} \Phi_F)}{\partial Z^t \partial \bar{Z}}.$$

Let

$$Q = (r-1) \log F'(t) + \log F''(t) + d \log(\nu + \lambda F'(t)) - rt,$$

$$P = \log \det(\partial \bar{\partial} \Phi_F) = Q + \log \det(\partial \bar{\partial} \phi) + \lambda r \phi,$$

$$\text{Ric}_{g_\phi} = -\frac{\partial^2}{\partial z^t \partial \bar{z}} \log \left(\det \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right) \right).$$

Note that

$$\begin{aligned} Q'(t) &= (r-1) \frac{F''(t)}{F'(t)} + \frac{F'''(t)}{F''(t)} + d \frac{\lambda F''(t)}{\nu + \lambda F'(t)} - r \\ &= \varphi'(x) + \left(\frac{r-1}{x} + \frac{\lambda d}{\nu + \lambda x} \right) \varphi(x) - r \\ &= \frac{(p\varphi)'}{p}(x) - r \end{aligned}$$

and

$$Q''(t) = \left(\frac{(p\varphi)'}{p} \right)'(x) \cdot \varphi(x),$$

so

$$\begin{aligned}
 & \frac{\partial^2 \log \det(\partial\bar{\partial}\Phi_F)}{\partial Z^t \partial \bar{Z}} \\
 = & Q'' \frac{\partial t}{\partial Z^t} \frac{\partial t}{\partial \bar{Z}} + Q' \frac{\partial^2 t}{\partial Z^t \partial \bar{Z}} + \begin{pmatrix} \frac{\partial^2 \log \det(\partial\bar{\partial}\phi)}{\partial z^t \partial \bar{z}} + \lambda r \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} & 0 \\ 0 & 0 \end{pmatrix} \\
 = & Q'' \begin{pmatrix} \lambda^2 \frac{\partial \phi}{\partial z^t} \frac{\partial \phi}{\partial \bar{z}} & \frac{\lambda}{\rho^2} \frac{\partial \phi}{\partial z^t} w \\ \frac{\lambda}{\rho^2} \bar{w}^t \frac{\partial \phi}{\partial \bar{z}} & \frac{1}{\rho^4} \bar{w}^t w \end{pmatrix} + Q' \begin{pmatrix} \lambda \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} & 0 \\ 0 & \frac{1}{\rho^2} I_r - \frac{\bar{w}^t w}{\rho^4} \end{pmatrix} + \begin{pmatrix} \frac{\partial^2 \log \det(\partial\bar{\partial}\phi)}{\partial z^t \partial \bar{z}} + \lambda r \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} & 0 \\ 0 & 0 \end{pmatrix} \\
 = & \begin{pmatrix} \lambda(Q' + r) \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} + \frac{\partial^2 \log \det(\partial\bar{\partial}\phi)}{\partial z^t \partial \bar{z}} + \lambda^2 Q'' \frac{\partial \phi}{\partial z^t} \frac{\partial \phi}{\partial \bar{z}} & \frac{\lambda Q''}{\rho^2} \frac{\partial \phi}{\partial z^t} w \\ \frac{\lambda Q''}{\rho^2} \bar{w}^t \frac{\partial \phi}{\partial \bar{z}} & \frac{Q'}{\rho^2} I_r + \frac{Q'' - Q'}{\rho^4} \bar{w}^t w \end{pmatrix},
 \end{aligned}$$

$$\frac{\partial f}{\partial Z^t} = f' \varphi \left(\lambda \frac{\partial \phi}{\partial z}, \frac{\bar{w}}{\rho^2} \right)^t, \quad \frac{\partial f}{\partial \bar{Z}} = f' \varphi \left(\lambda \frac{\partial \phi}{\partial \bar{z}}, \frac{w}{\rho^2} \right),$$

$$\frac{\partial^2 f}{\partial Z^t \partial \bar{Z}} = (f' \varphi)' \varphi \frac{\partial t}{\partial Z^t} \frac{\partial t}{\partial \bar{Z}} + f' \varphi \frac{\partial^2 t}{\partial Z^t \partial \bar{Z}} = a \varphi \begin{pmatrix} \lambda \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} + \lambda^2 \varphi' \frac{\partial \phi}{\partial z^t} \frac{\partial \phi}{\partial \bar{z}} & \frac{\lambda \varphi'}{\rho^2} \frac{\partial \phi}{\partial z^t} w \\ \frac{\lambda \varphi'}{\rho^2} \bar{w}^t \frac{\partial \phi}{\partial \bar{z}} & \frac{1}{\rho^2} I_r + \frac{\varphi' - 1}{\rho^4} \bar{w}^t w \end{pmatrix},$$

$$|df|_{g_F}^2 = 2 \operatorname{Tr} \left(T^{-1} \frac{\partial f}{\partial Z^t} \frac{\partial f}{\partial \bar{Z}} \right) = 2a^2 \varphi^2 \operatorname{Tr} \left(T^{-1} \frac{\partial t}{\partial Z^t} \frac{\partial t}{\partial \bar{Z}} \right) = 2a^2 \varphi$$

and

$$\begin{aligned}
 \Delta_{g_F} f &= \operatorname{Tr} \left(T^{-1} \frac{\partial^2 f}{\partial Z^t \partial \bar{Z}} \right) = a \varphi' \varphi \operatorname{Tr} \left(T^{-1} \frac{\partial t}{\partial Z^t} \frac{\partial t}{\partial \bar{Z}} \right) + a \varphi \operatorname{Tr} \left(T^{-1} \frac{\partial^2 t}{\partial Z^t \partial \bar{Z}} \right) \\
 &= a \varphi' + a \left(\frac{\lambda d}{\nu + \lambda x} + \frac{r-1}{x} \right) \varphi = a \frac{(p\varphi)'}{p}.
 \end{aligned}$$

Let

$$\begin{aligned}
 A &:= x \frac{\mu - 2f \Delta_{g_F} f + (2n-1) |df|_{g_F}^2}{f^2} + Q' - 2a(n-1) \frac{\varphi}{f} \\
 &= x \left(\frac{\mu}{f^2} + \frac{-2a(p\varphi)'}{pf} + \frac{2(2n-1)a^2 \varphi}{f^2} \right) + \frac{(p\varphi)'}{p} - r - 2a(n-1) \frac{\varphi}{f} \\
 &= \left(1 - \frac{2ax}{f} \right) \varphi' + \left(\frac{p'}{p} - \frac{2a(n-1)}{f} - \frac{2axp'}{pf} + \frac{2(2n-1)a^2 x}{f^2} \right) \varphi + \frac{\mu x}{f^2} - r,
 \end{aligned}$$

$$D := (\varphi - x) \frac{\mu - 2f \Delta_{g_F} f + (2n-1) |df|_{g_F}^2}{f^2} + Q'' - Q' - 2a(n-1) \frac{(\varphi' - 1)\varphi}{f},$$

$$\begin{aligned}
C &:= \varphi \frac{\mu - 2f\Delta_{g_F}f + (2n-1)|df|_{g_F}^2}{f^2} + Q'' - 2a(n-1)\frac{\varphi'\varphi}{f} \\
&= \left(\frac{\mu}{f^2} + \frac{-2a\left(\varphi' + \frac{p'}{p}\varphi\right)}{f} + \frac{2(2n-1)a^2\varphi}{f^2} + \varphi'' + \frac{p'}{p}\varphi' + \left(\frac{p'}{p}\right)'\varphi - 2a(n-1)\frac{\varphi'}{f} \right) \varphi \\
&= \left(\varphi''(x) + \left(\frac{p'}{p} - \frac{2an}{f}\right)\varphi' + \left(\frac{2(2n-1)a^2}{f^2} - \frac{2ap'}{pf} + \left(\frac{p'}{p}\right)'\right)\varphi + \frac{\mu}{f^2} \right) \varphi
\end{aligned}$$

and

$$\begin{aligned}
\gamma &:= \frac{\mu - 2f\Delta_{g_F}f + (2n-1)|df|_{g_F}^2}{f^2}(\nu + \lambda x) + \lambda(Q' + r) - \frac{2(n-1)a\lambda}{f}\varphi \\
&= \frac{\mu - 2f\Delta_{g_F}f + (2n-1)|df|_{g_F}^2}{f^2}\nu + \lambda r + \lambda A \\
&= \nu \left(-\frac{2a}{f}\varphi' + \left(\frac{2(2n-1)a^2}{f^2} - \frac{2ap'}{pf}\right)\varphi + \frac{\mu}{f^2} \right) + \lambda r + \lambda A \\
(2.9) \quad &= -\frac{2a\nu}{f} \left(\varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f}\right)\varphi - \frac{\mu}{2af} \right) + \lambda r + \lambda A \\
(2.10) \quad &= \frac{\lambda f - 2a(\nu + \lambda x)}{f} \left(\varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f}\right)\varphi \right) + \frac{\lambda a}{f}\varphi + \frac{\mu(\nu + \lambda x)}{f^2}.
\end{aligned}$$

Note that the equation (2.10) is equivalent to the equation (2.5).

The equation (2.8) is equivalent to the following system of equations

$$(2.11) \quad \frac{A}{\rho^2}I_r + \frac{D}{\rho^4}\bar{w}^t w = 0,$$

$$(2.12) \quad C \frac{\lambda}{\rho^2} \frac{\partial \phi}{\partial z^t} w = 0,$$

$$(2.13) \quad \gamma \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} + \frac{\partial^2 \log \det(\partial \bar{\partial} \phi)}{\partial z^t \partial \bar{z}} + C \lambda^2 \frac{\partial \phi}{\partial z^t} \frac{\partial \phi}{\partial \bar{z}} = 0.$$

Using (2.12), we get

$$C \frac{\lambda}{\rho^2} \frac{\partial \phi}{\partial z^t} w \bar{w}^t = C \lambda \frac{\partial \phi}{\partial z^t} = 0,$$

namely $C \frac{\partial \phi}{\partial z^t} = 0$. Then

$$\lambda \frac{dC}{dx} \varphi \frac{\partial \phi}{\partial z^t} \frac{\partial \phi}{\partial \bar{z}} + C \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} = 0,$$

which implies $C^2 \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} = 0$. Since $\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} > 0$, it follows that $C \equiv A + D = 0$.

Using (2.11) and

$$\det \left(\beta I_r - \frac{A}{\rho^2} I_r - \frac{D}{\rho^4} \bar{z}^t z \right) = \left(\beta - \frac{A}{\rho^2} \right)^{r-1} \left(\beta - \frac{A+D}{\rho^2} \right),$$

we have $A + D = 0$ for $r = 1$, and $A = 0$, $A + D = 0$ for $r > 1$. Thus we have (2.6) and (2.7).

Substituting $C = 0$ into (2.13), we obtain

$$\gamma \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} + \frac{\partial^2 \log \det(\partial \bar{\partial} \phi)}{\partial z^t \partial \bar{z}} = 0.$$

For the case of $d = 1$, taking the partial derivative with respect to the variable w for the above equation, by $\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} > 0$, we have $\frac{d\gamma}{dx} \varphi \frac{\bar{w}^t}{\rho^2} = 0$, thus $\frac{d\gamma}{dx} = 0$. This indicates that γ is a constant and g_ϕ is a Kähler-Einstein metric. That is, equation (2.4) holds.

For the case of $d > 1$, we have that γ is a constant and g_ϕ is a Kähler-Einstein metric.

(II – 2) In this part, we prove the following conclusions: when $\nu a = 0$, $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$ if and only if

$$(2.14) \quad \text{Ric}_{g_\phi} = \gamma g_{g_\phi}$$

and

$$(2.15) \quad \left(1 - \frac{2ax}{f}\right)\varphi' + \left(\frac{p'}{p} - \frac{2a(n-1)}{f} - \frac{2axp'}{pf} + \frac{2(2n-1)a^2x}{f^2}\right)\varphi + \frac{\mu x}{f^2} = \begin{cases} c, & r = 1, \\ r, & r > 1. \end{cases}$$

Where c and

$$(2.16) \quad \gamma = \frac{\mu\nu}{f^2} + \lambda \times \begin{cases} c, & r = 1, \\ r, & r > 1 \end{cases}$$

are constants.

When $\nu a = 0$, from (2.9) we get

$$\gamma = \frac{\mu\nu}{f^2} + \lambda r + \lambda A.$$

By γ and $\frac{\mu\nu}{f^2}$ are constants, then A is a constant. Namely

$$(2.17) \quad A \equiv \left(1 - \frac{2ax}{f}\right)\varphi' + \left(\frac{p'}{p} - \frac{2a(n-1)}{f} - \frac{2axp'}{pf} + \frac{2(2n-1)a^2x}{f^2}\right)\varphi + \frac{\mu x}{f^2} - r = \tilde{c}.$$

Now, we show that if $\nu a = 0$ and A is constant, then the equation (2.6) holds.

Let

$$Q_1 := \frac{\frac{p'}{p} - \frac{2a(n-1)}{f} - \frac{2axp'}{pf} + \frac{2(2n-1)a^2x}{f^2}}{1 - \frac{2ax}{f}} = -\frac{a}{ax-b} + \frac{r-1}{x} + \frac{d\lambda}{\lambda x + \nu} - \frac{a(2n-1)}{ax+b},$$

$$Q_2 := \frac{\frac{\mu x}{f^2} - r}{1 - \frac{2ax}{f}} = r + \frac{4abr - \mu}{2a(ax-b)} - \frac{\mu}{2a(ax+b)},$$

$$P_1 := \frac{p'}{p} - \frac{2an}{f} = \frac{d\lambda}{\lambda x + \nu} + \frac{r-1}{x} - \frac{2an}{ax+b},$$

$$P_2 := \frac{2(2n-1)a^2}{f^2} - \frac{2ap'}{pf} + \left(\frac{p'}{p}\right)',$$

$$P_3 := \frac{\mu}{f^2},$$

$$R := \varphi'' + P_1\varphi' + P_2\varphi + P_3,$$

$$S := \varphi' + Q_1\varphi + Q_2.$$

Then from

$$S = \varphi' + Q_1\varphi + Q_2,$$

we get

$$S' = \varphi'' + Q_1\varphi' + Q_1'\varphi + Q_2'.$$

Thus

$$S' - R = (Q_1 - P_1) \left(\varphi' + \frac{Q_1' - P_2}{Q_1 - P_1} \varphi + \frac{Q_2' - P_3}{Q_1 - P_1} \right),$$

where

$$Q_1 - P_1 = -\frac{2ab}{a^2x^2 - b^2}, \quad \frac{Q'_1 - P_2}{Q_1 - P_1} = Q_1 + \frac{d\nu a}{b(\nu + \lambda x)}, \quad \frac{Q'_2 - P_3}{Q_1 - P_1} = Q_2.$$

This shows that

$$R = S' - (Q_1 - P_1)S + \frac{2d\nu a^2}{(a^2x^2 - b^2)(\nu + \lambda x)}\varphi.$$

Therefore, if $\nu a = 0$ and $S = \frac{\tilde{c}}{1 - \frac{2ax}{f}}$, then $R = 0$. That is, the equation (2.17) implies the equation (2.6).

It is easy to verify that (2.15) is equivalent to (2.5) for $\nu a = 0$.

Let $c = r + \tilde{c}$, by (2.4), (2.5) and (2.6), we obtain (2.14), (2.15) and (2.16).

(II - 3) In this part, we prove the following conclusions: when $\nu a \neq 0$, $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$ if and only if $r = 1$,

$$(2.18) \quad \text{Ric}_{g_\phi} = \gamma g_{g_\phi}$$

and

$$(2.19) \quad \varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f} + \frac{\lambda a}{\lambda f - 2a(\nu + \lambda x)} \right) \varphi + \frac{\mu(\nu + \lambda x) - \gamma f^2}{f(\lambda f - 2a(\nu + \lambda x))} = 0,$$

where γ is a constant on M .

If $\nu a \neq 0$ and $r > 1$, then by (2.7) (that is $A = 0$) and

$$\begin{aligned} \gamma &= -\frac{2a\nu}{f} \left(\varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f} \right) \varphi - \frac{\mu}{2af} \right) + \lambda r + \lambda A \\ &= -\frac{2a\nu}{f} \left(\varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f} \right) \varphi - \frac{\mu}{2af} \right) + \lambda r \end{aligned}$$

is a constant, we have

$$(2.20) \quad \left(1 - \frac{2ax}{f} \right) \left(\varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f} \right) \varphi \right) + \frac{a}{f} \varphi + \frac{\mu x}{f^2} - r = 0,$$

$$(2.21) \quad \varphi' + \left(\frac{p'}{p} - \frac{(2n-1)a}{f} \right) \varphi = \frac{\mu}{2af} + \frac{(r\lambda - \gamma)f}{2a\nu}.$$

Substituting (2.21) into (2.20), we have

$$(2.22) \quad \varphi = \frac{\lambda r - \gamma}{2\nu} x^2 + rx + \frac{\gamma - \lambda r}{2a^2\nu} b^2 + \frac{br}{a} - \frac{\mu}{2a^2}.$$

Substituting (2.22) into (2.21), we have

$$a^2(r\lambda - \gamma)x^2 + 2ra^2\nu x + 2rab\nu - b^2r\lambda + b^2\gamma - \mu\nu \equiv 0$$

or

$$a\lambda(1-n)x^2 + ((r+1-2n)a\nu + (n-1)b\lambda)x + (r-1)b\nu \equiv 0.$$

This is impossible.

Now let $r = 1$ and $n = d + 1$, it follows from (2.10) that

$$\gamma = \left(-\lambda + \frac{2(b\lambda - a\nu)}{ax + b} \right) \varphi'(x) + \left(\frac{2a(2d+1)(a\nu - b\lambda)}{(ax + b)^2} + \frac{2\lambda a}{ax + b} + \frac{\lambda^2 d}{\lambda x + \nu} \right) \varphi(x) + \frac{\mu(\lambda x + \nu)}{(ax + b)^2}.$$

Since γ is a constant on M , taking the derivative of the above equation with respect to x , it follows that

$$-\frac{(\lambda x + 2\nu)a - b\lambda}{ax + b} \left\{ \varphi''(x) + \left(\frac{\lambda d}{\lambda x + \nu} - \frac{2a(d+1)}{ax + b} \right) \varphi'(x) + \frac{\mu}{(ax + b)^2} + \left(\frac{2(2d+1)a^2}{(ax + b)^2} - \frac{2ad\lambda}{(\nu + \lambda x)(ax + b)} - \frac{d\lambda^2}{(\nu + \lambda x)^2} \right) \varphi(x) \right\} = 0,$$

so

$$\varphi''(x) + \left(\frac{\lambda d}{\lambda x + \nu} - \frac{2a(d+1)}{ax + b} \right) \varphi'(x) + \frac{\mu}{(ax + b)^2} + \left(\frac{2(2d+1)a^2}{(ax + b)^2} - \frac{2ad\lambda}{(\nu + \lambda x)(ax + b)} - \frac{d\lambda^2}{(\nu + \lambda x)^2} \right) \varphi(x) = 0.$$

Using

$$\frac{p'}{p} - \frac{2an}{f} = \frac{\lambda d}{\lambda x + \nu} - \frac{2a(d+1)}{ax + b}, \quad \frac{\mu}{f^2} = \frac{\mu}{(ax + b)^2},$$

$$\frac{2(2n-1)a^2}{f^2} - \frac{2ap'}{pf} + \left(\frac{p'}{p} \right)' = \frac{2(2d+1)a^2}{(ax + b)^2} - \frac{2ad\lambda}{(\nu + \lambda x)(ax + b)} - \frac{d\lambda^2}{(\nu + \lambda x)^2},$$

it follows from γ is constant on M that (2.5) implies (2.6).

(II – 4) Finally, note that (2.15) equivalent to (2.5) for $\nu a = 0$, from (2.15) and (2.19), $\tilde{g} = \frac{1}{f^2} g_F$ is an Einstein metric on M (that is $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) if and only if equations (2.2) and (2.3) hold. \square

Remark 2.2. For $r = 1$, under the situation of Theorem 2.1, let $u = \nu + \lambda x$, $\theta = \frac{1}{2\lambda^2\varphi(x)} d^c u$. By

$$\omega_F = \frac{1}{2} dd^c(\nu\phi(z) + F(t)) = \nu\omega_\phi + \frac{1}{2} \left(F''(t) dt \wedge d^c t + F'(t) dd^c t \right),$$

we have

$$\begin{aligned} \omega_F &= \nu\omega_\phi + \frac{1}{2} \left(\varphi(x) \frac{dx}{\varphi(x)} \wedge \frac{d^c x}{\varphi(x)} + \lambda x dd^c \phi \right) \\ &= (\nu + \lambda x)\omega_\phi + \frac{1}{2\varphi(x)} dx \wedge d^c x \\ &= u\omega_\phi + \frac{1}{2\lambda^2\varphi(x)} du \wedge d^c u = u\omega_\phi + du \wedge \theta. \end{aligned}$$

From [2], if \tilde{g} is a conformally Einstein with conformal factor f , then

$$(2.23) \quad \begin{cases} \text{Ric}_{g_\phi} = \gamma g_\phi, \\ G''(u) - \left(\frac{2(n-1)q}{qu+p} + \frac{n}{u} \right) G'(u) + \frac{2n(n-1)q}{z(qu+p)} G(u) + 2\gamma u^{n-2} = 0, \end{cases}$$

where

$$G(u) = 2\lambda^2 u^{n-1} \varphi(x), \quad ax + b = qz + p, \quad q = \frac{a}{\lambda}, \quad p = \frac{b\lambda - a\nu}{\lambda}.$$

Using the variable u and the function G , (2.2) is reduced to

$$(2.24) \quad \begin{cases} \text{Ric}_{g_\phi} = \gamma g_\phi, \\ G'(u) - \left(\frac{(2n-1)q}{qu+p} + \frac{q}{qu-p} \right) G(u) - \frac{2\mu u^n - 2\gamma u^{n-1}(qu+p)^2}{(qu+p)(qu-p)} = 0. \end{cases}$$

The following proves that if G is a solution of (2.24), then G must be a solution of (2.23).

Let $H(u) = \frac{G(u)}{(qu+p)^{2n-1}(qu-p)}$. Equations (2.23) and (2.24) are equivalent to

$$(2.25) \quad \frac{d}{du} \frac{(qu-p)^2(qu+p)^{2n}H'(u)}{u^n} + 2\gamma \frac{(qu+p)(qu-p)}{u^2} = 0$$

and

$$(2.26) \quad H'(u) - \frac{2\mu u^n - 2\gamma u^{n-1}(qu+p)^2}{(qu+p)^{2n}(qu-p)^2} = 0.$$

respectively. It is easy to see that the solution H of the equation (2.26) must satisfies the equation (2.25).

Remark 2.3. For $\nu = 0, a \neq 0, r > 1$, under the situation of Theorem 2.1, let ω_{FS} be the Fubini-Study metric on $\mathbb{C}\mathbb{P}^{r-1}$, $u = x$, $\theta = \frac{1}{2\varphi(x)}d^c u$ and $\omega_0 = \lambda\omega_\phi + \omega_{FS}$. By

$$dt = \frac{dx}{\varphi(x)} = \frac{du}{\varphi(u)}, \quad dd^c t = dd^c(\lambda\phi + \log \|w\|^2) = 2\lambda\omega_\phi + 2\omega_{FS}$$

and

$$\omega_F = \frac{1}{2}dd^c(\nu\phi(z) + F(t)) = \nu\omega_\phi + \frac{1}{2}\left(F''(t)dt \wedge d^c t + F'(t)dd^c t\right),$$

we get

$$\begin{aligned} \omega_F &= \nu\omega_\phi + \frac{1}{2}\left(\varphi(x)\frac{dx}{\varphi(x)} \wedge \frac{d^c x}{\varphi(x)} + x(2\lambda\omega_\phi + 2\omega_{FS})\right) \\ &= (\nu + \lambda x)\omega_\phi + x\omega_{FS} + \frac{1}{2\varphi(x)}dx \wedge d^c x \\ &= u\omega_0 + du \wedge \theta. \end{aligned}$$

From [2], if \tilde{g} is a conformally Einstein with conformal factor $f = ax + b$, then

$$(2.27) \quad \begin{cases} \rho_0 = c\omega_0, \\ G''(u) - \left(\frac{2(n-1)q}{qu+p} + \frac{n}{u}\right)G'(u) + \frac{2n(n-1)q}{z(qu+p)}G(u) + 2cu^{n-2} = 0, \end{cases}$$

where

$$G(u) = 2u^{n-1}\varphi(x), \quad ax + b = qz + p, \quad q = a, \quad p = b.$$

Since $\rho_0 = \rho_\phi + \rho_{FS} = \rho_\phi + r\omega_{FS}$, it follows from $\rho_0 = c\omega_0$ that

$$c = r, \quad \rho_\phi = \gamma\omega_\phi = r\lambda\omega_\phi.$$

Using the variable u , (2.3) is reduced to

$$\begin{cases} \text{Ric}_{g_\phi} = r\lambda g_\phi, \\ \varphi'(u) + \left(\frac{n-1}{u} - \frac{(2n-1)q}{qu+p} - \frac{q}{qu-p}\right)\varphi(u) - \frac{\mu u - r(qu+p)^2}{(qu+p)(qu-p)} = 0. \end{cases}$$

That is

$$\frac{d}{du} \frac{2u^{n-1}\varphi(u)}{(qu+p)^{2n-1}(qu-p)} = \frac{2\mu u^n - 2ru^{n-1}(qu+p)^2}{(qu+p)^{2n}(qu-p)^2}$$

or

$$\frac{d}{du} \left(2u^{n-1}\varphi(u)\right) - \left(\frac{(2n-1)q}{qu+p} + \frac{q}{qu-p}\right) \left(2u^{n-1}\varphi(u)\right) - \frac{2\mu u^n - 2ru^{n-1}(qu+p)^2}{(qu+p)(qu-p)} = 0.$$

Which implies that $2u^{n-1}\varphi(u)$ is a solution of (2.27).

3. THE COMPLETENESS OF CONFORMALLY KÄHLER METRICS

Lemma 3.1. *Under the situation of Theorem 1.1, for a given point $z_0 \in M_0$, let*

$$M_{z_0} = \{u \in M : u \in \Pi^{-1}(z_0)\}, N_0 = \{u \in M : h(u, u) = 0\}.$$

then M_{z_0} is totally geodesic with respect to \tilde{g} . If $\nu = 1$ and $N_0 \neq \emptyset$, then N_0 is also a totally geodesic submanifold of M with respect to the metric \tilde{g} .

Proof. Given a point $z_0 \in M_0$, take a sufficiently small holomorphic neighborhood Ω (Ω is required to be holomorphic homeomorphic to a domain in \mathbb{C}^d , the following Ω is not distinguished from its holomorphic homeomorphism in \mathbb{C}^d and is still denoted as Ω) of z_0 in M such that the Kähler metric g_0 has a Kähler potential ϕ on Ω and the holomorphic bundle L_0 has a local trivializing holomorphic section $\sigma : \Omega \rightarrow L_0 \setminus \{0\}$ over Ω , thus $E = L_0^{\oplus r}$ has holomorphic coordinates $(z, w) \in \Omega \times \mathbb{C}^r$, so the Hermitian metric h of holomorphic vector bundles E can be expressed as

$$h(u, u) = e^{\lambda\phi(z)} \|w\|^2, u = (z, w\sigma(z)) \in E.$$

Therefore, the Kähler form of the Kähler metric g_F is expressed as $\omega_F = \sqrt{-1}\partial\bar{\partial}\Phi_F$, where

$$\Phi_F = \nu\phi(z) + F(t), t = \lambda\phi(z) + \log \rho^2, \rho = \|w\|, z \in \Omega \subset M_0, u = (z, w\sigma(z)) \in E.$$

For $U \in \mathcal{U}(r)$ (the unitary group of order r) and 1 is not the eigenvalue of unitary matrix U , it is easy to see that N_0 is the fixed point set of isometric mapping

$$(z, w\sigma(z)) \in M \mapsto (z, wU\sigma(z)) \in M,$$

so N_0 is a totally geodesic submanifold of (M, \tilde{g}) .

Let D be the Chern connection for the Kähler manifold (M, g) , \tilde{D} be the Levi-Civita connection for the Riemannian manifold (M, \tilde{g}) .

Setting

$$D \begin{pmatrix} \frac{\partial}{\partial Z^t} \\ \frac{\partial}{\partial \bar{Z}^t} \end{pmatrix} = \Omega_{\mathbb{C}} \otimes \begin{pmatrix} \frac{\partial}{\partial Z^t} \\ \frac{\partial}{\partial \bar{Z}^t} \end{pmatrix}, \tilde{D} \begin{pmatrix} \frac{\partial}{\partial Z^t} \\ \frac{\partial}{\partial \bar{Z}^t} \end{pmatrix} = \tilde{\Omega}_{\mathbb{C}} \otimes \begin{pmatrix} \frac{\partial}{\partial Z^t} \\ \frac{\partial}{\partial \bar{Z}^t} \end{pmatrix},$$

where

$$\begin{aligned} Z &= (z, w), \Omega_{\mathbb{C}} = \begin{pmatrix} (\partial T)T^{-1} & 0 \\ 0 & (\bar{\partial} \bar{T})\bar{T}^{-1} \end{pmatrix}, \\ \tilde{\Omega}_{\mathbb{C}} &= \begin{pmatrix} (\partial T)T^{-1} & 0 \\ 0 & (\bar{\partial} \bar{T})\bar{T}^{-1} \end{pmatrix} - \frac{df}{f} I_{2n} - \frac{1}{f} \begin{pmatrix} \frac{\partial f}{\partial Z^t} \\ \frac{\partial f}{\partial \bar{Z}^t} \end{pmatrix} \begin{pmatrix} dZ & d\bar{Z} \end{pmatrix} \\ (3.1) \quad &+ \frac{1}{f} \begin{pmatrix} 0 & T \\ \bar{T} & 0 \end{pmatrix} \begin{pmatrix} dZ^t \\ d\bar{Z}^t \end{pmatrix} \begin{pmatrix} \frac{\partial f}{\partial Z} & \frac{\partial f}{\partial \bar{Z}} \end{pmatrix} \begin{pmatrix} 0 & \bar{T}^{-1} \\ T^{-1} & 0 \end{pmatrix}, \end{aligned}$$

the definition of T can be referred to Lemma 2.1, I_{2n} is the identity matrix of order $2n$, and ∂ and $\bar{\partial}$ can be written as

$$\partial = dz \frac{\partial}{\partial z^t} + dw \frac{\partial}{\partial w^t}, \bar{\partial} = d\bar{z} \frac{\partial}{\partial \bar{z}^t} + d\bar{w} \frac{\partial}{\partial \bar{w}^t}.$$

For the expression of $\tilde{\Omega}_{\mathbb{C}}$, see [15].

There exists local coordinates (z_1, \dots, z_d) on a neighborhood of the point z_0 such that

$$\phi(z_0) = 0, \frac{\partial \phi}{\partial z}(z_0) = \frac{\partial \phi}{\partial \bar{z}}(z_0) = 0, \frac{\partial^2 \phi}{\partial z^t \partial \bar{z}}(z_0) = I_d \text{ (the identity matrix of order } d\text{)}$$

for the Kähler potential ϕ on the domain Ω . Then

$$\begin{aligned}\frac{\partial f}{\partial Z} &= \left(f' \lambda \frac{\partial \phi}{\partial z} \quad \frac{\partial f}{\partial w} \right) = \left(0 \quad \frac{\partial f}{\partial w} \right), \quad \frac{\partial f}{\partial \bar{Z}} = \left(f' \lambda \frac{\partial \phi}{\partial \bar{z}} \quad \frac{\partial f}{\partial \bar{w}} \right) = \left(0 \quad \frac{\partial f}{\partial \bar{w}} \right), \\ T &= \begin{pmatrix} (\nu + \lambda F') I_d & 0 \\ 0 & T_{22} \end{pmatrix}, \quad T^{-1} = \begin{pmatrix} \frac{1}{\nu + \lambda F'} I_d & 0 \\ 0 & T_{22}^{-1} \end{pmatrix}, \\ \partial_w T &= \begin{pmatrix} \lambda F'' \partial_w t I_d & 0 \\ 0 & \partial_w T_{22} \end{pmatrix}, \quad \bar{\partial}_w T = \begin{pmatrix} \lambda F'' \bar{\partial}_w t I_d & 0 \\ 0 & \bar{\partial}_w T_{22} \end{pmatrix}\end{aligned}$$

and

$$(\partial_w T) T^{-1} = \begin{pmatrix} \frac{\lambda F'' \partial_w t}{\nu + \lambda F'} I_d & 0 \\ 0 & (\partial_w T_{22}) T_{22}^{-1} \end{pmatrix}, \quad (\bar{\partial}_w T) \bar{T}^{-1} = \begin{pmatrix} \frac{\lambda F'' \bar{\partial}_w t}{\nu + \lambda F'} I_d & 0 \\ 0 & (\bar{\partial}_w T_{22}) \bar{T}_{22}^{-1} \end{pmatrix}$$

at $z = z_0$, where $\partial_w = dw \frac{\partial}{\partial w^t}$, $\bar{\partial}_w = d\bar{w} \frac{\partial}{\partial \bar{w}^t}$, $\partial_w t = \frac{1}{\rho^2} dw \bar{w}^t$ and $\bar{\partial}_w t = \frac{1}{\rho^2} w d\bar{w}^t$.

When $z = z_0$, $dz = 0$ and $d\bar{z} = 0$, by (3.1) we have

$$\tilde{D} \begin{pmatrix} \frac{\partial}{\partial w^t} \\ \frac{\partial}{\partial \bar{w}^t} \end{pmatrix} = \tilde{\Omega}_{\mathbb{C}}^w \otimes \begin{pmatrix} \frac{\partial}{\partial w^t} \\ \frac{\partial}{\partial \bar{w}^t} \end{pmatrix},$$

where

$$\begin{aligned}(3.2) \quad \tilde{\Omega}_{\mathbb{C}}^w &= \begin{pmatrix} (\partial_w T_{22}) T_{22}^{-1} & 0 \\ 0 & (\bar{\partial}_w T_{22}) \bar{T}_{22}^{-1} \end{pmatrix} - \frac{df}{f} I_{2r} - \frac{1}{f} \begin{pmatrix} \frac{\partial f}{\partial w^t} \\ \frac{\partial f}{\partial \bar{w}^t} \end{pmatrix} \begin{pmatrix} dw & d\bar{w} \end{pmatrix} \\ &+ \frac{1}{f} \begin{pmatrix} 0 & T_{22} \\ \bar{T}_{22} & 0 \end{pmatrix} \begin{pmatrix} dw^t \\ d\bar{w}^t \end{pmatrix} \begin{pmatrix} \frac{\partial f}{\partial w} & \frac{\partial f}{\partial \bar{w}} \end{pmatrix} \begin{pmatrix} 0 & \bar{T}_{22}^{-1} \\ T_{22}^{-1} & 0 \end{pmatrix}.\end{aligned}$$

On the other hand, let \hat{D} be the Levi-Civita connection of the induced metric of \tilde{g} on M_{z_0} , then

$$\hat{D} \begin{pmatrix} \frac{\partial}{\partial w^t} \\ \frac{\partial}{\partial \bar{w}^t} \end{pmatrix} = \hat{\Omega}_{\mathbb{C}} \otimes \begin{pmatrix} \frac{\partial}{\partial w^t} \\ \frac{\partial}{\partial \bar{w}^t} \end{pmatrix}.$$

By Lemma 2.1, we get $\hat{\Omega}_{\mathbb{C}} = \tilde{\Omega}_{\mathbb{C}}^w$, this implies that

$$\tilde{D}_{e_j} e_i = \hat{D}_{e_i} e_j$$

at $z = z_0$ for all $e_i, e_j \in \left\{ \frac{\partial}{\partial w_1}, \dots, \frac{\partial}{\partial w_r}, \frac{\partial}{\partial \bar{w}_1}, \dots, \frac{\partial}{\partial \bar{w}_r} \right\}$, thus M_{z_0} is a totally geodesic submanifold of (M, \tilde{g}) . □

Theorem 3.1. *Under the assumption of Theorem 1.1, let*

$$l = l(t_0) + \int_{x(t_0)}^{x(t)} \frac{1}{(ax+b)\sqrt{\varphi(x)}} dx, \quad x_0 < x(t_0) < x_1.$$

If $\lim_{x \rightarrow x_0^+} l = -\infty$ for M is any one of

$$E^*, \mathbb{B}_{L_0}^{r*}, E - \{u \in E : h(u, u) \leq 1\}, \mathbb{P}(E \oplus 1) - M_0, \mathbb{P}(E \oplus 1) - \{u \in E : h(u, u) \leq 1\};$$

and $\lim_{x \rightarrow x_1^-} l = +\infty$ for M is any one of $E, E^, \mathbb{B}_{L_0}^r, E - \{u \in E : h(u, u) \leq 1\}$ and $\mathbb{B}_{L_0}^{r*}$. Then the Riemannian metric \tilde{g} is complete on M .*

Remark 3.1. By Lemma 3.2 of [15] and Theorem 3.1, if the metric g_0 on M_0 is complete and fiber metrics of the metric \tilde{g} on M are complete, then the metric \tilde{g} must be complete on M .

Proof of Theorem 3.1. The proof is divided into two parts.

(i) Firstly, we calculate the square of the norm of the differential of a function on M with respect to the metric \tilde{g} .

Let $H = \psi(z) + p(t)$, $z \in M_0$, $t = \log h(u, u)$ and $u \in E$.

According to the proof of Lemma 3.1, the Kähler form of the Kähler metric g_F can be expressed locally as $\omega_F = \sqrt{-1}\partial\bar{\partial}\Phi_F$, where

$$\Phi_F = \nu\phi(z) + F(t), t = \lambda\phi(z) + \log \rho^2, \rho = \|w\|, z \in \Omega \subset M_0, u = (z, w\sigma(z)) \in E.$$

Since

$$\frac{\partial H}{\partial z} = \frac{\partial \psi}{\partial z} + \lambda p' \frac{\partial \phi}{\partial z}$$

and

$$\frac{\partial H}{\partial w} = p' \frac{\bar{w}}{\rho^2},$$

thus

$$\frac{\partial H}{\partial Z} = \left(\frac{\partial \psi}{\partial z} + \lambda p' \frac{\partial \phi}{\partial z}, p' \frac{\bar{w}}{\rho^2} \right), \|\mathrm{d}H\|_{\tilde{g}}^2 = 2 \frac{\partial H}{\partial \bar{Z}} \left(\frac{1}{f^2} T \right)^{-1} \frac{\partial H}{\partial Z^t},$$

where

$$Z = (z, w), T = \frac{\partial^2 \Phi_F}{\partial Z^t \partial \bar{Z}}.$$

By Lemma 2.1, we have

$$\begin{aligned} \frac{\partial H}{\partial \bar{z}} T_{11}^{-1} \frac{\partial H}{\partial z^t} &= \frac{1}{\nu + \lambda F'} \left(\frac{\partial \psi}{\partial \bar{z}} + \lambda p' \frac{\partial \phi}{\partial \bar{z}} \right) \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \left(\frac{\partial \psi}{\partial z^t} + \lambda p' \frac{\partial \phi}{\partial z^t} \right) \\ &= \frac{1}{2(\nu + \lambda F')} \|\mathrm{d}\psi + \lambda p' \mathrm{d}\phi\|_{g_0}^2, \end{aligned}$$

$$\begin{aligned} \frac{\partial H}{\partial \bar{z}} T_{12}^{-1} \frac{\partial H}{\partial w^t} &= -\frac{\lambda}{\nu + \lambda F'} \left(\frac{\partial \psi}{\partial \bar{z}} + \lambda p' \frac{\partial \phi}{\partial \bar{z}} \right) \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \frac{\partial \phi}{\partial z^t} w p' \frac{\bar{w}^t}{\rho^2} \\ &= -\frac{\lambda p'}{(\nu + \lambda F')} \left(\frac{\partial \psi}{\partial \bar{z}} + \lambda p' \frac{\partial \phi}{\partial \bar{z}} \right) \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \frac{\partial \phi}{\partial z^t}, \end{aligned}$$

$$\frac{\partial H}{\partial \bar{w}} T_{21}^{-1} \frac{\partial H}{\partial z^t} = -\frac{\lambda p'}{(\nu + \lambda F')} \frac{\partial \phi}{\partial \bar{z}} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \left(\frac{\partial \psi}{\partial z^t} + \lambda p' \frac{\partial \phi}{\partial z^t} \right),$$

$$\begin{aligned} \frac{\partial H}{\partial \bar{w}} T_{22}^{-1} \frac{\partial H}{\partial w^t} &= \frac{(p')^2}{\rho^4} w \left(\frac{r^2}{F'} I_r + \left(\frac{1}{F''} - \frac{1}{F'} \right) \bar{w}^t w + \frac{\lambda^2}{\nu + \lambda F'} \frac{\partial \phi}{\partial \bar{z}} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \frac{\partial \phi}{\partial z^t} \bar{w}^t w \right) \bar{w}^t \\ &= (p')^2 \left(\frac{1}{F'} + \left(\frac{1}{F''} - \frac{1}{F'} \right) + \frac{\lambda^2}{\nu + \lambda F'} \frac{\partial \phi}{\partial \bar{z}} \left(\frac{\partial^2 \phi}{\partial z^t \partial \bar{z}} \right)^{-1} \frac{\partial \phi}{\partial z^t} \right) \\ &= (p')^2 \left(\frac{1}{F''} + \frac{\lambda^2}{2(\nu + \lambda F')} \|\mathrm{d}\phi\|_{g_0}^2 \right). \end{aligned}$$

So

$$\begin{aligned}
\|dH\|_{\tilde{g}}^2 &= 2f^2 \frac{\partial H}{\partial \bar{Z}} T^{-1} \frac{\partial H}{\partial Z^t} \\
&= 2f^2 \left(\frac{1}{2(\nu + \lambda F')} \|d\psi + \lambda p' d\phi\|_{g_0}^2 - \frac{\lambda p'}{(\nu + \lambda F')} (d\psi + \lambda p' d\phi, d\phi)_{g_0} \right. \\
&\quad \left. + (p')^2 \left(\frac{1}{F''} + \frac{\lambda^2}{2(\nu + \lambda F')} \|d\phi\|_{g_0}^2 \right) \right) \\
&= 2f^2 \left(\frac{1}{2(\nu + \lambda F')} \|d\psi\|_{g_0}^2 + \frac{(p')^2}{F''} \right) \\
&= \frac{f^2}{\nu + \lambda F'} \|d\psi\|_{g_0}^2 + 2 \frac{(fp')^2}{F''}.
\end{aligned}$$

(ii) Secondly, we prove that the metric \tilde{g} is complete.

Since the metric g_0 is complete, it follows that there is a smooth proper function $\psi(z)$ on M_0 such that $\psi(z) > 1$ (otherwise, replace ψ with the function $c \log(\psi^2 + 4) + 1$) and $\|\psi\|_{g_0}^2 \leq 1$. For a sufficiently large positive integer m , let $p = (e^l + e^{-l})^m$, $H = \log(p(t) + \psi(z))$, then H is a smooth proper function on M , and

$$\|dH\|_{\tilde{g}}^2 = \frac{\|d(p + \psi)\|_{g_0}^2}{(p + \psi)^2} = \frac{1}{(p + \psi)^2} \frac{f^2}{\nu + \lambda F'} \|d\psi\|_{g_0}^2 + \frac{2}{(p + \psi)^2} \frac{(fp')^2}{F''}.$$

From

$$l = l(t_0) + \int_{x(t_0)}^{x(t)} \frac{1}{(ax + b)\sqrt{\varphi(x)}} dx = l(t_0) + \int_{t_0}^t \frac{\sqrt{F''(t)}}{f(t)} dt,$$

we get

$$f(t)l'(t) = \sqrt{F''(t)}.$$

Then

$$\frac{1}{(p + \psi)^2} \frac{(fp')^2}{F''} = \frac{p^2}{(p + \psi)^2} \left(\frac{p'}{p} \right)^2 \frac{f^2}{F''} \leq \left(\frac{p'}{p} \right)^2 \frac{f^2}{F''} = m^2 \left(\frac{e^l - e^{-l}}{e^l + e^{-l}} \right)^2 \frac{(fl')^2}{F''} \leq m^2.$$

Let

$$x_0 = \inf\{F'(t)\}, \quad x_1 = \sup\{F'(t)\}, \quad X_0 = \nu + \lambda x_0, \quad X_1 = \nu + \lambda x_1.$$

Then $x_0 < x_1$, $X_0 \geq 0$ and $X_1 \geq 0$.

If $0 < X_0, X_1 < +\infty$, then $\frac{f(x)^2}{\nu + \lambda x}$ is bounded on $[x_0, x_1]$, so by

$$\frac{1}{(p + \psi)^2} \frac{f^2}{\nu + \lambda F'} \|d\psi\|_{g_0}^2 \leq \frac{f^2}{p^2(\nu + \lambda F')} = \frac{f^2}{p^2(\nu + \lambda x)} \leq \frac{f^2}{\nu + \lambda x},$$

we get $\frac{1}{(p + \psi)^2} \frac{f^2}{\nu + \lambda F'} \|d\psi\|_{g_0}^2$ is bounded on M . Therefore, the metric \tilde{g} is complete on M .

If $f = \nu + \lambda x$ and $0 \leq X_0, X_1 < +\infty$, we also obtain that the metric \tilde{g} is complete on M .

For the case of $X_0 = 0$ or $X_1 = 0$ or $X_0 = +\infty$ or $X_1 = +\infty$. Without losing generality, we assume that $X_0 = 0$ or $+\infty$.

When $X_0 = 0$, let $u = \frac{1}{x - x_0}$ and $l_0 = l(t_0)$. By

$$\int_{x(t_0)}^{x_0} \frac{1}{(ax + b)\sqrt{\varphi(x)}} dx = -\infty,$$

there exists a positive rational number $k \geq 1$ and a positive real number δ such that $(ax+b)\sqrt{\varphi(x)} = (x-x_0)^k\psi(x)$, $\psi(x_0) > 0$, and the Taylor expansion of a function $\frac{1}{\psi(x)}$ at x_0 :

$$\frac{1}{\psi(x)} = \sum_{j=0}^{+\infty} A_j(x-x_0)^j$$

converges uniformly on the interval $[x_0, x_0 + \delta]$. So a series

$$\frac{1}{(ax+b)\sqrt{\varphi(x)}} = \sum_{j=0}^{[k]-1} \frac{A_j}{(x-x_0)^{k-j}} + \sum_{j=[k]}^{\infty} A_j(x-x_0)^{j-k},$$

converges uniformly on the closed subinterval of the interval $(x_0, x_0 + \delta)$, where $[k]$ is the largest integer not exceeding k .

Using

$$\nu + \lambda x = \lambda(x-x_0) \text{ (by } \nu + \lambda x_0 = 0),$$

$$\begin{aligned} l &= l_0 + \int_{x(t_0)}^x \frac{1}{(au+b)\sqrt{\varphi(u)}} du \quad (x(t_0) \in (x_0, x_0 + \delta), x \in (x_0, x_0 + \delta)) \\ &= C + \sum_{j=0}^{[k]-2} \frac{A_j}{(j+1-k)(x-x_0)^{k-j-1}} \\ &\quad + A_{[k]-1} \times \begin{cases} \log(x-x_0) \text{ (for } k = [k]) \\ \frac{1}{([k]-k)(x-x_0)^{k-[k]}} \text{ (for } k > [k]) \end{cases} + o((x-x_0)^{[k]+1-k}) \end{aligned}$$

and

$$e^{-l} = e^{-C+o((x-x_0)^{[k]+1-k})} e^{\sum_{j=0}^{[k]-2} \frac{A_j}{(k-j-1)(x-x_0)^{k-j-1}}} \times \begin{cases} \frac{1}{(x-x_0)^{A_{[k]-1}}} \text{ (for } k = [k]) \\ e^{\frac{A_{[k]-1}}{k-[k]} \frac{1}{(x-x_0)^{k-[k]}}} \text{ (for } k > [k]) \end{cases},$$

it follows that

$$e^{-l} \sim e^{-C} e^{\sum_{j=0}^{[k]-2} \frac{A_j u^{k-j-1}}{k-j-1}} \times \begin{cases} u^{A_{[k]-1}} \text{ (for } k = [k]) \\ e^{\frac{A_{[k]-1}}{k-[k]} u^{k-[k]}} \text{ (for } k > [k]) \end{cases} \quad (\text{as } u \rightarrow +\infty)$$

and

$$\frac{1}{p^2(\nu + \lambda x)} \sim \frac{u}{\lambda e^{-2ml}} \sim \frac{e^{2mC}}{\lambda} \times \begin{cases} u^{1-2mA_0} \text{ (for } k = 1) \\ \frac{u^{1-2mA_{[k]-1}}}{e^{2m \sum_{j=0}^{[k]-2} \frac{A_j u^{k-j-1}}{k-j-1}}} \text{ (for } k = [k] > 1) \\ \frac{u}{e^{2m \sum_{j=0}^{[k]-1} \frac{A_j u^{k-j-1}}{k-j-1}}} \text{ (for } k > [k] \geq 1) \end{cases}$$

as $u \rightarrow +\infty$, here C is a constant that depends on x_0 , l_0 and $x(t_0)$. By $f > 0$ and $\varphi > 0$ on M , we have $A_0 > 0$, so there exists a positive integer m such that $mA_0 > 1$. Thus

$$\frac{1}{(p+\psi)^2} \frac{f^2}{\nu + \lambda F'} \|\mathrm{d}\psi\|_{g_0}^2 \leq \frac{f^2}{p^2(\nu + \lambda F')} \rightarrow 0 \text{ as } x \rightarrow x_0^+.$$

When $X_0 = +\infty$, namely $x_0 = -\infty$ and $\lambda < 0$. If $f(x) = ax + b \equiv b$, then

$$\frac{1}{(p + \psi)^2} \frac{f^2}{\nu + \lambda F'} \|\mathrm{d}\psi\|_{g_0}^2 \leq \frac{f^2}{p^2(\nu + \lambda x)} \rightarrow 0 \text{ as } x \rightarrow -\infty.$$

If $f(x) = ax + b$ and $a < 0$, then by $\lim_{x \rightarrow x_0^+} l = -\infty$, $f > 0$ and $\varphi > 0$, there are constants c and C such that

$$\begin{aligned} \frac{1}{(ax + b)\sqrt{\varphi(x)}} &= \frac{1}{-x} \sum_{j=0}^{+\infty} \frac{A_j}{x^j} \quad (x \in (-\infty, c), A_0 > 0), \\ l &= l_0 + \int_{x(t_0)}^x \frac{1}{(au + b)\sqrt{\varphi(u)}} \mathrm{d}u \\ &= C - A_0 \log|x| + \sum_{j=1}^{+\infty} \frac{A_j}{jx^j} \\ &= C - A_0 \log|x| + o\left(\frac{1}{x}\right) \end{aligned}$$

and

$$\frac{f^2}{p^2(\nu + \lambda x)} \sim \frac{a^2 x}{\lambda e^{-2ml}} \sim \frac{a^2 e^{2mC}}{\lambda} \frac{x}{|x|^{2mA_0}}$$

as $x \rightarrow -\infty$. Since $A_0 > 0$, there exists a positive integer m such that $mA_0 > 1$, so

$$\frac{1}{(p + \psi)^2} \frac{f^2}{\nu + \lambda F'} \|\mathrm{d}\psi\|_{g_0}^2 \leq \frac{f^2}{p^2(\nu + \lambda x)} \rightarrow 0 \text{ as } x \rightarrow -\infty.$$

From the above conclusions, we have that

$$\|\mathrm{d}H\|_{\tilde{g}}^2 = \frac{\|\mathrm{d}(p + \psi)\|_g^2}{(p + \psi)^2} = \frac{1}{(p + \psi)^2} \frac{f^2}{\nu + \lambda F'} \|\mathrm{d}\psi\|_{g_0}^2 + \frac{1}{(p + \psi)^2} \frac{(fp')^2}{F''}.$$

is bounded on M , so by [20, 21] the metric \tilde{g} is complete on M . \square

4. PROOF OF THEOREM 1.1

Proof of Theorem 1.1. From Lemma 4.1 of [16], the metric g_F can be extended across $M = \mathbb{P}(E \oplus 1) - E$ if and only if $\varphi(x_1) = 0$ and $\varphi'(x_1) = -1$.

Since the metric g_0 is complete on M_0 , using Lemma 3.1 and Theorem 3.1, it follows that the metric \tilde{g} is complete on M if and only if its fiber metrics are complete.

From the proof of Lemma 3.1, the Kähler form of the Kähler metric g_F can be expressed locally as $\omega_F = \sqrt{-1} \partial \bar{\partial} \Phi_F$, where

$$\Phi_F = \nu \phi(z) + F(t), \quad t = \lambda \phi(z) + \log \rho^2, \quad \rho = \|w\|, \quad z \in \Omega \subset M_0, \quad u = (z, w\sigma(z)) \in E.$$

Therefore, the proof of Theorem 1.1 is converted to the proof of Theorem 4.1 below. \square

Theorem 4.1. *Let g_0 be a Kähler metric on a domain $\Omega \subset \mathbb{C}^d$ associated with the Kähler form $\omega_\phi = \sqrt{-1} \partial \bar{\partial} \phi$, namely $g_0(X, Y) = \omega_\phi(X, JY)$, where ϕ is globally defined the Kähler potential on the domain $\Omega \subset \mathbb{C}^d$, and J is the canonical complex structure on the complex Euclidean space. Suppose M represents any of the following sets*

$$\Omega \times \mathbb{C}^r, \quad \left\{ (z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda \phi(z)} \|w\|^2 \right\}, \quad \left\{ (z, w) \in \Omega \times \mathbb{C}^r : e^{\lambda \phi(z)} \|w\|^2 < 1 \right\}$$

and

$$\left\{ (z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)} \|w\|^2 < 1 \right\}, \left\{ (z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)} \|w\|^2 \right\}.$$

Set g_F is a Kähler metric on the domain M associated with the Kähler form $\omega_F = \sqrt{-1}\partial\bar{\partial}\Phi_F$, here

$$\Phi_F(z, w) = \nu\phi(z) + F(t), \quad t = \lambda\phi(z) + \log \|w\|^2, \quad \nu = 0, 1, \quad \lambda \neq 0.$$

Let

$$n = d + r, \quad x = F'(t), \quad \varphi(x) = F''(t), \quad f = ax + b > 0, \quad a \neq 0, \quad x_0 = \inf\{F'(t)\}, \quad x_1 = \sup\{F'(t)\}.$$

The following conclusions hold.

(I) For $\nu = 1$ and $b = 1$, $\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu\tilde{g}$) with the complete fibre metrics on the domain M if and only if

$$(4.1) \quad \begin{cases} \text{Ric}_{g_0} = (\lambda + \mu - 2a) g_0, \\ \varphi(x) = \frac{(1+ax)^{2d+1}(a\lambda x + 2a - \lambda)}{(1+\lambda x)^d} \int_0^x \frac{(1+\lambda u)^d (\mu(1+\lambda u) - (\lambda + \mu - 2a)(1+au)^2)}{(1+au)^{2d+2}(a\lambda u + 2a - \lambda)^2} du, \quad 0 \leq x < -\frac{1}{a}, \\ \mu \leq 0, a \leq \lambda, a < 0, r = 1, \end{cases}$$

here

$$M = \begin{cases} \{(z, w) \in \Omega \times \mathbb{C}^r : e^{\lambda\phi(z)} \|w\|^2 < 1\} \text{ for } \mu < 0, \\ \Omega \times \mathbb{C}^r \text{ for } \mu = 0. \end{cases}$$

(II)

(II-1) $\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu\tilde{g}$) with the complete fibre metrics on the domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)} \|w\|^2 < 1\}$ if and only if

$$(4.2) \quad \begin{cases} \text{Ric}_{g_0} = \mu \frac{\nu + \lambda x_0}{(b + ax_0)^2} g_0, \\ \varphi(x) = \mu \frac{(b+ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_0}^x \frac{(\nu + \lambda u)^d \left((\nu + \lambda u) - \frac{\nu + \lambda x_0}{(b + ax_0)^2} (b + au)^2 \right)}{(b + au)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ r = 1, \mu < 0, a < 0, \end{cases}$$

where

$$\begin{cases} -\infty \leq x_0 < x_1 = -\frac{b}{a} \text{ for } \lambda < 0 \text{ and } -\frac{b}{a} \leq -\frac{\nu}{\lambda}, \\ -\frac{\nu}{\lambda} \leq x_0 < x_1 = -\frac{b}{a} \text{ for } \lambda > 0 \text{ and } -\frac{\nu}{\lambda} < -\frac{b}{a}. \end{cases}$$

(II-2) $\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu\tilde{g}$) with the complete fibre metrics on the domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)} \|w\|^2\}$ if and only if

$$(4.3) \quad \begin{cases} \text{Ric}_{g_0} = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} g_0, \\ \varphi(x) = \mu \frac{(b+ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d \left((\nu + \lambda u) - \frac{\nu + \lambda x_1}{(b + ax_1)^2} (b + au)^2 \right)}{(b + au)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ r = 1, \mu < 0, a > 0, \end{cases}$$

where

$$\begin{cases} -\frac{b}{a} = x_0 < x_1 \leq +\infty \text{ for } \lambda > 0 \text{ and } -\frac{b}{a} \geq -\frac{\nu}{\lambda}, \\ -\frac{b}{a} = x_0 < x_1 \leq -\frac{\nu}{\lambda} \text{ for } \lambda < 0 \text{ and } -\frac{\nu}{\lambda} > -\frac{b}{a}. \end{cases}$$

(III) Let $r = 1$, $\varphi(x_1) = 0$ and $\varphi'(x_1) = -1$.

(III-1) $\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) with the fibre metrics complete at $t = -\infty$ on the domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2\}$ if and only if

$$(4.4) \quad \begin{cases} \text{Ric}_{g_0} = \gamma g_0, \\ \gamma = \frac{a\lambda x_1 + 2a\nu - b\lambda}{ax_1 + b}, \\ \varphi(x) = -\gamma \frac{(a\lambda x + 2a\nu - b\lambda)(ax + b)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d}{(au + b)^{2d}(a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ a > 0, \mu = 0, \end{cases}$$

where

$$\begin{cases} -\frac{b}{a} = x_0 < x_1 < +\infty \text{ for } \lambda > 0 \text{ and } -\frac{b}{a} \geq -\frac{\nu}{\lambda}, \\ -\frac{b}{a} = x_0 < x_1 \leq -\frac{\nu}{\lambda} \text{ for } \lambda < 0 \text{ and } -\frac{\nu}{\lambda} > -\frac{b}{a}. \end{cases}$$

(III-2) $\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) with the fibre metrics complete at $t = 0$ on the domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)}\|w\|^2\}$ if and only if

$$(4.5) \quad \begin{cases} \text{Ric}_{g_0} = \gamma g_0, \\ \gamma = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{ax_1 + b}, \\ \varphi(x) = \frac{(a\lambda x + 2a\nu - b\lambda)(ax + b)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(au + b)^{2d}(a\lambda u + 2a\nu - b\lambda)^2} du, \quad x_0 < x < x_1, \\ a > 0, \mu < 0, \end{cases}$$

where

$$\begin{cases} -\frac{b}{a} = x_0 < x_1 < +\infty \text{ for } \lambda > 0 \text{ and } -\frac{b}{a} \geq -\frac{\nu}{\lambda}, \\ -\frac{b}{a} = x_0 < x_1 \leq -\frac{\nu}{\lambda} \text{ for } \lambda < 0 \text{ and } -\frac{\nu}{\lambda} > -\frac{b}{a}. \end{cases}$$

(IV) Let $r > 1$, $\varphi(x_1) = 0$ and $\varphi'(x_1) = -1$.

(IV-1) $\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) with the fibre metrics complete at $t = -\infty$ on the domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2\}$ if and only if

$$(4.6) \quad \begin{cases} \text{Ric}_{g_0} = r\lambda g_0, \\ \varphi(x) = r \frac{(ax + b)^{2n-1}(ax - b)}{x^{n-1}} \int_x^{x_1} \frac{u^{n-1}}{(au + b)^{2n-2}(au - b)^2} du, \quad -\frac{b}{a} = x_0 < x < x_1 = -\frac{(r+1)b}{(r-1)a}, \\ a > 0, b < 0, \lambda > 0, \mu = \nu = 0. \end{cases}$$

(IV - 2) $\tilde{g} = \frac{1}{f^2}g_F$ is an Einstein metric (namely $\text{Ric}_{\tilde{g}} = \mu \tilde{g}$) with the fibre metrics complete at $t = 0$ on the domain $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)}\|w\|^2\}$ if and only if

$$(4.7) \quad \left\{ \begin{array}{l} \text{Ric}_{g_0} = r\lambda g_0, \\ \mu = \frac{(ax_1+b)(r(ax_1+b)-(ax_1-b))}{x_1} < 0, \\ \varphi(x) = \frac{(ax+b)^{2n-1}(ax-b)}{x^{n-1}} \int_{x_1}^x \frac{u^{n-1}(\mu u - r(b+au)^2)}{(au+b)^{2n}(au-b)^2} du, \quad -\frac{b}{a} = x_0 < x < x_1 < -\frac{(r+1)b}{(r-1)a}, \\ a > 0, b < 0, \lambda > 0, \nu = 0. \end{array} \right.$$

Proof. By the well-known theorems of Bonnet and Myers, it follows that non-compact manifolds do not admit any Einstein metric with positive scalar curvature. Since M is not compact, we get $\mu \leq 0$. The following proof is divided into four parts.

Part (I)

From the metric g_F is smooth at $t = -\infty$, it follows that

$$0 = x_0 = \lim_{t \rightarrow -\infty} x(t) < \sup x(t) = x_1, \quad \varphi(0) = 0, \quad \varphi'(0) = 1.$$

Using (2.2), we obtain $\gamma = \lambda + \mu - 2a$ and

$$(4.8) \quad \varphi(x) = \frac{(1+ax)^{2d+1}(2a-\lambda+a\lambda x)}{(1+\lambda x)^d} m(x),$$

where

$$(4.9) \quad m(x) = \int_0^x \frac{(1+\lambda u)^d (\mu(1+\lambda u) - (\lambda + \mu - 2a)(1+au)^2)}{(1+au)^{2d+2}(a\lambda u + 2a - \lambda)^2} du.$$

(I - i) For the case of $a = \lambda$.

By (4.8), we get

$$\begin{aligned} \varphi(x) &= \frac{1}{a}(1+ax)^{d+2} \int_0^x \frac{\mu - (\lambda + \mu - 2a)(1+au)}{(1+au)^{d+3}} du \\ &= \frac{1}{a^2}(1+ax)^{d+2} \left(\frac{\mu}{d+2} \left(1 - \frac{1}{(1+ax)^{d+2}}\right) - \frac{\lambda + \mu - 2a}{d+1} \left(1 - \frac{1}{(1+ax)^{d+1}}\right) \right) \\ &= \frac{1}{a^2} \left(\frac{\mu}{d+2} - \frac{\lambda + \mu - 2a}{d+1} \right) (1+ax)^{d+2} + \frac{\lambda + \mu - 2a}{(d+1)a^2} (1+ax) - \frac{\mu}{(d+2)a^2} \\ &= -\frac{\mu + (d+2)(\lambda - 2a)}{(d+2)(d+1)a^2} (1+ax)^{d+2} + \frac{\lambda + \mu - 2a}{(d+1)a^2} (1+ax) - \frac{\mu}{(d+2)a^2} \\ &= \frac{(d+2)a - \mu}{(d+2)(d+1)a^2} (1+ax)^{d+2} + \frac{\mu - a}{(d+1)a^2} (1+ax) - \frac{\mu}{(d+2)a^2}. \end{aligned}$$

If $a > 0$, by $\mu \leq 0$, then $\mu - (\lambda + \mu - 2a)(1 + ax) = a(1 + (a - \mu)x) \geq 0$ for $x \geq 0$, thus $\varphi(x) > 0$ for $x > 0$, so whether x_1 is finite or infinite, there is

$$\int_0^{x_1} \frac{dx}{(1 + ax)\sqrt{\varphi(x)}} < +\infty.$$

This indicates that the fiber metrics for \tilde{g} are incomplete.

If $a < 0$, by $1 + ax > 0$, then $0 < x < -\frac{1}{a}$, thus $\frac{\mu - (\lambda + \mu - 2a)(1 + ax)}{a} > 0$ for $0 < x < -\frac{1}{a}$. If $0 < x_1 < -\frac{1}{a}$, then

$$\int_0^{x_1} \frac{dx}{(1 + ax)\sqrt{\varphi(x)}} < +\infty,$$

hence $x_1 = -\frac{1}{a}$.

From $\varphi(-\frac{1}{a}) = -\frac{\mu}{(d+2)a^2} \geq 0$, we have

$$\int_0^{x_1} \frac{dx}{(1 + ax)\sqrt{\varphi(x)}} = +\infty,$$

This show that fiber metrics are complete.

Let

$$t = \int_{-\frac{1}{a}}^{x(t)} \frac{dv}{\varphi(v)}, \quad F(t) = \int_{-\frac{1}{a}}^{x(t)} \frac{v dv}{\varphi(v)}$$

for $\mu < 0$, and

$$t = \int_{-\frac{1}{2a}}^{x(t)} \frac{dv}{\varphi(v)}, \quad F(t) = \int_{-\frac{1}{2a}}^{x(t)} \frac{v dv}{\varphi(v)}$$

for $\mu = 0$.

It follows by $\varphi(0) = 0$, $\varphi'(0) = 1$ and $\varphi(-\frac{1}{a}) = -\frac{\mu}{(d+2)a^2}$ that

$$\lim_{x \rightarrow 0^+} t = -\infty, \quad \lim_{x \rightarrow (-\frac{1}{a})^-} t = 0$$

for $\mu < 0$, and

$$\lim_{x \rightarrow 0^+} t = -\infty, \quad \lim_{x \rightarrow (-\frac{1}{a})^-} t = +\infty$$

for $\mu = 0$.

(I – ii) For the case of $\lambda = 2a$.

Using

$$\gamma = \lambda + \mu - 2a = \mu, \mu(1 + \lambda x) - \gamma(1 + ax)^2 = -a^2 \mu x^2$$

and (4.8), we get

$$\varphi(x) = -\frac{\mu x(1 + ax)^{2d+1}}{2(1 + \lambda x)^d} \int_0^x \frac{(1 + \lambda u)^d}{(1 + au)^{2d+2}} du,$$

so

$$\varphi(0) = \varphi'(0) = 0.$$

This conflicts with the smoothness of the metric \tilde{g} at $t = -\infty$.

(I – iii) For the case of $\lambda \neq a$ and $\lambda \neq 2a$.

(I – iii – 1) If $\frac{\lambda-2a}{a\lambda} = \frac{1}{a} - \frac{2}{\lambda} < -\frac{1}{\lambda} < -\frac{1}{a} < 0$ or $-\frac{1}{a} < -\frac{1}{\lambda} < \frac{1}{a} - \frac{2}{\lambda} < 0$, that is $0 < \lambda < a$ or $0 < a < \lambda < 2a$.

By $\mu \leq 0$, then

$$\gamma = \lambda + \mu - 2a < 0, 2a - \lambda + a\lambda x = a\lambda \left(x + \frac{2}{\lambda} - \frac{1}{a} \right) > 0, 1 + \lambda x > 0, 1 + ax > 0$$

for $x \geq 0$.

Let

$$q(x) = \mu(1 + \lambda x) - \gamma(1 + ax)^2.$$

By

$$q(0) = \mu - \gamma = 2a - \lambda > 0, q'(x) = \lambda\mu - 2a\gamma(1 + ax) = (2a - \lambda)(2a - \mu) - 2a^2\gamma x > 0$$

for $x \geq 0$, it follows that $q(x) > 0$ for $x \geq 0$.

Using (4.8), we get $\varphi(x) > 0$ for $x > 0$ and $\varphi(x) \sim Cx^{d+2}$ as $x \rightarrow +\infty$, thus

$$\int_0^{x_1} \frac{dx}{(1 + ax)\sqrt{\varphi(x)}} < +\infty.$$

This indicates that the fiber metrics for \tilde{g} are incomplete.

(I – iii – 2) If $\frac{\lambda-2a}{a\lambda} = \frac{1}{a} - \frac{2}{\lambda} < -\frac{1}{\lambda} < 0 < -\frac{1}{a}$ or $0 < -\frac{1}{a} < -\frac{1}{\lambda} < \frac{\lambda-2a}{a\lambda}$, that is $a < 0 < \lambda$ or $a < \lambda < 0$, then

$$\lambda - 2a - a\lambda x > 0$$

for $0 \leq x \leq c := -\frac{1}{a}$.

Let

$$q(x) = \mu(1 + \lambda x) - (\lambda + \mu - 2a)(1 + ax)^2.$$

Using $\mu \leq 0$, $q(0) = 2a - \lambda < 0$ and $q(c) = \mu(1 - \frac{\lambda}{a}) \leq 0$, we obtain:

For $\gamma = \lambda + \mu - 2a = 0$, $q(x) \leq 0$ for $x \in (0, c)$;

For $\gamma = \lambda + \mu - 2a < 0$, an equation $q(x) = 0$ has no root in the interval $(0, c)$, so $q(x) \leq 0$ for $x \in (0, c)$;

For $\gamma = \lambda + \mu - 2a > 0$, $\mu(1 + ax) \leq 0$ and $\gamma(1 + ax)^2 > 0$ for $x \in (0, c)$, therefore $q(x) \leq 0$ for $x \in (0, c)$.

By (4.8), the above indicates that $\varphi(x) > 0$ for $x \in (0, c)$.

Since $1 + ax > 0$, it follows that $0 \leq x < -\frac{1}{a}$, so $0 < x_1 \leq -\frac{1}{a}$. By the fiber metrics for \tilde{g} are complete, that is

$$\int_0^{x_1} \frac{dx}{(1 + ax)\sqrt{\varphi(x)}} = +\infty,$$

we have $x_1 = -\frac{1}{a}$.

For $\mu < 0$,

$$\begin{aligned} \lim_{x \rightarrow c^-} \varphi(x) &= \frac{2(a - \lambda)}{(1 - \frac{\lambda}{a})^d} \lim_{x \rightarrow c^-} \frac{\int_0^x \frac{(1 + \lambda u)^d (\mu(1 + \lambda u) - (\lambda + \mu - 2a)(1 + au)^2)}{(1 + au)^{2d+2} (a\lambda u + 2a - \lambda)^2} du}{(1 + ax)^{-(2d+1)}} \\ &= \frac{2(a - \lambda)}{(1 - \frac{\lambda}{a})^d} \frac{(1 - \frac{\lambda}{a})^{d+1} \mu}{(2a - 2\lambda)^2} \frac{1}{-(2d + 1)a} \\ &= \frac{-\mu}{2(2d + 1)a^2} > 0. \end{aligned}$$

Let

$$t = \int_c^{x(t)} \frac{dv}{\varphi(v)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

it follows that

$$\lim_{x \rightarrow 0^+} t = -\infty, \quad \lim_{x \rightarrow c^-} t = 0,$$

which implies that the g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : e^{\lambda\phi(z)}\|w\|^2 < 1\}$. Using

$$\int_0^c \frac{1}{(1+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we obtain that the fiber metrics of \tilde{g} is complete on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : e^{\lambda\phi(z)}\|w\|^2 < 1\}$.

For $\mu = 0$,

$$\begin{aligned} \lim_{x \rightarrow c^-} \frac{\varphi(x)}{(1+ax)^2} &= \frac{2(a-\lambda)}{(1-\frac{\lambda}{a})^d} \lim_{x \rightarrow c^-} \frac{\int_0^x \frac{(1+\lambda u)^d (2a-\lambda)}{(1+au)^{2d} (a\lambda u + 2a-\lambda)^2} du}{(1+ax)^{-(2d-1)}} \\ &= \frac{2(a-\lambda)}{(1-\frac{\lambda}{a})^d} \frac{(1-\frac{\lambda}{a})^d (2a-\lambda)}{(2a-2\lambda)^2} \frac{1}{-(2d-1)a} \\ &= \frac{\lambda-2a}{2(2d-1)a^2(1-\frac{\lambda}{a})} > 0. \end{aligned}$$

Let

$$t = \int_{\frac{c}{2}}^{x(t)} \frac{dv}{\varphi(v)}, \quad F(t) = \int_{\frac{c}{2}}^{x(t)} \frac{v dv}{\varphi(v)},$$

it follows that

$$\lim_{x \rightarrow 0^+} t = -\infty, \quad \lim_{x \rightarrow c^-} t = +\infty.$$

Using

$$\int_{\frac{c}{2}}^c \frac{1}{(1+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we obtain that the fiber metrics of \tilde{g} is complete on $M = \Omega \times \mathbb{C}^r$.

(I - iii - 3) If $\frac{\lambda-2a}{a\lambda} = \frac{1}{a} - \frac{2}{\lambda} < 0 < -\frac{1}{\lambda} < -\frac{1}{a}$ or $-\frac{1}{a} < 0 < -\frac{1}{\lambda} < \frac{1}{a} - \frac{2}{\lambda}$, then $\lambda < 2a < 0$ or $\lambda < 0 < a$.

By $1+ax > 0$ and $1+\lambda x > 0$, we have $0 \leq x < -\frac{1}{\lambda}$, so $0 < x_1 \leq c := -\frac{1}{\lambda}$.

Since $\varphi(x) > 0$ for $0 = x_0 < x < x_1$, by (4.8) and (4.9), thus $m(x_1) \geq 0$.

When $x_1 = -\frac{1}{\lambda}$ and

$$m(x_1) = \int_0^{x_1} \frac{(1+\lambda u)^d (\mu(1+\lambda u) - (\lambda + \mu - 2a)(1+au)^2)}{(1+au)^{2d+2} (a\lambda u + 2a - \lambda)^2} du = 0,$$

from (4.8), we get

$$\lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{1+\lambda x} = \frac{\mu + \lambda - 2a}{(d+1)\lambda^2} < 0.$$

This conflicts with $\varphi(x) > 0$ for $0 < x < x_1$.

When $x_1 < -\frac{1}{\lambda}$, or $x_1 = -\frac{1}{\lambda}$ and $m(x_1) > 0$, by (4.8) we have

$$\int_0^{x_1} \frac{dx}{(1+ax)\sqrt{\varphi(x)}} < +\infty.$$

This indicates that the fiber metrics for \tilde{g} are incomplete.

(I – iii – 4) If $0 < \frac{\lambda-2a}{a\lambda} = \frac{1}{a} - \frac{2}{\lambda} < -\frac{1}{\lambda} < -\frac{1}{a}$ or $-\frac{1}{a} < -\frac{1}{\lambda} < 0 < \frac{\lambda-2a}{a\lambda}$, then $2a < \lambda < a < 0$ or $0 < 2a < \lambda$. Since $1 + ax > 0$ and $1 + \lambda x > 0$ for $0 = x_0 < x < x_1$, it follows that $0 < x_1 \leq -\frac{1}{\lambda}$ for $2a < \lambda < a < 0$.

Let

$$\psi(x) = d\gamma(1 + ax)^2 - (d + 1)\mu(1 + \lambda x).$$

If $x_1 = +\infty$, using $\varphi(x) \sim Cx^{d+2}$ as $x \rightarrow +\infty$, then

$$\int_0^{x_1} \frac{dx}{(1 + ax)\sqrt{\varphi(x)}} < +\infty,$$

this is in contradiction with the completeness of the metric \tilde{g} .

If $x_1 < +\infty$, from the completeness of \tilde{g} , we get $\varphi(x_1) = \varphi'(x_1) = 0$, thus by (2.1) and (2.2), we have

$$(4.10) \quad \varphi(x) = \frac{(1 + ax)^{2d+1}}{(1 + \lambda x)^d} \int_x^{x_1} (u - x) \frac{(1 + \lambda u)^{d-1} \psi(u)}{(1 + au)^{2d+3}} du \quad (\text{by } \varphi(x_1) = 0 \text{ and } \varphi'(x_1) = 0)$$

and

$$\mu(1 + \lambda x_1) - \gamma(1 + ax_1)^2 = 0 \quad (\text{by } \varphi(x_1) = 0 \text{ and } \varphi'(x_1) = 0).$$

Then

$$\gamma = \frac{\mu(1 + \lambda x_1)}{(1 + ax_1)^2} \leq 0$$

and

$$\psi(x) = \mu \left\{ d \frac{(1 + \lambda x_1)}{(1 + ax_1)^2} (1 + ax)^2 - (d + 1)(1 + \lambda x) \right\},$$

which implies that

$$\psi(x_1) = -\mu(1 + \lambda x_1) \geq 0.$$

By $\mu \leq 0$, $\gamma = \lambda + \mu - 2a$ and $\lambda - 2a > 0$, we have

$$\psi(0) = d\gamma - (d + 1)\mu = d(\lambda - 2a) - \mu > 0,$$

thus $\psi(x) \geq 0$ for $0 \leq x \leq x_1$. So from (4.10), we have $\varphi(0) > 0$, this conflicts with $\varphi(0) = 0$.

Part (II)**Part (II) for $r = 1$**

(II – i) For the case of $a < 0$ and $\lambda < 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $x < \min\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$, so $x_1 \leq \min\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$.
For $x_0 > -\infty$, according to the completeness of the fibre metric for \tilde{g} , namely

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we have $\varphi(x_0) = \varphi'(x_0) = 0$. Then by (2.2), we obtain $\gamma = \frac{\mu(\nu+\lambda x_0)}{(b+ax_0)^2}$ and

$$(4.11) \quad \varphi(x) = -\frac{(b+ax)^{2d+1}(b\lambda - 2a\nu - a\lambda x)}{(\nu + \lambda x)^d} \int_{x_0}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b+au)^2)}{(b+au)^{2d+2}(b\lambda - 2a\nu - a\lambda u)^2} du$$

for $x \in (x_0, x_1)$.

(II – i – 1) If $-\frac{b}{a} \leq -\frac{\nu}{\lambda} \leq \frac{b}{a} - \frac{2\nu}{\lambda}$.

For $-\infty < x_0 < x < x_1 \leq c := -\frac{b}{a}$, using

$$b\lambda - 2a\nu - a\lambda x > 0, \frac{\nu + \lambda x_0}{(b+ax_0)^2} > 0, \nu + \lambda x - \frac{\nu + \lambda x_0}{(b+ax_0)^2}(b+ax)^2 > 0,$$

we have

$$\mu(\nu + \lambda x) - \gamma(b+ax)^2 = \mu \left(\nu + \lambda x - \frac{\nu + \lambda x_0}{(b+ax_0)^2}(b+ax)^2 \right) < 0$$

for $x_0 < x < x_1 \leq c$ only if $\mu < 0$. So $\varphi(x) > 0$ for $x \in (x_0, x_1)$ only if $\mu < 0$.

Using

$$\lim_{x \rightarrow c^-} \varphi(x) = \begin{cases} \frac{-\mu}{2(2d+1)a^2} > 0 \text{ (for } -\frac{b}{a} < -\frac{\nu}{\lambda}), \\ \frac{-\mu}{(d+2)a^2} > 0 \text{ (for } -\frac{b}{a} = -\frac{\nu}{\lambda}) \end{cases} \quad (\text{by (2.2)})$$

and fiber metrics for \tilde{g} are complete, we have $x_1 = c$.

Let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow x_0^+} \frac{\varphi(x)}{(x-x_0)^2} = -\frac{\mu}{2(b+ax_0)^2} \text{ (by (4.11)),}$$

then

$$\lim_{x \rightarrow x_0^+} t(x) = -\infty, \quad \lim_{x \rightarrow c^-} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2 < 1\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty, \quad \int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete.

For $x_0 = -\infty$, from (2.2), we obtain

$$\frac{(\nu + \lambda x)^d \varphi(x)}{(ax + b)^{2d+1}(a\lambda x + 2a\nu - b\lambda)} = C_1 + \int_{\widetilde{x}_0}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(au + b)^2)}{(au + b)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du,$$

where

$$\widetilde{x}_0 := x_1 - 1, C_1 = \frac{(\nu + \lambda \widetilde{x}_0)^d \varphi(\widetilde{x}_0)}{(a\widetilde{x}_0 + b)^{2d+1}(a\lambda \widetilde{x}_0 + 2a\nu - b\lambda)}.$$

Since

$$-\infty < C_1 + \int_{\widetilde{x}_0}^{-\infty} \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(au + b)^2)}{(au + b)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du < +\infty$$

and $a\lambda x + 2a\nu - b\lambda < 0$ for $-\infty < x < x_1$, it follows that

$$\lim_{x \rightarrow -\infty} \frac{(\nu + \lambda x)^d \varphi(x)}{(ax + b)^{2d+1}(a\lambda x + 2a\nu - b\lambda)} = C \leq 0.$$

If $C < 0$, then $\varphi(x) \sim C_2 x^{d+2}$ as $x \rightarrow -\infty$, this is in contradiction with

$$\int_{-\infty}^{\widetilde{x}_0} \frac{dx}{(b + ax)\sqrt{\varphi(x)}} = +\infty,$$

thus $C = 0$, so

$$(4.12) \quad \varphi(x) = \frac{(b + ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{-\infty}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(b + au)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du$$

for $x \in (-\infty, x_1)$.

As $x \rightarrow -\infty$, by

$$-\frac{(b + ax)^{2d+1}(b\lambda - 2a\nu - a\lambda x)}{(\nu + \lambda x)^d} \sim \frac{a^{2d+2}}{\lambda^{d-1}} x^{d+2},$$

$$\frac{(\nu + \lambda x)^d (\mu(\nu + \lambda x) - \gamma(b + ax)^2)}{(b + ax)^{2d+2}(a\lambda x + 2a\nu - b\lambda)^2} \sim \begin{cases} -\frac{\lambda^{d-2}\gamma}{a^{2d+2}} \frac{1}{x^{d+2}}, \gamma \neq 0, \\ \frac{\lambda^{d-1}\mu}{a^{2d+4}} \frac{1}{x^{d+3}}, \gamma = 0 \end{cases}$$

and (4.12), we get

$$\lim_{x \rightarrow -\infty} \frac{\varphi(x)}{x} = \frac{\gamma}{(d+1)\lambda} \quad (\gamma \neq 0)$$

and

$$\lim_{x \rightarrow -\infty} \varphi(x) = -\frac{\mu}{(d+2)a^2} \quad (\gamma = 0).$$

By

$$\int_{-\infty}^{\widetilde{x}_0} \frac{dx}{(b + ax)\sqrt{\varphi(x)}} = +\infty,$$

it follows that $\gamma = 0$. From (4.12), so $\varphi(x) > 0$ for $x \in (-\infty, c)$ only $\mu < 0$.

If $x_1 < c$, then

$$\int_{x_1-1}^{x_1} \frac{1}{(b + ax)\sqrt{\varphi(x)}} dx < +\infty,$$

according to the completeness of fiber metrics for \widetilde{g} , we get $x_1 = c$

Let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

then

$$\lim_{x \rightarrow -\infty} t(x) = -\infty, \quad \lim_{x \rightarrow c^-} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2 < 1\}$. Since

$$\int_{-\infty}^{c-1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty, \quad \int_{c-1}^c \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric of \tilde{g} is complete.

(II – i – 2) If $\frac{b}{a} - \frac{2\nu}{\lambda} < -\frac{\nu}{\lambda} < -\frac{b}{a}$, then $x_0 < x_1 \leq -\frac{\nu}{\lambda} < -\frac{b}{a}$. The following we prove that this situation is impossible.

If $\frac{b}{a} - \frac{2\nu}{\lambda} = x_0 < x_1 \leq -\frac{\nu}{\lambda} < -\frac{b}{a}$, by $\varphi(x_0) = \varphi'(x_0) = 0$ and (2.2), we get

$$\gamma = -\frac{\mu\lambda^2}{4a(a\nu - b\lambda)}, \quad \mu(\nu + \lambda x) - \gamma(b + ax)^2 = -a^2\gamma(x - x_0)^2.$$

Thus

$$\varphi(x) = -\frac{a\gamma}{\lambda} \frac{(b + ax)^{2d+1}(x - x_0)}{(\nu + \lambda x)^d} \int_{x_0}^x \frac{(\nu + \lambda u)^d}{(b + au)^{2d+2}} du$$

and

$$\int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx < +\infty.$$

This is in contradiction with the completeness of fiber metrics of \tilde{g} .

If $-\infty < x_0 < x_1 \leq -\frac{\nu}{\lambda} < -\frac{b}{a}$ and $x_0 \neq \frac{b}{a} - \frac{2\nu}{\lambda}$, by the completeness of fiber metrics of \tilde{g} , we get $\varphi(x_0) = \varphi'(x_0) = \varphi(x_1) = \varphi'(x_1) = 0$, so

$$\gamma = \frac{\mu(\nu + \lambda x_0)}{(b + ax_0)^2} = \frac{\mu(\nu + \lambda x_1)}{(b + ax_1)^2} \leq 0,$$

$$\mu(\nu + \lambda x) - \gamma(ax + b)^2 = a^2\gamma(x - x_0)(x_1 - x)$$

and

$$\psi(x) := d\gamma(ax + b)^2 - (d+1)\mu(\nu + \lambda x) = -da^2\gamma(x - x_0)(x_1 - x) - \mu(\nu + \lambda x) \geq 0$$

for $x_0 < x < x_1$. Using (2.1), we have

$$\varphi(x) = \frac{(b + ax)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_0}^x (x - u) \frac{(\nu + \lambda u)^{d-1} \psi(u)}{(b + au)^{2d+3}} du \quad (\text{by } \varphi(x_0) = 0 \text{ and } \varphi'(x_0) = 0),$$

which implies that $\varphi(x_1) > 0$, this is in contradiction with the completeness of \tilde{g} .

If $x_0 = -\infty$, by (4.12), we have $\gamma = 0$ and $\mu < 0$. By $\varphi(x_1) = 0$, $\varphi'(x_1) = 0$ and (2.1), we have

$$\varphi(x) = \frac{(b + ax)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_1}^x (x - u) \frac{(\nu + \lambda u)^{d-1} \psi(u)}{(b + au)^{2d+3}} du$$

which implies that $\varphi(x) \sim C_3 x^{d+2}$ as $x \rightarrow -\infty$, this is in contradiction with the completeness of \tilde{g} .

(II – ii) For the case of $a > 0$ and $\lambda > 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $x > \max\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$, so $x_1 > x_0 \geq \max\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$.

If $x_1 < +\infty$, by

$$\int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we get $\varphi(x_1) = \varphi'(x_1) = 0$. Then by (2.2), we obtain $\gamma = \frac{\mu(\nu+\lambda x_1)}{(b+ax_1)^2}$ and

$$(4.13) \quad \varphi(x) = \frac{(b+ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(b + au)^{2d+2}(b\lambda - 2a\nu - a\lambda u)^2} du$$

for $x \in (x_0, x_1)$. It follows from $\varphi \neq 0$ that $\mu < 0$.

(II – ii – 1) If $-\frac{b}{a} \geq -\frac{\nu}{\lambda} \geq \frac{b}{a} - \frac{2\nu}{\lambda}$, then

$$a\lambda x + 2a\nu - b\lambda > 0, \frac{\nu + \lambda x_1}{(b + ax_1)^2} > 0, \nu + \lambda x - \frac{\nu + \lambda x_1}{(b + ax_1)^2}(b + ax)^2 > 0$$

for $c := -\frac{b}{a} \leq x_0 < x < x_1 < +\infty$. Thus

$$\mu(\nu + \lambda x) - \gamma(b + ax)^2 = \mu \left(\nu + \lambda x - \frac{\nu + \lambda x_0}{(b + ax_0)^2}(b + ax)^2 \right) < 0,$$

so $\varphi(x) > 0$ for $c < x < x_1$ from (4.13).

Using

$$\lim_{x \rightarrow c^+} \varphi(x) = \begin{cases} \frac{-\mu}{2(2d+1)a^2} > 0 \text{ (for } -\frac{b}{a} < -\frac{\nu}{\lambda}), \\ \frac{-\mu}{(d+2)a^2} > 0 \text{ (for } -\frac{b}{a} = -\frac{\nu}{\lambda}) \end{cases} \quad (\text{by (2.2)})$$

and fiber metrics for \tilde{g} are complete, we have $x_0 = c$.

Let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{(x - x_1)^2} = -\frac{\mu}{2(b + ax_1)^2},$$

then

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty, \quad \int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete.

For $x_1 = +\infty$, from (2.2), we obtain

$$\frac{(\nu + \lambda x)^d \varphi(x)}{(ax + b)^{2d+1}(a\lambda x + 2a\nu - b\lambda)} = C_1 + \int_{\tilde{x}_0}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(au + b)^2)}{(au + b)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du,$$

where

$$\tilde{x}_0 := x_0 + 1, \quad C_1 = \frac{(\nu + \lambda \tilde{x}_0)^d \varphi(\tilde{x}_0)}{(a\tilde{x}_0 + b)^{2d+1}(a\lambda \tilde{x}_0 + 2a\nu - b\lambda)}.$$

Since

$$-\infty < C_1 + \int_{\tilde{x}_0}^{+\infty} \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(au + b)^2)}{(au + b)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du < +\infty,$$

it follows that

$$\lim_{x \rightarrow +\infty} \frac{(\nu + \lambda x)^d \varphi(x)}{(ax + b)^{2d+1} (a\lambda x + 2a\nu - b\lambda)} = C \geq 0.$$

If $C > 0$, then $\varphi(x) \sim C_4 x^{d+2}$ ($C_4 > 0$) as $x \rightarrow +\infty$, this is in contradiction with

$$\int_{\tilde{x}_0}^{+\infty} \frac{dx}{(b + ax)\sqrt{\varphi(x)}} = +\infty,$$

thus $C = 0$, so

$$(4.14) \quad \varphi(x) = \frac{(b + ax)^{2d+1} (a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{+\infty}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(b + au)^{2d+2} (a\lambda u + 2a\nu - b\lambda)^2} du$$

for $x \in (x_0, +\infty)$.

As $x \rightarrow +\infty$,

$$\frac{(b + ax)^{2d+1} (a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \sim \frac{a^{2d+2}}{\lambda^{d-1}} x^{d+2}$$

and

$$\frac{(\nu + \lambda x)^d (\mu(\nu + \lambda x) - \gamma(b + ax)^2)}{(b + ax)^{2d+2} (a\lambda x + 2a\nu - b\lambda)^2} \sim \begin{cases} -\frac{\lambda^{d-2}\gamma}{a^{2d+2}} \frac{1}{x^{d+2}}, \gamma \neq 0, \\ \frac{\lambda^{d-1}\mu}{a^{2d+4}} \frac{1}{x^{d+3}}, \gamma = 0, \end{cases}$$

it follows from (4.14) that

$$\lim_{x \rightarrow +\infty} \frac{\varphi(x)}{x} = \frac{\gamma}{(d+1)\lambda} \quad (\gamma \neq 0)$$

and

$$\lim_{x \rightarrow +\infty} \varphi(x) = -\frac{\mu}{(d+2)a^2} \quad (\gamma = 0).$$

By

$$\int_{\tilde{x}_0}^{+\infty} \frac{dx}{(b + ax)\sqrt{\varphi(x)}} = +\infty,$$

it follows that $\gamma = 0$, then $\varphi(x) > 0$ for $x \in (x_0, +\infty)$ only $\mu < 0$ from (4.14).

If $x_0 > c$, then

$$\int_{x_0}^{x_0+1} \frac{1}{(b + ax)\sqrt{\varphi(x)}} dx < +\infty,$$

according to the completeness of fiber metrics for \tilde{g} , we get $x_0 = c$

Let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

then

$$\lim_{x \rightarrow +\infty} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)} \|w\|^2\}$. Since

$$\int_{c+1}^{+\infty} \frac{1}{(b + ax)\sqrt{\varphi(x)}} dx = +\infty, \quad \int_c^{c+1} \frac{1}{(b + ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric of \tilde{g} is complete.

(II – ii – 2) If $\frac{b}{a} - \frac{2\nu}{\lambda} > -\frac{\nu}{\lambda} > -\frac{b}{a}$, then $x_1 > x_0 \geq -\frac{\nu}{\lambda} > -\frac{b}{a}$. We prove that this situation is impossible.

If $\frac{b}{a} - \frac{2\nu}{\lambda} = x_1 > x_0 \geq -\frac{\nu}{\lambda} > -\frac{b}{a}$, by $\varphi(x_1) = \varphi'(x_1) = 0$ and (2.2), we get

$$\gamma = -\frac{\mu\lambda^2}{4a(av - b\lambda)}, \quad \mu(\nu + \lambda x) - \gamma(b + ax)^2 = -a^2\gamma(x - x_1)^2.$$

Thus

$$\varphi(x) = \frac{a\gamma}{\lambda} \frac{(b + ax)^{2d+1}(x_1 - x)}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d}{(b + au)^{2d+2}} du$$

and

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b + ax)\sqrt{\varphi(x)}} dx < +\infty.$$

This is in contradiction with the completeness of fiber metrics of \tilde{g} .

If $-\frac{b}{a} < -\frac{\nu}{\lambda} \leq x_0 < x_1 < +\infty$ and $x_1 \neq \frac{b}{a} - \frac{2\nu}{\lambda}$, by the completeness of fiber metrics of \tilde{g} , we get $\varphi(x_0) = \varphi'(x_0) = \varphi(x_1) = \varphi'(x_1) = 0$, so

$$\gamma = \frac{\mu(\nu + \lambda x_0)}{(b + ax_0)^2} = \frac{\mu(\nu + \lambda x_1)}{(b + ax_1)^2} \leq 0,$$

$$\mu(\nu + \lambda x) - \gamma(ax + b)^2 = a^2\gamma(x - x_0)(x_1 - x)$$

and

$$\psi(x) := d\gamma(ax + b)^2 - (d + 1)\mu(\nu + \lambda x) = -da^2\gamma(x - x_0)(x_1 - x) - \mu(\nu + \lambda x) \geq 0$$

for $x_0 < x < x_1$. Using (2.1), we have

$$\varphi(x) = \frac{(b + ax)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_1}^x (x - u) \frac{(\nu + \lambda u)^{d-1} \psi(u)}{(b + au)^{2d+3}} du \quad (\text{by } \varphi(x_1) = 0 \text{ and } \varphi'(x_1) = 0),$$

which implies that $\varphi(x_0) > 0$, this is in contradiction with the completeness of \tilde{g} .

If $x_1 = +\infty$, by (4.14), we have $\gamma = 0$ and $\mu < 0$. By $\varphi(x_0) = 0$, $\varphi'(x_0) = 0$ and (2.1), we have

$$\varphi(x) = \frac{(b + ax)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_0}^x (x - u) \frac{(\nu + \lambda u)^{d-1} \psi(u)}{(b + au)^{2d+3}} du$$

which implies that $\varphi(x) \sim C_5 x^{d+2}$ as $x \rightarrow +\infty$, this is in contradiction with the completeness of \tilde{g} .

(II – iii) For the case of $a > 0$ and $\lambda < 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $-\frac{b}{a} < x < -\frac{\nu}{\lambda}$, so $-\frac{b}{a} \leq x_0 < x_1 \leq -\frac{\nu}{\lambda} < \frac{b}{a} - \frac{2\nu}{\lambda}$.

By

$$\int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b + ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we get $\varphi(x_1) = \varphi'(x_1) = 0$. Then by (2.2), we obtain $\gamma = \frac{\mu(\nu + \lambda x_1)}{(b + ax_1)^2}$ and

$$(4.15) \quad \varphi(x) = \frac{(b + ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(b + au)^{2d+2}(b\lambda - 2a\nu - a\lambda u)^2} du$$

for $x \in (x_0, x_1)$. From (4.15) and $\varphi \neq 0$, it follows that $\mu < 0$.

Note that when $c := -\frac{b}{a} \leq x_0 < x < x_1$,

$$a\lambda x + 2a\nu - b\lambda > 0, \frac{\nu + \lambda x_1}{(b + ax_1)^2} \geq 0, \nu + \lambda x - \frac{\nu + \lambda x_1}{(b + ax_1)^2}(b + ax)^2 > 0,$$

so

$$\mu(\nu + \lambda x) - \gamma(b + ax)^2 = \mu \left(\nu + \lambda x - \frac{\nu + \lambda x_0}{(b + ax_0)^2}(b + ax)^2 \right) < 0,$$

thus $\varphi(x) > 0$ for $c < x < x_1$ by (4.15).

Using

$$\lim_{x \rightarrow c^+} \varphi(x) = \frac{-\mu}{2(2d+1)a^2} > 0$$

and fiber metrics for \tilde{g} are complete, we have $x_0 = c$.

Let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{(x - x_1)^2} = -\frac{\mu}{2(b + ax_1)^2},$$

then

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty, \quad \int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete.

(II – iv) For the case of $a < 0$ and $\lambda > 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $-\frac{\nu}{\lambda} < x < -\frac{b}{a}$, so $\frac{b}{a} - \frac{2\nu}{\lambda} < -\frac{\nu}{\lambda} \leq x_0 < x_1 \leq -\frac{b}{a}$.
By

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we get $\varphi(x_0) = \varphi'(x_0) = 0$. Then by (2.2), we obtain $\gamma = \frac{\mu(\nu+\lambda x_0)}{(b+ax_0)^2}$ and

$$(4.16) \quad \varphi(x) = \frac{(b+ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_0}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(b + au)^{2d+2}(b\lambda - 2a\nu - a\lambda u)^2} du$$

for $x \in (x_0, x_1)$. From (4.16) and $\varphi \neq 0$, it follows that $\mu < 0$.

Note that when $x_0 < x < c := -\frac{b}{a}$,

$$b\lambda - 2a\nu - a\lambda x > 0, \frac{\nu + \lambda x_0}{(b + ax_0)^2} \geq 0, \nu + \lambda x - \frac{\nu + \lambda x_0}{(b + ax_0)^2}(b + ax)^2 > 0,$$

so

$$\mu(\nu + \lambda x) - \gamma(b + ax)^2 = \mu \left(\nu + \lambda x - \frac{\nu + \lambda x_0}{(b + ax_0)^2}(b + ax)^2 \right) < 0$$

which implies that $\varphi(x) > 0$ for $x_0 < x < c$.

Using

$$\lim_{x \rightarrow c^-} \varphi(x) = \frac{-\mu}{2(2d+1)a^2} > 0$$

and fiber metrics for \tilde{g} are complete, we have $x_1 = c$.

Let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow x_0^+} \frac{\varphi(x)}{(x-x_0)^2} = -\frac{\mu}{2(b+ax_0)^2},$$

then

$$\lim_{x \rightarrow x_0^+} t(x) = -\infty, \quad \lim_{x \rightarrow c^-} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2 < 1\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty, \quad \int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete.

Part (II) for $r > 1$

Since $a \neq 0$, from (2.3), we get $\nu = 0$.

By Remark 2.1, when $\nu = 0$ and $r > 1$, there must be $\lambda > 0$.

(II – i) For the case of $a > 0$ and $\lambda > 0$.

By $ax + b > 0$ and $\nu + \lambda x = \lambda x > 0$, we get $x > \max\{-\frac{b}{a}, 0\}$, so $x_1 > x_0 \geq \max\{-\frac{b}{a}, 0\}$.

If $x_1 < +\infty$, by

$$\int_{\frac{x_0+x_1}{2}}^{x_1} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we get $\varphi(x_1) = \varphi'(x_1) = 0$. Then by (2.3), we obtain $\mu = \frac{r(ax_1+b)^2}{x_1} > 0$. This is in contradiction with $\mu \leq 0$.

For $x_1 = +\infty$, from (2.3), we obtain

$$\frac{x^{n-1}\varphi(x)}{(ax+b)^{2n-1}(ax-b)} = C_1 + \int_{\tilde{x}_0}^x \frac{u^{n-1}(\mu u - r(b+au)^2)}{(b+au)^{2n}(au-b)^2} du,$$

where

$$\tilde{x}_0 := x_0 + 1, \quad C_1 = \frac{\tilde{x}_0^{n-1}\varphi(\tilde{x}_0)}{(a\tilde{x}_0+b)^{2n-1}(a\tilde{x}_0-b)}.$$

Since

$$-\infty < C_1 + \int_{\tilde{x}_0}^{+\infty} \frac{u^{n-1}(\mu u - r(b+au)^2)}{(b+au)^{2n}(au-b)^2} du < +\infty,$$

it follows that

$$\lim_{x \rightarrow +\infty} \frac{x^{n-1}\varphi(x)}{(ax+b)^{2n-1}(ax-b)} = C.$$

If $C \neq 0$, then $\varphi(x) \sim C_2 x^{n+1}$ as $x \rightarrow +\infty$, this is in contradiction with

$$\int_{\tilde{x}_0}^{+\infty} \frac{dx}{(b+ax)\sqrt{\varphi(x)}} = +\infty,$$

thus $C = 0$, so

$$(4.17) \quad \varphi(x) = \frac{(ax+b)^{2n-1}(ax-b)}{x^{n-1}} \int_{+\infty}^x \frac{u^{n-1}(\mu u - r(b+au)^2)}{(b+au)^{2n}(au-b)^2} du$$

for $x \in (x_0, +\infty)$.

As $x \rightarrow +\infty$,

$$\frac{(ax+b)^{2n-1}(ax-b)}{x^{n-1}} \sim a^{2n} x^{n+1}$$

and

$$\frac{x^{n-1}(\mu x - r(b+ax)^2)}{(b+ax)^{2n}(ax-b)^2} \sim \frac{-r}{a^{2n}} \frac{1}{x^{n+1}}$$

it follows from (4.17) that

$$\lim_{x \rightarrow +\infty} \frac{\varphi(x)}{x} = \frac{r}{n}.$$

So

$$\int_{\tilde{x}_0}^{+\infty} \frac{dx}{(b+ax)\sqrt{\varphi(x)}} < +\infty,$$

this is in contradiction with the completeness of fiber metrics of \tilde{g} .

(II – ii) For the case of $a < 0$ and $\lambda > 0$.

By $ax+b > 0$ and $\nu + \lambda x = \lambda x > 0$, we get $0 < x < -\frac{b}{a}$, so $b > 0$ and $\frac{b}{a} < 0 \leq x_0 < x_1 \leq -\frac{b}{a}$.

By

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we get $\varphi(x_0) = \varphi'(x_0) = 0$. Then by (2.3), we obtain $\mu = \frac{r(ax_0+b)^2}{x_0} > 0$, this is in contradiction with $\mu \leq 0$.

Part (III)

By (2.2), using $\varphi(x_1) = 0$, we get

$$(4.18) \quad \varphi(x) = \frac{(a\lambda x + 2a\nu - b\lambda)(ax+b)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(au+b)^2)}{(au+b)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du$$

for $x_0 < x < x_1$. Then

$$-1 = \varphi'(x_1) = \lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x - x_1} = \frac{\mu(\lambda x_1 + \nu) - \gamma(ax_1 + b)^2}{(ax_1 + b)(a\lambda x_1 + 2a\nu - b\lambda)}.$$

So

$$\gamma = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b + ax_1}.$$

(III – i) For the case of $a < 0$ and $\lambda < 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $x < \min\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$, so $x_1 \leq \min\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$. For $x_0 > -\infty$, according to the completeness of the fibre metric for \tilde{g} , namely

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

we have $\varphi(x_0) = \varphi'(x_0) = 0$. Then by (2.2), we obtain $\gamma = \frac{\mu(\nu+\lambda x_0)}{(b+ax_0)^2} < 0$, $\mu < 0$ and

$$(4.19) \quad \varphi(x) = -\frac{(b+ax)^{2d+1}(b\lambda - 2a\nu - a\lambda x)}{(\nu + \lambda x)^d} \int_{x_0}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b+au)^2)}{(b+au)^{2d+2}(b\lambda - 2a\nu - a\lambda u)^2} du$$

for $x \in (x_0, x_1)$.

(III – i – 1) If $-\frac{b}{a} \leq -\frac{\nu}{\lambda} \leq \frac{b}{a} - \frac{2\nu}{\lambda}$.

Note that when $-\infty < x_0 < x < x_1 \leq c := -\frac{b}{a}$, by

$$b\lambda - 2a\nu - a\lambda x > 0, \frac{\nu + \lambda x_0}{(b+ax_0)^2} > 0, \nu + \lambda x - \frac{\nu + \lambda x_0}{(b+ax_0)^2}(b+ax)^2 > 0$$

and (4.19), we get $\varphi(x_1) > 0$. This is in contradiction with $\varphi(x_1) = 0$.

For $x_0 = -\infty$, according to the proof of (II – i – 1), we get $\gamma = 0$, $\mu < 0$ and

$$(4.20) \quad \varphi(x) = \frac{(b+ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{-\infty}^x \frac{\mu(\nu + \lambda u)^{d+1}}{(b+au)^{2d+2}(a\lambda u + 2a\nu - b\lambda)^2} du$$

for $x \in (-\infty, x_1)$, which implies that $\varphi(x_1) > 0$. This conflicts with $\varphi(x_1) = 0$.

(III – i – 2) If $\frac{b}{a} - \frac{2\nu}{\lambda} < -\frac{\nu}{\lambda} < -\frac{b}{a}$, then $\frac{b}{a} - \frac{2\nu}{\lambda} \leq x_0 < x_1 \leq -\frac{\nu}{\lambda} < -\frac{b}{a}$ or $x_0 < x_1 \leq \frac{b}{a} - \frac{2\nu}{\lambda} < -\frac{\nu}{\lambda} < -\frac{b}{a}$ or $x_0 < \frac{b}{a} - \frac{2\nu}{\lambda} < x_1 \leq -\frac{\nu}{\lambda} < -\frac{b}{a}$.

If $-\infty < x_0$, and $\frac{b}{a} - \frac{2\nu}{\lambda} < x_1 \leq -\frac{\nu}{\lambda} < -\frac{b}{a}$, by

$$\gamma = \mu \frac{\nu + \lambda x_1}{(b+ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b+ax_1} = \mu \frac{\nu + \lambda x_0}{(b+ax_0)^2} < 0,$$

we get

$$\mu(\nu + \lambda x) - \gamma(ax + b)^2 < 0$$

for $x_0 < x < x_1$. So

$$\psi(x) := d\gamma(ax + b)^2 - (d+1)\mu(\nu + \lambda x) = -d(\mu(\nu + \lambda x) - \gamma(ax + b)^2) - \mu(\nu + \lambda x) > 0$$

for $x_0 < x < x_1$. Using (2.1), we have

$$\varphi(x) = \frac{(b+ax)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_0}^x (x-u) \frac{(\nu + \lambda u)^{d-1} \psi(u)}{(b+au)^{2d+3}} du \quad (\text{by } \varphi(x_0) = 0 \text{ and } \varphi'(x_0) = 0),$$

which implies that $\varphi(x_1) > 0$, this conflicts with $\varphi(x_1) = 0$.

If $-\infty < x_0 < x_1 \leq c := \frac{b}{a} - \frac{2\nu}{\lambda} < -\frac{\nu}{\lambda} < -\frac{b}{a}$, by

$$\gamma = \frac{\mu(\nu + \lambda x_0)}{(b+ax_0)^2} < 0, \mu(\nu + \lambda c) - \gamma(ac + b)^2 = -\frac{\mu(\nu a - b\lambda)(a\lambda x_0 + 2\nu a - b\lambda)^2}{a(ax_0 + b)^2 \lambda^2} < 0$$

and

$$b\lambda - 2a\nu - a\lambda x > 0, \mu(\nu + \lambda x) - \gamma(ax + b)^2 < 0$$

for $x_0 < x < c$. From (4.19) we have $\varphi(x_1) > 0$, this conflicts with $\varphi(x_1) = 0$.

If $-\infty = x_0 < x_1 \leq -\frac{\nu}{\lambda} < -\frac{b}{a}$, by (4.12), we have $\gamma = 0$ and $\mu < 0$. Using

$$\gamma = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b + ax_1},$$

it follows that $\frac{b}{a} - \frac{2\nu}{\lambda} < x_1$. From (4.18), we get $\varphi(x) < 0$ for $x < \frac{b}{a} - \frac{2\nu}{\lambda}$, this conflicts with $\varphi(x) > 0$ for $x_0 < x < x_1$.

(III – ii) For the case of $a > 0$ and $\lambda > 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $x > \max\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$, so $x_1 > x_0 \geq \max\{-\frac{b}{a}, -\frac{\nu}{\lambda}\}$,

$$(4.21) \quad \varphi(x) = \frac{(b + ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_1}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(b + au)^{2d+2}(b\lambda - 2a\nu - a\lambda u)^2} du$$

for $x \in (x_0, x_1)$, and

$$\gamma = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b + ax_1}.$$

(III – ii – 1) If $-\frac{b}{a} \geq -\frac{\nu}{\lambda} \geq \frac{b}{a} - \frac{2\nu}{\lambda}$.

Let

$$q(x) = \mu(\nu + \lambda x) - \gamma(b + ax)^2.$$

Note that when $c := -\frac{b}{a} < x < x_1$, by $\mu \leq 0$, $q(-\frac{b}{a}) \leq 0$ and $q(x_1) < 0$, we get

$$a\lambda x + 2a\nu - b\lambda > 0,$$

$$\mu(\nu + \lambda x) - \gamma(b + ax)^2 \leq 0 \quad (\gamma \leq 0)$$

and

$$\mu(\nu + \lambda x) - \gamma(b + ax)^2 \leq -\gamma(b + ax)^2 \leq 0 \quad (\gamma > 0),$$

so $\varphi(x) > 0$ for $x \in (c, x_1)$ by (4.21).

Using

$$\lim_{x \rightarrow c^+} \varphi(x) = \begin{cases} \frac{-\mu}{2(2d+1)a^2} \quad (\text{for } \frac{b}{a} < \frac{\nu}{\lambda}), \\ \frac{-\mu}{(d+2)a^2} \quad (\text{for } \frac{b}{a} = \frac{\nu}{\lambda}), \end{cases}$$

$$\lim_{x \rightarrow c^+} \frac{\varphi(x)}{(b + ax)^2} = \frac{\gamma}{2(2d-1)a(a\nu - b\lambda)} > 0 \quad (\text{for } \mu = 0 \text{ and } \frac{b}{a} < \frac{\nu}{\lambda}),$$

$$\lim_{x \rightarrow c^+} \frac{\varphi(x)}{b + ax} = \frac{\gamma}{(d+1)a\lambda} > 0 \quad (\text{for } \mu = 0 \text{ and } \frac{b}{a} = \frac{\nu}{\lambda})$$

and fiber metrics for \tilde{g} are complete, we have $x_0 = c$.

For $\mu < 0$, let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow c^+} \varphi(x) = \frac{-\mu}{2(2d+1)a^2} \quad (\text{for } \frac{b}{a} < \frac{\nu}{\lambda}), \quad \lim_{x \rightarrow c^+} \varphi(x) = \frac{-\mu}{(d+2)a^2} \quad (\text{for } \frac{b}{a} = \frac{\nu}{\lambda})$$

and $\lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x_1 - x} = 1$, we have

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete at $x = x_0$.

For $\mu = 0$, let

$$t = \int_{\frac{x_0+x_1}{2}}^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_{\frac{x_0+x_1}{2}}^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow c^+} \frac{\varphi(x)}{(b+ax)^2} = \frac{\gamma}{2(2d-1)a(av-b\lambda)} > 0 \quad (\text{for } \frac{b}{a} < \frac{\nu}{\lambda}), \quad \lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x_1-x} = 1$$

and

$$\lim_{x \rightarrow c^+} \frac{\varphi(x)}{b+ax} = \frac{\gamma}{(d+1)a\lambda} > 0 \quad (\text{for } \frac{b}{a} = \frac{\nu}{\lambda}),$$

we get

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = -\infty.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete at $x = x_0$.

(III – ii – 2) If $\frac{b}{a} - \frac{2\nu}{\lambda} > -\frac{\nu}{\lambda} > -\frac{b}{a}$, then $\frac{b}{a} - \frac{2\nu}{\lambda} \geq x_1 > x_0 \geq -\frac{\nu}{\lambda} > -\frac{b}{a}$ or $x_1 > x_0 \geq \frac{b}{a} - \frac{2\nu}{\lambda} > -\frac{\nu}{\lambda} > -\frac{b}{a}$ or $x_1 > \frac{b}{a} - \frac{2\nu}{\lambda} > x_0 \geq -\frac{\nu}{\lambda} > -\frac{b}{a}$.

If $\frac{b}{a} - \frac{2\nu}{\lambda} = x_1 > x_0 \geq -\frac{\nu}{\lambda} > -\frac{b}{a}$, by $\mu \leq 0$, $\varphi(x_1) = 0$, $\varphi'(x_1) = -1$ and (2.2), we get

$$\gamma = \frac{\mu \lambda^2}{4a(b\lambda - a\nu)} \leq 0, \quad \mu(\nu + \lambda x) - \gamma(b+ax)^2 = -a^2\gamma(x-x_1)^2.$$

Thus

$$\varphi(x) = \frac{(b+ax)^{2d+1}(x_1-x)}{(\nu+\lambda x)^d} \left(\frac{(\nu+\lambda x_1)^d}{(b+ax_1)^{2d+1}} + \frac{a\gamma}{\lambda} \int_{x_1}^x \frac{(\nu+\lambda u)^d}{(b+au)^{2d+2}} du \right) > 0 \quad (x_0 \leq x < x_1)$$

and

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx < +\infty.$$

This is in contradiction with the completeness of fiber metrics of \tilde{g} .

If $\frac{b}{a} - \frac{2\nu}{\lambda} > x_1 > x_0 \geq -\frac{\nu}{\lambda} > -\frac{b}{a}$, then

$$\gamma = \mu \frac{\nu + \lambda x_1}{(b+ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b+ax_1}, \quad b\lambda - 2a\nu - a\lambda x > 0$$

for $x \leq x_1$. By the completeness of fiber metrics of \tilde{g} , we get $\varphi(x_0) = \varphi'(x_0) = 0$, thus by (2.2),

$$\gamma = \mu \frac{\nu + \lambda x_0}{(b+ax_0)^2} = \mu \frac{\nu + \lambda x_1}{(b+ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b+ax_1},$$

namely,

$$\mu \frac{\nu + \lambda x_0}{(b + ax_0)^2} - \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} = \frac{a\lambda x_1 + 2a\nu - b\lambda}{b + ax_1}.$$

Since

$$\frac{d}{dx} \frac{\nu + \lambda x}{(b + ax)^2} = \frac{b\lambda - 2a\nu - a\lambda x}{(ax + b)^3} > 0$$

for $x_0 \leq x \leq x_1$ and $\mu \leq 0$, it follows that

$$\mu \frac{\nu + \lambda x_0}{(b + ax_0)^2} - \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} \geq 0,$$

this conflicts with $\frac{a\lambda x_1 + 2a\nu - b\lambda}{b + ax_1} < 0$.

If $-\frac{b}{a} < -\frac{\nu}{\lambda} \leq x_0$ and $\frac{b}{a} - \frac{2\nu}{\lambda} < x_1$, by

$$\gamma = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b + ax_1} = \mu \frac{\nu + \lambda x_0}{(b + ax_0)^2} < 0,$$

we get

$$\mu(\nu + \lambda x) - \gamma(ax + b)^2 < 0$$

for $x_0 < x < x_1$. So

$$\psi(x) := d\gamma(ax + b)^2 - (d+1)\mu(\nu + \lambda x) = -d(\mu(\nu + \lambda x) - \gamma(ax + b)^2) - \mu(\nu + \lambda x) > 0$$

for $x_0 < x < x_1$. Using (2.1), we have

$$\varphi(x) = \frac{(b + ax)^{2d+1}}{(\nu + \lambda x)^d} \int_{x_0}^x (x - u) \frac{(\nu + \lambda u)^{d-1} \psi(u)}{(b + au)^{2d+3}} du \quad (\text{by } \varphi(x_0) = 0 \text{ and } \varphi'(x_0) = 0),$$

which implies that $\varphi(x_1) > 0$, this conflicts with $\varphi(x_1) = 0$.

(III – iii) For the case of $a > 0$ and $\lambda < 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $-\frac{b}{a} < x < -\frac{\nu}{\lambda}$, so $-\frac{b}{a} \leq x_0 < x_1 \leq -\frac{\nu}{\lambda} < \frac{b}{a} - \frac{2\nu}{\lambda}$.
Let

$$q(x) := \mu(\nu + \lambda x) - \gamma(b + ax)^2.$$

Note that when $c := -\frac{b}{a} \leq x_0 < x < x_1$, by

$$\gamma = \mu \frac{\nu + \lambda x_1}{(b + ax_1)^2} + \frac{a\lambda x_1 + 2a\nu - b\lambda}{b + ax_1}, \quad q(c) \leq 0, \quad q(x_1) < 0, \quad \mu \leq 0,$$

we have

$$q(x) = \mu(\nu + \lambda x) - \gamma(b + ax)^2 < 0, \quad a\lambda x + 2a\nu - b\lambda > 0,$$

for $x \in (c, x_1)$. From (4.18), so $\varphi(x) > 0$ for $x \in (c, x_1)$.

Using

$$\begin{aligned} \lim_{x \rightarrow c^+} \varphi(x) &= \frac{-\mu}{2(2d+1)a^2} > 0 \quad (\mu < 0), \\ \lim_{x \rightarrow c^+} \frac{\varphi(x)}{(b + ax)^2} &= \frac{\gamma}{2(2d-1)a(a\nu - b\lambda)} > 0 \quad (\mu = 0) \end{aligned}$$

and fiber metrics for \tilde{g} are complete, we have $x_0 = c$.

For $\mu < 0$, let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{vdv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x_1 - x} = 1,$$

then

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete.

For $\mu = 0$, let

$$t = \int_{\frac{x_0+x_1}{2}}^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_{\frac{x_0+x_1}{2}}^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow c^+} \frac{\varphi(x)}{(x-c)^2} = \frac{a\gamma}{2(2d-1)(a\nu - b\lambda)} > 0, \quad \lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x_1 - x} = 1,$$

then

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = -\infty.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete.

(III – iv) For the case of $a < 0$ and $\lambda > 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $-\frac{\nu}{\lambda} < x < -\frac{b}{a}$, so $\frac{b}{a} - \frac{2\nu}{\lambda} < -\frac{\nu}{\lambda} \leq x_0 < x_1 \leq -\frac{b}{a}$.

By $\varphi(x_0) = \varphi'(x_0) = 0$ and (2.2), we obtain $\gamma = \frac{\mu(\nu + \lambda x_0)}{(b + ax_0)^2}$ and

$$\varphi(x) = \frac{(b+ax)^{2d+1}(a\lambda x + 2a\nu - b\lambda)}{(\nu + \lambda x)^d} \int_{x_0}^x \frac{(\nu + \lambda u)^d (\mu(\nu + \lambda u) - \gamma(b + au)^2)}{(b + au)^{2d+2}(b\lambda - 2a\nu - a\lambda u)^2} du$$

for $x \in (x_0, x_1)$.

When $x_0 < x < c := -\frac{b}{a}$, from

$$b\lambda - 2a\nu - a\lambda x > 0, \quad \frac{\nu + \lambda x_0}{(b + ax_0)^2} \geq 0, \quad \nu + \lambda x - \frac{\nu + \lambda x_0}{(b + ax_0)^2}(b + ax)^2 > 0,$$

it follows that

$$\mu(\nu + \lambda x) - \gamma(b + ax)^2 = \mu \left(\nu + \lambda x - \frac{\nu + \lambda x_0}{(b + ax_0)^2}(b + ax)^2 \right) \leq 0.$$

Hence $\varphi(x) < 0$ for $\mu < 0$ and $x \in (x_0, c)$, and $\varphi(x) = 0$ for $\mu = 0$ and $x \in (x_0, c)$. This is in contradiction with $\varphi(x) > 0$ for $x \in (x_0, x_1)$.

Part (IV)

Since $a \neq 0$, from (2.3), we get $\nu = 0$.

By Remark 2.1, when $\nu = 0$ and $r > 1$, there must be $\lambda > 0$ and $x = F'(t) > 0$.

By (2.3), using $\varphi(x_1) = 0$, we get

$$(4.22) \quad \varphi(x) = \frac{(ax+b)^{2n-1}(ax-b)}{x^{n-1}} \int_{x_1}^x \frac{u^{n-1}(\mu u - r(b+au)^2)}{(b+au)^{2n}(au-b)^2} du$$

for $x \in (x_0, x_1)$. Then

$$-1 = \varphi'(x_1) = \lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x - x_1} = \frac{\mu x_1 - r(ax_1 + b)^2}{(ax_1 + b)(ax_1 - b)}.$$

So

$$\mu = \frac{(ax_1 + b)(r(ax_1 + b) - (ax_1 - b))}{x_1}.$$

Since $\mu \leq 0$, $x_1 > 0$ and $ax_1 + b > 0$, we have $r \leq \frac{ax_1 - b}{ax_1 + b}$.

(IV - i) For the case of $a > 0$ and $\lambda > 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $x > \max\{-\frac{b}{a}, 0\}$, so $x_1 > x_0 \geq \max\{-\frac{b}{a}, 0\}$.

(IV - i - 1) If $b \leq 0$, namely $-\frac{b}{a} \geq 0 \geq \frac{b}{a}$.

By $1 < r \leq \frac{ax_1 - b}{ax_1 + b}$, we have $b < 0$ and $x_1 \leq -\frac{(r+1)b}{(r-1)a}$.

Let

$$q(x) = \mu x - r(b + ax)^2.$$

Note that when $c := -\frac{b}{a} < x < x_1$, by $\mu \leq 0$, $q(-\frac{b}{a}) \leq 0$ and $q(x_1) < 0$, we get

$$ax - b > 0 \text{ and } \mu x - r(b + ax)^2 \leq -r(b + ax)^2 \leq 0,$$

so $\varphi(x) > 0$ for $x \in (c, x_1)$ by (4.22).

Using

$$\lim_{x \rightarrow c^+} \varphi(x) = \frac{-\mu}{2(2n-1)a^2} \geq 0,$$

$$\lim_{x \rightarrow c^+} \frac{\varphi(x)}{(b+ax)^2} = \frac{-r}{2(2n-3)ab} > 0 \text{ (for } \mu = 0\text{)}$$

and fiber metrics for \tilde{g} are complete, we have $x_0 = c$.

For $\mu < 0$, let

$$t = \int_c^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_c^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow c^+} \varphi(x) = \frac{-\mu}{2(2d+1)a^2} > 0$$

and $\lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x_1 - x} = 1$, we have

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = 0.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 1 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete at $x = x_0$.

For $\mu = 0$, let

$$t = \int_{\frac{x_0+x_1}{2}}^{x(t)} \frac{du}{\varphi(u)}, \quad F(t) = \int_{\frac{x_0+x_1}{2}}^{x(t)} \frac{v dv}{\varphi(v)},$$

by

$$\lim_{x \rightarrow c^+} \frac{\varphi(x)}{(b+ax)^2} = \frac{-r}{2(2n-3)ab} > 0 \quad \text{and} \quad \lim_{x \rightarrow x_1^-} \frac{\varphi(x)}{x_1-x} = 1,$$

we get

$$\lim_{x \rightarrow x_1^-} t(x) = +\infty, \quad \lim_{x \rightarrow c^+} t(x) = -\infty.$$

This shows that the metric g_F is defined on $M = \{(z, w) \in \Omega \times \mathbb{C}^r : 0 < e^{\lambda\phi(z)}\|w\|^2\}$. Since

$$\int_{x_0}^{\frac{x_0+x_1}{2}} \frac{1}{(b+ax)\sqrt{\varphi(x)}} dx = +\infty,$$

it follows that any fiber metric for \tilde{g} is complete at $x = x_0$.

(IV – i – 2) If $b > 0$, then

$$\mu = \frac{(ax_1 + b)(r(ax_1 + b) - (ax_1 - b))}{x_1} > 0.$$

This conflicts with $\mu \leq 0$.

(IV – ii) For the case of $a < 0$ and $\lambda > 0$.

By $ax + b > 0$ and $\nu + \lambda x > 0$, we get $0 < x < -\frac{b}{a}$, so $b > 0$ and $0 \leq x_0 < x_1 \leq -\frac{b}{a}$.

Using $\varphi(x_0) = \varphi'(x_0) = 0$ and (2.3), we obtain $\mu = \frac{r(ax_0+b)^2}{x_0} > 0$. This conflicts with $\mu \leq 0$. \square

Acknowledgement. This work is partially supported by NSFC (No.12071354) and the Scientific Research Fund of Leshan Normal University (No.DGZZ202024)..

REFERENCES

- [1] V. Apostolov, D. M. J. Calderbank, P. Gauduchon: The geometry of weakly self-dual Kähler surfaces, *Compositio Math.* 135 (2003), 279-322.
- [2] V. Apostolov, D. M. J. Calderbank and P. Gauduchon: Hamiltonian 2-forms in Kähler geometry, I general theory. *J. differential geometry*, 73 (2006), 359-412.
- [3] V. Apostolov, D. M. J. Calderbank and P. Gauduchon: Ambitoric geometry I, Einstein metrics and extremal ambikähler structures, *J. Reine Angew. Math.*, 721 (2016), 109-147.
- [4] V. Apostolov, D. M. J. Calderbank and P. Gauduchon: Ambitoric geometry II, extremal toric surfaces and Einstein 4-orbifolds, *Ann. Sci. Éc. Norm. Supér. (4)*, 48 (2015), 1075-1112.
- [5] V. Apostolov, G. Maschler: conformally Kähler, Einstein-Maxwell geometry, *J. Eur. Math. Soc.* 21 (2019), 1319-1360.

- [6] V. Apostolov, G. Maschler, and C. W. Tønnesen-Friedman: Weighted extremal Kähler metrics and the Einstein-Maxwell geometry of projective bundles, arXiv:1808.02813.
- [7] L. Bérard-Bergery, Sur de nouvelles variétés riemanniennes d'Einstein, (French) [Some new Einstein Riemannian manifolds] Institut Élie Cartan, 6, 1-60, Inst. Élie Cartan, 6, Univ. Nancy, Nancy, 1982.
- [8] A. L. Besse, Einstein Manifolds, Ergebnisse (3) 10, Springer-Verlag, Berlin-Heidelberg-New York, 1987.
- [9] X.-X. Chen, C. LeBrun and B. Weber: On conformally Kähler, Einstein manifolds, J. Amer. Math. Soc., 21 (2008), 1137-1168.
- [10] A. Derdzinski: Hermitian Einstein metrics, Global Riemannian Geometry, Durham, 1983 (ed. T. J. Willmore and N. Hitchin, Ellis Horwood, Chichester, 1984), 105-114.
- [11] A. Derdzinski and G. Maschler: Local classification of conformally-Einstein Kähler metrics in Higher dimensions, Proc. London Math. Soc., 87(3) (2003), 779-819.
- [12] A. Derdzinski and G. Maschler: Special Kähler-Ricci potentials on compact Kähler manifolds, J. reine angew. Math, 593 (2006), 73-116.
- [13] A. Derdzinski and G. Maschler: A moduli curve for compact conformally-Einstein Kähler manifolds, Compositio Math. 141 (2005), 1029-1080.
- [14] Z. M. Feng: The first two coefficients of the Bergman function expansions for Cartan-Hartogs domains, International Journal of Mathematics, **29** (2018), 1850043 (45 pages).
- [15] Z. M. Feng: Rotationally symmetric conformal Kähler, Einstein-Maxwell metrics, New York J. Math. 26 (2020), 334-361.
- [16] J. Fu, S.T. Yau, W. Zhou: On complete constant scalar curvature Kähler metrics with Poincaré-Mok-Yau asymptotic property, Communications in Analysis and Geometry, **24**(3)(2016): 521-557.
- [17] A. Futaki, H. Ono: Volume minimization and conformally Kähler, Einstein-Maxwell geometry, J. Math. Soc. Japan., 70(4) (2018), 1493-1521.
- [18] A. Futaki, H. Ono: conformally Einstein-Maxwell Kähler metrics and structure of the automorphism group, Mathematische Zeitschrift, 292 (2019), 571-589.
- [19] P. Gao, S.T. Yau, W. Zhou: Nonexistence for complete Kähler-Einstein metrics on some noncompact manifolds, Math. Ann., 369 (2017), 1271-1282.
- [20] William B. Gordon: An analytical criterion for the completeness of Riemannian manifolds, Proceedings of the American Mathematical Society, 37(1), 1973.
- [21] William B. Gordon: Corrections to "An Analytical Criterion for the Completeness of Riemannian Manifolds", Proceedings of the American Mathematical Society, 45(1), 1974.
- [22] A. Hwang, M. Singer: A momentum construction for circle-invariant Kähler metrics, Transactions of the American Mathematical Society, 354 (2002), 2285-2325.
- [23] C. Koca and C. W. Tønnesen-Friedman: Strongly Hermitian Einstein-Maxwell solutions on ruled surfaces, Ann. Glob. Annl. Geom., 50 (2016), 29-46.
- [24] A. Lahdili: Automorphisms and deformations of conformally Kähler, Einstein-Maxwell metrics, J. Geom. Anal., 29 (2019), 542-568.
- [25] A. Lahdili: conformally Kähler, Einstein-Maxwell metrics and boundedness of the modified Mabuchi functional, Int. Math. Res. Notices (IMRN), rny239 (2018).
- [26] A. Lahdili: Kähler metrics with constant weighted scalar curvature and weighted K-stability. Proceedings of the London Mathematical Society, 119(4)(2019), 1065-1114.
- [27] C. LeBrun: The Einstein-Maxwell equations, Kähler metrics, and Hermitian geometry, J. Geom. Phys., 91 (2015), 163-171.
- [28] C. LeBrun: The Einstein-Maxwell equations and conformally Kähler geometry, Commun. Math. Phys., 344 (2016), 621-653.
- [29] D. Page: A compact rotating gravitational instanton, Phys. Lett. B, 79 (1978), 235-238.
- [30] Wang A., Yin W.P., Zhang L.Y., Roos G.: The Kähler-Einstein metric for some Hartogs domains over bounded symmetric domains. Science in China Series A: Mathematics **49**(9), 1175-1210 (2006).