UPPER HEAT KERNEL ESTIMATES FOR NONLOCAL OPERATORS VIA ARONSON'S METHOD

MORITZ KASSMANN AND MARVIN WEIDNER

ABSTRACT. In his celebrated article, Aronson established Gaussian bounds for the fundamental solution to the Cauchy problem governed by a second order divergence form operator with uniformly elliptic coefficients. We extend Aronson's proof of upper heat kernel estimates to nonlocal operators whose jumping kernel satisfies a pointwise upper bound and whose energy form is coercive. A detailed proof is given in the Euclidean space and extensions to doubling metric measure spaces are discussed.

1. INTRODUCTION

1.1. **Background.** Let $a_{i,j} : [0,T] \times \mathbb{R}^d \to (0,\infty)$, $i, j = 1, \ldots, d$, be bounded, measurable coefficients which satisfy the usual uniform ellipticity condition. A celebrated result by D.G. Aronson from 1967 says that the fundamental solution $\Gamma(y, s; x, \eta)$ to the Cauchy problem

$$\begin{cases} \partial_t u - \partial_i (a_{i,j} \partial_j u) = 0, & \text{in } (\eta, T) \times \mathbb{R}^d, \\ u(\eta) = u_0 \end{cases}$$
(1.1)

satisfies the following two-sided estimate for all $0 \le \eta < s < T$, and $x, y \in \mathbb{R}^d$:

$$c_1 t^{-\frac{d}{2}} e^{-c_2 \frac{|x-y|^2}{s-\eta}} \le \Gamma(y,s;x,\eta) \le c_3 t^{-\frac{d}{2}} e^{-c_4 \frac{|x-y|^2}{s-\eta}},$$
(1.2)

where $c_1, c_2, c_3, c_4 > 0$ depend only on d and the ellipticity constants. In other words, the fundamental solution of the classical heat equation $\partial_t u - \Delta u = 0$ is an upper and lower bound of Γ up to multiplicative constants, see [Aro67], [Aro68]. In this sense, the bounds (1.2) are robust in the class of second order divergence form operators with bounded, measurable, uniformly elliptic coefficients.

Aronson's proof is closely related to the so-called DeGiorgi-Nash-Moser theory for parabolic differential operators of second order with bounded, measurable and uniformly elliptic coefficients. The proof heavily relies on Hölder regularity estimates and the parabolic Harnack inequality for solutions to (1.1), see also [AS67].

[Aro68] has initiated several research studies on estimates for fundamental solutions to parabolic equations in various contexts. An important feature of this research is that it connects partial differential equations with geometry. This is due to the sensitivity of the heat kernel to the geometric properties of the underlying space. This phenomenon becomes apparent in the celebrated works [CLY81], [LY86], where the method of Aronson was generalized to prove heat kernel estimates on complete Riemannian manifolds with nonnegative Ricci curvature. Some

²⁰¹⁰ Mathematics Subject Classification. 35K08, 60J25, 31C25, 47D07, 39B62.

Key words and phrases. nonlocal operator, Dirichlet form, fundamental solution, heat kernel estimate.

Moritz Kassmann and Marvin Weidner gratefully acknowledge financial support by the German Research Foundation (SFB 1283 - 317210226 resp. GRK 2235 - 282638148).

of their arguments have been further refined and generalized in [Dav89], where an integral estimate for the heat kernel was established that is useful for proving the upper bound in (1.2). We refer the interested reader to [PEm84], [Gri09], [Li12] and the references therein for more detailed expositions on this topic.

1.2. Main results. The goal of this article is to extend Aronson's proof of upper heat kernel estimates to integro-differential operators of the form

$$L_t u(x) = \text{ p.v.} \int_{\mathbb{R}^d} (u(y) - u(x))k(t; x, y) \mathrm{d}y, \quad t \in (0, T), \ x \in \mathbb{R}^d.$$

Such operators are determined by a jumping kernel $k : (0,T) \times \mathbb{R}^d \times \mathbb{R}^d \to [0,\infty]$ which is assumed to be symmetric, i.e., k(t;x,y) = k(t;y,x) and satisfies a pointwise upper bound

$$k(t;x,y) \le \Lambda |x-y|^{-d-\alpha}, \quad t \in (0,T), \ x,y \in \mathbb{R}^d, \tag{k} \leq 0$$

for some given constant $\Lambda > 0$, and $\alpha \in (0, 2), 0 < T \leq \infty$. Moreover, we assume that there is $\lambda > 0$ such that for any ball $B \subset \mathbb{R}^d$ and every $v \in H^{\alpha/2}(B)$:

$$\int_{B} \int_{B} (v(x) - v(y))^{2} k(t; x, y) \mathrm{d}y \mathrm{d}x \ge \lambda [v]_{H^{\alpha/2}(B)}^{2}, \quad t \in (0, T).$$
 (\mathcal{E}_{\geq})

 (\mathcal{E}_{\geq}) can be thought of as a coercivity assumption on k and is substantially weaker than a pointwise lower bound. We refer the reader to Section 2 for a more detailed discussion and to Section 4 where we explain a possible extension of our method and replace (\mathcal{E}_{\geq}) by a Faber-Krahn inequality.

We are now ready to state the main result of this article in the aforementioned setup. For a possible extension to doubling metric measure spaces and jumping kernels of mixed type, we refer to Theorem 4.1.

Theorem 1.1. Let $k : (0,T) \times \mathbb{R}^d \times \mathbb{R}^d \to [0,\infty]$ be symmetric and assume (k_{\leq}) , (\mathcal{E}_{\geq}) . Let $p(y,s;x,\eta)$ be the fundamental solution to the Cauchy problem

$$\begin{cases} \partial_t u - L_t u = 0, & in(\eta, T) \times \mathbb{R}^d, \\ u(\eta) = u_0 \in L^2(\mathbb{R}^d), \end{cases}$$
(1.3)

where $\eta \in [0,T)$. Then there exists a constant c > 0 depending on $d, \alpha, \lambda, \Lambda$ such that for every $0 \le \eta < s < T$, and $x, y \in \mathbb{R}^d$:

$$p(y,s;x,\eta) \le c(s-\eta)^{-\frac{d}{\alpha}} \left(1 + \frac{|x-y|^{\alpha}}{s-\eta}\right)^{-\frac{d+\alpha}{\alpha}}.$$
(1.4)

Estimate (1.4) states that the fundamental solution to (1.3) possesses the same upper bound as the fundamental solution to the fractional heat equation $\partial_t u + (-\Delta)^{\alpha/2} u = 0$, see [BG60].

Different versions of Theorem 1.1 already exist in the literature. Let us give a brief account on the history of heat kernel bounds for nonlocal operators. Two-sided estimates of the form (1.4) have been established by [BL02], [CK03] using a probabilistic approach. They assume that the jumping kernel is pointwise comparable to $|x - y|^{-d-\alpha}$, $\alpha \in (0, 2)$, from above and below. Their analysis of the upper heat kernel estimate heavily relies on [CKS87], where Davies' method was extended to a more general setup, including jump processes. Building upon this, [BGK09] derived (1.4) assuming (k_{\leq}) and a Nash inequality.

In a series of articles, see [GHL14], [GHH17], [GHH18], the analysis of heat kernel estimates was extended to metric measure spaces with walk dimension greater than 2. The authors were able to characterize upper heat kernel estimates, as well as two-sided heat kernel estimates in terms of equivalent conditions on the jumping kernels and the geometry of the underlying space. Their approach does not use the underlying stochastic process and is based on certain comparison inequalities of the corresponding heat semigroups relying on the parabolic maximum principle. Note that Davies' method was extended to jumping kernels with jump index $\alpha > 2$ in [MSC19], [HL18]. The aforementioned results assume certain homogeneity of the doubling measure space and do not deal with mixed-type jumping kernels.

In [CK08], [CKKW21], [CKW21] upper and two-sided heat kernel estimates were investigated on doubling metric measures spaces for jumping kernels of mixed type. We would like to draw the reader's attention to Theorem 1.15 in [CKW21]. In the case $\alpha < 2$, it states that upper heat kernel estimates of the form (1.4) are equivalent to a pointwise upper bound on the jumping kernel and a Faber-Krahn inequality, which can be understood as an implicit lower bound. A major difference between our approach and [CKW21] is that our method relies on purely analytic arguments, while [CKW21] makes essential use of the corresponding stochastic process. In Theorem 4.1 we extend our approach to doubling metric measure spaces and jumping kernels of mixed type. Let us mention that we prove on-diagonal heat kernel estimates with the help of a parabolic $L^{\infty} - L^1$ -estimate, see Lemma 4.2. This rather straightforward approach allows us to avoid truncation methods and the usage of the iteration techniques of [Kig04].

In contrast to our setup, all jumping kernels in the results discussed above are assumed to be time-homogeneous. Note that it would require substantial effort to extend methods based on stochastic processes to situations with time-dependent jumping kernels. Heat kernel estimates for time-inhomogeneous jumping kernels were established in [MM13a], [MM13b] where the authors assume pointwise upper and lower bounds on the jumping kernel. Note that assuming pointwise lower bounds is more restrictive than (\mathcal{E}_{\geq}) . The focus of these works lies on the treatment of an additional divergence-free drift of first order.

There are further results on heat kernel estimates for nonlocal operators, which are related to Theorem 1.1. For example, sharp two-sided estimates for jump processes on \mathbb{R}^d with upper scaling index not strictly less than 2 are established in [BKKL19]. In [KKK], heat kernel estimates for a certain class of jump processes with singular jumping measures are proved.

1.3. Strategy of proof. A main insight of Aronson's proof for second order differential operators is the observation that solutions u to the Cauchy problem (1.1) satisfy the weighted L^2 -estimate

$$\sup_{t \in (\eta,s)} \int_{\mathbb{R}^d} H(t,x) u^2(t,x) \mathrm{d}x \le \int_{\mathbb{R}^d} H(\eta,x) u_0^2(x) \mathrm{d}x \tag{1.5}$$

for $0 \leq \eta < s < T$, whenever H satisfies

$$C|\nabla H^{1/2}|^2 \le -\partial_t H, \quad \text{in } (\eta, s) \times \mathbb{R}^d,$$
(1.6)

for a given number C > 0 depending on the ellipticity constants. (1.6) is closely related to the famous Li-Yau inequality:

$$|\nabla \log w|^2 \le \frac{d}{2t} + \partial_t \log w. \tag{1.7}$$

In fact, a direct computation reveals that the Gauss-Weierstrass kernel $w(t,x) = t^{-\frac{d}{2}}e^{-\frac{|x|^2}{4t}}$ satisfies (1.7) with equality. By a scaling argument, it becomes evident that (1.6) holds true for $H(t,x) = (C[t])^{\frac{d}{2}}w(C[t],x-y) = \exp(-\frac{|x-y|^2}{4C[t]})$, where $[t] := 2(s-\eta) - (t-\eta)$ and $y \in \mathbb{R}^d$ can be chosen arbitrarily.

This insight suggests that some qualitative information on the decay of solutions to (1.1) is

encoded in the weighted L^2 -estimate (1.5). Indeed, by combining (1.5) with a localized $L^{\infty} - L^2$ estimate, as it was proved by Moser ([Mos64], [Mos67], [Mos71]), one can deduce a qualitative estimate in terms of H, on how quick solutions to the Cauchy problem increase inside a ball that lies outside the support of the initial data. From such estimate, it is not difficult to deduce

$$\left(\int_{\mathbb{R}^d \setminus B_{\sigma}(y)} \Gamma^2(y, s; z, \eta) \mathrm{d}z\right)^{1/2} \le c(s - \eta)^{-\frac{d}{4}} e^{-\frac{\sigma^2}{32C(s - \eta)}}$$

for every $\sigma > 0$ and $0 \le \eta < s < T$ with $s - \eta \le \sigma^2$. Together with the on-diagonal estimate $\Gamma(y, s; x, \eta) \le c(s - \eta)^{-\frac{d}{2}}$, one deduces the upper bound in (1.2) via a standard argument.

We would like to point out that (1.7) is at the core of the works [LY86], [CLY81], where (1.7) was used to derive a parabolic Harnack inequality on Riemannian manifolds with nonnegative Ricci-curvature and to establish its equivalence to Gaussian heat kernel bounds.

Next, let us summarize how we adapt Aronson's proof to integro-differential operators. First, we require a nonlocal analog of (1.5). We prove that there exist functions H that satisfy

$$C\Gamma^{\alpha}_{\rho}(H^{1/2}, H^{1/2}) \le -\partial_t H, \quad \text{in } (\eta, s) \times \mathbb{R}^d$$
(1.8)

given $C, \rho > 0$. Here, Γ_{ρ}^{α} denotes the ρ -truncated carré du champ operator of order $\alpha \in (0, 2)$, which is defined as follows:

$$\Gamma^{\alpha}_{\rho}(f,f)(x) = \int_{B_{\rho}(x)} (f(x) - f(y))^2 |x - y|^{-d-\alpha} \mathrm{d}y, \quad x \in \mathbb{R}^d.$$

By a careful choice of a function H satisfying (1.8), we to deduce a nonlocal analog of (1.5) for solutions to the ρ -truncated Cauchy problem, see Lemma 3.1. The corresponding integrodifferential operator only takes into account differences up to distance ρ . In order to prove an a priori bound for the corresponding fundamental solution p_{ρ} (see Theorem 3.4), we derive a parabolic $L^{\infty} - L^2$ -estimate in the spirit of [Str19], see Lemma 2.4. The estimate involves a nonlocal, truncated tail-term which requires special treatment. In a final step, we obtain the desired upper heat kernel estimate (1.4) for p by gluing together short and long jumps. Such argument is by now standard in the theory of jump processes.

Last, we explain several difficulties that occur when avoiding the detour via the truncated jumping kernel.

First, the corresponding $L^{\infty} - L^2$ -estimate involves a non-truncated tail-term which cannot be controlled without any further assumptions on k.

Second, finding suitable weight functions H which satisfy a non-truncated version of (1.8) is a challenging task in the light of the following observation: The corresponding nonlocal analog of the Li-Yau inequality, which would imply an estimate of the form (1.5) for solutions to (1.3), reads as follows:

$$\frac{\Gamma^{\alpha}(w_{\alpha}^{1/2}, w_{\alpha}^{1/2})}{w_{\alpha}} \le \frac{d}{\alpha t} + \partial_t \log(w_{\alpha}).$$
(1.9)

However, one can show that the fundamental solution $w_{\alpha}(t,x)$ to $\partial_t u + (-\Delta)^{\alpha/2} u = 0$ does not satisfy (1.9). Let us give a quick proof of this fact. First, note that w_{α} is a radial function and satisfies

$$\frac{d}{\alpha t}w_{\alpha} + \partial_t w_{\alpha} = -\frac{|x|}{\alpha t}\partial_{|x|}w_{\alpha}$$

For a proof of this identity, we refer to (2.5) in [Vaz18]. Consequently,

$$\frac{d}{\alpha t} + \partial_t \log w_\alpha(t,0) = 0,$$

but this is a contradiction to (1.9) since $\Gamma^{\alpha}(w_{\alpha}^{1/2}, w_{\alpha}^{1/2})(t, 0) > 0$ for t > 0.

1.4. **Outline.** This article is separated into five sections. In Section 2 we present several auxiliary results that we need in our proof. Section 3 contains the derivation of the upper heat kernel bounds and proves Theorem 1.1. In Section 4, we explain how our method can be applied to jumping kernels of mixed type on metric measure spaces. In Section 5, we provide a proof of a gluing lemma which differs from [GH08] due to the time-inhomogeneity of the jumping kernels under consideration.

2. Preliminaries

In this section we provide several auxiliary results that will be required for the proof of Theorem 1.1 in Section 3.

Let $k: (0,T) \times \mathbb{R}^d \times \mathbb{R}^d \to [0,\infty]$ be a symmetric jumping kernel satisfying the pointwise upper bound (k_{\leq}) and the coercivity condition (\mathcal{E}_{\geq}) .

Let us make a few comments on assumption (\mathcal{E}_{\geq}) : First of all, (\mathcal{E}_{\geq}) can be regarded as a nonlocal substitute of the classical uniform ellipticity condition for local operators. In fact, it is considerably weaker than a pointwise lower bound on the jumping kernel, since (\mathcal{E}_{\geq}) allows for jumping kernels that might degenerate in certain directions, as for example kernels that are supported on double cones. We refer the interested reader to [CS20] for an investigation of such condition.

A coercivity assumption like (\mathcal{E}_{\geq}) is crucial to our approach since it is needed for the $L^{\infty} - L^2$ -estimate (Lemma 2.4) and also the on-diagonal heat kernel bound (Theorem 2.3). In the literature, lower bounds on jumping kernels are often introduced through functional inequalities, e.g. in [BGK09], or [CKW21] where the authors assume a Nash -, or a Faber-Krahn inequality. We point out that such assumption would have been possible also in this work, since the proofs of Lemma 2.4, Theorem 2.3 can be changed accordingly (see Section 4). For a discussion on the equivalence of Nash - and Faber-Krahn inequalities, we refer the reader to [CKW21].

For any $\rho > 0$, we define the truncated jumping kernel k_{ρ} via

$$k_{\rho}(t; x, y) = k(t; x, y) \mathbb{1}_{\{|x-y| < \rho\}}(x, y).$$

The associated integro-differential operator L_t^{ρ} is defined as

$$L_t^{\rho}u(x) = \text{ p.v.} \int_{\mathbb{R}^d} (u(y) - u(x))k_{\rho}(t; x, y) \mathrm{d}y.$$

Definition 2.1. We say that a function $u \in L^2_{loc}((\eta, T); H^{\alpha/2}(\mathbb{R}^d))$ with $\partial_t u \in L^1_{loc}((\eta, T); L^2_{loc}(\mathbb{R}^d))$ solves the Cauchy problem associated with k in $(\eta, T) \times \mathbb{R}^d$:

$$\begin{cases} \partial_t u - L_t u = 0, & in (\eta, T) \times \mathbb{R}^d, \\ u(\eta) = u_0 \in L^2(\mathbb{R}^d), \end{cases}$$
(2.1)

if for every $\phi \in H^{\alpha/2}(\mathbb{R}^d)$ with $\operatorname{supp}(\phi)$ compact, it holds

$$\int_{\mathbb{R}^d} \partial_t u(t, x) \phi(x) \mathrm{d}x + \mathcal{E}_t(u(t), \phi) = 0, \quad a.e. \ t \in (\eta, T),$$
(2.2)

$$\|u(t) - u_0\|_{L^2(\mathbb{R}^d)} \to 0, \quad \text{as } t \searrow \eta,$$
(2.3)

where we write

$$\mathcal{E}_t(u,v) = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (u(t,x) - u(t,y))(v(t,x) - v(t,y))k(t;x,y) \mathrm{d}y \mathrm{d}x$$
(2.4)

for the family of energy forms $(\mathcal{E}_t)_{t \in (\eta,T)}$ associated with k.

Solutions to the ρ -truncated Cauchy problem associated with k in $(\eta, T) \times \mathbb{R}^d$

$$\begin{cases} \partial_t u - L_t^{\rho} u = 0, & \text{in } (\eta, T) \times \mathbb{R}^d, \\ u(\eta) &= u_0 \in L^2(\mathbb{R}^d), \end{cases}$$
(2.5)

are defined in an analogous way, replacing k by k_{ρ} .

Throughout this article, we will assume that the fundamental solutions to the Cauchy problem (2.1) and the ρ -truncated Cauchy problem (2.5) exist, i.e., that the solution u to (2.1) and u_{ρ} to (2.5) are unique, and have the representation

$$u(s,y) = \int_{\mathbb{R}^d} p(y,s;x,\eta) u_0(x) \mathrm{d}x, \quad u_\rho(s,y) = \int_{\mathbb{R}^d} p_\rho(y,s;x,\eta) u_0(x) \mathrm{d}x, \quad s \in (\eta,T), \ y \in \mathbb{R}^d,$$
(2.6)

where $p, p_{\rho} : (0, T) \times \mathbb{R}^d \times [0, T) \times \mathbb{R}^d \to [0, \infty]$ satisfy the following properties for all $0 \le \eta < t < s < T, x, y \in \mathbb{R}^d$:

$$p(y,s;x,\eta) = p(x,s;y,\eta) > 0, \quad p_{\rho}(y,s;x,\eta) = p_{\rho}(x,s;y,\eta) > 0, \tag{2.7}$$

$$\int_{\mathbb{R}^d} p(y,s;x,\eta) \mathrm{d}x = 1, \quad \int_{\mathbb{R}^d} p_\rho(y,s;x,\eta) \mathrm{d}x = 1, \tag{2.8}$$

$$p(y,s;x,\eta) = \int_{\mathbb{R}^d} p(y,s;z,t) p(z,t;x,\eta) dz, \quad p_{\rho}(y,s;x,\eta) = \int_{\mathbb{R}^d} p_{\rho}(y,s;z,t) p_{\rho}(z,t;x,\eta) dz.$$
(2.9)

In the following, we will denote the unique solutions to (2.1) and (2.5) by $P_{\eta,s}u_0$, and $P_{\eta,s}^{\rho}u_0$. $(P_{\eta,s})_{s\in[\eta,T)}$, and $(P_{\eta,s}^{\rho})_{s\in[\eta,T)}$ are called the heat semigroups associated with k, and k_{ρ} .

In the time-homogeneous case, i.e., when k does not depend on t, the existence of $(P_{\eta,s})_{s\in[\eta,T)}$ and $(P_{\eta,s}^{\rho})_{s\in[\eta,T)}$ is guaranteed by symmetric Dirichlet form theory. The existence of the fundamental solution classically follows from so-called ultracontractivity estimates for the heat semigroup which are a consequence of Nash's inequality. For time-inhomogeneous jumping kernels k which satisfy the following pointwise lower bound for some $\lambda > 0$

$$k(t;x,y) \ge \lambda |x-y|^{-d-\alpha}, \ t \in (0,T), \ x,y \in \mathbb{R}^d,$$
 (2.10)

the existence of the fundamental solutions p and p_{ρ} was proved in [MM13a] by approximation of k through a sequence of smooth jumping kernels for which the desired properties follow from the theory of pseudo-differential operators. A similar result is proved in [Kom88], [Kom95] but under an additional smoothness assumption on t.

The following result explains the connection between p and p_{ρ} and is crucial to our approach. It is frequently used in the derivation of upper heat kernel bounds for alpha-stable like processes and goes back to a probabilistic construction carried out in [Mey75]. An analytic proof via the parabolic maximum principle is derived in [GH08] (see also [GHL14]). Since both proofs are known only in the time-homogeneous case, we will provide a modified version of the argument in [GH08] in the appendix.

Lemma 2.2. Assume that k satisfies (k_{\leq}) , (\mathcal{E}_{\geq}) . Then there exists c > 0 such that for every $\rho > 0, 0 \leq \eta < s < T$, and $x, y \in \mathbb{R}^d$ it holds

$$p(y,s;x,\eta) \le p_{\rho}(y,s;x,\eta) + c(s-\eta)\rho^{-d-\alpha},$$
(2.11)

$$p_{\rho}(y,s;x,\eta) \le e^{c\rho^{-\alpha}(s-\eta)}p(y,s;x,\eta).$$

$$(2.12)$$

Next, we provide the so-called on-diagonal bound for the heat kernels p, and p_{ρ} .

Theorem 2.3 (on-diagonal bound). Assume that k satisfies (k_{\leq}) , (\mathcal{E}_{\geq}) . Then there exists c > 0 depending on $d, \alpha, \lambda, \Lambda$ such that for every $\rho > 0$, $0 \leq \eta < s < T$, and $x, y \in \mathbb{R}^d$ it holds

$$p(y,s;x,\eta) \le c(s-\eta)^{-\frac{d}{\alpha}},\tag{2.13}$$

$$p_{\rho}(y,s;x,\eta) \le c e^{c\rho^{-\alpha}(s-\eta)}(s-\eta)^{-\frac{a}{\alpha}}.$$
(2.14)

There are at least two ways to prove Theorem 2.3. One approach classically goes via Nash inequalities (see [CKW21]) and can be traced back to Nash's famous work [Nas58]. This proof also works in the time-inhomogeneous setup (see [MM13a]).

Another way to establish on-diagonal bounds goes via $L^{\infty} - L^1$ -estimates. For this, we refer to Lemma 4.2, where such estimate is proved in a more general setup. Observe that $(t, z) \mapsto$ $p(y, t; z, \eta)$ solves $\partial_t u - Lu = 0$ in $(\eta, T) \times \mathbb{R}^d$ for every $y \in \mathbb{R}^d$. Therefore, by the $L^{\infty} - L^1$ estimate (4.12), for every $0 \le \eta < s < T$ and $x, y \in \mathbb{R}^d$:

$$p(y,s;x,\eta) \le c(s-\eta)^{-\frac{d}{\alpha}} \sup_{t \in (\eta,s)} \int_{\mathbb{R}^d} p(y,t;z,\eta) \mathrm{d}z \le c(s-\eta)^{-\frac{d}{\alpha}},$$

where we used (2.8) in the last step. (2.14) follows from Lemma 2.2.

The remainder of this section is devoted to proving an $L^{\infty} - L^2$ -estimate for solutions to $\partial_t u - L_t^{\rho} u = 0$ in a time-space cylinder $I_R(t_0) \times B_R(x_0)$, where $t_0 \in (0,T)$, $x_0 \in \mathbb{R}^d$, and $I_R(t_0) := (t_0 - R^{\alpha}, t_0) \subset (\eta, T)$.

Since we consider only truncated jumping kernels, it is possible to estimate the tail-term by an L^2 -norm over a ball with radius ρ , without any pointwise lower bounds of k or a UJS-type condition, see [BBK09]. This is at the cost of a suboptimal scaling constant in the resulting estimate, which luckily does not affect the proof of the final heat kernel estimate.

A similar result for solutions to elliptic equations was obtained in [CKW21]. We follow the strategy outlined in [Str19] which is based on a nonlocal adaptation of De Giorgi's iteration technique (see [DCKP14],[DCKP16]) but present the proof in all details due to our special treatment of the tail term.

Lemma 2.4 (truncated $L^{\infty} - L^2$ -estimate). Assume that k satisfies (k_{\leq}) , (\mathcal{E}_{\geq}) . There exists a constant C > 0 depending on $d, \alpha, \lambda, \Lambda$ such that for every $t_0 \in (0, T)$, $x_0 \in \mathbb{R}^d$, and $\rho, R > 0$ with $R \leq \rho/2 \wedge t_0^{1/\alpha}$, and every subsolution u to $\partial_t u - L_t^{\rho} u = 0$ in $I_R(t_0) \times B_R(x_0)$ it holds:

$$\sup_{I_{R/2}(t_0) \times B_{R/2}(x_0)} u \le C \left(\frac{\rho^{\alpha}}{R^{\alpha}}\right)^{\frac{d}{2\alpha}} R^{-\frac{d}{2}} \sup_{t \in I_R(t_0)} \left(\int_{B_{2\rho}(x_0)} u^2(t, x) \mathrm{d}x\right)^{1/2}.$$
 (2.15)

For the definition of a subsolution to $\partial_t u - L_t^{\rho} u = 0$ in $I_R(t_0) \times B_R(x_0)$, we refer to the appendix.

Proof. To keep notation simple, from now on we will write $I_R := I_R(t_0)$, and $B_R(x_0) := B_R$. From the usual Caccioppoli inequality, see [Str19], we know that for every r, R with $0 < r \le R \le \rho/2$ and every l > 0, it holds

$$\begin{split} \sup_{t \in I_R} \int_{B_R} w_l^2(t, x) \mathrm{d}x + \int_{I_R} \int_{B_R} \int_{B_R} (w_l(s, x) - w_l(s, y))^2 k_\rho(s; x, y) \mathrm{d}y \mathrm{d}x \mathrm{d}s \\ & \leq c_1 \left(\sigma(R, r) \int_{I_{R+r}} \int_{B_{R+r}} w_l^2(t, x) \mathrm{d}x \mathrm{d}t + \|w_l\|_{L^1(I_{R+r} \times B_{R+r})} \sup_{t \in I_{R+r}} \sup_{x \in B_{R+\frac{r}{2}}} \int_{B_{R+r}^c} w_l(t, y) k_\rho(t; x, y) \mathrm{d}y \right) \end{split}$$

where $c_1 > 0$ is a constant, $w_l = (u - l)_+$, and $\sigma(r, R) = r^{-(\alpha \vee 1)}(R + r)^{(\alpha \vee 1) - \alpha}$. Define $\kappa = 1 + \frac{\alpha}{d}$, and $A(l, R) := |\{(t, x) \in I_R \times B_R : u(t, x) > l\}|$. Then, by (\mathcal{E}_{\geq}) and the fractional Sobolev inequality:

$$\begin{split} \int_{I_R} \int_{B_R} w_l^2(t,x) \mathrm{d}x \mathrm{d}t &\leq c_2 |A(l,R)|^{\frac{1}{\kappa'}} \left(\int_{I_R} \int_{B_R} w_l^{2\kappa}(s,x) \mathrm{d}x \mathrm{d}s \right)^{\frac{1}{\kappa}} \\ &\leq c_3 |A(l,R)|^{\frac{1}{\kappa'}} \left(\left(\sup_{t \in I_R} \int_{B_R} w_l^2(t,x) \mathrm{d}x \right)^{\kappa-1} \int_{I_R} \left(\int_{B_R} w_l^{\frac{2d}{d-\alpha}}(s,x) \mathrm{d}x \right)^{\frac{d-\alpha}{d}} \mathrm{d}s \right)^{\frac{1}{\kappa}} \\ &\leq c_4 |A(l,R)|^{\frac{1}{\kappa'}} \left(\sigma(R,r) \int_{I_{R+r}} \int_{B_{R+r}} w_l^2(t,x) \mathrm{d}x \mathrm{d}t \right. \\ &\quad + \|w_l\|_{L^1(I_{R+r} \times B_{R+r})} \sup_{t \in I_{R+r}} \sup_{x \in B_{R+\frac{r}{2}}} \int_{B_{R+r}} w_l(t,y) k_{\rho}(t;x,y) \mathrm{d}y \right) \end{split}$$

for some $c_2, c_3, c_4 > 0$. Observe that for some constant $c_5 > 0$ by (k_{\leq})

$$\sup_{t\in I_{R+r}}\sup_{x\in B_{R+\frac{r}{2}}}\int_{B_{R+r}^c}w_l(t,y)k_\rho(t;x,y)\mathrm{d}y \le c_5r^{-\alpha}\left(\frac{\rho^{\alpha}}{r^{\alpha}}\right)^{\frac{d}{\alpha}}\sup_{t\in I_{R+r}}f_{B_{2\rho}}|u(t,y)|\mathrm{d}y$$

Let us now fix $R \in (0, \rho/2]$ and define sequences $l_i = M(1-2^{-i})$, for M > 0 to be defined later, $r_i = 2^{-i-1}R$, $R_{i+1} = R_i - r_{i+1}$, $R_0 := R$, $A_i = \int_{I_{R_i}} \int_{B_{R_i}} w_{l_i}^2(t, x) dx dt$. Note that by definition: $R/2 = \lim_{i \to \infty} R_i < \cdots < R_2 < R_1 < R_0 = R$, and $l_i \nearrow M$, and $\sigma(r_i, R_i) \le c_6 R^{-\alpha} 2^{2i}$. Then, we deduce from the two lines above:

$$\begin{split} A_{i} &\leq c_{7} \frac{1}{(l_{i} - l_{i-1})^{\frac{2}{\kappa'}}} \left(\sigma(R_{i}, r_{i}) + \frac{r_{i}^{-\alpha} \left(\frac{\rho^{\alpha}}{r_{i}^{\alpha}}\right)^{\frac{d}{\alpha}} \sup_{t \in I_{R_{i-1}}} f_{B_{2\rho}} |u(t, x)| \mathrm{d}x}{l_{i} - l_{i-1}} \right) A_{i-1}^{1 + \frac{1}{\kappa'}} \\ &\leq c_{8} \frac{2^{\frac{2i}{\kappa'}}}{M^{\frac{2}{\kappa'}}} \left(\frac{2^{2i}}{R^{\alpha}} + \frac{2^{i(d+\alpha+1)}\rho^{d}}{MR^{d+\alpha}} \sup_{t \in I_{R}} f_{B_{2\rho}} |u(t, x)| \mathrm{d}x} \right) A_{i-1}^{1 + \frac{1}{\kappa'}} \\ &\leq c_{9} \frac{2^{\gamma i}}{M^{\frac{2}{\kappa'}}R^{\alpha}} \left(1 + \frac{\left(\frac{\rho^{\alpha}}{R^{\alpha}}\right)^{\frac{d}{\alpha}} \sup_{t \in I_{R}} f_{B_{2\rho}} |u(t, x)| \mathrm{d}x}{M} \right) A_{i-1}^{1 + \frac{1}{\kappa'}} \end{split}$$

where $c_7, c_8, c_9 > 0$, and $\gamma = 2 + \frac{2}{\kappa'} + d + \alpha > 0$. Let us now choose

$$M := \left(\frac{\rho^{\alpha}}{R^{\alpha}}\right)^{\frac{d}{\alpha}} \sup_{t \in I_R} \oint_{B_{2\rho}} |u(t,x)| \mathrm{d}x + C^{\frac{\kappa'^2}{2}} c_{10}^{\frac{\kappa'}{2}} R^{-\frac{\alpha\kappa'}{2}} A_0^{1/2},$$

where $c_{10} = 2c_9$, $C = 2^{\gamma} > 1$. Consequently, it holds

$$A_{i} \leq \frac{c_{10}}{M^{\frac{2}{\kappa'}}R^{\alpha}} C^{i} A_{i-1}^{1+\frac{1}{\kappa'}}, \quad A_{0} \leq C^{-\kappa'^{2}} \left(\frac{c_{10}}{R^{\alpha}M^{\frac{2}{\kappa'}}}\right)^{-\kappa'}.$$

By Lemma 7.1 in [Giu03], we obtain

$$\sup_{I_{R/2} \times B_{R/2}} u \leq M = \left(\frac{\rho^{\alpha}}{R^{\alpha}}\right)^{\frac{d}{\alpha}} \sup_{t \in I_R} f_{B_{2\rho}} |u(t,x)| dx + C^{\frac{\kappa'^2}{2}} c_{10}^{\frac{\kappa'}{2}} R^{-\frac{\alpha\kappa'}{2}} A_0^{1/2}$$
$$\leq c_{11} \left(\frac{\rho^{\alpha}}{R^{\alpha}}\right)^{\frac{d}{\alpha}} \sup_{t \in I_R} f_{B_{2\rho}} |u(t,x)| dx + c_{11} \left(f_{I_R} f_{B_R} u^2(t,x) dx dt\right)^{1/2}$$
$$\leq c_{12} \left(\frac{\rho^{\alpha}}{R^{\alpha}}\right)^{\frac{d}{2\alpha}} R^{-\frac{d}{2}} \sup_{t \in I_R} \left(\int_{B_{2\rho}} u^2(t,x) dx\right)^{1/2},$$

for some $c_{11}, c_{12} > 0$, as desired, where we used $R \leq \rho/2$.

3. Nonlocal Aronson Method

In this section we prove Theorem 1.1. As is standard for proofs of heat kernel bounds for nonlocal operators, we first establish bounds for the heat kernel corresponding to the truncated jumping kernel and derive the estimate for the original jumping kernel by gluing together short and long jumps with the help of Lemma 2.2 in a second step. The following lemma is a nonlocal version of (1.5).

The following lemma is a nonlocal version of (1.3).

Lemma 3.1. Assume that k satisfies (k_{\leq}) and let $\rho > 0$, $0 \leq \eta < s < T$. Let $u \in L^{\infty}((\eta, T) \times \mathbb{R}^d)$ be a solution to the ρ -truncated Cauchy problem (2.5) in $(\eta, T) \times \mathbb{R}^d$. Then there exists a constant $C = C(\Lambda)$ such that for every bounded function $H : [\eta, s] \times \mathbb{R}^d \to [0, \infty)$ satisfying

•
$$C\Gamma^{\alpha}_{\rho}(H^{1/2}, H^{1/2}) \leq -\partial_t H \text{ in } (\eta, s) \times \mathbb{R}^d$$

•
$$H^{1/2} \in L^2((\eta, s); H^{\alpha/2}(\mathbb{R}^d)),$$

the following estimate holds true:

$$\sup_{t \in (\eta,s)} \int_{\mathbb{R}^d} H(t,x) u^2(t,x) \mathrm{d}x \le \int_{\mathbb{R}^d} H(\eta,x) u_0^2(x) \mathrm{d}x.$$
(3.1)

Proof. Let $R \geq 2$ and $\gamma_R \in C_c^{\infty}(\mathbb{R}^d)$ such that $\gamma_R \equiv 1$ in $B_{R-1}(0)$, $\gamma_R \equiv 0$ in $\mathbb{R}^d \setminus B_R(0)$, $0 \leq \gamma_R \leq 1$, $|\nabla \gamma_R| \leq 2$. Consequently, $\Gamma_{\rho}^{\alpha}(\gamma_R, \gamma_R)$ satisfies:

$$\Gamma^{\alpha}_{\rho}(\gamma_R,\gamma_R)(x) \leq \begin{cases} 0, & x \in B_{\frac{R-1}{2}}(0), \\ c_1, & x \in \mathbb{R}^d \setminus B_{\frac{R-1}{2}}(0) \end{cases}$$
(3.2)

for some constant $c_1 > 0$. We test the equation for u with the test function $\phi = \gamma_R^2 H u$ and integrate in time over (η, τ) , where $\tau \in (\eta, s)$, and obtain:

$$\int_{\eta}^{\tau} \int_{\mathbb{R}^d} (\partial_t u) \phi \mathrm{d}x \mathrm{d}t + \int_{\eta}^{\tau} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (u(t,x) - u(t,z)) (\phi(t,x) - \phi(t,z)) k_{\rho}(t;x,z) \mathrm{d}z \mathrm{d}x \mathrm{d}t = 0.$$

Note that ϕ is a valid test function since for every $t \in (\eta, s)$ by assumption it holds $\gamma_R H(t)u(t) \in H^{\alpha/2}(\mathbb{R}^d)$. From $(\partial_t u)u = \frac{1}{2}\partial_t(u^2)$ and integration by parts:

$$\begin{split} \int_{\mathbb{R}^d} u^2(\tau, x) \gamma_R^2(x) H(\tau, x) \mathrm{d}x + 2 \int_{\eta}^{\tau} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (u(t, x) - u(t, z)) (\phi(t, x) - \phi(t, z)) k_{\rho}(t; x, z) \mathrm{d}z \mathrm{d}x \mathrm{d}t \\ &= \int_{\mathbb{R}^d} u_0^2(x) \gamma_R^2(x) H(\eta, x) \mathrm{d}x + \int_{\eta}^{\tau} \int_{\mathbb{R}^d} u^2(t, x) \gamma_R^2(x) \partial_t H(t, x) \mathrm{d}x \mathrm{d}t. \end{split}$$

We treat the nonlocal term by making use of the following algebraic inequality:

$$(u_1 - u_2)(\gamma_1^2 H_1 u_1 - \gamma_2^2 H_2 u_2) \ge (\gamma_1 H_1^{1/2} u_1 - \gamma_2 H_2^{1/2} u_2)^2 - c_2 \left((\gamma_1 - \gamma_2)^2 (H_1 + H_2) (u_1^2 + u_2^2) + (H_1^{1/2} - H_2^{1/2})^2 (\gamma_1^2 + \gamma_2^2) (u_1^2 + u_2^2) \right),$$

where $c_2 > 0$. Its proof is based on the following two observations:

$$(u_1 - u_2)(\gamma_1^2 H_1 u_1 - \gamma_2^2 H_2 u_2) = (\gamma_1 H_1^{1/2} u_1 - \gamma_2 H_2^{1/2} u_2)^2 - u_1 u_2 (\gamma_1 H_1^{1/2} - \gamma_2 H_2^{1/2})^2,$$

$$u_1 u_2 (\gamma_1 H_1^{1/2} - \gamma_2 H_2^{1/2})^2 \le c_2 (u_1^2 + u_2^2) \left((\gamma_1 - \gamma_2)^2 (H_1 + H_2) + (\gamma_1^2 + \gamma_2^2) (H_1^{1/2} - H_2^{1/2})^2 \right).$$

Moreover, note that for $R > 4\rho$ it holds $\gamma_R^2(z) = \gamma_R^2(x) = 1$ for every $x \in B_{\frac{R-1}{2}}(0)$ and $z \in B_\rho(x)$. By symmetry of k, (k_{\leq}) and the observations from above, we deduce:

$$\begin{split} \sup_{\tau \in (\eta,s)} &\int_{\mathbb{R}^d} u^2(\tau,x) \gamma_R^2(x) H(\tau,x) dx \leq \int_{\mathbb{R}^d} u_0^2(x) \gamma_R^2(x) H(\eta,x) dx \\ &+ \int_{\eta}^s \int_{\mathbb{R}^d} u^2(t,x) \gamma_R^2(x) \left(2c_2 \Lambda \Gamma_{\rho}^{\alpha} (H^{1/2}, H^{1/2})(t,x) + \partial_t H(t,x) \right) dx dt \\ &+ 2c_2 \Lambda \int_{\eta}^s \int_{\mathbb{R}^d} \int_{B_{\rho}(x)} u^2(t,x) \gamma_R^2(z) (H^{1/2}(t,x) - H^{1/2}(t,z))^2 |x-y|^{-d-\alpha} dz dx dt \\ &+ 2c_2 \Lambda \int_{\eta}^s \int_{\mathbb{R}^d} u^2(t,x) H(t,x) \Gamma_{\rho}^{\alpha} (\gamma_R,\gamma_R) (x) dx dt \\ &+ 2c_2 \Lambda \int_{\eta}^s \int_{\mathbb{R}^d} \int_{B_{\rho}(x)} u^2(t,z) H(t,x) (\gamma_R(x) - \gamma_R(z))^2 |x-y|^{-d-\alpha} dz dx dt \\ &\leq \int_{\mathbb{R}^d} u_0^2(x) \gamma_R^2(x) H(\eta,x) dx \\ &+ \int_{\eta}^s \int_{\mathbb{R}^d} u^2(t,x) \gamma_R^2(x) \left(4c_2 \Lambda \Gamma_{\rho}^{\alpha} (H^{1/2}, H^{1/2})(t,x) + \partial_t H(t,x) \right) dx dt \\ &+ 2c_2 \Lambda \|u\|_{\infty}^2 \int_{\eta}^s \int_{\mathbb{R}^d \setminus B_{\frac{R-1}{2}(0)}} \Gamma_{\rho}^{\alpha} (H^{1/2}, H^{1/2})(t,x) dx dt \\ &+ 4c_2 \Lambda \|u\|_{\infty}^2 \int_{\eta}^s \int_{\mathbb{R}^d} \Gamma_{\rho}^{\alpha} (\gamma_R,\gamma_R)(x) H(t,x) dx dt. \end{split}$$

Now, assume that H satisfies the assumption with $C = 4c_2\Lambda$. Then, using also (3.2), the following holds true:

$$\begin{split} \left(4c_2\Lambda\Gamma_{\rho}^{\alpha}(H^{1/2},H^{1/2})(t,x)+\partial_t H(t,x)\right) &\leq 0, \quad t\in(\eta,s), \ x\in\mathbb{R}^d,\\ \int_{\eta}^s \int_{\mathbb{R}^d\setminus B_{\frac{R-1}{2}}(0)} \Gamma_{\rho}^{\alpha}(H^{1/2},H^{1/2})(t,x)\mathrm{d}x\mathrm{d}t\to 0, \quad \mathrm{as}\ R\to\infty,\\ \int_{\eta}^s \int_{\mathbb{R}^d} \Gamma_{\rho}^{\alpha}(\gamma_R,\gamma_R)(x)H(t,x)\mathrm{d}x\mathrm{d}t &\leq c_1 \int_{\eta}^s \int_{\mathbb{R}^d\setminus B_{\frac{R-1}{2}}(0)} H(t,x)\mathrm{d}x\mathrm{d}t\to 0, \quad \mathrm{as}\ R\to\infty. \end{split}$$

Upon the observation that $\gamma_R \to 1$, as $R \to \infty$, it follows

$$\sup_{\tau \in (\eta,s)} \int_{\mathbb{R}^d} u^2(\tau,x) H(\tau,x) \mathrm{d}x \le \int_{\mathbb{R}^d} u_0^2(x) H(\eta,x) \mathrm{d}x,$$

as desired.

Our next goal is to establish the following auxiliary estimate:

Theorem 3.2. Assume that k satisfies (k_{\leq}) , (\mathcal{E}_{\geq}) . Let $y \in \mathbb{R}^d$, $\sigma, \rho > 0$, $\eta \geq 0$. Let $u_0 \in L^2(\mathbb{R}^d)$ be such that $u_0 \equiv 0$ in $B_{\sigma}(y)$. Assume that $u \in L^{\infty}((\eta, T) \times \mathbb{R}^d)$ is a weak solution to $\partial_t u - L_t^{\rho} u = 0$ in $(\eta, T) \times \mathbb{R}^d$. Then there exist $\nu > 1, C > 0$ depending on $d, \alpha, \lambda, \Lambda$ such that for every $s \in (\eta, T)$ with $s - \eta \leq \frac{1}{4\nu}\rho^{\alpha}$:

$$|u(s,y)| \le C(s-\eta)^{-\frac{d}{2\alpha}} 2^{\frac{\sigma}{6\rho}} \left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)^{-\frac{\sigma}{6\rho} + \frac{1}{2} + \frac{d}{2\alpha}} \|u_0\|_{L^2(\mathbb{R}^d)}.$$

The idea to prove Theorem 3.2 is to find a suitable function H such that Lemma 3.1 is applicable. In the following, we present a suitable such function.

Given $y \in \mathbb{R}^d$, $\rho > 0$, $0 \le \eta < s < T$, $\nu > 1$ with $s - \eta \le \frac{1}{4\nu}\rho^{\alpha}$, we define $H_{y,\rho,\eta,s,\nu} = H$: $[\eta, s] \times \mathbb{R}^d \to [0, \infty)$ via

$$H(t,x) := \left(\frac{\rho^{\alpha}}{\nu[2(s-\eta) - (t-\eta)]}\right)^{-1} \wedge \left(\frac{\rho^{\alpha}}{\nu[2(s-\eta) - (t-\eta)]}\right)^{-\frac{|x-y|}{3\rho}} = e^{-\log\left(\frac{\rho^{\alpha}}{\nu[2(s-\eta) - (t-\eta)]}\right)\left(\frac{|x-y|}{3\rho} \vee 1\right)}.$$

Lemma 3.3. For every C > 0, there exists $\nu = \nu(d, \alpha, C) > 1$ such that for every $y \in \mathbb{R}^d$, $\rho > 0, 0 \le \eta < s < T$ with $s - \eta \le \frac{1}{4\nu}\rho^{\alpha}$, the function $H_{y,\rho,\eta,s,\nu} = H$ defined above satisfies

$$C\Gamma^{\alpha}_{\rho}(H^{1/2}, H^{1/2}) \leq -\partial_t H, \quad in \ (\eta, s) \times \mathbb{R}^d, \tag{3.3}$$

$$H^{1/2} \in L^2((\eta, s); H^{\alpha/2}(\mathbb{R}^d)).$$
 (3.4)

Proof. Let $y \in \mathbb{R}^d$, $\rho > 0$, $0 \le \eta < s < T$, with $s - \eta \le \frac{1}{4\nu}\rho^{\alpha}$, where $\nu > 1$ to be chosen later. Recall:

$$H(t,x) := \left(\frac{\rho^{\alpha}}{\nu[2(s-\eta) - (t-\eta)]}\right)^{-1} \wedge \left(\frac{\rho^{\alpha}}{\nu[2(s-\eta) - (t-\eta)]}\right)^{-\frac{|x-y|}{3\rho}} = e^{-\log\left(\frac{\rho^{\alpha}}{\nu[2(s-\eta) - (t-\eta)]}\right)\left(\frac{|x-y|}{3\rho} \vee 1\right)}.$$
(3.5)

Note that by assumption, $\frac{\rho^{\alpha}}{\nu[2(s-\eta)-(t-\eta)]} > 1$ for every $t \in [\eta, s]$. Let $t \in [\eta, s]$ be fixed. We split the proof of (3.3) into three cases.

Case 1: $|x - y| \le 2\rho$. In this case, trivially $\Gamma^{\alpha}_{\rho}(H^{1/2}, H^{1/2})(t, x) = 0$, and

$$-\partial_t H(t,x) = -\partial_t \left(\frac{\nu[2(s-\eta) - (t-\eta)]}{\rho^{\alpha}} \right) = \nu \rho^{-\alpha} > 0$$

Therefore, (3.3) holds true for any $\nu > 0$.

Case 2: $2\rho \leq |x-y| \leq 3\rho$. In this case, $-\partial_t H(t,x) = \nu \rho^{-\alpha}$, as in Case 1. Moreover, let us fix $x_0 \in B_\rho(x)$ with $|x_0-y| = 3\rho$ such that for every $z \in \mathbb{R}^d \setminus B_{3\rho}(y)$ it holds $|x_0-z| \leq 2|x-z|$. This holds true, e.g. if one chooses x_0 as a point on $\partial B_{3\rho}(y)$ that minimizes dist $(x, \mathbb{R}^d \setminus B_{3\rho}(y))$, since then by triangle inequality: $|x_0-z| \leq |x_0-x| + |x-z| \leq 2|x-z|$. Note that $H(t,x) = H(t,x_0)$ and $B_\rho(x) \subset B_{2\rho}(x_0)$. Therefore:

$$\begin{split} \Gamma_{\rho}^{\alpha}(H^{1/2}, H^{1/2})(t, x) &= \int_{B_{\rho}(x) \setminus B_{3\rho}(y)} \left(H^{1/2}(t, x) - H^{1/2}(t, z) \right)^{2} |x - z|^{-d - \alpha} \mathrm{d}z \\ &\leq c_{1} \int_{B_{2\rho}(x_{0}) \setminus B_{3\rho}(y)} \left(H^{1/2}(t, x_{0}) - H^{1/2}(t, z) \right)^{2} |x_{0} - z|^{-d - \alpha} \mathrm{d}z \\ &\leq \int_{B_{2\rho}(x_{0}) \setminus B_{3\rho}(y)} |\nabla H^{1/2}(t, x_{0})|^{2} |x_{0} - z|^{2 - d - \alpha} \mathrm{d}z \\ &\leq c_{2} |\nabla H^{1/2}(t, x_{0})|^{2} \rho^{2 - \alpha} \\ &= c_{2} \left((6\rho)^{-1} \log \left(\frac{\rho^{\alpha}}{\nu[2(s - \eta) - (t - \eta)]} \right) e^{-\log \left(\frac{\rho^{\alpha}}{\nu[2(s - \eta) - (t - \eta)]} \right)^{2} \left(\frac{\rho^{\alpha}}{\nu[2(s - \eta) - (t - \eta)]} \right)^{2} \rho^{2 - \alpha} \\ &= c_{3} \rho^{-\alpha} \left(\log \left(\frac{\rho^{\alpha}}{\nu[2(s - \eta) - (t - \eta)]} \right) \left(\frac{\rho^{\alpha}}{\nu[2(s - \eta) - (t - \eta)]} \right)^{-1/2} \right)^{2} \end{split}$$

for some $c_1, c_2, c_3, c_4 > 0$. We used the inequality $\log(a) \leq a^{1/2}$ in the last step. Moreover, note that the estimate $(H^{1/2}(t, x_0) - H^{1/2}(t, z))^2 \leq |\nabla H^{1/2}(t, x_0)|^2 |x_0 - z|^2$ is correct since $\sup_{z \in \mathbb{R}^d \setminus B_{3\rho}(y)} |\nabla H^{1/2}(t, z)| = |\nabla H^{1/2}(t, x_0)|$ due to $x_0 \in \partial B_{3\rho}(y)$. Therefore, (3.3) holds true in this case for any $\nu > c_4 C$.

Case 3: $|x - y| > 3\rho$. In this case,

$$-\partial_t H(t,x) = \frac{|x-y|}{3\rho[2(s-\eta) - (t-\eta)]} e^{-\log\left(\frac{\rho^{\alpha}}{\nu[2(s-\eta) - (t-\eta)]}\right)\left(\frac{|x-y|}{3\rho}\right)}.$$

Moreover:

$$\begin{split} \Gamma_{\rho}^{\alpha}(H^{1/2}, H^{1/2})(t, x) &= \int_{B_{\rho}(x)} \left(H^{1/2}(t, x) - H^{1/2}(t, z) \right)^{2} |x - z|^{-d - \alpha} \mathrm{d}z \\ &\leq \int_{B_{\rho}(x)} \left(e^{-\log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}\right) \left(\frac{|x - y|}{6\rho}\right)} - e^{-\log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}\right) \left(\frac{|z - y|}{6\rho}\right)} \right)^{2} |x - z|^{-d - \alpha} \mathrm{d}z \\ &\leq \int_{B_{\rho}(x)} \sup_{z \in B_{\rho}(x)} \left| \nabla e^{-\log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}\right) \left(\frac{|z - y|}{6\rho}\right)} \right|^{2} |x - z|^{2 - d - \alpha} \mathrm{d}z \\ &\leq c_{5} \left((6\rho)^{-1} \log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}\right) e^{-\log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}\right) \left(\frac{|x - y|}{6\rho}\right)} \right)^{2} \rho^{2 - \alpha} \\ &= c_{6} \left[\log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}}\right) \right]^{2} \left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}}\right)^{1/3} e^{-\log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}\right) \left(\frac{|x - y|}{3\rho}\right)} \\ &\leq c_{7} \frac{1}{2\rho\nu^{[2(s - \eta) - (t - \eta)]}} e^{-\log\left(\frac{\rho^{\alpha}}{\nu^{[2(s - \eta) - (t - \eta)]}\right) \left(\frac{|x - y|}{3\rho}\right)} \end{split}$$

for some $c_5, c_6, c_7 > 0$. In the third inequality, we used $|z - y| \ge |x - y| - \rho$, and in the second to last step we applied the estimate $\log(a) \le ca^{1/3}$. This holds with c > 0 independent of a > 1. Therefore, by choosing $\nu > c_7 C$, (3.3) is satisfied also in this case.

Together, we have proved (3.3). Finally, note that $\Gamma^{\alpha}_{\rho}(H^{1/2}, H^{1/2}) \in L^1((\eta, s) \times \mathbb{R}^d)$ since for

 $|x-y| \ge 3\rho$, we computed above

$$\Gamma^{\alpha}_{\rho}(H^{1/2}, H^{1/2})(t, x) \le |x - y| c^{-\frac{|x - y|}{3\rho}},$$

where c > 1 is a constant that might depend on η, s, ρ . This proves (3.4).

Having at hand the function H defined in (3.5), it is possible to establish Theorem 3.2.

Proof of Theorem 3.2. The idea is to apply Lemma 3.1 with H as in Lemma 3.3. It follows that for every $y \in \mathbb{R}^d$, $0 \le \eta < s < T$ with $s - \eta \le \frac{1}{4\nu}\rho^{\alpha}$:

$$\sup_{\tau \in (\eta,s)} \int_{B_{2\rho}(y)} u^2(\tau,x) H(\tau,x) \mathrm{d}x \le \sup_{\tau \in (\eta,s)} \int_{\mathbb{R}^d} u^2(\tau,x) H(\tau,x) \mathrm{d}x \le \int_{\mathbb{R}^d \setminus B_\sigma(y)} u_0^2(x) H(\eta,x) \mathrm{d}x.$$

Consequently,

$$\sup_{\tau \in (\eta,s)} \int_{B_{2\rho}(y)} u^2(\tau,x) \mathrm{d}x \le \left(\frac{\sup_{x \in \mathbb{R}^d \setminus B_{\sigma}(y)} H(\eta,x)}{\inf_{\tau \in (\eta,s), x \in B_{2\rho}(y)} H(\tau,x)} \right) \|u_0\|_{L^2(\mathbb{R}^d)}^2.$$

By the truncated $L^{\infty} - L^2$ -estimate Lemma 2.4, applied with $R = (s - \eta)^{1/\alpha}$, $t_0 = s$, $x_0 = y$:

$$\sup_{(\eta',s)\times B_{\frac{1}{2}(s-\eta)^{1/\alpha}}(y)} u \le c_1(s-\eta)^{-\frac{d}{2\alpha}} \left(\frac{\rho^{\alpha}}{s-\eta}\right)^{\frac{d}{2\alpha}} \sup_{\tau\in(\eta,s)} \left(\int_{B_{2\rho}(y)} u^2(\tau,x) \mathrm{d}x\right)^{1/2} \\ \le c_1(s-\eta)^{-\frac{d}{2\alpha}} \left(\frac{\rho^{\alpha}}{s-\eta}\right)^{\frac{d}{2\alpha}} \left(\frac{\sup_{x\in\mathbb{R}^d\setminus B_{\sigma}(y)} H(\eta,x)}{\inf_{\tau\in(\eta,s),x\in B_{2\rho}(y)} H(\tau,x)}\right)^{1/2} \|u_0\|_{L^2(\mathbb{R}^d)}$$

for some $c_1 > 0$, where $\eta' := s - 2^{-\alpha}(s - \eta) \in (\eta, s)$. Note that there exist $c_2, c_3 > 0$ such that for $x \in \mathbb{R}^d \setminus B_{\sigma}(y)$ it holds

$$H(\eta, x) \le c_2 \left(\frac{\rho^{\alpha}}{2\nu(s-\eta)}\right)^{-\frac{\sigma}{3\rho}}$$

and for $(\tau, x) \in [\eta, s] \times B_{2\rho}(y)$ we have:

$$H(\tau, x) \ge e^{-\log\left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)\left(\frac{2\rho}{6\rho} \lor 1\right)} \ge c_3 e^{-\log\left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)} \ge c_3 \left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)^{-1}$$

This follows directly from the definition of H and $s - \eta \leq \frac{1}{4\nu}\rho^{\alpha}$. Together, we obtain

$$|u(s,y)| \le c_4(s-\eta)^{-\frac{d}{2\alpha}} 2^{\frac{\sigma}{6\rho}} \left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)^{-\frac{\sigma}{6\rho}+\frac{1}{2}+\frac{d}{2\alpha}} \|u_0\|_{L^2(\mathbb{R}^d)}$$

as desired.

for some $c_4 > 0$, as desired.

Having proved Theorem 3.2, we are now in the position to establish upper off-diagonal bounds for $p_{\rho}(y, s; x, \eta)$:

Theorem 3.4. Assume that k satisfies (k_{\leq}) , (\mathcal{E}_{\geq}) . Then there exists c > 0 depending on $d, \alpha, \lambda, \Lambda$ such that for every $\rho > 0$, $0 \leq \eta < s < T$, and $x, y \in \mathbb{R}^d$ with $s - \eta \leq \frac{1}{4\nu}\rho^{\alpha}$:

$$p_{\rho}(y,s;x,\eta) \le c(s-\eta)^{-\frac{d}{\alpha}} 2^{\frac{|x-y|}{12\rho}} \left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)^{-\frac{|x-y|}{12\rho} + \frac{1}{2} + \frac{d}{2\alpha}}.$$
(3.6)

Proof. Note that the on-diagonal bound (2.14) and (2.8) immediately imply for every $0 \le \eta < s < T$ with $s - \eta \le \frac{1}{4\nu} \rho^{\alpha}$ and $x, y \in \mathbb{R}^d$:

$$\left(\int_{\mathbb{R}^d} p_\rho^2(z,s;x,\eta) \mathrm{d}z\right)^{1/2} \le c_1(s-\eta)^{-\frac{d}{2\alpha}}$$
(3.7)

for some $c_1 > 0$. On the other hand, from Theorem 3.2, it follows for every $0 \le \eta < s < T$ with $s - \eta \le \frac{1}{4\nu}\rho^{\alpha}$ and $x, y \in \mathbb{R}^d$:

$$\left(\int_{\mathbb{R}^d \setminus B_{\sigma}(y)} p_{\rho}^2(y,s;z,\eta) \mathrm{d}z\right)^{1/2} \le c_2(s-\eta)^{-\frac{d}{2\alpha}} 2^{\frac{\sigma}{6\rho}} \left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)^{-\frac{\sigma}{6\rho}+\frac{1}{2}+\frac{d}{2\alpha}}$$
(3.8)

for some $c_2 > 0$. To see this, one observes that $u(t,x) = \int_{\mathbb{R}^d \setminus B_{\sigma}(y)} p_{\rho}(x,t;z,\eta) p_{\rho}(y,s;z,\eta) dz$ satisfies the assumptions of Theorem 3.2 with

$$u_0(x) = p_\rho(y,s;x,\eta) \mathbb{1}_{\{|x-y| > \sigma\}}(x), \quad u(s,y) = \int_{\mathbb{R}^d \setminus B_\sigma(y)} p_\rho^2(y,s;z,\eta) \mathrm{d}z.$$

To prove (3.6), let us fix $0 \le \eta < s < T$ with $s - \eta \le \frac{1}{4\nu}\rho^{\alpha}$, and $x, y \in \mathbb{R}^d$. Then we define $\sigma = \frac{1}{2}|x-y|$ and compute, using (2.9):

$$p_{\rho}(y,s;x,\eta) = \int_{\mathbb{R}^d} p_{\rho}(y,s;z,(s-\eta)/2) p_{\rho}(z,(s-\eta)/2;x,\eta) dz$$

=
$$\int_{\mathbb{R}^d \setminus B_{\sigma}(y)} p_{\rho}(y,s;z,(s-\eta)/2) p_{\rho}(z,(s-\eta)/2;x,\eta) dz$$

+
$$\int_{B_{\sigma}(y)} p_{\rho}(y,s;z,(s-\eta)/2) p_{\rho}(z,(s-\eta)/2;x,\eta) dz$$

=
$$J_1 + J_2.$$

For J_1 , we compute, using (3.7), (3.8):

$$J_{1} \leq \left(\int_{\mathbb{R}^{d} \setminus B_{\sigma}(y)} p_{\rho}^{2}(y,s;z,(s-\eta)/2) \mathrm{d}z\right)^{1/2} \left(\int_{\mathbb{R}^{d} \setminus B_{\sigma}(y)} p_{\rho}^{2}(z,(s-\eta)/2;x,\eta) \mathrm{d}z\right)^{1/2} \\ \leq c_{3}(s-\eta)^{-\frac{d}{\alpha}} 2^{\frac{|x-y|}{12\rho}} \left(\frac{\rho^{\alpha}}{\nu(s-\eta)}\right)^{-\frac{|x-y|}{12\rho} + \frac{1}{2} + \frac{d}{2\alpha}}$$

for some $c_3 > 0$. For J_2 , observe that $B_{\sigma}(y) \subset \mathbb{R}^d \setminus B_{\sigma}(x)$, and therefore by (2.7):

$$J_{2} \leq \left(\int_{\mathbb{R}^{d} \setminus B_{\sigma}(x)} p_{\rho}^{2}(y,s;z,(s-\eta)/2) \mathrm{d}z \right)^{1/2} \left(\int_{\mathbb{R}^{d} \setminus B_{\sigma}(x)} p_{\rho}^{2}(z,(s-\eta)/2;x,\eta) \mathrm{d}z \right)^{1/2} \\ \leq c_{4}(s-\eta)^{-\frac{d}{\alpha}} 2^{\frac{|x-y|}{12\rho}} \left(\frac{\rho^{\alpha}}{\nu(s-\eta)} \right)^{-\frac{|x-y|}{12\rho} + \frac{1}{2} + \frac{d}{2\alpha}}$$

for some $c_4 > 0$. Together, we obtain the desired result.

Bounds for the heat kernel corresponding to the truncated jumping kernel
$$p_{\rho}$$
 imply bounds
for p with the help of the gluing lemma Lemma 2.2. The underlying argument is known
among probabilists as 'Meyer's decomposition'. For an analytic proof relying on the parabolic
maximum principle we refer to the appendix.

We are now ready to provide the proof of our main result Theorem 1.1:

Proof of Theorem 1.1. Let $x, y \in \mathbb{R}^d$ be fixed. By (2.13) it suffices to prove that for some constants $c_0, c_1 > 0$ and $s - \eta \leq c_0 |x - y|^{\alpha}$ it holds

$$p(y,s;x,\eta) \le c_1 \frac{s-\eta}{|x-y|^{d+\alpha}}.$$

By Lemma 2.2, and (k_{\leq}) we know that for every $\rho > 0, 0 \leq \eta < s < T$:

$$p(y,s;x,\eta) \le p_{\rho}(y,s;x,\eta) + c_{2}(s-\eta) \|k-k\mathbb{1}_{\{|x-y|\le\rho\}}\|_{\infty} \le p_{\rho}(y,s;x,\eta) + c_{3}(s-\eta)\rho^{-d-\alpha}$$
(3.9)

for some $c_2, c_3 > 0$. We choose $\rho = \frac{|x-y|}{12} \left(\frac{d+\alpha}{\alpha} + \frac{1}{2} + \frac{d}{2\alpha}\right)^{-1}$. Then by (3.6) and (3.9) it holds for $s - \eta \leq \frac{1}{4\nu}\rho^{\alpha}$:

$$p(y,s;x,\eta) \le c_4(s-\eta)|x-y|^{-d-\alpha},$$

where $c_4 > 0$, as desired.

4. EXTENSION: JUMPING KERNELS OF MIXED TYPE ON METRIC MEASURE SPACES

In this section, we discuss a possible extension of the nonlocal Aronson method to jumping kernels of mixed type. Moreover, we work on a general doubling metric measure space.

Let (M, d) be a locally compact, separable metric space, and let μ be a positive Radon measure with full support. We assume that (M, d, μ) satisfies the volume doubling property, i.e., there exists $C > 0, d \in \mathbb{N}$ such that

$$\frac{\mu(B_R(x))}{\mu(B_r(x))} \le C\left(\frac{R}{r}\right)^d, \quad x \in M, \ 0 < r \le R.$$
(VD)

Note that as a consequence of (VD), for every $\delta > 0$ there exist $c_1, c_2 > 0$ such that for every $R > 0, x, y \in M$ with $d(x, y) \leq \delta R$: $c_1 \mu(B_R(x)) \leq \mu(B_R(y)) \leq c_2 \mu(B_R(x))$.

Moreover, let $\phi: [0,\infty) \to [0,\infty)$ be strictly increasing with $\phi(0) = 0, \phi(1) = 1$ and

$$C^{-1}\left(\frac{R}{r}\right)^{\alpha_1} \le \frac{\phi(R)}{\phi(r)} \le C\left(\frac{R}{r}\right)^{\alpha_2}, \quad 0 < r \le R,$$
(4.1)

for some constant C > 0 and $0 < \alpha_1 \le \alpha_2 < 2$. For a detailed discussion of the setup, we refer to [CKW21].

Consider symmetric iumping kernels $k \cdot (0, T) \times M \times M \rightarrow \mathbb{R}$ satisfying for

onsider symmetric jumping kernels
$$k: (0,T) \times M \times M \to \mathbb{R}$$
 satisfying for some $\Lambda > 0$

$$k(t;x,y) \le \Lambda \mu(B_{d(x,y)}(x))^{-1} \phi(d(x,y))^{-1}, \ x,y \in M,$$
 (k_{\le})

and assume that there is $\mathcal{F} \subset L^2(M,\mu)$ such that $(\mathcal{E}_t,\mathcal{F})$ is a regular Dirichlet form on $L^2(M,\mu)$ for every $t \in (0,T)$, where for every $u, v \in \mathcal{F}$,

$$\mathcal{E}_t(u,v) = \int_M \int_M (u(x) - u(y))(v(x) - v(y))k(t;x,y) \mathrm{d}x \mathrm{d}y$$

is defined in the usual way. For simplicity, we write $dx := \mu(dx)$.

Moreover, we assume that the Faber-Krahn inequality holds true, i.e., that there exist $c, \nu > 0$ such that for all $t \in (0,T)$, R > 0, $x_0 \in M$, $D \subset B_R(x_0)$ and every $u \in \mathcal{F}$ with $u \equiv 0$ in $M \setminus D$:

$$\mathcal{E}_t(u,u) \ge c\phi(R)^{-1} \left(\frac{\mu(B_R(x_0))}{\mu(D)}\right)^{\nu} \|u\|_{L^2(D)}^2.$$
 (FK)

Let $p(y, s; x, \eta)$ be the fundamental solution to the Cauchy problem associated with k.

Theorem 4.1. Let (M, d, μ) and ϕ be as above, and assume (VD), (4.1). Assume that k satisfies (k_{\leq}) and (FK). Then there exists c > 0 such that for every $0 \leq \eta < s < T$, $x, y \in M$:

$$p(y,s;x,\eta) \le c \left[\mu(B_{\phi^{-1}(s-\eta)}(x))^{-1} \wedge \frac{s-\eta}{\mu(B_{d(x,y)}(x))\phi(d(x,y))} \right].$$
(4.2)

We remark that variants of Theorem 4.1 for time-homogeneous jumping kernels of mixed type on doubling metric measure spaces can be found in several articles, e.g., [CK08], [CKW21], [CKKW21].

Proof. First of all, we observe that by (VD), (4.1), (k_{\leq}) :

$$\int_{B_R(x)} d(x,y)^2 k(t;x,y) dy \le cR^2 \phi(R)^{-1}, \qquad \int_{M \setminus B_R(x)} k(t;x,y) dy \le c\phi(R)^{-1}.$$
(4.3)

for every $t \in (0,T)$, $x \in M$, R > 0. For a proof, see [CKW21]. Given $y \in M$, $\rho > 0$, $0 \le \eta < s < T$, $\nu > 1$ with $s - \eta \le \frac{1}{4\nu}\phi(\rho)$, we define $H_{y,\rho,\eta,s,\nu} = H : [\eta,s] \times M \to [0,\infty)$ via

$$H(t,x) := \left(\frac{\phi(\rho)}{\nu[2(s-\eta) - (t-\eta)]}\right)^{-1} \wedge \left(\frac{\phi(\rho)}{\nu[2(s-\eta) - (t-\eta)]}\right)^{-\frac{d(x,y)}{3\rho}}$$

With the help of (4.3), (4.1) it is easy to check along the lines of Lemma 3.3 that H satisfies the assumptions of Lemma 3.1, namely for every C > 0 there exists $\nu > 1$ such that

$$C\Gamma^{\phi}_{\rho}(H^{1/2}, H^{1/2}) \leq -\partial_t H \quad \text{in } (\eta, s) \times M, \quad H^{1/2} \in L^2((\eta, s); \mathcal{F}),$$

where

$$\Gamma^{\phi}_{\rho}(H^{1/2}, H^{1/2})(t, x) = \int_{B_{\rho}(x)} (H^{1/2}(t, x) - H^{1/2}(t, z))^2 \mu(B_{d(x, z)}(x))^{-1} \phi(d(x, z))^{-1} \mathrm{d}z.$$

Next, we observe that by (VD), (4.3), (4.1), (k_{\leq}) , and (FK) the following $L^{\infty} - L^2$ -estimate holds true for local subsolutions u to $\partial_t u - L_t^{\rho} u = 0$ in $I_R(t_0) \times B_R(x_0)$, where $I_R(t_0) = (t_0 - \phi(R), t_0)$:

$$\sup_{I_{R/2}(t_0) \times B_{R/2}(x_0)} u \le C \left(\frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x_0))}\right)^{\frac{1}{2}} \mu(B_R(x_0))^{-\frac{1}{2}} \sup_{t \in I_R(t_0)} \left(\int_{B_{2\rho}(x_0)} u^2(t, x) \mathrm{d}x\right)^{1/2}.$$
 (4.4)

We give a proof of this fact later, see Lemma 4.2.

Applying H to Lemma 3.1, we obtain from (4.4) and the definition of H:

$$|u(s,y)| \le c_1 \left(\frac{\mu(B_{\rho}(y))}{\mu(B_{\phi^{-1}(s-\eta)}(y))}\right)^{\frac{1}{2}} \mu(B_{\phi^{-1}(s-\eta)}(y))^{-\frac{1}{2}} 2^{\frac{\sigma}{6\rho}} \left(\frac{\phi(\rho)}{\nu(s-\eta)}\right)^{-\frac{\sigma}{6\rho}+\frac{1}{2}} \|u_0\|_{L^2(D)}$$
(4.5)

for $c_1 > 0$, where u, u_0 are as in Lemma 3.1, and $0 \le \eta < s < T$ with $s - \eta \le \frac{1}{4\nu}\phi(\rho), \sigma > 0$. Moreover, the following on-diagonal estimates hold for every $0 \le \eta < s < T, x, y \in M, \rho > 0$:

$$p(y,s;x,\eta) \le c\mu (B_{\phi^{-1}(s-\eta)}(x))^{-1},$$
(4.6)

$$p_{\rho}(y,s;x,\eta) \le c e^{c(s-\eta)\phi(\rho)^{-1}} \mu(B_{\phi^{-1}(s-\eta)}(x))^{-1}, \tag{4.7}$$

for some c > 0. These estimates are proved in Lemma 5.1 and Section 4.4 in [CKW21] using a stochastic approach. A more direct proof, using only analysis tools, goes via the $L^{\infty} - L^{1}$ estimate (4.12). In fact, given $0 \leq \eta < s < T$, $x, y \in M$, we apply (4.12) to $(t, z) \mapsto p(y, t; z, \eta)$, choosing $R := \phi^{-1}(s - \eta), x_0 := x, t_0 := s$. Using (2.8), we obtain

$$p(y,s;x,\eta) \le c\mu(B_{\phi^{-1}(s-\eta)}(x))^{-1} \sup_{t \in I_{\phi^{-1}(s-\eta)}(s)} \int_M p(y,t;z,\eta) \mathrm{d}z \le c\mu(B_{\phi^{-1}(s-\eta)}(x))^{-1},$$

as desired. (4.7) is a direct consequence of (4.6) in the light of (5.7) and (4.3). Note that the proof of (5.7) is written for $M = \mathbb{R}^d$ but works in the same way in the current setup. Combining (4.5) and (4.7), we derive as in Theorem 3.4 for every $0 \leq \eta < s < T$ with $s - \eta \leq \frac{1}{4\nu}\phi(\rho), x, y \in M$:

$$p_{\rho}(y,s;x,\eta) \le c_2 \left(\frac{\rho}{\phi^{-1}(s-\eta)}\right)^{\frac{d}{2}} \mu(B_{\phi^{-1}(s-\eta)}(x))^{-\frac{1}{2}} \mu(B_{\phi^{-1}(s-\eta)}(y))^{-\frac{1}{2}} 2^{\frac{d(x,y)}{12\rho}} \left(\frac{\phi(\rho)}{\nu(s-\eta)}\right)^{-\frac{d(x,y)}{12\rho}+\frac{1}{2}}$$

for $c_2 > 0$. Finally, we explain how to deduce off-diagonal bounds for p. We get from (5.6):

$$p(y,s;x,\eta) \le p_{\rho}(y,s;x,\eta) + \int_{\eta}^{s} P_{\tau}^{\rho} K_{\rho}(y) \mathrm{d}\tau.$$

$$(4.8)$$

We choose $\rho = \frac{d(x,y)}{12} \left(\frac{d+\alpha_1}{\alpha_1} + \frac{1}{2} + \frac{d}{2\alpha_1} \right)^{-1}$ and obtain by (VD), (4.1):

$$p_{\rho}(y,s;x,\eta) \leq c_{2} \left(\frac{\rho}{\phi^{-1}(s-\eta)}\right)^{\frac{d}{2}} \left[\mu(B_{\phi^{-1}(s-\eta)}(x))\mu(B_{\phi^{-1}(s-\eta)}(y))\right]^{-\frac{1}{2}} 2^{\frac{d(x,y)}{12\rho}} \left(\frac{\phi(\rho)}{\nu(s-\eta)}\right)^{-\frac{d(x,y)}{12\rho}+\frac{1}{2}} \\ \leq c_{3} \left(\frac{d(x,y)}{\phi^{-1}(s-\eta)}\right)^{\frac{d}{2}} \left[\mu(B_{\phi^{-1}(s-\eta)}(x))\mu(B_{\phi^{-1}(s-\eta)}(y))\right]^{-\frac{1}{2}} \left(\frac{\phi(d(x,y))}{s-\eta}\right)^{-\frac{3d}{2\alpha_{1}}-1} \\ \leq c_{4} \left(\frac{\phi^{-1}(\phi(d(x,y)))}{\phi^{-1}(s-\eta)}\right)^{\frac{d}{2}} \left(\frac{\mu(B_{\phi^{-1}(\phi(d(x,y)))}(x))\mu(B_{\phi^{-1}(s-\eta)}(y))}{\mu(B_{\phi^{-1}(s-\eta)}(x))\mu(B_{\phi^{-1}(s-\eta)}(y))}\right)^{\frac{1}{2}} \frac{\left(\frac{\phi(d(x,y))}{s-\eta}\right)^{-\frac{3d}{2\alpha_{1}}-1}}{\mu(B_{d(x,y)}(x))} \\ \leq c_{5} \left(\frac{\phi(d(x,y))}{s-\eta}\right)^{\frac{3d}{2\alpha_{1}}} \left(\frac{\phi(d(x,y))}{s-\eta}\right)^{-\frac{3d}{2\alpha_{1}}-1}}{\mu(B_{d(x,y)}(x))^{-1}} \\ = c_{5} \frac{s-\eta}{\mu(B_{d(x,y)}(x))\phi(d(x,y))}$$

$$(4.9)$$

for $c_3, c_4, c_5 > 0$. Next, we estimate $\int_n^s P_\tau^\rho K_\rho(y) d\tau$. For this, we compute by (k_{\leq}) and (VD):

$$\int_{\eta}^{s} P_{\tau}^{\rho} K_{\rho}(y) d\tau = \sum_{k=1}^{\infty} \int_{\eta}^{s} P_{\tau}^{\rho} \left[\mathbbm{1}_{B_{ck\rho}(y) \setminus B_{c(k-1)\rho}(y)} K_{\rho} \right](y) d\tau$$

$$\leq c_{6} \phi(d(x,y))^{-1} \sum_{k=1}^{\infty} \int_{\eta}^{s} P_{\tau}^{\rho} \left[\mathbbm{1}_{B_{ck\rho}(y) \setminus B_{c(k-1)\rho}(y)} \mu(B_{\rho}(\cdot))^{-1} \right](y) d\tau$$

$$\leq c_{7} \mu(B_{\rho}(x))^{-1} \phi(d(x,y))^{-1} \sum_{k=1}^{\infty} k^{d} \int_{\eta}^{s} P_{\tau}^{\rho} \mathbbm{1}_{B_{ck\rho}(y) \setminus B_{c(k-1)\rho}(y)}(y) d\tau$$

for $c_6, c_7 > 0$, and $c > 3 + \frac{6d}{\alpha_1}$. Using (4.5), (VD) and (4.1), we estimate for $\tau \in (\eta, s), k \ge 2$:

$$k^{d} P^{\rho}_{\tau} \mathbb{1}_{B_{ck\rho}(y) \setminus B_{c(k-1)\rho}(y)}(y) \le c_{8} k^{d} \left(\frac{\mu(B_{k\rho}(y))}{\mu(B_{\phi^{-1}(\tau-\eta)}(y))} \right) 2^{\frac{c(k-1)}{6}} \left(\frac{\phi(\rho)}{\nu(\tau-\eta)} \right)^{-\frac{c(k-1)}{6} + \frac{1}{2}}$$

$$\leq c_9 k^{2d} 2^{\frac{c(k-1)}{6}} \left(\frac{\phi(\rho)}{\nu(\tau-\eta)}\right)^{-\frac{c(k-1)}{6} + \frac{1}{2} + \frac{d}{\alpha_1}} \\ \leq c_{10} k^{2d} 2^{\frac{c(k-1)}{6}} 4^{-\frac{c(k-1)}{6} + \frac{1}{2} + \frac{d}{\alpha_1}}$$

for $c_8, c_9, c_{10} > 0$, where we use $s - \eta \leq \frac{1}{4\nu}\phi(\rho)$. From (2.8), it follows

$$\int_{\eta}^{s} P_{\tau}^{\rho} K_{\rho}(y) d\tau \leq c_{11} \mu(B_{\rho}(x))^{-1} \phi(d(x,y))^{-1} \int_{\eta}^{s} \left(P_{\tau}^{\rho} \mathbb{1}_{B_{c\rho}(y)}(y) + \sum_{k=2}^{\infty} k^{2d} 2^{\frac{ck}{6}} 4^{-\frac{ck}{6}} \right) d\tau \\
\leq c_{12} \frac{s - \eta}{\mu(B_{d(x,y)}(x)) \phi(d(x,y))}$$
(4.10)

for $c_{11}, c_{12} > 0$. Combining (4.8), (4.9), (4.10) we obtain the desired off-diagonal estimate for $s - \eta \leq \frac{1}{4\nu}\phi(\rho)$. Together with the on-diagonal estimate (4.6), we deduce the desired result. \Box

Remark. Note that the proof of Theorem 4.1 does not require the scaling argument from [CK08] since we are working with an on-diagonal estimate and an $L^{\infty} - L^2$ -estimate that take into account the parabolic scaling of the corresponding equation, see also [CKKW21].

A central ingredient in the proof of Theorem 4.1 is the $L^{\infty} - L^2$ -estimate (4.4) for subsolutions to $\partial_t u - L_t^{\rho} u = 0$. Its proof is similar to the proof of Lemma 2.4. However we will provide some details since this seems to be the first time that (FK) is used for nonlocal parabolic $L^{\infty} - L^2$ estimates. Moreover, we provide an $L^{\infty} - L^1$ -estimate for subsolutions to $\partial_t u - L_t u = 0$ which allows us to give a direct proof of the on-diagonal upper heat kernel estimate. In the elliptic case, $L^{\infty} - L^2$ - and $L^{\infty} - L^1$ -estimates are established via (FK) for example in [CKW21].

Lemma 4.2. Let (M, d, μ) and k be as in Theorem 4.1. Then there exists $C_1 > 0$ such that for every $t_0 \in (0,T)$, $x_0 \in M$, $\rho, R > 0$ with $R \leq \rho/2 \wedge \phi^{-1}(t_0)$ and every subsolution u to $\partial_t u - L_t^{\rho} u = 0$ in $I_R(t_0) \times B_R(x_0)$ it holds:

$$\sup_{I_{R/2}(t_0) \times B_{R/2}(x_0)} u \le C_1 \left(\frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x_0))}\right)^{\frac{1}{2}} \mu(B_R(x_0))^{-\frac{1}{2}} \sup_{t \in I_R(t_0)} \left(\int_{B_{2\rho}(x_0)} u^2(t, x) \mathrm{d}x\right)^{1/2}.$$
 (4.11)

Moreover, there exists $C_2 > 0$ such that for every $t_0 \in (0,T)$, $x_0 \in M$, $R \leq \phi^{-1}(t_0)$ and every subsolution u to $\partial_t u - L_t u = 0$ in $I_R(t_0) \times B_R(x_0)$, with $u \geq 0$ in $I_R(t_0) \times B_R(x_0)$, it holds:

$$\sup_{I_{R/2}(t_0) \times B_{R/2}(x_0)} u \le C_2 \mu (B_R(x_0))^{-1} \sup_{t \in I_R(t_0)} \int_M |u(t,x)| \mathrm{d}x.$$
(4.12)

We refer to Section 5 for the definition of a subsolution. $H^{\alpha/2}(\mathbb{R}^d)$ should be replaced by \mathcal{F} .

Proof. First, we prove (4.11). Let l > k > 0, $0 < r \le R \le \rho/2$, $A(l,R) := |\{(t,x) \in I_R \times B_R(x_0) : u(t,x) > l\}|$. Let u be a subsolution to $\partial_t u - L_t^{\rho} u = 0$. First, observe that for every $t \in I_{R+r}$:

$$\sup_{x \in B_R(x_0)} \int_{B_{\rho}(x) \setminus B_{R+\frac{r}{2}}(x_0)} |u(t,y)| k(t;x,y) \mathrm{d}y \le c_1 \sigma'(R,r) \oint_{B_{2\rho}(x_0)} |u(t,y)| \mathrm{d}y,$$

where $c_1 > 0$, $\sigma'(R, r) := \sup_{x \in B_R(x_0)} \phi(r)^{-1} \frac{\mu(B_{\rho}(x_0))}{\mu(B_r(x))}$ and we used (k_{\leq}) . By Caccioppoli's inequality and (4.3), every subsolution u to $\partial_t u - L_t^{\rho} u = 0$ in $I_R \times B_R(x_0)$ satisfies:

$$\sup_{t \in I_R} \left| A\left(l, R + \frac{r}{2}\right) \right| \le \frac{1}{2} (l-k)^{-2} \sup_{t \in I_R} \int_{B_{R+\frac{r}{2}}(x_0)} w_{\frac{k+l}{2}}^2(t, x) \mathrm{d}x$$

$$\leq c_2(l-k)^{-2} \left(\sigma(R,r) + \frac{\sigma'(R,r)}{l-k} \sup_{t \in I_{R+r}} \oint_{B_{2\rho}(x_0)} |u(t,x)| \mathrm{d}x \right) \int_{I_{R+r}} \int_{B_{R+r}(x_0)} w_k^2(t,x) \mathrm{d}x \mathrm{d}t,$$

where $c_2 > 0$, $\sigma(R, r) = \phi(r)^{-1} \vee (\phi(R+r) - \phi(R))^{-1}$ and

$$\begin{split} \phi \left(R + \frac{r}{2}\right)^{-1} \int_{I_R} \left(\frac{\mu(B_{R+\frac{r}{2}}(x_0))}{A(l,R+\frac{r}{2})}\right)^{\nu} \int_{B_R(x_0)} w_l^2(t,x) \mathrm{d}x \mathrm{d}t \le c_3 \int_{I_R} \mathcal{E}_t(\tau w_l(t),\tau w_l(t)) \mathrm{d}t \\ \le c_4 \left(\sigma(R,r) + \frac{\sigma'(R,r)}{l-k} \sup_{t \in I_{R+r}} \int_{B_{2\rho}(x_0)} |u(t,x)| \mathrm{d}x\right) \int_{I_{R+r}} \int_{B_{R+r}(x_0)} w_k^2(t,x) \mathrm{d}x \mathrm{d}t, \end{split}$$

where $c_3, c_4 > 0, \tau \in C_c^{\infty}(\mathbb{R}^d)$ is an arbitrary function such that $\tau \equiv 1$ in $B_R(x_0), \tau \equiv 0$ in $B_{R+\frac{r}{2}}(x_0), \|\nabla \tau\|_{\infty} \leq 4r^{-1}$, and we used (FK). By combination of the foregoing two estimates, we obtain for some $c_5 > 0$:

$$\int_{I_R} \int_{B_R(x_0)} w_l^2(t, x) dx dt \le c_5 \left(\int_{I_{R+r}} \int_{B_{R+r}(x_0)} w_k^2(t, x) dx dt \right)^{1+\nu} \times \frac{\phi(R + \frac{r}{2})}{\mu(B_{R+\frac{r}{2}}(x_0))^{\nu}} (l-k)^{-2\nu} \left(\sigma(R, r) + \frac{\sigma'(R, r)}{l-k} \sup_{t \in I_{R+r}} \int_{B_{2\rho}(x_0)} |u(t, x)| dx \right)^{1+\nu}.$$

$$(4.13)$$

Let us now fix $R \in (0, \rho/2]$ and define sequences $l_i = M(1 - 2^{-i})$, for M > 0 to be defined later, $r_i = 2^{-i-1}R$, $R_{i+1} = R_i - r_{i+1}$, $R_0 := R$, $A_i = \int_{I_{R_i}} \int_{B_{R_i}(x_0)} w_{l_i}^2(t, x) dx dt$. We deduce:

$$A_{i} \leq c_{6}(l_{i}-l_{i-1})^{-2\nu} \frac{\phi(R_{i})}{\mu(B_{R_{i-1}}(x_{0}))^{\nu}} \left(\sigma(R_{i},r_{i}) + \frac{\sigma'(R_{i},r_{i})}{l_{i}-l_{i-1}} \sup_{t \in I_{R_{i-1}}} \oint_{B_{2\rho}(x_{0})} |u(t,x)| \mathrm{d}x\right)^{1+\nu} A_{i-1}^{1+\nu}$$
$$\leq c_{7} 2^{\gamma i} M^{-2\nu} (\phi(R)\mu(B_{R}(x_{0})))^{-\nu} \left(1 + M^{-1} \frac{\mu(B_{\rho}(x_{0}))}{\mu(B_{R}(x_{0}))} \sup_{t \in I_{R}} \oint_{B_{2\rho}(x_{0})} |u(t,x)| \mathrm{d}x\right)^{1+\nu} A_{i-1}^{1+\nu}$$

for some $\gamma, c_6, c_7 > 0$, using that $\sigma(R_i, r_i) \leq c_8 2^{\gamma_1 i} \phi(R)^{-1}, \ \sigma'(R_i, r_i) \leq c_9 2^{\gamma_2 i} \phi(R)^{-1} \frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x_0))}$ for some $c_8, c_9, \gamma_1, \gamma_2 > 0$. The latter follows from the fact that for all $x \in B_R(x_0)$: $\frac{\mu(B_{\rho}(x_0))}{\mu(B_{r_i}(x))} = \frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x))} \frac{\mu(B_R(x))}{\mu(B_{r_i}(x))} \leq c_{10} 2^{\gamma_3 i} \frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x_0))}, \text{ where } c_{10}, \gamma_3 > 0.$ Let us choose $c_{11} = c_7 2^{1+\gamma}$, and

$$M := \frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x_0))} \sup_{t \in I_R} \oint_{B_{2\rho}(x_0)} |u(t,x)| \mathrm{d}x + 2^{-\frac{\gamma}{2\nu^2}} c_{11}^{\frac{1}{2\nu}} (\phi(R)\mu(B_R(x_0)))^{-\frac{1}{2}} A_0^{1/2} d_0^{1/2} d$$

Hence:

$$A_{i} \leq (c_{11}M^{-2\nu}(\phi(R)\mu(B_{R}(x_{0})))^{-\nu})2^{\gamma i}A_{i-1}^{1+\nu}, \quad A_{0} \leq 2^{-\frac{\gamma}{\nu^{2}}}(c_{11}M^{-2\nu}(\phi(R)\mu(B_{R}(x_{0})))^{-\nu})^{-\frac{1}{\nu}},$$

and we can apply Lemma 7.1 in [Giu03] to deduce that for some $c_{12} > 0$:

$$\sup_{I_{R/2} \times B_{R/2}(x_0)} u \leq \frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x_0))} \sup_{t \in I_R} \oint_{B_{2\rho}(x_0)} |u(t,x)| dx + c_7 \left(\oint_{I_R} \oint_{B_R(x_0)} u^2(t,x) dx dt \right)^{1/2}$$
$$\leq c_{12} \left(\frac{\mu(B_{\rho}(x_0))}{\mu(B_R(x_0))} \right)^{\frac{1}{2}} (\mu(B_R(x_0))^{-\frac{1}{2}} \sup_{t \in I_R} \left(\int_{B_{2\rho}(x_0)} u^2(t,x) dx \right)^{1/2}.$$

This proves (4.11). Let us now demonstrate how to prove (4.12). Let u be a subsolution to $\partial_t u - L_t u = 0$. First, we provide a different estimate of the tail term. For every $t \in I_{R+r}$:

$$\sup_{x \in B_R(x_0)} \int_{M \setminus B_{R+\frac{r}{2}}(x_0)} |u(t,y)| k(t;x,y) \mathrm{d}y \le c_{13} \widetilde{\sigma}'(R,r) \int_M |u(t,y)| \mathrm{d}y,$$

where $\tilde{\sigma}'(R, r) := \sup_{x \in B_R(x_0)} \mu(B_r(x))^{-1} \phi(r)^{-1}$ and we applied (k_{\leq}) . As in (4.13), we get:

$$\int_{I_R} \int_{B_R(x_0)} w_l^2(t, x) \mathrm{d}x \mathrm{d}t \le c_{14} \left(\int_{I_{R+r}} \int_{B_{R+r}(x_0)} w_k^2(t, x) \mathrm{d}x \mathrm{d}t \right)^{1+\nu} \times \frac{\phi(R + \frac{r}{2})}{\mu(B_{R+\frac{r}{2}}(x_0))^{\nu}} (l-k)^{-2\nu} \left(\sigma(R, r) + \frac{\widetilde{\sigma}'(R, r)}{l-k} \sup_{t \in I_{R+r}} \int_M |u(t, x)| \mathrm{d}x \right)^{1+\nu}$$

for some $c_{14} > 0$. From now on, let $\overline{R} > 0$ be fixed. Moreover, let $0 < \overline{R}/2 \le r < R \le \overline{R}$ and define sequences $l_i = M(1 - 2^{-i})$, for M > 0 to be defined later, $r_i = 2^{-i-1}(R - r)$, $R_{i+1} = R_i - r_{i+1}, R_0 := R, A_i = \int_{I_{R_i}} \int_{B_{R_i}(x_0)} w_{l_i}^2(t, x) dx dt$ and deduce

$$A_{i} \leq c_{16} \frac{2^{\gamma i}}{M^{2\nu}} \frac{\phi(R)}{\mu(B_{R}(x_{0}))^{\nu}} \left(\frac{R}{R-r} \frac{1}{\phi(R-r)} + \frac{1}{M} \left(\frac{R}{R-r}\right)^{d} \frac{\phi(R-r)^{-1}}{\mu(B_{R}(x_{0}))} \sup_{t \in I_{R}} \int_{M} |u(t,x)| \mathrm{d}x\right)^{1+\nu} A_{i-1}^{1+\nu}$$

$$\leq c_{17} \frac{2^{\gamma i}}{M^{2\nu}} \frac{\left(\frac{R}{R-r}\right)^{\alpha_{2}+1}}{(\phi(R-r)\mu(B_{R}(x_{0})))^{\nu}} \left(1 + \frac{1}{M} \frac{\left(\frac{R}{R-r}\right)^{d-1}}{\mu(B_{R}(x_{0}))} \sup_{t \in I_{R}} \int_{M} |u(t,x)| \mathrm{d}x\right)^{1+\nu} A_{i-1}^{1+\nu},$$

for $c_{16}, c_{17}, \gamma > 0$, using (4.1) and that by (VD): $\mu(B_{R_i}(x_0)) \ge \mu(B_{\overline{R}/2}(x_0)) \ge c_{18}\mu(B_{\overline{R}}(x_0)) \ge c_{18}\mu(B_{\overline{R}}(x_0)) \ge c_{18}\mu(B_{\overline{R}}(x_0)), \sigma(R_i, r_i) \le c_{19}2^{\gamma_4 i}\frac{R}{R-r}\phi(R-r)^{-1}, \widetilde{\sigma}'(R_i, r_i) \le c_{20}2^{\gamma_5 i}\phi(R-r)^{-1}\left(\frac{R}{R-r}\right)^d\mu(B_R(x_0))^{-1}.$ The latter follows from the fact that for all $x \in B_R(x_0)$: $\mu(B_{r_i}(x))^{-1} = \frac{\mu(B_R)(x)}{\mu(B_{r_i}(x))}\mu(B_R(x))^{-1} \le c_{21}2^{\gamma_6 i}\left(\frac{R}{R-r}\right)^d\mu(B_R(x_0))^{-1}.$ We choose $c_{22} = c_{17}2^{1+\gamma}$ and

$$M := \frac{\left(\frac{R}{R-r}\right)^{d-1}}{\mu(B_R(x_0))} \sup_{t \in I_R} \int_M |u(t,x)| \mathrm{d}x + 2^{-\frac{\gamma}{2\nu^2}} c_{22}^{\frac{1}{2\nu}} \left[\frac{\left(\frac{R}{R-r}\right)^{\alpha_2+1}}{(\phi(R-r)\mu(B_R(x_0)))^{\nu}} \right]^{\frac{1}{2\nu}} A_0^{1/2}$$

and deduce by arguments analogous to those in the first part of the proof:

$$\sup_{I_r \times B_r(x_0)} u \leq \left(\frac{R}{R-r}\right)^{d-1} \frac{1}{\mu(B_R(x_0))} \sup_{t \in I_R} \int_M |u(t,x)| dx + c_{23} \left[\frac{\left(\frac{R}{R-r}\right)^{\alpha_2+1}}{(\phi(R-r)\mu(B_R(x_0)))^{\nu}}\right]^{\frac{1}{2\nu}} \left(\int_{I_R} \int_{B_R(x_0)} u^2(t,x) dx dt\right)^{1/2} = I_1 + I_2,$$

where $c_{23} > 0$. We further estimate

$$I_2 \le \left(\frac{R}{R-r}\right)^{\frac{\alpha_2(1+\nu)+1}{2\nu}} \sup_{t \in I_R} \left(\oint_{B_R(x_0)} u^2(t,x) \mathrm{d}x \right)^{1/2}$$

$$\leq \frac{1}{2} \sup_{I_R \times B_R(x_0)} u + c_{24} \left(\frac{R}{R-r}\right)^{\frac{\alpha_2(1+\nu)+1}{\nu}} \sup_{t \in I_R} \oint_{B_R(x_0)} |u(t,x)| \mathrm{d}x$$

where $c_{24} > 0$ and we applied (4.1) and Hölder's and Young's inequality. Together, we obtain

$$\sup_{I_r \times B_r(x_0)} u \le \frac{1}{2} \sup_{I_R \times B_R(x_0)} u + c_{25} \left(\frac{R}{R-r}\right)^{\delta} \frac{1}{\mu(B_R(x_0))} \sup_{t \in I_R} \int_M |u(t,x)| \mathrm{d}x$$

for $c_{25} > 0$ and $\delta := d - 1 \lor \frac{\alpha_2(1+\nu)+1}{\nu}$. We can apply Lemma 1.1 in [GG82] to the estimate above and deduce that there exists $c_{26} > 0$ such that for every $0 < \overline{R}/2 \le r < R \le \overline{R}$:

$$\sup_{I_r \times B_r(x_0)} u \le c_{26} \left(\frac{\overline{R}}{R-r}\right)^{\delta} \frac{1}{\mu(B_{\overline{R}}(x_0))} \sup_{t \in I_{\overline{R}}} \int_M |u(t,x)| \mathrm{d}x.$$
(4.14)

Choosing $r = \overline{R}/2, R = \overline{R}$ implies the desired result (4.12).

5. Appendix

The main goal of this section is to give a proof of Lemma 2.2 via analysis methods. We mainly follow the strategy carried out in [GH08] modifying some of their arguments due to the time-inhomogeneity of the jumping kernel.

First, we introduce the notion of a subsolution to $\partial_t u - L_t u = 0$ in $I \times \Omega$ for some open interval $I \subset (\eta, T)$ and some open set $\Omega \subset \mathbb{R}^d$.

Definition 5.1. Let $\Omega \subset \mathbb{R}^d$ be open and bounded. We say that a function $u \in L^2_{loc}(I; H^{\alpha/2}(\mathbb{R}^d))$ with $\partial_t u \in L^1_{loc}(I; L^2_{loc}(\Omega))$ is a subsolution to

$$\partial_t u - L_t u = 0, \quad in \ I \times \Omega, \tag{5.1}$$

if for every $\phi \in H^{\alpha/2}(\mathbb{R}^d)$ with $\phi \equiv 0$ in $\mathbb{R}^d \setminus \Omega$ and $\phi \geq 0$:

$$\int_{\mathbb{R}^d} \partial_t u(t, x) \phi(x) \mathrm{d}x + \mathcal{E}_t(u(t), \phi) \le 0, \quad a.e. \ t \in I.$$
(5.2)

In this case we say that u solves $\partial_t u - L_t u \leq 0$ in $I \times \Omega$. u is a solution to (5.1) if (5.2) holds for any $\phi \in H^{\alpha/2}(\mathbb{R}^d)$ with $\phi \equiv 0$ in $\mathbb{R}^d \setminus \Omega$. (Sub-)solutions to the corresponding ρ -truncated problem are defined accordingly, replacing k by k_{ρ} .

The main ingredient in the proof of Lemma 2.2 is the following parabolic maximum principle, which is an analog of Proposition 4.11 in [GH08]:

Lemma 5.2 (parabolic maximum principle). Let $\Omega \subset \mathbb{R}^d$ be open. Assume that k satisfies $(k_{\leq}), (\mathcal{E}_{\geq})$. Assume that u solves

$$\begin{cases} \partial_t u - L_t u \leq 0, \ in \ (\eta, T) \times \Omega, \\ u_+(t) \in H_{\Omega}^{\alpha/2}(\mathbb{R}^d), \quad \forall t \in (\eta, T), \\ u_+(t) \to 0, \ in \ L^2(\Omega), \ as \ t \searrow \eta. \end{cases}$$
(5.3)

Then $u \leq 0$ a.e. in $(\eta, T) \times \Omega$. The same result holds for subsolutions to $\partial_t u - L_t^{\rho} u = 0$.

Proof. By assumptions (k_{\leq}) , (\mathcal{E}_{\geq}) and dominated convergence theorem, we have for $f \in L^2(\mathbb{R}^d)$ $\lambda[f]^2_{H^{\alpha/2}(\mathbb{R}^d)} \leq \mathcal{E}_t(f, f) \leq \Lambda[f]^2_{H^{\alpha/2}(\mathbb{R}^d)}, \quad \forall t \in (0, T).$

Thus, $(\mathcal{E}_t, H^{\alpha/2}(\mathbb{R}^d))$ is a regular Dirichlet form for every $t \in (0, T)$. Along the lines of Lemma 4.3 in [GH08] one can prove

$$\int_{\mathbb{R}^d} \partial_t u(t,x) \phi(u(t,x)) \mathrm{d}x \le 0, \quad \text{a.e. } t \in (\eta,T)$$

where $\phi \in C^{\infty}(\mathbb{R})$ such that $\phi \equiv 0$ in $(-\infty, 0]$, $\phi > 0$ on $(0, \infty)$, and $0 \le \phi' \le 1$. From here, the remainder of the proof follows along the lines of Proposition 4.11 in [GH08]. The proof for subsolutions to $\partial_t u - L_t^{\rho} u = 0$ is carried out via similar arguments, using that for some c > 0:

$$[f]_{H^{\alpha/2}(\mathbb{R}^d)}^2 \le c\rho^{-\alpha} \|f\|_{L^2(\mathbb{R}^d)}^2 + \mathcal{E}_t^{\rho}(f, f), \quad t \in (0, T).$$
(5.4)

In the following, for $f \in L^2(\Omega)$, we will denote by $(s, x) \mapsto P_{\eta,s}^{\Omega} f(x)$ the solution to $\partial_t u - L_t u = 0$ in $(\eta, T) \times \Omega$ with $\|P_{\eta,s}^{\Omega} f - f\|_{L^2(\Omega)} \to 0$ as $s \searrow \eta$, and $P_{\eta,s}^{\Omega} f \equiv 0$ in $\mathbb{R}^d \setminus \Omega$. We define $(s, x) \mapsto P_{\eta,s}^{\Omega,\rho} f(x)$ to be solution to the corresponding ρ -truncated problem.

In order to prove Lemma 2.2, we need an approximation result for $P_{\eta,s}f$. In the time-homogeneous case, its proof is given in Lemma 4.17 in [GH08]. However, their argument does not work in our situation due to the lack of a resolvent operator associated with $P_{\eta,s}$.

Lemma 5.3. Let $(\Omega_n)_{n\in\mathbb{N}}$ be an increasing sequence of open subsets of \mathbb{R}^d with $\bigcup_{n\in\mathbb{N}} \Omega_n = \mathbb{R}^d$, and $\eta \geq 0$. Assume that k satisfies (k_{\leq}) , (\mathcal{E}_{\geq}) . Then for every $f \in L^2(\mathbb{R}^d)$ it holds $P_{\eta,s}^{\Omega_n} f \to P_{\eta,s} f$, $P_{\eta,s}^{\Omega_n,\rho} f \to P_{\eta,s}^{\rho} f$ pointwise a.e. for a.e. $s \in (\eta, T)$, and every $\rho > 0$.

Proof. We denote $u_n(s) := P_{\eta,s}^{\Omega_n} f$. By the parabolic maximum principle it follows that $u_n(s) \leq u_{n+1}(s) \leq \cdots \leq P_{\eta,s} f$. Consequently, there exists a function $u \in L^2((\eta, T) \times \mathbb{R}^d)$ such that $u_n(s) \nearrow u(s) \leq P_{\eta,s} f$ a.e., and by dominated convergence we have $||u_n - u||_{L^2((\eta, T) \times \mathbb{R}^d)} \to 0$. It remains to prove $u(s) = P_{\eta,s} f$. Note that by (5.2) it is sufficient to prove that $u_n \to u$ in $L^2((\eta, T); H^{\alpha/2}(\mathbb{R}^d))$. Let us test the equation for u_n with $\phi = u_n$, integrate over (η, T) . Then:

$$\int_{\mathbb{R}^d} u_n^2(T, x) \mathrm{d}x + 2 \int_{\eta}^T \mathcal{E}_t(u_n(t), u_n(t)) \mathrm{d}t \le \int_{\mathbb{R}^d} f^2(x) \mathrm{d}x$$

By uniform boundedness of $||u_n||_{L^2((\eta,T)\times\mathbb{R}^d)}$, we conclude that also $||u_n||_{L^2((\eta,T);H^{\alpha/2}(\mathbb{R}^d))}$ is uniformly bounded, which implies $u_n \rightharpoonup u$ in $L^2((\eta,T);H^{\alpha/2}(\mathbb{R}^d))$, as desired.

The desired result for $P_{\eta,s}^{\Omega_n,\rho}$ follows by the same argument, using (5.4) to conclude boundedness in $L^2((\eta,T); H^{\alpha/2}(\mathbb{R}^d))$.

Proof of Lemma 2.2. Let $f \in L^2(\mathbb{R}^d)$ with $f \ge 0$, and $\Omega_n \subset \mathbb{R}^d$ as in Lemma 5.3. We proceed as in the proof of Proposition 4.4 in [GHL14] to deduce for a.e. $0 \le \eta < s < T$ and $n \in \mathbb{N}$:

$$P_{\eta,s}^{\Omega_n} f \le P_{\eta,s}^{\Omega_n,\rho} f + \int_{\eta}^{s} \left[\int_{\mathbb{R}^d} P_{\eta,\tau}^{\Omega_n,\rho} f(y) K_{\rho}(y) \mathrm{d}y \right] \mathrm{d}\tau,$$
(5.5)

where $K_{\rho}(y) = \sup_{z \in \mathbb{R}^d, t \in (\eta, T)} k(t; z, y) \mathbb{1}_{\{|z-y| \ge \rho\}}(y)$. The proof of (5.5) directly follows by application of the parabolic maximum principle Lemma 5.2 for L_t to

$$u(s,x) = P_{\eta,s}^{\Omega_n} f(x) - P_{\eta,s}^{\Omega_n,\rho} f(x) - \phi_n(x) \int_{\eta}^{s} \left[\int_{\mathbb{R}^d} P_{\eta,\tau}^{\Omega_n,\rho} f(y) K_{\rho}(y) \mathrm{d}y \right] \mathrm{d}\tau,$$

where $\phi_n \in C_c^{\infty}(\mathbb{R}^d)$ with $0 \leq \phi_n \leq 1$ and $\phi_n \equiv 1$ in Ω_n . Taking $n \to \infty$ in (5.5) implies

$$P_{\eta,s}f \le P_{\eta,s}^{\rho}f + \int_{\eta}^{s} \left[\int_{\mathbb{R}^{d}} P_{\eta,\tau}^{\rho} K_{\rho}(y) f(y) \mathrm{d}y \right] \mathrm{d}\tau,$$

where we used Lemma 5.3 and (2.7). By (2.6) it is easy to deduce for a.e. $0 \le \eta < s < T$ and a.e. $x, y \in \mathbb{R}^d$:

$$p(y,s;x,\eta) \le p_{\rho}(y,s;x,\eta) + \int_{\eta}^{s} P_{\eta,\tau}^{\rho} K_{\rho}(y) \mathrm{d}\tau.$$
(5.6)

(2.11) is now a direct consequence of (2.8) and (k_{\leq}) . The proof of (2.12) follows by carrying out the same arguments as in the proof of Proposition 4.4 in [GHL14] with

$$u(s,x) = P_{\eta,s}^{\Omega_n,\rho} f(x) - P_{\eta,s}^{\Omega_n} f(x) e^{(s-\eta)K'_{\rho}},$$

where $K'_{\rho} := \sup_{(t,x) \in (0,T) \times \mathbb{R}^d} \int_{\mathbb{R}^d \setminus B_{\rho}(x)} k(t;x,y) dy$. This time, we apply the parabolic maximum principle Lemma 5.2 for L^{ρ}_t and obtain for a.e. $0 \le \eta < s < T$ and a.e. $x, y \in \mathbb{R}^d$:

$$p_{\rho}(y,s;x,\eta) \le p(y,s;x,\eta)e^{(s-\eta)K'_{\rho}}.$$
 (5.7)

Note that by Hölder regularity of weak solutions to (1.3), it is possible to obtain the desired results for every η, s, x, y .

References

- [Aro67] D. G. Aronson. Bounds for the fundamental solution of a parabolic equation. Bull. Amer. Math. Soc., 73:890–896, 1967.
- [Aro68] D. G. Aronson. Non-negative solutions of linear parabolic equations. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3), 22:607–694, 1968.
- [AS67] D. G. Aronson and James Serrin. Local behavior of solutions of quasilinear parabolic equations. Arch. Rational Mech. Anal., 25:81–122, 1967.
- [BBK09] Martin T. Barlow, Richard F. Bass, and Takashi Kumagai. Parabolic Harnack inequality and heat kernel estimates for random walks with long range jumps. *Math. Z.*, 261(2):297–320, 2009.
- [BG60] R. M. Blumenthal and R. K. Getoor. Some theorems on stable processes. Trans. Amer. Math. Soc., 95:263–273, 1960.
- [BGK09] Martin T. Barlow, Alexander Grigor'yan, and Takashi Kumagai. Heat kernel upper bounds for jump processes and the first exit time. J. Reine Angew. Math., 626:135–157, 2009.
- [BKKL19] Joohak Bae, Jaehoon Kang, Panki Kim, and Jaehun Lee. Heat kernel estimates for symmetric jump processes with mixed polynomial growths. *Ann. Probab.*, 47(5):2830–2868, 2019.
- [BL02] Richard F. Bass and David A. Levin. Transition probabilities for symmetric jump processes. *Trans. Amer. Math. Soc.*, 354(7):2933–2953, 2002.
- [CK03] Zhen-Qing Chen and Takashi Kumagai. Heat kernel estimates for stable-like processes on d-sets. Stochastic Process. Appl., 108(1):27–62, 2003.
- [CK08] Zhen-Qing Chen and Takashi Kumagai. Heat kernel estimates for jump processes of mixed types on metric measure spaces. *Probab. Theory Related Fields*, 140(1-2):277–317, 2008.
- [CKKW21] Zhen-Qing Chen, Panki Kim, Takashi Kumagai, and Jian Wang. Heat kernel upper bounds for symmetric Markov semigroups. J. Funct. Anal., 281(4):Paper No. 109074, 40, 2021.
- [CKS87] E. A. Carlen, S. Kusuoka, and D. W. Stroock. Upper bounds for symmetric Markov transition functions. Ann. Inst. H. Poincaré Probab. Statist., 23(2, suppl.):245–287, 1987.
- [CKW21] Zhen-Qing Chen, Takashi Kumagai, and Jian Wang. Stability of heat kernel estimates for symmetric non-local Dirichlet forms. *Mem. Amer. Math. Soc.*, 271(1330):v+89, 2021.
- [CLY81] Siu Yuen Cheng, Peter Li, and Shing Tung Yau. On the upper estimate of the heat kernel of a complete Riemannian manifold. Amer. J. Math., 103(5):1021–1063, 1981.
- [CS20] Jamil Chaker and Luis Silvestre. Coercivity estimates for integro-differential operators. *Calc. Var. Partial Differential Equations*, 59(4):Paper No. 106, 20, 2020.
- [Dav89] E. B. Davies. *Heat kernels and spectral theory*, volume 92 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1989.
- [DCKP14] Agnese Di Castro, Tuomo Kuusi, and Giampiero Palatucci. Nonlocal Harnack inequalities. J. Funct. Anal., 267(6):1807–1836, 2014.
- [DCKP16] Agnese Di Castro, Tuomo Kuusi, and Giampiero Palatucci. Local behavior of fractional pminimizers. Ann. Inst. H. Poincaré Anal. Non Linéaire, 33(5):1279–1299, 2016.
- [GG82] Mariano Giaquinta and Enrico Giusti. On the regularity of the minima of variational integrals. Acta Math., 148:31–46, 1982.
- [GH08] Alexander Grigoryan and Jiaxin Hu. Off-diagonal upper estimates for the heat kernel of the Dirichlet forms on metric spaces. *Invent. Math.*, 174(1):81–126, 2008.

- [GHH17] Alexander Grigor'yan, Eryan Hu, and Jiaxin Hu. Lower estimates of heat kernels for non-local Dirichlet forms on metric measure spaces. J. Funct. Anal., 272(8):3311–3346, 2017.
- [GHH18] Alexander Grigor'yan, Eryan Hu, and Jiaxin Hu. Two-sided estimates of heat kernels of jump type Dirichlet forms. Adv. Math., 330:433–515, 2018.
- [GHL14] Alexander Grigor'yan, Jiaxin Hu, and Ka-Sing Lau. Estimates of heat kernels for non-local regular Dirichlet forms. *Trans. Amer. Math. Soc.*, 366(12):6397–6441, 2014.
- [Giu03] Enrico Giusti. Direct methods in the calculus of variations. World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [Gri09] Alexander Grigor'yan. Heat kernel and analysis on manifolds, volume 47 of AMS/IP Studies in Advanced Mathematics. American Mathematical Society, Providence, RI; International Press, Boston, MA, 2009.
- [HL18] Jiaxin Hu and Xuliang Li. The Davies method revisited for heat kernel upper bounds of regular Dirichlet forms on metric measure spaces. *Forum Math.*, 30(5):1129–1155, 2018.
- [Kig04] Jun Kigami. Local Nash inequality and inhomogeneity of heat kernels. Proc. London Math. Soc. (3), 89(2):525–544, 2004.
- [KKK] Moritz Kassmann, Kyung-Youn Kim, and Takashi Kumagai. Heat kernel bounds for nonlocal operators with singular kernels. ArXiv e-print 1910.04242, to appear in J. Math. Pures Appl.
- [Kom88] Takashi Komatsu. Continuity estimates for solutions of parabolic equations associated with jump type Dirichlet forms. Osaka J. Math., 25(3):697–728, 1988.
- [Kom95] Takashi Komatsu. Uniform estimates for fundamental solutions associated with non-local Dirichlet forms. Osaka J. Math., 32(4):833–860, 1995.
- [Li12] Peter Li. Geometric analysis, volume 134 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2012.
- [LY86] Peter Li and Shing-Tung Yau. On the parabolic kernel of the Schrödinger operator. Acta Math., 156(3-4):153–201, 1986.
- [Mey75] P. A. Meyer. Renaissance, recollements, mélanges, ralentissement de processus de Markov. Ann. Inst. Fourier (Grenoble), 25(3-4):xxiii, 465–497, 1975.
- [MM13a] Yasunori Maekawa and Hideyuki Miura. On fundamental solutions for non-local parabolic equations with divergence free drift. *Adv. Math.*, 247:123–191, 2013.
- [MM13b] Yasunori Maekawa and Hideyuki Miura. Upper bounds for fundamental solutions to non-local diffusion equations with divergence free drift. J. Funct. Anal., 264(10):2245–2268, 2013.
- [Mos64] Jürgen Moser. A Harnack inequality for parabolic differential equations. Comm. Pure Appl. Math., 17:101–134, 1964.
- [Mos67] Jürgen Moser. Correction to: "A Harnack inequality for parabolic differential equations". Comm. Pure Appl. Math., 20:231–236, 1967.
- [Mos71] Jürgen Moser. On a pointwise estimate for parabolic differential equations. Comm. Pure Appl. Math., 24:727–740, 1971.
- [MSC19] Mathav Murugan and Laurent Saloff-Coste. Heat kernel estimates for anomalous heavy-tailed random walks. Ann. Inst. Henri Poincaré Probab. Stat., 55(2):697–719, 2019.
- [Nas58] John Nash. Continuity of solutions of parabolic and elliptic equations. Amer. J. Math., 80:931–954, 1958.
- [PEm84] F. O. Porper and S. D. Eidel' man. Two-sided estimates of the fundamental solutions of secondorder parabolic equations and some applications of them. Uspekhi Mat. Nauk, 39(3(237)):107–156, 1984.
- [Str19] Martin Strömqvist. Harnack's inequality for parabolic nonlocal equations. Ann. Inst. H. Poincaré Anal. Non Linéaire, 36(6):1709–1745, 2019.
- [Vaz18] Juan Luis Vazquez. Asymptotic behaviour for the fractional heat equation in the Euclidean space. Complex Var. Elliptic Equ., 63(7-8):1216–1231, 2018.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, POSTFACH 100131, D-33501 BIELEFELD

Email address: moritz.kassmann@uni-bielefeld.de

 $\mathit{URL}:$ www.math.uni-bielefeld.de/ \sim kassmann

Fakultät für Mathematik, Universität Bielefeld, Postfach 100131, D-33501 Bielefeld

Email address: mweidner@math.uni-bielefeld.de