

Explicit solution of divide-and-conquer dividing by a half recurrences with polynomial independent term

Tomás M. Coronado, Arnau Mir, Francesc Rosselló

Department of Mathematics and Computer Science,
University of the Balearic Islands, E-07122 Palma, Spain
and

Balearic Islands Health Research Institute (IdISBa), E-07010 Palma, Spain

November 25, 2021

Abstract

Divide-and-conquer dividing by a half recurrences, of the form

$$x_n = a \cdot x_{\lceil n/2 \rceil} + a \cdot x_{\lfloor n/2 \rfloor} + p(n), \quad n \geq 2,$$

appear in many areas of applied mathematics, from the analysis of algorithms to the optimization of phylogenetic balance indices. The Master Theorems that solve these equations do not provide the solution’s explicit expression, only its big- Θ order of growth. In this paper we give an explicit expression—in terms of the binary decomposition of n —for the solution x_n of a recurrence of this form, with given initial condition x_1 , when the independent term $p(n)$ is a polynomial in $\lceil n/2 \rceil$ and $\lfloor n/2 \rfloor$.

1 Introduction

Divide-and-conquer dividing by a half recurrences, of the form

$$x_n = a \cdot x_{\lceil n/2 \rceil} + a \cdot x_{\lfloor n/2 \rfloor} + p(n), \quad n \geq 2, \tag{1}$$

appear in many areas of applied mathematics. As evidence, for instance, the *On-line Encyclopedia of Integer Sequences* (OEIS, <https://oeis.org>) contains more than 200 integer sequences with the keyword “divide and conquer”, many of which satisfy a recurrence like (1) [35].

The most popular and best studied setting where such recurrences arise is in Computer Science, and more specifically in the analysis of balanced divide and conquer algorithms. This type of algorithms solve a problem by splitting its input into two or more parts of the same size, solving (recursively) the problem on these parts, and finally combining these solutions into a solution for the global instance [2, §2.6–7]. Typical examples of this strategy are the heapsort and mergesort algorithms and several fast integer and matrix multiplication methods. When the input is split into two parts of the same size (up to a unit of difference, when the size of the original input is odd) and both parts contribute equally to the final solution, the algorithm’s running time t_n on an instance of size n satisfies a recurrence of the form

$$t_n = a \cdot t_{\lceil n/2 \rceil} + a \cdot t_{\lfloor n/2 \rfloor} + p(n)$$

where the independent term $p(n)$, called in this context the *toll function*, represents the cost of combining the solutions of the subproblems into a solution for the original problem.

Our interest in this type of recurrences stems from the study of phylogenetic balance indices. A *phylogenetic tree*, the standard representation of the joint evolutionary history of a group of extant species (or other Operational Phylogenetic Units, OPU, like genes, languages, or myths),

is, from the formal point of view, a leaf-labeled rooted tree. In a phylogenetic tree, its leaves represent the species under study, its internal nodes represent their common ancestors, the root represents the most recent common ancestor of all of them, and the arcs represent direct descent through mutations [12, 32]. A phylogenetic tree is *bifurcating*, or *fully resolved*, when every internal node has two direct descendants, or *children*.

Biologists use the *shape* of phylogenetic trees, that is, their raw branching structure, to deduce information on the forces beneath the speciation and extinction processes that have taken place [18]. A popular set of tools used in the analysis of phylogenetic tree shapes are the *shape indices* [10], and among them the *balance indices*, which measure the propensity of the direct descendants of any given node in a tree to have the same number of descendant leaves. Many balance indices have been proposed in the literature: see, for instance, [6, 12, 16, 17, 24, 25, 26, 27, 31, 33] and the recent survey [15] and the references therein. Let us recall here the four indices that appear below in the Examples section:

- The *Sackin index* $S(T)$ of a tree T [31, 33] is the sum of its leaves' *depths* (that is, of the lengths of the paths from the root to the leaves).
- The *Colless index* $C(T)$ of a bifurcating tree T [6] is the sum, over all its internal nodes v , of the absolute value of the difference between the numbers of descendant leaves of the pair of children of v .
- The *Total Cophenetic index* $\Phi(T)$ of a tree T [26] is the sum, over all its pairs of internal nodes, of the depth of their lowest common ancestor.
- The *rooted Quartet index* $rQI(T)$ of a bifurcating tree T [9] is the number of its subtrees of 4 leaves that are fully symmetric.

Any sensible balance index should classify, for each number of leaves n , as most unbalanced trees the *rooted caterpillars* with n leaves K_n —the bifurcating trees such that all their internal nodes have a leaf child—and as most balanced bifurcating trees the *maximally balanced trees* with n leaves B_n —where, for each internal node, the numbers of descendant leaves of its pair of children differ at most in 1—(cf. Fig. 1). The four indices described above satisfy this condition [15].




Figure 1: A rooted caterpillar with 7 leaves (left) and a maximally balanced with 7 leaves (right).

Now, given a balance index I , it is unreasonable to expect it to allow for the comparison of the balance of trees with different numbers of leaves. A possible way to circumvent this difficulty is to normalize it to $[0, 1]$ for every number of leaves n by subtracting its minimum value for this number of leaves and dividing by its range of values on the space of trees with n leaves [33, 34]. To perform this normalization for bifurcating trees, it is necessary to know the value of I for the rooted caterpillars K_n and the maximally balanced trees B_n . It turns out that, while computing $I(K_n)$ is usually easy, to compute $I(B_n)$ one is led in most cases to solve a recurrence of the form

$$I(B_n) = I(B_{\lceil n/2 \rceil}) + I(B_{\lfloor n/2 \rfloor}) + p_I(n)$$

with $p_I(n)$ a function that, in some sense, measures the contribution of the root to the value of I . For the four aforementioned indices, this independent term is a polynomial in $\lceil n/2 \rceil$ and $\lfloor n/2 \rfloor$: for the Sackin index, $p_S(n) = n = \lceil n/2 \rceil + \lfloor n/2 \rfloor$; for the Colless index, $p_C(n) = \lceil n/2 \rceil - \lfloor n/2 \rfloor$;

for the Total Cophenetic index, $p_{\Phi}(n) = \binom{n}{2} = \binom{\lceil n/2 \rceil + \lfloor n/2 \rfloor}{2}$; and for the rooted Quartet index, $p_{rQI}(n) = \binom{\lceil n/2 \rceil}{2} \binom{\lfloor n/2 \rfloor}{2}$ [15]. At the moment of writing this paper, the values of $S(B_n)$ and $C(B_n)$ were known (see Examples 2 and 3 in Section 3), but not those of $\Phi(B_n)$ or $rQI(B_n)$.

In the context of the analysis of algorithms, divide-and-conquer recurrences like (1) and more general ones are “solved” by finding the big- Θ order of growth of x_n using some *Master Theorem*. The original Master Theorem was obtained by Bentley, Haken, and Saxe [4] and extended in [7, §4.5–6], and since then several improved versions have been obtained by other researchers: see, for instance, [3, 11, 20, 23, 30, 37, 38, 39]. Typically, a Master Theorem deduces the asymptotic behaviour of a solution x_n of (1) from that of the independent term of the recurrence (and, in more general recurrences, from the number of parts into which the input is divided, which we fix here to 2, and the contribution of each subproblem to the general problem, which we assume here to be equal and represented by the coefficient a). Thus, Master Theorems do not provide explicit solutions of recurrences, only their asymptotic behaviour.

In the analysis of algorithms, knowing the growing order of the computational cost of an algorithm on an instance of size n is usually enough. But in other applications, like for instance in order to normalize balance indices as we explained above, an explicit expression for the solution is needed. Many specific divide-and-conquer recurrences are explicitly solved when needed, like for instance the cost of the mergesort algorithm in the worst case, solved as Ex. 34 in [19, Ch. 3], but no general solution is known. To our knowledge, the only attempt to find an explicit solution of Eqn. (1) is made by Hwang, Janson and Tsai [22] by proving that, when $a = 1$ and under very general conditions on the independent term $p(n)$, the solution x_n has the form

$$x_n = nP(\log_2(n)) + F(n) - Q(n)$$

with P continuous and 1-periodic and F, Q of precise growing orders, and giving explicit expressions for P, F, Q in terms of series expansions.

In this paper we consider the more restrictive case when $p(n)$ is a polynomial in $\lceil n/2 \rceil$ and $\lfloor n/2 \rfloor$. In this case, we are able to give an explicit finite formula for x_n in terms of the binary decomposition of n . Although for our applications the case when $a = 1$ was enough, our formula works for the arbitrary a case.

The rest of this paper is organized as follows. In Section 2 we state our main result and in Section 3 we use it to solve several examples. As a proof of concept, in some of them we obtain formulas that were already known, but we also include several examples for which no explicit formula was known so far. Finally, the bulk of the paper (Section 4) is devoted to prove our main result.

We have implemented the formula provided by our main theorem in Python using SymPy, a Python library for symbolic mathematics. This implementation is available at the GitHub repository https://github.com/biocom-uib/divide_and_conquer.

Acknowledgments. This research was partially supported by the Spanish Ministry of Science, Innovation and Universities and the European Regional Development Fund through project PGC2018-096956-B-C43 (FEDER/MICINN/AEI). We thank G. Valiente for some useful suggestions.

2 Main result

For every $n \in \mathbb{N}_{\geq 1}$, let its binary decomposition be $n = \sum_{j=1}^{s_n} 2^{q_j(n)}$, with $0 \leq q_1(n) < \dots < q_{s_n}(n)$, and then, for every $i \geq 1$, let $M_i(n) = \sum_{j=i}^{s_n} 2^{q_j(n)}$. Notice that $M_1(n) = n$ and $M_{s_n} = 2^{q_{s_n}(n)}$. In order to simplify the notations, we set $q_0(n) = 0$ and $M_0(n) = M_1(n) + 2^{q_0(n)} = n + 1$.

Additionally, for every $d, n \in \mathbb{N}$ and $x \in \mathbb{R}_{\neq 0}$ let

$$T(d, n, x) = \sum_{k=0}^{n-1} k^d x^k$$

(with the convention, which we use throughout this paper, that $0^0 = 1$). In particular,

$$T(0, n, x) = \begin{cases} n & \text{if } x = 1 \\ \frac{x^n - 1}{x - 1} & \text{if } x \neq 1 \end{cases} \quad T(1, n, x) = \begin{cases} \binom{n}{2} & \text{if } x = 1 \\ \frac{nx^n(x-1) - x(x^n-1)}{(x-1)^2} & \text{if } x \neq 1 \end{cases}$$

For every $m \in \mathbb{N}$, let B_m denote the m -th Bernoulli number of the first kind [5]. In particular, $B_0 = 1$, $B_1 = -1/2$, and $B_{2k+1} = 0$ for every $k \geq 1$.

For $d = 0, 1$, for every $a \in \mathbb{R}$, for every $m \in \mathbb{N}$, and for every $n \in \mathbb{N}_{\geq 1}$, let

$$\begin{aligned} \alpha_n^{(d,m)}(a) &= \frac{1}{2(m+1)} \sum_{i=1}^{s_n} \sum_{j=0}^m \binom{m+1}{j} B_j 2^j M_i(n)^{m+1-j} (T(d, q_i(n), a2^{j-m}) - T(d, q_{i-1}(n), a2^{j-m})) \\ &\quad + \sum_{i=1}^{s_n-1} q_i(n)^d (a2^{-m})^{q_i(n)} (n - M_i(n)) M_{i+1}(n)^m - T(d, q_{s_n}(n), 2a) \cdot \delta_{m=0} \end{aligned}$$

with $\delta_{m=0} = 1$ if $m = 0$ and $\delta_{m=0} = 0$ if $m > 0$.

For every $a \in \mathbb{R}_{>0}$ and for every $r, t \in \mathbb{N}$, let $\ell_{a,t} = \log_2(a) - t$ and

$$\delta_\ell(a, r, t) = \begin{cases} 1 & \text{if } r > 0, \text{ and } \ell_{a,t} \in \{0, \dots, r-1\} \\ 0 & \text{otherwise} \end{cases}$$

If $a \leq 0$, then $\ell_{a,t}$ is undefined and $\delta_\ell(a, r, t) = 0$.

The main result in this paper is the following:

Theorem. *Let $a \in \mathbb{R}_{\neq 0}$ and let $P(x, y) = \sum_{r,t \geq 0} b_{r,t} x^r y^t \in \mathbb{R}[x, y]$ be a bivariate polynomial. Then, the solution of the recurrence equation*

$$x_n = a \cdot x_{\lceil n/2 \rceil} + a \cdot x_{\lfloor n/2 \rfloor} + P(\lceil n/2 \rceil, \lfloor n/2 \rfloor), \quad n \geq 2,$$

with initial condition x_1 is

$$x_n = \sum_{r,t} b_{r,t} x_n^{(r,t)}(a) + ((2a)^{q_{s_n}(n)} + (2a-1)(na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)}))x_1$$

with, for every $n \geq 2$,

$$\begin{aligned} x_n^{(r,t)}(a) &= \sum_{k=1}^{r+t} \left(\sum_{\substack{i=k \\ i \neq t + \ell_{a,t} + 1}}^{r+t} \frac{\binom{r}{i-t-1} \binom{i}{k} B_{i-k}}{i(2^{i-1} - a)} \right) n^k + \frac{1}{a-1} \cdot \delta_{r>0, t=0, a \neq 1} \\ &\quad + \left(1 - \sum_{\substack{l=0 \\ l \neq \ell_{a,t}}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} \right) (T(0, q_{s_n}(n), 2a) + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)}) \\ &\quad + \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{\substack{l=i-t+1 \\ l \neq \ell_{a,t}}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} \right) \alpha_n^{(0,i)}(a) \\ &\quad + \frac{\delta_\ell(a, r, t)}{a} \binom{r}{\ell_{a,t}} \left(T(1, q_{s_n}(n), 2a) + (na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)}) q_{s_n}(n) \right. \\ &\quad \left. + \sum_{i=0}^{t+\ell_{a,t}-1} \binom{t+\ell_{a,t}}{i} \alpha_n^{(1,i)}(a) \right) \end{aligned}$$

where $\delta_{r>0, t=0, a \neq 1} = 1$ if $r > 0$, $t = 0$, and $a \neq 1$, and $\delta_{r>0, t=0, a \neq 1} = 0$ otherwise.

In particular:

(a) If $r = 0$,

$$\begin{aligned}
x_n^{(0,t)}(a) &= \sum_{i=0}^{t-1} 2^{-i} \binom{t}{i} \alpha_n^{(0,i)}(a) + T(0, q_{s_n}(n), 2a) + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} \\
&= \begin{cases} \sum_{i=0}^{t-1} 2^{-i} \binom{t}{i} \alpha_n^{(0,i)}(1/2) + q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1 & \text{if } a = \frac{1}{2} \\ \sum_{i=0}^{t-1} 2^{-i} \binom{t}{i} \alpha_n^{(0,i)}(a) + \frac{(2a)^{q_{s_n}(n)} - 1}{2a - 1} + n \cdot a^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} & \text{if } a \neq \frac{1}{2} \end{cases}
\end{aligned}$$

(b) If $r \geq 1$ and $t = 0$:

(b.1) If $a = 1/2$,

$$\begin{aligned}
x_n^{(r,0)}(1/2) &= 2 \sum_{k=1}^r \left(\sum_{j=k}^r \frac{\binom{r}{j-1} \binom{j}{k} B_{j-k}}{j(2^j - 1)} \right) n^k - 2 \\
&\quad + \left(1 - 2 \sum_{l=0}^{r-1} \frac{\binom{r}{l}}{2^{l+1} - 1} \right) (q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1) \\
&\quad - \sum_{i=0}^{r-1} \left(2^{-i} \binom{r}{i} + 2 \sum_{l=i+1}^{r-1} \frac{\binom{r}{l} \binom{l}{i}}{2^{l+1} - 1} \right) \alpha_n^{(0,i)}(1/2)
\end{aligned}$$

(b.2) If $a = 1$,

$$\begin{aligned}
x_n^{(r,0)}(1) &= \sum_{k=2}^r \left(\sum_{j=k}^r \frac{\binom{r}{j-1} \binom{j}{k} B_{j-k}}{j(2^{j-1} - 1)} \right) n^k + \left(q_{s_n}(n) + 1 + \sum_{j=1}^{r-1} \frac{\binom{r}{j} (B_j - 1)}{2^j - 1} \right) n \\
&\quad + 1 + \sum_{j=1}^{r-1} \frac{\binom{r}{j}}{2^j - 1} - 2^{q_{s_n}(n)+1} - \sum_{i=0}^{r-1} \left(2^{-i} \binom{r}{i} + \sum_{l=i+1}^{r-1} \frac{\binom{r}{l} \binom{l}{i}}{2^l - 1} \right) \alpha_n^{(0,i)}(1)
\end{aligned}$$

(b.3) If $a \notin \{1/2, 1, \dots, 2^{r-1}\}$

$$\begin{aligned}
x_n^{(r,0)}(a) &= \sum_{k=1}^r \left(\sum_{j=k}^r \frac{\binom{r}{j-1} \binom{j}{k} B_{j-k}}{j(2^{j-1} - a)} \right) n^k + \frac{1}{a - 1} \\
&\quad + \left(1 - \sum_{l=0}^{r-1} \frac{\binom{r}{l}}{2^l - a} \right) \left(\frac{(2a)^{q_{s_n}(n)} - 1}{2a - 1} + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} \right) \\
&\quad - \sum_{i=0}^{r-1} \left(2^{-i} \binom{r}{i} + \sum_{l=i+1}^{r-1} \frac{\binom{r}{l} \binom{l}{i}}{2^l - a} \right) \alpha_n^{(0,i)}(a)
\end{aligned}$$

(b.4) If $a = 2^\ell$ with $\ell \in \{1, \dots, r-1\}$,

$$\begin{aligned}
x_n^{(r,0)}(a) &= \sum_{k=1}^r \left(\sum_{\substack{i=k \\ i \neq \ell+1}}^r \frac{\binom{r}{i-1} \binom{i}{k} B_{i-k}}{i(2^{i-1} - a)} \right) n^k + \frac{1}{a-1} \\
&+ \left(1 - \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^l - a} \right) \left(\frac{(2a)^{q_{s_n}(n)} - 1}{2a-1} + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} \right) \\
&- \sum_{i=0}^{r-1} \left(2^{-i} \binom{r}{i} + \sum_{\substack{l=i+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{l}{i}}{2^l - a} \right) \alpha_n^{(0,i)}(a) \\
&+ \frac{1}{a} \binom{r}{\ell} \left(\frac{((2a-1)q_{s_n}(n) - 2a)(2a)^{q_{s_n}(n)} + 2a}{(2a-1)^2} + q_{s_n}(n)a^{q_{s_n}(n)}(n - 2^{q_{s_n}(n)}) \right) \\
&+ \frac{1}{a} \binom{r}{\ell} \sum_{i=0}^{\ell-1} \binom{\ell}{i} \alpha_n^{(1,i)}(a)
\end{aligned}$$

(c) If $r \geq 1$ and $t \geq 1$:

(c.1) If $a = 1/2$,

$$\begin{aligned}
x_n^{(r,t)}(1/2) &= 2 \sum_{k=1}^{r+t} \left(\sum_{j=k}^{r+t} \frac{\binom{r}{j-t-1} \binom{j}{k} B_{j-k}}{j(2^j - 1)} \right) n^k \\
&+ \left(1 - 2 \sum_{l=0}^{r-1} \frac{\binom{r}{l}}{2^{t+l+1} - 1} \right) \left(q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1 \right) \\
&+ \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - 2 \sum_{l=i-t+1}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l+1} - 1} \right) \alpha_n^{(0,i)}(1/2)
\end{aligned}$$

(c.2) If $a = 1$,

$$\begin{aligned}
x_n^{(r,t)}(1) &= \sum_{k=2}^{r+t} \left(\sum_{j=k}^{r+t} \frac{\binom{r}{j-t-1} \binom{j}{k} B_{j-k}}{j(2^{j-1} - 1)} \right) n^k \\
&+ \left(1 + \sum_{j=1}^{r+t-1} \frac{\binom{r}{j-t} (B_j - 1)}{2^j - 1} \right) n + \sum_{l=0}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - 1} - 1 \\
&+ \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{l=i-t+1}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - 1} \right) \alpha_n^{(0,i)}(1)
\end{aligned}$$

(c.3) If $a \notin \{1/2, 1, 2^t, \dots, 2^{r+t-1}\}$,

$$\begin{aligned}
x_n^{(r,t)}(a) &= \sum_{k=1}^{r+t} \left(\sum_{j=k}^{r+t} \frac{\binom{r}{j-t-1} \binom{j}{k} B_{j-k}}{j(2^{j-1} - a)} \right) n^k \\
&+ \left(1 - \sum_{l=0}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} \right) \left(\frac{(2a)^{q_{s_n}(n)} - 1}{2a-1} + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} \right) \\
&+ \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{l=i-t+1}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} \right) \alpha_n^{(0,i)}(a)
\end{aligned}$$

(c.4) If $a = 2^{t+\ell}$, for some $\ell \in \{0, \dots, r-1\}$,

$$\begin{aligned}
x_n^{(r,t)}(a) &= \sum_{k=1}^{r+t} \left(\sum_{\substack{j=k \\ j \neq t+\ell+1}}^{r+t} \frac{\binom{r}{j-t-1} \binom{j}{k} B_{j-k}}{j(2^{j-1} - a)} \right) n^k \\
&+ \frac{1}{a} \binom{r}{\ell} \left(\frac{q_{s_n}(n)(2a)^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)+1} + 2a}{(2a-1)^2} + q_{s_n}(n)a^{q_{s_n}(n)}(n - 2^{q_{s_n}(n)}) \right) \\
&+ \left(1 - \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} \right) \left(\frac{(2a)^{q_{s_n}(n)} - 1}{2a-1} + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} \right) \\
&+ \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} \right) - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} \alpha_n^{(0,i)}(a) \\
&+ \frac{1}{a} \binom{r}{\ell} \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} \alpha_n^{(1,i)}(a)
\end{aligned}$$

3 Examples

In this section we gather several applications of our main theorem. The sequences' identifiers, when available, refer to the OEIS.

Example 1. When the independent term in the divide-and-conquer recurrence is constant,

$$x_n = a \cdot x_{\lceil n/2 \rceil} + a \cdot x_{\lfloor n/2 \rfloor} + c$$

with $c \in \mathbb{R}$, the solution with initial condition x_1 is

$$\begin{aligned}
x_n &= c \cdot x_n^{(0,0)}(a) + ((2a)^{q_{s_n}(n)} + (2a-1)(na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)}))x_1 \\
&= \begin{cases} c(q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1) + x_1 & \text{if } a = \frac{1}{2} \\ ((2a-1)x_1 + c) \left(na^{q_{s_n}(n)} - \frac{2a-2}{2a-1} (2a)^{q_{s_n}(n)} \right) - \frac{c}{2a-1} & \text{if } a \neq \frac{1}{2} \end{cases}
\end{aligned}$$

In particular, if $a = 1$,

$$x_n = (x_1 + c)n - c.$$

Example 2. The minimum total Sackin index S_n of a rooted bifurcating tree with n leaves (the binary entropy function: sequence A003314) satisfies the recurrence

$$S_n = S_{\lceil n/2 \rceil} + S_{\lfloor n/2 \rfloor} + n = S_{\lceil n/2 \rceil} + S_{\lfloor n/2 \rfloor} + \lceil n/2 \rceil + \lfloor n/2 \rfloor$$

with $S_1 = 0$. Then, according to our main theorem,

$$S_n = x_n^{(1,0)}(1) + x_n^{(0,1)}(1)$$

where

$$\begin{aligned}
x_n^{(1,0)}(1) &= (q_{s_n}(n) + 1)n + 1 - 2^{q_{s_n}(n)+1} - \alpha_n^{(0,0)}(1) \\
x_n^{(0,1)}(1) &= \alpha_n^{(0,0)}(1) + n - 1
\end{aligned}$$

and therefore

$$\begin{aligned}
S_n &= (q_{s_n}(n) + 1)n + 1 - 2^{q_{s_n}(n)+1} - \alpha_n^{(0,0)}(1) + \alpha_n^{(0,0)}(1) + n - 1 \\
&= (q_{s_n}(n) + 2)n - 2^{q_{s_n}(n)+1}
\end{aligned}$$

in agreement with previously published formulas: cf. [14, 22].

Example 3. The minimum Colless index c_n of a rooted bifurcating tree with n leaves (sequence A296062) satisfies the recurrence

$$c_n = c_{\lceil n/2 \rceil} + c_{\lfloor n/2 \rfloor} + \left\lceil \frac{n}{2} \right\rceil - \left\lfloor \frac{n}{2} \right\rfloor$$

with $c_1 = 0$. Therefore, according to our main theorem

$$\begin{aligned} c_n &= x_n^{(1,0)}(1) - x_n^{(0,1)}(1) \\ &= (q_{s_n}(n) + 1)n + 1 - 2^{q_{s_n}(n)+1} - \alpha_n^{(0,0)}(1) - (\alpha_n^{(0,0)}(1) + n - 1) \\ &= q_{s_n}(n)n - 2^{q_{s_n}(n)+1} + 2 - 2\alpha_n^{(0,0)}(1) \end{aligned}$$

Now,

$$\begin{aligned} \alpha_n^{(0,0)}(1) &= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(q_i(n) - q_{i-1}(n)) + \sum_{i=1}^{s_n-1} (n - M_i(n)) - 2^{q_{s_n}(n)} + 1 \quad (2) \\ &= \sum_{i=1}^{s_n} 2^{q_i(n)-1} q_i(n) + (s_n - 1)n - \sum_{j=1}^{s_n} \sum_{i=j}^{s_n} 2^{q_i(n)} + 1 \\ &= \sum_{i=1}^{s_n} 2^{q_i(n)-1} q_i(n) + (s_n - 1)n - \sum_{i=1}^{s_n} i 2^{q_i(n)} + 1 \\ &= (s_n - 1)n + \sum_{i=1}^{s_n} 2^{q_i(n)-1} (q_i(n) - 2i) + 1 \quad (3) \end{aligned}$$

and hence, finally,

$$\begin{aligned} c_n &= q_{s_n}(n)n - 2^{q_{s_n}(n)+1} + 2 - 2 \left((s_n - 1)n + \sum_{i=1}^{s_n} 2^{q_i(n)-1} (q_i(n) - 2i) + 1 \right) \\ &= \sum_{i=1}^{s_n-1} 2^{q_i(n)} (q_{s_n}(n) - q_i(n) - 2(s_n - i - 1)) \end{aligned}$$

as it was proved in [8] by showing by induction that this sequence satisfies the recurrence above.

Remark 1. The solution of

$$x_n = a \cdot x_{\lceil n/2 \rceil} + a \cdot x_{\lfloor n/2 \rfloor} + (\lceil n/2 \rceil - \lfloor n/2 \rfloor)$$

and of

$$x_n = a \cdot x_{\lceil n/2 \rceil} + a \cdot x_{\lfloor n/2 \rfloor} + (\lceil n/2 \rceil - \lfloor n/2 \rfloor)^m$$

for $m \geq 1$, is the same. Therefore, for every $m \geq 1$,

$$x^{(1,0)}(a) - x^{(0,1)}(a) = \sum_{p=0}^m \binom{m}{p} (-1)^{m-p} x^{(p,m-p)}(a)$$

Example 4. The sequence $x_n = n^2$ satisfies the equation

$$x_n = 2x_{\lceil n/2 \rceil} + 2x_{\lfloor n/2 \rfloor} + \lfloor n/2 \rfloor - \lceil n/2 \rceil$$

with $x_1 = 1$. Therefore,

$$x_n = x_n^{(0,1)}(2) - x_n^{(1,0)}(2) + 3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)}$$

Now,

$$\begin{aligned} x_n^{(0,1)}(2) &= \alpha_n^{(0,0)}(2) + \frac{1}{3} (3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\ x_n^{(1,0)}(2) &= -n + 1 + \frac{2}{3} (3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) - \alpha_n^{(0,0)}(2) \end{aligned}$$

and

$$\begin{aligned}
\alpha_n^{(0,0)}(2) &= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) ((2^{q_i(n)} - 1) - (2^{q_i(n)-1} - 1)) + \sum_{i=1}^{s_n-1} 2^{q_i(n)} (n - M_i(n)) - \frac{4^{q_i(n)} - 1}{3} \\
&= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) (2^{q_i(n)} - 2^{q_i(n)-1}) + \sum_{i=1}^{s_n} 2^{q_i(n)} (n - M_i(n)) - \frac{1}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&= n^2 - \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) 2^{q_i(n)} - \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) 2^{q_i(n)-1} - \frac{1}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&= n^2 - \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) 2^{q_i(n)} - \frac{1}{2} \sum_{i=0}^{s_n} M_{i+1}(n) 2^{q_i(n)} - \frac{1}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&= n^2 - \frac{1}{2} \sum_{i=1}^{s_n} 2^{q_i(n)} (M_i(n) + M_{i+1}(n)) - \frac{1}{2} n - \frac{1}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&= n^2 - \frac{1}{2} \sum_{i=1}^{s_n} (M_i(n) - M_{i+1}(n)) (M_i(n) + M_{i+1}(n)) - \frac{1}{2} n - \frac{1}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&= n^2 - \frac{1}{2} \sum_{i=1}^{s_n} (M_i(n)^2 - M_{i+1}(n)^2) - \frac{1}{2} n - \frac{1}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&= \frac{1}{2} n^2 - \frac{1}{2} n - \frac{1}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1)
\end{aligned}$$

and therefore

$$\begin{aligned}
x_n &= \alpha_n^{(0,0)}(2) + \frac{1}{3} (3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&\quad - \left(-n + 1 + \frac{2}{3} (3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) - \alpha_n^{(0,0)}(2) \right) + 3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} \\
&= 2\alpha_n^{(0,0)}(2) + \frac{2}{3} (3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) + n \\
&= n^2 - n - \frac{2}{3} (3n2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) + \frac{2}{3} (3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) + n \\
&= n^2
\end{aligned}$$

indeed.

Example 5. The sequence λ_n of Lebesgue constants of the Walsh system satisfies the recurrence

$$\lambda_n = \frac{1}{2} \lambda_{\lceil n/2 \rceil} + \frac{1}{2} \lambda_{\lfloor n/2 \rfloor} + \frac{1}{2} \left\lceil \frac{n}{2} \right\rceil - \frac{1}{2} \left\lfloor \frac{n}{2} \right\rfloor$$

with $\lambda_1 = 1$ [13, 21]. Therefore,

$$\lambda_n = \frac{1}{2} x_n^{(1,0)}(1/2) - \frac{1}{2} x_n^{(0,1)}(1/2) + 1$$

where

$$\begin{aligned}
x_n^{(1,0)}(1/2) &= 2n - q_{s_n}(n) - n \cdot 2^{-q_{s_n}(n)} - 1 - \alpha_n^{(0,0)}(1/2) \\
x_n^{(0,1)}(1/2) &= \alpha_n^{(0,0)}(1/2) + q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1 \\
\alpha_n^{(0,0)}(1/2) &= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) (T(0, q_i(n), 1/2) - T(0, q_{i-1}(n), 1/2)) + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) - q_{s_n}(n) \\
&= \sum_{i=1}^{s_n} M_i(n) (2^{-q_{i-1}(n)} - 2^{-q_i(n)}) + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) - q_{s_n}(n) \\
&= \sum_{i=1}^{s_n} M_i(n) 2^{-q_{i-1}(n)} - \sum_{i=1}^{s_n} M_i(n) 2^{-q_i(n)} + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) - q_{s_n}(n) \\
&= \sum_{i=0}^{s_n} M_{i+1}(n) 2^{-q_i(n)} - \sum_{i=1}^{s_n} M_i(n) 2^{-q_i(n)} + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) - q_{s_n}(n) \\
&= n + \sum_{i=1}^{s_n} (M_{i+1}(n) - M_i(n)) 2^{-q_i(n)} + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) - q_{s_n}(n) \\
&= n - \sum_{i=1}^{s_n} 2^{q_i(n)} 2^{-q_i(n)} + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) - q_{s_n}(n) \\
&= n - s_n + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) - q_{s_n}(n)
\end{aligned}$$

Therefore,

$$\begin{aligned}
\lambda_n &= \frac{1}{2} (2n - q_{s_n}(n) - n \cdot 2^{-q_{s_n}(n)} - 1 - \alpha_n^{(0,0)}(1/2)) \\
&\quad - \frac{1}{2} (\alpha_n^{(0,0)}(1/2) + q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1) + 1 \\
&= -\alpha_n^{(0,0)}(1/2) + n - q_{s_n}(n) - n \cdot 2^{-q_{s_n}(n)} + 1 \\
&= -n + s_n - \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) + q_{s_n}(n) + n - q_{s_n}(n) - n \cdot 2^{-q_{s_n}(n)} + 1 \\
&= s_n - \sum_{i=1}^{s_n} 2^{-q_i(n)} (n - M_i(n))
\end{aligned}$$

in agreement with [13, Eqn. (5.5)].

Remark 2. As R. Stephan points out in [35], the OEIS contains many sequences defined by recurrences of the form

$$\begin{cases} a_{2n} = C \cdot a_n + C \cdot a_{n-1} + P(n) \\ a_{2n+1} = 2C \cdot a_n + Q(n) \end{cases}$$

for some real number C and functions P, Q . It is straightforward to check then that the sequence $x_n := a_{n-1}$ satisfies, for $n \geq 2$, the recurrence (cf. [22])

$$x_n = C \cdot x_{\lceil n/2 \rceil} + C \cdot x_{\lfloor n/2 \rfloor} + Q(\lfloor n/2 \rfloor - 1) + (\lceil n/2 \rceil - \lfloor n/2 \rfloor)(P(\lfloor n/2 \rfloor) - Q(\lfloor n/2 \rfloor - 1)).$$

If P and Q are polynomials, these sequences are covered by our main theorem.

Example 6. Let a_n denote the sequence A005536. By the second last entry in the last table of [35] and the last remark, the sequence $x_n = a_{n-1}$, for $n \geq 1$, satisfies the recurrence

$$x_n = -x_{\lceil n/2 \rceil} - x_{\lfloor n/2 \rfloor} + \lfloor n/2 \rfloor$$

with $x_1 = 0$; cf. also [22, Ex 7.1]. By our main theorem, this sequence is

$$x_n = x_n^{(0,1)}(-1)$$

where

$$\begin{aligned} x_n^{(0,1)}(-1) &= \alpha_n^{(0,0)}(-1) - \frac{1}{3}((-2)^{q_{s_n}(n)} - 1) + (-1)^{q_{s_n}(n)}n - (-2)^{q_{s_n}(n)} \\ &= \alpha_n^{(0,0)}(-1) - \frac{4}{3} \cdot (-2)^{q_{s_n}(n)} + \frac{1}{3} + (-1)^{q_{s_n}(n)}n \\ \alpha_n^{(0,0)}(-1) &= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) (T(0, q_i(n), -1) - T(0, q_{i-1}(n), -1)) \\ &\quad + \sum_{i=1}^{s_n-1} (-1)^{q_i(n)} (n - M_i(n)) - T(0, q_{s_n}(n), -2) \\ &= \frac{1}{4} \sum_{i=1}^{s_n} M_i(n) ((-1)^{q_{i-1}(n)} - (-1)^{q_i(n)}) + \sum_{i=1}^{s_n-1} (-1)^{q_i(n)} (n - M_i(n)) + \frac{1}{3}((-2)^{q_{s_n}(n)} - 1) \\ &= \frac{1}{4} \sum_{i=1}^{s_n} M_i(n) (-1)^{q_{i-1}(n)} - \frac{1}{4} \sum_{i=1}^{s_n} M_i(n) (-1)^{q_i(n)} + \sum_{i=1}^{s_n-1} (-1)^{q_i(n)} (n - M_i(n)) \\ &\quad + \frac{1}{3}((-2)^{q_{s_n}(n)} - 1) \\ &= \frac{1}{4} \sum_{i=0}^{s_n} M_{i+1}(n) (-1)^{q_i(n)} - \frac{1}{4} \sum_{i=1}^{s_n} M_i(n) (-1)^{q_i(n)} + \sum_{i=1}^{s_n-1} (-1)^{q_i(n)} (n - M_i(n)) \\ &\quad + \frac{1}{3}((-2)^{q_{s_n}(n)} - 1) \\ &= \frac{1}{4}n - \frac{1}{4} \sum_{i=1}^{s_n} (-2)^{q_i(n)} + \sum_{i=1}^{s_n-1} (-1)^{q_i(n)} (n - M_i(n)) + \frac{1}{3}((-2)^{q_{s_n}(n)} - 1) \end{aligned}$$

and hence

$$\begin{aligned} x_n &= \frac{1}{4}n - \frac{1}{4} \sum_{i=1}^{s_n} (-2)^{q_i(n)} + \sum_{i=1}^{s_n-1} (-1)^{q_i(n)} (n - M_i(n)) + \frac{1}{3}((-2)^{q_{s_n}(n)} - 1) \\ &\quad - \frac{4}{3} \cdot (-2)^{q_{s_n}(n)} + \frac{1}{3} + (-1)^{q_{s_n}(n)}n \\ &= \frac{1}{4}n - \frac{1}{4} \sum_{i=1}^{s_n} (-2)^{q_i(n)} + \sum_{i=1}^{s_n} (-1)^{q_i(n)} (n - M_i(n)) \\ &= \frac{1}{4}n - \frac{1}{4} \sum_{i=1}^{s_n} (-2)^{q_i(n)} + \sum_{i=1}^{s_n} \left((-1)^{q_i(n)} \sum_{j=1}^{i-1} 2^{q_j(n)} \right) \\ &= \frac{1}{4} \sum_{i=1}^{s_n} 2^{q_i(n)} \left(1 - (-1)^{q_i(n)} + 4 \sum_{j=i+1}^{s_n} (-1)^{q_j(n)} \right) \end{aligned}$$

which is a simpler expression than the one given in [22, Ex 7.1].

Example 7. Let a_n denote the sequence A087733. By Remark 2, the sequence $x_n = a_{n-1}$, for $n \geq 1$, satisfies the recurrence

$$x_n = -x_{\lceil n/2 \rceil} - x_{\lfloor n/2 \rfloor} + \lfloor n/2 \rfloor \lceil n/2 \rceil$$

with $x_1 = 0$. By our main theorem, this sequence is

$$\begin{aligned}
x_n &= x_n^{(1,1)}(-1) \\
&= \sum_{k=1}^2 \left(\sum_{j=k}^2 \frac{\binom{1}{j-2} \binom{j}{k} B_{j-k}}{j(2^{j-1} + 1)} \right) n^k \\
&\quad + \left(1 - \sum_{l=0}^0 \frac{\binom{1}{l}}{2^{1+l} + 1} \right) \left(\frac{(-2)^{q_{s_n}(n)} - 1}{-3} + n(-1)^{q_{s_n}(n)} - (-2)^{q_{s_n}(n)} \right) \\
&\quad + \sum_{i=0}^1 \left(2^{-i} \binom{2}{i} - 2^{-i+1} \binom{1}{i-1} - \sum_{l=i}^0 \frac{\binom{1}{l} \binom{1+l}{i}}{2^{1+l} + 1} \right) \alpha_n^{(0,i)}(-1) \\
&= \frac{1}{6} n^2 - \frac{1}{6} n + \frac{2}{9} (1 + 3(-1)^{q_{s_n}(n)} n - 4(-2)^{q_{s_n}(n)}) + \frac{2}{3} \alpha_n^{(0,0)}(-1) \\
&= \frac{1}{6} n^2 - \frac{1}{6} n + \frac{2}{9} (1 + 3(-1)^{q_{s_n}(n)} n - 4(-2)^{q_{s_n}(n)}) \\
&\quad + \frac{2}{3} \left[\frac{1}{4} n - \frac{1}{4} \sum_{i=1}^{s_n} (-2)^{q_i(n)} + \sum_{i=1}^{s_n-1} (-1)^{q_i(n)} (n - M_i(n)) + \frac{1}{3} ((-2)^{q_{s_n}(n)} - 1) \right] \\
&\text{(cf. Example 6)} \\
&= \frac{1}{6} n^2 - \frac{1}{6} \sum_{i=1}^{s_n} (-2)^{q_i(n)} + \frac{2}{3} \sum_{i=1}^{s_n} (-1)^{q_i(n)} (n - M_i(n)) \\
&= \frac{1}{6} n^2 - \frac{1}{6} \sum_{i=1}^{s_n} (-2)^{q_i(n)} + \frac{2}{3} \sum_{i=1}^{s_n-1} \left(2^{q_i(n)} \sum_{j=i+1}^{s_n} (-1)^{q_j(n)} \right)
\end{aligned}$$

To our knowledge, no closed expression had been published so far for this sequence.

Example 8. The minimum total cophenetic index Φ_n of a rooted bifurcating tree with n leaves (sequence A174605) satisfies the recurrence

$$\begin{aligned}
\Phi_n &= \Phi_{\lceil n/2 \rceil} + \Phi_{\lfloor n/2 \rfloor} + \binom{\lceil n/2 \rceil}{2} + \binom{\lfloor n/2 \rfloor}{2} \\
&= \Phi_{\lceil n/2 \rceil} + \Phi_{\lfloor n/2 \rfloor} + \frac{1}{2} \left\lceil \frac{n}{2} \right\rceil^2 - \frac{1}{2} \left\lceil \frac{n}{2} \right\rceil + \frac{1}{2} \left\lfloor \frac{n}{2} \right\rfloor^2 - \frac{1}{2} \left\lfloor \frac{n}{2} \right\rfloor
\end{aligned}$$

with initial condition $\Phi_1 = 0$ [26]. Therefore

$$\Phi_n = \frac{1}{2} (x_n^{(2,0)}(1) + x_n^{(0,2)}(1) - x_n^{(1,0)}(1) - x_n^{(0,1)}(1)) = \frac{1}{2} (x_n^{(2,0)}(1) + x_n^{(0,2)}(1) - S_n)$$

where

$$S_n = x_n^{(1,0)}(1) + x_n^{(0,1)}(1) = (q_{s_n}(n) + 2)n - 2^{q_{s_n}(n)+1}$$

was already computed in Example 2 and

$$\begin{aligned}
x_n^{(2,0)}(1) &= n^2 + (q_{s_n}(n) - 2)n + 3 - 2^{q_{s_n}(n)+1} - 3\alpha_n^{(0,0)}(1) - \alpha_n^{(0,1)}(1) \\
x_n^{(0,2)}(1) &= \alpha_n^{(0,0)}(1) + \alpha_n^{(0,1)}(1) + n - 1
\end{aligned}$$

Therefore

$$\begin{aligned}
\Phi_n &= \frac{1}{2} \left(n^2 + (q_{s_n}(n) - 2)n + 3 - 2^{q_{s_n}(n)+1} - 3\alpha_n^{(0,0)}(1) - \alpha_n^{(0,1)}(1) \right. \\
&\quad \left. + \alpha_n^{(0,0)}(1) + \alpha_n^{(0,1)}(1) + n - 1 - ((q_{s_n}(n) + 2)n - 2^{q_{s_n}(n)+1}) \right) \\
&= \frac{1}{2} (n^2 - 3n + 2 - 2\alpha_n^{(0,0)}(1)) \\
&= \binom{n-1}{2} - \left(s_n n + \sum_{i=1}^{s_n} 2^{q_i(n)-1} (q_i(n) - 2i) - n + 1 \right) \\
&\text{(by Eqn. (3))} \\
&= \binom{n}{2} - s_n n - \sum_{i=1}^{s_n} 2^{q_i(n)-1} (q_i(n) - 2i)
\end{aligned}$$

No closed expression had been published so far for this sequence.

Example 9. The maximum rooted quartet index ρ_n of a rooted bifurcating tree with n leaves (sequence A300445) satisfies the recurrence

$$\begin{aligned}
\rho_n &= \rho_{\lceil n/2 \rceil} + \rho_{\lfloor n/2 \rfloor} + \binom{\lceil n/2 \rceil}{2} \binom{\lfloor n/2 \rfloor}{2} \\
&= \rho_{\lceil n/2 \rceil} + \rho_{\lfloor n/2 \rfloor} + \frac{1}{4} \left(\binom{\lceil n/2 \rceil}{2} \binom{\lfloor n/2 \rfloor}{2} - \lceil n/2 \rceil^2 \lfloor n/2 \rfloor - \lfloor n/2 \rfloor^2 \lceil n/2 \rceil + \lceil n/2 \rceil^2 \lfloor n/2 \rfloor^2 \right)
\end{aligned}$$

with initial condition $\rho_1 = 0$ [9]. Therefore

$$\rho_n = \frac{1}{4} (x_n^{(1,1)}(1) - x_n^{(2,1)}(1) - x_n^{(1,2)}(1) + x_n^{(2,2)}(1))$$

where

$$\begin{aligned}
x_n^{(1,1)}(1) &= \frac{1}{2} n^2 + (1 + B_1 - 1)n + (1 - 0 - 1)\alpha_n^{(0,0)}(1) + (1 - 1 - 0)\alpha_n^{(0,1)}(1) = \binom{n}{2} \\
x_n^{(2,1)}(1) &= \sum_{k=2}^3 \left(\sum_{j=k}^3 \frac{\binom{j-2}{k} \binom{j}{k} B_{j-k}}{j(2^{j-1} - 1)} \right) n^k + \left(1 + \sum_{l=1}^2 \frac{\binom{l-1}{2} (B_l - 1)}{2^l - 1} \right) n + \sum_{l=0}^1 \frac{\binom{2}{l}}{2^{1+l} - 1} - 1 \\
&\quad + \sum_{i=0}^2 \left(2^{-i} \binom{3}{i} - 2^{-i+1} \binom{2}{i-1} - \sum_{l=i}^1 \frac{\binom{2}{l} \binom{1+l}{i}}{2^{1+l} - 1} \right) \alpha_n^{(0,i)}(1) \\
&= \frac{2}{9} n^3 + \frac{1}{6} n^2 - \frac{19}{18} n + \frac{2}{3} - \frac{2}{3} \alpha_n^{(0,0)}(1) - \frac{5}{6} \alpha_n^{(0,1)}(1) - \frac{1}{4} \alpha_n^{(0,2)}(1) \\
x_n^{(1,2)}(1) &= \sum_{k=2}^3 \left(\sum_{j=k}^3 \frac{\binom{j-3}{k} \binom{j}{k} B_{j-k}}{j(2^{j-1} - 1)} \right) n^k + \left(1 + \sum_{l=1}^2 \frac{\binom{l-2}{1} (B_l - 1)}{2^l - 1} \right) n + \sum_{l=0}^0 \frac{\binom{1}{l}}{2^{2+l} - 1} - 1 \\
&\quad + \sum_{i=0}^2 \left(2^{-i} \binom{3}{i} - 2^{-i+1} \binom{1}{i-2} - \sum_{l=i-1}^0 \frac{\binom{1}{l} \binom{2+l}{i}}{2^{2+l} - 1} \right) \alpha_n^{(0,i)}(1) \\
&= \frac{1}{9} n^3 - \frac{1}{6} n^2 + \frac{13}{18} n - \frac{2}{3} + \frac{2}{3} \alpha_n^{(0,0)}(1) + \frac{5}{6} \alpha_n^{(0,1)}(1) + \frac{1}{4} \alpha_n^{(0,2)}(1) \\
x_n^{(2,2)}(1) &= \sum_{k=2}^4 \left(\sum_{j=k}^4 \frac{\binom{j-3}{k} \binom{j}{k} B_{j-k}}{j(2^{j-1} - 1)} \right) n^k + \left(1 + \sum_{l=1}^3 \frac{\binom{l-2}{2} (B_l - 1)}{2^l - 1} \right) n + \sum_{l=0}^1 \frac{\binom{2}{l}}{2^{2+l} - 1} - 1 \\
&\quad + \sum_{i=0}^3 \left(2^{-i} \binom{4}{i} - 2^{-i+1} \binom{2}{i-2} - \sum_{l=i-1}^1 \frac{\binom{2}{l} \binom{2+l}{i}}{2^{2+l} - 1} \right) \alpha_n^{(0,i)}(1) \\
&= \frac{1}{14} n^4 - \frac{2}{63} n^3 - \frac{2}{21} n^2 + \frac{55}{126} n - \frac{8}{21} + \frac{8}{21} \alpha_n^{(0,0)}(1) + \frac{10}{21} \alpha_n^{(0,1)}(1) + \frac{1}{7} \alpha_n^{(0,2)}(1)
\end{aligned}$$

Therefore,

$$\begin{aligned}
\rho_n &= \frac{1}{4} \left[\binom{n}{2} - \left(\frac{2}{9}n^3 + \frac{1}{6}n^2 - \frac{19}{18}n + \frac{2}{3} - \frac{2}{3}\alpha_n^{(0,0)}(1) - \frac{5}{6}\alpha_n^{(0,1)}(1) - \frac{1}{4}\alpha_n^{(0,2)}(1) \right) \right. \\
&\quad - \left(\frac{1}{9}n^3 - \frac{1}{6}n^2 + \frac{13}{18}n - \frac{2}{3} + \frac{2}{3}\alpha_n^{(0,0)}(1) + \frac{5}{6}\alpha_n^{(0,1)}(1) + \frac{1}{4}\alpha_n^{(0,2)}(1) \right) \\
&\quad \left. + \left(\frac{1}{14}n^4 - \frac{2}{63}n^3 - \frac{2}{21}n^2 + \frac{55}{126}n - \frac{8}{21} + \frac{8}{21}\alpha_n^{(0,0)}(1) + \frac{10}{21}\alpha_n^{(0,1)}(1) + \frac{1}{7}\alpha_n^{(0,2)}(1) \right) \right] \\
&= \frac{1}{504} [(n-3)(n-2)(n-1)(9n+8) + 48\alpha_n^{(0,0)}(1) + 60\alpha_n^{(0,1)}(1) + 18\alpha_n^{(0,2)}(1)] \quad (4)
\end{aligned}$$

Now,

$$\begin{aligned}
\alpha_n^{(0,0)}(1) &= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(q_i(n) - q_{i-1}(n)) + \sum_{i=1}^{s_n-1} (n - M_i(n)) - 2^{q_{s_n}(n)} + 1 \quad (\text{by Eqn. (2)}) \\
&= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(q_i(n) - q_{i-1}(n)) + \sum_{i=1}^{s_n} (n - M_i(n)) - n + 1 \\
\alpha_n^{(0,1)}(1) &= \frac{1}{4} \sum_{i=1}^{s_n} \left(M_i(n)^2 (T(0, q_i(n), 1/2) - T(0, q_{i-1}(n), 1/2)) + 4B_1 M_i(n)(q_i(n) - q_{i-1}(n)) \right) \\
&\quad + \sum_{i=1}^{s_n-1} 2^{-q_i(n)} (n - M_i(n)) M_{i+1}(n) \\
&= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)^2 \left(2^{-q_{i-1}(n)} - 2^{-q_i(n)} \right) - \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(q_i(n) - q_{i-1}(n)) + \sum_{i=1}^{s_n} 2^{-q_i(n)} M_{i+1}(n) \\
\alpha_n^{(0,2)}(1) &= \frac{1}{6} \sum_{i=1}^{s_n} \left(M_i(n)^3 (T(0, q_i(n), 1/4) - T(0, q_{i-1}(n), 1/4)) \right. \\
&\quad \left. + 6B_1 M_i(n)^2 (T(0, q_i(n), 1/2) - T(0, q_{i-1}(n), 1/2)) + 12B_2 M_i(n)(q_i(n) - q_{i-1}(n)) \right) \\
&\quad + \sum_{i=1}^{s_n-1} 4^{-q_i(n)} (n - M_i(n)) M_{i+1}(n)^2 \\
&= \frac{2}{9} \sum_{i=1}^{s_n} M_i(n)^3 (4^{-q_{i-1}(n)} - 4^{-q_i(n)}) - \sum_{i=1}^{s_n} M_i(n)^2 (2^{-q_{i-1}(n)} - 2^{-q_i(n)}) \\
&\quad + \frac{1}{3} \sum_{i=1}^{s_n} M_i(n)(q_i - q_{i-1}) + \sum_{i=1}^{s_n} 4^{-q_i(n)} M_{i+1}(n)^2 (n - M_i(n))
\end{aligned}$$

and then, substituting these expressions in Eqn. (4), we finally obtain

$$\begin{aligned}
\rho_n &= \frac{1}{252} \left[3(9n+8) \binom{n-1}{3} - 24n + 24 \right. \\
&\quad \left. + 2 \sum_{i=1}^{s_n} M_i(n)^3 (4^{-q_{i-1}(n)} - 4^{-q_i(n)}) + 6 \sum_{i=1}^{s_n} M_i(n)^2 (2^{-q_{i-1}(n)} - 2^{-q_i(n)}) \right. \\
&\quad \left. + \sum_{i=1}^{s_n} (n - M_i(n)) (24 + 30 \cdot 2^{-q_i(n)} M_{i+1}(n) + 9 \cdot 4^{-q_i(n)} M_{i+1}(n)^2) \right]
\end{aligned}$$

This expression can be simplified as follows. Notice that

$$\begin{aligned}
& 2 \sum_{i=1}^{s_n} M_i(n)^3 (4^{-q_{i-1}(n)} - 4^{-q_i(n)}) + 6 \sum_{i=1}^{s_n} M_i(n)^2 (2^{-q_{i-1}(n)} - 2^{-q_i(n)}) \\
&= 2 \left(\sum_{i=0}^{s_n} M_{i+1}(n)^3 4^{-q_i(n)} - \sum_{i=1}^{s_n} M_i(n)^3 4^{-q_i(n)} \right) \\
&\quad + 6 \left(\sum_{i=0}^{s_n-1} M_{i+1}(n)^2 2^{-q_i(n)} - \sum_{i=1}^{s_n} M_i(n)^2 2^{-q_i(n)} \right) \\
&= 2n^3 + 2 \sum_{i=1}^{s_n} 4^{-q_i(n)} (M_{i+1}(n)^3 - M_i(n)^3) \\
&\quad + 6n^2 + 6 \sum_{i=1}^{s_n} 2^{-q_i(n)} (M_{i+1}(n)^2 - M_i(n)^2) \\
&= 2n^3 + 6n^2 + 2 \sum_{i=1}^{s_n} 4^{-q_i(n)} (-3M_i(n)^2 2^{q_i(n)} + 3M_i(n) 4^{q_i(n)} - 8^{q_i(n)}) \\
&\quad + 6 \sum_{i=1}^{s_n} 2^{-q_i(n)} (-2M_i(n) 2^{q_i(n)} + 4^{q_i(n)}) \\
&= 2n^3 + 6n^2 - 2 \sum_{i=1}^{s_n} (3M_i(n)^2 2^{-q_i(n)} + 3M_i(n) - 2 \cdot 2^{q_i(n)}) \\
&= 2n^3 + 6n^2 + 4n - 6 \sum_{i=1}^{s_n} M_i(n) (M_i(n) 2^{-q_i(n)} + 1)
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{i=1}^{s_n} (n - M_i(n)) (24 + 30 \cdot 2^{-q_i(n)} M_{i+1}(n) + 9 \cdot 4^{-q_i(n)} M_{i+1}(n)^2) \\
&= \sum_{i=1}^{s_n} (n - M_i(n)) (24 + 30 \cdot 2^{-q_i(n)} (M_i(n) - 2^{q_i(n)}) + 9 \cdot 4^{-q_i(n)} (M_i(n) - 2^{q_i(n)})^2) \\
&= \sum_{i=1}^{s_n} (n - M_i(n)) (3 + 12 \cdot 2^{-q_i(n)} M_i(n) + 9 \cdot 4^{-q_i(n)} M_i(n)^2) \\
&= 3n \sum_{i=1}^{s_n} (1 + 4 \cdot 2^{-q_i(n)} M_i(n) + 3 \cdot 4^{-q_i(n)} M_i(n)^2) \\
&\quad - 3 \sum_{i=1}^{s_n} M_i(n) (1 + 4 \cdot 2^{-q_i(n)} M_i(n) + 3 \cdot 4^{-q_i(n)} M_i(n)^2)
\end{aligned}$$

and hence, finally

$$\begin{aligned}
\rho_n &= \frac{1}{252} \left[3(9n+8) \binom{n-1}{3} - 24n + 24 + 2n^3 + 6n^2 + 4n \right. \\
&\quad - 6 \sum_{i=1}^{s_n} M_i(n) (M_i(n) 2^{-q_i(n)} + 1) + 3n \sum_{i=1}^{s_n} (1 + 4 \cdot 2^{-q_i(n)} M_i(n) + 3 \cdot 2^{-2q_i(n)} M_i(n)^2) \\
&\quad \left. - 3 \sum_{i=1}^{s_n} M_i(n) (1 + 4 \cdot 2^{-q_i(n)} M_i(n) + 3 \cdot 2^{-2q_i(n)} M_i(n)^2) \right] \\
&= \frac{1}{504} \left[9n^4 - 42n^3 + 63n^2 - 6n + 6n \sum_{i=1}^{s_n} (1 + M_i(n) 2^{-q_i(n)}) (1 + 3M_i(n) 2^{-q_i(n)}) \right. \\
&\quad \left. - 18 \sum_{i=1}^{s_n} M_i(n) (1 + M_i(n) 2^{-q_i(n)})^2 \right]
\end{aligned}$$

where notice that $M_i(n) 2^{-q_i(n)} = \lfloor n/2^{q_i(n)} \rfloor$. Again, no closed expression was known so far for this sequence, either.

Example 10. Let $a_{1,n}$ and $a_{3,n}$ denote the sequences A006581 and A006583, respectively. By Remark 2, the sequences $\sigma_{i,n} = a_{i,n-1}$, for $i = 1, 3$, are the solutions with $\sigma_{1,1} = \sigma_{3,1} = 0$ of the recurrences

$$\sigma_{i,n} = 2\sigma_{i,\lceil n/2 \rceil} + 2\sigma_{i,\lfloor n/2 \rfloor} + g_i(n)$$

where

$$\begin{aligned}
g_1(n) &= \left\lfloor \frac{n}{2} \right\rfloor \left(\left\lceil \frac{n}{2} \right\rceil - \left\lfloor \frac{n}{2} \right\rfloor \right) = \left\lfloor \frac{n}{2} \right\rfloor \left\lceil \frac{n}{2} \right\rceil - \left\lfloor \frac{n}{2} \right\rfloor^2 \\
g_3(n) &= 6 \left(\left\lfloor \frac{n}{2} \right\rfloor - 1 \right) + \left(\left\lceil \frac{n}{2} \right\rceil - \left\lfloor \frac{n}{2} \right\rfloor \right) \left(5 \left\lfloor \frac{n}{2} \right\rfloor - 4 - 6 \left\lfloor \frac{n}{2} \right\rfloor + 6 \right) \\
&= 4 \left\lfloor \frac{n}{2} \right\rfloor + 2 \left\lceil \frac{n}{2} \right\rceil - 6 - g_1(n)
\end{aligned}$$

Therefore,

$$\begin{aligned}
\sigma_{1,n} &= x_n^{(1,1)}(2) - x_n^{(0,2)}(2) \\
\sigma_{3,n} &= 4x_n^{(0,1)}(2) + 2x_n^{(1,0)}(2) - 6x_n^{(0,0)}(2) - \sigma_{1,n}
\end{aligned}$$

where

$$\begin{aligned}
x_n^{(0,0)}(2) &= \frac{1}{3}(3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \quad (\text{cf. Example 1}) \\
x_n^{(0,1)}(2) &= \alpha_n^{(0,0)}(2) + x_n^{(0,0)}(2) \\
x_n^{(0,2)}(2) &= \alpha_n^{(0,0)}(2) + \alpha_n^{(0,1)}(2) + x_n^{(0,0)}(2) \\
x_n^{(1,0)}(2) &= -n + \frac{2}{3}(3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) + 1 - \alpha_n^{(0,0)}(2) \\
&= 2x_n^{(0,0)}(2) - n + 1 - \alpha_n^{(0,0)}(2) \\
x_n^{(1,1)}(2) &= \sum_{k=1}^2 \left(\sum_{\substack{j=k \\ j \neq 2}}^2 \frac{\binom{1}{j-2} \binom{j}{k} B_{j-k}}{j(2^{j-1} - 2)} \right) n^k \\
&\quad + \frac{1}{2} \left(\frac{(q_{s_n}(n) - 1)4^{q_{s_n}(n)+1} - q_{s_n}(n)4^{q_{s_n}(n)} + 4}{9} + nq_{s_n}(n)2^{q_{s_n}(n)} - q_{s_n}(n)4^{q_{s_n}(n)} \right) \\
&\quad + \left(1 - \sum_{\substack{l=0 \\ l \neq 0}}^0 \frac{\binom{1}{l}}{2^{1+l} - 2} \right) \frac{1}{3}(3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&\quad + \sum_{i=0}^1 \left(2^{-i} \binom{2}{i} - 2^{-i+1} \binom{1}{i-1} - \sum_{\substack{l=i \\ l \neq 0}}^0 \frac{\binom{1}{l} \binom{1+l}{i}}{2^{1+l} - 2} \right) \alpha_n^{(0,i)}(2) + \frac{1}{2} \sum_{i=0}^0 \binom{1}{i} \alpha_n^{(1,i)}(a) \\
&= \frac{1}{18}(9nq_{s_n}(n)2^{q_{s_n}(n)} - 6q_{s_n}(n)4^{q_{s_n}(n)} - 4 \cdot 4^{q_{s_n}(n)} + 4) \\
&\quad + \frac{1}{3}(3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) + \alpha_n^{(0,0)}(2) + \frac{1}{2}\alpha_n^{(1,0)}(2)
\end{aligned}$$

and

$$\begin{aligned}
\alpha_n^{(0,0)}(2) &= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(T(0, q_i(n), 2) - T(0, q_{i-1}(n), 2)) + \sum_{i=1}^{s_n-1} 2^{q_i(n)}(n - M_i(n)) - T(0, q_{s_n}(n), 4) \\
&= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(2^{q_i(n)} - 2^{q_{i-1}(n)}) + \sum_{i=1}^{s_n-1} 2^{q_i(n)}(n - M_i(n)) - \frac{1}{3}(4^{q_{s_n}(n)} - 1) \\
\alpha_n^{(0,1)}(2) &= \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)^2(q_i(n) - q_{i-1}(n)) - \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(2^{q_i(n)} - 2^{q_{i-1}(n)}) \\
&\quad + \sum_{i=1}^{s_n-1} (n - M_i(n))M_{i+1}(n) \\
\alpha_n^{(1,0)}(2) &= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(T(1, q_i(n), 2) - T(1, q_{i-1}(n), 2)) + \sum_{i=1}^{s_n-1} q_i(n)2^{q_i(n)}(n - M_i(n)) \\
&\quad - T(1, q_{s_n}(n), 4) \\
&= \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)((q_i(n) - 2)2^{q_i(n)} - (q_{i-1}(n) - 2)2^{q_{i-1}(n)}) \\
&\quad + \sum_{i=1}^{s_n-1} q_i(n)2^{q_i(n)}(n - M_i(n)) - \frac{1}{9}((3q_{s_n}(n) - 4)4^{q_{s_n}(n)} + 4)
\end{aligned}$$

Therefore, finally,

$$\begin{aligned}
\sigma_{1,n} &= x_n^{(1,1)}(2) - x_n^{(0,2)}(2) \\
&= \frac{1}{18}(9nq_{s_n}(n)2^{q_{s_n}(n)} - 6q_{s_n}(n)4^{q_{s_n}(n)} - 4 \cdot 4^{q_{s_n}(n)} + 4) \\
&\quad + \frac{1}{3}(3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) + \alpha_n^{(0,0)}(2) + \frac{1}{2}\alpha_n^{(1,0)}(2) \\
&\quad - \alpha_n^{(0,0)}(2) - \alpha_n^{(0,1)}(2) - \frac{1}{3}(3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) \\
&= \frac{1}{18}(9nq_{s_n}(n)2^{q_{s_n}(n)} - 6q_{s_n}(n)4^{q_{s_n}(n)} - 4 \cdot 4^{q_{s_n}(n)} + 4) + \frac{1}{2}\alpha_n^{(1,0)}(2) - \alpha_n^{(0,1)}(2) \\
&= \frac{1}{18}(9nq_{s_n}(n)2^{q_{s_n}(n)} - 6q_{s_n}(n)4^{q_{s_n}(n)} - 4 \cdot 4^{q_{s_n}(n)} + 4) \\
&\quad + \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)((q_i(n) - 2)2^{q_i(n)} - (q_{i-1}(n) - 2)2^{q_{i-1}(n)}) \\
&\quad + \frac{1}{2} \sum_{i=1}^{s_n-1} q_i(n)2^{q_i(n)}(n - M_i(n)) - \frac{1}{18}(3q_{s_n}(n)4^{q_{s_n}(n)} - 4 \cdot 4^{q_{s_n}(n)} + 4) \\
&\quad - \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)^2(q_i(n) - q_{i-1}(n)) + \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)(2^{q_i(n)} - 2^{q_{i-1}(n)}) \\
&\quad - \sum_{i=1}^{s_n-1} (n - M_i(n))M_{i+1}(n) \\
&= \frac{1}{2}(nq_{s_n}(n)2^{q_{s_n}(n)} - q_{s_n}(n)4^{q_{s_n}(n)}) + \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)(q_i(n)2^{q_i(n)} - q_{i-1}(n)2^{q_{i-1}(n)}) \\
&\quad - \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)^2(q_i(n) - q_{i-1}(n)) + \frac{1}{2} \sum_{i=1}^{s_n-1} q_i(n)2^{q_i(n)}(n - M_i(n)) \\
&\quad - \sum_{i=1}^{s_n-1} (n - M_i(n))M_{i+1}(n) \\
&= \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)(q_i(n)2^{q_i(n)} - q_{i-1}(n)2^{q_{i-1}(n)}) - \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)^2(q_i(n) - q_{i-1}(n)) \\
&\quad + \frac{1}{2} \sum_{i=1}^{s_n} q_i(n)2^{q_i(n)}(n - M_i(n)) - \sum_{i=1}^{s_n} (n - M_i(n))M_{i+1}(n) \\
&= \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)q_i(n)(2^{q_i(n)} - M_i(n)) + \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)q_{i-1}(n)(M_i(n) + 2^{q_{i-1}(n)}) \\
&\quad - \frac{1}{2} \sum_{i=1}^{s_n} M_i(n)q_{i-1}(n)2^{q_{i-1}(n)} + \frac{1}{2} \sum_{i=1}^{s_n} q_i(n)2^{q_i(n)}(n - M_{i+1}(n) - 2^{q_i(n)}) \\
&\quad - \sum_{i=1}^{s_n} (n - M_i(n))M_{i+1}(n) \\
&= -\frac{1}{4} \sum_{i=1}^{s_n} M_i(n)q_i(n)M_{i+1}(n) + \frac{1}{4} \sum_{i=1}^{s_n} M_i(n)q_{i-1}(n)M_{i-1}(n) \\
&\quad - \frac{1}{2} \sum_{i=0}^{s_n} M_{i+1}(n)q_i(n)2^{q_i(n)} - \frac{1}{2} \sum_{i=1}^{s_n} q_i(n)2^{q_i(n)}M_{i+1}(n) \\
&\quad + \frac{1}{2} \sum_{i=1}^{s_n} q_i(n)2^{q_i(n)}(n - 2^{q_i(n)}) - \sum_{i=1}^{s_n} (n - M_i(n))M_{i+1}(n)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{4} \sum_{i=1}^{s_n} M_i(n) q_i(n) M_{i+1}(n) + \frac{1}{4} \sum_{i=0}^{s_n-1} M_{i+1}(n) q_i(n) M_i(n) \\
&\quad - \sum_{i=1}^{s_n} q_i(n) 2^{q_i(n)} M_{i+1}(n) + \frac{1}{2} \sum_{i=1}^{s_n} q_i(n) 2^{q_i(n)} (n - 2^{q_i(n)}) - \sum_{i=1}^{s_n} (n - M_i(n)) M_{i+1}(n) \\
&= \frac{1}{2} \sum_{i=1}^{s_n} q_i(n) 2^{q_i(n)} (n - 2^{q_i(n)} - 2M_{i+1}(n)) - \sum_{i=1}^{s_n} (n - M_i(n)) M_{i+1}(n)
\end{aligned}$$

and

$$\begin{aligned}
\sigma_{3,n} &= 4x_n^{(0,1)}(2) + 2x_n^{(1,0)}(2) - 6x_n^{(0,0)}(2) - \sigma_{1,n} \\
&= 4(\alpha_n^{(0,0)}(2) + x_n^{(0,0)}(2)) + 2(2x_n^{(0,0)}(2) - n + 1 - \alpha_n^{(0,0)}(2)) - 6x_n^{(0,0)}(2) - \sigma_{1,n} \\
&= 2\alpha_n^{(0,0)}(2) + 2x_n^{(0,0)}(2) - 2n + 2 - \sigma_{1,n} \\
&= \sum_{i=1}^{s_n} M_i(n)(2^{q_i(n)} - 2^{q_{i-1}(n)}) + 2 \sum_{i=1}^{s_n-1} 2^{q_i(n)}(n - M_i(n)) - \frac{2}{3}(4^{q_{s_n}(n)} - 1) \\
&\quad + \frac{2}{3}(3n \cdot 2^{q_{s_n}(n)} - 2 \cdot 4^{q_{s_n}(n)} - 1) - 2n + 2 - \sigma_{1,n} \\
&= \sum_{i=1}^{s_n} M_i(n)(2^{q_i(n)} - 2^{q_{i-1}(n)}) + 2 \sum_{i=1}^{s_n} 2^{q_i(n)}(n - M_i(n)) - 2n + 2 - \sigma_{1,n} \\
&= -\sum_{i=1}^{s_n} M_i(n) 2^{q_{i-1}(n)} + n^2 + \sum_{i=1}^{s_n} 2^{q_i(n)}(n - M_i(n)) - 2n + 2 - \sigma_{1,n} \\
&= 2 \binom{n-1}{2} - \sigma_{1,n}
\end{aligned}$$

Again, no closed formulas for these sequences had been published so far.

Remark 3. Notice that $x_{2^m}^{(r,t)} = 2ax_{2^{m-1}}^{(r,t)} + 2^{(m-1)(r+t)}$ and therefore, since $x_1^{(r,t)} = 0$,

$$\begin{aligned}
x_{2^m}^{(r,t)}(a) &= \sum_{k=0}^{m-1} (2a)^k 2^{(r+t)(m-k-1)} = 2^{(r+t)(m-1)} \sum_{k=0}^{m-1} (2^{-r-t+1}a)^k \\
&= 2^{(r+t)(m-1)} T(0, m, 2^{-r-t+1}a) = \begin{cases} 2^{(r+t)(m-1)} m & \text{if } a = 2^{r+t-1}, \text{ i.e., if } \ell = r-1 \\ \frac{(2a)^m - 2^{m(r+t)}}{2a - 2^{r+t}} & \text{if } a \neq 2^{r+t-1} \end{cases}
\end{aligned}$$

We have checked that our general formula for $x_n^{(r,t)}(a)$ satisfies this equality when $n = 2^m$ with *Mathematica*.

4 Proof of the main result

4.1 Some notations

1. Throughout this paper we shall use the following notations related to binary decompositions of natural numbers. For every $n \in \mathbb{N}$, we shall write its binary decomposition as

$$n = \sum_{j=1}^{s_n} 2^{q_j(n)}, \quad \text{with } 0 \leq q_1(n) < \dots < q_{s_n}(n);$$

if $n = 0$, we set $s_0 = 0$. With these notations, s_n is the *binary weight* of n , that is, the number of 1's in the binary representation of n , and, if $n \geq 1$, $q_{s_n}(n) = \lfloor \log_2(n) \rfloor$. In order to simplify the notations, we shall set $q_0(n) = 0$.

Notice that, for every $M \in \mathbb{N}_{\geq 2}$ and for every natural number $1 \leq p < 2^{q_1(M)}$,

$$M + p = \sum_{i=1}^{s_p} 2^{q_i(p)} + \sum_{i=1}^{s_M} 2^{q_i(M)}$$

is the binary decomposition of $M + p$, with $0 \leq q_1(p) < \dots < q_{s_p}(p) < q_1(M) < \dots < q_{s_M}(M)$, and hence $s_{M+p} = s_p + s_M$ and

$$q_j(M + p) = \begin{cases} q_j(p) & \text{for } j = 1, \dots, s_p \\ q_{j-s_p}(M) & \text{for } j = s_p + 1, \dots, s_{M+p} \end{cases} \quad (5)$$

For every $n \in \mathbb{N}$ and $i = 1, \dots, s_n$, let

$$M_i(n) = \sum_{j=i}^{s_n} 2^{q_j(n)} = 2^{q_i(n)} \left\lfloor \frac{n}{2^{q_i(n)}} \right\rfloor.$$

2. For every $n \in \mathbb{N}$, let $\varphi_0(n) = \lfloor n/2 \rfloor$ and $\varphi_1(n) = \lceil n/2 \rceil$ and, for every $m \geq 1$ and for every sequence $b_m \dots b_0 \in \{0, 1\}^{m+1}$, let

$$\varphi_{b_m \dots b_0}(n) = \varphi_{b_m}(\varphi_{b_{m-1} \dots b_0}(n)).$$

By Thompson's Rounding Lemma [36], for every sequence $b_m \dots b_0 \in \{0, 1\}^{m+1}$,

$$\varphi_{b_m \dots b_0}(n) = \left\lfloor \frac{n + \sum_{i=0}^m b_i 2^i}{2^{m+1}} \right\rfloor. \quad (6)$$

3. For every $m \in \mathbb{N}$, let B_m denote the m -th Bernoulli number of the first kind. Recall that these Bernoulli numbers can be defined, starting with $B_0 = 1$, by means of the recurrence

$$\sum_{k=0}^m \binom{m+1}{k} B_k = 0, \quad m \geq 1. \quad (7)$$

Let moreover $B_m(x) \in \mathbb{Q}[x]$ be the Bernoulli polynomial of degree m , defined by

$$B_m(x) = \sum_{k=0}^m \binom{m}{k} B_k x^{m-k}.$$

We list below several well-known properties of the Bernoulli numbers and polynomials that we shall use in our proofs, frequently without any further notice. For these and other properties of the Bernoulli numbers and polynomials, see for instance [1, Ch. 23].

$$B_1 = -1/2 \text{ and } B_{2i+1} = 0 \text{ for every } i \geq 1 \quad (8)$$

$$B_m = B_m(0) = (-1)^m B_m(1) \quad (9)$$

$$B_m(x+y) = \sum_{k=0}^m \binom{m}{k} B_k(x) y^{m-k} \quad (10)$$

In particular,

$$B_m(2) = B_m(1+1) = \sum_{k=0}^m \binom{m}{k} B_k(1) \quad (\text{by (10)})$$

$$= \sum_{k=0}^m \binom{m}{k} (-1)^k B_k \quad (\text{by (9)})$$

$$= \sum_{k=0}^m \binom{m}{k} B_k - 2mB_1 = \sum_{k=0}^m \binom{m}{k} B_k + m \quad (\text{by (8)})$$

$$= B_m(1) + m = (-1)^m B_m + m \quad (\text{again by (9)}) \quad (11)$$

4. For every $d \in \mathbb{N}$, $n \in \mathbb{N}_{\geq 1}$, and $x \in \mathbb{R} \setminus \{0\}$, let

$$T(d, n, x) = \sum_{k=0}^{n-1} k^d x^k$$

The value of $T(d, n, 1)$ is given by *Faulhaber's formula* [19, (6.78)]: for every $d \in \mathbb{N}$ and $n \in \mathbb{N}_{\geq 1}$,

$$\sum_{k=0}^{n-1} k^d = \frac{1}{d+1} \sum_{j=0}^d \binom{d+1}{j} B_j n^{d+1-j}$$

We shall also use this formula in the following, equivalent way:

$$\sum_{k=1}^{n-1} k^d = \frac{1}{d+1} \left(\sum_{j=0}^{d+1} \binom{d+1}{j} B_j n^{d+1-j} + (-1)^d B_{d+1} \right) \quad (12)$$

The equivalence stems from the fact that if $d \geq 1$, $(-1)^d B_{d+1} = -B_{d+1}$, while when $d = 0$, $(-1)^d B_{d+1} = B_1 = -1/2$.

When $x \neq 1$, the double sequence T satisfies the recurrence

$$T(d, n, x) = \frac{x}{1-x} \sum_{p=0}^{d-1} \binom{d}{p} T(p, n, x) - \frac{x^n}{1-x} n^d, \quad d \geq 1.$$

In particular, when $x \neq 1$,

$$T(1, n, x) = \frac{x}{1-x} (T(0, n, x) - nx^{n-1}) \quad (13)$$

We shall use these double sequences $T(d, n, x)$ in order to unify some notations and proofs, but, as we have already encountered in the statement of our main result, we actually only need to know closed formulas for them when $d = 0, 1$:

$$\begin{aligned} T(0, n, 1) &= n, & T(0, n, x) &= \frac{x^n - 1}{x - 1} \quad \text{for } x \neq 1 \\ T(1, n, 1) &= \binom{n}{2}, & T(1, n, x) &= \frac{nx^n(x-1) - x(x^n-1)}{(x-1)^2} \quad \text{for } x \neq 1 \end{aligned} \quad (14)$$

4.2 Some technical lemmas

We begin by providing closed formulas for several sums that we shall often encounter in our computations.

Lemma 4. *For every $d \in \mathbb{N}$, $n \in \mathbb{N}_{\geq 1}$, and $x \in \mathbb{R} \setminus \{0\}$:*

$$\sum_{k=1}^{n-1} q_{s_k}(k)^d x^{q_{s_k}(k)} = T(d, q_{s_n}(n), 2x) + nq_{s_n}(n)^d x^{q_{s_n}(n)} - q_{s_n}(n)^d (2x)^{q_{s_n}(n)}$$

Proof.

$$\begin{aligned} \sum_{k=1}^{n-1} q_{s_k}(k)^d x^{q_{s_k}(k)} &= \sum_{k=1}^{2^{q_{s_n}(n)}-1} q_{s_k}(k)^d x^{q_{s_k}(k)} + \sum_{k=2^{q_{s_n}(n)}}^{n-1} q_{s_k}(k)^d x^{q_{s_k}(k)} \\ &= \sum_{j=0}^{q_{s_n}(n)-1} j^d 2^j x^j + (n - 2^{q_{s_n}(n)}) q_{s_n}(n)^d x^{q_{s_n}(n)} \end{aligned}$$

□

In particular:

- If $x \neq 1/2$,

$$\begin{aligned} \sum_{k=1}^{n-1} x^{q_{s_k}(k)} &= \frac{(2x)^{q_{s_n}(n)} - 1}{2x - 1} + n \cdot x^{q_{s_n}(n)} - (2x)^{q_{s_n}(n)} \\ \sum_{k=1}^{n-1} q_{s_k}(k) x^{q_{s_k}(k)} &= \frac{q_{s_n}(n)(2x)^{q_{s_n}(n)}}{2x - 1} - \frac{2x((2x)^{q_{s_n}(n)} - 1)}{(2x - 1)^2} + nq_{s_n}(n)x^{q_{s_n}(n)} - q_{s_n}(n)(2x)^{q_{s_n}(n)} \end{aligned}$$

- If $x = 1/2$,

$$\begin{aligned} \sum_{k=1}^{n-1} x^{q_{s_k}(k)} &= q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1 \\ \sum_{k=1}^{n-1} q_{s_k}(k) x^{q_{s_k}(k)} &= \frac{q_{s_n}(n)(q_{s_n}(n) - 3)}{2} + nq_{s_n}(n)2^{-q_{s_n}(n)} \end{aligned}$$

Lemma 5. For every $d \in \mathbb{N}$, $n \in \mathbb{N}_{\geq 1}$, and $x \in \mathbb{R} \setminus \{0\}$,

$$\sum_{k=0}^{n-1} \sum_{i=1}^{s_k} q_i(k)^d x^{q_i(k)} = \sum_{i=1}^{s_n} 2^{q_i(n)-1} T(d, q_i(n), x) + \sum_{i=1}^{s_n} q_i(n)^d x^{q_i(n)} (n - M_i(n))$$

Proof. Let $x \in \mathbb{R} \setminus \{0\}$. For every $d \in \mathbb{N}$ and $n \in \mathbb{N}_{\geq 1}$, set

$$a_n^{(d)} := \sum_{k=0}^{n-1} \sum_{i=1}^{s_k} q_i(k)^d x^{q_i(k)}.$$

Then,

$$\begin{aligned} a_n^{(d)} &= \sum_{k=0}^{M_2(n)-1} \sum_{i=1}^{s_k} q_i(k)^d x^{q_i(k)} + \sum_{k=M_2(n)}^{M_2(n)+2^{q_1(n)}-1} \sum_{i=1}^{s_k} q_i(k)^d x^{q_i(k)} \\ &= a_{M_2(n)}^{(d)} + \sum_{p=0}^{2^{q_1(n)}-1} \sum_{i=1}^{s_{M_2(n)+p}} q_i(M_2(n) + p)^d x^{q_i(M_2(n)+p)} \\ &= a_{M_2(n)}^{(d)} + \sum_{p=0}^{2^{q_1(n)}-1} \left(\sum_{i=1}^{s_p} q_i(p)^d x^{q_i(p)} + \sum_{i=1}^{s_{M_2(n)}} q_i(M_2(n))^d x^{q_i(M_2(n))} \right) \quad (\text{by (5)}) \\ &= a_{M_2(n)}^{(d)} + \sum_{p=0}^{2^{q_1(n)}-1} \sum_{i=1}^{s_p} q_i(p)^d x^{q_i(p)} + 2^{q_1(n)} \sum_{i=2}^{s_n} q_i(n)^d x^{q_i(n)} \\ &= a_{M_2(n)}^{(d)} + a_{2^{q_1(n)}}^{(d)} + 2^{q_1(n)} \sum_{i=2}^{s_n} q_i(n)^d x^{q_i(n)}. \end{aligned}$$

So, a simple argument by induction shows that, for every $2 \leq r \leq s_n$

$$a_n^{(d)} = a_{M_r(n)}^{(d)} + \sum_{j=1}^{r-1} a_{2^{q_j(n)}}^{(d)} + \sum_{j=1}^{r-1} \left(2^{q_j(n)} \sum_{i=j+1}^{s_n} q_i(n)^d x^{q_i(n)} \right).$$

Taking $r = s_n$, we obtain

$$\begin{aligned}
a_n^{(d)} &= \sum_{j=1}^{s_n} a_{2^{q_j(n)}}^{(d)} + \sum_{j=1}^{s_n-1} \left(2^{q_j(n)} \sum_{i=j+1}^{s_n} q_i(n)^d x^{q_i(n)} \right) \\
&= \sum_{j=1}^{s_n} a_{2^{q_j(n)}}^{(d)} + \sum_{i=2}^{s_n} \left(q_i(n)^d x^{q_i(n)} \sum_{j=1}^{i-1} 2^{q_j(n)} \right) \\
&= \sum_{j=1}^{s_n} a_{2^{q_j(n)}}^{(d)} + \sum_{i=2}^{s_n} q_i(n)^d x^{q_i(n)} (n - M_i(n))
\end{aligned} \tag{15}$$

Now, for every $l \in \mathbb{N}$, we have that

$$\begin{aligned}
a_{2^l}^{(d)} &= \sum_{k=0}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d x^{q_i(k)} = \sum_{t=0}^{l-1} t^d \cdot x^t \cdot \#\{(k, i) \mid 0 \leq k \leq 2^l - 1, q_i(k) = t\} \\
&= 2^{l-1} \sum_{t=0}^{l-1} t^d \cdot x^t = 2^{l-1} T(d, l, x)
\end{aligned} \tag{16}$$

because each $\#\{(k, i) \mid 0 \leq k \leq 2^l - 1, q_i(k) = t\}$ is simply the number of binary words $w \in \{0, 1\}^l$ with an 1 in the position $t + 1$ starting from the right, which is 2^{l-1} .

Using this expression in Eqn. (15), and taking into account that $M_1(n) = n$, we obtain the formula in the statement. \square

In particular:

- If $x \neq 1$,

$$\sum_{k=0}^{n-1} \sum_{i=1}^{s_k} x^{q_i(k)} = \sum_{i=1}^{s_n} x^{q_i(n)} (n - M_i(n)) + \sum_{i=1}^{s_n} 2^{q_i(n)} \cdot \frac{x^{q_i(n)} - 1}{2(x-1)} \tag{17}$$

$$\sum_{k=0}^{n-1} \sum_{j=1}^{s_k} q_j(k) x^{q_j(k)} = \sum_{i=1}^{s_n} q_i(n) x^{q_i(n)} (n - M_i(n)) + \sum_{i=1}^{s_n} 2^{q_i(n)} \cdot \frac{(x-1)q_i(n)x^{q_i(n)} - x^{q_i(n)+1} + x}{2(x-1)^2} \tag{18}$$

- When $x = 1$,

$$\sum_{k=0}^{n-1} s_k = \sum_{k=1}^{n-1} \sum_{i=1}^{s_k} 1^{q_i(k)} = \sum_{i=1}^{s_n} 2^{q_i(n)-1} (q_i(n) + 2(s_n - i)) \tag{19}$$

$$\sum_{k=1}^{n-1} \sum_{j=1}^{s_k} q_j(k) = \sum_{i=1}^{s_n} 2^{q_i(n)-1} \left(\binom{q_i(n)}{2} + 2 \sum_{j=i+1}^{s_n} q_j(n) \right) \tag{20}$$

Lemma 6. Let $(z_{l,p})_{(l,p) \in \mathbb{N}^2}$ be a double sequence satisfying:

(a) For every $p \in \mathbb{N}$, $z_{0,p} = 0$

(b) For every $l, p > 0$,

$$z_{l,p} = 2z_{l-1,p} + \sum_{q=0}^{p-1} \binom{p}{q} 2^{(p-q)(l-1)} z_{l-1,q}$$

Then, for every $l \geq 0$ and for every $p > 0$,

$$z_{l,p} = -\frac{2^l}{p+1} \sum_{j=1}^{l-1} \left(\sum_{i=1}^p \binom{p+1}{i} \right) (2^i - 1) 2^{(p-i)l + (i-1)j} B_i \Big) z_{j,0}.$$

Proof. Let us iterate several times the recurrence defining our double sequence:

$$\begin{aligned}
z_{l,p} &= 2z_{l-1,p} + \sum_{q=0}^{p-1} \binom{p}{q} 2^{(p-q)(l-1)} z_{l-1,q} \\
&= 2 \left(2z_{l-2,p} + \sum_{q=0}^{p-1} \binom{p}{q} 2^{(p-q)(l-2)} z_{l-2,q} \right) + \sum_{q=0}^{p-1} \binom{p}{q} 2^{(p-q)(l-1)} z_{l-1,q} \\
&= 2^2 z_{l-2,p} + \sum_{q=0}^{p-1} \binom{p}{q} \left(2^{(p-q)(l-2)+1} z_{l-2,q} + 2^{(p-q)(l-1)} z_{l-1,q} \right) \\
&= 2^2 \left(2z_{l-3,p} + \sum_{q=0}^{p-1} \binom{p}{q} 2^{(p-q)(l-3)} z_{l-3,q} \right) + \sum_{q=0}^{p-1} \binom{p}{q} \left(2^{(p-q)(l-2)+1} z_{l-2,q} + 2^{(p-q)(l-1)} z_{l-1,q} \right) \\
&= 2^3 z_{l-3,p} + \sum_{q=0}^{p-1} \binom{p}{q} \left(2^{(p-q)(l-3)+2} z_{l-3,q} + 2^{(p-q)(l-2)+1} z_{l-2,q} + 2^{(p-q)(l-1)} z_{l-1,q} \right)
\end{aligned}$$

Reasoning in this way, and recalling that $z_{0,q} = 0$ for every q , it is easy to prove by induction on l that

$$z_{l,p} = \sum_{q=0}^{p-1} \binom{p}{q} \left(\sum_{j=1}^{l-1} 2^{(p-q)j+(l-1-j)} z_{j,q} \right) = 2^{l-1} \sum_{q=0}^{p-1} \binom{p}{q} \left(\sum_{j=1}^{l-1} 2^{(p-q-1)j} z_{j,q} \right) \quad (21)$$

We have computed explicitly $z_{l,p}$ for several $p \geq 1$ using this recurrence, in order to look for a

pattern. The results have been:

$$\begin{aligned}
z_{l,2} &= 2^{l-1} \sum_{j=1}^{l-1} (2^l - 2 \cdot 2^{j-1}) z_{j,0} \\
z_{l,3} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{2l} - 3 \cdot 2^{l+j-1}) z_{j,0} \\
z_{l,4} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{3l} - 4 \cdot 2^{2l+j-1} + 8 \cdot 2^{3(j-1)}) z_{j,0} \\
z_{l,5} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{4l} - 5 \cdot 2^{3l+j-1} + 20 \cdot 2^{l+3(j-1)}) z_{j,0} \\
z_{l,6} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{5l} - 6 \cdot 2^{4l+j-1} + 40 \cdot 2^{2l+3(j-1)} - 96 \cdot 2^{5(j-1)}) z_{j,0} \\
z_{l,7} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{6l} - 7 \cdot 2^{5l+j-1} + 70 \cdot 2^{3l+3(j-1)} - 336 \cdot 2^{l+5(j-1)}) z_{j,0} \\
z_{l,8} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{7l} - 8 \cdot 2^{6l+j-1} + 112 \cdot 2^{4l+3(j-1)} - 896 \cdot 2^{2l+5(j-1)} + 2176 \cdot 2^{7(j-1)}) z_{j,0} \\
z_{l,9} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{8l} - 9 \cdot 2^{7l+j-1} + 168 \cdot 2^{5l+3(j-1)} - 2016 \cdot 2^{3l+5(j-1)} + 9792 \cdot 2^{l+7(j-1)}) z_{j,0} \\
z_{l,10} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{9l} - 10 \cdot 2^{8l+j-1} + 240 \cdot 2^{6l+3(j-1)} - 4032 \cdot 2^{4l+5(j-1)} + 32640 \cdot 2^{2l+7(j-1)} \\
&\quad - 79360 \cdot 2^{9(j-1)}) z_{j,0} \\
z_{l,11} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{10l} - 11 \cdot 2^{9l+j-1} + 330 \cdot 2^{7l+3(j-1)} - 7392 \cdot 2^{5l+5(j-1)} + 89760 \cdot 2^{3l+7(j-1)} \\
&\quad - 436480 \cdot 2^{l+9(j-1)}) z_{j,0}, \\
z_{l,12} &= 2^{l-1} \sum_{j=1}^{l-1} (2^{11l} - 12 \cdot 2^{10l+j-1} + 440 \cdot 2^{8l+3(j-1)} - 12672 \cdot 2^{6l+5(j-1)} + 215424 \cdot 2^{4l+7(j-1)} \\
&\quad - 1745920 \cdot 2^{2l+9(j-1)} + 4245504 \cdot 2^{11(j-1)}) z_{j,0}.
\end{aligned}$$

These results hint that

$$z_{l,p} = 2^{l-1} \sum_{j=1}^{l-1} \left(2^{(p-1)l} + \sum_{i=1}^{p-1} a_i^{(p)} \binom{p}{i} 2^{(p-1-i)l+i(j-1)} \right) z_{j,0}.$$

with $a_{2i}^{(p)} = 0$ if $i > 0$,

$$a_1^{(p)} = -1, a_3^{(p)} = 2, a_5^{(p)} = -16, a_7^{(p)} = 272, a_9^{(p)} = -7936, \dots$$

These values are consistent with

$$a_i^{(p)} = -\frac{1}{i+1} \cdot 2^{i+1} (2^{i+1} - 1) B_{i+1}, \quad \text{for } 1 \leq i \leq p-1$$

So, this leads us to conjecture that, if $p > 0$,

$$\begin{aligned}
z_{l,p} &= 2^{l-1} \sum_{j=1}^{l-1} \left(2^{(p-1)l} - \sum_{i=1}^{p-1} \frac{1}{i+1} \cdot 2^{i+1} (2^{i+1} - 1) B_{i+1} \binom{p}{i} \right) 2^{(p-1-i)l+i(j-1)} z_{j,0} \\
&= 2^{l-1} \sum_{j=1}^{l-1} \left(2^{(p-1)l} - \sum_{i=1}^{p-1} \frac{1}{p+1} \cdot 2^{i+1} (2^{i+1} - 1) B_{i+1} \binom{p+1}{i+1} \right) 2^{(p-1-i)l+i(j-1)} z_{j,0} \\
&= 2^{l-1} \sum_{j=1}^{l-1} \left(2^{(p-1)l} - \sum_{i=2}^p \frac{1}{p+1} \cdot 2^i (2^i - 1) B_i \binom{p+1}{i} \right) 2^{(p-i)l+(i-1)(j-1)} z_{j,0} \\
&= 2^{l-1} \sum_{j=1}^{l-1} \left(2^{(p-1)l} - \sum_{i=1}^p \frac{1}{p+1} \cdot 2^i (2^i - 1) B_i \binom{p+1}{i} \right) 2^{(p-i)l+(i-1)(j-1)} \\
&\quad + \frac{1}{p+1} \cdot 2 B_1 (p+1) 2^{(p-1)l} z_{j,0} \\
&= 2^{l-1} \sum_{j=1}^{l-1} \left(2^{(p-1)l} - \sum_{i=1}^p \frac{1}{p+1} \cdot 2^i (2^i - 1) B_i \binom{p+1}{i} \right) 2^{(p-i)l+(i-1)(j-1)} - 2^{(p-1)l} z_{j,0} \\
&= -\frac{2^l}{p+1} \sum_{j=1}^{l-1} \left(\sum_{i=1}^p (2^i - 1) \binom{p+1}{i} \right) 2^{(p-i)l+(i-1)j} B_i z_{j,0} \tag{22}
\end{aligned}$$

and we proceed to prove this equality by induction on l .

The case when $l = 1$ is true because the right hand sum of (22) is empty and, by (21), $z_{1,p} = 0$. Assume now that the equality (22) is true for $z_{1,p}, \dots, z_{l-1,p}$ and every $p > 0$, and let us prove it for $z_{l,p}$. By (21) and the induction hypothesis,

$$\begin{aligned}
2^{1-l} z_{l,p} &= \sum_{q=0}^{p-1} \binom{p}{q} \sum_{k=1}^{l-1} 2^{(p-q-1)k} z_{k,q} \\
&= \sum_{k=1}^{l-1} 2^{(p-1)k} z_{k,0} \\
&\quad - \sum_{q=1}^{p-1} \binom{p}{q} \sum_{k=1}^{l-1} 2^{(p-q-1)k} \cdot 2^{k-1} \sum_{j=1}^{k-1} \left(\sum_{i=1}^q \frac{2^i (2^i - 1) B_i}{q+1} \binom{q+1}{i} \right) 2^{(q-i)k+(i-1)(j-1)} z_{j,0} \\
&\text{(by the induction hypothesis)} \\
&= \sum_{j=1}^{l-1} 2^{(p-1)j} z_{j,0} \\
&\quad - \sum_{j=1}^{l-2} \left[\sum_{q=1}^{p-1} \binom{p}{q} \sum_{k=j+1}^{l-1} 2^{(p-q)k-1} \sum_{i=1}^q \frac{2^i (2^i - 1) B_i}{q+1} \binom{q+1}{i} \right] 2^{(q-i)k+(i-1)(j-1)} z_{j,0} \\
&= \sum_{j=1}^{l-1} \left[2^{(p-1)j} - \sum_{q=1}^{p-1} \binom{p}{q} \sum_{k=j+1}^{l-1} 2^{(p-q)k-1} \sum_{i=1}^q \frac{2^i (2^i - 1) B_i}{q+1} \binom{q+1}{i} \right] 2^{(q-i)k+(i-1)(j-1)} z_{j,0}
\end{aligned}$$

and then the coefficient of $z_{j,0}$, for $j \leq l-1$, in $2^{-l+1} z_{l,p}$, which we want to prove to be

$$-\frac{2}{p+1} \sum_{i=1}^p (2^i - 1) B_i \binom{p+1}{i} 2^{(p-i)l+(i-1)j}$$

is

$$\begin{aligned}
& 2^{(p-1)j} - \sum_{q=1}^{p-1} \binom{p}{q} \sum_{k=j+1}^{l-1} 2^{(p-q)k-1} \sum_{i=1}^q \frac{2^i(2^i-1)B_i}{q+1} \binom{q+1}{i} 2^{(q-i)k+(i-1)(j-1)} \\
&= 2^{(p-1)j} - \sum_{i=1}^{p-1} \sum_{q=i}^{p-1} \sum_{k=j+1}^{l-1} \frac{(2^i-1)B_i}{q+1} \binom{p}{q} \binom{q+1}{i} 2^{(p-i)k+(i-1)j} \\
&= 2^{(p-1)j} - \sum_{i=1}^{p-1} \sum_{q=i}^{p-1} \sum_{k=j+1}^{l-1} \frac{(2^i-1)B_i}{p+1} \binom{p+1}{q+1} \binom{q+1}{i} 2^{(p-i)k+(i-1)j} \\
&= 2^{(p-1)j} - \sum_{i=1}^{p-1} \sum_{q=i}^{p-1} \sum_{k=j+1}^{l-1} \frac{(2^i-1)B_i}{p+1} \binom{p+1}{i} \binom{p+1-i}{q+1-i} 2^{(p-i)k+(i-1)j} \\
&= 2^{(p-1)j} - \sum_{i=1}^{p-1} \frac{2^{(i-1)j}(2^i-1)B_i}{p+1} \binom{p+1}{i} \left(\sum_{q=i}^{p-1} \binom{p+1-i}{q+1-i} \right) \left(\sum_{k=j+1}^{l-1} 2^{(p-i)k} \right) \\
&= 2^{(p-1)j} - \sum_{i=1}^{p-1} \frac{2^{(i-1)j}(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(2^{p-i}-1)} \frac{2^{(p-i)l} - 2^{(p-i)(j+1)}}{2^{p-i} - 1} \\
&= 2^{(p-1)j} - \sum_{i=1}^{p-1} \frac{(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(i-1)j+1} (2^{(p-i)l} - 2^{(p-i)(j+1)}) \\
&= 2^{(p-1)j} - \sum_{i=1}^{p-1} \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(p-i)l+(i-1)j} + \sum_{i=1}^{p-1} \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{p-i+(p-1)j} \\
&= 2^{(p-1)j} - \sum_{i=0}^p \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(p-i)l+(i-1)j} + \sum_{i=0}^p \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{p-i+(p-1)j} \\
&= 2^{(p-1)j} - \sum_{i=0}^p \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(p-i)l+(i-1)j} + \sum_{i=0}^p \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{p-i+(p-1)j} \\
&= 2^{(p-1)j} - \sum_{i=0}^p \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(p-i)l+(i-1)j} + \frac{2^{p+1+(p-1)j}}{p+1} \sum_{i=0}^p \binom{p+1}{i} B_i \\
&\quad - \frac{2^{(p-1)j}}{p+1} \sum_{i=0}^p \binom{p+1}{i} 2^{p+1-i} B_i \\
&= 2^{(p-1)j} - \sum_{i=0}^p \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(p-i)l+(i-1)j} - \frac{2^{(p-1)j}}{p+1} (B_{p+1}(2) - B_{p+1}) \\
&= - \sum_{i=0}^p \frac{2(2^i-1)B_i}{p+1} \binom{p+1}{i} 2^{(p-i)l+(i-1)j} \quad (\text{by Eqn. (11)})
\end{aligned}$$

as we wanted to prove. \square

For every $l, m, p, d \in \mathbb{N}$, with $p < m$, and for every given $a \in \mathbb{R} \setminus \{0\}$, let

$$\gamma_l^{(d,p,m)} = \sum_{k=1}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d (2^{-m}a)^{q_i(k)} M_{i+1}(k)^p$$

In the last part of our computations we shall use an alternative expression for $\gamma_l^{(d,p,m)}$ which we derive now as an application of the last lemma.

We have that $\gamma_0^{(d,p,m)} = 0$ by definition, and, by Eqn. (16),

$$\gamma_l^{(d,0,m)} = \sum_{k=1}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d (a2^{-m})^{q_i(k)} = 2^{l-1} T(d, l, a2^{-m}) \quad (23)$$

The case when $l, p > 0$ is covered by the next lemma.

Lemma 7. *If $l, p > 0$, then*

$$\gamma_l^{(d,p,m)} = \frac{a^{l-1} 2^{-(m-1)(l-1)+pl}}{p+1} \sum_{t=1}^{l-1} (l-t-1)^d (a^{-1} 2^{m-p-1})^t (B_{p+1}(2^t) - B_{p+1})$$

Proof. If $l, p > 0$,

$$\begin{aligned} \gamma_l^{(d,p,m)} &= \gamma_{l-1}^{(d,p,m)} + \sum_{k=2^{l-1}}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d (a 2^{-m})^{q_i(k)} M_{i+1}(k)^p \\ &= \gamma_{l-1}^{(d,p,m)} + \sum_{k=2^{l-1}}^{2^l-1} \sum_{i=1}^{s_k-1} q_i(k)^d (a 2^{-m})^{q_i(k)} \left(M_{i+1}(k - 2^{l-1}) + 2^{l-1} \right)^p \\ &= \gamma_{l-1}^{(d,p,m)} + \sum_{k=0}^{2^{l-1}-1} \sum_{i=1}^{s_k} q_i(k)^d (a 2^{-m})^{q_i(k)} \left(M_{i+1}(k) + 2^{l-1} \right)^p \\ &= \gamma_{l-1}^{(d,p,m)} + \sum_{k=1}^{2^{l-1}-1} \sum_{i=1}^{s_k} q_i(k)^d (a 2^{-m})^{q_i(k)} \sum_{q=0}^p \binom{p}{q} M_{i+1}(k)^q 2^{(p-q)(l-1)} \\ &= \gamma_{l-1}^{(d,p,m)} + \sum_{q=0}^p \binom{p}{q} 2^{(p-q)(l-1)} \sum_{k=1}^{2^{l-1}-1} \sum_{i=1}^{s_k} q_i(k)^d (a 2^{-m})^{q_i(k)} M_{i+1}(k)^q \\ &= \gamma_{l-1}^{(d,p,m)} + \sum_{q=0}^p \binom{p}{q} 2^{(p-q)(l-1)} \gamma_{l-1}^{(d,q,m)} = 2 \gamma_{l-1}^{(d,p,m)} + \sum_{q=0}^{p-1} \binom{p}{q} 2^{(p-q)(l-1)} \gamma_{l-1}^{(d,q,m)} \end{aligned}$$

Then, by Lemma 6 and Eqn. (23),

$$\begin{aligned}
\gamma_i^{(d,p,m)} &= -\frac{2^l}{p+1} \sum_{j=1}^{l-1} \left(\sum_{i=1}^p \binom{p+1}{i} (2^i - 1) 2^{(p-i)l+(i-1)j} B_i \right) \gamma_j^{(d,0,m)} \\
&= -\frac{2^l}{p+1} \sum_{j=1}^{l-1} \left[\sum_{i=1}^p \binom{p+1}{i} (2^i - 1) 2^{(p-i)l+(i-1)j} B_i 2^{j-1} \sum_{t=0}^{j-1} t^d (a2^{-m})^t \right] \\
&= -\frac{2^{pl+l-1}}{p+1} \sum_{i=1}^p \left[\binom{p+1}{i} (2^i - 1) 2^{-il} \left(\sum_{t=0}^{l-2} t^d (a2^{-m})^t \sum_{j=t+1}^{l-1} 2^{ij} \right) \right] B_i \\
&= -\frac{2^{pl+l-1}}{p+1} \sum_{i=1}^p \left[\binom{p+1}{i} (2^i - 1) 2^{-il} \left(\sum_{t=0}^{l-2} t^d (a2^{-m})^t \cdot \frac{2^{il} - 2^{i(t+1)}}{2^i - 1} \right) \right] B_i \\
&= -\frac{2^{pl+l-1}}{p+1} \sum_{i=1}^p \binom{p+1}{i} B_i \cdot \left(\sum_{t=0}^{l-2} t^d (a2^{-m})^t \right) \\
&\quad + \frac{2^{pl+l-1}}{p+1} \sum_{i=1}^p \binom{p+1}{i} 2^{-i(l-1)} \left(\sum_{t=0}^{l-2} t^d (a2^{i-m})^t \right) B_i \\
&= \frac{2^{pl+l-1}}{p+1} B_0 \cdot \left(\sum_{t=0}^{l-2} t^d (a2^{-m})^t \right) + \frac{2^{pl+l-1}}{p+1} \sum_{i=1}^p \binom{p+1}{i} 2^{-i(l-1)} \left(\sum_{t=0}^{l-2} t^d (a2^{i-m})^t \right) B_i \\
&= \frac{2^{pl+l-1}}{p+1} \sum_{i=0}^p \binom{p+1}{i} 2^{-i(l-1)} \left(\sum_{t=0}^{l-2} t^d (a2^{i-m})^t \right) B_i \\
&= \frac{2^{pl+l-1}}{p+1} \sum_{i=0}^p \binom{p+1}{i} 2^{-i(l-1)} \left(\sum_{t=1}^{l-1} (l-1-t)^d (a2^{i-m})^{l-1-t} \right) B_i \\
&= \frac{a^{l-1} 2^{-(m-1)(l-1)+pl}}{p+1} \sum_{t=1}^{l-1} (l-t-1)^d (a^{-1} 2^{m-p-1})^t \left(\sum_{i=0}^p \binom{p+1}{i} 2^{(p+1-i)t} B_i \right) \\
&= \frac{a^{l-1} 2^{-(m-1)(l-1)+pl}}{p+1} \sum_{t=1}^{l-1} (l-t-1)^d (a^{-1} 2^{m-p-1})^t (B_{p+1}(2^t) - B_{p+1})
\end{aligned}$$

as we claimed. □

Corollary 8. For every $l, m, p, d \in \mathbb{N}$, with $p < m$,

$$\begin{aligned}
\gamma_l^{(d,p,m)} &= \frac{a^{l-1} 2^{-(m-1)(l-1)+pl}}{p+1} \sum_{t=1}^{l-1} (l-t-1)^d (a^{-1} 2^{m-p-1})^t (B_{p+1}(2^t) - B_{p+1}) \\
&\quad + (a2^{-(m-1)})^{l-1} (l-1)^d \cdot \delta_{p=0, l>0}
\end{aligned}$$

where $\delta_{p=0, l>0} = 1$ if $p = 0$ and $l > 0$, and $\delta_{p=0, l>0} = 0$ otherwise.

Proof. When $l = 0$, both sides of the equality in the statement are equal to 0, and the last lemma

covers the case when $p > 0$ and $l > 0$. Finally, when $p = 0$ and $l > 0$, by (23),

$$\begin{aligned}
\gamma_l^{(d,0,m)} &= 2^{l-1} \sum_{t=0}^{l-1} t^d (a2^{-m})^t = 2^{l-1} \sum_{t=0}^{l-1} (l-1-t)^d (a2^{-m})^{l-1-t} \\
&= a^{l-1} 2^{-(m-1)(l-1)} \left(\sum_{t=1}^{l-1} (l-1-t)^d (a^{-1}2^m)^t + (l-1)^d \right) \\
&= a^{l-1} 2^{-(m-1)(l-1)} \left(\sum_{t=1}^{l-1} (l-1-t)^d (a^{-1}2^{m-1})^t (B_1(2^t) - B_1) + (l-1)^d \right) \\
&= \frac{a^{l-1} 2^{-(m-1)(l-1)+0 \cdot l}}{0+1} \sum_{t=1}^{l-1} (l-t-1)^d (a^{-1}2^{m-0-1})^t (B_{0+1}(2^t) - B_{0+1}) + (a2^{-(m-1)})^{l-1} (l-1)^d
\end{aligned}$$

□

4.3 Statement of the problem

Let $P(x, y) = \sum_{r,t \geq 0} b_{r,t} x^r y^t \in \mathbb{R}[x, y]$ be a bivariate polynomial and $a \in \mathbb{R} \setminus \{0\}$, and consider the recurrence equation

$$x_n = a \cdot x_{\lceil n/2 \rceil} + a \cdot x_{\lfloor n/2 \rfloor} + P(\lceil n/2 \rceil, \lfloor n/2 \rfloor), \quad n \geq 2. \quad (24)$$

Lemma 9. *If $(x_n^0)_n$ is the solution of (24) with initial condition $x_1^0 = 0$, then*

$$\bar{x}_n = x_n^0 + ((2a)^{q_{s_n}(n)} + (2a-1)(na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)}))x_1$$

is its solution with initial condition x_1

Proof. Since $\bar{x}_1 = x_1$, we must prove that $(\bar{x}_n)_n$ satisfies (24). Since $(x_n^0)_n$ already satisfies it, it is enough to check that, for every $n \geq 2$,

$$\begin{aligned}
&(2a)^{q_{s_n}(n)} + (2a-1)(na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)}) \\
&= a((2a)^{q_{s_{\lceil n/2 \rceil}}(\lceil n/2 \rceil)} + (2a-1)(\lceil n/2 \rceil a^{q_{s_{\lceil n/2 \rceil}}(\lceil n/2 \rceil)} - (2a)^{q_{s_{\lceil n/2 \rceil}}(\lceil n/2 \rceil)})) \\
&\quad + a((2a)^{q_{s_{\lfloor n/2 \rfloor}}(\lfloor n/2 \rfloor)} + (2a-1)(\lfloor n/2 \rfloor a^{q_{s_{\lfloor n/2 \rfloor}}(\lfloor n/2 \rfloor)} - (2a)^{q_{s_{\lfloor n/2 \rfloor}}(\lfloor n/2 \rfloor)}))
\end{aligned}$$

The simplest way to do it is by distinguishing two cases; to simplify the notations, we denote in the rest of this proof $q_{s_n}(n)$ by q .

(i) If $n \neq 2^{q+1} - 1$, then $q_{s_{\lceil n/2 \rceil}}(\lceil n/2 \rceil) = q_{s_{\lfloor n/2 \rfloor}}(\lfloor n/2 \rfloor) = q - 1$ and then

$$\begin{aligned}
&a((2a)^{q-1} + (2a-1)(\lceil n/2 \rceil a^{q-1} - (2a)^{q-1}) + (2a)^{q-1} + (2a-1)(\lfloor n/2 \rfloor a^{q-1} - (2a)^{q-1})) \\
&= a(2(2a)^{q-1} + (2a-1)(na^{q-1} - 2(2a)^{q-1})) = (2a)^q + (2a-1)(na^q - (2a)^q)
\end{aligned}$$

(ii) If $n = 2^{q+1} - 1$, then $\lceil n/2 \rceil = 2^q$ and $\lfloor n/2 \rfloor = 2^q - 1$, and then

$$\begin{aligned}
&a((2a)^q + (2a-1)(2^q a^q - (2a)^q) + (2a)^{q-1} + (2a-1)((2^q - 1)a^{q-1} - (2a)^{q-1})) \\
&= 2^{q+1} a^{q+1} - (2a-1)a^q = (2a)^q + (2a-1)((2^{q+1} - 1)a^q - (2a)^q)
\end{aligned}$$

□

Therefore, in the rest of this paper we shall only be concerned with the solution of (24) with initial condition $x_1 = 0$. Now, if, for every $(r, t) \in \mathbb{N}^2$ such that $b_{r,t} \neq 0$, we know the solution $(x_n^{(r,t)})_n$ of the recurrence

$$x_n^{(r,t)} = a \cdot x_{\lceil n/2 \rceil}^{(r,t)} + a \cdot x_{\lfloor n/2 \rfloor}^{(r,t)} + \left\lceil \frac{n}{2} \right\rceil^r \cdot \left\lfloor \frac{n}{2} \right\rfloor^t, \quad n \geq 2, \quad (25)$$

with initial condition $x_1^{(r,t)} = 0$, then the solution of (24) with initial condition $x_1 = 0$ will be

$$x_n = \sum_{r,t} b_{r,t} x_n^{(r,t)}.$$

So, we are finally led to consider, for any $r, t \in \mathbb{N}$, the equation (25). Let $x_n^{(r,t)}$ be its solution with $x_1^{(r,t)} = 0$.

4.4 The sequence of differences of consecutive terms $y_n^{(r,t)}$

Let $y_n^{(r,t)} = x_n^{(r,t)} - x_{n-1}^{(r,t)}$, with $y_1^{(r,t)} = 0$, and

$$x_n^{(r,t)} = \sum_{k=1}^n y_k^{(r,t)}. \quad (26)$$

Notice that

$$y_2^{(r,t)} = x_2^{(r,t)} - x_1^{(r,t)} = a \cdot x_1^{(r,t)} + a \cdot x_1^{(r,t)} + 1 - x_1^{(r,t)} = 1.$$

Let us derive a recurrence for $y_n^{(r,t)}$, for $n \geq 3$. Taking into account the parity of n , we have:

- For every $m \geq 2$,

$$\begin{aligned} y_{2m}^{(r,t)} &= x_{2m}^{(r,t)} - x_{2m-1}^{(r,t)} \\ &= 2a \cdot x_m^{(r,t)} + m^{r+t} - a \cdot x_m^{(r,t)} - a \cdot x_{m-1}^{(r,t)} - m^r (m-1)^t = a \cdot y_m^{(r,t)} + m^{r+t} - m^r (m-1)^t \\ &= a \cdot y_{\lceil (2m)/2 \rceil}^{(r,t)} + \lceil (2m)/2 \rceil^r \lfloor (2m)/2 \rfloor^t - \lfloor (2m)/2 \rfloor^r (\lceil (2m)/2 \rceil - 1)^t \end{aligned}$$

- For every $m \geq 1$,

$$\begin{aligned} y_{2m+1}^{(r,t)} &= x_{2m+1}^{(r,t)} - x_{2m}^{(r,t)} \\ &= a \cdot x_{m+1}^{(r,t)} + a \cdot x_m^{(r,t)} + (m+1)^r m^t - 2a \cdot x_m^{(r,t)} - m^{r+t} = a \cdot y_{m+1}^{(r,t)} + (m+1)^r m^t - m^{r+t} \\ &= a \cdot y_{\lceil (2m+1)/2 \rceil}^{(r,t)} + \lceil (2m+1)/2 \rceil^r \lfloor (2m+1)/2 \rfloor^t - \lfloor (2m+1)/2 \rfloor^r (\lceil (2m+1)/2 \rceil - 1)^t \end{aligned}$$

So, in summary, $y_n^{(r,t)}$ satisfies the recurrence

$$y_n^{(r,t)} = a \cdot y_{\varphi_1(n)}^{(r,t)} + \varphi_1(n)^r \varphi_0(n)^t - \varphi_0(n)^r (\varphi_1(n) - 1)^t, \quad n \geq 3.$$

Therefore, for every $n \geq 3$ and $m \geq 1$,

$$y_n^{(r,t)} = a^m \cdot y_{\underbrace{\varphi_{1 \dots 1}(n)}_m}^{(r,t)} + \sum_{k=0}^{m-1} a^k \left(\varphi_{\underbrace{1 \dots 1}_k}(n)^r \varphi_{\underbrace{0 \dots 1}_k}(n)^t - \varphi_{\underbrace{0 \dots 1}_k}(n)^r (\varphi_{\underbrace{1 \dots 1}_k}(n) - 1)^t \right)$$

where, by Eqn. (6), for every $k \geq 0$

$$\begin{aligned} \varphi_{\underbrace{1 \dots 1}_k}(n) &= \left\lfloor \frac{n + 2^{k+1} - 1}{2^{k+1}} \right\rfloor = \left\lfloor \frac{n-1}{2^{k+1}} \right\rfloor + 1 \\ \varphi_{\underbrace{0 \dots 1}_k}(n) &= \left\lfloor \frac{n + 2^k - 1}{2^{k+1}} \right\rfloor = \left\lfloor \frac{n-1}{2^{k+1}} + \frac{1}{2} \right\rfloor \end{aligned}$$

This implies that, for every $n \geq 3$,

$$y_n^{(r,t)} = a^{L_n} y_2^{(r,t)} + \sum_{k=1}^{L_n} a^{k-1} \left(\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor \right)^r \left[\frac{n-1}{2^k} + \frac{1}{2} \right]^t - \left[\frac{n-1}{2^k} + \frac{1}{2} \right]^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \right)$$

where L_n is such that $2 = \underbrace{\varphi_1 \dots 1}_{L_n}(n) = 1 + \lfloor (n-1)/2^{L_n} \rfloor$ that is, $L_n = q_{s_{n-1}}(n-1)$. Thus, in summary:

$$y_n^{(r,t)} = a^{q_{s_{n-1}}(n-1)} + \sum_{k=1}^{q_{s_{n-1}}(n-1)} a^{k-1} \left[\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor\right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \right] \quad (27)$$

Next result gives an expression for $y_n^{(r,t)}$ that will be more suitable to our purposes. To simplify the notations, in its statement and henceforth, for every $d, m, n \in \mathbb{N}$, we set

$$S_n^{(d,m)} = \sum_{j=1}^{s_n-1} q_j(n)^d (2^{-m}a)^{q_j(n)} M_{j+1}(n)^m.$$

Moreover, we shall let $\ell = \log_2(a) - t$ when $a > 0$, and we shall use the following Kronecker delta:

$$\delta_\ell = \begin{cases} 1 & \text{if } a > 0, r > 0, \text{ and } \ell \in \{0, \dots, r-1\} \\ 0 & \text{otherwise} \end{cases}$$

Proposition 10. *For every $n \geq 2$,*

$$\begin{aligned} y_n^{(r,t)} &= a^{q_{s_{n-1}}(n-1)} + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} ((n-1)^{t+l} - a^{q_{s_{n-1}}(n-1)}) \\ &\quad + \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} \right) S_{n-1}^{(0,i)} \\ &\quad + \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{j=0}^{s_{n-1}-1} M_{j+1}(n-1)^{t+\ell} (q_{j+1}(n-1) - q_j(n-1)) \end{aligned}$$

Proof. Let $n \in \mathbb{N}_{geq 2}$. To simplify the notations, in this proof we denote s_{n-1} by s and each $q_i(n-1)$ and $M_i(n-1)$, $i = 1, \dots, s$, by q_i and M_i , respectively. Recall that $\ell = \log_2(a) - t$ if $a > 0$. This number will play a role in sums of geometric sequences of the form $\sum_{k=1}^m (2^{-(t+l)}a)^k$ with $l \in \mathbb{N}$, which yield two different expressions depending on whether $l = \ell$, that is, $2^{-(t+l)}a = 1$, or not.

It is easy to check that when $n = 2$ the right-hand side in the formula for $y_n^{(r,t)}$ given in the statement is 1, using that $s_1 = 1$ and $q_{s_1}(1) = 0$. Assume now that $n \geq 3$. In this case, by (27),

$$y_n^{(r,t)} = \widehat{y}_n + a^{q_s}$$

where

$$\widehat{y}_n = \sum_{k=1}^{q_s} a^{k-1} \left(\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor\right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \right)$$

Let us compute this sum. To begin with, notice that:

- If $k \leq q_1$,

$$\begin{aligned} &\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor\right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \\ &= \left(1 + \frac{n-1}{2^k}\right)^r \left(\frac{n-1}{2^k}\right)^t - \left(\frac{n-1}{2^k}\right)^r \left(\frac{n-1}{2^k}\right)^t \\ &= (1 + 2^{-k}M_1)^r (2^{-k}M_1)^t - (2^{-k}M_1)^{r+t} = \sum_{l=0}^{r-1} \binom{r}{l} (2^{-k}M_1)^{t+l} \end{aligned}$$

- If $q_j + 1 < k \leq q_{j+1}$, for some $j = 1, \dots, s-1$,

$$\begin{aligned}
& \left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor\right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \\
&= \left(1 + \left\lfloor \sum_{i=1}^s 2^{q_i-k} \right\rfloor\right)^r \left\lfloor \sum_{i=1}^s 2^{q_i-k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \sum_{i=1}^s 2^{q_i-k} + \frac{1}{2} \right\rfloor^r \left\lfloor \sum_{i=1}^s 2^{q_i-k} \right\rfloor^t \\
&= \left(1 + \sum_{i=j+1}^s 2^{q_i-k}\right)^r \left(\sum_{i=j+1}^s 2^{q_i-k}\right)^t - \left(\sum_{i=j+1}^s 2^{q_i-k}\right)^r \left(\sum_{i=j+1}^s 2^{q_i-k}\right)^t \\
&= (1 + 2^{-k} M_{j+1})^r (2^{-k} M_{j+1})^t - (2^{-k} M_{j+1})^{r+t} = \sum_{l=0}^{r-1} \binom{r}{l} (2^{-k} M_{j+1})^{t+l}
\end{aligned}$$

- If $k = q_j + 1$, for some $j = 1, \dots, s-1$,

$$\begin{aligned}
& \left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor\right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \\
&= \left(1 + \left\lfloor \sum_{i=1}^s 2^{q_i-q_j-1} \right\rfloor\right)^r \left\lfloor \sum_{i=1}^s 2^{q_i-q_j-1} + \frac{1}{2} \right\rfloor^t - \left\lfloor \sum_{i=1}^s 2^{q_i-q_j-1} + \frac{1}{2} \right\rfloor^r \left\lfloor \sum_{i=1}^s 2^{q_i-q_j-1} \right\rfloor^t \\
&= \left(1 + \sum_{i=j+1}^s 2^{q_i-q_j-1}\right)^r \left(\sum_{i=j+1}^s 2^{q_i-q_j-1} + 1\right)^t - \left(\sum_{i=j+1}^s 2^{q_i-q_j-1} + 1\right)^r \left(\sum_{i=j+1}^s 2^{q_i-q_j-1}\right)^t \\
&= (1 + 2^{-q_j-1} M_{j+1})^{r+t} - (1 + 2^{-q_j-1} M_{j+1})^r (2^{-q_j-1} M_{j+1})^t \\
&= \sum_{i=0}^{r+t-1} \binom{r+t}{i} (2^{-q_j-1} M_{j+1})^i - \sum_{l=0}^{r-1} \binom{r}{l} (2^{-q_j-1} M_{j+1})^{t+l}
\end{aligned}$$

Then,

$$\begin{aligned}
& \sum_{k=1}^{q_1} a^{k-1} \left(\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor\right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \right) \\
&= \sum_{k=1}^{q_1} a^{k-1} \sum_{l=0}^{r-1} \binom{r}{l} (2^{-k} M_1)^{t+l} = \frac{1}{a} \sum_{l=0}^{r-1} \binom{r}{l} M_1^{t+l} \sum_{k=1}^{q_1} (2^{-(t+l)} a)^k = (*)
\end{aligned}$$

Now, if $\ell \notin \{0, \dots, r-1\}$, then $2^{-(t+l)} a \neq 1$ for every $l = 0, \dots, r-1$ and hence

$$(*) = \sum_{l=0}^{r-1} \binom{r}{l} M_1^{t+l} \cdot \frac{1 - (2^{-(t+l)} a)^{q_1}}{2^{t+l} - a}$$

while if $\ell \in \{0, \dots, r-1\}$, then

$$\begin{aligned}
(*) &= \frac{1}{a} \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \binom{r}{l} M_1^{t+l} \sum_{k=1}^{q_1} (2^{-(t+l)} a)^k + \frac{1}{a} \binom{r}{\ell} M_1^{t+\ell} \sum_{k=1}^{q_1} 1^k \\
&= \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \binom{r}{l} M_1^{t+l} \cdot \frac{1 - (2^{-(t+l)} a)^{q_1}}{2^{t+l} - a} + \frac{1}{a} \binom{r}{\ell} M_1^{t+\ell} q_1
\end{aligned}$$

We shall summarize these two cases in a single expression by writing

$$(*) = \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \binom{r}{l} M_1^{t+l} \cdot \frac{1 - (2^{-(t+l)} a)^{q_1}}{2^{t+l} - a} + \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} M_1^{t+\ell} q_1. \quad (28)$$

Notice that if $r = 0$, then $(*) = 0$ and, since in this case $\delta_\ell = 0$, the right-hand side expression of (28) is indeed 0. Without making use of δ_ℓ , (28) would not be true when $r = 0$. In the next sum we shall apply a similar argument to split each sum $\sum_{l=0}^{r-1}$ into a sum $\sum_{\substack{l=0 \\ l \neq \ell}}^{r-1}$ and a term that must only be considered when $r > 0$ and $\ell \in \{0, \dots, r-1\}$, and hence it appears multiplied by δ_ℓ .

For every $j = 1, \dots, s-1$:

$$\begin{aligned}
& \sum_{k=q_j+1}^{q_{j+1}} a^{k-1} \left(\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor \right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \right) \\
&= \sum_{k=q_j+2}^{q_{j+1}} a^{k-1} \sum_{l=0}^{r-1} \binom{r}{l} (2^{-k} M_{j+1})^{t+l} + a^{q_j} \sum_{i=0}^{r+t-1} \binom{r+t}{i} (2^{-q_j-1} M_{j+1})^i - a^{q_j} \sum_{l=0}^{r-1} \binom{r}{l} (2^{-q_j-1} M_{j+1})^{t+l} \\
&= \frac{1}{a} \sum_{l=0}^{r-1} \binom{r}{l} M_{j+1}^{t+l} \sum_{k=q_j+1}^{q_{j+1}} (2^{-(t+l)} a)^k + \frac{1}{a} \sum_{i=0}^{r+t-1} \binom{r+t}{i} M_{j+1}^i (2^{-i} a)^{q_j+1} - \frac{2}{a} \sum_{l=0}^{r-1} \binom{r}{l} M_{j+1}^{t+l} (2^{-(t+l)} a)^{q_j+1} \\
&= \frac{1}{a} \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \binom{r}{l} M_{j+1}^{t+l} \sum_{k=q_j+1}^{q_{j+1}} (2^{-(t+l)} a)^k + \delta_\ell \frac{1}{a} \binom{r}{\ell} M_{j+1}^{t+\ell} \sum_{k=q_j+1}^{q_{j+1}} 1^k + \frac{1}{a} \sum_{i=0}^{r+t-1} \binom{r+t}{i} M_{j+1}^i (2^{-i} a)^{q_j+1} \\
&\quad - \frac{2}{a} \sum_{l=0}^{r-1} \binom{r}{l} M_{j+1}^{t+l} (2^{-(t+l)} a)^{q_j+1} \\
&= \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \binom{r}{l} M_{j+1}^{t+l} \cdot \frac{(2^{-(t+l)} a)^{q_j} - (2^{-(t+l)} a)^{q_j+1}}{2^{t+l} - a} + \delta_\ell \frac{1}{a} \binom{r}{\ell} M_{j+1}^{t+\ell} (q_{j+1} - q_j) \\
&\quad + \frac{1}{a} \sum_{i=0}^{r+t-1} \binom{r+t}{i} M_{j+1}^i (2^{-i} a)^{q_j+1} - \frac{2}{a} \sum_{l=0}^{r-1} \binom{r}{l} M_{j+1}^{t+l} (2^{-(t+l)} a)^{q_j+1}
\end{aligned}$$

Then (and using $q_0 = 0$)

$$\begin{aligned}
\hat{y}_n &= \sum_{k=1}^{q_s} a^{k-1} \left(\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor \right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \right) \\
&= \sum_{j=0}^{s-1} \sum_{k=q_j+1}^{q_{j+1}} a^{k-1} \left(\left(1 + \left\lfloor \frac{n-1}{2^k} \right\rfloor \right)^r \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^t - \left\lfloor \frac{n-1}{2^k} + \frac{1}{2} \right\rfloor^r \left\lfloor \frac{n-1}{2^k} \right\rfloor^t \right) \\
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} M_1^{t+\ell} q_1 + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \binom{r}{l} M_1^{t+l} \cdot \frac{1 - (2^{-(t+l)} a)^{q_1}}{2^{t+l} - a} \\
&\quad + \sum_{j=1}^{s-1} \left[\delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} M_{j+1}^{t+\ell} (q_{j+1} - q_j) + \frac{1}{a} \sum_{i=0}^{r+t-1} \binom{r+t}{i} M_{j+1}^i (2^{-i} a)^{q_j+1} \right. \\
&\quad \left. + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \binom{r}{l} M_{j+1}^{t+l} \cdot \frac{(2^{-(t+l)} a)^{q_j} - (2^{-(t+l)} a)^{q_j+1}}{2^{t+l} - a} - \frac{2}{a} \sum_{l=0}^{r-1} \binom{r}{l} M_{j+1}^{t+l} (2^{-(t+l)} a)^{q_j+1} \right] \\
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{j=0}^{s-1} M_{j+1}^{t+\ell} (q_{j+1} - q_j) + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} \sum_{j=0}^{s-1} M_{j+1}^{t+l} \cdot ((2^{-(t+l)} a)^{q_j} - (2^{-(t+l)} a)^{q_j+1}) \\
&\quad + \sum_{i=0}^{r+t-1} \binom{r+t}{i} \sum_{j=1}^{s-1} 2^{-i} M_{j+1}^i (2^{-i} a)^{q_j} - 2 \sum_{i=t}^{r+t-1} \binom{r}{i-t} \sum_{j=1}^{s-1} 2^{-i} M_{j+1}^i (2^{-i} a)^{q_j} \tag{29}
\end{aligned}$$

Now,

$$\begin{aligned}
& \sum_{j=0}^{s-1} ((2^{-(t+l)}a)^{q_j} - (2^{-(t+l)}a)^{q_{j+1}}) M_{j+1}^{t+l} \\
&= M_1^{t+l} + \sum_{j=1}^{s-1} (2^{-(t+l)}a)^{q_j} (M_{j+1}^{t+l} - M_j^{t+l}) - (2^{-(t+l)}a)^{q_s} M_s^{t+l} \\
&= (n-1)^{t+l} + \sum_{j=1}^{s-1} (2^{-(t+l)}a)^{q_j} (M_{j+1}^{t+l} - (M_{j+1} + 2^{q_j})^{t+l}) - (2^{-(t+l)}a)^{q_s} (2^{q_s})^{t+l} \\
&= (n-1)^{t+l} - a^{q_s} - \sum_{j=1}^{s-1} (2^{-(t+l)}a)^{q_j} \sum_{i=0}^{t+l-1} \binom{t+l}{i} M_{j+1}^i 2^{(t+l-i)q_j} \\
&= (n-1)^{t+l} - a^{q_s} - \sum_{i=0}^{t+l-1} \binom{t+l}{i} \sum_{j=1}^{s-1} M_{j+1}^i (2^{-i}a)^{q_j} \tag{30}
\end{aligned}$$

Using this in Eqn. (29), we obtain

$$\begin{aligned}
\widehat{y}_n &= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{j=0}^{s-1} M_{j+1}^{t+\ell} (q_{j+1} - q_j) \\
&+ \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} ((n-1)^{t+l} - a^{q_s} - \sum_{i=0}^{t+l-1} \binom{t+l}{i} \sum_{j=1}^{s-1} M_{j+1}^i (2^{-i}a)^{q_j}) \\
&+ \sum_{i=0}^{r+t-1} \binom{r+t}{i} \sum_{j=1}^{s-1} 2^{-i} M_{j+1}^i (2^{-i}a)^{q_j} - 2 \sum_{i=t}^{r+t-1} \binom{r}{i-t} \sum_{j=1}^{s-1} 2^{-i} M_{j+1}^i (2^{-i}a)^{q_j} \\
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{j=0}^{s-1} M_{j+1}^{t+\ell} (q_{j+1} - q_j) + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} ((n-1)^{t+l} - a^{q_s}) \\
&- \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \sum_{i=0}^{t+l-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} S_{n-1}^{(0,i)} + \sum_{i=0}^{r+t-1} \binom{r+t}{i} 2^{-i} S_{n-1}^{(0,i)} - 2 \sum_{i=t}^{r+t-1} \binom{r}{i-t} 2^{-i} S_{n-1}^{(0,i)} \\
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{j=0}^{s-1} M_{j+1}^{t+\ell} (q_{j+1} - q_j) + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} ((n-1)^{t+l} - a^{q_s}) \\
&- \sum_{i=0}^{r+t-1} \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} S_{n-1}^{(0,i)} + \sum_{i=0}^{r+t-1} \binom{r+t}{i} 2^{-i} S_{n-1}^{(0,i)} - 2 \sum_{i=0}^{r+t-1} \binom{r}{i-t} 2^{-i} S_{n-1}^{(0,i)} \\
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{j=0}^{s-1} M_{j+1}^{t+\ell} (q_{j+1} - q_j) + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a} ((n-1)^{t+l} - a^{q_s}) \\
&+ \sum_{i=0}^{r+t-1} (2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t}) - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} S_{n-1}^{(0,i)}
\end{aligned}$$

as we claimed. \square

Remark 11. When $r = t = 0$, the formula in the last proposition simply says $y_n^{(r,t)} = a^{q_{s_{n-1}}(n-1)}$.

In the expression for $y_n^{(r,t)}$ given in the last proposition, the term

$$\delta_\ell \cdot \frac{1}{a} \sum_{j=0}^{s_{n-1}-1} M_{j+1}(n-1)^{t+\ell} (q_{j+1}(n-1) - q_j(n-1))$$

is different from 0 only when $r > 0$ and $a \in \{2^t, \dots, 2^{r+t-1}\}$. When $t + \ell = 0$, that is, when $a = 1$, the sum in it simplifies to

$$\sum_{j=0}^{s_{n-1}-1} M_{j+1}(n-1)^0 (q_{j+1}(n-1) - q_j(n-1)) = q_{s_{n-1}}(n-1) \quad (31)$$

And notice that this sum contributes in this $a = 1$ case to $y_n^{(r,t)}$ only when $r > 0$ and $a = 1 \in \{2^t, \dots, 2^{r+t-1}\}$, that is, when $r > 0$ and $t = 0$. When $t + \ell > 0$, and recalling that $2^{t+\ell} = a$, this sum is

$$\begin{aligned} & \sum_{j=0}^{s_{n-1}-1} M_{j+1}(n-1)^{t+\ell} (q_{j+1}(n-1) - q_j(n-1)) \quad (32) \\ &= \sum_{j=1}^{s_{n-1}-1} q_j(n-1) (M_j(n-1)^{t+\ell} - M_{j+1}(n-1)^{t+\ell}) + 2^{(t+\ell)q_{s_{n-1}}(n-1)} q_{s_{n-1}}(n-1) \\ &= \sum_{j=1}^{s_{n-1}-1} q_j(n-1) ((M_{j+1}(n-1) + 2^{q_j(n-1)})^{t+\ell} - M_{j+1}(n-1)^{t+\ell}) + a^{q_{s_{n-1}}(n-1)} q_{s_{n-1}}(n-1) \\ &= a^{q_{s_{n-1}}(n-1)} q_{s_{n-1}}(n-1) + \sum_{j=1}^{s_{n-1}-1} q_j(n-1) \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} M_{j+1}(n-1)^i 2^{q_j(n-1)(t+\ell-i)} \\ &= a^{q_{s_{n-1}}(n-1)} q_{s_{n-1}}(n-1) + \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} \sum_{j=1}^{s_{n-1}-1} q_j(n-1) M_{j+1}(n-1)^i (a 2^{-i})^{q_j(n-1)} \\ &= a^{q_{s_{n-1}}(n-1)} q_{s_{n-1}}(n-1) + \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} S_{n-1}^{(1,i)} \quad (33) \end{aligned}$$

where, for every $n, i \in \mathbb{N}$,

$$S_n^{(1,i)} = \sum_{j=1}^{s_n-1} q_j(n) (2^{-i} a)^{q_j(n)} M_{j+1}(n)^i.$$

Notice that when $\delta_\ell = 1$ and $a = 1$, Eqn. (33) becomes Eqn. (31).

4.5 Proof of the main result, up to computing the α 's

For every $d, n, m \in \mathbb{N}$,

$$\alpha_n^{(d,m)} = \sum_{k=1}^{n-1} S_k^{(d,m)} = \sum_{k=1}^{n-1} \sum_{j=1}^{s_k-1} q_j(k)^d (2^{-m} a)^{q_j(k)} M_{j+1}(k)^m$$

By Proposition 10 and Eqn. (26), we have

$$\begin{aligned}
x_n^{(r,t)} &= \sum_{k=2}^n y_k^{(r,t)} = \sum_{k=1}^{n-1} \left[a^{q_{s_k}(k)} + \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{j=0}^{s_k-1} M_{j+1}(k)^{t+\ell} (q_{j+1}(k) - q_j(k)) + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l}-a} \cdot k^{t+l} \right. \\
&\quad \left. - \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l}-a} \cdot a^{q_{s_k}(k)} + \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l}-a} \right) S_k^{(0,i)} \right] \\
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{k=1}^{n-1} \sum_{j=0}^{s_k-1} M_{j+1}(k)^{t+\ell} (q_{j+1}(k) - q_j(k)) + \left(1 - \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l}-a} \right) \left(\sum_{k=1}^{n-1} a^{q_{s_k}(k)} \right) \\
&\quad + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l}-a} \sum_{k=1}^{n-1} k^{t+l} + \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l}-a} \right) \sum_{k=1}^{n-1} S_k^{(0,i)} \\
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{k=1}^{n-1} \sum_{j=0}^{s_k-1} M_{j+1}(k)^{t+\ell} (q_{j+1}(k) - q_j(k)) + \left(1 - \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l}-a} \right) \left(\sum_{k=1}^{n-1} a^{q_{s_k}(k)} \right) \\
&\quad + \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{(t+l+1)(2^{t+l}-a)} \left(\sum_{j=0}^{t+l+1} \binom{t+l+1}{j} B_j n^{t+l+1-j} + (-1)^{t+l} B_{t+l+1} \right) \\
&\quad + \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l}-a} \right) \alpha_n^{(0,i)}
\end{aligned}$$

(by Faulhaber's Formula (12))

$$\begin{aligned}
&= \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \sum_{k=1}^{n-1} \sum_{j=0}^{s_k-1} M_{j+1}(k)^{t+\ell} (q_{j+1}(k) - q_j(k)) + \left(1 - \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l}-a} \right) \left(\sum_{k=1}^{n-1} a^{q_{s_k}(k)} \right) \\
&\quad + \sum_{k=1}^{r+t} \left(\sum_{\substack{i=k \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1} \binom{i}{k} B_{i-k}}{i(2^{i-1}-a)} \right) n^k + \sum_{\substack{i=t+1 \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1}}{i(2^{i-1}-a)} (B_i - (-1)^i B_i) \\
&\quad + \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l}-a} \right) \alpha_n^{(0,i)} \tag{34}
\end{aligned}$$

because

$$\begin{aligned}
& \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{(t+l+1)(2^{t+l}-a)} \left(\sum_{j=0}^{t+l+1} \binom{t+l+1}{j} B_j n^{t+l+1-j} + (-1)^{t+l} B_{t+l+1} \right) \\
&= \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{(t+l+1)(2^{t+l}-a)} \left(\sum_{k=0}^{t+l+1} \binom{t+l+1}{k} B_{t+l+1-k} n^k + (-1)^{t+l} B_{t+l+1} \right) \\
&= \sum_{\substack{i=t+1 \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1}}{i(2^{i-1}-a)} \left(\sum_{k=0}^i \binom{i}{k} B_{i-k} n^k - (-1)^i B_i \right) \\
&= \sum_{k=0}^{r+t} \left(\sum_{\substack{i=k \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1} \binom{i}{k} B_{i-k}}{i(2^{i-1}-a)} \right) n^k - \sum_{\substack{i=t+1 \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1}}{i(2^{i-1}-a)} (-1)^i B_i \\
&= \sum_{\substack{i=0 \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1}}{i(2^{i-1}-a)} B_i + \sum_{k=1}^{r+t} \left(\sum_{\substack{i=k \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1} \binom{i}{k} B_{i-k}}{i(2^{i-1}-a)} \right) n^k - \sum_{\substack{i=t+1 \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1}}{i(2^{i-1}-a)} (-1)^i B_i \\
&= \sum_{k=1}^{r+t} \left(\sum_{\substack{i=k \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1} \binom{i}{k} B_{i-k}}{i(2^{i-1}-a)} \right) n^k + \sum_{\substack{i=t+1 \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1}}{i(2^{i-1}-a)} (B_i - (-1)^i B_i)
\end{aligned}$$

Now, in Eqn. (34) we have that, by Lemma 4,

$$\begin{aligned}
\sum_{k=1}^{n-1} a^{q_{s_k}(k)} &= T(0, q_{s_n}(n), 2a) + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} \\
&= \begin{cases} q_{s_n}(n) + n \cdot 2^{-q_{s_n}(n)} - 1 & \text{if } a = 1/2 \\ \frac{(2a)^{q_{s_n}(n)} - 1}{2a - 1} + n \cdot a^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)} & \text{if } a \neq 1/2 \end{cases}
\end{aligned}$$

and, by (8),

$$\sum_{\substack{i=t+1 \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1}}{i(2^{i-1}-a)} (B_i - (-1)^i B_i) = \begin{cases} 1/(a-1) & \text{if } r > 0, t = 0, a \neq 1 \\ 0 & \text{otherwise} \end{cases}$$

Concerning the first term of Eqn. (34), which only contributes to $x_n^{(r,t)}$ when $r > 0$ and $a = 2^{t+\ell}$ with $\ell \in \{0, \dots, r-1\}$, by (33) and Lemma 4 this sum can be expressed as

$$\begin{aligned}
& \sum_{k=1}^{n-1} \sum_{j=0}^{s_k-1} M_{j+1}(k)^{t+\ell} (q_{j+1}(k) - q_j(k)) = \sum_{k=1}^{n-1} \left(a^{q_{s_k}(k)} q_{s_k}(k) + \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} S_k^{(1,i)} \right) \\
&= \sum_{k=1}^{n-1} a^{q_{s_k}(k)} q_{s_k}(k) + \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} \sum_{k=1}^{n-1} S_k^{(1,i)} \\
&= T(1, q_{s_n}(n), 2a) + nq_{s_n}(n)a^{q_{s_n}(n)} - q_{s_n}(n)(2a)^{q_{s_n}(n)} + \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} \alpha_n^{(1,i)}
\end{aligned}$$

In summary, this proves the following theorem. The expressions given in Section 2 are obtained by rewriting the formula in this theorem for the different relevant cases of the exponents r, t and the coefficient a .

Theorem 12. For every $n \geq 2$,

$$\begin{aligned}
x_n^{(r,t)} &= \left(1 - \sum_{\substack{l=0 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l}}{2^{t+l} - a}\right) (T(0, q_{s_n}(n), 2a) + na^{q_{s_n}(n)} - (2a)^{q_{s_n}(n)}) \\
&+ \sum_{k=1}^{r+t} \left(\sum_{\substack{i=k \\ i \neq t+\ell+1}}^{r+t} \frac{\binom{r}{i-t-1} \binom{i}{k} B_{i-k}}{i(2^{i-1} - a)} \right) n^k + \frac{1}{a-1} \cdot \delta_{r>0, t=0, a \neq 1} \\
&+ \sum_{i=0}^{r+t-1} \left(2^{-i} \binom{r+t}{i} - 2^{-i+1} \binom{r}{i-t} - \sum_{\substack{l=i-t+1 \\ l \neq \ell}}^{r-1} \frac{\binom{r}{l} \binom{t+l}{i}}{2^{t+l} - a} \right) \alpha_n^{(0,i)} \\
&+ \delta_\ell \cdot \frac{1}{a} \binom{r}{\ell} \left(T(1, q_{s_n}(n), 2a) + nq_{s_n}(n)a^{q_{s_n}(n)} - q_{s_n}(n)(2a)^{q_{s_n}(n)} + \sum_{i=0}^{t+\ell-1} \binom{t+\ell}{i} \alpha_n^{(1,i)} \right)
\end{aligned}$$

where $\delta_{r>0, t=0, a \neq 1} = 1$ if $r > 0$, $t = 0$, and $a \neq 1$, and it is 0 otherwise.

So, this proves our main result, up to the computation of the terms

$$\alpha_n^{(d,i)} = \sum_{k=1}^{n-1} \sum_{j=1}^{s_k-1} q_j(k)^d (2^{-i}a)^{q_j(k)} M_{j+1}(k)^i, \quad d = 0, 1.$$

And notice that we only need to compute the $\alpha^{(1,i)}$'s when $r \geq 1$ and $a = 2^{t+\ell}$, for some $\ell \in \{0, \dots, r-1\}$ with $t + \ell > 0$. In particular, in the computation of $\alpha^{(1,i)}$ we can assume that $a \neq 1, 1/2$, in order to avoid discussing cases that are irrelevant for our main result.

4.6 Computation of the α 's

Recall that, for every $d, m \in \mathbb{N}$ and $n \in \mathbb{N}_{\geq 1}$, we set

$$S_n^{(d,m)} = \sum_{j=1}^{s_n-1} q_j(n)^d (2^{-m}a)^{q_j(n)} M_{j+1}(n)^m$$

and

$$\alpha_n^{(d,m)} = \sum_{k=1}^{n-1} S_k^{(d,m)} = \sum_{k=1}^{n-1} \sum_{j=1}^{s_k-1} q_j(k)^d (a2^{-m})^{q_j(k)} M_{j+1}(k)^m$$

We are interested in closed formulas for $\alpha_n^{(d,m)}$ when $d = 0$ as well as when $d = 1$ and specific values of a which in particular exclude the cases $a = 1, 1/2$.

When $m = 0$, by Lemmas 4 and 5 we have that

$$\begin{aligned}
\alpha_n^{(d,0)} &= \sum_{k=1}^{n-1} \sum_{j=1}^{s_k-1} q_j(k)^d a^{q_j(k)} = \sum_{k=1}^{n-1} \sum_{j=1}^{s_k} q_j(k)^d a^{q_j(k)} - \sum_{k=1}^{n-1} q_{s_k}(k)^d a^{q_{s_k}(k)} \\
&= \sum_{i=1}^{s_n} 2^{q_i(n)-1} T(d, q_i(n), a) + \sum_{i=1}^{s_n} q_i(n)^d a^{q_i(n)} (n - M_i(n)) \\
&\quad - T(d, q_{s_n}(n), 2a) - (n - 2^{q_{s_n}(n)}) q_{s_n}(n)^d a^{q_{s_n}(n)} \\
&= \sum_{i=1}^{s_n} 2^{q_i(n)-1} T(d, q_i(n), a) + \sum_{i=1}^{s_n-1} q_i(n)^d a^{q_i(n)} (n - M_i(n)) - T(d, q_{s_n}(n), 2a) \quad (35)
\end{aligned}$$

and then, in particular,

$$\begin{aligned} \alpha_n^{(0,0)} &= \sum_{i=1}^{s_n} 2^{q_i(n)-1} T(0, q_i(n), a) + \sum_{i=1}^{s_n-1} a^{q_i(n)} (n - M_i(n)) - T(0, q_{s_n}(n), 2a) \\ &= \begin{cases} \sum_{j=1}^{s_n} 2^{q_j(n)-1} (q_j(n) - 2j) + (s_n - 1)n + 1 & \text{if } a = 1 \\ \sum_{j=1}^{s_n-1} 2^{-q_j(n)} (n - M_j(n)) + n - q_{s_n} - s_n & \text{if } a = 1/2 \\ \sum_{j=1}^{s_n-1} a^{q_j(n)} (n - M_j(n)) + \frac{1}{2(a-1)} \left(\sum_{j=1}^{s_n} (2a)^{q_j(n)} - n \right) - \frac{(2a)^{q_{s_n}(n)} - 1}{2a-1} & \text{if } a \neq 1, 1/2 \end{cases} \end{aligned}$$

and, when $a \neq 1, 1/2$,

$$\begin{aligned} \alpha_n^{(1,0)} &= \sum_{i=1}^{s_n} 2^{q_i(n)-1} T(1, q_i(n), a) + \sum_{i=1}^{s_n-1} q_i(n) a^{q_i(n)} (n - M_i(n)) - T(1, q_{s_n}(n), 2a) \\ &= \sum_{i=1}^{s_n-1} q_i(n) a^{q_i(n)} (n - M_i(n)) + \frac{1}{2(a-1)} \sum_{i=1}^{s_n} (2a)^{q_i(n)} q_i(n) - \frac{a}{2(a-1)^2} \sum_{i=1}^{s_n} (2a)^{q_i(n)} \\ &\quad - \frac{1}{2a-1} (2a)^{q_{s_n}(n)} q_{s_n}(n) + \frac{2a}{(2a-1)^2} (2a)^{q_{s_n}(n)} + \frac{a}{2(a-1)^2} n - \frac{2a}{(2a-1)^2} \end{aligned}$$

Therefore, in the sequel we only consider the case $m > 0$.

Lemma 13. For every $m \in \mathbb{N}_{\geq 1}$, $d \in \mathbb{N}$, $M \in \mathbb{N}_{\geq 1}$, and $0 \leq l \leq q_1(M)$,

$$\alpha_{M+2^l}^{(d,m)} = \alpha_M^{(d,m)} + \alpha_{2^l}^{(d,m)} + \sum_{p=0}^{m-1} \binom{m}{p} M^{m-p} \gamma_l^{(d,p,m)} + 2^l S_M^{(d,m)}$$

Proof. Let $M = \sum_{i=1}^s 2^{q_i}$, with $s \geq 1$ and $q_s > \dots > q_1 > 0$, and let $0 \leq l \leq q_1$, so that $s_M = s$ and $q_i(M) = q_i$ for each $i = 1, \dots, s$. Then

$$\begin{aligned} \alpha_{M+2^l}^{(d,m)} &= \sum_{k=1}^{M-1} \sum_{i=1}^{s_k-1} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^m + \sum_{k=M}^{M+2^l-1} \sum_{i=1}^{s_k-1} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^m \\ &= \alpha_M^{(d,m)} + \sum_{k=M}^{M+2^l-1} \sum_{i=1}^{s_k-1} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^m \end{aligned}$$

where

$$\begin{aligned}
& \sum_{k=M}^{M+2^l-1} \sum_{i=1}^{s_k-1} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^m \\
&= \sum_{k=M}^{M+2^l-1} \sum_{i=1}^{s_k-s} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^m + \sum_{k=M}^{M+2^l-1} \sum_{i=s_k-s+1}^{s_k-1} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^m \\
&= \sum_{k=M}^{M+2^l-1} \sum_{i=1}^{s_k-s} q_i(k)^d (a2^{-m})^{q_i(k)} (M_{i+1}(k-M) + M)^m + \sum_{k=M}^{M+2^l-1} \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} M_{i+1}(M)^m \\
&= \sum_{k=0}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d (a2^{-m})^{q_i(k)} (M_{i+1}(k) + M)^m + 2^l \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} M_{i+1}(M)^m \\
&= \sum_{k=1}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d (a2^{-m})^{q_i(k)} \sum_{p=0}^m \binom{m}{p} M_{i+1}(k)^p M^{m-p} + 2^l S_M^{(d,m)} \\
&= \sum_{p=0}^{m-1} \binom{m}{p} M^{m-p} \sum_{k=1}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^p + \sum_{k=1}^{2^l-1} \sum_{i=1}^{s_k} q_i(k)^d (a2^{-m})^{q_i(k)} M_{i+1}(k)^m \\
&\quad + 2^l S_M^{(d,m)} \\
&= \sum_{p=0}^{m-1} \binom{m}{p} M^{m-p} \gamma_l^{(d,p,m)} + \alpha_{2^l}^{(d,m)} + 2^l S_M^{(d,m)}
\end{aligned}$$

yielding the expression in the statement. \square

Corollary 14. For every $m \in \mathbb{N}_{\geq 1}$ and for every $l \geq 0$,

$$\alpha_{2^l}^{(d,m)} = \frac{2^{ml+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} 2^{-p(l-1)} B_p \cdot T(d, l-1, 2^{p-m} a)$$

Proof. Taking $M = 2^l$ in the last lemma (and recalling that $S_{2^l}^{(d,m)} = 0$ because $s_{2^l} = 1$), we obtain

$$\alpha_{2^{l+1}}^{(d,m)} = 2\alpha_{2^l}^{(d,m)} + \sum_{p=0}^{m-1} \binom{m}{p} 2^{l(m-p)} \gamma_l^{(d,p,m)}$$

The solution of this recurrence with $\alpha_{2^0}^{(d,m)} = \alpha_1^{(d,m)} = 0$ is

$$\alpha_{2^l}^{(d,m)} = \sum_{k=1}^{l-1} 2^{l-k-1} \sum_{p=0}^{m-1} \binom{m}{p} 2^{k(m-p)} \gamma_k^{(d,p,m)}$$

Then, using Corollary 8,

$$\begin{aligned}
\alpha_{2^l}^{(d,m)} &= \sum_{k=1}^{l-1} 2^{l-k-1} \sum_{p=0}^{m-1} \binom{m}{p} 2^{k(m-p)} \frac{a^{k-1} 2^{-(m-1)(k-1)+pk}}{p+1} \sum_{t=1}^{k-1} (k-t-1)^d (a^{-1} 2^{m-p-1})^t (B_{p+1}(2^t) - B_{p+1}) \\
&\quad + \sum_{k=1}^{l-1} 2^{l-k-1} 2^{km} (a 2^{-(m-1)})^{k-1} (k-1)^d \\
&= \frac{2^{m+l-2}}{m+1} \sum_{t=1}^{l-2} \left[\left(\sum_{p=0}^{m-1} \binom{m+1}{p+1} \right) 2^{(m-p-1)t} (B_{p+1}(2^t) - B_{p+1}) \right] \left(\sum_{k=t+1}^{l-1} (k-t-1)^d a^{k-t-1} \right) \\
&\quad + 2^{m+l-2} \sum_{k=1}^{l-1} (k-1)^d a^{k-1} \\
&= \frac{2^{m+l-2}}{m+1} \sum_{t=1}^{l-2} \left[\left(\sum_{p=1}^m \binom{m+1}{p} \right) 2^{(m-p)t} (B_p(2^t) - B_p) \right] \left(\sum_{k=0}^{l-t-2} k^d a^k \right) + 2^{m+l-2} \sum_{k=0}^{l-2} k^d a^k \\
&= \frac{2^{m+l-2}}{m+1} \sum_{t=0}^{l-2} \left[\left(\sum_{p=0}^m \binom{m+1}{p} \right) 2^{(m-p)t} (B_p(2^t) - B_p) \right] T(d, l-t-1, a)
\end{aligned}$$

by Eqn. (9). Now,

$$\begin{aligned}
&\sum_{p=0}^m \binom{m+1}{p} 2^{(m-p)t} (B_p(2^t) - B_p) \\
&= 2^{-t} \sum_{p=0}^m \binom{m+1}{p} 2^{t(m+1-p)} B_p(2^t) - \sum_{p=0}^m \binom{m+1}{p} 2^{(m-p)t} B_p \\
&= 2^{-t} (B_{m+1}(2^{t+1}) - B_{m+1}(2^t)) - \sum_{p=0}^m \binom{m+1}{p} 2^{(m-p)t} B_p \\
&= 2^{-t} \sum_{p=0}^{m+1} \binom{m+1}{p} 2^{(m+1-p)(t+1)} B_p - 2^{-t} \sum_{p=0}^{m+1} \binom{m+1}{p} 2^{(m+1-p)t} B_p - \sum_{p=0}^m \binom{m+1}{p} 2^{(m-p)t} B_p \\
&= 2 \sum_{p=0}^m \binom{m+1}{p} (2^{(m-p)(t+1)} - 2^{(m-p)t}) B_p
\end{aligned}$$

and therefore

$$\begin{aligned}
\alpha_{2^i}^{(d,m)} &= \frac{2^{m+l-2}}{m+1} \sum_{t=0}^{l-2} \left[\left(2 \sum_{p=0}^m \binom{m+1}{p} (2^{(m-p)(t+1)} - 2^{(m-p)t}) B_p \right) T(d, l-t-1, a) \right] \\
&= \frac{2^{m+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} B_p \cdot \left[\sum_{t=0}^{l-2} 2^{(m-p)(t+1)} T(d, l-t-1, a) - \sum_{t=0}^{l-2} 2^{(m-p)t} T(d, l-t-1, a) \right] \\
&= \frac{2^{m+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} B_p \cdot \left[\sum_{t=1}^{l-1} 2^{(m-p)t} T(d, l-t, a) - \sum_{t=0}^{l-2} 2^{(m-p)t} T(d, l-t-1, a) \right] \\
&= \frac{2^{m+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} B_p \cdot \left[\sum_{t=1}^{l-1} 2^{(m-p)t} (l-t-1)^d a^{l-t-1} + \sum_{t=1}^{l-2} 2^{(m-p)t} T(d, l-t-1, a) \right. \\
&\quad \left. - \sum_{t=0}^{l-2} 2^{(m-p)t} T(d, l-t-1, a) \right] \\
&= \frac{2^{m+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} B_p \cdot \left[\sum_{t=1}^{l-1} 2^{(m-p)t} (l-t-1)^d a^{l-t-1} - T(d, l-1, a) \right] \\
&= \frac{2^{m+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} B_p \cdot \left(\sum_{t=1}^{l-1} 2^{(m-p)t} (l-t-1)^d a^{l-t-1} \right) \quad (\text{by Eqn. (7)}) \\
&= \frac{2^{ml+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} 2^{-p(l-1)} B_p \cdot \left(\sum_{t=1}^{l-1} (l-t-1)^d (2^{p-m} a)^{l-t-1} \right) \\
&= \frac{2^{ml+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} 2^{-p(l-1)} B_p \cdot \left(\sum_{k=0}^{l-2} k^d (2^{p-m} a)^k \right) \\
&= \frac{2^{ml+l-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} 2^{-p(l-1)} B_p \cdot T(d, l-1, 2^{p-m} a)
\end{aligned}$$

as we claimed. \square

Proposition 15. For every $n, m \geq 1$,

$$\begin{aligned}
\alpha_n^{(d,m)} &= \frac{1}{2(m+1)} \sum_{i=1}^{s_n} \sum_{j=0}^m \binom{m+1}{j} B_j 2^j M_i(n)^{m+1-j} (T(d, q_i(n), a 2^{j-m}) - T(d, q_{i-1}(n), a 2^{j-m})) \\
&\quad + \sum_{i=1}^{s_n-1} q_i(n)^d (a 2^{-m})^{q_i(n)} (n - M_i(n)) M_{i+1}(n)^m
\end{aligned}$$

Proof. To simplify the notations, let $s = s_n$ and $q_i = q_i(n)$ and $M_i = M_i(n) = \sum_{h=i}^s 2^{q_h}$, for every $i = 1, \dots, s$. Then, Lemma 13 and a simple argument by induction implies that

$$\begin{aligned}
\alpha_n^{(d,m)} &= \sum_{k=1}^s \alpha_{2^{q_k}}^{(d,m)} + \sum_{p=0}^{m-1} \binom{m}{p} \sum_{i=1}^{s-1} M_{i+1}^{m-p} \gamma_{q_i}^{(d,p,m)} + \sum_{k=1}^{s-1} 2^{q_k} S_{M_{k+1}}^{(d,m)} \\
&= \sum_{k=1}^s \alpha_{2^{q_k}}^{(d,m)} + \sum_{p=0}^{m-1} \binom{m}{p} \sum_{i=1}^{s-1} M_{i+1}^{m-p} \gamma_{q_i}^{(d,p,m)} + \sum_{k=1}^{s-1} 2^{q_k} \sum_{i=k+1}^{s-1} q_i^d (a 2^{-m})^{q_i} M_{i+1}^m \\
&= \sum_{k=1}^s \alpha_{2^{q_k}}^{(d,m)} + \sum_{p=0}^{m-1} \binom{m}{p} \sum_{i=1}^{s-1} M_{i+1}^{m-p} \gamma_{q_i}^{(d,p,m)} + \sum_{i=2}^{s-1} q_i^d (a 2^{-m})^{q_i} M_{i+1}^m \sum_{k=1}^{i-1} 2^{q_k} \\
&= \sum_{k=1}^s \alpha_{2^{q_k}}^{(d,m)} + \sum_{p=0}^{m-1} \binom{m}{p} \sum_{i=1}^{s-1} M_{i+1}^{m-p} \gamma_{q_i}^{(d,p,m)} + \sum_{i=1}^{s-1} q_i^d (a 2^{-m})^{q_i} (n - M_i) M_{i+1}^m \quad (36)
\end{aligned}$$

where, with the convention that $M_{s+1}^0 = 1$,

$$\begin{aligned}
\sum_{k=1}^s \alpha_{2^{q_k}}^{(d,m)} &= \sum_{k=1}^s \frac{2^{mq_k+q_k-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} 2^{-p(q_k-1)} B_p \cdot T(d, q_k - 1, 2^{p-m} a) \\
&= \sum_{k=1}^s \frac{2^{mq_k+q_k-1}}{m+1} \sum_{p=0}^m \binom{m+1}{p} 2^{-p(q_k-1)} B_p \cdot \left(\sum_{t=0}^{q_k-2} t^d (2^{p-m} a)^t \right) \\
&= \frac{2^m}{m+1} \sum_{p=0}^m \binom{m+1}{p} B_p \cdot \left(\sum_{k=1}^s 2^{(m-p+1)(q_k-1)} \sum_{t=0}^{q_k-2} t^d (2^{p-m} a)^t \right) \\
&= \frac{2^m}{m+1} \sum_{p=0}^{m-1} \binom{m+1}{m+1} \binom{m+1}{p} B_p \cdot \left(\sum_{k=1}^s M_{k+1}^{m-m} 2^{(m-p+1)(q_k-1)} \sum_{t=0}^{q_k-2} t^d (2^{p-m} a)^t \right) \\
&\quad + 2^{m-1} B_m \cdot \left(\sum_{k=1}^s 2^{q_k} \sum_{t=0}^{q_k-2} t^d a^t \right)
\end{aligned}$$

and, by Corollary 8, setting $\delta_{q_1 \geq 1} = \min\{q_1, 1\}$

$$\begin{aligned}
\sum_{p=0}^{m-1} \binom{m}{p} \sum_{i=1}^{s-1} M_{i+1}^{m-p} \gamma_{q_i}^{(d,p,m)} &= \sum_{p=0}^{m-1} \binom{m}{p} \sum_{i=1}^s M_{i+1}^{m-p} \gamma_{q_i}^{(d,p,m)} \\
&= \sum_{p=0}^{m-1} \binom{m}{p} \sum_{i=1}^s M_{i+1}^{m-p} \frac{a^{q_i-1} 2^{-(m-1)(q_i-1)+pq_i}}{p+1} \sum_{t=1}^{q_i-1} (q_i - t - 1)^d (a^{-1} 2^{m-p-1})^t (B_{p+1}(2^t) - B_{p+1}) \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1 - 1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i - 1)^d \\
&= \frac{1}{m+1} \sum_{p=0}^{m-1} \binom{m+1}{p+1} 2^p \sum_{i=1}^s M_{i+1}^{m-p} \sum_{t=1}^{q_i-1} (q_i - t - 1)^d (2^{p-m+1} a)^{q_i-t-1} (B_{p+1}(2^t) - B_{p+1}) \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1 - 1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i - 1)^d \\
&= \frac{1}{m+1} \sum_{p=0}^{m-1} \binom{m+1}{p+1} 2^p \sum_{i=1}^s M_{i+1}^{m-p} \sum_{t=0}^{q_i-2} t^d (2^{p-m+1} a)^t (B_{p+1}(2^{q_i-t-1}) - B_{p+1}) \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1 - 1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i - 1)^d \\
&= \frac{1}{m+1} \sum_{p=0}^{m-1} \binom{m+1}{p+1} 2^p \sum_{i=1}^s M_{i+1}^{m-p} \sum_{t=0}^{q_i-2} t^d 2^{(p-m+1)t} a^t \sum_{j=0}^p \binom{p+1}{j} 2^{(p+1-j)(q_i-t-1)} B_j \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1 - 1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i - 1)^d \\
&= \frac{1}{m+1} \sum_{j=0}^{m-1} \sum_{p=j}^{m-1} \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (a 2^{-(m-j)})^t B_j \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1 - 1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i - 1)^d
\end{aligned}$$

Thus, returning back to (36) (and still with the convention that $M_{s+1}^0 = 1$), we have

$$\begin{aligned}
\alpha_n^{(d,m)} &= \frac{2^m}{m+1} \sum_{j=0}^{m-1} \binom{m+1}{m+1} \binom{m+1}{j} \left(\sum_{i=1}^s M_{i+1}^{m-m} 2^{(m-j+1)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (2^{j-m} a)^t \right) B_j \\
&\quad + 2^{m-1} B_m \cdot \left(\sum_{i=1}^s 2^{q_i} \sum_{t=0}^{q_i-2} t^d a^t \right) \\
&\quad + \frac{1}{m+1} \sum_{j=0}^{m-1} \sum_{p=j}^{m-1} \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (a 2^{-(m-j)})^t B_j \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1-1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i-1)^d \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a 2^{-m})^{q_i} (n - M_i) M_{i+1}^m \\
&= \frac{1}{m+1} \sum_{j=0}^{m-1} \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (a 2^{-(m-j)})^t B_j \\
&\quad + 2^{m-1} B_m \cdot \left(\sum_{i=1}^s 2^{q_i} \sum_{t=0}^{q_i-2} t^d a^t \right) \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1-1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i-1)^d \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a 2^{-m})^{q_i} (n - M_i) M_{i+1}^m \\
&= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (a 2^{-(m-j)})^t B_j \\
&\quad + M_2^m (a 2^{-(m-1)})^{q_1-1} (q_1-1)^d \delta_{q_1 \geq 1} + \sum_{i=2}^s M_{i+1}^m (a 2^{-(m-1)})^{q_i-1} (q_i-1)^d \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a 2^{-m})^{q_i} (n - M_i) M_{i+1}^m \tag{37}
\end{aligned}$$

Now notice that, for every $i = 1, \dots, s$

$$\begin{aligned}
&\frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} (q_i-1)^d (a 2^{-(m-j)})^{q_i-1} B_j \\
&= \frac{1}{m+1} (q_i-1)^d a^{q_i-1} \sum_{p=0}^m \binom{m+1}{p+1} M_{i+1}^{m-p} 2^{(p+1-m)(q_i-1)+p} \sum_{j=0}^p \binom{p+1}{j} B_j \\
&= \frac{1}{m+1} (q_i-1)^d a^{q_i-1} (m+1) M_{i+1}^m 2^{-(m-1)(q_i-1)} \quad (\text{by Eqn. (7)}) \\
&= (q_i-1)^d M_{i+1}^m (a 2^{-(m-1)})^{q_i-1}
\end{aligned}$$

Therefore, if $q_1 \geq 1$, (37) becomes

$$\begin{aligned}
\alpha_n^{(d,m)} &= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (a2^{-(m-j)})^t B_j \\
&\quad + \sum_{i=1}^s M_{i+1}^m (a2^{-(m-1)})^{q_i-1} (q_i-1)^d + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n-M_i) M_{i+1}^m \\
&= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-1} t^d (a2^{-(m-j)})^t B_j \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n-M_i) M_{i+1}^m
\end{aligned}$$

while, if $q_1 = 0$,

$$\begin{aligned}
\alpha_n^{(d,m)} &= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (a2^{-(m-j)})^t B_j \\
&\quad + \sum_{i=2}^s M_{i+1}^m (a2^{-(m-1)})^{q_i-1} (q_i-1)^d + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n-M_i) M_{i+1}^m \\
&= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=2}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-2} t^d (a2^{-(m-j)})^t B_j \\
&\quad + \sum_{i=2}^s M_{i+1}^m (a2^{-(m-1)})^{q_i-1} (q_i-1)^d + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n-M_i) M_{i+1}^m \\
&= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=2}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-1} t^d (a2^{-(m-j)})^t B_j \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n-M_i) M_{i+1}^m \\
&= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-1} t^d (a2^{-(m-j)})^t B_j \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n-M_i) M_{i+1}^m
\end{aligned}$$

So, in summary, for every $n \geq 1$,

$$\begin{aligned}
\alpha_n^{(d,m)} &= \frac{1}{m+1} \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-1} t^d (a2^{-(m-j)})^t B_j \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n-M_i) M_{i+1}^m \tag{38}
\end{aligned}$$

It remains to simplify the first term in the right-hand side expression:

$$\begin{aligned}
& \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{p+1} \binom{p+1}{j} 2^p \sum_{i=1}^s M_{i+1}^{m-p} 2^{(p+1-j)(q_i-1)} \sum_{t=0}^{q_i-1} t^d (a2^{-(m-j)})^t B_j \\
&= \sum_{i=1}^s \sum_{t=0}^{q_i-1} t^d a^t \sum_{j=0}^m \sum_{p=j}^m \binom{m+1}{j} \binom{m+1-j}{m-p} M_{i+1}^{m-p} 2^{(p+1-j)q_i+j-1-(m-j)t} B_j \\
&= \sum_{i=1}^s \sum_{t=0}^{q_i-1} t^d a^t \sum_{j=0}^m \sum_{p=0}^{m-j} \binom{m+1}{j} \binom{m+1-j}{p} M_{i+1}^p 2^{(m-p+1-j)q_i+j-1-(m-j)t} B_j \\
&= \sum_{i=1}^s \sum_{t=0}^{q_i-1} t^d a^t \sum_{j=0}^m \left(\sum_{p=0}^{m-j} \binom{m+1-j}{p} M_{i+1}^p 2^{(m+1-j-p)q_i} \right) \binom{m+1}{j} 2^{j-1-(m-j)t} B_j \\
&= \sum_{i=1}^s \sum_{t=0}^{q_i-1} t^d a^t \sum_{j=0}^m (M_i^{m+1-j} - M_{i+1}^{m+1-j}) \binom{m+1}{j} 2^{j-1-(m-j)t} B_j \\
&= \frac{1}{2} \sum_{j=0}^m \binom{m+1}{j} B_j 2^j \sum_{i=1}^s (M_i^{m+1-j} - M_{i+1}^{m+1-j}) T(d, q_i, a2^{j-m}) \\
&= \frac{1}{2} \sum_{j=0}^m \binom{m+1}{j} B_j 2^j \sum_{i=1}^s M_i^{m+1-j} (T(d, q_i, a2^{j-m}) - T(d, q_{i-1}, a2^{j-m}))
\end{aligned}$$

So, returning back to Eqn. (38), we finally obtain

$$\begin{aligned}
\alpha_n^{(d,m)} &= \frac{1}{2(m+1)} \sum_{i=1}^s \sum_{j=0}^m \binom{m+1}{j} B_j 2^j M_i^{m+1-j} (T(d, q_i, a2^{j-m}) - T(d, q_{i-1}, a2^{j-m})) \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n - M_i) M_{i+1}^m
\end{aligned}$$

as we claimed. \square

Remark 16. Recall from Eqn. (35) that

$$\alpha_n^{(d,0)} = \sum_{i=1}^{s_n} 2^{q_i(n)-1} T(d, q_i(n), a) + \sum_{i=1}^{s_n-1} q_i(n)^d a^{q_i(n)} (n - M_i(n)) - T(d, q_{s_n}(n), 2a)$$

Now, if we set $m = 0$ in the right-hand side term of the expression for $\alpha_n^{(d,m)}$, when $m > 0$, established in the last proposition, we obtain

$$\begin{aligned}
& \frac{1}{2} \sum_{i=1}^{s_n} M_i(n) (T(d, q_i(n), a) - T(d, q_{i-1}(n), a)) + \sum_{i=1}^{s_n-1} q_i(n)^d a^{q_i(n)} (n - M_i(n)) \\
&= \frac{1}{2} \sum_{i=1}^{s_n} T(d, q_i(n), a) (M_i(n) - M_{i+1}(n)) + \sum_{i=1}^{s_n-1} q_i(n)^d a^{q_i(n)} (n - M_i(n)) \\
&= \sum_{i=1}^{s_n} T(d, q_i(n), a) 2^{q_i(n)-1} + \sum_{i=1}^{s_n-1} q_i(n)^d a^{q_i(n)} (n - M_i(n)) \\
&= \alpha_n^{(d,0)} + T(d, q_{s_n}(n), 2a)
\end{aligned}$$

Therefore, we can combine Eqn. (35) and Proposition 15 into a single formula:

$$\begin{aligned}
\alpha_n^{(d,m)} &= \frac{1}{2(m+1)} \sum_{i=1}^s \sum_{j=0}^m \binom{m+1}{j} B_j 2^j M_i^{m+1-j} (T(d, q_i, a2^{j-m}) - T(d, q_{i-1}, a2^{j-m})) \\
&\quad + \sum_{i=1}^{s-1} q_i^d (a2^{-m})^{q_i} (n - M_i) M_{i+1}^m - T(d, q_{s_n}(n), 2a) \cdot \delta_{m=0}
\end{aligned}$$

with $\delta_{m=0} = 1$ if $m = 0$ and $\delta_{m=0} = 0$ if $m > 0$.

Remark 17. When $n = 2^l$, the formula in Proposition 15 says

$$\begin{aligned}\alpha_{2^l}^{(d,m)} &= \frac{1}{2(m+1)} \sum_{j=0}^m \binom{m+1}{j} B_j 2^{j+(m+1-j)l} T(d, l, a2^{j-m}) \\ &= \frac{2^{(m+1)l-1}}{m+1} \sum_{j=0}^m \binom{m+1}{j} B_j 2^{-j(l-1)} T(d, l, a2^{j-m})\end{aligned}$$

and this expression is equivalent to the one established in Corollary 14,

$$\alpha_{2^l}^{(d,m)} = \frac{2^{(m+1)l-1}}{m+1} \sum_{j=0}^m \binom{m+1}{j} B_j 2^{-j(l-1)} T(d, l-1, a2^{j-m}),$$

because

$$\sum_{j=0}^m \binom{m+1}{j} B_j 2^{-j(l-1)} (a2^{j-m})^{l-1} = 2^{-m(l-1)} \sum_{j=0}^m \binom{m+1}{j} B_j = 0$$

References

- [1] M. Abramowitz, I. A. S. Stegun. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables* (10th printing). National Bureau of Standards Applied Mathematics Series, 55 (1972).
- [2] A. V. Aho, J. E. Hopcroft, J. D. Ullman. *The design and analysis of computer algorithms*. Addison-Wesley Publishing Company (1974).
- [3] M. Akra, L. Bazzi. On the solution of linear recurrence equations. *Computational Optimization and Applications* 10 (1998), 195–210.
- [4] J. L. Bentley, D. Haken, J. B. Saxe. A general method for solving divide-and-conquer recurrences. *ACM SIGACT News* 12 (1980), 36–44,
- [5] Bernoulli numbers. *Encyclopedia of Mathematics*, URL: http://encyclopediaofmath.org/index.php?title=Bernoulli_numbers&oldid=32791
- [6] D. Colless. Review of “Phylogenetics: the theory and practice of phylogenetic systematics”. *Systematic Zoology* 31 (1982), 100–104.
- [7] T. H. Cormen, C. E. Leiserson, R. L. Rivest, C. Stein. *Introduction to Algorithms* (4th Edition). MIT Press (2009)
- [8] T. M. Coronado, M. Fischer, L. Herbst, F. Rosselló, K. Wicke. On the minimum value of the Colless index and the bifurcating trees that achieve it, *Journal of Mathematical Biology* 80 (2020), 1993–2054.
- [9] T. M. Coronado, A. Mir, F. Rosselló, G. Valiente. A balance index for phylogenetic trees based on rooted quartets. *Journal of Mathematical Biology* 79 (2019), 1105–1148.
- [10] M. Dehmer, F. Emmert-Streib (Eds.). *Quantitative graph theory*. CRC Press (2014).
- [11] M. Drmota, W. Szpankowski. A master theorem for discrete divide and conquer recurrences. *Journal of the ACM* 60 (2013), 1-49.
- [12] J. Felsenstein. *Inferring Phylogenies*. Sinauer Associates (2004).
- [13] N. J. Fine. On the Walsh functions. *Transactions of the American Mathematical Society* 65 (1949), 372-414.

- [14] M. Fischer. Extremal values of the Sackin tree balance index. *Annals of Combinatorics* 25 (2021), 515–541.
- [15] M. Fischer, L. Herbst, S. Kersting, L. Kühn, K. Wicke. Tree balance indices: a comprehensive survey. arXiv:2109.12281 (2021).
- [16] M. Fisher, V. Liebscher. On the balance of unrooted trees. *Journal of Graph Algorithms and Applications*, 25 (2021), 133–150.
- [17] G. Fusco, Q. C. Cronk. A new method for evaluating the shape of large phylogenies. *Journal of Theoretical Biology* 175 (1995), 235–243.
- [18] D. J. Futuyma (Ed. Chair). Executive document: Evolution, science and society: Evolutionary biology and the National Research Agenda. *The American Naturalist* 158 (2001), S4.
- [19] R. Graham, D. Knuth, O. Patashnik. *Concrete Mathematics* (2nd edition). Addison-Wesley (1994).
- [20] M. Guedj. A Simple but Powerful Extension of the Master Theorem for Divide-and-Conquer Sequences. hal-03329411 (2021).
- [21] H. K. Hwang. Asymptotic expansion for the Lebesgue constants of the Walsh system. *Journal of Computational and Applied Mathematics* 71 (1996), 237–243.
- [22] H. K. Hwang, S. Janson, T. H. Tsai. Exact and asymptotic solutions of a divide-and-conquer recurrence dividing at half: Theory and applications. *ACM Transactions on Algorithms* 13 (2017), Article 47, DOI: <https://doi.org/10.1145/3127585>.
- [23] M. Kao. Multiple-size divide-and-conquer recurrences. *ACM SIGACT News* 28 (1997), 67–69
- [24] M. Kirkpatrick, M. Slatkin. Searching for evolutionary patterns in the shape of a phylogenetic tree. *Evolution* 47 (1993), 1171–1181.
- [25] A. McKenzie, M. Steel. Distributions of cherries for two models of trees. *Mathematical Biosciences* 164 (2000), 81–92.
- [26] A. Mir, F. Rosselló, L. Rotger. A new balance index for phylogenetic trees. *Mathematical Biosciences*, 241 (2013), 125–136.
- [27] A. Mir, L. Rotger, F. Rosselló. Sound Colless-like balance indices for multifurcating trees. *PLoS ONE* 13 (2018), e0203401.
- [28] J. Mosterín. *Conceptos y teorías en la ciencia*. Alianza Editorial (1984).
- [29] G. Orosi. A Simple Derivation Of Faulhaber’s Formula. *Applied Mathematics E-Notes*, 18 (2018): 124–126.
- [30] S. Roura. Improved Master Theorems for Divide-and-Conquer Recurrences. *Journal of the ACM* 48 (2001), 170–205.
- [31] M. J. Sackin. “Good” and “bad” phenograms. *Systematic Zoology* 21 (1972), 225–226.
- [32] C. Semple and M. Steel. *Phylogenetics*. Oxford University Press (2003).
- [33] K. Shao, R. Sokal. Tree balance. *Systematic Zoology* 39 (1990), 266–276.
- [34] E. Stam. Does imbalance in phylogenies reflect only bias? *Evolution* 56 (2002), 1292–1295.
- [35] R. Stephan. Some Divide and Conquer Sequences with (relatively) Simple Ordinary Generating Functions. OEIS (2014), <https://oeis.org/somedcgf.html>.

- [36] B.C. Thompson. Solving recurrence relations involving floor and ceiling functions. *Electronics Letters*, 30 (1994), 1391–1393.
- [37] R. M. Verma. A general method and a master theorem for divide-and-conquer recurrences with applications. *Journal of Algorithms* 16 (1994), 67–79.
- [38] R. M. Verma. General techniques for analyzing recursive algorithms with applications. *SIAM Journal on Computing*, 26 (1997), 568–581.
- [39] X. Wang, Q. Fu. A frame for general divide-and-conquer recurrences. *Information Processing Letters* 59 (1996), 45–51