

# GEOMETRIC CHARACTERIZATIONS OF THE STRICT HADAMARD DIFFERENTIABILITY OF SETS

ABDERRAHIM JOURANI AND MOUSTAPHA SENE

ABSTRACT. Let  $S$  be a closed subset of a Banach space  $X$ . Assuming that  $S$  is epi-Lipschitzian at  $\bar{x}$  in the boundary  $\text{bdry } S$  of  $S$ , we show that  $S$  is strictly Hadamard differentiable at  $\bar{x}$  IFF the Clarke tangent cone  $T(S, \bar{x})$  to  $S$  at  $\bar{x}$  contains a closed hyperplane IFF the Clarke tangent cone  $T(\text{bdry } S, \bar{x})$  to  $\text{bdry } S$  at  $\bar{x}$  is a closed hyperplane. Moreover when  $X$  is of finite dimension,  $Y$  is a Banach space and  $g : X \mapsto Y$  is a locally Lipschitz mapping around  $\bar{x}$ , we show that  $g$  is strictly Hadamard differentiable at  $\bar{x}$  IFF  $T(\text{graph } g, (\bar{x}, g(\bar{x})))$  is isomorphic to  $X$  IFF the set-valued mapping  $x \mapsto K(\text{graph } g, (x, g(x)))$  is continuous at  $\bar{x}$  and  $K(\text{graph } g, (\bar{x}, g(\bar{x})))$  is isomorphic to  $X$ , where  $K(A, a)$  denotes the contingent cone to a set  $A$  at  $a \in A$ .

## 1. INTRODUCTION

Let  $X$  be a real normed vector space and  $S$  be a closed subset with  $\bar{x} \in S$ . A vector  $v \in X$  is said to be a *Clarke tangent vector* to  $S$  at  $\bar{x}$  if for any neighborhood  $V$  of  $v$  there exist a neighborhood  $U$  of  $\bar{x}$  and some  $\lambda > 0$  such that for all  $(t, x) \in ]0, \lambda[ \times (U \cap S)$  we have

$$(x + tV) \cap S \neq \emptyset.$$

The characterization in terms of sequences can be stated as follows :  $v \in X$  is a Clarke tangent vector to  $S$  at  $\bar{x}$  if and only if for any sequence  $(x_n)_{n \in \mathbb{N}}$  of  $S$  converging to  $\bar{x}$  and any sequence of positive reals  $(t_n)_{n \in \mathbb{N}}$  converging to 0, there exists a sequence  $(v_n)_{n \in \mathbb{N}}$  in  $X$  converging to  $v$  such that  $x_n + t_n v_n \in S$  for all  $n \in \mathbb{N}$  sufficiently large. The collection  $T(S, \bar{x})$  of all such vectors is a closed convex cone which is called the Clarke tangent cone to  $S$  at  $\bar{x}$  (see [7]). When  $\bar{x} \notin S$  one puts,  $T(S, \bar{x}) = \emptyset$ .

The Clarke tangent cone is involved in many geometrical representation of sets such as locally compact and definable sets in  $\mathbb{R}^n$ . For instance in [2] the authors proved that a nonempty locally compact subset  $S \subset \mathbb{R}^n$  is a  $C^1$ -manifold if and only if the Clarke tangent cone and the upper paratangent cone to  $S$  coincide at every point. More historical characterizations of  $C^1$ -manifolds in Euclidean spaces by tangent cones are reviewed in [1]. More

---

2000 *Mathematics Subject Classification*. Primary 49J52, 46N10, 58C20; Secondary 34A60.

*Key words and phrases*. Epi-Lipschitzian set, Compactly epi-Lipschitzian set, Clarke Tangent cone, Contingent cone, Clarke subdifferential.

recently, the authors in [20] have established a geometrical characterization of definable sets in terms of tangent cones.

It is our purpose in this paper to provide geometrical characterizations of the strict Hadamard differentiability of epi-Lipschitz and non epi-Lipschitz sets by tangent cones. Indeed we show that an epi-Lipschitz set  $S$  at  $\bar{x}$  is strictly Hadamard differentiable at  $\bar{x}$  if and only if the Clarke tangent cone to  $S$  at  $\bar{x}$  contains a closed hyperplane if and only if the Clarke tangent cone  $T(\text{bdry } S, \bar{x})$  to the boundary of  $S$  at  $\bar{x}$  is a closed hyperplane. Moreover in finite dimension, we establish that a mapping  $g : X \rightarrow Y$  is strictly Hadamard differentiable at  $\bar{x}$  if and only if  $T(\text{graph } g, (\bar{x}, g(\bar{x})))$  is isomorphic to  $X$ . Further corollaries are provided throughout the paper.

The paper is organized as follows. Section 2 is devoted to some nonsmooth analysis tools. In section 3, we present epi-Lipschitz sets and define the concepts of strictly Hadamard differentiability of sets. Then, we study the inverse image of an epi-Lipschitz set by a strictly Hadamard differentiable mapping. Section 4 and 5 are respectively devoted to the strict Hadamard differentiability of epi-Lipschitz and non epi-Lipschitz sets.

## 2. TOOLS OF NONSMOOTH ANALYSIS

Let  $S$  be a subset of a vector normed space  $X$ . The Clarke normal cone to  $S$  at  $\bar{x} \in S$  is the negative polar of  $T(S, \bar{x})$ , that is

$$N(S, \bar{x}) = \{x^* \in X^* : \langle x^*, h \rangle \leq 0, \forall h \in T(S, \bar{x})\}$$

where  $X^*$  denotes the topological dual space of  $X$  and  $\langle \cdot, \cdot \rangle$  is the pairing between  $X$  and  $X^*$ . Having defined this object, we may now introduce the Clarke subdifferential. Let  $f : X \mapsto \mathbb{R} \cup \{+\infty\}$  be a lower semicontinuous function and  $\bar{x} \in X$ , with  $f(\bar{x}) < \infty$ . The Clarke subdifferential of  $f$  at  $\bar{x}$  is given ([8]) by

$$\partial f(\bar{x}) = \{x^* \in X^* : (x^*, -1) \in N(\text{epi } f, (\bar{x}, f(\bar{x})))\}$$

where  $\text{epi } f$  denotes the epigraph of  $f$ , that is,

$$\text{epi } f = \{(x, r) \in X \times \mathbb{R} : f(x) \leq r\}.$$

It is natural to put  $\partial f(\bar{x}) = \emptyset$  whenever  $f(\bar{x}) = \infty$ . When  $f$  is locally Lipschitz around  $\bar{x}$ , the Clarke subdifferential has the following equivalent definition

$$\partial f(\bar{x}) = \{x^* \in X^* : \langle x^*, h \rangle \leq \limsup_{\substack{t \rightarrow 0^+ \\ x \rightarrow \bar{x}}} \frac{f(x + th) - f(x)}{t} \quad \forall h \in X\}.$$

It is known ([8]), that whenever  $f$  is locally Lipschitz around  $\bar{x}$ , the following equivalence holds true

$$\partial f(\bar{x}) \text{ is a singleton IFF } f \text{ is strictly Hadamard differentiable at } \bar{x}.$$

We recall that a mapping  $g : X \mapsto Y$ , with  $Y$  a vector normed space, is strictly Hadamard differentiable at  $\bar{x}$  if there exists a linear continuous

mapping  $Dg(\bar{x}) : X \rightarrow Y$  such that

$$\lim_{\substack{t \rightarrow 0^+ \\ x \rightarrow \bar{x} \\ u \rightarrow h}} \frac{g(x+tu) - g(x)}{t} = Dg(\bar{x})(h)$$

The mapping  $Dg(\bar{x})$  is called the strict Hadamard derivative of  $g$  at  $\bar{x}$ .

In order to apply Graves Theorem, we need an other concept of differentiability. We recall ([22]) that a mapping  $g : X \mapsto Y$ , with  $Y$  a vector normed space, is strictly Fréchet differentiable at  $\bar{x}$  if

$$\lim_{\substack{x \rightarrow \bar{x} \\ x' \rightarrow \bar{x}}} \frac{g(x) - g(x') - Dg(\bar{x})(x - x')}{\|x - x'\|} = 0$$

where  $Dg(\bar{x})$  denotes the strict Fréchet derivative of  $g$  at  $\bar{x}$ .

It is not difficult to see that each strictly Fréchet differentiable mapping is strictly Hadamard differentiable and both concepts coincide in finite dimensional spaces for locally Lipschitz mappings. This equivalence is no longer true in the infinite dimensional situation. Borwein [5] have established an equivalence of both concepts for locally Lipschitz functions with an additional condition.

### 3. EPI-LIPSCHITZ SETS

Let  $S$  be a subset of a normed space  $X$ . Following Rockafellar (see. [22, 23]),  $S$  is said to be epi-Lipschitzian at  $\bar{x} \in S$  in a direction  $\bar{v} \neq 0$  if there exists  $\gamma > 0$  such that the following inclusion holds

$$S \cap B(\bar{x}, \gamma) + tB(\bar{v}, \gamma) \subset S \quad \forall t \in ]0, \gamma[.$$

If  $S$  is epi-Lipschitzian at any of its points then it is called epi-Lipschitzian. Assume that  $S$  is epi-Lipschitzian at  $\bar{x}$  in a direction  $\bar{v} \neq 0$ . In the case of finite dimensional real normed spaces, Rockafellar (cf. [24, § 4]) proved the existence of a closed hyperplane  $E$  such that  $X = E \oplus \mathbb{R}\bar{v}$ , a neighbourhood  $\Omega$  of  $\bar{x}$  in  $X$  and a function  $f : E \rightarrow \mathbb{R}$  locally Lipschitzian near  $\pi_E(\bar{x})$  (where  $\bar{x} = \pi_E(\bar{x}) + \pi_{\bar{v}}(\bar{x})$  with  $\pi_E(\bar{x}) \in E$  and  $\pi_{\bar{v}}(\bar{x}) \in \mathbb{R}\bar{v}$  uniquely defined) such that

$$\Omega \cap S = \Omega \cap \{u + r\bar{v} : u \in E, r \in \mathbb{R}, f(u) \leq r\}.$$

Further he observed in [25, § 5] that if  $S$  is closed then the arguments in [24] are also true in the case of infinite dimensional real normed space. Having this, one can write

$$(3.1) \quad \Omega \cap S = \Omega \cap \varphi(\text{epi } f) \quad \text{where} \quad \varphi(u, r) := u + r\bar{v}$$

This means that  $S$  can locally be represented as the epigraph of locally Lipschitz function  $f$ .

$$(3.2) \quad \textit{We call } f \textit{ a local representation of } S \textit{ near } \bar{x}.$$

In [24], Rockafellar showed that

$$\text{a closed set } S \subset X \text{ is epi-Lipschitzian at } \bar{x} \text{ IFF } \text{int}T(S, \bar{x}) \neq \emptyset$$

provided that  $X$  is of finite dimension. Borwein and Strojwas [3] established that this characterization remains valid in infinite dimensional Banach spaces provided that  $S$  is compactly epi-Lipschitzian at  $\bar{x}$ . Following Borwein and Strojwas [3], a set  $S \subset X$  is said to be compactly epi-Lipschitzian (CEL) at  $\bar{x}$  if there exist  $\gamma > 0$  and a norm-compact set  $H$  such that

$$(3.3) \quad S \cap B(\bar{x}, \gamma) + tB(0, 1) \subset S + tH, \forall t \in [0, \gamma].$$

It is clear that this class of sets encompasses that of epi-Lipschitz sets. Note that in finite dimensions, every nonempty set is CEL.

Further representations of closed epi-Lipschitz sets in terms of sub-level of Lipschitz functions are established in Czarnecki and Thibault [10]. We recall some of them in the following.

Let  $X$  be a real normed space and  $S \subset X$ . The *signed distance function* to  $S$  is defined by

$$\Delta_S(x) := d_S(x) - d_{S^c}(x) \quad \text{for all } x \in X,$$

where  $S^c = X \setminus S$  and

$$d_S(x) := \inf_{u \in S} \|u - x\|$$

is the distance function to the set  $S$ . It is easily seen that

$$\text{cl}S = \{x \in X : \Delta_S(x) \leq 0\} \quad \text{and} \quad \text{bdry} S = \{x \in X : \Delta_S(x) = 0\}.$$

Moreover it is known (see e.g., [13]) that  $\Delta_S$  is 1-Lipschitzian on  $X$ . Czarnecki and Thibault [10] established the following local sublevel representation.

**Theorem 3.1.** [10] *Let  $S$  be a subset of a real normed space  $X$  and  $\bar{x} \in S \cap \text{bdry} S$ . Assume with  $\bar{S} := \text{cl}S$  that  $\text{int}\bar{S} \cap U \subset S$  for some neighbourhood  $U$  of  $\bar{x}$ . Then, the set  $S$  is epi-Lipschitzian at  $\bar{x}$  if and only if  $0 \notin \partial\Delta_S(\bar{x})$ .*

This allows to state the following known result.

**Corollary 3.2.** *Let  $S$  be a subset of a normed space  $X$  which is epi-Lipschitzian at  $\bar{x} \in S \cap \text{bdry} S$ . Then, the Clarke normal cone of  $S$  at  $\bar{x}$  can be described as*

$$N(S, \bar{x}) = \mathbb{R}_+ \partial\Delta_S(\bar{x}),$$

where  $\mathbb{R}_+ := [0, +\infty[$ .

**Theorem 3.3.** [10] *Let  $g : X \rightarrow \mathbb{R}$  be a Lipschitz continuous function near  $\bar{x}$  of the real normed space  $X$  and let  $S := \{x \in X : g(x) \leq 0\}$ . Assume that  $\bar{x} \in \text{bdry} S$  and  $0 \notin \partial g(\bar{x})$ . Then  $S$  is epi-Lipschitzian at  $\bar{x}$ .*

**Lemma 3.4.** [10] *Let  $S$  be a subset of a normed space  $X$  which is epi-Lipschitzian at  $\bar{x} \in S \cap \text{bdry} S$ . Then, the following equality holds*

$$T(\text{bdry} S, \bar{x}) = T(S, \bar{x}) \cap -T(S, \bar{x}),$$

which yields in particular that  $T(\text{bdry} S, \bar{x})$  is a closed vector subspace of  $X$ .

Using the Ioffe-approximate subdifferential  $\partial_A$  (see [14] and [15]), the author [18] established the following result.

**Proposition 3.5** (Corollary 4.1 [18]). *Let  $X$  be a Banach space and  $g : X \rightarrow \mathbb{R} \cup \{+\infty\}$  be a lower semicontinuous function around  $\bar{x}$ , with  $g(\bar{x}) = 0$ . Let  $S := \{x \in \mathbb{R}^n : g(x) \leq 0\}$ . Assume that  $\bar{x} \in \text{bdry } S$  and  $0 \notin \partial_{Ag}(\bar{x})$ . Then  $S$  is CEL at  $\bar{x}$  provided that  $\text{epig}$  is CEL at  $(\bar{x}, 0)$ .*

It is well known that if  $\bar{x}$  is an interior point of  $S$  then  $T(S, \bar{x}) = X$ . Unfortunately the reverse implication is not true in general. To see this, take  $X$  a separable Hilbert space with orthonormal basis  $(b_n)_n$  and consider (see. [4]) the set  $S = \text{clco}\{\frac{b_n}{2^n}, -\frac{b_n}{2^n}\}$  with  $\bar{x} = 0$ . Then  $\text{int}S = \emptyset$  and  $T(S, \bar{x}) = X$ . It is shown in [16] that the reverse implication holds for the class of compactly epi-Lipschitz (CEL) sets. The main argument used in [16] is based on the following result established in [17].

**Proposition 3.6** (Corollary 2.7 in [17]). *Let  $S \subset X$  be a closed CEL set at  $\bar{x}$  in the boundary of  $S$ . Then there exists  $x^* \in X^*$ , with  $x^* \neq 0$ , such that  $x^* \in N(S, \bar{x})$ .*

Let us prove the following theorem about epi-Lipschitz property of image and inverse image of epi-Lipschitz sets in Banach spaces

**Proposition 3.7** (Image and inverse image of epi-Lipschitz sets). *Let  $X$  and  $Y$  be Banach spaces,  $S \subset Y$  be a closed set and  $\varphi : X \rightarrow Y$  be a strictly Fréchet differentiable mapping at  $\bar{x}$  with a surjective derivative  $D\varphi(\bar{x})$ . Then the set  $C := \varphi^{-1}(S)$  is epi-Lipschitzian at  $\bar{x} \in \varphi^{-1}(S)$  IFF  $S$  is epi-Lipschitz at  $\varphi(\bar{x})$ .*

*Proof.* Suppose that  $C$  is epi-Lipschitzian at  $\bar{x}$ . Since  $\varphi$  is strictly Fréchet differentiable at  $\bar{x}$  with a surjective derivative  $D\varphi(\bar{x})$ , Theorem 1.3 in [11] ensures the existence of  $\alpha > 0$  and  $r > 0$  such that

$$(3.4) \quad B(\varphi(x), t) \subset \varphi(B(x, \alpha t)) \forall x \in B(\bar{x}, r), \forall t \in [0, r].$$

On the other hand since the set  $C := \varphi^{-1}(S)$  is epi-Lipschitzian at  $\bar{x}$  there exist  $\bar{v} \in X$ ,  $\bar{v} \neq 0$ , and  $s > 0$  such that

$$(3.5) \quad C \cap B(\bar{x}, s) + tB(\bar{v}, s) \subset C \forall t \in [0, s]$$

Moreover  $\varphi$  strictly Fréchet differentiable at  $\bar{x}$  implies that for all  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$(3.6) \quad \|\varphi(x) - \varphi(y) - D\varphi(\bar{x})(x - y)\| \leq \varepsilon \|x - y\| \forall x, y \in B(\bar{x}, 2\delta).$$

We may assume that  $\alpha \geq 1$ ,  $r = s$  and  $\alpha(\varepsilon\|\bar{v}\| + 2\delta) \leq \frac{r}{2}$ . Pick  $\delta_1 > 0$ , with  $\alpha\delta_1 \leq \frac{\delta}{2}$ , and let  $y \in B(\varphi(\bar{x}), \delta_1) \cap S$  and  $b \in B(0, \delta) \subset Y$ . Relation (3.4) ensures the existence of  $x \in B(\bar{x}, \alpha\delta_1)$  such that  $y = \varphi(x)$ , so  $x \in C \cap B(\bar{x}, \alpha\delta_1)$ .

Now using (3.6), we obtain

$$\forall t \in ]0, \frac{\delta}{\|\bar{v}\|}], \quad \|\varphi(x + t\bar{v}) - \varphi(x) - tD\varphi(\bar{x})\bar{v}\| \leq \varepsilon t\|\bar{v}\|$$

and hence

$$\varphi(x) + t(D\varphi(\bar{x})\bar{v} + b) \in B(\varphi(x + t\bar{v}), t(\varepsilon\|\bar{v}\|) + \delta).$$

By (3.4), there exists  $b' \in B(0, \alpha(\varepsilon\|\bar{v}\|) + \delta) \subset X$  such that

$$\varphi(x) + t(D\varphi(\bar{x})\bar{v} + b) = \varphi(x + t(\bar{v} + b')).$$

Invoking relation (3.5), we get  $x + t(\bar{v} + b') \in C$ , equivalently,  $\varphi(x + t(\bar{v} + b')) \in S$  and hence

$$\varphi(x) + t(D\varphi(\bar{x})\bar{v} + b) \in S.$$

In summary, we have

$$\forall t \in ]0, \frac{\delta}{\|\bar{v}\|}], \quad B(\varphi(\bar{x}), \delta_1) \cap S + tB(D\varphi(\bar{x})\bar{v}, \delta) \subset S.$$

By taking  $\gamma = \min\{\frac{\delta}{\|\bar{v}\|}, \delta_1\}$  we deduce that  $S$  is epi-Lipschitzian at  $\varphi(\bar{x})$ .

Conversely, suppose that  $S$  is epi-Lipschitzian at  $\varphi(\bar{x})$ . As  $\varphi$  is strictly Fréchet differentiable at  $\bar{x}$  with surjective derivative  $D\varphi(\bar{x})$ , there exists  $\alpha > 0$  such that for all  $\varepsilon > 0$  there exists  $\delta > 0$  satisfying

$$(3.7) \quad \varphi(B(\bar{x}, \delta)) \subset B(\varphi(\bar{x}), \varepsilon),$$

$$(3.8) \quad \|\varphi(x) - \varphi(x') - D\varphi(\bar{x})(x - x')\| \leq \varepsilon\|x - x'\| \forall x, x' \in B(\bar{x}, \delta),$$

and

$$(3.9) \quad B(0, 1) \subset D\varphi(\bar{x})(B(0, \alpha)).$$

By assumption, there exists  $\gamma > 0$  and  $w \in Y$  such that

$$(3.10) \quad \forall t \in ]0, \gamma], \quad B(\varphi(\bar{x}), \gamma) \cap S + tB(w, \gamma) \subset S.$$

Using the surjectivity of  $D\varphi(\bar{x})$  there exists  $v \in X$  such that  $w = D\varphi(\bar{x})(v)$ . So, let  $x \in B(\bar{x}, \delta) \cap C$ ,  $t \in ]0, \delta[$  and  $u \in B(v, \delta)$ . Relations (3.8), (3.7) and (3.9) ensure the following inclusions

$$\begin{aligned} \varphi(x + tu) &\in \varphi(x) + tD\varphi(\bar{x})(u) + B(0, \varepsilon t\|u\|) \quad (\text{by (3.8)}) \\ &\subset B(\varphi(\bar{x}), \varepsilon) \cap S + tD\varphi(\bar{x})(u) + B(0, \varepsilon t\|u\|) \quad (\text{by (3.7)}) \\ &\subset B(\varphi(\bar{x}), \varepsilon) \cap S + tD\varphi(\bar{x})(B(u, \alpha\varepsilon\|u\|)) \quad (\text{by (3.9)}) \\ &\subset B(\varphi(\bar{x}), \varepsilon) \cap S + t(B(D\varphi(\bar{x})(u), \alpha\varepsilon\|u\|) \cdot \|D\varphi(\bar{x})\|) \\ &\subset S \quad (\text{because } \varepsilon \text{ is arbitrary and (3.10)}). \end{aligned}$$

This shows that  $C$  is epi-Lipschitzian at  $\bar{x}$ . The proof is completed.  $\square$

**Remark 3.8.** This proposition tells us that we may replace the function  $\varphi$  in (3.1) by any strictly Fréchet differentiable mapping  $\varphi : E \times \mathbb{R} \rightarrow X$  at  $\bar{y} := (\bar{u}, f(\bar{u})) \in \varphi^{-1}(\bar{x})$  with a surjective derivative  $D\varphi(\bar{y})$ . Where  $S$  is as in the first part of (3.1).

**Remark 3.9.** Note that this proposition is not true even if  $\varphi$  is a linear non surjective operator. Indeed, let ([5])  $X = Y = l^2$  be the Hilbert space of square summable sequences, with  $(e_k)$  its canonical orthonormal base and let the operator  $\varphi : l^2 \rightarrow l^2$  be defined by

$$\varphi\left(\sum x_i e_i\right) = \sum 2^{1-i} x_i e_i.$$

It is easily seen that for all  $i$ ,  $e_i \in \text{Im}\varphi$  and for all  $k \geq 1$ ,  $\sum_{i=1}^k 2^{1-i} e_i \in \text{Im}\varphi$  but  $\sum 2^{1-i} e_i \notin \text{Im}\varphi$  (this shows that  $\text{Im}(\varphi)$  is not closed). Then  $\varphi$  is not surjective. Further  $\text{Im}(\varphi)$  is a proper dense subspace of  $l^2$ . So that the space  $X$  is epi-Lipschitzian at 0, but not  $\varphi(X)$ .

#### 4. STRICT HADAMARD DIFFERENTIABILITY OF EPI-LIPSCHITZ SETS

This section is devoted to the local representation of epi-Lipschitz set as the epigraph of strictly Hadamard differentiable function. As a motivation of the following result we start with the following observation. Let  $S = \text{epi } f$  be the epigraph of the function  $f$  defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

and let  $\bar{x} = (0, 0)$ . Then  $f$  is Lipschitz near 0 and  $T(S, \bar{x}) = \text{epi} |\cdot|$ . This shows that  $T(S, \bar{x})$  does not contain any hyperplane.

In the sequel, we will establish that if  $f$  is Lipschitz near  $\bar{x}$  then  $T(\text{epi } f, (\bar{x}, f(\bar{x})))$  contains a closed hyperplane if and only if  $f$  is strictly Hadamard differentiable at  $\bar{x}$ .

**Definition 4.1.** We say that an epi-Lipschitz set  $S$  at  $\bar{x} \in S$  is strictly Hadamard differentiable at  $\bar{x}$  if its local representation  $f$  is strictly Hadamard differentiable at  $\bar{x}$ .

**Theorem 4.2.** Let  $X$  be a real Banach space and  $S \subset X$  be a closed set containing  $\bar{x}$ . Suppose that  $S$  is epi-Lipschitz at  $\bar{x}$ . Then  $S$  is strictly Hadamard differentiable at  $\bar{x}$  if and only if  $T(S, \bar{x})$  contains a closed hyperplane.

*Proof.* Since  $S$  is epi-Lipschitz at  $\bar{x}$ , there exist

- a closed hyperplane  $E$  and  $\bar{v} \in X \setminus \{0\}$  such that  $X = E \oplus \mathbb{R}\bar{v}$ ,
- a Lipschitz continuous mapping  $f : E \rightarrow \mathbb{R}$  near  $\bar{u} = \pi_E \bar{x}$  and
- a strictly Fréchet differentiable mapping  $\varphi : E \times \mathbb{R} \rightarrow X$  at  $\bar{y} = (\bar{u}, f(\bar{u}))$  with a surjective derivative  $D\varphi(\bar{y})$

such that

$$(4.1) \quad S \cap \Omega = \Omega \cap \varphi(\text{epi } f)$$

Since  $f$  is Lipschitz continuous around  $\bar{u}$ , it is Clarke subdifferentiable at  $\bar{u}$ . Moreover, by (4.1) and the fact that  $D\varphi(\bar{u}, f(\bar{u}))$  is surjective, we have (see for example Theorem 1.17 in [21])

$$(4.2) \quad N(\text{epi } f, (\bar{u}, f(\bar{u}))) = [D\varphi(\bar{y})]^* N(S, \bar{x})$$

where  $[D\varphi(\bar{y})]^*$  denotes the adjoint linear mapping of  $D\varphi(\bar{y})$ .

This implies that

$$(4.3) \quad \partial f(\bar{u}) \times \{-1\} \subset [D\varphi(\bar{y})]^* N(S, \bar{x}).$$

Now suppose that  $T(S, \bar{x})$  contains a closed hyperplane. Let us show that  $f$  is strictly Hadamard differentiable at  $\bar{u}$ .

Taking the polar of  $T(S, \bar{x})$  and using (4.3) we have

$$\partial f(\bar{u}) \times \{-1\} \subset \mathbb{R}_+[D\varphi(\bar{y})]^*(a^*)$$

for some  $a^* \neq 0$ . Let  $[D\varphi(\bar{y})]^*(a^*) = (a_1^*, a_2^*)$  with  $a_1^* \in E^*$  and  $a_2^* \in \mathbb{R}$ , we have

$$\partial f(\bar{u}) \times \{-1\} \subset \mathbb{R}_+(a_1^*, a_2^*).$$

Now for  $u^* \in \partial f(\bar{u})$ , there exists  $\lambda > 0$  such that  $(u^*, -1) = \lambda(a_1^*, a_2^*)$ .

This implies that  $a_2^* = -\frac{1}{\lambda}$ . Moreover  $u^*$  is constant with  $u^* = -\frac{a_1^*}{a_2^*}$ . That is

$$\partial f(\bar{u}) = \left\{ -\frac{a_1^*}{a_2^*} \right\}.$$

Since  $f$  is Lipschitz continuous in an open neighbourhood of  $\bar{u}$  and  $\partial f(\bar{u})$  is a singleton then  $f$  is strictly Hadamard differentiable at  $\bar{u}$ .

Conversely suppose that  $f$  is strictly Hadamard differentiable at  $\bar{u}$ . Let us show that  $T(S, \bar{x})$  contains a closed hyperplane.

Relation (4.2) ensures that  $N(S, \bar{x}) = \mathbb{R}_+[D\varphi(\bar{y})]^{*-1}(\nabla f(\bar{u}), -1)$ . Since  $D\varphi(\bar{y})$  is surjective, the set  $[D\varphi(\bar{y})]^{*-1}(\nabla f(\bar{u}), -1)$  is a singleton. Put  $a^* = [D\varphi(\bar{y})]^{*-1}(\nabla f(\bar{u}), -1)$ . Then  $a^* \neq 0$  and

$$T(S, \bar{x}) = \{h \in X : \langle a^*, h \rangle \leq 0\}$$

which clearly contains a hyperplane.  $\square$

In what follows, we shall prove that the strict Hadamard differentiability of  $S$  at  $\bar{x}$  is equivalent to saying that  $T(\text{bdry } S, \bar{x})$  is a closed hyperplane. We begin with the next lemma.

**Lemma 4.3.** *Let  $S$  be a closed subset of a real Banach (resp. real normed vector) space  $X$ . Assume that  $S$  is CEL (resp. epi-Lipschitzian) at  $\bar{x} \in \text{bdry } S$ . If  $T(S, \bar{x})$  contains a closed hyperplane  $H$ , then  $T(S, \bar{x}) \cap (-T(S, \bar{x})) = H$  (resp.  $T(\text{bdry } S, \bar{x}) = H$ ).*

*Proof.* Let  $X$  be a real Banach space. Suppose that  $S$  is CEL at  $\bar{x}$  and that  $T(S, \bar{x})$  contains a closed hyperplane  $H$ . Assume that there exists  $h \in T(S, \bar{x}) \cap (-T(S, \bar{x}))$  such that  $h \notin H$ . By the Banach strict separation there exist  $x^* \neq 0$  and  $\alpha \neq 0$  such that

$$\langle x^*, h \rangle > \alpha > \langle x^*, v \rangle, \forall v \in H.$$

This implies that  $\langle x^*, v \rangle = 0$  for all  $v \in H$  and  $\alpha > 0$ . Consequently we have  $x^* \in H^\perp$ . Since  $H \subset T(S, \bar{x})$  we have  $N(S, \bar{x}) \subset H^\perp$ . This implies that

$$N(S, \bar{x}) \subset \mathbb{R}x^*.$$

Moreover since  $\bar{x} \in \text{bdry } S$  and  $S$  is CEL at  $\bar{x}$ , Proposition 3.6 ensures that  $N(S, \bar{x}) \neq \{0\}$ . Therefore

$$x^* \in N(S, \bar{x}) \quad \text{or} \quad -x^* \in N(S, \bar{x}).$$



Because  $h \in T(S, \bar{x}) \cap (-T(S, \bar{x}))$ , any one of the two last conditions imply that  $\langle x^*, h \rangle = 0$  which contradicts the fact that  $\alpha > 0$ . Therefore

$$T(S, \bar{x}) \cap (-T(S, \bar{x})) = H.$$

Now let  $X$  be a real normed space. Suppose that  $S$  is epi-Lipschitzian and that  $T(S, \bar{x})$  contains a closed hyperplane  $H$ . Assume that there exists  $h \in T(S, \bar{x}) \cap (-T(S, \bar{x}))$  such that  $h \notin H$ . Following the above arguments with the Banach strict separation theorem there exists some  $x^* \neq 0$  such that

$$N(S, \bar{x}) \subset \mathbb{R}x^*.$$

Moreover Theorem 3.1 and Corollary 3.2 imply that  $N(S, \bar{x}) \neq \{0\}$ . Similar arguments as above give that  $T(S, \bar{x}) \cap (-T(S, \bar{x})) = H$ . Therefore Lemma 3.4 imply that  $T(\text{bdry } S, \bar{x}) = H$ .  $\square$

So, we have the following result which follows immediately from the above lemma and Theorem 4.2.

**Theorem 4.4.** *Let  $S$  be a closed subset of the real Banach space  $X$  which is epi-Lipschitzian at  $\bar{x} \in \text{bdry } S$ . Then  $S$  is strictly Hadamard differentiable at  $\bar{x}$  if and only if  $T(\text{bdry } S, \bar{x})$  is a closed hyperplane.*

We complete this section by the two following corollaries of Theorem 4.2.

**Corollary 4.5.** *Let  $S$  be a closed subset in a real Banach space  $X$ . Assume that  $S$  is epi-Lipschitzian at  $\bar{x} \in \text{bdry } S$ . Then  $S$  is strictly Hadamard differentiable at  $\bar{x}$  if and only if  $\partial\Delta_S(\bar{x})$  is contained in a segment  $]0, a^*]$  for some  $a^* \in X^*$ .*

*Proof.* Suppose that  $S$  is strictly Hadamard differentiable at  $\bar{x}$ . Then by Theorem 4.2, there exists a closed hyperplane  $H$  such that  $H \subset T(S, \bar{x})$ . This implies that

$$N(S, \bar{x}) \subset \mathbb{R}x^* \quad \text{for some } x^* \in X^* \setminus \{0\}.$$

This and Corollary 3.2 imply that

$$\mathbb{R}_+\partial\Delta_S(\bar{x}) \subset \mathbb{R}x^*.$$

Moreover by Theorem 3.1, we have  $0 \notin \partial\Delta_S(\bar{x})$ . This and the fact  $\partial\Delta_S(\bar{x})$  is bounded implies that

$$\partial\Delta_S(\bar{x}) \subset ]0, a^*] \quad \text{for some } a^* \neq 0.$$

Conversely assume that  $\partial\Delta_S(\bar{x}) \subset ]0, a^*]$  for some  $a^* \neq 0$ . Then we have

$$\mathbb{R}_+\partial\Delta_S(\bar{x}) \subset \mathbb{R}a^*.$$

This implies that

$$\langle a^* \rangle^\perp \subset T(S, \bar{x}).$$

That is  $T(S, \bar{x})$  contains a closed hyperplane. Therefore by Theorem 4.2,  $S$  is strictly Hadamard differentiable at  $\bar{x}$ .  $\square$

**Corollary 4.6.** *Let  $g : X \rightarrow \mathbb{R}$  be a Lipschitz continuous function near  $\bar{x}$  in the real Banach space  $X$  and let  $S := \{x \in X : g(x) \leq 0\}$ . Assume that  $\bar{x} \in \text{bdry } S$  and  $0 \notin \partial g(\bar{x})$ . Then  $S$  is strictly Hadamard differentiable at  $\bar{x}$  if and only if there exists some  $a^* \neq 0$  such that*

$$\partial g(\bar{x}) \subset ]0, a^*].$$

*Proof.* Suppose that  $S$  is strictly Hadamard differentiable. By Theorem 3.3, we have  $S$  is epi-Lipschitzian at  $\bar{x}$ . Therefore by Theorem 4.2, there exists a closed hyperplane  $H$  such that  $H \subset T(S, \bar{x})$ . This implies that

$$N(S, \bar{x}) \subset \mathbb{R}x^* \quad \text{for some } x^* \in X^* \setminus \{0\}.$$

That is

$$\mathbb{R}_+ \partial g(\bar{x}) \subset \mathbb{R}x^*.$$

Since  $\partial g(\bar{x})$  is bounded and not containing 0 we have

$$\partial g(\bar{x}) \subset ]0, a^*] \quad \text{for some } a^* \in X^* \setminus \{0\}.$$

Conversely if  $\partial g(\bar{x}) \subset ]0, a^*]$  for some  $a^* \neq 0$ . Since  $0 \notin \partial g(\bar{x})$ , Theorem 3.3 implies that  $S$  is epi-Lipschitzian at  $\bar{x}$ . Moreover

$$N(S, \bar{x}) \subset \mathbb{R}a^*.$$

This implies that  $T(S, \bar{x})$  contains a closed hyperplane. Therefore Theorem 4.2 implies that  $S$  is strictly Hadamard differentiable at  $\bar{x}$ .  $\square$

## 5. STRICT HADAMARD DIFFERENTIABILITY OF NON EPI-LIPSCHITZ SETS

This section is concerned with the strict Hadamard differentiability of non epi-Lipschitz sets in finite dimensions. We start with the next theorem.

**Theorem 5.1.** *Let  $X$  be a finite dimensional real vector space and  $S \subset X$  be a closed subset. Assume that  $S$  is non epi-Lipschitzian at  $\bar{x} \in S$  and that  $T(S, \bar{x})$  contains a closed hyperplane  $H$ . Then  $T(S, \bar{x})$  coincides with  $H$ .*

*Proof.* Assume that  $\text{Int } T(S, \bar{x}) = \emptyset$  and that there exists a closed hyperplane  $H$  such that  $H \subset T(S, \bar{x})$ . Let  $h \in T(S, \bar{x})$  such that  $h \notin H$ . By the Banach separation theorem there exists  $x^* \in X$  and  $\alpha \in \mathbb{R}$  such that

$$(5.1) \quad \langle x^*, h \rangle > \alpha \geq \langle x^*, u \rangle, \quad \forall u \in H.$$

This implies that

$$(5.2) \quad \langle x^*, u \rangle = 0, \quad \forall u \in H \quad \text{and} \quad \alpha \geq 0.$$

From (5.1) and (5.2) we have respectively:

$$x^* \notin N(S, \bar{x}) \quad \text{and} \quad x^* \in H^\perp.$$

Let  $a^* \neq 0$  be a vector in  $X$  such that  $H^\perp = \mathbb{R}a^*$ . Assume without loss of generality that  $x^* \in \mathbb{R}_+ a^*$ . Since  $N(S, \bar{x}) \subset H^\perp$  and  $x^* \notin N(S, \bar{x})$ , we deduce that  $N(S, \bar{x}) = \mathbb{R}_- a^*$ . Consequently

$$T(S, \bar{x}) = \{h \in X : \langle a^*, h \rangle \geq 0\}.$$

This is a contradiction because  $\text{Int } T(S, \bar{x}) = \emptyset$ . Therefore  $T(S, \bar{x}) = H$ .  $\square$

The following theorem gives a geometric characterization of the strict Hadamard differentiability of mappings.

**Theorem 5.2.** *Let  $X$  be a finite dimensional real vector space and  $Y$  a real Banach space. Let  $g : X \mapsto Y$  be a mapping which is locally Lipschitzian around  $\bar{x}$  and  $\varphi : X \times Y \rightarrow Z$  be a strictly Hadamard differentiable mapping at  $(\bar{x}, \bar{y})$  with a bijective derivative  $D\varphi(\bar{x}, \bar{y})$  where  $Z$  is a Banach space and  $\bar{y} = g(\bar{x})$ . Let  $S \subset Z$  be a closed set such that  $S \cap \Omega = \varphi(\text{graph } g) \cap \Omega$  with  $\Omega \subset Z$  an open set containing  $\bar{z} := \varphi(\bar{x}, \bar{y})$ . Then the following assertions are equivalent*

- 1)  $g$  is strictly Hadamard differentiable at  $\bar{x}$ ,
- 2)  $T(S, \bar{z})$  is isomorphic to  $X$ .

*Proof.* 1)  $\Rightarrow$  2) : Suppose that  $g$  is strictly Hadamard differentiable at  $\bar{x}$ . Then the following equality holds

$$(5.3) \quad T(\text{graph } (g), (\bar{x}, g(\bar{x}))) = \text{graph } Dg(\bar{x}).$$

Indeed, let  $(h, k) \in T(\text{graph } (g), (\bar{x}, g(\bar{x})))$ . Then for all  $t_n \rightarrow 0^+$  there exists  $h_n \rightarrow h$  and  $k_n \rightarrow k$  such that

$$g(\bar{x} + t_n h_n) = g(\bar{x}) + t_n k_n, \text{ for } n \text{ sufficiently large.}$$

Using the differentiability and the local Lipschitzness of  $g$  at  $\bar{x}$ , we get

$$Dg(\bar{x})h = \lim_{n \rightarrow +\infty} \frac{g(\bar{x} + t_n h) - g(\bar{x})}{t_n} = k.$$

Conversely, let  $h \in X$ ,  $(x_n) \subset X$ , with  $\lim_{n \rightarrow +\infty} x_n = \bar{x}$  and  $t_n \rightarrow 0^+$ . Then, by the strict Hadamard differentiability of  $g$

$$k := Dg(\bar{x})h = \lim_{n \rightarrow +\infty} \frac{g(x_n + t_n h) - g(x_n)}{t_n}.$$

Put  $k_n = \frac{g(x_n + t_n h) - g(x_n)}{t_n}$ . Then  $(x_n, g(x_n)) + t_n(h, k_n) \in \text{graph } g$  for all  $n$ . Thus,  $(h, k) \in T(\text{graph } g, (\bar{x}, g(\bar{x})))$ .

Since  $D\varphi(\bar{x}, \bar{y})$  is bijective, we have

$$(5.4) \quad T(S, \varphi(\bar{x}, \bar{y})) = D\varphi(\bar{x}, \bar{y})(T(\text{graph } (g), (\bar{x}, g(\bar{x}))))).$$

Now we define the mapping  $\psi : X \mapsto T(S, \varphi(\bar{x}, \bar{y}))$  by

$$\psi(h) = D\varphi(\bar{x}, \bar{y})(h, Dg(\bar{x})h).$$

Using relation (5.3), it is easy to see that  $\psi$  is an isomorphism from  $X$  into  $T(S, \varphi(\bar{x}, \bar{y}))$ .

2)  $\Rightarrow$  1): Let  $\xi : X \mapsto T(S, \varphi(\bar{x}, \bar{y}))$  be an isomorphism. Using relation (5.4), the mapping  $\psi := D\varphi(\bar{x}, \bar{y})^{-1} \circ \xi$  is an isomorphism from  $X$  to  $T(\text{graph } (g), (\bar{x}, g(\bar{x})))$ . Let  $\psi_1 : X \mapsto X$  and  $\psi_2 : X \mapsto Y$  be two linear mappings such that  $\psi(x) = (\psi_1(x), \psi_2(x))$  for all  $x \in X$ . We have for all  $h \in X$ ,  $(\psi_1(h), \psi_2(h)) \in T(\text{graph } (g), (\bar{x}, g(\bar{x})))$ . Let  $h \in X$ ,  $x_n \rightarrow \bar{x}$

and  $t_n \rightarrow 0^+$ . Then there exist  $h_n \rightarrow \psi_1(h)$  and  $k_n \rightarrow \psi_2(h)$  such that  $(x_n, g(x_n)) + t_n(h_n, k_n) \in \text{graph } g$ , for all integer  $n$  sufficiently large. So

$$(5.5) \quad \lim_{n \rightarrow +\infty} \frac{g(x_n + t_n \psi_1(h)) - g(x_n)}{t_n} = \psi_2(h).$$

We claim that the kernel,  $\ker \psi_1$ , of  $\psi_1$  is reduced to the singleton  $\{0\}$ . Because  $\psi$  is an isomorphism, it is easy to see that  $\ker \psi_1 \cap \ker \psi_2 = \{0\}$ . Let  $h \in \ker \psi_1$ . Then relation (5.5) ensures that  $\psi_2(h) = 0$  and hence  $h = 0$ . This means that  $\psi_1$  is an isomorphism. Put  $\gamma = \psi_2 \circ \psi_1^{-1}$ . We claim that  $\gamma$  is the strict Hadamard derivative of  $g$  at  $\bar{x}$ , that is,

$$\lim_{\substack{x \rightarrow \bar{x} \\ x' \rightarrow \bar{x}}} \frac{g(x) - g(x') - \gamma(x - x')}{\|x - x'\|} = 0.$$

So suppose the contrary, that is, there exists  $\varepsilon > 0$  such that

$$\limsup_{\substack{x \rightarrow \bar{x} \\ x' \rightarrow \bar{x}}} \left\| \frac{g(x) - g(x') - \gamma(x - x')}{\|x - x'\|} \right\| \geq \varepsilon.$$

Let  $u_n \rightarrow \bar{x}$  and  $u'_n \rightarrow \bar{x}$  be such that

$$\lim_{n \rightarrow +\infty} \left\| \frac{g(u_n) - g(u'_n) - \gamma(u_n - u'_n)}{\|u_n - u'_n\|} \right\| \geq \varepsilon.$$

Take  $t_n := \|u_n - u'_n\|$  and  $w_n = \frac{u_n - u'_n}{\|u_n - u'_n\|}$ . Since  $X$  is of finite dimension, extracting subsequence if necessary, we may assume that  $w_n \rightarrow w$ , with  $\|w\| = 1$ . So

$$\lim_{n \rightarrow +\infty} \left\| \frac{g(u'_n + t_n w_n) - g(u'_n) - t_n \gamma(w_n)}{t_n} \right\| \geq \varepsilon$$

or equivalently (because  $g$  is locally Lipschitzian around  $\bar{x}$ )

$$\lim_{n \rightarrow +\infty} \left\| \frac{g(u'_n + t_n w) - g(u'_n) - t_n \gamma(w_n)}{t_n} \right\| \geq \varepsilon.$$

Since  $\psi_1$  is an isomorphism, there exists a unique  $h \in X$  such that  $w = \psi_1(h)$ . Using relation (5.5), we obtain a contradiction with the last inequality. The proof is then completed.  $\square$

Let us stress in the following remark an equivalence result of the strict Hadamard differentiability for sets in finite dimensional spaces regardless of whether the set is epi-Lipschitzian or not.

**Remark 5.3.** Let  $X$  be a finite dimensional space and  $g : X \mapsto \mathbb{R}$  be a locally Lipschitz function around  $\bar{x}$ . Then  $g$  is strictly Hadamard differentiable at  $\bar{x}$  IFF  $T(\text{graph } g, (\bar{x}, g(\bar{x})))$  is isomorphic to  $X$  and here  $X \times \{0\}$  is considered as an hyperplane of  $X \times \mathbb{R}$ .

Therefore, let  $S \subset X$  be closed subset having  $g$  as its local representation. Then  $S$  is strictly Hadamard differentiable at  $\bar{x}$  if and only if  $T(S, \bar{x})$  is isomorphic to  $X$ .

Now, we may establish a geometrical characterization in the spirit of [2] involving the contingent or Bouligand cone. We recall that the contingent cone to a closed set  $S$  at  $\bar{x} \in S$  is defined as the following limit superior of the set-differential quotient

$$K(S, \bar{x}) := \limsup_{t \rightarrow 0^+} \frac{S - \bar{x}}{t}.$$

When  $X$  is a finite dimensional space the contingent cone and the Clarke tangentcone are connected in the following way(see [9] and [23])

$$(5.6) \quad T(S, \bar{x}) = \liminf_{\substack{S \\ x \rightarrow \bar{x}}} K(S, x).$$

We say that a set-valued mapping  $F : X \rightrightarrows Y$  is continuous at  $\bar{x}$  if

$$\lim_{x \rightarrow \bar{x}} F(x) = F(\bar{x})$$

where

$$\lim_{x \rightarrow \bar{x}} F(x) := \limsup_{x \rightarrow \bar{x}} F(x) = \liminf_{x \rightarrow \bar{x}} F(x).$$

Here

$$\limsup_{x \rightarrow \bar{x}} F(x) := \{h \in X : \liminf_{x \rightarrow \bar{x}} d_{F(x)}(h) = 0\}$$

and

$$\liminf_{x \rightarrow \bar{x}} F(x) := \{h \in X : \lim_{x \rightarrow \bar{x}} d_{F(x)}(h) = 0\}.$$

**Corollary 5.4.** *Let  $X, Y$  and  $Z$  be finite dimensional spaces,  $g : X \mapsto Y$  be a mapping which is locally Lipschitzian around  $\bar{x}$  and  $\varphi : X \times Y \rightarrow Z$  be a strictly Hadamard differentiable mapping at  $(\bar{x}, \bar{y})$  with a bijective derivative  $D\varphi(\bar{x}, \bar{y})$  and put  $\bar{y} = g(\bar{x})$ . Let  $S \subset Z$  be a closed set such that  $S \cap \Omega = \varphi(\text{graph } g) \cap \Omega$ , where  $\Omega \subset Z$  is an open set containing  $\bar{z} := \varphi(\bar{x}, \bar{y})$ . Then the following assertions are equivalent*

- 1)  $g$  is strictly Hadamard differentiable at  $\bar{x}$ ,
- 2') The set-valued mapping  $z \rightrightarrows K(S, z)$  is continuous at  $\bar{z}$  with respect to  $S$  and  $K(S, \bar{z})$  is isomorphic to  $X$ .

*Proof.* Using relation (5.6), it is easy to see that 2') implies the assertion 2) of Theorem 5.2. So it is enough to prove the implication 1)  $\implies$  2'). Without loss of generality, we may assume that  $\varphi = id_{X \times Y}$  and  $Z = X \times Y$ , where  $id_{X \times Y}$  is the identity mapping of  $X \times Y$ . It is easy to see that

$$(5.7) \quad K(\text{graph } g, (\bar{x}, g(\bar{x}))) = \text{graph } Dg(\bar{x}).$$

We will establish the following inclusions

$$\limsup_{x \rightarrow \bar{x}} K(\text{graph } g, (x, g(x))) \subset K(\text{graph } g, (\bar{x}, g(\bar{x}))) \subset \liminf_{x \rightarrow \bar{x}} K(\text{graph } g, (x, g(x))).$$

So let  $(u, v) \in \limsup_{x \rightarrow \bar{x}} K(\text{graph } g, (x, g(x)))$ . Then there are sequences  $x_n \rightarrow$

$\bar{x}$  and  $(u_n, v_n) \rightarrow (u, v)$  such that for  $n$  large enough  $(u_n, v_n) \in K(\text{graph } g, (x_n, g(x_n)))$ .

Thus for each integer  $n$  there exist sequences  $t_k^n \rightarrow 0^+$  and  $(u_k^n, v_k^n) \rightarrow (u_n, v_n)$  such that  $(x_n, g(x_n)) + t_k^n(u_k^n, v_k^n) \in \text{graph } g$  for all  $k$  sufficiently

large. Extracting a diagonal subsequence, we may assume that for some subsequence of integers  $(k_n)_n$ , we have  $\lim_{n \rightarrow +\infty} t_{k_n}^n = 0$ ,  $\lim_{n \rightarrow +\infty} (u_{k_n}^n, v_{k_n}^n) = (u, v)$  and  $(x_n, g(x_n)) + t_{k_n}^n (u_{k_n}^n, v_{k_n}^n) \in \text{graph } g$  for all  $n$  sufficiently large. Thus, since  $g$  is strictly Hadamard differentiable at  $\bar{x}$ ,

$$Dg(\bar{x})u = \lim_{n \rightarrow +\infty} \frac{g(x_n + t_{k_n}^n u_{k_n}^n) - g(x_n)}{t_{k_n}^n} = v.$$

This last equality ensures that  $(u, v) \in K(\text{graph } g, (\bar{x}, g(\bar{x})))$ .

To prove the second inclusion

$$K(\text{graph } g, (\bar{x}, g(\bar{x}))) \subset \liminf_{x \rightarrow \bar{x}} K(\text{graph } g, (x, g(x))),$$

it suffices to use relations (5.7), (5.6) and (5.3).  $\square$

**Acknowledgment.** The authors would like to thank the referee for useful comments which led to an improvement of their original manuscript. Abderrahim Jourani is partially supported by the EIPHI Graduate School (contract ANR-17-EURE-0002).

#### REFERENCES

- [1] F. Bigolin and S. N. Golo, *A historical account on characterizations of  $C^1$ -manifolds in Euclidean spaces by tangent cones*, J. Math. Anal. Appl., 412(2014) 63-76.
- [2] F. Bigolin and G.H. Greco, *Geometric characterizations of  $C^1$ -manifolds in Euclidean spaces by tangent cones*, J. Math. Anal. Appl., 396 (2012) 145-163.
- [3] J. M. Borwein and H. M. Strojwas, *Proximal analysis and boundaries of closed sets in Banach spaces part II: Applications*, Can. J. Math., 39 (1987) 428-472.
- [4] J. M. Borwein, *Epi-Lipschitz-Like sets in Banach space: Theorems and examples*, Nonlinear Analysis TMA, 11 (1987)1207-1217.
- [5] J. M. Borwein, *Minimal cuscos and subgradients of Lipschitz functions*, Fixed Point Theory and its Applications, (J.-B. Baillon and M. Thera, eds.), Pitman Lecture Notes in Math., vol. 252, Longman, Essex, 1991.
- [6] M. Brokate, *A regularity condition for optimization in Banach spaces : Counter examples*, Appl. Math. Optim., 6 (1980), 189-192.
- [7] F. H. Clarke, *Optimization and nonsmooth analysis*, Wiley-interscience, New York, 1983.
- [8] F. H. Clarke, *Generalized gradients and applications*, Trans. Amer. Math. Soc., 205 (1975) 247-262.
- [9] B. Cornet, *Contributions à la théorie mathématique de mécanisme dynamiques d'allocation des ressources*. Thèse, Université de Paris IX, 1981.
- [10] M. O. Czarnecki and L. Thibault, *Sublevel representation of epi-Lipschitz sets and other properties*, Math. Prog., 168 (2018) 555-569
- [11] A. L. Dontchev, *The Graves Theorem Revisited*, J. Convex Anal., 3 (1996) 45-53.
- [12] M. Ghomi and R. Howard (2013), *Tangent cones and regularity of real hypersurfaces*, Journal für die reine und angewandte Mathematik (Crelles Journal), 2014(697), pp. 221-247. Retrieved 17 Sep. 2018, from doi:10.1515/crelle-2013-0091.
- [13] J. -B. Hiriart-Urruty, *Tangent cones, generalized gradients and mathematical programming in Banach spaces*, Math. Oper. Res., 4 (1979) 79-97.
- [14] A. D. Ioffe, *Approximate subdifferentials and applications 2: Functions on locally convex spaces*, Mathematika, 33 (1986) 111-128.

- [15] A. D. Ioffe, *Approximate subdifferentials and applications 3: Metric theory*, *Matematika*, 36 (1989) 1-38
- [16] A. Jourani, *The role of locally compact cones in nonsmooth analysis*, *Communications on Applied Nonlinear Analysis*, 5 (1998) 1-35.
- [17] A. Jourani, *Necessary conditions for extremality and separation theorems with application to multiobjective optimization*, *Optimization*, 44 (1998) 327-350
- [18] A. Jourani, *On a class of compactly epi-Lipschitzian sets*, *Nonlinear Analysis TMA*, 54 (2003) 471-483
- [19] Jourani A. and M. Sène, *Characterization of the Clarke regularity of subanalytic sets*, *Proc. Amer. Math. Soc.*, 146 (2018) 1639-1649.
- [20] K. Kurdyka , Olivier Le Gal, V. N. Nguyen Xuan, *Tangent cones and  $C^1$  regularity of definable set*, *J. Math. Anal. Appl.*, 457 (2018) 978–990.
- [21] B. S. Mordukhovich, *Variational Analysis and Generalized Differentiation I : Basic Theory*, Grundlehren vol 333, 331 Springer, Berlin (2006).
- [22] G. Peano, *Sur la définition de la dérivée*, *Mathesis*, 2 (1892) 12-14.
- [23] J. P. Penot, *A characterization of tangential regularity*, *Nonlinear Anal. TMA*, 5 (1981), 625-643.
- [24] R. T. Rockafellar, *Clarke's tangent cone and the boundaries of closed sets in  $\mathbb{R}^n$* , *Nonlinear Anal. TMA*, 3 (1979) 145-154.
- [25] R. T. Rockafellar, *Generalized direction derivatives and subgradients of nonconvex functions*, *Can. J. Math.*, 32 (1980) 257-280.

(A. Jourani), UNIVERSITÉ DE BOURGOGNE FRANCHE-COMTÉ, INSTITUT DE MATHÉMATIQUES DE BOURGOGNE, UMR 5584, CNRS, , 21000 DIJON, FRANCE,  
*Email address:* `abderrahim.jourani@u-bourgogne.fr`

(M. Sene) MOUSTAPHA SÈNE, UNIVERSITÉ GASTON BERGER, DÉPARTEMENT DE MATHÉMATIQUES, SAINT-LOUIS, SÉNÉGAL,  
*Email address:* `moustapha2.sene@ugb.edu.sn`