

Minimizers of L^2 -Subcritical Inhomogeneous Variational Problems with A Spatially Decaying Nonlinearity

Yongshuai Gao*, Yujin Guo[†] and Shuang Wu[‡]

*School of Mathematics and Statistics,
Hubei key Laboratory of Mathematical Sciences,
Central China Normal University,
Wuhan 430079, People's Republic of China*

December 2, 2021

Abstract

We study the minimizers of L^2 -subcritical inhomogeneous variational problems with spatially decaying nonlinear terms, which contain $x = 0$ as a singular point. The limit concentration behavior of minimizers is proved as $M \rightarrow \infty$ by establishing the refined analysis of the spatially decaying nonlinear term.

Keywords: L^2 -subcritical variational problems; Spatially decaying nonlinearity; Minimizers; Mass concentration

1 Introduction

In this paper, we consider the minimizers of the following L^2 -subcritical constraint inhomogeneous variational problem

$$I(M) := \inf_{\{u \in \mathcal{H}, \|u\|_2^2 = 1\}} E_M(u), \quad M > 0, \quad (1.1)$$

where the Gross-Pitaevskii (GP) energy functional $E_M(u)$ contains a spatially decaying nonlinearity and is defined by

$$E_M(u) := \int_{\mathbb{R}^N} (|\nabla u|^2 + V(x)|u|^2) dx - \frac{2M^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|u|^{p+1}}{|x|^b} dx, \quad N \geq 1, \quad (1.2)$$

*Email: ysgao@mails.ccnu.edu.cn.

[†]Email: yguo@ccnu.edu.cn. Y. J. Guo is partially supported by NSFC under Grant 11931012.

[‡]Email: swu@mails.ccnu.edu.cn.

and the space \mathcal{H} is defined as

$$\mathcal{H} := \left\{ u(x) \in H^1(\mathbb{R}^N) : \int_{\mathbb{R}^N} V(x)|u(x)|^2 < \infty \right\}$$

with the associated norm $\|u\|_{\mathcal{H}} = \left\{ \int_{\mathbb{R}^N} (|\nabla u(x)|^2 + |u(x)|^2 + V(x)|u(x)|^2) dx \right\}^{\frac{1}{2}}$. Here positive constants $b > 0$ and $p > 0$ of (1.1) satisfy

$$0 < b < \min\{2, N\}, \quad 1 < p < 1 + \frac{4-2b}{N}, \quad \text{where } N \geq 1, \quad (1.3)$$

so that $E_M(u)$ admits $x = 0$ as a singular point in its nonlinear term. We always assume that the trapping potential $V(x) \geq 0$ satisfies

(V). $V(x) \in L_{loc}^\infty(\mathbb{R}^N) \cap C_{loc}^\alpha(\mathbb{R}^N)$ with $\alpha \in (0, 1)$, $\{x \in \mathbb{R}^N : V(x) = 0\} = \{0\}$ and $\lim_{|x| \rightarrow \infty} V(x) = \infty$.

The variational problem (1.1) arises in various physical contexts, including the propagation of a laser beam in the optical fiber, Bose-Einstein condensates (BECs), and nonlinear optics (cf. [1, 3, 28]), where the constant $M > 0$ often represents the attractive interaction strength, and $V(x) \geq 0$ denotes an external potential. The variational problem (1.1) and its associated elliptic equation have attracted a lot of attentions over the past few years, due to the appearance of the singular point $x = 0$ in the nonlinear term, see [2, 8, 9, 12, 13, 15, 30] and the references therein.

When $b = 0$, (1.1) is a homogeneous constraint variational problem, for which there are many existing results (1.1) (cf. [6, 14, 18–20, 22, 26, 27, 29, 34]), including the existence and nonexistence of minimizers, and their quantitative properties of all kinds. More precisely, when $p > 1 + \frac{4}{N}$, one can use the energy estimates to obtain the nonexistence of minimizers for (1.1) with $b = 0$ as soon as $M > 0$ (cf. [6, 7]), which is essentially in the L^2 -supercritical case. However, if $p = 1 + \frac{4}{N}$, then (1.1) with $b = 0$ reduces to the L^2 -critical case, which was addressed widely by the second author and his collaborators, see [18–20, 22] and the references therein. As for the case where $1 < p < 1 + \frac{4}{N}$, (1.1) with $b = 0$ is in the L^2 -subcritical case and admits generally minimizers for all $M \in (0, \infty)$. In this case, the uniqueness, symmetry breaking and concentration behavior of minimizers were investigated recently as $M \rightarrow \infty$, see [21, 26, 29] and the references therein.

When $b \neq 0$, the variational problem (1.1) contains the inhomogeneous nonlinear term $m(x)|u|^{p+1}$, where $m(x) = \frac{1}{|x|^b}$ admits $x = 0$ as a singular point. We remark that the inhomogeneous L^2 -constraint variational problems were analyzed recently in [10, 11, 29] and the references therein. However, as far as we know, the above mentioned works handle mainly with the inhomogeneous nonlinear term $m(x)|u|^{p+1}$ where $m(x)$ satisfies $m(x) \in L^\infty(\mathbb{R}^N)$ without any singular point. On the other hand,

Ardila and Dinh obtained recently in [2] the existence of minimizers and the stability of the standing waves, for which they studied the associated constraint variational problem (1.1), in the L^2 -subcritical case where the harmonic potential satisfies $V(x) = \gamma^2|x|^2$ ($\gamma > 0$), $b > 0$ and $p > 0$ satisfy (1.3).

Under the assumptions (V) and (1.3), we comment that it is standard to obtain the existence of minimizers for $I(M)$ for all $M > 0$, see [2, Theorem 1.8] and the related argument. Motivated by above mentioned works, in this paper we mainly study the limit behavior of minimizers u_M for $I(M)$ as $M \rightarrow \infty$, and the main purpose of this paper is to investigate the impact of the singular point $x = 0$ on the behavior of u_M as $M \rightarrow \infty$.

We now assume that u_M is a minimizer of $I(M)$ for any $M > 0$. It then follows from the variational theory that u_M satisfies the following Euler-Lagrange equation

$$-\Delta u_M + V(x)u_M - M^{\frac{p-1}{2}} \frac{u_M^p}{|x|^b} = \mu_M u_M \quad \text{in } \mathbb{R}^N, \quad (1.4)$$

where $\mu_M \in \mathbb{R}^N$ is a suitable Lagrange multiplier associated to u_M . By the form of the energy functional $E_M(\cdot)$, one can obtain from [25, Theorem 6.17] that $E_M(u) = E_M(|u|)$ holds for any $u \in \mathcal{H}$, which implies that $|u_M|$ is also a minimizer of $I(M)$. By the strong maximum principle, one can further derive from (1.4) that $|u_M| > 0$ holds in \mathbb{R}^N . Therefore, u_M must be either positive or negative. Without loss of generality, in the following we only consider positive minimizers $u_M > 0$ of $I(M)$.

Under the assumption (1.3), we next recall the following sharp Gagliardo-Nirenberg (GN) inequality (cf. [13, Theorem 1.2]):

$$\int_{\mathbb{R}^N} \frac{|u|^{p+1}}{|x|^b} dx \leq C_{GN}^{-1} \|\nabla u\|_2^{\frac{N(p-1)+b}{2}+b} \|u\|_2^{p+1-\frac{N(p-1)}{2}-b}, \quad u \in H^1(\mathbb{R}^N), \quad (1.5)$$

where $C_{GN} > 0$ satisfies

$$C_{GN} = \left(\frac{N(p-1) + 2b}{2(p+1) - N(p-1) - 2b} \right)^{\frac{N(p-1)+2b}{4}} \frac{2(p+1) - N(p-1) - 2b}{2(p+1)} \|w\|_2^{p-1}, \quad (1.6)$$

and w is the unique positive radially symmetric solution (cf. [4, 15, 16, 24, 33]) of

$$-\Delta w + w - \frac{w^p}{|x|^b} = 0 \quad \text{in } \mathbb{R}^N, \quad w \in H^1(\mathbb{R}^N). \quad (1.7)$$

The equality in (1.5) is achieved at $u = w$. Moreover, w satisfies the following Pohozaev identity

$$\|\nabla w\|_2^2 = \frac{N(p-1) + 2b}{2(p+1)} \int_{\mathbb{R}^N} \frac{|w|^{p+1}}{|x|^b} dx = \frac{N(p-1) + 2b}{2(p+1) - N(p-1) - 2b} \|w\|_2^2. \quad (1.8)$$

Recall also from [15, Theorem 2.2] that there exist positive constants $\delta > 0$ and $C > 0$ such that $w(x)$ satisfies

$$w(x), |\nabla w(x)| \leq Ce^{-\delta|x|} \text{ as } |x| \rightarrow \infty. \quad (1.9)$$

All above properties of w are often used in the refined analysis of minimizers for $I(M)$ as $M \rightarrow \infty$.

Using above notations, the main result of the present paper can be stated as the following theorem.

Theorem 1.1. *Under the assumptions (V) and (1.3), let u_k be a positive minimizer of $I(M_k)$, where $M_k \rightarrow \infty$ as $k \rightarrow \infty$. Then there exists a subsequence, still denoted by $\{u_k\}$, of $\{u_k\}$ such that u_k satisfies*

$$w_k(x) := \epsilon_k^{\frac{N}{2}} u_k(\epsilon_k x) \rightarrow \frac{w(x)}{\sqrt{a^*}} \text{ uniformly in } L^\infty(\mathbb{R}^N) \text{ as } k \rightarrow \infty, \quad (1.10)$$

where $\epsilon_k := \left(\frac{M_k}{a^*}\right)^{-\frac{p-1}{4-N(p-1)-2b}} > 0$, $a^* := \|w\|_2^2 > 0$, and $w > 0$ is the unique positive solution of (1.7). Moreover, u_k decays exponentially in the sense that for sufficiently large $k > 0$,

$$w_k(x) \leq Ce^{-\sqrt{\theta}|x|} \text{ and } |\nabla w_k(x)| \leq Ce^{-\theta|x|} \text{ as } |x| \rightarrow \infty, \quad (1.11)$$

where $0 < \theta < 1$ and $C > 0$ are independent of $k > 0$.

The proof of Theorem 1.1 shows essentially that as $M_k \rightarrow \infty$, u_k prefers to concentrate near the singular point $x = 0$ of $I(M_k)$, instead of a minimum point for $V(x)$. The proof of Theorem 1.1 depends on the refined estimate of $\int_{\mathbb{R}^N} V(x)|u_k|^2 dx$ as $k \rightarrow \infty$, for which we shall consider the following constraint variational problem without the trap:

$$\tilde{I}(M) := \inf_{\{u \in H^1(\mathbb{R}^N), \|u\|_2^2 = 1\}} \tilde{E}_M(u), \quad N \geq 1, \quad (1.12)$$

where $\tilde{E}_M(u)$ is defined by

$$\tilde{E}_M(u) := \int_{\mathbb{R}^N} |\nabla u|^2 dx - \frac{2M^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|u|^{p+1}}{|x|^b} dx. \quad (1.13)$$

By deriving the energy estimates between $\tilde{I}(M_k)$ and $I(M_k)$ as $M_k \rightarrow \infty$, we shall verify that $I(M_k) - \tilde{I}(M_k) \rightarrow 0$ as $M_k \rightarrow \infty$, which further implies that $\int_{\mathbb{R}^N} V(x)|u_k|^2 dx \rightarrow 0$ as $M_k \rightarrow \infty$. Furthermore, the L^∞ -uniform convergence of (1.10), which is established by analyzing delicately the singular nonlinear term of $I(M_k)$, seems crucial in the further refined investigations on the minimizers of $I(M_k)$ as $M_k \rightarrow \infty$.

This paper is organized as follows. Section 2 is devoted to the refined energy estimates of $I(M)$ as $M \rightarrow \infty$, based on which we shall complete in Section 3 the proof of Theorem 1.1 on the limit behavior of minimizers for $I(M)$ as $M \rightarrow \infty$.

2 Energy estimates of $I(M)$

This section is devoted to establishing the energy estimates of $I(M)$ as $M \rightarrow \infty$ by analyzing the energy of $\tilde{I}(M)$ defined in (1.12). Employing the concentration-compactness principle, one can deduce that $\tilde{I}(M)$ admits minimizers for any $M \in (0, \infty)$, see, e.g., [6, 26, 27]. Moreover, without loss of generality, we may consider positive minimizers of $\tilde{I}(M)$ defined in (1.12). We start with the following energy estimates of $\tilde{I}(M)$.

Lemma 2.1. *Under the assumption (1.3), assume that $V(x)$ satisfies (V), and let \tilde{u}_M be a positive minimizer of $\tilde{I}(M)$. Then for any $M > 0$,*

$$\tilde{I}(M) = -\lambda_0 \left(\frac{M}{a^*} \right)^{\frac{2(p-1)}{4-N(p-1)-2b}}, \quad (2.1)$$

and

$$\tilde{u}_M(x) = \frac{1}{\sqrt{a^*}} \tilde{\alpha}_M^{\frac{N}{2}} w(\tilde{\alpha}_M x), \quad (2.2)$$

where $\tilde{\alpha}_M := \left(\frac{M}{a^*} \right)^{\frac{p-1}{4-N(p-1)-2b}} > 0$, $\lambda_0 := -\frac{N(p-1)+2b-4}{2(p+1)-N(p-1)-2b} > 0$ and $a^* := \|w\|_2^2$. Here $w > 0$ is the unique positive solution of the equation (1.7).

Proof. Assume that \tilde{u}_M is a positive minimizer of $\tilde{I}(M)$ and \tilde{u}_1 is a positive minimizer of $\tilde{I}(1)$. We claim that for any $M > 0$,

$$\tilde{I}(M) = M^{\frac{2(p-1)}{4-N(p-1)-2b}} \tilde{I}(1) \text{ and } \tilde{u}_M(x) = \alpha_M^{\frac{N}{2}} \tilde{u}_1(\alpha_M x), \quad (2.3)$$

where $\alpha_M := M^{\frac{p-1}{4-N(p-1)-2b}} > 0$. Indeed, setting $\tilde{w}_1(x) := \alpha_M^{-\frac{N}{2}} \tilde{u}_M(\alpha_M^{-1}x)$, one can deduce from (1.12) that

$$\begin{aligned} \tilde{I}(M) &= \tilde{E}_M(\tilde{u}_M) = \int_{\mathbb{R}^N} |\nabla \tilde{u}_M|^2 dx - \frac{2M^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|\tilde{u}_M|^{p+1}}{|x|^b} dx \\ &= \alpha_M^2 \int_{\mathbb{R}^N} |\nabla \tilde{w}_1|^2 dx - \frac{2M^{\frac{p-1}{2}}}{p+1} \cdot \alpha_M^{\frac{N(p+1)}{2}} \cdot \alpha_M^{-N} \cdot \alpha_M^b \int_{\mathbb{R}^N} \frac{|\tilde{w}_1|^{p+1}}{|x|^b} dx \\ &= M^{\frac{2(p-1)}{4-N(p-1)-2b}} \left[\int_{\mathbb{R}^N} |\nabla \tilde{w}_1|^2 dx - \frac{2}{p+1} \int_{\mathbb{R}^N} \frac{|\tilde{w}_1|^{p+1}}{|x|^b} dx \right] \\ &\geq M^{\frac{2(p-1)}{4-N(p-1)-2b}} \tilde{I}(1). \end{aligned} \quad (2.4)$$

Similarly, setting $\tilde{w}_M(x) := \alpha_M^{\frac{N}{2}} \tilde{u}_1(\alpha_M x)$ as a test function of $\tilde{I}(M)$, one can get that

$$\tilde{I}(M) \leq \tilde{E}_M(\tilde{w}_M) = M^{\frac{2(p-1)}{4-N(p-1)-2b}} \tilde{I}(1). \quad (2.5)$$

Following (2.4) and (2.5), we conclude that the first equality of (2.3) holds. Furthermore, one can check that \tilde{w}_1 is a minimizer of $\tilde{I}(1)$ and \tilde{w}_M is a minimizer of $\tilde{I}(M)$. This proves the second equality of (2.3). Therefore, the claim (2.3) holds true.

We next prove that for any $M > 0$,

$$\tilde{I}(1) = -\lambda_0(a^*)^{-\frac{2(p-1)}{4-N(p-1)-2b}}, \quad \text{where } \lambda_0 := -\frac{N(p-1) + 2b - 4}{2(p+1) - N(p-1) - 2b} > 0, \quad (2.6)$$

and

$$\tilde{u}_1(x) = (a^*)^{-\frac{2-b}{4-N(p-1)-2b}} w\left((a^*)^{-\frac{p-1}{4-N(p-1)-2b}} x\right). \quad (2.7)$$

Consider a test function $0 < \tilde{v}_0 \in H^1(\mathbb{R}^N)$ satisfying $\|\tilde{v}_0\|_2^2 = 1$. Set $\tilde{v}_\epsilon(x) := \epsilon^{\frac{N}{2}} \tilde{v}_0(\epsilon x)$, where $\epsilon > 0$ is small enough. One can get that for sufficiently small $\epsilon > 0$,

$$\begin{aligned} \tilde{I}(1) &\leq \tilde{E}_1(\tilde{v}_\epsilon) = \int_{\mathbb{R}^N} |\nabla \tilde{v}_\epsilon|^2 dx - \frac{2}{p+1} \int_{\mathbb{R}^N} \frac{|\tilde{v}_\epsilon|^{p+1}}{|x|^b} dx \\ &= \epsilon^2 \int_{\mathbb{R}^N} |\nabla \tilde{v}_0|^2 dx - \frac{2\epsilon^{\frac{N(p-1)}{2}+b}}{p+1} \int_{\mathbb{R}^N} \frac{|\tilde{v}_0|^{p+1}}{|x|^b} dx < 0, \end{aligned} \quad (2.8)$$

due to the assumption (1.3). Let $\tilde{u}_1 > 0$ be a positive minimizer of $\tilde{I}(1)$. Then \tilde{u}_1 satisfies the following Euler-Lagrange equation

$$-\Delta \tilde{u}_1(x) = \tilde{\mu}_1 \tilde{u}_1(x) + \frac{\tilde{u}_1^p(x)}{|x|^b} \quad \text{in } \mathbb{R}^N, \quad (2.9)$$

where $\tilde{\mu}_1 \in \mathbb{R}$ is the Lagrangian multiplier associated to \tilde{u}_1 . Applying (2.8) and (2.9), we get that

$$\begin{aligned} \tilde{\mu}_1 &= \int_{\mathbb{R}^N} |\nabla \tilde{u}_1|^2 dx - \int_{\mathbb{R}^N} \frac{|\tilde{u}_1|^{p+1}}{|x|^b} dx \\ &= \tilde{I}(1) - \frac{p-1}{p+1} \int_{\mathbb{R}^N} \frac{|\tilde{u}_1|^{p+1}}{|x|^b} dx < 0. \end{aligned} \quad (2.10)$$

Since $w > 0$ is the unique positive solution of (1.7), one can conclude from (1.7) and (2.9) that

$$\tilde{u}_1(x) = (-\tilde{\mu}_1)^{\frac{2-b}{2(p-1)}} w\left((-\tilde{\mu}_1)^{\frac{1}{2}} x\right),$$

where $\tilde{\mu}_1 < 0$ holds by (2.10). Moreover, since

$$1 = \|\tilde{u}_1\|_2^2 = (-\tilde{\mu}_1)^{\frac{4-2b-N(p-1)}{2(p-1)}} \|w\|_2^2 = (-\tilde{\mu}_1)^{\frac{4-2b-N(p-1)}{2(p-1)}} a^*,$$

one can derive that

$$\tilde{\mu}_1 = -(a^*)^{\frac{2(1-p)}{4-2b-N(p-1)}} < 0 \quad \text{and} \quad \tilde{u}_1(x) = (a^*)^{\frac{b-2}{4-2b-N(p-1)}} w\left((a^*)^{\frac{1-p}{4-2b-N(p-1)}} x\right) > 0.$$

Hence, (2.7) is proved. On the other hand, substituting (2.7) and (1.8) into (1.13), we

get that

$$\begin{aligned}
\tilde{I}(1) &= \tilde{E}_1(\tilde{u}_1) = \int_{\mathbb{R}^N} |\nabla \tilde{u}_1|^2 dx - \frac{2}{p+1} \int_{\mathbb{R}^N} \frac{|\tilde{u}_1|^{p+1}}{|x|^b} dx \\
&= (a^*)^{\frac{2(b-2)}{4-2b-N(p-1)}} \cdot (a^*)^{\frac{2(1-p)}{4-2b-N(p-1)}} \cdot (a^*)^{\frac{N(p-1)}{4-2b-N(p-1)}} \int_{\mathbb{R}^N} |\nabla w|^2 dx \\
&\quad - (a^*)^{\frac{(b-2)(p+1)}{4-2b-N(p-1)}} \cdot (a^*)^{\frac{N(p-1)}{4-2b-N(p-1)}} \cdot (a^*)^{\frac{(1-p)b}{4-2b-N(p-1)}} \cdot \frac{2}{p+1} \int_{\mathbb{R}^N} \frac{|w|^{p+1}}{|x|^b} dx \\
&= (a^*)^{\frac{2(b-2)+(p-1)(N-2)}{4-2b-N(p-1)}} \cdot \frac{N(p-1)+2b}{2(p+1)-N(p-1)-2b} \cdot a^* \\
&\quad - (a^*)^{\frac{2(b-2)+(p-1)(N-2)}{4-2b-N(p-1)}} \cdot \frac{2}{p+1} \cdot \frac{2(p+1)}{2(p+1)-N(p-1)-2b} \cdot a^* \\
&= (a^*)^{-\frac{2(p-1)}{4-2b-N(p-1)}} \cdot \frac{N(p-1)+2b-4}{2(p+1)-N(p-1)-2b} \\
&= -\lambda_0 (a^*)^{-\frac{2(p-1)}{4-2b-N(p-1)}},
\end{aligned}$$

which thus implies that (2.6) holds.

We finally conclude from (2.3)–(2.7) that for any $M > 0$,

$$\tilde{I}(M) = M^{\frac{2(p-1)}{4-N(p-1)-2b}} \cdot (a^*)^{-\frac{2(p-1)}{4-N(p-1)-2b}} \cdot (-\lambda_0) = -\lambda_0 \left(\frac{M}{a^*} \right)^{\frac{2(p-1)}{4-N(p-1)-2b}},$$

and

$$\begin{aligned}
\tilde{u}_M(x) &= \alpha_M^{\frac{N}{2}} \tilde{u}_1(\alpha_M x) = \alpha_M^{\frac{N}{2}} (a^*)^{-\frac{2-b}{4-N(p-1)-2b}} w \left((a^*)^{-\frac{p-1}{4-N(p-1)-2b}} \cdot \alpha_M x \right) \\
&= M^{\frac{N(p-1)}{2[4-N(p-1)-2b]}} \cdot (a^*)^{-\frac{2-b}{4-N(p-1)-2b}} \cdot w \left(\left(\frac{M}{a^*} \right)^{\frac{p-1}{4-N(p-1)-2b}} x \right) \\
&= \frac{1}{\sqrt{a^*}} \tilde{\alpha}_M^{\frac{N}{2}} w(\tilde{\alpha}_M x).
\end{aligned}$$

Therefore, the proof of Lemma 2.1 is completed. \square

Applying Lemma 2.1, we now establish the energy estimates of $I(M)$.

Lemma 2.2. *Under the assumption (1.3), assume that $V(x)$ satisfies (V). Then we have*

$$\lim_{M \rightarrow \infty} \frac{I(M)}{\left(\frac{M}{a^*} \right)^{\frac{2(p-1)}{4-N(p-1)-2b}}} = -\lambda_0, \quad (2.11)$$

where $a^* := \|w\|_2^2$, $\lambda_0 := -\frac{N(p-1)+2b-4}{2(p+1)-N(p-1)-2b} > 0$, and $w > 0$ is the unique positive solution of the equation (1.7).

Proof. We first establish the lower bound of $I(M)$ as $M \rightarrow \infty$. Let $u_M > 0$ be a

positive minimizer of $I(M)$. Under the assumption (V), we get from (1.12) that

$$\begin{aligned}
I(M) &= \int_{\mathbb{R}^N} |\nabla u_M|^2 dx + \int_{\mathbb{R}^N} V(x) u_M^2(x) dx - \frac{2M^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|u_M|^{p+1}}{|x|^b} dx \\
&\geq \int_{\mathbb{R}^N} |\nabla u_M|^2 dx - \frac{2M^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|u_M|^{p+1}}{|x|^b} dx \\
&\geq \tilde{I}(M) = -\lambda_0 \left(\frac{M}{a^*} \right)^{\frac{2(p-1)}{4-(p-1)-2b}} \quad \text{as } M \rightarrow \infty,
\end{aligned} \tag{2.12}$$

where $\lambda_0 > 0$ is as in Lemma 2.1. This thus implies the lower bound of $I(M)$ as $M \rightarrow \infty$.

We next estimate the upper bound of $I(M)$ as $M \rightarrow \infty$. Define

$$u_\tau(x) := \frac{A_\tau \tau^{\frac{N}{2}}}{\|w\|_2} w(\tau x) \varphi(x), \quad \tau > 0, \tag{2.13}$$

where $0 \leq \varphi(x) \in C^\infty(\mathbb{R}^N)$ is a cut-off function satisfying

$$\varphi(x) = \begin{cases} 1, & |x| \leq 1; \\ 0, & |x| \geq 2, \end{cases}$$

$w > 0$ is the unique positive solution of (1.7), and $A_\tau > 0$ is a suitable constant such that $\|u_\tau(x)\|_2^2 = 1$. Applying the exponential decay of w in (1.9), one can check that as $\tau \rightarrow \infty$,

$$1 \leq A_\tau^2 = \frac{\|w\|_2^2}{\int_{\mathbb{R}^N} w^2(x) \varphi^2\left(\frac{x}{\tau}\right) dx} \leq 1 + \frac{\int_{B_\tau^c} w^2(x) dx}{\int_{B_\tau} w^2(x) dx} \leq 1 + C e^{-2\delta\tau}, \tag{2.14}$$

where $C > 0$ is independent of $\tau > 0$. Substituting (2.13) into (1.2) and applying the exponential decay of w in (1.9) and the identity (1.8), direct calculations yield that as $\tau \rightarrow \infty$,

$$\begin{aligned}
\int_{\mathbb{R}^N} |\nabla u_\tau|^2 dx &= \frac{A_\tau^2}{\|w\|_2^2} \int_{\mathbb{R}^N} \left| \varphi\left(\frac{x}{\tau}\right) \tau \nabla w(x) + w(x) \nabla \varphi\left(\frac{x}{\tau}\right) \right|^2 dx \\
&= \frac{A_\tau^2}{\|w\|_2^2} \int_{\mathbb{R}^N} \left[\tau^2 \varphi^2\left(\frac{x}{\tau}\right) |\nabla w(x)|^2 + w^2(x) |\nabla \varphi\left(\frac{x}{\tau}\right)|^2 \right. \\
&\quad \left. + 2\tau \nabla w(x) \varphi\left(\frac{x}{\tau}\right) \nabla \varphi\left(\frac{x}{\tau}\right) w(x) \right] dx \\
&\leq (1 + C e^{-2\delta\tau}) \frac{N(p-1) + 2b}{2(p+1) - N(p-1) - 2b} \tau^2,
\end{aligned} \tag{2.15}$$

$$\begin{aligned}
\int_{\mathbb{R}^N} V(x)u_\tau^2(x)dx &= \frac{A_\tau^2}{\|w\|_2^2} \int_{\mathbb{R}^N} V\left(\frac{x}{\tau}\right)w^2(x)\varphi^2\left(\frac{x}{\tau}\right)dx \\
&= \frac{A_\tau^2}{\|w\|_2^2} \int_{B_{2\tau}(0)} V\left(\frac{x}{\tau}\right)w^2(x)\varphi^2\left(\frac{x}{\tau}\right)dx \\
&\leq \frac{A_\tau^2}{\|w\|_2^2} \left[\int_{0 \leq |x| \leq \sqrt{\tau}} V\left(\frac{x}{\tau}\right)w^2(x)dx + Ce^{-\delta\sqrt{\tau}} \right],
\end{aligned} \tag{2.16}$$

and

$$\begin{aligned}
\int_{\mathbb{R}^N} \frac{|u_\tau|^{p+1}}{|x|^b} dx &= \frac{A_\tau^{p+1} \tau^{\frac{N}{2}(p+1)}}{\|w\|_2^{p+1}} \int_{\mathbb{R}^N} \frac{|w(\tau x)\varphi(x)|^{p+1}}{|x|^b} dx \\
&= \frac{A_\tau^{p+1} \tau^{\frac{N}{2}(p-1)+b}}{\|w\|_2^{p+1}} \int_{\mathbb{R}^N} \frac{|w(x)\varphi\left(\frac{x}{\tau}\right)|^{p+1}}{|x|^b} dx \\
&\geq \tau^{\frac{N}{2}(p-1)+b} \frac{2(p+1)}{2(p+1) - N(p-1) - 2b} (a^*)^{\frac{1-p}{2}} - Ce^{-\delta(p+1)\tau} \tau^{\frac{N(p-1)}{2}}.
\end{aligned} \tag{2.17}$$

It then follows from (2.15)–(2.17) that as $\tau \rightarrow \infty$,

$$I(M) \leq E_M(u_\tau) \leq \frac{N(p-1) + 2b}{2(p+1) - N(p-1) - 2b} \tau^2 - \tau^{\frac{N}{2}(p-1)+b} \frac{4\left(\frac{M}{a^*}\right)^{\frac{p-1}{2}}}{2(p+1) - N(p-1) - 2b} + V(0) + Ce^{-\delta\sqrt{\tau}}.$$

Setting $\tau = \left(\frac{M}{a^*}\right)^{\frac{p-1}{4-N(p-1)-2b}}$ into the above estimate, it then gives that as $M \rightarrow \infty$,

$$I(M) \leq -\lambda_0 \left(\frac{M}{a^*}\right)^{\frac{2(p-1)}{4-N(p-1)-2b}} + Ce^{-\delta\sqrt{\tau}} = -\lambda_0 \left(\frac{M}{a^*}\right)^{\frac{2(p-1)}{4-N(p-1)-2b}} + o(1), \tag{2.18}$$

where $\lambda_0 := -\frac{N(p-1)+2b-4}{2(p+1)-N(p-1)-2b} > 0$. Thus, (2.11) follows from (2.12) and (2.18), which completes the proof of Lemma 2.2. \square

3 Proof of Theorem 1.1

In this section, we shall complete the proof of Theorem 1.1 on the limit behavior of minimizers for (1.1) by the blow-up analysis. We first establish the following lemma.

Lemma 3.1. *Under the assumption (1.3), assume that $V(x)$ satisfies (V), and let u_M be a positive minimizer of $I(M)$. Then we have*

$$\int_{\mathbb{R}^N} V(x)u_M^2(x)dx \rightarrow 0 \quad \text{as } M \rightarrow \infty. \tag{3.1}$$

Proof. The key of proving (3.1) is to verify that

$$I(M) - \tilde{I}(M) \rightarrow 0 \quad \text{as } M \rightarrow \infty. \tag{3.2}$$

Indeed, if (3.2) holds, then one derive from (1.12) and (1.13) that

$$\int_{\mathbb{R}^N} V(x)u_M^2(x)dx = I(M) - \tilde{E}_M(u_M) \leq I(M) - \tilde{I}(M) \rightarrow 0 \text{ as } M \rightarrow \infty, \quad (3.3)$$

which thus implies that (3.1) holds.

We now prove (3.2). By Lemma 2.1, we deduce from (2.18) that

$$I(M) \leq \tilde{I}(M) + o(1) \text{ as } M \rightarrow \infty. \quad (3.4)$$

On the other hand, it follows from (1.12) and (1.13) that

$$I(M) - \tilde{I}(M) \geq E_M(u_M) - \tilde{E}_M(u_M) = \int_{\mathbb{R}^N} V(x)u_M^2(x)dx \geq 0 \text{ as } M \rightarrow \infty. \quad (3.5)$$

Therefore, the estimate (3.2) now follows from (3.4) and (3.5), and we are done. \square

Motivated by [20, 22, 32], we next establish the following lemma.

Lemma 3.2. *Under the assumption (1.3), assume that $V(x)$ satisfies (V), and let u_k be a positive minimizer of $I(M_k)$, where $M_k \rightarrow \infty$ as $k \rightarrow \infty$. Define*

$$w_k(x) := \epsilon_k^{\frac{N}{2}} u_k(\epsilon_k x), \quad (3.6)$$

where $\epsilon_k := \left(\frac{M_k}{a^*}\right)^{-\frac{p-1}{4-N(p-1)-2b}} \rightarrow 0$ as $k \rightarrow \infty$. Then there exists a subsequence, still denoted by $\{w_k\}$, of $\{u_k\}$ such that

$$w_k(x) \rightarrow \frac{w(x)}{\sqrt{a^*}} \text{ strongly in } H^1(\mathbb{R}^N) \text{ as } k \rightarrow \infty, \quad (3.7)$$

where $a^* := \|w\|_2^2$ and $w > 0$ is the unique positive solution of (1.7).

Proof. We first prove that there exist some positive constants C_1, C_2, C'_1, C'_2 , which are independent of k , such that as $k \rightarrow \infty$,

$$0 < C_1 \leq \|\nabla w_k\|_2^2 \leq C_2 \text{ and } 0 < C'_1 \leq \int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx \leq C'_2. \quad (3.8)$$

Indeed, using Lemmas 2.2 and 3.1, we deduce from (1.2) and (3.6) that as $k \rightarrow \infty$,

$$\begin{aligned} \epsilon_k^2 I(M_k) &= \epsilon_k^2 \left(\int_{\mathbb{R}^N} |\nabla u_k|^2 dx + \int_{\mathbb{R}^N} V(x)u_k^2 dx - \frac{2M_k^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|u_k|^{p+1}}{|x|^b} dx \right) \\ &= \left(\int_{\mathbb{R}^N} |\nabla w_k|^2 dx - \frac{2(a^*)^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx + o(\epsilon_k^2) \right) \rightarrow -\lambda_0 < 0, \end{aligned} \quad (3.9)$$

which implies that

$$\frac{2(a^*)^{\frac{p-1}{2}} \int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx}{p+1 \int_{\mathbb{R}^N} |\nabla w_k|^2 dx} \rightarrow 1 \text{ as } k \rightarrow \infty. \quad (3.10)$$

By contradiction, assume that $\|\nabla w_k\|_2^2 \rightarrow \infty$ as $k \rightarrow \infty$. Define $\gamma_k^2 := \|\nabla w_k\|_2^2 > 0$ and $v_k(x) = \gamma_k^{-\frac{N}{2}} w_k(\gamma_k^{-1}x)$, so that $\gamma_k^2 \rightarrow \infty$ as $k \rightarrow \infty$. It then follows that $\|v_k\|_2^2 = 1$ and $\|\nabla v_k\|_2^2 = 1$ for all $k \geq 1$. Further, we deduce from the GN inequality (1.5) that

$$\int_{\mathbb{R}^N} \frac{|v_k|^{p+1}}{|x|^b} dx \leq C_{GN}^{-1} \|\nabla v_k\|_2^{\frac{N(p-1)}{2}+b} \|v_k\|_2^{p+1-\frac{N(p-1)}{2}-b} = C_{GN}^{-1}, \quad (3.11)$$

where $C_{GN} > 0$ is given in (1.6). Under the assumption (1.3), since $\|v_k\|_2^2 = \|\nabla v_k\|_2^2 = 1$ for all $k \geq 1$, it follows from (3.11) that

$$\begin{aligned} \frac{2(a^*)^{\frac{p-1}{2}} \int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx}{p+1 \int_{\mathbb{R}^N} |\nabla w_k|^2 dx} &= \frac{2(a^*)^{\frac{p-1}{2}} \int_{\mathbb{R}^N} \frac{|v_k|^{p+1}}{|x|^b} dx}{p+1 \|\nabla v_k\|_2^2} \gamma_k^{-\frac{4-N(p-1)-2b}{2}} \\ &\leq \frac{2(a^*)^{\frac{p-1}{2}}}{p+1} C_{GN}^{-1} \gamma_k^{-\frac{4-N(p-1)-2b}{2}} \rightarrow 0 \quad \text{as } k \rightarrow \infty, \end{aligned} \quad (3.12)$$

which however contradicts to (3.10). Hence, we conclude that $\|\nabla w_k\|_2^2 \leq C_2$ holds uniformly as $k \rightarrow \infty$. Applying the GN inequality (1.5) and the fact that $\|w_k\|_2^2 = 1$, we deduce from above that $\int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx \leq C'_2$ holds uniformly as $k \rightarrow \infty$. On the other hand, one can obtain from (3.9) that $\int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx \geq C'_1$ holds uniformly as $k \rightarrow \infty$, together with (1.5), which then imply that $\|\nabla w_k\|_2^2 \geq C_1$ holds uniformly as $k \rightarrow \infty$. We therefore conclude that (3.8) holds.

From (3.6) and (3.8), we deduce that w_k is bounded uniformly in $H^1(\mathbb{R}^N)$, which implies that there exist a subsequence, still denoted by $\{w_k\}$, of $\{w_k\}$ and $0 \leq w_0 \in H^1(\mathbb{R}^N)$ such that

$$w_k \rightharpoonup w_0 \geq 0 \quad \text{weakly in } H^1(\mathbb{R}^N) \quad \text{as } k \rightarrow \infty. \quad (3.13)$$

We now prove that $w_0 \not\equiv 0$. Motivated by [2, 15], we first claim that

$$\int_{\mathbb{R}^N} |x|^{-b} w_k^{p+1} dx \rightarrow \int_{\mathbb{R}^N} |x|^{-b} w_0^{p+1} dx \quad \text{as } k \rightarrow \infty. \quad (3.14)$$

Actually, we have

$$\begin{aligned} &\left| \int_{\mathbb{R}^N} |x|^{-b} w_k^{p+1} dx - \int_{\mathbb{R}^N} |x|^{-b} w_0^{p+1} dx \right| \\ &\leq \int_{\mathbb{R}^N} |x|^{-b} |w_k^{p+1} - w_0^{p+1}| dx \\ &= \int_{B_R} |x|^{-b} |w_k^{p+1} - w_0^{p+1}| dx + \int_{\mathbb{R}^N \setminus B_R} |x|^{-b} |w_k^{p+1} - w_0^{p+1}| dx := A_k + B_k, \end{aligned}$$

where $R > 0$ is arbitrary. Under the assumption (1.3), we have $1 < p < 1 + \frac{4-2b}{N}$ and

$$1 + \frac{4-2b}{N} < 1 + \frac{4-2b}{N-2}, \quad \text{if } N \geq 3.$$

By Hölder inequality, we then have

$$\begin{aligned}
A_k &= \int_{B_R} |x|^{-b} |w_k^{p+1} - w_0^{p+1}| dx \\
&\leq \left(\int_{B_R} |x|^{-br} dx \right)^{\frac{1}{r}} \left(\int_{B_R} |w_k^{p+1} - w_0^{p+1}|^t dx \right)^{\frac{1}{t}} \\
&\leq C \left(\int_{B_R} |w_k^{p+1} - w_0^{p+1}|^t dx \right)^{\frac{1}{t}},
\end{aligned} \tag{3.15}$$

where $\frac{1}{r} + \frac{1}{t} = 1$, $t > 1$ and $r > 1$ satisfies $\frac{b}{N} < \frac{1}{r}$. Note that $\frac{1}{t} = 1 - \frac{1}{r} < \frac{N-b}{N}$. Consider $p_1 > 0$ and $q_1 > 0$ satisfying

$$\frac{p}{p_1} + \frac{1}{q_1} = \frac{1}{t} < \frac{N-b}{N}, \tag{3.16}$$

which then yields from (3.15) that

$$\begin{aligned}
A_k &\leq C \left(\int_{B_R} |w_k^{p+1} - w_0^{p+1}|^t dx \right)^{\frac{1}{t}} \\
&\leq C \left(\|w_k\|_{L^{p_1}(B_R)}^p + \|w_0\|_{L^{p_1}(B_R)}^p \right) \|w_k - w_0\|_{L^{q_1}(B_R)}.
\end{aligned} \tag{3.17}$$

Similar to [2, Theorem 1.5], choose suitable constants $p_1 > 0$ and $q_1 > 0$ satisfying (3.16), so that

$$\|w_k\|_{L^{p_1}(B_R)}^p + \|w_0\|_{L^{p_1}(B_R)}^p \leq C \quad \text{and} \quad \|w_k - w_0\|_{L^{q_1}(B_R)} \rightarrow 0 \quad \text{as } k \rightarrow \infty,$$

where $C > 0$ is independent of $k > 0$. This further implies from (3.17) that

$$A_k \rightarrow 0 \quad \text{as } k \rightarrow \infty, \tag{3.18}$$

On the other hand, for any $\epsilon > 0$, there exists $R \geq \epsilon^{-\frac{1}{b}}$ such that

$$\begin{aligned}
B_k &= \int_{\mathbb{R}^N \setminus B_R} |x|^{-b} |w_k^{p+1} - w_0^{p+1}| dx \\
&\leq \epsilon \int_{\mathbb{R}^N \setminus B_R} (w_k^{p+1} + w_0^{p+1}) dx \leq C\epsilon \quad \text{as } k \rightarrow \infty,
\end{aligned}$$

due to Sobolev's embedding theorem and the uniform boundedness of w_k in $H^1(\mathbb{R}^N)$. Since $\epsilon > 0$ is arbitrary, we conclude from above that the claim (3.14) holds true. Following (3.8) and (3.14), one can deduce that $w_0 \not\equiv 0$.

Next, we prove that $\|w_0\|_2^2 = 1$. By contradiction, we assume that $\|w_0\|_2^2 = l$, where $l \in (0, 1)$. Set $w_l := \frac{w_0}{\sqrt{l}}$. By (3.13), we may assume that $w_k \rightarrow w_0$ *a.e.* in \mathbb{R}^N as $k \rightarrow \infty$. Using the Brézis-Lieb lemma, we obtain that

$$\|\nabla w_k\|_2^2 = \|\nabla w_0\|_2^2 + \|\nabla(w_k - w_0)\|_2^2 + o(1) \quad \text{as } k \rightarrow \infty. \tag{3.19}$$

From (2.1), (2.11), (3.6), (3.14) and (3.19), we derive that as $k \rightarrow \infty$,

$$\begin{aligned}
-\lambda_0 &= \lim_{k \rightarrow \infty} \epsilon_k^2 I(M_k) \\
&= \lim_{k \rightarrow \infty} \epsilon_k^2 \left(\int_{\mathbb{R}^N} |\nabla u_k|^2 dx + \int_{\mathbb{R}^N} V(x) u_k^2 dx - \frac{2M_k^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|u_k|^{p+1}}{|x|^b} dx \right) \\
&\geq \lim_{k \rightarrow \infty} \left(\int_{\mathbb{R}^N} |\nabla w_k|^2 dx - \frac{2(a^*)^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx \right) \\
&= \int_{\mathbb{R}^N} |\nabla w_0|^2 dx + \lim_{k \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla(w_k - w_0)|^2 dx - \frac{2(a^*)^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|w_0|^{p+1}}{|x|^b} dx \quad (3.20) \\
&\geq \int_{\mathbb{R}^N} |\nabla w_0|^2 dx - \frac{2(a^*)^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|w_0|^{p+1}}{|x|^b} dx \\
&= l \left[\int_{\mathbb{R}^N} |\nabla w_l|^2 dx - \frac{2(a^*l)^{\frac{p-1}{2}}}{p+1} \int_{\mathbb{R}^N} \frac{|w_l|^{p+1}}{|x|^b} dx \right] \\
&> l\tilde{I}(a^*) = -l\lambda_0 < 0,
\end{aligned}$$

which is a contradiction. Hence, $\|w_0\|_2^2 = 1$ holds true.

Since $\|w_k\|_2^2 = \|w_0\|_2^2 = 1$, we have

$$w_k(x) \rightarrow w_0(x) \quad \text{strongly in } L^2(\mathbb{R}^N) \text{ as } k \rightarrow \infty. \quad (3.21)$$

By the weak lower semicontinuity, (2.1) and (3.14), we then derive from (3.9) that

$$\nabla w_k(x) \rightarrow \nabla w_0(x) \quad \text{strongly in } L^2(\mathbb{R}^N) \text{ as } k \rightarrow \infty. \quad (3.22)$$

Note from (3.9) that $\{w_k\}$ is a minimizing sequence of $\tilde{I}(a^*)$. One then deduces from (3.14) and (3.22) that w_0 is a minimizer of $\tilde{I}(a^*)$. By (2.2), we obtain that $w_0(x) = \frac{w(x)}{\sqrt{a^*}}$. Combining (3.21) and (3.22), we obtain that

$$w_k(x) \rightarrow w_0(x) = \frac{w(x)}{\sqrt{a^*}} \quad \text{strongly in } H^1(\mathbb{R}^N) \text{ as } k \rightarrow \infty, \quad (3.23)$$

which gives (3.7). The lemma is thus proved. \square

Applying above lemmas, we are now ready to prove Theorem 1.1.

Proof of Theorem 1.1: 1. We first prove the exponential decay (1.11). Let $u_k > 0$ be a minimizer of $I(M_k)$, and consider the sequence $\{w_k\}$ defined in Lemma 3.2, where $M_k \rightarrow \infty$ as $k \rightarrow \infty$. We claim that there exists a subsequence, still denoted by $\{w_k\}$, of $\{w_k\}$ such that

$$w_k(x) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty \text{ uniformly for sufficiently large } k > 0. \quad (3.24)$$

Indeed, one can derive from (3.7) that for any $2 \leq \alpha < 2^*$,

$$\int_{|x| \geq \gamma} |w_k|^\alpha dx \rightarrow 0 \quad \text{as } \gamma \rightarrow \infty \text{ uniformly for sufficiently large } k > 0. \quad (3.25)$$

On the other hand, it follows from (1.4) and (3.6) that w_k satisfies the following equation

$$-\Delta w_k + \epsilon_k^2 V(\epsilon_k x) w_k - (a^*)^{\frac{p-1}{2}} \frac{w_k^p}{|x|^b} = \mu_k \epsilon_k^2 w_k \quad \text{in } \mathbb{R}^N, \quad (3.26)$$

where $\mu_k \in \mathbb{R}$ is the Lagrange multiplier. Applying (1.8), (3.6) and (3.14), we deduce from Lemma 2.2 that

$$\begin{aligned} \epsilon_k^2 \mu_k &= \epsilon_k^2 \left(I(M_k) - \frac{p-1}{p+1} M_k^{\frac{p-1}{2}} \int_{\mathbb{R}^N} \frac{|u_k|^{p+1}}{|x|^b} dx \right) \\ &= \epsilon_k^2 I(M_k) - \frac{p-1}{p+1} (a^*)^{\frac{p-1}{2}} \int_{\mathbb{R}^N} \frac{|w_k|^{p+1}}{|x|^b} dx \\ &\rightarrow -1 \quad \text{as } k \rightarrow \infty. \end{aligned} \quad (3.27)$$

Using (3.27), we derive from (3.26) that as $k \rightarrow \infty$,

$$-\Delta w_k - c(x) w_k \leq 0 \quad \text{in } \mathbb{R}^N, \quad \text{where } c(x) = (a^*)^{\frac{p-1}{2}} \frac{w_k^{p-1}(x)}{|x|^b}. \quad (3.28)$$

Furthermore, one can check from Hölder inequality that

$$c(x) \in L^t(\mathbb{R}^N), \quad \text{where } t \in \left(\frac{2N}{N(p-1) + 2b}, \frac{2N}{(N-2)(p-1) + 2b} \right).$$

Applying De Giorgi-Nash-Moser theory ([23, Theorem 4.1]) to (3.28), we deduce that

$$\max_{B_1(\xi)} w_k(x) \leq C \left(\int_{B_2(\xi)} |w_k(x)|^\alpha dx \right)^{\frac{1}{\alpha}} \quad \text{for sufficiently large } k > 0, \quad (3.29)$$

where $\xi \in \mathbb{R}^N$ is arbitrary, and $C > 0$ depends only on the bound of $\|c(x)\|_{L^t(B_2(\xi))}$. Thus, (3.24) follows from (3.25) and (3.29).

Due to the smallness of $|x|^{-b}$ for large $|x| > 0$, we now derive from (3.24), (3.26) and (3.27) that there exists a sufficiently large constant $R > 0$, which is independent of k , such that as $k \rightarrow \infty$,

$$-\Delta w_k(x) + \theta w_k(x) \leq 0 \quad \text{in } \mathbb{R}^N \setminus B_R(0), \quad (3.30)$$

where $0 < \theta < 1$ is independent of k . By the comparison principle [5, Theorem 6.4.2], we obtain from (3.30) that as $k \rightarrow \infty$,

$$w_k(x) \leq C e^{-\sqrt{\theta}|x|} \quad \text{in } \mathbb{R}^N \setminus B_R(0), \quad (3.31)$$

which implies the exponential decay (1.11) for w_k as $k \rightarrow \infty$. Moreover, under the assumption (V), since the term $|x|^{-b}$ is small for large $|x|$, applying the local elliptic estimate (cf. (3.15) in [17]) yields from (3.31) that as $k \rightarrow \infty$,

$$|\nabla w_k(x)| \leq C e^{-\theta|x|} \quad \text{for } |x| > R,$$

which thus gives the exponential decay (1.11) for ∇w_k as $k \rightarrow \infty$. This proves (1.11).

2. We next prove that (1.10) holds true. Recall from Lemma 3.2 that

$$w_k(x) \rightarrow \frac{w(x)}{\sqrt{a^*}} \text{ in } H^1(\mathbb{R}^N) \text{ as } k \rightarrow \infty, \quad (3.32)$$

where the convergence holds for the whole sequence $\{w_k(x)\}$, due to the uniqueness of $w(x) > 0$. Following (3.32), the L^∞ -uniform convergence (1.10) for the case $N = 1$ can be directly obtained by applying Sobolev's embedding theorem $H^1(\mathbb{R}) \hookrightarrow L^\infty(\mathbb{R})$.

We now prove the L^∞ -uniform convergence (1.10) for the case $N \geq 2$. Rewrite (3.26) as

$$-\Delta w_k(x) = G_k(x) \text{ in } H^1(\mathbb{R}^N), \quad (3.33)$$

where

$$G_k(x) := \mu_k \epsilon_k^2 w_k - \epsilon_k^2 V(\epsilon_k x) w_k + (a^*)^{\frac{p-1}{2}} \frac{w_k^p}{|x|^b}.$$

Since w_k is bounded uniformly in $H^1(\mathbb{R}^N)$ as $k \rightarrow \infty$, we derive from (3.29) that

$$w_k \text{ is bounded uniformly in } L^\infty(\mathbb{R}^N). \quad (3.34)$$

Using Hölder inequality, we deduce that for any $R > 0$,

$$\int_{B_R(0)} \left| \frac{w_k^p(x)}{|x|^b} \right|^r dx \leq \left(\int_{B_R(0)} \frac{1}{|x|^{brt}} dx \right)^{\frac{1}{t}} \left(\int_{B_R(0)} |w_k^p|^{rt'} dx \right)^{\frac{1}{t'}} < \infty, \quad (3.35)$$

where $r \in (1, \frac{N}{b})$, $\frac{1}{t} + \frac{1}{t'} = 1$, and $t' = 1 + \max\left\{\frac{N}{N-br}, \frac{2}{pr}\right\}$. We then obtain from (3.35) that

$$w_k^p(x) |x|^{-b} \in L_{loc}^r(\mathbb{R}^N) \text{ for any } r \in (1, \frac{N}{b}), \quad (3.36)$$

which implies that $G_k(x)$ is bounded uniformly in $L_{loc}^r(\mathbb{R}^N)$. For any large $R > 0$, it thus follows from [17, Theorem 9.11] that

$$\|w_k(x)\|_{W^{2,r}(B_R)} \leq C \left(\|w_k(x)\|_{L^r(B_{R+1})} + \|G_k(x)\|_{L^r(B_{R+1})} \right), \quad (3.37)$$

where $C > 0$ is independent of $k > 0$ and $R > 0$. By the compactness of the embedding $W^{2,r}(B_R) \hookrightarrow L^\infty(B_R)$ for $2r > N$, cf. [17, Theorem 7.26], we conclude that there exists a subsequence, still denoted by $\{w_k\}$, of $\{w_k\}$ such that

$$w_k(x) \rightarrow \tilde{w}_0(x) \text{ uniformly in } L^\infty(B_R) \text{ as } k \rightarrow \infty. \quad (3.38)$$

Since $R > 0$ is arbitrary, we obtain from (3.32) that

$$w_k(x) \rightarrow \frac{w(x)}{\sqrt{a^*}} \text{ uniformly in } L_{loc}^\infty(\mathbb{R}^N) \text{ as } k \rightarrow \infty. \quad (3.39)$$

On the other hand, we deduce from (1.9) and (1.11) that for any $\epsilon > 0$, there exists a constant $R_\epsilon > 0$, independent of $k > 0$, such that

$$|w_k(x)|, \left| \frac{w(x)}{\sqrt{a^*}} \right| < \frac{\epsilon}{4} \quad \text{for any } |x| > R_\epsilon,$$

which implies that

$$\sup_{|x| > R_\epsilon} \left| w_k(x) - \frac{w(x)}{\sqrt{a^*}} \right| \leq \sup_{|x| > R_\epsilon} (|w_k(x)| + \left| \frac{w(x)}{\sqrt{a^*}} \right|) \leq \frac{\epsilon}{2}.$$

Recall from (3.39) that for sufficiently large $k > 0$,

$$\sup_{|x| \leq R_\epsilon} \left| w_k(x) - \frac{w(x)}{\sqrt{a^*}} \right| \leq \frac{\epsilon}{2}.$$

We now conclude from above that the L^∞ -uniform convergence (1.10) holds true for all $N \geq 2$. The proof of Theorem 1.1 is therefore complete. \square

References

- [1] Agrawal, G.P.: *Nonlinear Fiber Optics*. Academic Press, Cambridge (2007)
- [2] Ardila, A.H., Dinh, V.D.: Some qualitative studies of the focusing inhomogeneous Gross-Pitaevskii equation. *Z. Angew. Math. Phys.* **71**, 79 (2020)
- [3] Baym, G., Pethick, C.J.: Ground state properties of magnetically trapped Bose-Einstein condensate rubidium gas. *Phys. Rev. Lett.* **76**, 6–9 (1996)
- [4] Berestycki, H., Lions, P.L.: Nonlinear scalar field equations, I, Existence of a ground state. *Arch. Rational Mech. Anal.* **82**, 313–346 (1983)
- [5] Cao, D.M., Peng, S.J., Yan, S.: *Singularly Perturbed Methods for Nonlinear Elliptic Problems*. Cambridge University Press, New York (2021).
- [6] Cazenave, T.: *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics, Vol 10. American Mathematical Society, Courant Institute of Mathematical Sciences (2003)
- [7] Cazenave, T., Lions, P.L.: Orbital stability of standing waves for some nonlinear Schrödinger equations. *Comm. Math. Phys.* **85**, no. 4, 549–561 (1982)
- [8] Combet, V., Genoud, F.: Classification of minimal mass blow-up solutions for an L^2 critical inhomogeneous NLS. *J. Evol. Equ.* **16**, 483–500 (2016)

- [9] de Bouard, A., Fukuizumi, R.: Stability of standing waves for nonlinear Schrödinger equations with inhomogeneous nonlinearities. *Ann. Henri Poincaré* **6**, 1157–1177 (2005)
- [10] Deng, Y.B., Guo, Y.J., Lu, L.: On the collapse and concentration of Bose-Einstein condensates with inhomogeneous attractive interactions. *Calc. Var. Partial Differ. Equ.* **54**, 99–118 (2015)
- [11] Deng, Y.B., Guo, Y.J., Lu, L.: Threshold behavior and uniqueness of ground states for mass critical inhomogeneous Schrödinger equations. *J. Math. Phys.* **59**, 011503 (2015)
- [12] Dinh, V.D.: Blowup of H^1 solutions for a class of the focusing inhomogeneous nonlinear Schrödinger equation. *Nonlinear Anal.* **174**, 169–188 (2018)
- [13] Farah, L.G.: Global well-posedness and blow-up on the energy space for the inhomogeneous nonlinear Schrödinger equation. *J. Evol. Equ.* **16** (1), 193–208 (2016)
- [14] Fukuizumi, R.: Stability and instability of standing waves for the Schrödinger equation with harmonic potential. *Discrete Contin. Dyn. Syst.* **7**, 525–544 (2000)
- [15] Genoud, F., Stuart, C.A.: Schrödinger equations with a spatially decaying nonlinearity: existence and stability of standing waves. *Discrete Contin. Dyn. Syst.* **21**, 137–186 (2008)
- [16] Gidas, B., Ni, W.M. and Nirenberg, L.: Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n , mathematical analysis and applications part A. *Adv. Math. Suppl. Stud.* **7**, 369–402 (1981)
- [17] Gilbarg, D., Trudinger, N.S.: Elliptic Partial Differential Equations, 2nd. Belin: Springer, (1997)
- [18] Guo, Y.J., Lin, C.S. and Wei, J.C.: Local uniqueness and refined spike profiles of ground states for two-dimensional attractive Bose-Einstein condensates. *SIAM J. Math. Anal.* **49** (5), 3671–3715 (2017)
- [19] Guo, Y.J., Seiringer, R.: On the mass concentration for Bose-Einstein condensates with attractive interactions. *Lett. Math. Phys.* **104**, 141–156 (2014)
- [20] Guo, Y.J., Wang, Z.Q., Zeng, X.Y. and Zhou, H.S.: Properties for ground states of attractive Gross-Pitaevskii equations with multi-well potentials. *Nonlinearity* **31**, 957–979 (2018)

- [21] Guo, Y.J., Zeng, X.Y. and Zhou, H.S.: Concentration behavior of standing waves for almost mass critical Schrödinger equations. *J. Differential Equations* **256**, 2079–2100 (2014)
- [22] Guo, Y.J., Zeng, X.Y. and Zhou, H.S.: Energy estimates and symmetry breaking in attractive Bose-Einstein condensates with ring-shaped potentials. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **33**, 809–828 (2016)
- [23] Han, Q. Lin, F.: Elliptic Partial Differential Equations, 2nd, Courant Lecture Notes in Mathematics, Vol. 1. *Courant Institute of Mathematical Science/AMS*, New York (2011)
- [24] Li, Y., Ni, W.M.: Radial symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n . *Comm. Partial Diff. Eqns.* **18**, 1043–1054 (1993)
- [25] Lieb, E. H., Loss, M.: Analysis, 2nd ed., Graduate Studies in Mathematics Vol. 14 *American Mathematical Society*, (2001).
- [26] Lions, P.L.: The concentration-compactness principle in the calculus of variations. The locally compact case I. *Ann. Inst. H. Poincaré. Anal. Non Linéaire* **1**, 109–145 (1984)
- [27] Lions, P.L.: The concentration-compactness principle in the calculus of variations. The locally compact case II. *Ann. Inst. H. Poincaré. Anal. Non Linéaire* **1**, 223–283 (1984)
- [28] Liu, C.S., Tripathi, V. K. Laser guiding in an axially nonuniform plasma channel. *Phys. Plasmas* **1** 3100–3103 (1994).
- [29] Maeda, M.: On the symmetry of the ground states of nonlinear Schrödinger equation with potential. *Adv. Nonlinear Stud.* **10**, 895–925 (2010)
- [30] Saanouni, T.: Remarks on the inhomogeneous fractional nonlinear Schrödinger equation. *J. Math. Phys.* **57** (8), 081503 (2016)
- [31] Stuart, C.A.: Bifurcation in $L^p(\mathbb{R}^N)$ for a semilinear elliptic equation. *Proc. London Math. Soc.* **57**, 511–541 (1988)
- [32] Wang, X.F.: On concentration of positive bound states of nonlinear Schrödinger equations. *Commun. Math. Phys.* **153**, 229–244 (1993)
- [33] Yanagida, E.: Uniqueness of positive radial solutions of $\Delta u + g(r)u + h(r)u^p = 0$ in \mathbb{R}^n . *Arch. Rational Mech. Anal.* **115**, 257–274 (1991)
- [34] Zhang, J.: Stability of standing waves for nonlinear Schrödinger equations with unbounded potentials. *Z. Angew. Math. Phys.* **51**, 498–503 (2000)