

GENERALIZED TORSION FOR HYPERBOLIC 3-MANIFOLD GROUPS WITH ARBITRARY LARGE RANK

TETSUYA ITO, KIMIHIKO MOTEGI, AND MASAKAZU TERAGAITO

ABSTRACT. Let G be a group and g a non-trivial element in G . If some non-empty finite product of conjugates of g equals to the trivial element, then g is called a *generalized torsion element*. To the best of our knowledge, we have no hyperbolic 3-manifold groups with generalized torsion elements whose rank is explicitly known to be greater than two. The aim of this short note is to demonstrate that for a given integer $n > 1$ there are infinitely many closed hyperbolic 3-manifolds M_n which enjoy the property: (i) the Heegaard genus of M_n is n , (ii) the rank of $\pi_1(M_n)$ is n , and (iii) $\pi_1(M_n)$ has a generalized torsion element. Furthermore, we may choose M_n as homology lens spaces and so that the order of the generalized torsion element is arbitrarily large.

1. INTRODUCTION

Let G be a group and let g be a non-trivial element in G . If some non-empty finite product of conjugates of g equals to the identity, then g is called a *generalized torsion element*. In particular, any non-trivial torsion element is a generalized torsion element. As a natural generalization of the order of torsion element, the *order* of the generalized torsion element g is defined as

$$\min\{k \geq 2 \mid \exists x_1, \dots, x_k \in G \text{ s.t. } (x_1 g x_1^{-1})(x_2 g x_2^{-1}) \cdots (x_k g x_k^{-1}) = 1\}.$$

A group G is said to be *bi-orderable* if G admits a strict total ordering $<$ which is invariant under multiplication from the left and right. That is, if $g < h$, then $agb < ahb$ for any $g, h, a, b \in G$. In this paper, the trivial group $\{1\}$ is considered to be bi-orderable.

It is easy to see that a bi-orderable group does not have a generalized torsion element. Thus the existence of generalized torsion element is an obstruction for a group to be bi-orderable. It is known that the converse does not hold in general [9, Chapter 4]. However, for 3-manifold groups (fundamental groups of 3-manifolds), one may expect that the converse does hold, and in [7] the authors proposed the conjecture: a 3-manifold group is bi-orderable if and only if it has no generalized torsion element.

The first named author has been partially supported by JSPS KAKENHI Grant Number JP19K03490, 21H04428.

The second named author has been partially supported by JSPS KAKENHI Grant Number JP19K03502, 21H04428, and Joint Research Grant of Institute of Natural Sciences at Nihon University for 2021.

The third named author has been partially supported by JSPS KAKENHI Grant Number JP20K03587.

2020 *Mathematics Subject Classification*. Primary 57K10, 57M05, 57M07, 57M50 Secondary 20F05, 20F60, 06F15

The conjecture holds for Seifert fiber spaces and Sol manifolds [7]. In particular, there is a Seifert fiber space M such that $\pi_1(M)$ has a generalized torsion element and the rank of $\pi_1(M)$ is arbitrarily large.

Very recently Sekino [13] proves that the conjecture holds for once punctured torus bundles. In particular, he gives tunnel number two once punctured torus bundles whose fundamental group has a generalized torsion element.

Furthermore, since taking connected sum or gluing along incompressible tori preserves generalized torsion, we may also construct non-Seifert fibered 3-manifolds M so that $\pi_1(M)$ has a generalized torsion element and arbitrarily large rank. See also [4] for a behavior of generalized torsion elements under prime decomposition and torus decomposition.

Let us restrict our attention to hyperbolic 3-manifold groups. Two-generator, one-relator hyperbolic 3-manifold groups with generalized torsion elements are given in [10, 14, 15, 8]. On the other hand, to the best of our knowledge, we have no hyperbolic 3-manifold groups with generalized torsion elements whose rank is explicitly known to be greater than two.

From a viewpoint of combinatorial group theory, it seems to be difficult to handle groups with more than two generators. The aim of this short note is to demonstrate the following by using a geometric interpretation of generalized torsion elements.

Theorem 1.1. *For a given integer $n > 1$ there are infinitely many closed hyperbolic 3-manifolds (precisely homology lens spaces) M_n which enjoy the property: (i) the Heegaard genus of M_n is n , (ii) the rank of $\pi_1(M_n)$ is n , and (iii) $\pi_1(M_n)$ has a generalized torsion element. Furthermore, we may choose M_n so that the order of the generalized torsion element is arbitrarily large.*

2. SINGULAR SPANNING DISK AND GENERALIZED TORSION

We recall a useful construction to produce a generalized torsion element. A *singular spanning disk* of a knot K in S^3 is a smooth map $\Phi : D^2 \rightarrow S^3$ (or, its image) such that $\Phi(\partial D) = K$ and that K intersects $\Phi(\text{int}D)$ transversely in finitely many points. Each intersection point $\Phi(\text{int}D \cap K)$ has a sign according to the orientations. We call $\Phi(D)$ a (p, q) -*singular spanning disk* if K intersects $\Phi(\text{int}D)$ positively in p points and negatively in q points.

For a knot $K \subset S^3$, we denote the 3-manifold obtained by p/q -surgery on K by $K(p/q)$. Without loss of generality, we may assume $p \geq 0$.

We recall the following result [5].

Theorem 2.1. *Let K be a knot in S^3 . If K has a $(n, 0)$ -singular spanning disk, then the image of a meridian μ in $\pi_1(K(p/q))$ is a generalized torsion element of order p whenever $\frac{p}{q} \geq n$. Similarly, if K has a $(0, n)$ -singular spanning disk, then the image of a meridian μ in $\pi_1(K(p/q))$ is a generalized torsion element of order p whenever $\frac{p}{q} \leq -n$.*

A singular disk D which has only clasp singularity as illustrated in Figure 2.1 is said to be *coherent* if K intersects $\text{int}D$ in the same direction. A typical and fundamental example of $(n, 0)$ - or $(0, n)$ -singular spanning disk is a *coherent clasp disk*.




FIGURE 2.1. Clasp disk

3. PROOF OF THEOREM 1.1

We denote the Heegaard genus of a 3-manifold M by $g(M)$. We prove Theorem 1.1 by giving a family of concrete examples below.

Let K_n be a Montesinos knot with rational tangles R_1, \dots, R_{n-1}, R_n , where $R_i = [2, -2, 2]$ for $i = 1, \dots, n-1$ and $R_n = [-6, 4]$. Then K_n is a Montesinos knot

$$M \left(\frac{\beta_1}{\alpha_1}, \dots, \frac{\beta_{n-1}}{\alpha_{n-1}}, \frac{\beta_n}{\alpha_n} \right), \text{ where } \frac{\beta_i}{\alpha_i} = \frac{5}{3} \text{ (} 1 \leq i \leq n-1 \text{), and } \frac{\beta_n}{\alpha_n} = \frac{17}{6}.$$




FIGURE 3.1. $(2n + 4, 0)$ -singular spanning disk for K_n

We collect some desired properties of K_n as several claims.

Claim 3.1. *The tunnel number of K_n is $n - 1$, i.e. $g(E(K_n)) = n$.*

Proof. Since $\gcd(\alpha_1, \dots, \alpha_n) = 3 \neq 1$, Theorem 0.1(1) in [6] shows that the tunnel number of K_n is $n - 1$, i.e. $g(E(K_n)) = n$. \square

Claim 3.2. *Assume that p is even. Then $g(K_n(p/q)) = \text{rank } \pi_1(K_n(p/q)) = n$.*

Proof. Since p is even and $\gcd(\alpha_1, \dots, \alpha_n) = 3 \neq 1$, we apply [6, Theorem 0.1(3)] to obtain the desired result. \square

Claim 3.3. $\pi_1(K_n(p/q))$ has a generalized torsion element of order p whenever $p/q \geq 2n - 4$.

Proof. As shown in Figure 3.1, we have a singular spanning disk $\Phi: D \rightarrow S^3$ such that $\Phi(\partial D) = K_n$ and K_n intersects $\Phi(\text{int}D)$ positively in $2(n-1) + 6 = 2n + 4$ points, i.e. K_n bounds a $(2n + 4, 0)$ -singular spanning disk. Hence, it follows from Theorem 2.1 that $\pi_1(K_n(p/q))$ has a generalized torsion element of order p whenever $p/q \geq 2n + 4$. \square

Claim 3.4. K_n is a hyperbolic knot.

Proof. Following [11], K_n is either a hyperbolic knot or a torus knot. If $n \geq 3$, then the tunnel number of K_n is $n - 1 \geq 2$ (Claim 3.1). Since any nontrivial torus knot has tunnel number one [1], K_n is a hyperbolic knot.

When $n = 2$, the tunnel number of K_n is one. However, it is easy to see that K_n is a two-bridge knot $C(2, -2, 6, -7, 1)$ in the Conway notation. By the classification of two-bridge knots, K_n cannot be a two-bridge torus knot $T_{2,q}$ for any odd integer q . (See also [2].) \square

Claim 3.5. $K_n(p/q)$ is hyperbolic for all but finitely many $p/q \in \mathbb{Q}$. In particular, if $n \geq 4$, then $K_n(p/q)$ is hyperbolic for all $p/q \in \mathbb{Q}$.

Proof. Since K_n is hyperbolic (Claim 3.4), the first assertion follows from Thurston's hyperbolic Dehn surgery theorem [16, 17]. The second assertion follows from [18]. \square

It follows from Claims 3.1, 3.2, 3.3 and 3.5 that for a given integer $n > 1$, we may choose an even integer p so that a homology lens space $M_n = K_n(p/q)$ enjoys the required property.

This completes a proof of Theorem 1.1. \square

Remark 3.6. Since $\pi_1(K_n(p/q))$ has a generalized torsion element, it is not bi-orderable.

Currently, when we restrict to hyperbolic case, our technique of constructing a generalized torsion element based on singular clasp disk only can produce a non-integral homology sphere; a generalized torsion obtained by this method is a nontrivial torsion element of its homology.

Question 3.7. Does there exist a hyperbolic 3-manifold with torsion-free 1st homology group whose fundamental group has arbitrarily large rank and generalized torsion? In particular, does there exist a hyperbolic integral homology sphere whose fundamental group has arbitrarily large rank and generalized torsion?

Recall that in our current knowledge, as for a bi-orderability of a knot group, the Alexander polynomial plays a fundamental role; when K is fibered (more generally, when $[G(K), G(K)]$ is residually torsion-free-nilpotent) and all the roots of its Alexander polynomial $\Delta_K(t)$ is positive real, then $G(K)$ is bi-orderable, hence in particular, $G(K)$ has no generalized torsion.

Conversely, when K is rationally homologically fibered, i.e. $\deg \Delta_K(t) = 2g(K)$, and $\Delta_K(t)$ has no positive real roots then $G(K)$ is not bi-orderable [3]. To prove non-bi-orderability, we usually construct a generalized torsion element in the meta-abelian quotient $G(K)/[[G(K), G(K)], [G(K), G(K)]]$, with a help of the Alexander

polynomial, providing a clue to find a generalized torsion element of $G(K)$ itself. Thus, from a bi-orderability point of view, the following counterpart of Question 3.7 is interesting.

Question 3.8. *Does there exist a hyperbolic knot with trivial Alexander polynomial whose fundamental group has arbitrarily large rank and generalized torsion?*

REFERENCES

- [1] M. Boileau, M. Rost and H. Zieschang; On Heegaard decompositions of torus knot exteriors and related Seifert fibred spaces, *Math. Ann.* **279** (1988) 553–581.
- [2] F. Bonahon and L. Siebenmann; New Geometric Splittings of Classical Knots, and the Classification and Symmetries of Arborescent Knots, freely available at <http://www-bcf.usc.edu/~fbonahon>.
- [3] T. Ito; Alexander polynomial obstruction of bi-orderability for rationally homologically fibered knot groups, *New York J. Math.* **23** (2017), 497–503.
- [4] T. Ito, K. Motegi and M. Teragaito; *Generalized torsion and decomposition of 3-manifolds*, *Proc. Amer. Math. Soc.* **147** (2019), 4999–5008.
- [5] T. Ito, K. Motegi, and M. Teragaito; *Generalized torsion and Dehn filling*, *Topology Appl.* **301** (2021), Paper No.107515.
- [6] M. Lustig and Y. Moriah; generalized Montesinos knots, tunnels and \mathcal{N} -torsion, *Math. Ann.* **295** (1992), 167–189.
- [7] K. Motegi and M. Teragaito; Generalized torsion elements and bi-orderability of 3-manifold groups, *Canad. Math. Bull.* **60** (2017), 830–844.
- [8] K. Motegi and M. Teragaito; Generalized torsion for knots with arbitrarily high genus (arXiv:2106.14449), to appear in *Canad. Math. Bull.*
- [9] R. Mura and A. Rhemtulla; Orderable groups, *Lecture Notes in Pure and Applied Mathematics*, Vol. **27**. Marcel Dekker, Inc., New York-Basel, 1977.
- [10] G. Naylor and D. Rolfsen; Generalized torsion in knot groups, *Canad. Math. Bull.* **59** (2016), 182–189.
- [11] U. Oertel; Closed incompressible surfaces in complements of star links, *Pacific J. Math.* **111** (1984), 209–230.
- [12] H. Schubert; Knoten mit zwei Brücken, *Math. Z.* **65** (1956), 133–170.
- [13] N. Sekino; Generalized torsions in once punctured torus bundles, arXiv:2110.13411
- [14] M. Teragaito; Generalized torsion elements in the knot groups of twist knots, *Proc. Amer. Math. Soc.* **144** (2016), no.6, 2677–2682.
- [15] M. Teragaito; Generalized torsion elements and hyperbolic links, *J. Knot Theory Ramifications* **29** (2020), no.11, 2050079 (9 pages)
- [16] W.P. Thurston; The geometry and topology of 3-manifolds, *Lecture notes*, Princeton University, 1979.
- [17] W.P. Thurston; Three dimensional manifolds, Kleinian groups and hyperbolic geometry, *Bull. Amer. Math. Soc.* **6** (1982), 357–381.
- [18] Y-Q. Wu; Dehn surgery on arborescent knots, *J. Diff. Geom.* **43** (1996), 171–197.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN
Email address: `tetitoh@math.kyoto-u.ac.jp`

DEPARTMENT OF MATHEMATICS, NIHON UNIVERSITY, 3-25-40 SAKURAJOSUI, SETAGAYA-KU, TOKYO 156-8550, JAPAN
Email address: `motegi.kimihiko@nihon-u.ac.jp`

DEPARTMENT OF MATHEMATICS AND MATHEMATICS EDUCATION, HIROSHIMA UNIVERSITY, 1-1-1 KAGAMIYAMA, HIGASHI-HIROSHIMA, 739-8524, JAPAN
Email address: `teragai@hiroshima-u.ac.jp`