

# ON THE CIRCUITS OF SPLITTING MATROIDS REPRESENTABLE OVER $GF(p)$

PRASHANT MALAVADKAR<sup>1</sup>, UDAY JAGDALE<sup>2</sup> AND SACHIN GUNJAL<sup>3</sup>

School of Mathematics and Statistics, MIT-World Peace University, Pune 411 038, India.

e-mail: 1. [prashant.malavadkar@mitwpu.edu.in](mailto:prashant.malavadkar@mitwpu.edu.in), 2. [uday.jagdale@mitwpu.edu.in](mailto:uday.jagdale@mitwpu.edu.in),

3. [sachin.gunj@mitwpu.edu.in](mailto:sachin.gunj@mitwpu.edu.in)

## Abstract

We extend the splitting operation from binary matroids (Raghunathan et al., 1998) to  $p$ -matroids, where  $p$ -matroids refer to matroids representable over  $GF(p)$ . We also characterize circuits, bases, and independent sets of the resulting matroid. Sufficient conditions to yield Eulerian  $p$ -matroids from Eulerian and non-Eulerian  $p$ -matroids by applying the splitting operation are obtained. A class of connected  $p$ -matroids that gives connected  $p$ -matroids under the splitting operation is characterized.

**Keywords:** connected matroid; splitting operation; Eulerian matroid; circuits

**AMS Subject Classification:** 05B35; 05C50; 05C83

## 1 Introduction

Fleischer [2] characterized Eulerian graphs using the notion of splitting away a pair of edges from a vertex of degree at least three. Raghunathan et al. [7] extended the splitting operation from graphs to binary matroids, and characterization of Eulerian binary matroids was obtained using this operation as follows:

**Theorem 1.1.** *A binary matroid  $M$  on a set  $E$  is Eulerian if and only if  $M$  can be transformed by repeated applications of the splitting operation into a matroid in which  $E$  is a circuit.*

A sufficient condition was provided by Shikare [9] to obtain connected binary matroids from a 4-connected binary matroids using the splitting operation:

**Theorem 1.2.** *If  $M$  is a 4-connected binary matroid on at least 9 elements and  $a, b$  are distinct elements of  $M$ , then  $M_{a,b}$  is a connected binary matroid.*

The splitting operation and its various properties were discussed in [1, 3, 4, 5, ?, 8, 12, 14].

In this paper, simple and coloopless matroids representable over  $GF(p)$  are discussed. A matroid  $M$  is called a  $p$ -matroid if  $M \cong M[A]$ , where  $M[A]$  is the vector matroid of matrix  $A$  of size  $m \times n$  over the field  $F = GF(p)$  for some prime  $p$ . We denote the

set of column labels of  $M$  (viz. the ground set of  $M$ ) by  $E$ , the set of circuits of  $M$  by  $\mathcal{C}(M)$ , and the set of independent sets of  $M$  by  $\mathcal{I}(M)$ . For undefined and standard terminologies, refer to Oxley [6].

In the present paper, we define the splitting operation on  $p$ -matroids and give a characterization of the circuits of the resulting  $p$ -matroid. We give a sufficient condition for the  $p$ -matroid to be connected after the splitting operation. A sufficient condition to yield Eulerian  $p$ -matroid from Eulerian  $p$ -matroid after the splitting operation is also obtained.

## 2 Splitting Operation on $p$ -Matroids

Let  $M$  be a  $p$ -matroid on ground set  $E$ ,  $\{a, b\} \subset E$  and  $\alpha$  is a non-zero element of  $GF(p)$ . We define splitting operation on a  $p$ -matroid  $M$  as follows.

**Definition 2.1.** Let  $M \cong M[A]$  be a  $p$ -matroid on ground set  $E$ ,  $\{a, b\} \subset E$ , and  $\alpha \neq 0$  in  $GF(p)$ . The matrix  $A_{a,b}$  is constructed from  $A$  by appending an extra row to  $A$  which has coordinates equal to  $\alpha$  in the columns corresponding to the elements  $a, b$  and zero elsewhere. Define the splitting matroid  $M_{a,b}$  to be the vector matroid  $M[A_{a,b}]$ . The transformation of  $M$  to  $M_{a,b}$  is called the splitting operation.

**Example 2.2.** Consider the vector matroid  $M[A]$  of a matrix  $A$  over a field  $GF(3)$ .

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \end{pmatrix} \quad \mathbf{A}_{3,5} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 2 & 0 & 2 & 0 & 0 & 0 \end{pmatrix}$$

For  $a = 3$ ,  $b = 5$  and  $\alpha = 2$ , the matrix  $A_{3,5}$ , represents the splitting matroid  $M_{3,5}$ .

The collection of circuits of  $M$  and  $M_{3,5}$  is given in the following table.

Circuits of $M$	Circuits of $M_{3,5}$
$\{1, 2, 4, 8\}, \{1, 2, 6, 7\}, \{3, 5, 6, 8\}, \{4, 6, 7, 8\}$	$\{1, 2, 4, 8\}, \{1, 2, 6, 7\}, \{3, 5, 6, 8\}, \{4, 6, 7, 8\}$
$\{1, 2, 4, 5, 7\}, \{3, 4, 5, 7\}, \{3, 4, 8\}$	$\{1, 2, 3, 4, 5, 7\}, \{3, 4, 5, 6, 7\}, \{3, 4, 5, 7, 8\}$
$\{1, 2, 3\}, \{4, 5, 6\}, \{5, 7, 8\}, \{3, 6, 7\}$	$\{1, 2, 3, 4, 5, 6\}, \{1, 2, 3, 5, 7, 8\}$
$\{1, 2, 5, 6, 8\}$	—

It is interesting to observe that unlike the splitting in binary matroids, circuits  $\{1, 2, 3, 4, 5, 7\}$ ,  $\{3, 4, 5, 6, 7\}$ ,  $\{3, 4, 5, 7, 8\}$  of  $M_{3,5}$  are neither circuits of  $M$  nor disjoint union of circuits of  $M$ .

**Remark 2.3.** Let  $\{a, b\}$  be a cocircuit of  $M$ . Then  $M_{a,b} \cong M$ .

**Remark 2.4.** *Observe that  $\text{rank}(A) \leq \text{rank}(A_{a,b}) \leq \text{rank}(A) + 1$ , which implies that  $\text{rank}(M) \leq \text{rank}(M_{a,b}) \leq \text{rank}(M) + 1$ .*

**Remark 2.5.** *When  $p = 2$ , the splitting operation coincides with the splitting operation for binary matroids introduced by Raghunathan et al.[7].*

In the following discussion, we describe circuits of  $M_{a,b}$  with respect to circuits of  $M$ .

**Lemma 2.6.** *If  $C \in \mathcal{C}(M)$  and  $C \cap \{a, b\} = \emptyset$ , then  $C$  is a circuit of  $M_{a,b}$ .*

*Proof.* Observe that  $C$  is a dependent set of  $M_{a,b}$ . If  $C$  is not a circuit of  $M_{a,b}$ , then  $C$  contains a circuit  $C'$  of  $M_{a,b}$ . Note that  $C'$  is a dependent set of  $M$  and  $C' \subset C$  which is not possible. Therefore  $C$  is a circuit of  $M_{a,b}$ .  $\square$

Let  $C_k = \{u_1, u_2, \dots, u_l : u_i \in E, i = 1, 2, \dots, l\}$  be a circuit of  $M$  for some positive integers  $k, l$  and  $|C_k \cap \{a, b\}| = 2$ . Suppose, without loss of generality,  $u_1 = a, u_2 = b$ . If there are non-zero constants  $a_1^k, a_2^k, \dots, a_l^k$  in  $GF(p)$  such that  $\sum_{i=1}^{i=l} a_i^k u_i \equiv 0 \pmod{p}$  and  $a_1^k + a_2^k \equiv 0 \pmod{p}$  then we call  $C_k$  a  $p$ -circuit of  $M$ . However, if  $|C_k \cap \{a, b\}| = 1$  or  $|C_k \cap \{a, b\}| = 2$  and  $a_1^k + a_2^k \not\equiv 0 \pmod{p}$ , then we call  $C_k$  an  $np$ -circuit of  $M$ .

We denote  $\mathcal{C}_0 = \{C \in \mathcal{C}(M) : C \text{ is a } p\text{-circuit or } C \cap \{a, b\} = \emptyset\}$ .

**Remark 2.7.** *Let  $C$  and  $\{a, b\}$  be a circuit and a cocircuit of a  $p$ -matroid  $M$ , respectively. If  $p = 2$ , then every circuit of  $M$  is a circuit of  $M_{a,b}$ . But for  $p > 2$  every circuit of  $M$  may not be a circuit of  $M_{a,b}$ . Note that  $|C \cap \{a, b\}| \neq 1$ , therefore  $|C \cap \{a, b\}|$  is even. If  $a, b \in C$  and  $C$  is an  $np$ -circuit of  $M$ , then  $C \in \mathcal{I}(M_{a,b})$ .*

**Lemma 2.8.** *Let  $M$  be a  $p$ -matroid on ground set  $E$  and  $\{a, b\} \subset E$ . If  $C_1$  and  $C_2$  are disjoint  $np$ -circuits of  $M$  such that  $a \in C_1, b \in C_2$ , then  $C_1 \cup C_2$  is a dependent set of  $M_{a,b}$ .*

*Proof.* Let  $C_1 = \{u_1, u_2, \dots, u_l\}$  and  $C_2 = \{v_1, v_2, \dots, v_k\}$ . Without loss of generality, assume  $u_1 = a$  and  $v_1 = b$ . Since  $C_1$  and  $C_2$  are circuits, there exist non-zero constants  $\alpha_1, \alpha_2, \dots, \alpha_l$  and  $\beta_1, \beta_2, \dots, \beta_k$  such that

$$\alpha_1 u_1 + \alpha_2 u_2 + \dots + \alpha_l u_l \equiv 0 \pmod{p} \quad (1)$$

$$\beta_1 v_1 + \beta_2 v_2 + \dots + \beta_k v_k \equiv 0 \pmod{p}. \quad (2)$$

Multiplying equation (1) by  $\alpha_1^{-1}$  and equation (2) by  $-\beta_1^{-1}$  we get

$$u_1 + \alpha'_2 u_2 + \dots + \alpha'_l u_l \equiv 0 \pmod{p},$$

$$-v_1 + \beta'_2 v_2 + \dots + \beta'_k v_k \equiv 0 \pmod{p}.$$

Thus we get non-zero scalars  $1, \alpha'_i, i = 2, 3, \dots, l$  and  $-1, \beta'_j, j = 2, 3, \dots, k$  such that

$$u_1 + \alpha'_2 u_2 + \dots + \alpha'_l u_l + (-1)v_1 + \beta'_2 v_2 + \dots + \beta'_k v_k \equiv 0$$

and

$$\text{coeff.}(u_1) + \text{coeff.}(v_1) \equiv 0 \pmod{p}.$$

Therefore  $C_1 \cup C_2$  is a dependent set of  $M_{a,b}$ .  $\square$

**Remark 2.9.** Let  $C_1, C_2$  be disjoint  $np$ -circuits of  $M$  such that  $a \in C_1, b \in C_2$  and  $I \neq \phi$  be an independent set of  $M$ . Then  $C_1 \cup C_2 \cup I$  can not be a circuit of  $M_{a,b}$  as it contains the dependent set  $C_1 \cup C_2$  of  $M_{a,b}$ .

Consider subsets of  $E$  of the type  $C \cup I$  where  $C = \{u_1, u_2, \dots, u_l\}$  is an  $np$ -circuit of  $M$  which is disjoint from an independent set  $I = \{v_1, v_2, \dots, v_k\}$  and  $\{a, b\} \subset (C \cup I)$ . We say  $C \cup I$  is  $p$ -dependent if it contains no member of  $\mathcal{C}_0$  and there are non-zero constants  $\alpha_1, \alpha_2, \dots, \alpha_l$  and  $\beta_1, \beta_2, \dots, \beta_k$  such that  $\sum_{i=1}^l \alpha_i u_i + \sum_{j=1}^k \beta_j v_j = 0 \pmod{p}$  and  $\text{coeff.}(a) + \text{coeff.}(b) = 0 \pmod{p}$ .

Theorem 2.10 characterizes the circuits of the splitting matroid  $M_{a,b}$  with respect to circuits of  $M$ .

**Theorem 2.10.** Let  $M$  be a  $p$ -matroid on ground set  $E$  and  $\{a, b\} \subset E(M)$ . Then  $\mathcal{C}(M_{a,b}) = \mathcal{C}_0 \cup \mathcal{C}_1 \cup \mathcal{C}_2$  where

$$\mathcal{C}_0 = \{C \in \mathcal{C}(M) : C \text{ is a } p\text{-circuit or } C \cap \{a, b\} = \phi\};$$

$$\mathcal{C}_1 = \text{Set of minimal members of } \{C \cup I : (C \cup I) \text{ is } p\text{-dependent and there are no disjoint } np\text{-circuits } C'_1, C'_2 \text{ such that } C'_1 \cup C'_2 \subset (C \cup I)\}; \text{ and}$$

$$\mathcal{C}_2 = \text{Set of minimal members of } \{C_1 \cup C_2 : C_1, C_2 \in \mathcal{C}(M), a \in C_1, b \in C_2, C_1 \cap C_2 = \phi \text{ and there is no } C \in \mathcal{C}_0 \cup \mathcal{C}_1 \text{ such that } C \subset C_1 \cup C_2\}.$$

*Proof.* By the Definition 2.1 and Lemma 2.8, it is clear that every element of  $\mathcal{C}_0 \cup \mathcal{C}_1 \cup \mathcal{C}_2$  is a circuit of  $M_{a,b}$ . Conversely, let  $C \in \mathcal{C}(M_{a,b})$ , then  $C$  is a dependent set of  $M$ .

**Case I:** If  $C$  is a minimally dependent set of  $M$ , then either  $C \cap \{a, b\} = \phi$  or  $C$  is a  $p$ -circuit of  $M$ . That is,  $C \in \mathcal{C}_0$ .

**Case II:** If  $C$  is not a minimally dependent set of  $M$ , then it contains a minimally dependent set, say  $C_1$ . Note that  $C_1 \notin \mathcal{C}_0$ . Thus  $C$  contains no member of  $\mathcal{C}_0$ .

If  $a, b \in C_1$ , then  $C \setminus C_1$  is an independent set of  $M$ ; otherwise  $C \setminus C_1$  contains a circuit, say  $C'$ , such that  $C' \in \mathcal{C}_0$  which is not possible. Thus  $C = C_1 \cup I$  where  $C_1$  is an  $np$ -circuit,  $I = C \setminus C_1 \in \mathcal{I}(M)$  and it contains no member of  $\mathcal{C}_0$ . Therefore  $C$  is a minimal  $p$ -dependent set of  $M$  and it does not contain a disjoint union of two  $np$ -circuits. Thus  $C \in \mathcal{C}_1$ .

If  $a \in C_1$ ,  $b \in C \setminus C_1$  and  $C \setminus C_1$  is an independent set of  $M$ , denoted by  $I$ , then  $C = C_1 \cup I$ . Note that  $C_1 \cup I$  is a minimal  $p$ -dependent set of  $M$  and it does not contain a disjoint union of two  $np$ -circuits. Thus  $C \in \mathcal{C}_1$ .

If  $a \in C_1$ ,  $b \in C \setminus C_1$  and  $C \setminus C_1$  is dependent set of  $M$ , then there is a circuit, say  $C_2$ , contained in  $C \setminus C_1$  and  $b \in C_2$ . Therefore  $C = C_1 \cup C_2 \cup I$  where both  $C_1$  and  $C_2$  are  $np$ -circuits and  $I = C \setminus (C_1 \cup C_2)$  is an independent set of  $M$ . By Remark 2.9,  $I = \phi$ . Further,  $C = C_1 \cup C_2$  does not contain  $p$ -dependent set of  $M$  and a disjoint union of two  $np$ -circuits. Therefore  $C \in \mathcal{C}_2$ .

□

In the Example 2.2,  $\mathcal{C}(M_{3,5}) = \mathcal{C}_0 \cup \mathcal{C}_1 \cup \mathcal{C}_2$  where

$$\mathcal{C}_0 = \{\{1, 2, 4, 8\}, \{1, 2, 6, 7\}, \{3, 5, 6, 8\}, \{4, 6, 7, 8\}\};$$

$$\mathcal{C}_1 = \{\{1, 2, 3, 4, 5, 7\}, \{3, 4, 5, 6, 7\}, \{3, 4, 5, 7, 8\}\};$$

$$\mathcal{C}_2 = \{\{1, 2, 3, 4, 5, 6\}, \{1, 2, 3, 5, 7, 8\}\}.$$

## 2.1 Independent sets, Bases and Rank function of $M_{a,b}$

Now we describe independent sets, bases and the rank function of  $M_{a,b}$  in terms of independent sets, bases and the rank function of  $M$ , respectively.

Let  $\mathcal{I}_0 = \mathcal{I}(M)$  and  $\mathcal{I}_1 = \{C \cup I : (C \cup I) \text{ is not } p\text{-dependent and it contains no union of two disjoint } np\text{-circuits}\}$ .

**Proposition 2.11.**  $\mathcal{I}(M_{a,b}) = \mathcal{I}_0 \cup \mathcal{I}_1$ .

*Proof.* Clearly  $\mathcal{I}_0 \cup \mathcal{I}_1 \subseteq \mathcal{I}(M_{a,b})$ . Conversely, assume  $S \in \mathcal{I}(M_{a,b})$ . If  $S$  is an independent set of  $M$ , then  $S \in \mathcal{I}_0$ . If  $S$  is dependent set of  $M$ , then it contains an  $np$ -circuit of  $M$ . In the light of Lemma 2.8,  $S$  does not contain union of two disjoint  $np$ -circuits. Therefore  $S = C \cup I$  for some  $C \in \mathcal{C}(M)$ ,  $I \in \mathcal{I}(M)$  and  $C \cup I$  is not  $p$ -dependent. Hence  $S \in \mathcal{I}_1$ .

□

We use  $r$  and  $r'$  to denote the rank functions of matroid  $M$  and  $M_{a,b}$  respectively.

**Theorem 2.12.** *Let  $M$  be a  $p$ -matroid and  $\{a, b\} \subset E$ . If  $M$  contains an  $np$ -circuit, then  $\mathcal{B}(M_{a,b}) = \{B \cup e : B \in \mathcal{B}(M), e \notin B \text{ and } B \cup e \text{ contains neither } p\text{-circuit nor } p\text{-dependent set}\}$ .*

*Proof.* Let  $B \in \mathcal{B}(M)$  and  $e \notin B$ . If  $B \cup e$  contains no  $p$ -circuit and no  $p$ -dependent set, then  $B \cup e$  is an independent set of  $M_{a,b}$ . Moreover,  $B \cup e$  is a maximal independent set of  $M_{a,b}$  because  $r'(M_{a,b}) \leq r(M) + 1$ .

Conversely, let  $B'$  be a basis of  $M_{a,b}$ . Note that  $B'$  is an independent subset of  $M$  if and only if  $M$  contains no  $np$ -circuit. Since  $M$  contains an  $np$ -circuit,  $B'$  is a dependent set of  $M$ . If  $B'$  contains a  $p$ -circuit, then  $B'$  becomes dependent set of  $M_{a,b}$ , a contradiction. Therefore  $B'$  contains an  $np$ -circuit, say  $C$ . Thus  $B' = C \cup I$  where  $I = B' \setminus C$ . Note that  $I$  is an independent set of  $M$ , otherwise there is a  $np$ -circuit, say  $C_1$ , of  $M$  contained in  $I$  and  $C \cup C_1$  gives a dependent subset of  $B'$  in  $M_{a,b}$ , a contradiction.

Let  $e \in C$  and  $B = B' \setminus e = (C \setminus e) \cup I$ . Now if  $B$  is a dependent set of  $M$ , then  $r[(C \setminus e) \cup I] < r(M)$ . It implies  $r[(C \cup I)] < r(M)$  and  $r'(C \cup I) \leq r(M)$ . That is,  $r'(B') \leq r(M)$ , a contradiction. Therefore  $B = B' \setminus e$  is an independent set of  $M$ . Moreover,  $|B| = r(M)$ , which implies that  $B$  is a basis of  $M$ . Thus  $B' = B \cup e$ .  $\square$

In the following corollary, we provide the rank function of  $M_{a,b}$  with respect to the rank function of  $M$ .

**Corollary 2.13.** *Suppose  $S \subseteq E(M_{a,b})$ . Then*

$$\begin{aligned} r'(S) &= r(S), & \text{if } S \text{ contains no } np\text{-circuit of } M; \text{ and} \\ &= r(S) + 1, & \text{if } S \text{ contains an } np\text{-circuit of } M. \end{aligned} \quad (3)$$

*Proof.* Let  $B$  and  $B'$  be the bases of  $M|_S$  and  $M_{a,b}|_S$ , respectively, then  $r(S) = |B|$  and  $r'(S) = |B'|$ . If  $S$  contains no  $np$ -circuit, then  $|B'| = |B|$ . Therefore  $r(S) = r'(S)$ . If  $S$  contains an  $np$ -circuit, then by the Theorem 2.12,  $B' = B \cup e$  for some  $B \in \mathcal{B}(M|_S)$ ,  $e \in (S \setminus B)$  and  $B \cup e$  contains a unique  $np$ -circuit which is the fundamental circuit of  $e$  with respect to  $B$ . Therefore  $r'(S) = r'(M_{a,b}|_S) = |B'| = |B \cup e| = |B| + 1 = r(S) + 1$ .  $\square$

**Corollary 2.14.** *Let  $M$  be a  $p$ -matroid and  $\{a, b\} \subset E$ , then*

1.  $r'(M_{a,b}) = r(M)$  if and only if every circuit  $C \in \mathcal{C}(M)$  is a  $p$ -circuit. Moreover,  $M \cong M_{a,b}$ .
2. If  $M$  contains an  $np$ -circuit, then  $r'(M_{a,b}) = r(M) + 1$ .

**Remark 2.15.** *Let  $M$  be a  $p$ -matroid and  $\{a, b\} \subset E$ . If  $M$  contains an  $np$ -circuit, then  $\{a, b\}$  is a cocircuit of  $M_{a,b}$ . Assume the contrary, that is,  $\{a, b\}$  is not a cocircuit of  $M_{a,b}$ . Then there is a basis  $B'$  of  $M_{a,b}$  such that  $B' \cap \{a, b\} = \emptyset$ . By Theorem 2.12,  $B' = B \cup e$  for some  $B \in \mathcal{B}(M)$  and  $e \in E \setminus B$ . Now  $B' = B \cup e$  contains a unique  $np$ -circuit of  $M$  which is a contradiction to the assumption that there is a basis  $B'$  of  $M_{a,b}$  such that  $B' \cap \{a, b\} = \emptyset$ . Therefore,  $\{a, b\}$  is a cocircuit of  $M_{a,b}$ .*

### 3 Connectivity of Splitting Matroids

Let  $M$  be a matroid having ground set  $E$ . The 1-separation of the matroid  $M$  is a partition  $(S, T)$  of  $E$  such that,  $|S|, |T| \geq 1$  and  $r(S) + r(T) - r(M) < 1$ . We say  $M$

is connected if  $M$  has no 1-separation.

Observe that in Example 2.2, the matroid  $M$  as well as the splitting matroid  $M_{3,5}$  are connected. In general, the connectivity of a  $p$ -matroid is not closed under the splitting operation, which is observed in the following example.

**Example 3.1.** Consider the vector matroid  $M \cong M[A]$  represented by the matrix  $A$  over the field  $GF(3)$ .

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \mathbf{A}_{1,4} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Here the matroid  $M$  is connected. But the splitting matroid  $M_{1,4} \cong M[A_{1,4}]$  is not connected.

**Remark 3.2.** Let  $M$  be a  $p$ -matroid on ground set  $E$  and  $\{a, b\} \subset E$ . If  $M$  contains no  $np$ -circuit, then  $M_{a,b} \cong M$ . In this case,  $M$  is connected if and only if  $M_{a,b}$  is connected.

In the next result, a sufficient condition is provided for the matroid  $M_{a,b}$  to be connected.

**Lemma 3.3.** Let  $M$  be a connected  $p$ -matroid and  $\{a, b\} \subset E(M)$ . If for every proper subset  $S$  of  $E(M)$  with  $|S| \geq 1$ , either  $S$  or  $T = E(M) \setminus S$  contains an  $np$ -circuit of  $M$ , then  $M_{a,b}$  is connected.

*Proof.* Note that,  $M$  is a connected  $p$ -matroid, which implies that  $M$  has no 1-separation. On the contrary, assume  $M_{a,b}$  is not connected. That is,  $M_{a,b}$  has 1-separation, say  $(S, T)$ . Therefore

$$r'(S) + r'(T) - r'(M_{a,b}) < 1.$$

If  $S$  and  $T$  both contains  $np$ -circuits then, by lemma 2.13, we have

$$r(S) + 1 + r(T) + 1 - r(M) - 1 = r(S) + r(T) - r(M) + 1 < 1.$$

Thus we get a contradiction to the fact  $r(S) + r(T) - r(M) \geq 0$ . Further, if only one of  $S$  or  $T$ , say  $S$ , contains an  $np$ -circuit, then we have

$$r'(S) + r'(T) - r'(M_{a,b}) = r(S) + 1 + r(T) - r(M) - 1 < 1.$$

That is

$$r(S) + r(T) - r(M) < 1.$$

Thus  $(S, T)$  gives a 1-separation of  $M$  which is not possible.  $\square$

For  $n \geq 2$ , a matroid  $M$  is said to be vertically  $n$ -connected if for any positive integer  $k < n$ , there does not exist a partition  $(S, T)$  of  $E(M)$ , such that  $r(S) + r(T) - r(M) < k$ , and  $r(S), r(T) \geq k$ .

**Theorem 3.4.** *Let  $M$  be a connected, vertically 3-connected, simple  $p$ -matroid and  $\{a, b\} \subset E$ . Then  $M_{a,b}$  is connected  $p$ -matroid if and only if for every  $e \in E$  there is an  $np$ -circuit of  $M$  not containing  $e$ .*

*Proof.* On the contrary, suppose that  $M_{a,b}$  is not connected. And  $(S, T)$  is a 1-separation of  $E(M_{a,b})$ . Consequently,  $|S|, |T| \geq 1$  and  $r'(S) + r'(T) - r'(M_{a,b}) \leq 0$ . Following are the possible cases:

**Case I:**  $|S| = 1$

Suppose  $S = \{x\}$  where  $x \in E$ . As  $M$  contains an  $np$ -circuit  $C$  such that  $\{x\} \cap C = \emptyset$ , which gives  $C \subseteq T$ . By Corollary 2.13,  $r'(T) = r(T) + 1$ . We get  $r(S) + r(T) + 1 - r(M) - 1 \leq 0$ . Consequently,  $r(S) + r(T) - r(M) \leq 0$  and  $|S|, |T| \geq 1$ , which forms a 1-separation of  $M$ , a contradiction.

**Case II:**  $|S|, |T| > 1$

If either  $S$  or  $T$  contains an  $np$ -circuit then using Lemma 3.3 we conclude that  $M_{a,b}$  is connected.

Suppose both  $S$  and  $T$  do not contain an  $np$ -circuit. Then  $r'(S) + r'(T) - r'(M_{a,b}) \leq 0$ . Using Corollary 2.13 we get,  $r(S) + r(T) - r(M) - 1 \leq 0$ . That is  $r(S) + r(T) - r(M) \leq 1$ . Since  $M$  is simple,  $r(S), r(T) \geq 2$ . It implies  $(S, T)$  is a vertical 2-separation of  $M$ , a contradiction.

If each of  $S$  and  $T$  contains an  $np$ -circuit, then  $r'(S) + r'(T) - r'(M_{a,b}) \leq 0$ . Again by Corollary 2.13 we get  $r(S) + 1 + r(T) + 1 - r(M) - 1 \leq 0$ . That is  $r(S) + r(T) - r(M) \leq -1$ . Consequently,  $(S, T)$  is a 1 separation of  $M$ , which is a contradiction.

Therefore  $M_{a,b}$  has no 1-separation. We conclude that  $M_{a,b}$  is a connected  $p$ -matroid.

To check the necessity of the condition, suppose that  $M_{a,b}$  is connected. On the contrary, assume that there is an element  $e \in E$  which is contained in every  $np$ -circuit of  $M$ . Let  $S = \{e\}$  and  $T = E \setminus S$ , then  $T$  contains no  $np$ -circuit of  $M$ . Thus  $r'(S) = 1$ ,  $r'(T) = r(T) = r(M)$  and  $|S|, |T| \geq 1$ . Further,  $r'(S) + r'(T) - r'(M_{a,b}) = 1 + r(T) - r(M) - 1 = 0$ . Thus  $(S, T)$  forms a 1-separation of  $M_{a,b}$ , a contradiction. Therefore we conclude that for every  $e \in E(M)$  there is an  $np$ -circuit of  $M$  not containing  $e$ .  $\square$

A matroid  $M$  is connected if and only if, for every pair of distinct elements of  $E$ , there is a circuit containing both. This result is used to prove the following proposition.

**Proposition 3.5.** *Let  $M$  be a non-connected  $p$ -matroid on ground set  $E$  with exactly two connected components  $M_1, M_2$ , and  $\{a, b\} \subset E$ . Let  $a \in M_1, b \in M_2$ . If  $C$  and  $C'$  are  $np$ -circuits of  $M$  contained in  $M_1$  and  $M_2$ , respectively, then  $C \cup C'$  is a circuit of  $M_{a,b}$ .*



*Proof.* By Lemma 2.8,  $C \cup C'$  is a dependent set of  $M_{a,b}$ . Therefore there is a circuit  $C_1$  of  $M_{a,b}$  such that  $C_1 \subseteq C \cup C'$ . Note that  $C_1$  is a dependent set of  $M$ .

Suppose  $C_1 \subset C \cup C'$ . Then  $C_1 \not\subseteq C$  and  $C_1 \not\subseteq C'$  because  $C$  and  $C'$  are minimal dependent sets of  $M$ . Let  $C_1 \cap C \neq \phi$  and  $C_1 \cap C' \neq \phi$ . Then we have the following two cases.

**Case I:** Let  $C_1 \cap \{a, b\} = \phi$ . Then  $C_1$  is a  $p$ -circuit containing elements of  $M_1$  and  $M_2$  which imply that  $M$  is a connected matroid, a contradiction.

**Case II:** Let  $C_1 \cap \{a, b\} \neq \phi$ . Then  $a, b \in C_1$  because  $\{a, b\}$  is a co-circuit of  $M_{a,b}$ . Since  $M$  is not connected,  $C_1$  can not be a circuit of  $M$ . Thus there is a circuit, say  $C_2$ , of  $M$  such that  $C_2 \subset C_1$ . Now by similar argument as above,  $C_2 \not\subseteq C$  and  $C_2 \not\subseteq C'$ . Consequently,  $C_2 \cap C \neq \phi$  and  $C_2 \cap C' \neq \phi$ . Thus  $C_2$  is a circuit of  $M$  containing elements of  $M_1$  and  $M_2$ . Therefore  $M$  is connected which is a contradiction.

Hence, in either case, we get a contradiction. Thus  $C_1 = C \cup C'$ . □

**Corollary 3.6.** *Let  $M$  be a non-connected  $p$ -matroid on ground set  $E$  with exactly two connected components  $M_1, M_2$  and  $\{a, b\} \subset E$ . Let  $a \in M_1, b \in M_2$ . Then  $M_{a,b}$  is a connected  $p$ -matroid.*

*Proof.* It is enough to show that for every pair  $x, y \in E$ , there is a circuit of  $M_{a,b}$  containing  $x$  and  $y$ .

**Case I:** Let  $x, y \in M_1$ . Then there is a circuit, say  $C$ , of  $M$  containing  $x, y$ . If  $C$  is a  $p$ -circuit, then it is the desired circuit of  $M_{a,b}$ . If  $C$  is an  $np$ -circuit of  $M$ , then  $a \in C$ . Let  $C'$  be an  $np$ -circuit of  $M$  contained in  $M_2$ . Then by Proposition 3.5,  $C \cup C'$  is a circuit of  $M_{a,b}$  containing  $x$  and  $y$ .

**Case II:** Let  $x \in M_1$  and  $y \in M_2$ . As  $M_1, M_2$  are connected there are circuits, say  $C_1, C_2$ , containing  $x, a$  and  $y, b$ , respectively, in  $M$ . Note that  $C_1, C_2$  are  $np$ -circuits. By Proposition 3.5,  $C_1 \cup C_2$  is a circuit of  $M_{a,b}$  containing  $x$  and  $y$ . □

## 4 Applications

Eulerian  $p$ -matroids are not closed under the splitting operation as observed in the following example.

**Example 4.1.**  $M = M[A]$  is Eulerian  $p$ -matroid over  $GF(3)$  but the splitting matroid  $M_{1,2} = M[A_{1,2}]$  is not Eulerian  $p$ -matroid.

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 0 \end{pmatrix} \quad \mathbf{A}_{1,2} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Let  $M$  be Eulerian matroid on ground set  $E$ . Then there are disjoint circuits  $C_1, C_2, \dots, C_k$  of  $M$  such that  $E = C_1 \cup C_2 \cup \dots \cup C_k$ . Let  $\{a, b\} \subset E$  and  $M_{a,b}$  be the splitting matroid. We say that the collection  $\tilde{\mathcal{C}} = \{C_1, C_2, \dots, C_k\}$  is a  $p$ -decomposition of  $M$  if  $a, b \in C_i$ , for some  $i \in \{1, 2, \dots, k\}$ , and  $C_i$  is a  $p$ -circuit or  $a \in C_i, b \in C_j$  for some  $i \neq j$  in  $\{1, 2, \dots, k\}$  and  $(C_i \cup C_j) \in \mathcal{C}_2$ .

In the following result we give a sufficient condition to yield Eulerian  $p$ -matroids from Eulerian  $p$ -matroids after the splitting operation.

**Proposition 4.2.** *Let  $M$  be a  $p$ -matroid on ground set  $E$  and  $\{a, b\} \subset E$ . If  $M$  is Eulerian matroid having a  $p$ -decomposition, then  $M_{a,b}$  is Eulerian.*

*Proof.* Let  $\tilde{\mathcal{C}} = \{C_1, C_2, \dots, C_k\}$  be a  $p$ -decomposition of  $M$ . Then

$$E = C_1 \cup C_2 \cup \dots \cup C_k.$$

**Case I:** If  $a, b \in C_i$  some  $i \in \{1, 2, \dots, k\}$  and  $C_i$  is a  $p$ -circuit. Then the same collection  $\tilde{\mathcal{C}}$  is a disjoint circuit decomposition of  $E(M_{a,b})$ . Therefore  $M_{a,b}$  is Eulerian matroid.

**Case II:** Assume, without loss of generality,  $a \in C_1, b \in C_2$ . By hypothesis,  $\tilde{\mathcal{C}}$  is a  $p$ -decomposition, therefore  $(C_1 \cup C_2)$  is a circuit of  $M_{a,b}$ . Denote  $(C_1 \cup C_2)$  by  $C$ . Then the collection  $\{C, C_3, \dots, C_k\}$  is a disjoint circuit decomposition of  $M_{a,b}$ . Hence  $M_{a,b}$  is Eulerian matroid. □

**Proposition 4.3.** *If  $M_{a,b}$  is Eulerian  $p$ -matroid having a disjoint circuit decomposition which contains no member of  $\mathcal{C}_1$ , then  $M$  is Eulerian.*

*Proof.* Let  $E(M_{a,b}) = C_1 \cup C_2 \cup \dots \cup C_k$  be a disjoint circuit decomposition which contains no member of  $\mathcal{C}_1$ . If  $C_i \in \mathcal{C}_0, C_j \in \mathcal{C}_2$  for some  $i, j \in \{1, 2, \dots, k\}$ , then  $C_i \in \mathcal{C}(M)$  and  $C_j = C_j^1 \cup C_j^2$  where  $C_j^1, C_j^2 \in \mathcal{C}(M)$  and  $C_j^1 \cap C_j^2 = \emptyset$ . Therefore

$$E(M) = C_1 \cup C_2 \cup \dots \cup C_i \cup \dots \cup C_j^1 \cup C_j^2 \cup \dots \cup C_l$$

is a disjoint circuit decomposition of  $E(M)$ . Therefore  $M$  is Eulerian. □

Applying Proposition 4.2 and 4.3 for  $p = 2$ , we obtain the following result of Raghunthan et al. [7] for binary matroids.

**Corollary 4.4.** *Let  $M$  be a binary matroid and  $a, b \in E$ . Then  $M$  is Eulerian if and only if  $M_{a,b}$  is Eulerian.*

Note that, the splitting operation on non-Eulerian  $p$ -matroids may yield Eulerian  $p$ -matroids.

**Example 4.5.** *Consider the vector matroid  $M \cong M[A]$  represented by the matrix  $A$  over field  $GF(5)$ .*

$$\mathbf{A} = \begin{matrix} & 1 & 2 & 3 & 4 & 5 \\ \begin{pmatrix} 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 \end{pmatrix} & & & & & \end{matrix} \quad \mathbf{A}_{3,5} = \begin{matrix} & 1 & 2 & 3 & 4 & 5 \\ \begin{pmatrix} 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \end{pmatrix} & & & & & \end{matrix}$$

*The circuits of  $M$  are  $\{1, 2, 3, 4\}, \{1, 2, 5\}, \{3, 4, 5\}$ . Note that  $M$  is non-Eulerian matroid. For  $a = 3$ ,  $b = 5$  and  $\alpha = 1$ ,  $A_{3,5}$  represents the splitting matroid  $M_{3,5}$ . Here the matroid  $M_{3,5}$  has only one circuit  $\{1, 2, 3, 4, 5\}$ . Therefore it is Eulerian matroid over  $GF(5)$ .*

In the following result, we provide a sufficient condition for the splitting operation on non-Eulerian  $p$ -matroids to yield Eulerian  $p$ -matroids.

**Proposition 4.6.** *Let  $M$  be non-Eulerian  $p$ -matroid on ground set  $E$ . Then  $E$  can be written as follows:*

$$E = C_1 \cup C_2 \cup \dots \cup C_k \cup I$$

*where  $C_i \in \mathcal{C}(M)$ ,  $I \in \mathcal{I}(M)$ ,  $C_i \cap C_j = \emptyset$  when  $i \neq j \in \{1, 2, \dots, k\}$  and  $C_i \cap I = \emptyset$  for all  $i \in \{1, 2, \dots, k\}$ . Let  $\{a, b\} \subset E$ . If there is a circuit  $C_i$  such that  $C_i \cup I \in \mathcal{C}_1$ , then  $M_{a,b}$  is Eulerian Matroid.*

An Eulerian  $p$ -matroid  $M$  on  $E$  can be transformed by repeated applications of splitting operations into a matroid in which  $E$  is a circuit, which is observed in the following example.

**Example 4.7.** *The matrix  $A$  gives  $GF(3)$  representation of the Eulerian matroid  $S(5, 6, 12)$ . Let  $M$  denote the vector matroid of  $A$ . Consider the following sequence of the splitting operations:  $M^1 = M_{1,6}$ ,  $M^2 = M_{1,4}^1$ ,  $M^3 = M_{1,3}^2$ ,  $M^4 = M_{3,9}^3$ , and  $M^5 = M_{3,11}^4$ .*

$$A = \begin{pmatrix} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 2 & 2 & 2 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 2 & 2 & 2 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 2 & 1 & 0 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 2 & 2 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 1 & 0 & 0 \end{pmatrix}$$

The matroid  $M^5$  is Eulerian in which  $E(M)$  forms a circuit. We observe that each matroid  $M$  and  $M^k, k \in \{1, 2, 3, 4\}$ , have a  $p$ -decomposition. Therefore by Proposition 4.2 the matroids  $M^k, k \in \{1, 2, \dots, 5\}$  are Eulerian.

In the light of Example 4.5, the Theorem 1.1 by Raghunathan et al. [7] does not hold, in general, for  $p$ -matroids,  $p > 2$ .

For  $p > 2$ , it remains open to prove or disprove the existence of a sequence of splitting operations on Eulerian  $p$ -matroid  $M$  which transforms the ground set  $E(M)$  into a circuit. The following algorithm transforms  $E(M)$  into a circuit, if a sequence of splitting operations, described earlier, exists; otherwise,  $E(M)$  can not be transformed into a circuit.

**Algorithm:**

Let  $M$  be Eulerian  $p$ -matroid,

**Step 1 :** Find all the circuits of  $M$ .

**Step 2 :** List all the circuit decompositions of  $M$  as  $\{\mathcal{D}_1, \mathcal{D}_2, \dots, \mathcal{D}_k\}$ .

**Step 3 :** If there is a pair  $a, b \in E(M)$  such that  $\mathcal{D}_i$  is a  $p$ -decomposition of  $M$ , and  $|\mathcal{D}_i| > 1$ , for some  $i \in [k]$ , then the splitting matroid  $M_{a,b}$  is Eulerian by Proposition 4.2. Replace  $M$  by  $M_{a,b}$  and go to **Step 1**.

If no such pair  $a, b \in E(M)$  exists, then go to **Step 4**.

**Step 4 :** Then  $E(M)$  is the circuit of  $M$  or  $E(M)$  can not be transformed into a circuit.

## Acknowledgment

This paper is dedicated to the memory of T.T. Raghunathan who recently passed away on December 18, 2021.

### References

- [1] Y. M. Borse, S. B. Dhotre, On connected splitting matroids, *Southeast Asian Bull. Math.* **34** (2010) 807-811.
- [2] H. Fleischner, Eulerian graphs and related topics, Part 1, Vol 1, North Holland, Amsterdam (1990).
- [3] A. D. Mills, On the cocircuits of a splitting matroid, *Ars Combinatoria* **89** (2008) pp.243-253.
- [4] P. P. Malavadkar, M.M. Shikare and S. B. Dhotre, A characterization of n-connected splitting matroids, *Asian-European J. Math.* **7** (4)(2014) 1-7.
- [5] P. P. Malavadkar, A study of splitting operations for binary matroids and related results, Ph.D. Thesis, Savitribai Phule Pune University (2015).
- [6] J. G. Oxley, Matroid theory, *Oxford University Press*, Oxford (1992).
- [7] T. T. Raghunathan, M. M. Shikare, B. N. Waphare, Splitting in a binary matroid, *Discrete Mathematics* **184** (1998) 267-271.
- [8] M. M. Shikare, G. Azadi, Determination of bases of a splitting matroid, *European J. Combin.* **24** (2003) 45-52.
- [9] M. M. Shikare, Splitting Lemma for binary matroids, *Southeast Asian Bull. Math.* **32** (2008) 151-159.
- [10] W. T. Tutte, Lectures on matroids, *J. Res. Nat. Bur. Standards* **B69** (1965) 1-47.
- [11] W. T. Tutte, Connectivity in matroids, *Canad. J. Math.* **18** (1966) 1301-1324.
- [12] D. K. Wagner, Bipartite and Eulerian minors, *European J. Combin.* **74**(2018) 1-10.
- [13] D. J. A. Welsh, Euler and bipartite matroids, *J. Combin. Theory* **6** (1969) 375-377.
- [14] Yaokun Wu, Even poset and a parity result for binary linear code, *Linear Algebra and its Appl.* **418**(2006) 591-594.