Subgroup Analysis for Longitudinal data via Semiparametric Additive Mixed Effect Model

Xiaolin Bo, Weiping Zhang Department of Statistics and Finance, University of Science and Technology of China

Abstract

In this paper, we propose a general subgroup analysis framework based on semiparametric additive mixed effect models in longitudinal analysis, which can identify subgroups on each covariate and estimate the corresponding regression functions simultaneously. In addition, the proposed procedure is applicable for both balanced and unbalanced longitudinal data. A backfitting combined with k-means algorithm is developed to estimate each semiparametric additive component across subgroups and detect subgroup structure on each covariate respectively. The actual number of groups is estimated by minimizing a Bayesian information criteria. The numerical studies demonstrate the efficacy and accuracy of the proposed procedure in identifying the subgroups and estimating the regression functions. In addition, we illustrate the usefulness of our method with an application to PBC data and provide a meaningful partition of the population.

Keywords: subgroup identification, additive model, mixed effect, backfitting

1 Introduction

Subgroup analysis has emerged as important drug development tool with the demand of precision medicine emerging and rising (Foster et al., 2011; Schwalbe et al., 2017). Consequently, there is an increasing need to distinguish homogeneous subgroups of individuals, detect the diverse patterns in the subpopulations, model the relationships between the response variable and predictors differently across the subpopulations and make the best personalized predictions for individuals belonging to different subgroups. Thus, virous statistical methods have been developed for subgroup identification in longitudinal data, such as decision trees, mixture models, regularization methods and change point methods etc.

Since the seminal book on classification and regression trees (CART) by (Breiman et al., 1984), tree-based methods become widely used for subgroup identification. In general, a tree recursively partitions the subjects into binary nodes until specific stopping rule is met and in this way subgroups are yielded. Sela and Simonoff (2012) developed the RE-EM Tree procedure, fitting a mixed effect model by regarding the fixed effect as a regression tree and iteratively, estimating the random effect and the fixed effect like EM algorithm rather than traditional maximum likelihood estimation. Loh and Zheng (2013) extended GUIDE algorithm to longitudinal and multiresponse data, by means of treating each longitudinal data series as a curve and using chi-squared tests of the residual curve patterns to select a variable to split each node of the tree. Model-based recursive partitioning method developed by Zeileis et al. (2008) and Seibold et al. (2016), fits a parametric model in each node, with splitting variable chosen by independence tests and parameter values estimated as solutions to the score equations, which is the partial derivatives of the log-likelihood. Wei et al. (2020) proposed interaction tree for longitudinal trajectories, combining mixed effect models with regression splines to model the nonlinear progression patterns among repeated measures, and identify subgroups with differential treatment effects for two-sample comparisons in longitudinal randomized clinical trials.

Furthermore, the growth mixture modeling methods (Fraley and Raftery, 2002; Song et al., 2007; Jung and Wickrama, 2008), have been widely utilized to identify and predict latent subpopulations for longitudinal data. For example, McNicholas and Murphy (2010) and McNicholas (2016) developed a model-based clustering method (called longclust) method for balanced longitudinal data. Shen and Qu (2020) proposed a structured mixed-effects approach for longitudinal data to model subgroup distribution and identify subgroup membership simultaneously. In general, such approaches require to know the underlying distribution of data and the number of mixture components in advance. Comparably, clustering regression curves can be done to find subgroups. Abraham et al. (2003) proposed a clustering procedure which consists of two stages: fitting the functional data by B-splines and partitioning the estimated model coefficients using a k-means algorithm. They also shown their procedure possessing strong consistency. Ma et al. (2006) and Coffey et al. (2014) adopted the smoothing spline and penalized spline approximations under the mixed effect framework respectively to model time-course gene expression data and detect subpopulations. In addition, some distance-based clustering methods have been proposed to cluster the

trajectories of longitudinal data, for instance, Genolini and Falissard (2010) combined generalized Fréchet distance with k-means to achieve this goal. Lv et al. (2020) proposed a two-step classification algorithm which compares the L_2 -distances between kernel estimates of nonparametric functions to estimate parameters of group memberships and the number of subgroups simultaneously. Zhang et al. (2019a) proposed a quantile-regression-based clustering method for panel data by using a similar idea of k-means clustering to identify subgroups with heterogeneous slopes at a single quantile level or across multiple quantiles.

There has also been a line of work on regularization methods. For example, Ma and Huang (2017) proposed a concave pairwise fusion learning method to identify subgroups whose heterogeneity is driven by unobserved latent factors and thus can be represented by subject-specific intercepts. Zhang et al. (2019b) employed penalized median regression to detect subgroups automatically and achieve robustness against outliers and heteroscedasticity in random errors. Lu et al. (2021) proposed a subgroup identification method based on concave fusion penalization and median regression for longitudinal data with dropouts. However, these aforementioned methods mainly focused on dividing the individuals into several groups according to the intercept or the whole list of all regression coefficients, not on detecting subgroups on each covariate separately. For this purpose, Li et al. (2019) proposed an estimation procedure combining the likelihood method and the change point detection with the binary segmentation algorithm under partial linear models.

There is a clear need to relax the parametric assumption posed in Li et al. (2019) as model misspecification may result in biased estimation. An attractive approach is the semiparametric additive mixed effect model, which retains the flexibility of the nonparametric model but avoids the curse of dimensionality of a fully nonparametric model. In this paper, we propose a very general framework to recognize the subgroup structure on each covariate and estimate the regression functions in each subgroup simultaneously, based on semiparametric additive mixed effect model. Specifically, using the densely observed data for each individual, we give a initial estimates of the parametric part and additive components in the model, pretending that there is no subgroups in the population. Then we adopt the backfitting and k-means algorithm to estimate each semiparametric additive component across subgroups and detect subgroup structure on each covariate. The utilization of mixed effect enables us capture the within-subject correlation among longitudinal measurements, while additive nonparametric components are helpful for us to characterize the nonlinear relationships between covariates and the response.

The major contributions in this paper can be outlined as follows. First, we propose a very general framework for identifying subgroups on multiple covariates, which possesses the flexibility and interpretability of semiparametric additive mixed effect model. In addition, the proposed method can detect subgroups on each covariate, consequently it could make us more clear about which covariate contributes to the existence of subgroups among population. Second, the proposed model is applicable for both balanced and unbalanced longitudinal data. Third, the proposed procedure holds some theoretical properties and computationally simplicity, our simulation studies also indicate the fine efficiency of our approach.

The rest of this paper is organized as follows. Section 2 elaborates the proposed subgroup identification methods. Section 3 discusses the asymptotic properties. Section 4 conducts simulation studies to evaluate the proposed approach. Application to the PBC dataset is presented in Section 5. Finally, Section 6 concludes this paper by summarizing the main findings and outlining future research.

2 Methods

2.1 The model

Consider the following longitudinal dataset $(Y_{it}, X_{it}, Z_{it}, S_i)$, $i = 1, ..., n, t = 1, ..., n_i$ collected from n independent individuals, where Y_{it} is the response variable for the ith subject at the ith follow-up of total ith measurements, ith i

$$Y_{it} = S_i'\beta + f_1(X_{it1}) + \ldots + f_p(X_{itp}) + Z_{it}'b_i + \varepsilon_{it}, \quad i = 1, \ldots, n, \quad t = 1, \ldots, n_i,$$
 (1)

where

$$f_{j}(X_{itj}) = \begin{cases} f_{1j}(X_{itj}), & \text{when } i \in \Omega_{1,j} \\ f_{2j}(X_{itj}), & \text{when } i \in \Omega_{2,j} \\ \vdots & \vdots \\ f_{m_{j}j}(X_{itj}), & \text{when } i \in \Omega_{m_{j},j}. \end{cases}$$

$$(2)$$

with $\{\Omega_{kj}: 1 \leq k \leq m_j\}, j=1,\ldots,p$ representing an unknown partition of the subject index set $\{i: 1 \leq i \leq n\}$ on covariate X_j . Note that the number of subgroups m_j is unknown either. The traditional random-effects model assumes that the random effects follow a certain distribution, for instance, a normal distribution, and focuses on the variance component estimation of the random effects. However, we do not impose any distribution assumption on b_i , but instead assume that the random effects have mean zero and variance $\sigma_b^2 > 0$. In addition, ε_{it} is the random error with zero mean and variance σ_ε^2 . Under this model, the trajectory of the *i*th subject over time is represented by the linear regression part, the group-specific unknown additive functions and the subject-specific random effect. Without loss of generality, we assume that all $E[f_j(\cdot)] = 0$ to prevent identifiability problems.

Note that model (1) is very flexible, it retains the flexibility of the nonparametric model but avoids the need to model a fully nonparametric model. The linear part possesses the ease of interpretability on the baseline covariates, the random effect term represents the heterogeneity between individuals, and the additive nonparametric part reflect the nonlinear relationships between response and each covariate. The proposed model (1) also allows subgroups exist on each covariate, thus for the jth predictor, we may have m_j different nonparametric additive components and thus form m_j different subgroups. The subjects

with the same f_j share a similar nonparametric dependence between the response and the jth predictor. It can be seen that the detected subgroups for different j may have completer, partial, or zero overlap, so there may have at least $\max_j m_j$ subgroups and at most $\sum_{j=1}^p m_j$ subgroups, where each subgroup has a distinct set of additive components. In the following, we use |A| to denote the cardinality of a set A.

2.2 Subgroup identification algorithm

2.2.1 Initial estimates

In this section, we describe the procedure for initial estimates of the additive components. Let $\mathbf{Y}_i = (Y_{i1}, \dots, Y_{in_i})'$, $\mathbf{Z}_i = (Z_{i1}, \dots, Z_{in_i})'$, $\boldsymbol{\varepsilon}_i = (\varepsilon_{i1}, \dots, \varepsilon_{in_i})'$, and let $\mathbf{S}_i = \mathbf{1}_{n_i} S_i'$, where $\mathbf{1}_{n_i}$ is a $n_i \times 1$ vector with entries equal to one. Based on B-spline approximation, for each subject i, model (1) can be written in matrix notation as

$$Y_i \approx S_i \beta + B_{i1} \gamma_{i1} + \ldots + B_{ip} \gamma_{ip} + Z_i b_i + \varepsilon_i, \quad i = 1, \ldots, n,$$
 (3)

where $\mathbf{B}_{ij} = (B_{i1j}, \dots, B_{in_ij})'$ are B-spline basis functions, for $j = 1, \dots, p$,

$$\gamma_{ij} = \begin{cases}
\gamma_{1j}, & \text{when } i \in \Omega_{1,j} \\
\gamma_{2j}, & \text{when } i \in \Omega_{2,j} \\
\vdots & \vdots \\
\gamma_{m_j j}, & \text{when } i \in \Omega_{m_j,j}.
\end{cases}$$
(4)

In addition, let Σ_i and V_i denote the true and assumed working covariance of Y_i , where $V_i = A_i^{1/2} R_i A_i^{1/2}$. Here, A_i is a $n_i \times n_i$ diagonal matrix of the marginal variance of Y_i , and R_i is the corresponding working correlation matrix. In this situation, V_i is assumed to depend on a nuisance finite dimensional parameter vector $\tilde{\boldsymbol{\theta}}$.

In order to identify the subgroups on each covariate, we first pretend that all individuals come from the same group, or that there are no subgroups in the population, and obtain an initial estimate by fitting model (3). The initial estimates can be obtained by approximating the additive components $f_1(\cdot), \ldots, f_p(\cdot)$ through a spline basis expansion and then employing an extension of the standard GEE. Since smoothing spline has higher computational cost, here we implement B-spline, which not only maintains comparable performance in estimating, but also reduces computation complexity.

The initial estimates of nonparametric functions, $\widehat{f}_1^{(0)}, \ldots, \widehat{f}_p^{(0)}$, or to be more specific, the initial estimates of B-spline coefficients $\widehat{\gamma}_1^{(0)}, \ldots, \widehat{\gamma}_p^{(0)}$ will be input into the subgroup identification procedure in the next subsection. Huang et al. (2007) has shown that the estimate of B-spline coefficient is efficient if the covariance structure is correctly specified and it is still consistent and asymptotically normal even if the covariance structure is misspecified. The explicit derivation and proof can refer to Huang et al. (2007), so we will not discuss it in detail here.

2.2.2 Backfitting and subgroup pursuit

Given model (1), we can employ backfitting algorithm to fit additive models by iteratively solving

$$E[Y - S\beta - \sum_{k \neq j} f_k(X_k) | X_j] = f_j(X_j), j = 1, \dots, p,$$
(5)

and at each stage we replace the conditional expectation of the partial residuals with a univariate smoother. We first set

$$W_{itj} = Y_{it} - S_i \widehat{\beta} - \sum_{k \neq j} \widehat{f}_k^{(0)}(X_{itk}), j = 1, \dots, p,$$
 (6)

thus the problem can be transformed to univariate nonparametric regression. Various methods have been proposed to solve univariate nonparametric regression issue, for instance, kernel method, local polynomial regression and splines, and we are going to utilize the B-spline in each stage, since it has nice properties of efficiency and flexibility (De Boor and De Boor, 1978; Lorentz and DeVore, 1993).

For j = 1, ..., p, we aim to divide n subjects into m_j groups such that subjects with homogeneity being partitioned into the same group. We first assume that each subject has its own regression function between X_j and W_j . In other words, hypothesizing that for each subject i, we have

$$W_{itj} = m_{ij}(X_{itj}) + Z'_{it}b_i + \varepsilon_{itj}, \quad t = 1, \dots, n_i, \tag{7}$$

and then cluster functions $\{m_{ij}(\cdot), i = 1, ..., n\}$ into m_j groups.

To fit the nonparametric univariate function in (7), let $x \in [a, b]$ and let $(\xi_0 =)a < \xi_1 < \xi_2 < \ldots < \xi_{J_n} < b (= \xi_{J_n+1})$ be a subdivision by J_n distinct points on [a, b], and the spline function s(x) is a polynomial of degree d on any interval $[\xi_{i-1}, \xi_i]$ which has d-1 continuous derivations on (a, b). On the basis of B-spline, we can write a spline as $s(x, \gamma) = \sum_{l=1}^{J_n+d+1} \gamma_l B_l(x)$, where $\gamma = (\gamma_1, \ldots, \gamma_{J_n+d+1})'$ is the vector of spline coefficients. Noting that $\mathbf{W}_{ij} = (W_{i1j}, \ldots, W_{in_ij})'$, and $\mathbf{B}_{ij} = \{B_l(x_{itj})\}_{t=1,\ldots,J_n+d+1}^{l=1,\ldots,J_n+d+1}$, also supposing that $\mathbf{B}'_{ij}\mathbf{B}_{ij}$ is non-singular, the spline coefficients are estimated by

$$\widehat{\gamma}_{ij} = \underset{\gamma_{ij}}{\operatorname{argmin}} \frac{1}{n_i} \sum_{t=1}^{n_i} (W_{itj} - s(x_{itj}, \gamma_{ij}))^2 = [\mathbf{B}'_{ij} \mathbf{V}_i \mathbf{B}_{ij}]^{-1} \mathbf{B}'_{ij} \mathbf{V}_i \mathbf{W}_{ij}. \tag{8}$$

Thanks to the fine properties of B-spline that the n functions we estimate share the same degree and knots, as well the same basis functions $(B_1, \ldots, B_{J_n+d+1})$, each coordinate γ_{ij} has the same meaning for each function $m_{ij}, i = 1, \ldots, n$. Thus the set of functions $\{m_{1j}(\cdot), \ldots, m_{nj}(\cdot)\}$ is summarized by $\{\gamma_{1j}, \ldots, \gamma_{nj}\}$, a set of vectors of \mathbb{R}^{J_n+d+1} , and we just need to partition their coefficients γ_{ij} (Abraham et al., 2003).

For the set $\{\gamma_{1j}, \ldots, \gamma_{nj}\}$, we can easily establish the $n \times n$ distance matrix, most of time based on Euclidean distance. Numerous methods have been proposed to deal with this kind of clustering problem based on dissimilarity measure, such as k-means (MacQueen

et al., 1967), fuzzy clustering (Bellman et al., 1966; Ruspini, 1969), hierarchical clustering (Johnson, 1967), model-based clustering (Fraley and Raftery, 2002). In the literature, k-means is one of the most popular clustering methods to make objects within clusters mostly homogeneous and objects between clusters mostly heterogeneous by minimizing the objective function

$$\sum_{s=1}^{k} \sum_{i \in \Omega_s} \|z_i - \mu_s\|_2^2, \tag{9}$$

where μ_s is the center of cluster Ω_s and z_i is the *i*th object.

After partitioning $\{i: 1 \leq i \leq n\}$ into \hat{m}_j subgroups on X_j , we conduct generalized least squares to the subjects who belong to the group $\widehat{\Omega}_{k,j}, k = 1, \ldots, \hat{m}_j$ to obtain $\widehat{\gamma}_{kj}^{(r)}$ in the rth iteration by fitting

$$W_{ij} \approx B_{ij}\gamma_{kj} + Z_i b_i + \varepsilon_i, \quad i \in \widehat{\Omega}_{k,j}.$$
 (10)

We repeat this backfitting procedure for $j=1,\ldots,p$ until the estimated subgroups do not change anymore, and thus the final identified subgroup structure yields. The final estimator of the additive components are denoted by $\widehat{f}_1,\ldots,\widehat{f}_p$, and the final estimated B-spline coefficients are denoted by $\widehat{\gamma}_1,\ldots,\widehat{\gamma}_p$.

2.2.3 Determining the number of clusters

Another concerning issue is determining the number of clusters m_j , as k-means procedure requires to pre-specify the number of clusters m_j . In general, model-selection criteria such as Akaike Information Criterion or Bayesian Information Criterion (Schwarz, 1978) can be used to decide m_j to avoid over-fitting.

AIC and BIC are both methods of assessing model fit penalized for the number of estimated parameters, but they could give different results for estimating the number of clusters in a dataset. In cluster analysis, BIC tends to be preferred to AIC in estimating the number of clusters because it uses a larger penalty term and hence can recommend fewer clusters and it's mathematical formulation is more meaningful in this context (Pelleg and Moore, 2000), whereas AIC is more general. BIC can be written as

$$BIC(k) = -2log(\hat{L}) + log(n)k, \tag{11}$$

where $log(\hat{L})$ is the log-likelihood of the model, k is the total number of parameters and n is the sample size. An attractive property of the BIC is its consistency: the BIC selects the correct model with a probability that goes to 1 as n grows large (Zhang et al., 2010; Bai et al., 2018). Let \hat{m}_j be the minimizer of BIC for choosing m_j , we show that $\hat{m}_j \stackrel{P}{\rightarrow} m_j$ in the next section. In summary, the main steps of our subgroup identification algorithm is summarized in Algorithm 1.

Algorithm 1 Subgroup identification based on backfitting

Input:

 Y_{it} : the tth observation of the ith individual of response variable;

 X_{it} , the tth observation of the ith individual of time dependent variables;

 Z_{it} , the tth observation of the ith individual of covariates associated with random effects; S_i , baseline covariates of the ith individual;

Output:

```
group index \widehat{\Omega}_{1j}, \ldots, \widehat{\Omega}_{m_ij}, j = 1, \ldots, p;
      estimated coefficient \widehat{\beta} and estimated function \widehat{f}_1,\ldots,\widehat{f}_p;
  1: Initialize: fit Y_{it} = S'_i \beta + f_1(X_{it1}) + \ldots + f_p(X_{itp}) + Z'_{it} b_i + \varepsilon_{it}, i = 1, \ldots, n, t = 1, \ldots, n_i;
 2: return initial estimate \widehat{f}_1^{(0)}, \dots, \widehat{f}_p^{(0)}, \widehat{\beta};
  3: for j = 1, ..., p do
          set W_{itj} = Y_{it} - S_i \widehat{\beta} - \sum_{k \neq j} \widehat{f}_k(X_{itk});
  4:
           for i = 1, \ldots, n do
  5:
               solve univariate nonparametric regression W_{itj} = m_{ij}(X_{itj}) + Z'_{it}b_i + \varepsilon_{it} with B-spline;
  6:
               return B-spline coefficients \hat{\gamma}_{ij};
  7:
  8:
           end for
           clustering \{\widehat{\gamma}_{1j}, \dots, \widehat{\gamma}_{nj}\} with k-means;
 9:
           partition \{i \in 1, ..., n\} to group \widehat{\Omega}_{1,j}, ..., \widehat{\Omega}_{\widehat{m}_i,j};
10:
           update \widehat{f}_j = \widehat{f}_{1j}I(i \in \widehat{\Omega}_{1,j}) + \widehat{f}_{2j}I(i \in \widehat{\Omega}_{2,j}) + \ldots + \widehat{f}_{\widehat{m}_i,j}I(i \in \widehat{\Omega}_{\widehat{m}_i,j});
11:
13: repeat iteration line (3-12) until subgroups \{\widehat{\Omega}_{1j}, \dots, \widehat{\Omega}_{\widehat{m}_j j}, j = 1, \dots, p\} converge;
14: return subgroup index \{\widehat{\Omega}_{1j}, \dots, \widehat{\Omega}_{\widehat{m}_j j}, j = 1, \dots, p\} and estimated \widehat{f}_1, \dots, \widehat{f}_p, \widehat{\beta}.
```

3 Asymptotic properties

In this section, we establish the asymptotic properties of the proposed estimator. Let $L_2(\mathcal{X})$ be the space of all square integrable functions on $\mathcal{X} = [0,1]$, and $||f||_2^2 = \int_0^1 f(x)^2 dx$ for any $f \in L_2(\mathcal{X})$. Denote $||f||^2 = E[f(X)^2]$ and $||f||_n^2 = \frac{1}{n} \sum_{i=1}^n f(X_i)^2$ as the theoretical and empirical norms respectively, where X_i is a random sample from \mathcal{X} . For any set \mathcal{G} , $|\mathcal{G}|$ represents the cardinal of \mathcal{G} . For unbalanced dataset, we define $n_0 = \min_i \{n_i\}$, and J_n is the number of internal knots. And several regularity conditions are required to establish the asymptotic properties.

(A1) The function $f_i(\cdot) \in C^r[0,1] (i=1,\ldots,n)$ for some $r \geq 1$.

(A2) Let $a = \xi_0 < \xi_1 < \dots < \xi_{J_n} < \xi_{J_{n+1}} = b$ be a partition of [a, b] into $J_n + 1$ subintervals, the knot sequences $\{\xi_l\}_{l=0}^{J_n+1}$ have bounded mesh ratio, which means for some constant C

$$\frac{\max_{0 \le l \le J_n} |\xi_{l+1} - \xi_l|}{\min_{0 \le l \le J_n} |\xi_{l+1} - \xi_l|} \le C.$$

(A3) The design points $\{x_{it}\}(i=1,\ldots,n;t=1,\ldots,T_i)$ follow an absolutely continuous density function g_X , and there exist constants a_1 and a_2 such that $0 < a_1 \le \min_{x \in \mathcal{X}} g_X(x) \le$

 $\max_{x \in \mathcal{X}} g_X(x) \le a_2 < \infty.$

(A4) Assume that $N_g = O(N)$, where $N_{gj} = \sum_{i \in \Omega_{g,j}} n_i$ for $g = 1, ..., m_j$, and $N_{0j} = min(N_{1j}, ..., N_{m_j j}), N = \sum_{i=1}^n n_i$.

(A5) The eigenvalues of $\Sigma = \text{diag}(\Sigma_1, \dots, \Sigma_n)$ and $V = \text{diag}(V_1, \dots, V_n)$ are bounded away from zero to infinity.

Assumptions(A1)-(A3) are standard conditions for the nonparametric B-spline smoothing functions, and (A4) indicates that we require the cluster size to grow as the sample size increases.

Theorem 1. Under Assumption (A1)-(A5), as $n \to \infty$, and given a sufficiently large N_{0j} such that $J_n = O(N_{0j}^{\varsigma})$ with $0 < \varsigma < 1$, then the oracle estimator $\tilde{\gamma}_j$ satisfying

$$\|\tilde{\gamma}_j - \gamma_j\|_2^2 = O_p(J_n/N_{0j} + J_n^{-2r}), j = 1, \dots, p,$$

where $\tilde{\gamma}_i$ denotes estimated spline coefficients when the true subgroup membership is known.

Remark 1. The result of Theorem 1 implies that the convergence rate of the oracle approximation $\tilde{\gamma}_j$, $j=1,\ldots,p$ is faster than the B-spline estimator $\hat{\gamma}_j$. The convergence rate of oracle estimator assures that when prior knowledge on the true subgroup memberships is known, more information from each cluster with sufficient number of repeated measurements can be used. And the proof of Theorem 1 is similar to the proof of Lemma 4, which we will present in the Appendix.

Theorem 2. Under Assumption (A1)-(A5), as $n \to \infty$, and given a sufficiently large n_0 such that $J_n = O(n_0^{\varsigma})$ with $0 < \varsigma < 1$,, then the estimated additive components \hat{f}_j satisfying

$$\|\widehat{f}_j - f_j\|_2^2 = O_p(J_n/n_0 + J_n^{-2r}).$$

Remark 2. Theorem 2 shows that the convergence rate of the proposed additive estimator \hat{f}_j is of the same order as B-spline coefficients estimator $\hat{\gamma}_{ij}^{(r)}$ in backfitting procedure. Theorem 2 holds given a sufficiently large number of repeated measurements, however in practice, it does not have to be such large, in simulations if n_i is bigger than 7 or 8, we will have a nice result, more than 10 times repeated measurements would be better. The proof of Theorem 2 is given in the Appendix.

Theorem 3. Under the same conditions in Theorem 2, as $n_0 \to \infty$ and $\|\widehat{\gamma}_{ij} - \gamma_{ij}\|_2^2 \to 0$, we have

$$P(\widehat{\Omega}_i = \Omega_i) \to 1,$$

where $\widehat{\Omega}_j = \{\widehat{\Omega}_{1,j}, \dots, \widehat{\Omega}_{m_j,j}\}$ is the estimated subgrouping membership on covariate X_j , $\Omega_j = \{\Omega_{1,j}, \dots, \Omega_{m_j,j}\}$ is the true subgrouping membership, and $\widehat{\gamma}_{ij}$ is the estimate of γ_{ij} .

Remark 3. Theorem 3 indicates that when there is a sufficient number of repeated measurements for each subject, the proposed method can identify the true subgroup with probability tending to 1.

4 Numerical studies

In this section, we conduct Monte Carlo simulations on several examples to investigate the performance of the proposed subgroup identification procedure, and also compare the proposed method with some existing methods. Balanced and unbalanced longitudinal datasets are both considered, the situations of no subgroup (only one group), two subgroups are generated in different settings. Three cases are set up with the true additive components to be the combination of linear functions, the combination of nonlinear functions, and also the combination of linear functions and nonlinear function. For each simulation setting, we conduct the experiment for sample size n = 30, 50, 100, 200 and repeat 100 times for all cases, as well the working correlation matrix is chosen to be first-order autoregressive and exchangeable structure respectively (AR(0.3), AR(0.5), EX(0.3), EX(0.5)).

As for the evaluation of subgroup identification, we consider employing the accuracy percentage (%), Rand Index (RI) (Rand, 1971) and Normalized Mutual Information (NMI) (Vinh et al., 2010) to assess the identification performance, as they can measure how close the estimated grouping structure approaches the true structure. These three values are ranging between 0 and 1, with larger value indicating a higher score of similarity between the two groups. The accuracy percentage(%) is defined as the proportion of subjects that are correctly identified. And the definition of Rand Index is

$$RI = \frac{TP + TN}{TP + FP + FN + TN},$$

where TP (true positive) is the number of pairs fo subjects that are from different subgroups and assigned to different clusters, TN (true negative) is the number of pairs that are in the same subgroup and assigned to the cluster, FP (false positive) means the number of pairs that in the same subgroup but assigned to different clusters, FN (false positive) denotes the numbers of pairs which are from different subgroups but assigned to the same cluster. Assuming that $A = \{A_1, A_2, \ldots\}$ and $B = \{B_1, B_2, \ldots\}$ are two sets of disjoint group of $\{1, 2, \ldots, n\}$, the NMI is defined as

$$NMI(A, B) = \frac{2I(A, B)}{H(A) + H(B)},$$

where

$$I(A, B) = \sum_{i,j} \frac{|A_i \cap B_j|}{n} \log \left(\frac{n |A_i \cap B_j|}{|A_i||B_j|} \right)$$

is the mutual information between the two groups, and the entropy of A is defined as following

$$H(A) = \sum_{i} \frac{|A_i|}{n} \log \left(\frac{n}{|A_i|}\right).$$

Furthermore, to investigate the estimation accuracy, we calculate the mean squared error (MSE) of the proposed approach and report the ratio of oracle MSE to MSE of our

method. A ratio closer to 1 indicates a better estimation performance. And the following formula is utilized to find corresponding MSE of the estimated \hat{y}_{it} :

$$MSE = \frac{1}{n} \sum_{i=1}^{n} \frac{1}{n_i} \sum_{t=1}^{n_i} (\hat{y}_{it} - y_{it})^2.$$

Case I: In this setting, we consider the situation both f_1 and f_2 are linear functions,

$$y_{it} = f_1(x_{it1}) + f_2(x_{it2}) + Z'_{it}b_i + \varepsilon_{it}, i = 1, \dots, n, t = 1, \dots, n_i$$

where the true subgroup structure is

$$f_1(x) = \begin{cases} 3x - 1.5, & 1 \le i \le n/2 \\ -5x + 2.5, & n/2 < i \le n \end{cases}, f_2(x) = \begin{cases} 1.25x - 0.625, & 1 \le i \le n/2 \\ -6x + 3, & n/2 < i \le n \end{cases}$$

The covariates X_{it1} , X_{it2} are both generated from U(0,1), $Z_{it} \sim N(0,1)$. The vector of random effects $b_i \sim N(0,0.2)$ and ε_{it} are generated from N(0,0.1). We compare our method with two existing approaches, the model-based clustering and classification for longitudinal data method (longclust) (McNicholas, 2016), which is based on gaussian model mixture to detect subgroups, and the k-means for longitudinal data method (kml) (Genolini and Falissard, 2010), which implements k-means clustering based on generalized Fréchet distance to achieve subgrouping. Both longclust approach and kml approach are not suitable for unbalanced data, so in this setting we generate balanced data with $n_i = 15, i = 1, \ldots, n$.

Table 1 shows that the accuracy percentage (%), NMI and Rand Index of the proposed method are equal to 1, which means the proposed method has a excellent performance in the linear setting considering the sample size n=30,50 are relatively small. In addition, for kml method, the accuracy percentage and Rand Index are around 50%, indicating it only correctly identified half of the sample. And the performance of longclust method is quite different under different sample sizes, when the sample size is larger than 100, it has a quite well performance, but for n=30,50, its performance become unsatisfactory, indicating its inapplicability to small sample size. As longclust and kml methods only use the information of response variable and time, and the proposed method includes other covariates and random effect to modeling the trajectories and pursuing subgroups, the proposed method can better characterize the structure of longitudinal data.

Case II: In this setting, we consider the scenario of unbalanced data, and let both f_1 and f_2 to be nonlinear functions,

$$y_{it} = f_1(x_{it1}) + f_2(x_{it2}) + Z'_{it}b_i + \varepsilon_{it}, i = 1, \dots, n, t = 1, \dots, n_i,$$

where the true subgroup structure is

$$f_1(x) = \begin{cases} -1.75 \arctan(5(x - 0.6)) - 0.415, & \text{when } i \text{ is odd} \\ 2.5(1 - ((x - 0.75)/0.8)^2)^4 - 1.363, & \text{when } i \text{ is even} \end{cases}$$

$$f_2 = \begin{cases} 2sin(\pi x) - \frac{\pi}{4}, & 1 \le i \le n \\ 2cos(\pi x), & n/2 < i \le n \end{cases}$$

Table 1: Results for Case I. The accuracy percentage (%), NMI, Rand Index for the proposed method, longclust method and kml method with working correlation matrix to be AR(0.3), AR(0.5), EX(0.3) and EX(0.5). The sample size is chosen to be 30, 50, 100 and 200 respectively. The values in the parentheses are the standard deviations.

	n	proposed method				- longclust	kml
	11	AR(0.3)	AR(0.5)	EX(0.3)	EX(0.5)	Tongerust	KIIII
	30	1	1	1	1	0.5496	0.5112
		(0)	(0)	(0)	(0)	(0.0040)	(0.0120)
	50	1	1	1	1	0.600	0.5206
07		(0)	(0)	(0)	(0)	(0.0030)	(0.0060)
%	100	1	1	1	1	0.9330	0.5397
		(0)	(0)	(0)	(0)	(0.0024)	(0.0030)
	200	1	1	1	1	0.9410	0.5667
		(0)	(0)	(0)	(0)	(0.0025)	(0)
	30	1	1	1	1	0.1016	0.0050
		(0)	(0)	(0)	(0)	(0.0210)	(0.0127)
	50	1	1	1	1	$0.1505^{'}$	0.0191
NIMI		(0)	(0)	(0)	(0)	(0.012)	(0.0048)
NMI	100	1	1	1	1	0.6936	0.0369
		(0)	(0)	(0)	(0)	(0.0080)	(0.0017)
	200	1	1	1	1	0.7067	0.0374
		(0)	(0)	(0)	(0)	(0.0030)	(0)
Rand Index	30	1	1	1	1	0.5000	0.4852
		(0)	(0)	(0)	(0)	(0.0062)	(0.0183)
	50	1	1	1	1	0.5014	0.4912
		(0)	(0)	(0)	(0)	(0.0041)	(0.0117)
	100	1	1	1	1	0.8770	0.4953
		(0)	(0)	(0)	(0)	(0.0024)	(0.0008)
	200	1	1	1	1	0.8916	0.5026
		(0)	(0)	(0)	(0)	(0.0020)	(0.0004)

The covariates X_{it1}, X_{it2} are both generated from $U(0,1), Z_{it} \sim N(0,1)$. The vector of random effects $b_i \sim N(0,0.2)$ and ε_{it} are generated from N(0,0.1). n_i is a random integer generated between 10 and 20. Since the kml method and longclust method are not suitable for this scenario, we only present the simulation result of the proposed approach.

Table 2 shows the simulation results for Case II. For each setup, we present the accuracy percentage on each covariate $(\%_{X_1}, \%_{X_2})$ and total accuracy percentage $(\%_{total})$, as well as the NMI and Rand Index. Generally, when sample size n increases, the performance of subgroup identification and estimation also shows an increasing trend. Each performance evaluation index is close to 1, which reflects the good clustering accuracy of the proposed method. Moreover, the MSE values of the proposed method are comparable to the oracle values, indicating our approach is able to detect subgroups and give a good estimate simultaneously.

Case III: In this setting, we consider the scenario that f_1 and f_3 are nonlinear functions, and f_2 are linear function, and we include variable U as baseline covariate.

$$y_{it} = u_i \beta + f_1(x_{it1}) + f_2(x_{it2}) + f_3(x_{it3}) + b_i + \varepsilon_{it}, i = 1, \dots, n, t = 1, \dots, n_i,$$

where the true subgroup structure is

$$f_1(x) = 2\cos(\pi x), \quad i = 1, \dots, n$$

$$f_2(x) = \begin{cases} 3x - 1.5, & \text{when } i \text{ is odd} \\ -5x + 2.5, & \text{when } i \text{ is even} \end{cases}$$

$$f_3(x) = \begin{cases} -1.75 \times \arctan(5(x - 0.6)) - 0.415, & 1 \le i \le n/2 \\ 2.5 \times (1 - ((x - 0.75)/0.8)^2)^4 - 1.363, & n/2 < i \le n \end{cases}.$$

The covariates $X_{it1}, X_{it2}, X_{it3}$ are generated from $U(0, 1), U_i \sim B(1, 0.5)$, well the vector of random effects $b_i \sim N(0, 0.2)$ and ε_{it} are generated from N(0, 0.1). n_i is a random integer generated between 10 and 20.

Table 3 presents the accuracy percentage (%), NMI and Rand Index of our method when the true model is set with a mixture of linear functions and nonlinear functions, and when there may not exist subgroup. All these indexes are close to 1, demonstrating the good performance of the proposed method. Furthermore, we can see from Table 3 that the MSE values of the proposed method are close to the MSE of oracle estimators. The simulation of this scenario suggests that whether the relationship between the covariates and response variable is nonlinear or linear function, or whether there exist subgroups or not, the proposed approach could successfully identify the subgroup membership and then give consistent estimate of each additive component.

As the preceding examples, the proposed method is not only suitable for balanced data, but also applicable of unbalanced scenario. Furthermore, besides identifying subgroups with high accuracy, the proposed method also gives good estimation of additive components. In a word, these results confirm that the proposed approach has good performances on finding subgroups of heterogeneous trajectories.

Table 2: Results for Case II. The accuracy percentage on X_1 ($\%_{X_1}$), the accuracy percentage on X_2 ($\%_{X_2}$), the accuracy percentage of total population ($\%_{\text{total}}$), NMI, Rand Index and the ratio of oracle MSE to MSE of the proposed method with working correlation matrix to be AR(0.3), AR(0.5), EX(0.3) and EX(0.5). The sample size is chosen to be 30, 50, 100 and 200 respectively. The values in the parentheses are the standard deviations.

	n	$\%_{X_1}$	$\%_{X_2}$	$\%_{total}$	NMI	Rand Index	Ratio
	30	0.9863	0.9924	0.9850	0.9665	0.9859	0.9551
$\mathbf{AD}(0.2)$		(0.0268)	(0.0144)	(0.0269)	(0.0513)	(0.0240)	(0.2054)
AR(0.3)	50	0.9882	0.9932	0.9870	0.9692	0.9868	0.9667
		(0.0148)	(0.0116)	(0.0169)	(0.0419)	(0.0169)	(0.2002)
	100	0.9906	0.9945	0.9905	0.9696	0.9904	0.9744
		(0.0093)	(0.0069)	(0.0098)	(0.0295)	(0.0098)	(0.1544)
	200	0.9913	0.9953	0.9910	0.9711	0.9910	0.9843
		(0.0072)	(0.0067)	(0.0075)	(0.0240)	(0.0074)	(0.0808)
	30	0.9850	0.9920	0.9833	0.9639	0.9837	0.9524
AR(0.5)		(0.0203)	(0.0157)	(0.0209)	(0.0447)	(0.0204)	(0.1986)
AIC(0.5)	50	0.9906	0.9936	0.9884	0.9642	0.9886	0.9612
		(0.0134)	(0.0152)	(0.0173)	(0.0381)	(0.0166)	(0.1343)
	100	0.9912	0.9942	0.9892	0.9675	0.9888	0.9631
		(0.0094)	(0.0099)	(0.0114)	(0.0309)	(0.0118)	(0.1199)
	200	0.9914	0.9946	0.9902	0.9724	0.9895	0.9701
		(0.0063)	(0.0055)	(0.00062)	(0.0219)	(0.0066)	(0.0645)
	30	0.9970	0.9980	0.9970	0.9930	0.9970	0.9985
$\mathrm{EX}(0.3)$		(0.0096)	(0.0079)	(0.0096)	(0.0211)	(0.0096)	(0.0287)
	50	0.9977	0.9987	0.9977	0.9934	0.9977	0.9920
		(0.0046)	(0.0034)	(0.0047)	(0.0142)	(0.0046)	(0.0263)
	100	0.9983	0.9992	0.9983	0.9940	0.9983	0.9927
		(0.0042)	(0.0030)	(0.0042)	(0.0112)	(0.0041)	(0.0231)
	200	0.9991	0.9995	0.9991	0.9976	0.9991	0.9953
		(0.0034)	(0.0021)	(0.0034)	(0.0112)	(0.0034)	(0.0195)
	30	0.9970	0.9983	0.9970	0.9918	0.9970	0.9880
$\mathrm{EX}(0.5)$		(0.0096)	(0.0073)	(0.0096)	(0.0210)	(0.0095)	(0.0417)
121(0.0)	50	0.9975	0.9984	0.9975	0.9934	0.9974	0.9888
		(0.0067)	(0.0054)	(0.0067)	(0.0172)	(0.0067)	(0.0379)
	100	0.9976	0.9988	0.9976	0.9935	0.9977	0.9891
		(0.0037)	(0.0025)	(0.0037)	(0.0118)	(0.0037)	(0.0229)
	200	0.9986	0.9993	0.9986	0.9958	0.9986	0.9931
		(0.0034)	(0.0023)	(0.0034)	(0.0109)	(0.0034)	(0.0203)

Table 3: Results for Case III. The accuracy percentage on X_1 (% $_{X_1}$), the accuracy percentage on X_2 (% $_{X_2}$), the accuracy percentage on X_3 (% $_{X_3}$), the accuracy percentage of total population (% $_{\text{total}}$), NMI, Rand Index and the ratio of oracle MSE to MSE of the proposed method with working correlation matrix to be AR(0.3), AR(0.5), EX(0.3) and EX(0.5). The sample size is chosen to be 30, 50, 100 and 200 respectively. The values in the parentheses are the standard deviations.

	n	$\%_{X_1}$	$\%_{X_2}$	$\%_{X_3}$	$\%_{total}$	NMI	Rand Index	Ratio
	30	1	0.9907	0.9893	0.9880	0.9745	0.9885	0.9210
AD(0.2)		(0)	(0.0158)	(0.0206)	(0.0209)	(0.0435)	(0.0198)	(0.1471)
AR(0.3)	50	1	0.9927	0.9927	0.9924	0.9745	0.9927	0.9285
		(0)	(0.0111)	(0.0145)	(0.0152)	(0.0339)	(0.0142)	(0.1417)
	100	1	0.9933	0.9932	0.9925	0.9789	0.9926	0.9519
		(0)	(0.0089)	(0.0093)	(0.0094)	(0.0281)	(0.0093)	(0.1359)
	200	1	0.9942	0.9934	0.9930	0.9817	0.9930	0.9765
		(0)	(0.0056)	(0.0056)	(0.0058)	(0.0189)	(0.0058)	(0.0906)
	30	1	0.9918	0.9920	0.9833	0.9639	0.9837	0.9524
AR(0.5)		(0)	(0.0145)	(0.0157)	(0.0209)	(0.0447)	(0.0204)	(0.1986)
AIt(0.5)	50	1	0.9940	0.9936	0.9884	0.9642	0.9886	0.9612
		(0)	(0.0121)	(0.0152)	(0.0173)	(0.0381)	(0.0166)	(0.1343)
	100	1	0.9953	0.9942	0.9892	0.9675	0.9888	0.9631
		(0)	(0.0061)	(0.0099)	(0.0114)	(0.0309)	(0.0118)	(0.1199)
	200	1	0.9955	0.9946	0.9902	0.9724	0.9895	0.9701
		(0)	(0.0063)	(0.0055)	(0.00062)	(0.0219)	(0.0066)	(0.0645)
	30	1	0.9986	0.9989	0.9986	0.9930	0.9951	0.9948
EX(0.3)		(0)	(0.0051)	(0.0035)	(0.0051)	(0.0211)	(0.0129)	(0.0231)
$\mathbf{E}\mathbf{A}(0.3)$	50	1	0.9986	0.9992	0.9986	0.9934	0.9957	0.9949
		(0)	(0.0046)	(0.0035)	(0.0047)	(0.0142)	(0.0123)	(0.0196)
	100	1	0.9986	0.9993	0.9986	0.9940	0.9965	0.9955
		(0)	(0.0041)	(0.0035)	(0.0041)	(0.0112)	(0.0101)	(0.0149)
	200	1	0.9993	0.9997	0.9993	0.9976	0.9986	0.9977
		(0)	(0.0025)	(0.0021)	(0.0026)	(0.0112)	(0.0090)	(0.0110)
	30	1	0.9983	0.9987	0.9964	0.9953	0.9969	0.9911
EX(0.5)		(0)	(0.0072)	(0.0065)	(0.074)	(0.0142)	(0.0088)	(0.0371)
L 2 (0.0)	50	1	0.9984	0.9991	0.9983	0.9957	0.9983	0.9939
		(0)	(0.0056)	(0.0033)	(0.0056)	(0.0127)	(0.0055)	(0.0305)
	100	1	0.9985	0.9994	0.9985	0.9961	0.9985	0.9948
		(0)	(0.0042)	(0.0031)	(0.0042)	(0.0096)	(0.0042)	(0.0193)
	200	1	0.9989	0.9994	0.9989	0.9984	0.99989	0.9962
		(0)	(0.0027)	(0.0018)	(0.0027)	(0.0072)	(0.0027)	(0.0103)

5 Application

In this section, we apply the proposed method to primary biliary cirrhosis (PBC) data collected between 1974 and 1984, which is available in R package "joineRML". PBC is a chronic disease characterized by inflammatory destruction of the small bile ducts within the liver, which eventually leads to cirrhosis of the liver, followed by death. Patients often present abnormalities in their blood tests, such as elevated and gradually increased serum bilirubin. Characterizing the patterns of time courses of bilirubin levels is medical interest. A total of 424 patients, referred to Mayo Clinic during that ten-year interval, met eligibility criteria for the randomized placebo-controlled trial of the drug D-penicilamine, 312 formal study participants, and 106 eligible nonenrolled subjects. This dataset contains multiple laboratory results, but only on the first 312 patients, among which 140 had died and the rest were censored and the sex ratio is at least 9:1 (women to men). Some baseline covariates such as age, gender, drug use indicator, were recorded at the beginning of the study. And various biomarkers were collected longitudinally, containing presence of hepatomegaly, spiders, edema and ascites, serum bilirubin, albumin, alkaline, phosphatase, serum glutamic-oxaloacetic transaminase (SGOT), platelets per cubic, prothrombin time and histologic stage of disease, etc. We would not study the problem of missing data in this paper, and the covariate cholesterol includes too much missing data, so we will ignore this covariate. Considering that we will employ splines in the subgrouping procedure, subjects whose repeated measurements less than or equal to 8 times will not be included, thus in our analysis 62 patients with total 687 records will be utilized.

Biomedical research indicates that serum bilirubin concentration is a primary indicator to help assess and track the absence of liver diseases. It is usually normal at diagnosis (0.1-1 mg/dl) but rises with histological disease progression (Talwalkar and Lindor, 2003). Therefore, we focus on modeling the relationship between serum bilirubin and other covariates of interest. And we set log-transformed serum bilirubin level as the response variable, since the original level has positive observed values (Murtaugh et al., 1994).

There are some issues we intend to investigate (i) how the bilirubin levels evolve over time; (ii) how the evolution of bilirubin levels is related to other biomarkers; (iii) whether there exist any subgroups with different additive components among subjects and if there exist subgroups, how they differ from each other. Based on previous researches (Su et al., 2008; Ding and Wang, 2008; Tang et al., 2019), time-independent variables, age(transformed to dummy variable) at registration, gender (male and female), drug type (placebo and D-pencillamine) are included in our model, as well the time-dependent variables, time, SGOT and prothrombin. We analyse the PBC dataset based on the following semiparametric additive mixed effect model:

$$log(bili_{it}) = \alpha + \beta_1 gender_i + \beta_2 drug_i + \beta_3 age_i + f_1(time_{it}) + f_2(log(SGOT_{it})) + f_3(log(prothrombin_{it})) + b_i,$$

where $i = 1, ..., n, t = 1, ..., n_i$. b_i is random effect, and covariate *time* is rescaled to [0, 1], gender is binary gender indicator with 1 for female, drug is binary treatment indicator with 1 for D-pencillamine. After applying the proposed subgroup identification approach to PBC




Figure 1: Plot of fitted f(time) versus time for subjects in Group 1, Group 2 and Group 3.

dataset, 3 subgroups have been found on covariate time, and the BIC value of 3 subgroups reported by our algorithm is obviously less than other situations. Meanwhile, no subgroups have been detected on variables SGOT and prothrombin. There are 34 individuals in $\widehat{\Omega}_{1,1}$, 12 individual in $\widehat{\Omega}_{2,1}$ and 16 individuals in $\widehat{\Omega}_{3,1}$. Figure 1 presents the diverse functions of $\widehat{f}_1(time_{it})$, and it is obvious that these three functions have completely different trends. The time effect for people in subgroup 1 is always negative for $t \in [0,1]$, whereas the time effect for patients in subgroup 2 developed rapidly, indicating the severe condition of their illness, meanwhile the time effect for individuals in subgroup 3 developed towards a worse situation too, but not such severe like patients in subgroup 2.

The differences of the important biomarkers between three subgroups are shown in Figure 2 and Figure 3. We summarize the discrete biomarkers in histograms and the continuous biomarkers in box plots. We also conduct a t-test to compare the differences, and the result is displayed in Table 4. We could see that for some binary symptoms, such as ascites, hepatomegaly, spiders and edema, there are significant differences between these subgroups, demonstrating the heterogeneity of the subgroups detected by the proposed procedure. In addition, for the laboratory indexes collected longitudinally, like logserBilir, albumin, alkaline, SGOT, prothrombin, there are also distinct differences between each subgroup. In particular, the average level of response variable logserBilir for subgroup 2 is obviously higher than the patients in other groups, revealing their severity of illness, and this is consistent with our finding of $\hat{f}_1(time)$ in Figure 1. The estimated model for each group is as follows:

$$\begin{split} \widehat{logbili}_{it} = & 0.833 - 0.662 gender_i + 0.18 drug_i - 0.177 age_i + \widehat{f}_{11}(time_{it}) I(i \in \widehat{\Omega}_{1,1}) \\ & + \widehat{f}_{21}(time_{it}) I(i \in \widehat{\Omega}_{2,1}) + \widehat{f}_{31}(time_{it}) I(i \in \widehat{\Omega}_{3,1}) \\ & + \widehat{f}_{2}(log(SGOT_{it})) + \widehat{f}_{3}(log(prothrombin_{it})) + \widehat{b}_i, \end{split}$$




Figure 2: The frequency of some baseline covariates and biomarkers (ascities, hepatomegaly, spiders, edema, drug, age) for subjects in different subgroups.

where

$$\begin{split} f_{11} &= -0.692B_{0,5} - 0.446B_{1,5} - 0.505B_{2,5} - 0.055B_{3,5} - 0.138B_{4,5} - 0.119B_{5,5}, \\ f_{21} &= 0.141B_{0,5} + 0.365B_{1,5} + 1.402B_{2,5} + 1.152B_{3,5} + 1.861B_{4,5} + 1.324B_{5,5}, \\ f_{31} &= -0.097B_{0,5} - 0.265B_{1,5} + 0.188B_{2,5} + 0.147B_{3,5} + 0.755B_{4,5} + 0.492B_{5,5}, \\ f_{2} &= -1.759B_{0,5} - 1.057B_{1,5} - 0.692B_{2,5} - 0.171B_{3,5} + 0.682B_{4,5} + 0.082B_{5,5}, \\ f_{3} &= -0.072B_{0,5} - 0.034B_{1,5} - 0.111B_{2,5} + 0.318B_{3,5} + 1.150B_{4,5} + 1.275B_{5,5}. \end{split}$$

Throughout our method, the overall subjects have been partitioned to three segmentations. In this procedure, we take multiple covariates into consideration, and reveal the different time effect across three subgroups. In addition, we give the estimates of regression functions of each additive component in the model, which could capture the functional relationship between covariates and the response variable. The results provided by our approach may lead to more accurate subgrouping rule and may be helpful in making personalized medical decision for the patients who suffer from the disease.




Figure 3: Box plots of biomarkers (logserBilir, albumin, alkaline, SGOT, platelets, prothrombin) for subjects in different subgroups.

Table 4: The result of multiple t-test of the important biomarkers for individuals in different groups. μ_1, μ_2 and μ_3 represent the mean of biomarkers of each subgroup. p-value_{ij} is the p-value of multiple hypothesis test between group i and group j.

Biomarkers	μ_1	μ_2	μ_3	$\operatorname{p-value}_{12}$	$\operatorname{p-value}_{13}$	$\operatorname{p-value}_{23}$
ascites	0.0131	0.1374	0.0751	$7.6 imes10^{-8}$	0.0039	0.0139
hepatomegaly	0.2689	0.5115	0.4277	$1.3 imes10^{-6}$	$4.7 imes 10^{-4}$	0.1237
spiders	0.1802	0.5115	0.1387	4.7×10^{-15}	0.26	$8.8 imes10^{-15}$
edema	0.0535	0.1718	0.2312	$2.8 imes10^{-5}$	3.1×10^{-12}	0.055
logserBilir	-0.2152	1.5681	0.2108	$<2 imes10^{-16}$	1.7×10^{-12}	$<2 imes10^{-16}$
albumin	3.6022	3.2886	3.5040	$1.4 imes 10^{-9}$	0.0291	$3.2 imes 10^{-4}$
alkaline	1106.57	1955.98	943.60	$2.8 imes10^{-16}$	0.073	$<2 imes10^{-16}$
SGOT	4.4776	4.9641	4.3792	$<2 imes10^{-16}$	0.03	$<2 imes10^{-16}$
prothrombin	2.3594	2.4185	2.3722	$2.1 imes10^{-12}$	0.087	$2.3 imes10^{-6}$

6 Discussion

This article introduced a novel framework of subgroup identification for longitudinal data, based on semiparametric additive mixed effect model. We aim at finding subgroups on each covariate to reflect the diverse relationship between each covariate with the response variable. It is of great interest to describe the various association between covariates and the response variable, which could reveal how each covariate attributes to the subgroups. Numerical studies indicate that the proposed approach is effective in identifying subgroups and estimating the nonparametric regression functions simultaneously, for both balanced data and unbalanced data.

As continuous response has been considered in this work, one potential future work is to extend the proposed framework to discrete longitudinal outcomes. Second, when the dimension of covariates is high, it would cost more time to adopt the proposed method. So another open issue for future research is extending the framework to high dimensional additive model (Panagiotelis and Smith, 2008; Meier et al., 2009; Fan et al., 2011). In addition, dealing with the dataset where interactions between covariates exist is also a possible direction. These issues are challenging but deserve further exploration.

References

Abraham, C., Cornillon, P.-A., Matzner-Løber, E., and Molinari, N. (2003). Unsupervised curve clustering using b-splines. *Scandinavian journal of statistics*, 30(3):581–595.

Bai, Z., Choi, K. P., and Fujikoshi, Y. (2018). Consistency of aic and bic in estimating the number of significant components in high-dimensional principal component analysis. *The Annals of Statistics*, 46(3):1050–1076.

- Bellman, R., Kalaba, R., and Zadeh, L. (1966). Abstraction and pattern classification. Journal of Mathematical Analysis and Applications, 13(1):1–7.
- Breiman, L., Friedman, J., Stone, C. J., and Olshen, R. A. (1984). *Classification and regression trees.* CRC press.
- Coffey, N., Hinde, J., and Holian, E. (2014). Clustering longitudinal profiles using p-splines and mixed effects models applied to time-course gene expression data. *Computational Statistics & Data Analysis*, 71:14–29.
- De Boor, C. and De Boor, C. (1978). A practical guide to splines, volume 27. springer-verlag New York.
- Ding, J. and Wang, J.-L. (2008). Modeling longitudinal data with nonparametric multiplicative random effects jointly with survival data. *Biometrics*, 64(2):546–556.
- Donoho, D. and Jin, J. (2004). Higher criticism for detecting sparse heterogeneous mixtures. *The Annals of Statistics*, 32(3):962–994.
- Fan, J., Feng, Y., and Song, R. (2011). Nonparametric independence screening in sparse ultra-high-dimensional additive models. *Journal of the American Statistical Association*, 106(494):544–557.
- Foster, J. C., Taylor, J. M., and Ruberg, S. J. (2011). Subgroup identification from randomized clinical trial data. *Statistics in medicine*, 30(24):2867–2880.
- Fraley, C. and Raftery, A. E. (2002). Model-based clustering, discriminant analysis, and density estimation. *Journal of the American statistical Association*, 97(458):611–631.
- Genolini, C. and Falissard, B. (2010). Kml: k-means for longitudinal data. *Computational Statistics*, 25(2):317–328.
- Huang, J. Z., Zhang, L., and Zhou, L. (2007). Efficient estimation in marginal partially linear models for longitudinal/clustered data using splines. *Scandinavian Journal of Statistics*, 34(3):451–477.
- Johnson, S. C. (1967). Hierarchical clustering schemes. *Psychometrika*, 32(3):241–254.
- Jung, T. and Wickrama, K. A. (2008). An introduction to latent class growth analysis and growth mixture modeling. *Social and personality psychology compass*, 2(1):302–317.
- Li, J., Yue, M., and Zhang, W. (2019). Subgroup identification via homogeneity pursuit for dense longitudinal/spatial data. *Statistics in medicine*, 38(17):3256–3271.
- Loh, W.-Y. and Zheng, W. (2013). Regression trees for longitudinal and multiresponse data. *The Annals of Applied Statistics*, pages 495–522.

- Lorentz, G. and DeVore, R. (1993). Constructive approximation, polynomials and splines approximation.
- Lu, W., Qin, G., Zhu, Z., and Tu, D. (2021). Multiply robust subgroup identification for longitudinal data with dropouts via median regression. *Journal of Multivariate Analysis*, 181:104691.
- Lv, Y., Zhu, X., Zhu, Z., and Qu, A. (2020). Nonparametric cluster analysis on multiple outcomes of longitudinal data. *Statistica Sinica*, 30(4):1829–1856.
- Ma, P., Castillo-Davis, C. I., Zhong, W., and Liu, J. S. (2006). A data-driven clustering method for time course gene expression data. *Nucleic acids research*, 34(4):1261–1269.
- Ma, S. and Huang, J. (2017). A concave pairwise fusion approach to subgroup analysis. Journal of the American Statistical Association, 112(517):410–423.
- MacQueen, J. et al. (1967). Some methods for classification and analysis of multivariate observations. In *Proceedings of the fifth Berkeley symposium on mathematical statistics and probability*, volume 1, pages 281–297. Oakland, CA, USA.
- McNicholas, P. D. (2016). Model-based clustering. Journal of Classification, 33(3):331–373.
- McNicholas, P. D. and Murphy, T. B. (2010). Model-based clustering of longitudinal data. *Canadian Journal of Statistics*, 38(1):153–168.
- Meier, L., Van de Geer, S., and Bühlmann, P. (2009). High-dimensional additive modeling. *The Annals of Statistics*, 37(6B):3779–3821.
- Murtaugh, P. A., Dickson, E. R., Van Dam, G. M., Malinchoc, M., Grambsch, P. M., Langworthy, A. L., and Gips, C. H. (1994). Primary biliary cirrhosis: prediction of short-term survival based on repeated patient visits. *Hepatology*, 20(1):126–134.
- Panagiotelis, A. and Smith, M. (2008). Bayesian identification, selection and estimation of semiparametric functions in high-dimensional additive models. *Journal of Econometrics*, 143(2):291–316.
- Pelleg, D. and Moore, A. W. (2000). X-means: Extending k-means with efficient estimation of the number of clusters. In *In Proceedings of the 17th International Conf. on Machine Learning*, pages 727–734. Morgan Kaufmann.
- Rand, W. M. (1971). Objective criteria for the evaluation of clustering methods. *Journal* of the American Statistical association, 66(336):846–850.
- Roussas, G. G. and Ioannides, D. (1987). Moment inequalities for mixing sequences of random variables. *Stochastic Analysis and Applications*, 5(1):60–120.
- Ruspini, E. H. (1969). A new approach to clustering. Information and control, 15(1):22-32.

- Schwalbe, E., Lindsey, J., Nakjang, S., and et al. (2017). Novel molecular subgroups for clinical classification and outcome prediction in childhoodmedulloblastoma: a cohort study. *Lancet Oncology*, 18(7):958–971.
- Schwarz, G. (1978). Estimating the dimension of a model. *The annals of statistics*, pages 461–464.
- Seibold, H., Zeileis, A., and Hothorn, T. (2016). Model-based recursive partitioning for subgroup analyses. *The international journal of biostatistics*, 12(1):45–63.
- Sela, R. J. and Simonoff, J. S. (2012). Re-em trees: a data mining approach for longitudinal and clustered data. *Machine learning*, 86(2):169–207.
- Shen, J. and Qu, A. (2020). Subgroup analysis based on structured mixed-effects models for longitudinal data. *Journal of Biopharmaceutical Statistics*, 30(4):607–622.
- Shen, X., Wolfe, D., and Zhou, S. (1998). Local asymptotics for regression splines and confidence regions. *The annals of statistics*, 26(5):1760–1782.
- Song, J. J., Lee, H.-J., Morris, J. S., and Kang, S. (2007). Clustering of time-course gene expression data using functional data analysis. *Computational biology and chemistry*, 31(4):265–274.
- Su, X., Zhou, T., Yan, X., Fan, J., and Yang, S. (2008). Interaction trees with censored survival data. The international journal of biostatistics, 4(1).
- Talwalkar, J. A. and Lindor, K. D. (2003). Primary biliary cirrhosis. *The Lancet*, 362(9377):53–61.
- Tang, C. Y., Zhang, W., and Leng, C. (2019). Discrete longitudinal data modeling with a mean-correlation regression approach. *Statistica Sinica*, 29(2):853–876.
- Vinh, N. X., Epps, J., and Bailey, J. (2010). Information theoretic measures for clusterings comparison: Variants, properties, normalization and correction for chance. *The Journal of Machine Learning Research*, 11:2837–2854.
- Wei, Y., Liu, L., Su, X., Zhao, L., and Jiang, H. (2020). Precision medicine: Subgroup identification in longitudinal trajectories. *Statistical methods in medical research*, 29(9):2603–2616.
- Zeileis, A., Hothorn, T., and Hornik, K. (2008). Model-based recursive partitioning. *Journal of Computational and Graphical Statistics*, 17(2):492–514.
- Zhang, T. and Lin, G. (2021). Generalized k-means in glms with applications to the outbreak of covid-19 in the united states. *Computational Statistics & Data Analysis*, 159:107217.
- Zhang, Y., Li, R., and Tsai, C.-L. (2010). Regularization parameter selections via generalized information criterion. *Journal of the American Statistical Association*, 105(489):312–323.

Zhang, Y., Wang, H. J., and Zhu, Z. (2019a). Quantile-regression-based clustering for panel data. *Journal of Econometrics*, 213(1):54–67. Annals: In Honor of Roger Koenker.

Zhang, Y., Wang, H. J., and Zhu, Z. (2019b). Robust subgroup identification. *Statistica Sinica*, 29(4):1873–1889.

Zhu, Z., Fung, W. K., and He, X. (2008). On the asymptotics of marginal regression splines with longitudinal data. *Biometrika*, 95(4):907–917.

Appendix

Lemma 1. Without loss of generality, we consider the following B-spline basis functions that span G, that is, $B_k = J_n^{1/2} S_k$, $k = 1, ..., J_n$, where $\{S_k\}_{k=1}^{J_n}$ are the B-splines defined in Chapter 5 of Lorentz and DeVore (1993). It follows from Theorem 4.2 f Lorentz and DeVore (1993) that

$$|M_1||\gamma||_2^2 \le \int \{\sum_{k=1}^{J_n} B_k(t)\gamma_k\}^2 dt \le M_2||\gamma||_2^2$$

for some constant $0 < M_1 < M_2 < \infty$, where $\gamma = (\gamma_1, \ldots, r_{J_n})'$.

Lemma 2. For each i, there exist some constants $0 < M_1 < M_2 < \infty$ such that, except on an event whose probability tends to zero, all the eigenvalues of $\mathbf{X}_i'\mathbf{X}_i/n_i$ fall between M_1 and M_2 .

The proof of Lemma 2 can be referred to Theorem 7.3 of Roussas and Ioannides (1987).

Lemma 3. Define $f_i^*(t) = \mathbf{B}(t)\gamma^* \in G$ such that $||f_i^*(t) - f_i(t)||_2^2 = \inf_{g \in G} ||g(t) - f_i(t)||_2^2 \triangleq \varpi_i = J_n^{-2r}$ if $f_i \in \mathcal{H}_r$, which follows the result of page 149 of De Boor and De Boor (1978).

Lemma 4. Under Conditions (A1)-(A5), as $n \to \infty$ and if $J_n \to \infty$, $J_n^4 = o(n_0)$, then the B-splines coefficients we import into k-means satisfy

$$\|\widehat{\gamma}_{ij}^{(1)} - \gamma_{ij}\|_{2}^{2} = O_{p}(J_{n}/n_{i} + J_{n}^{-2r}),$$

where $i=1,\ldots,n, j=1,\ldots,p,\ \gamma_{ij}$ denotes the true B-splines coefficients for the j additive component of the i subject, and $\widehat{\gamma}_{ij}^{(r)}$ represents the estimate of γ_{ij} at the first iteration.

Remark 4. In Lemma 4, we investigate the convergence property of our initial estimator, which means that in the first iteration, the B-spline coefficients we input into k-means algorithm is close to the true coefficients as long as the repeated measurements of each subjects are sufficiently large, and this is consistent with our simulation studies. Moreover, we have noticed that the average mean squared error of our estimated coefficients is determined by two parts. The first part consists of average variance and approximation bias from the random effects, and the second part represents average squared shrinkage bias. Lemma 4 guarantees that the estimated subgroups $\widehat{\Omega}_{k,j}$ equal to the true subgroups $\Omega_{k,j}$ with probability 1.

Proof of Lemma 4. Denote $U_i = (S_i, X_i)$, following the idea of Huang et al. (2007), the initial estimates are

$$\begin{pmatrix} \widehat{\beta} \\ \widehat{\gamma} \end{pmatrix} = (\sum_{i=1}^n \boldsymbol{U}_i' \boldsymbol{V}_i^{-1} \boldsymbol{U}_i)^{-1} \sum_{i=1}^n \boldsymbol{U}_i' \boldsymbol{V}_i^{-1} \boldsymbol{Y}_i.$$

Let

$$\sum_{i=1}^{n} U_i' V_i^{-1} U_i = \begin{pmatrix} \sum_{i=1}^{n} S_i' V_i^{-1} S_i & \sum_{i=1}^{n} S_i' V_i^{-1} X_i \\ \sum_{i=1}^{n} X_i' V_i^{-1} S_i & \sum_{i=1}^{n} X_i' V_i^{-1} S_i \end{pmatrix} \triangleq \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix},$$

and it follows that well-known block matrix forms of matrix inverse that

$$\begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix}^{-1} = \begin{pmatrix} H^{11} & H^{12} \\ H^{21} & H^{22} \end{pmatrix} = \begin{pmatrix} H_{11 \cdot 2}^{-1} & -H_{11 \cdot 2}^{-1} H_{12} H_{22}^{-1} \\ -H_{22 \cdot 1}^{-1} H_{21} H_{11}^{-1} & H_{22 \cdot 1}^{-1} \end{pmatrix},$$

where $H_{11\cdot 2}=H_{12}H_{2}2^{-1}H_{21}$ and $H_{11\cdot 2}=H_{22}-H_{21}H_{11}^{-1}H_{12}$. Consequently,

$$\widehat{\beta} = H^{11} \left\{ \sum_{i=1}^{n} \mathbf{S}_{i}' \mathbf{V}_{i}^{-1} \mathbf{Y}_{i} - H_{12} H_{22}^{-1} \sum_{i=1}^{n} \mathbf{X}_{i}' \mathbf{V}_{i}^{-1} \mathbf{Y}_{i} \right\},\,$$

$$\widehat{\gamma}^{(0)} = H^{22} \left\{ \sum_{i=1}^{n} \mathbf{X}_{i}' \mathbf{V}_{i}^{-1} \mathbf{Y}_{i} - H_{21} H_{11}^{-1} \sum_{i=1}^{n} \mathbf{S}_{i} \mathbf{V}_{i}^{-1} \mathbf{Y}_{i} \right\},\,$$

where $\hat{\gamma}^{(0)} = (\hat{\gamma}_1^{(0)}, \dots, \hat{\gamma}_p^{(0)})'$. Next we are going to prove that the initial estimate of B-splines coefficients are bounded with probability and let us take j = 1 as example in the following proof. In the backfitting procedure, we define

$$oldsymbol{W}_{i1} = oldsymbol{Y}_i - oldsymbol{S}_i \widehat{eta} - \sum_{k=2}^p oldsymbol{B}_{ik} \hat{\gamma}_k^{(0)},$$

where $X_{ik} = (X_{i1k}, \dots, X_{in_ik})$, and update $f_{i1} = S(W_{i1}|x_{i1})$ with B-splines, which means in this situation we fit

$$egin{aligned} m{W}_{i1} &= f_{i1}(m{X}_{i1}) + m{Z}_i b_i + m{arepsilon}_i \ &pprox m{B}_{i1} r_{i1} + m{Z}_i b_i + m{arepsilon}_i. \end{aligned}$$

So the initial estimates we import into k-means algorithm are

$$\widehat{\gamma}_{i1}^{(1)} = (B'_{i1}V_i^{-1}B_{i1})^{-1}B'_{i1}V_i^{-1}W_{i1}.$$

Thus,

$$\begin{aligned} &\|\widehat{\gamma}_{i1}^{(1)} - \gamma_{i1}^*\|_2^2 = \|(\boldsymbol{B}_{i1}'\boldsymbol{V}_i^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}'\boldsymbol{V}_i^{-1}\boldsymbol{W}_{i1} - \gamma_{i1}^*\|_2^2 \\ &= \|(\boldsymbol{B}_{i1}'\boldsymbol{V}_i^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}'\boldsymbol{V}_i^{-1}(\boldsymbol{Y}_i - \boldsymbol{S}_i\widehat{\beta} - \sum_{k=2}^p \boldsymbol{B}_{ik}\widehat{\gamma}_k^{(0)}) - \gamma_{i1}^*\|_2^2 \end{aligned}$$

$$= \|(\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}(\boldsymbol{S}_{i}(\beta-\widehat{\beta})+f_{i1}+\sum_{k=2}^{p}(f_{ik}-\boldsymbol{B}_{ik}\widehat{\gamma}_{k}^{(0)})+\boldsymbol{Z}_{i}b_{i}+\varepsilon_{i})-\gamma_{i1}^{*}\|_{2}^{2}$$

$$\leq \|(\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}\boldsymbol{S}_{i}(\beta-\widehat{\beta})\|_{2}^{2}+\|(\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}f_{i1}-\gamma_{i1}^{*}\|_{2}^{2}$$

$$+\|(\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}\sum_{k=2}^{p}(f_{ik}-\boldsymbol{B}_{ik}\widehat{\gamma}_{k}^{(0)})\|_{2}^{2}+\|(\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}^{\prime}\boldsymbol{V}_{i}^{-1}(\boldsymbol{Z}_{i}b_{i}+\varepsilon_{i})\|_{2}^{2}$$

$$\triangleq I_{1}+I_{2}+I_{3}+I_{4}.$$

For I_1 , it is obviously that

$$E(I_{1}) = E(\|(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}S_{i}(\beta-\widehat{\beta})\|_{2}^{2})$$

$$= E((\beta-\widehat{\beta})'\boldsymbol{S}'_{i}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{S}_{i}(\beta-\widehat{\beta}))$$

$$= tr\{E\{(\beta-\widehat{\beta})'\boldsymbol{S}'_{i}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{S}_{i}(\beta-\widehat{\beta})\}\}$$

$$= tr\{E\{\boldsymbol{S}_{i}(\beta-\widehat{\beta})(\beta-\widehat{\beta})'\boldsymbol{S}'_{i}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\}\}$$

$$\leq \lambda_{max}\{\boldsymbol{S}_{i}E((\beta-\widehat{\beta})(\beta-\widehat{\beta})')\boldsymbol{S}'_{i}\}tr\{\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\}$$

$$= O(J_{n}/n_{i}).$$

Following Lemma A1 of Zhu et al. (2008), since $\lambda_{max}\{S_iE((\beta-\widehat{\beta})(\beta-\widehat{\beta})')S_i'\} < C_1$, $\lambda_{max}(B_{i1}'V_i^{-1}V_i^{-1}B_{i1}) < C_2$, and $\lambda_{max}((B_{i1}'V_i^{-1}B_{i1})^{-1}) < C_3$, by defining $F_{i1} = \frac{1}{n_i}B_{i1}'V_i^{-1}B_{i1}$, we have $||F_{i1}^{-1}||_{\infty} = O(J_n)$. According to Lemma 6.5 of Shen et al. (1998), we obtain the above inequalities, thus $I_1 = O_p(J_n/n_i)$. For I_2 , we have

$$I_{2} = \|(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}f_{i1} - \gamma_{i1}\|_{2}^{2}$$

$$= \|(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}(f_{i1}^{*} + f_{i1} - f_{i1}^{*}) - \gamma_{i1}^{*})\|_{2}^{2}$$

$$= \|(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}(f_{i1} - f_{i1}^{*})\|_{2}^{2}$$

$$= tr\{(f_{i1} - f_{i1}^{*})'\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}(f_{i1} - f_{i1}^{*})\}$$

$$\leq \lambda_{max}(\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1})trace\{(f_{i1} - f_{i1}^{*})(f_{i1} - f_{i1}^{*})'\}$$

$$= \lambda_{max}(\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1})\|f_{i1} - f_{i1}^{*}\|_{2}^{2}.$$

Since $\lambda_{max}(\boldsymbol{V}_i^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}_{i1}'\boldsymbol{V}_i^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}_{i1}'\boldsymbol{V}_i^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}_{i1}'\boldsymbol{V}_i^{-1}) < C$, following from the result of De Boor and De Boor (1978), we know that $||f_{i1} - f_{i1}^*||_2^2 = J_n^{-2r}$, and then $I_2 = O_p(J_n^{-2r})$. Next, for I_3 ,

$$E(I_{3}) = E\{\|(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\sum_{k=2}^{p}(f_{ik}-\boldsymbol{B}_{ik}\widehat{\gamma}_{k}^{(0)})\|_{2}^{2}\}$$

$$= tr\{E\{\sum_{k=2}^{p}(f_{ik}-\boldsymbol{B}_{ik}\widehat{\gamma}_{k}^{(0)})'\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\sum_{k=2}^{p}(f_{ik}-\boldsymbol{B}_{ik}\widehat{\gamma}_{k}^{(0)})\}\}$$

$$= tr\{E\{\sum_{k=2}^{p}(f_{ik}-\boldsymbol{B}_{ik}\widehat{\gamma}_{k}^{(0)})\sum_{k=2}^{p}(f_{ik}-\boldsymbol{B}_{ik}\widehat{\gamma}_{k}^{(0)})'\}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}(\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\boldsymbol{B}_{i1})^{-1}\boldsymbol{B}'_{i1}\boldsymbol{V}_{i}^{-1}\}$$

$$\leq \lambda_{max} \left(\sum_{k=2}^{p} (f_{ik} - \boldsymbol{B}_{ik} \widehat{\gamma}_{k}^{(0)}) \sum_{k=2}^{p} (f_{ik} - \boldsymbol{B}_{ik} \widehat{\gamma}_{k}^{(0)})' \right) \\
\times tr \left\{ \boldsymbol{V}_{i}^{-1} \boldsymbol{B}_{i1} (\boldsymbol{B}_{i1}' \boldsymbol{V}_{i}^{-1} \boldsymbol{B}_{i1})^{-1} (\boldsymbol{B}_{i1}' \boldsymbol{V}_{i}^{-1} \boldsymbol{B}_{i1})^{-1} \boldsymbol{B}_{i1}' \boldsymbol{V}_{i}^{-1} \right\} \\
= \lambda_{max} \left(\sum_{k=2}^{p} (f_{ik} - \boldsymbol{B}_{ik} \widehat{\gamma}_{k}^{(0)}) \sum_{k=2}^{p} (f_{ik} - \boldsymbol{B}_{ik} \widehat{\gamma}_{k}^{(0)})' \right) \lambda_{max} (\boldsymbol{B}_{i1}' \boldsymbol{V}_{i}^{-1} \boldsymbol{V}_{i}^{-1} \boldsymbol{B}_{i1}) \\
\times \lambda_{max} \left((\boldsymbol{B}_{i1}' \boldsymbol{V}_{i}^{-1} \boldsymbol{B}_{i1})^{-1} \right) tr \left\{ n_{i}^{-1} \boldsymbol{F}_{i}^{-1} \right\} \\
= O(J_{n}/n_{i})$$

Similarly to the proof of I_1 , since $||F_{i1}^{-1}||_{\infty} = O(J_n)$ and $\lambda_{max}(\sum_{k=2}^p (f_{ik} - \boldsymbol{B}_{ik} \widehat{\gamma}_k^{(0)}) \sum_{k=2}^p (f_{ik} - \boldsymbol{B}_{ik} \widehat{\gamma}_k^{(0)})')$, $\lambda_{max}(\boldsymbol{B}'_{i1}\boldsymbol{V}_i^{-1}\boldsymbol{B}_{i1})$, $\lambda_{max}((\boldsymbol{B}'_{i1}\boldsymbol{V}_i^{-1}\boldsymbol{B}_{i1})^{-1})$ are bounded away from zero to infinity, we can obtain $I_3 = O_p(J_n/n_i)$.

Furthermore, for I_4 , by Lemma 4 and the bounded assumption on the eigenvalues of V, it is easy to verify that there exist two constants $0 < C_1 \le C_2 < \infty$, such that

$$C_1 \frac{1}{n_i^2} E((\boldsymbol{Z}_i b_i + \boldsymbol{\varepsilon}_i)' \boldsymbol{V}_i^{-1} \boldsymbol{B}_{i1} \boldsymbol{B}'_{i1} \boldsymbol{V}_i^{-1} (\boldsymbol{Z}_i b_i + \boldsymbol{\varepsilon}_i)) \leq E(I_4),$$

$$E(I_4) \leq C_2 \frac{1}{n_i^2} E((\boldsymbol{Z}_i b_i + \boldsymbol{\varepsilon}_i)' \boldsymbol{V}_i^{-1} \boldsymbol{B}_{i1} \boldsymbol{B}'_{i1} \boldsymbol{V}_i^{-1} (\boldsymbol{Z}_i b_i + \boldsymbol{\varepsilon}_i)).$$

According to the operation properties of the trace and expectation, we have

$$E((\mathbf{Z}_{i}b_{i}+\boldsymbol{\varepsilon}_{i})'\mathbf{V}_{i}^{-1}\mathbf{B}_{i1}\mathbf{B}_{i1}'\mathbf{V}_{i}^{-1}(\mathbf{Z}_{i}b_{i}+\boldsymbol{\varepsilon}_{i})) = tr\{E((\mathbf{Z}_{i}b_{i}+\boldsymbol{\varepsilon}_{i})'\mathbf{V}_{i}^{-1}\mathbf{B}_{i1}\mathbf{B}_{i1}'\mathbf{V}_{i}^{-1}(\mathbf{Z}_{i}b_{i}+\boldsymbol{\varepsilon}_{i}))\}$$

$$= E(tr\{\mathbf{B}_{i1}'\mathbf{V}_{i}^{-1}(\mathbf{Z}_{i}b_{i}+\boldsymbol{\varepsilon}_{i})(\mathbf{Z}_{i}b_{i}+\boldsymbol{\varepsilon}_{i})'\mathbf{V}_{i}^{-1}\mathbf{B}_{i1}\}) = E(tr\{\mathbf{B}_{i1}'\mathbf{V}_{i}^{-1}\boldsymbol{\Sigma}_{i}\mathbf{V}_{i}^{-1}\mathbf{B}_{i1}\})$$

$$= tr\{E(\mathbf{B}_{i1}'\mathbf{V}_{i}^{-1}\boldsymbol{\Sigma}_{i}\mathbf{V}_{i}^{-1}\mathbf{B}_{i1})\} = O_{p}(n_{i}J_{p}).$$

Hence, we obtain $I_4 = O_p(J_n/n_i)$.

Consequently, combining the I_1, I_2, I_3, I_4 , we have

$$\|\widehat{\gamma}_{i1}^{(1)} - \gamma_{i1}\|_{2}^{2} \le \|\widehat{\gamma}_{i1}^{(1)} - \gamma_{i1}^{*}\|_{2}^{2} + \|\gamma_{i1}^{*} - \gamma_{i1}\|_{2}^{2} = O_{p}(J_{n}/n_{i} + J_{n}^{-2r}).$$

This finishes the proof.

Lemma 5. If $logc/logn \rightarrow 0$ as $n_0 \rightarrow \infty$, then we have

$$\hat{m}_i \stackrel{P}{\to} m_i$$

for j = 1, ..., p, where \hat{m}_j is the number of groups selected by BIC, m_j is the true number of groups on covariate X_j , and $c = \max_{\substack{j=1,...,p\\k=1,...,\hat{m}_j}} |\Omega_{k,j}|$.

Remark 5. Lemma 5 implies that when the true number of groups m_j is unknown, BIC can be utilized to determine the number of clusters and the estimated number of groups \hat{m}_j selected by BIC converges to m_j in probability. This is consistent with some findings for BIC in tuning parameter determinations, for instance, Zhang et al. (2010) and Bai et al. (2018).

Proof of Lemma 5. If $\hat{m}_j < m_j$, then we can find at least one pair of γ_i and $\gamma_{i'}$, such that they are not in the same cluster but they are grouped to the same cluster. By the Lemma 1 of Zhang and Lin (2021), for

$$BIC(\hat{m}_i) = -2log(\hat{L}) + log(n) \times \hat{m}_i \times df,$$

where df represents the degree of freedom of B-splines, the first item on the right-hand side goes to ∞ with rate n_0 . It is faster than the rate of BIC under $\hat{m}_j = m_j$, implying that $P(\hat{m}_j < m_j) = 0$ as $n_0 \to \infty$. Therefore, we only need to prove that $\hat{m}_j > m_j$ leads to $BIC(\hat{m}_j) > BIC(m_j)$. Most of time, the degree of freedom of B-splines is not less than 5, and it suffices to show that for

$$P(BIC(m_j) < BIC(\hat{m}_j)) = P(-2(\hat{l}(m_j) - \hat{l}(\hat{m}_j)) < log(n) \times df \times (\hat{m}_j - m_j))$$

according to Donoho and Jin (2004), the limiting distribution of $-2(\hat{l}(m_j) - \hat{l}(\hat{m}_j))$ is χ^2 distribution. The right-hand side of above formula is tending to 1 as $n \to \infty$. Hence we conclude that $P(BIC(m_j) < BIC(\hat{m}_j)) \to 1$ when $\hat{m}_j > m_j$, indicating that $\hat{m}_j \stackrel{P}{\to} m_j$. This finishes the proof.

Proof of Theorem 1. When the true subgroup memberships $\Omega_{1,j}, \ldots, \Omega_{m_j,j}, j = 1, \ldots, p$ are known, the oracle approximation for $\tilde{\gamma}_j$ can be easily obtained. Taking j = 1 as an example, for $i \in \Omega_{1,j}$, we have

$$\tilde{\gamma}_{11} = (\boldsymbol{B}_1' \boldsymbol{V}_1^{-1} \boldsymbol{B}_1)^{-1} \boldsymbol{B}_1' \boldsymbol{V}_1^{-1} (\boldsymbol{Y}_1 - \boldsymbol{S}_1 \widehat{\beta} - \sum_{j=2}^p \boldsymbol{B}_j \gamma_j^{(0)}),$$

where $\mathbf{B}_1 = (B'_{11}, \dots, B'_{N_{1,1}1})'$ is the B-spline design matrix, $\mathbf{V}_1 = diag(V_{11}, \dots, V_{N_{1,1}1})$ is the working correlation matrix of whom belonging to $\Omega_{1,1}$.

Similarly to the proof of Lemma 4, we can show that under Assumptions (A1)-(A5),

$$\|\tilde{\gamma}_j - \gamma_j\|_2^2 \le O_p(J_n/N_0 + J_n^{-2r}), j = 1, \dots, p.$$

This finishes the proof.

Proof of Theorem 2. When the algorithm converges, which means when $\widehat{\Omega}_{j}^{(n)} = \widehat{\Omega}_{j}^{(n-1)}$ for $\forall j = 1, \ldots, p$, the proposed method stops. Assuming that for covariate X_{j} , there are \hat{m}_{j} groups being detected, $\widehat{\Omega}_{1,j}, \ldots, \widehat{\Omega}_{\hat{m}_{j},j}$. And the number of membership of each subgroup are $n_{1,j}, \ldots, n_{\hat{m}_{j},j}$ respectively, where $\sum_{k=1}^{\hat{m}_{j}} n_{k,j} = n$.

Let $B_{1,j}, \ldots, B_{\hat{m}_j,j}$ denote the B-spline matrix of the $\widehat{\Omega}_{1,j}, \ldots, \widehat{\Omega}_{\hat{m}_j,j}$ group respectively, and $S_{1,j}, \ldots, S_{\hat{m}_j,j}$ denote the design matrix for the individuals of group $\widehat{\Omega}_{1,j}, \ldots, \widehat{\Omega}_{\hat{m}_j,j}$. So $\widehat{\gamma}_1, \ldots, \widehat{\gamma}_p$, the estimates of $\gamma_1, \ldots, \gamma_p$ after convergence are obtained as

$$\widehat{\gamma}_{k,j} = (\boldsymbol{B}'_{k,j} \boldsymbol{V}_{k,j}^{-1} \boldsymbol{B}_{k,j})^{-1} \boldsymbol{B}'_{k,j} \boldsymbol{V}_{k,j}^{-1} (y_{k,j} - \boldsymbol{S}_{k,j} \widehat{\beta} - \sum_{l \neq j} \boldsymbol{B}_{k,l} \widehat{\gamma}_{k,l}), \quad k = 1, \dots, \hat{m}_j, j = 1, \dots, p$$

Similarly to the proof the Lemma 4, we can easily conclude that

$$\|\widehat{\gamma}_{k,j} - \gamma_{k,j}\|_2^2 = O_p(J_n/n_{0k} + J_n^{-2r}),$$

where $n_{0k} = min\{n_i, i \in \widehat{\Omega}_{k,j}\}$. As a result, we have

$$\|\widehat{\gamma}_j - \gamma_j\|_2^2 = O_p(J_n/n_0 + J_n^{-2r}),$$

which leads to

$$\|\widehat{f}_j - f_j\|_2^2 = O_p(J_n/n_0 + J_n^{-2r}).$$

This finishes the proof.

Proof of Theorem 3. According to Theorem 1 of Abraham et al. (2003), if n is sufficiently large, as long as $\|\widehat{\gamma}_{ij}^{(1)} - \gamma_{ij}\|_2^2 \to 0$, for the K-means procedure, the unique minimizer of objective function

$$\mu_n = \sum_{j=1}^k \sum_{i \in G_j} \|\widehat{\gamma}_i - c_j\|_2^2$$

is arbitrarily close to the unique minimizer of objective function

$$\mu = \sum_{j=1}^{k} \sum_{i \in G_j} \|\gamma_i - c_j\|_2^2.$$

With Theorem 1 of Abraham et al. (2003) ensuring the consistency of K-means, we can establish that as $n \to \infty$ and $\|\widehat{\gamma}_{ij}^{(\infty)} - \gamma_{ij}\|_2^2 \to 0$, the proposed method can identify the true subgroup structure with probability tending to 1. Hence, we have

$$P(\widehat{\Omega}_j = \Omega_j) \to 1.$$

This finishes the proof.