

Dynamical analysis and cosmological evolution in Weyl integrable gravity

Andronikos Paliathanasis^{1,2,*}

¹*Institute of Systems Science, Durban University of Technology, Durban 4000, South Africa*

²*Instituto de Ciencias Físicas y Matemáticas,*

Universidad Austral de Chile, Valdivia 5090000, Chile

(Dated: December 2, 2021)

We investigate the cosmological evolution for the physical parameters in Weyl integrable gravity in a Friedmann–Lemaître–Robertson–Walker universe with zero spatially curvature. For the matter component, we assume that it is an ideal gas, and of the Chaplygin gas. From the Weyl integrable gravity a scalar field is introduced by a geometric approach which provides an interaction with the matter component. We calculate the stationary points for the field equations and we study their stability properties. Furthermore, we solve the inverse problem for the case of an ideal gas and prove that the gravitational field equations can follow from the variation of a Lagrangian function. Finally, variational symmetries are applied for the construction of analytic and exact solutions.

PACS numbers: 98.80.-k, 95.35.+d, 95.36.+x

Keywords: Cosmological dynamics; Weyl integrable theory; scalar field; interaction

1. INTRODUCTION

The cosmological constant component in the Einstein–Hilbert Action Integral, is the simplest dark energy candidate to describe of the recent acceleration phase of the universe, as it is provided by the cosmological observations [1]. In the so-called Λ CDM cosmology the universe is considered to be homogeneous and isotropic, described by the Friedmann–Lemaître–Robertson–Walker (FLRW) geometry with spatially flat term, where the matter

*Electronic address: anpaliat@phys.uoa.gr

component consists of the cosmological constant and a pressureless fluid source which attributes the dark matter component of the universe. The gravitational field equations are of second-order and can be integrated explicitly. Indeed, the field equations can be reduced to that of the one-dimensional “hyperbolic oscillator”. However, as the cosmological observations are improved, Λ -cosmology loses the important position in the “armoury” of cosmologists. For an interesting discussion on the subject we refer the reader to the recent review [2]. Furthermore, because of the simplicity of the field equations in Λ -cosmology, the cosmological constant term cannot provide a solution for the description of the complete cosmological evolution and history.

In order to solve these problems, cosmologists have introduced various solutions in the literature by introducing new degrees of freedom in the field equations. Time-varying Λ term, scalar fields and fluids with time-varying equation of state parameters, like the Chaplygin gases have been proposed to modify the energy-momentum tensor of the field equations [3–8]. On the other hand, a different approach is inspired by the modification of the Einstein-Hilbert Action integral, and leads to the family of theories known as alternative/modified theories of gravity [9–11]. Another interesting consideration is the interaction between the various components consisted of the energy momentum tensor [12]. Interaction in the dark components of the cosmological model, that is, between, the dark energy and the dark matter terms is supported by cosmological observations [13–16].

For a given proposed dark energy mode model, there are systematic methods for the investigation of the physical properties of the model. The derivation of exact and analytic solutions is an essential approach because analytic techniques can be used for the investigation of the cosmological viability of the model [17–20]. Furthermore, from the analysis of the asymptotic dynamics, that is, of the determination of the stationary points, the complete cosmological history can be constructed [21–23]. Indeed, constraints for the free parameters of a given model can be constructed through the analysis of the stationary points and the specific requirements for the stability of the stationary points [24–28].

In this piece of work, we study the evolution of the cosmological dynamics for the theory known as Weyl integrable gravity (WIG) [29–35]. In WIG a scalar field is introduced into the Einstein-Hilbert Action Integral by a geometric construction approach. Indeed, in Riemannian geometry the basic geometric object is the covariant derivative ∇_μ and the metric tensor $g_{\mu\nu}$, such that it has no metricity component, i.e. $\nabla_\kappa g_{\mu\nu} = 0$ [36]. In Weyl geometry

the fundamental geometric objects are the gauge vector field ω_μ and the metric tensor $g_{\mu\nu}$, such that $\tilde{\nabla}_\kappa g_{\mu\nu} = \omega_\kappa g_{\mu\nu}$, where now $\tilde{\nabla}_\mu$ notes the covariant derivative with respect to the affine connection $\tilde{\Gamma}_{\mu\nu}^\kappa$ which is defined as $\tilde{\Gamma}_{\mu\nu}^\kappa = \Gamma_{\mu\nu}^\kappa - \omega_{(\mu}\delta_{\nu)}^\kappa + \frac{1}{2}\omega^\kappa g_{\mu\nu}$. When ω_μ is defined by a scalar field ϕ , $\tilde{\Gamma}_{\mu\nu}^\kappa$ describes the affine connection for the conformal metric $\tilde{g}_{\mu\nu} = \phi g_{\mu\nu}$. The field equations of the WIG in the vacuum are equivalent to that of General Relativity with a massless scalar field, with positive or negative energy density. However, when a matter source is introduced, interaction terms appear as a natural consequence of the geometry of the theory [36]. In geometric terms of interaction context, we investigate the dynamics of the cosmological field equations so that we construct the cosmological history and investigate the viability of the theory. Furthermore, the integrability property for the field equations is investigated by using the method of variational symmetries for the determination of conservation laws.

In Section 2 we present the basic elements for the WIG theory. Furthermore, we write the field equations for our cosmological model in a spatially flat FLRW background space. In Section 3 we present the main results of our analysis in which we discuss the asymptotic dynamics for the field equations in the cases for which the matter source is an ideal gas, or a Chaplygin gas. Moreover, we investigate the dynamics in the presence of the cosmological constant term. In Section 4 we show that the field equations have a minisuperspace description when the matter source is an ideal gas. Specifically, we solve the inverse problem and we construct a point-like Lagrangian which describes the cosmological field equations. With the use of the variational symmetries we determine a conservation law and we present the analytic solution for the field equations by using the Hamilton-Jacobi approach. Our results are summarized in Section 5.

2. WEYL INTEGRABLE GRAVITY

Consider the two conformal related metric tensors $g_{\mu\nu}$, $\tilde{g}_{\mu\nu}$ such that $\tilde{g}_{\mu\nu} = \phi g_{\mu\nu}$. The Christoffel symbols of the two conformal related metrics are related as

$$\tilde{\Gamma}_{\mu\nu}^\kappa = \Gamma_{\mu\nu}^\kappa - \phi_{,(\mu}\delta_{\nu)}^\kappa + \frac{1}{2}\phi^{,\kappa} g_{\mu\nu}. \quad (1)$$

In Weyl geometry the fundamental objects are the metric tensor $g_{\mu\nu}$ and the covariant

derivative $\tilde{\nabla}_\mu$ defined by the Christoffel symbols $\tilde{\Gamma}_{\mu\nu}^\kappa$. Hence, the curvature tensor is defined

$$\tilde{\nabla}_\nu \left(\tilde{\nabla}_\mu u_\kappa \right) - \tilde{\nabla}_\mu \left(\tilde{\nabla}_\nu u_\kappa \right) = \tilde{R}_{\kappa\lambda\mu\nu} u^\lambda. \quad (2)$$

Consequently, the Ricci tensors of the two conformal metrics are related as follow

$$\tilde{R}_{\mu\nu} = R_{\mu\nu} - \tilde{\nabla}_\nu \left(\tilde{\nabla}_\mu \phi \right) - \frac{1}{2} \left(\tilde{\nabla}_\mu \phi \right) \left(\tilde{\nabla}_\nu \phi \right) - \frac{1}{2} g_{\mu\nu} \left(\frac{1}{\sqrt{-g}} \tilde{\nabla}_\nu \tilde{\nabla}_\mu \left(g^{\mu\nu} \sqrt{-g} \phi \right) - g^{\mu\nu} \left(\tilde{\nabla}_\mu \phi \right) \left(\tilde{\nabla}_\nu \phi \right) \right), \quad (3)$$

thus the Ricci scalar

$$\tilde{R} = R - \frac{3}{\sqrt{-g}} \tilde{\nabla}_\nu \tilde{\nabla}_\mu \left(g^{\mu\nu} \sqrt{-g} \phi \right) + \frac{3}{2} \left(\tilde{\nabla}_\mu \phi \right) \left(\tilde{\nabla}_\nu \phi \right). \quad (4)$$

In WIG the the fundamental Action Integral is defined by using the Weyl Ricci scalar \tilde{R} and the scalar field ϕ by the expression

$$S_W = \int dx^4 \sqrt{-g} \left(\tilde{R} + \xi \left(\tilde{\nabla}_\nu \left(\tilde{\nabla}_\mu \phi \right) \right) g^{\mu\nu} - \Lambda \right), \quad (5)$$

where ξ is a coupling constant. From (5) we observe that ϕ is a massless scalar field. However, in a more general consideration a potential function may be considered.

From the Action Integral (5) the Weyl-Einstein equations are as [36]

$$\tilde{G}_{\mu\nu} + \tilde{\nabla}_\nu \left(\tilde{\nabla}_\mu \phi \right) - (2\xi - 1) \left(\tilde{\nabla}_\mu \phi \right) \left(\tilde{\nabla}_\nu \phi \right) + \xi g_{\mu\nu} g^{\kappa\lambda} \left(\tilde{\nabla}_\kappa \phi \right) \left(\tilde{\nabla}_\lambda \phi \right) - \Lambda g_{\mu\nu} = 0, \quad (6)$$

where $\tilde{G}_{\mu\nu}$ is the Weyl Einstein tensor. By using the Riemannian Einstein tensor $G_{\mu\nu}$, the Weyl-Einstein field equations (6) become [36]

$$G_{\mu\nu} - \lambda \left(\phi_{,\mu} \phi_{,\nu} - \frac{1}{2} g_{\mu\nu} \phi^{,\kappa} \phi_{,\kappa} \right) - \Lambda g_{\mu\nu} = 0, \quad (7)$$

where λ is defined as $2\lambda \equiv 4\xi - 3$. Equations (7) are nothing else than the field equations of Einstein's General Relativity with a massless scalar field. When $\lambda > 0$, the scalar field ϕ is a quintessence while, when $\lambda < 0$, ϕ is a phantom field [36]

Moreover, for the equation of motion of the scalar field ϕ , the Klein-Gordon equation is [36]

$$\left(\tilde{\nabla}_\nu \left(\tilde{\nabla}_\mu \phi \right) \right) g^{\mu\nu} + 2g^{\mu\nu} \left(\tilde{\nabla}_\mu \phi \right) \left(\tilde{\nabla}_\nu \phi \right) = 0, \quad (8)$$

or by using the Riemannian covariant derivative ∇_μ , expression (8) is written in the usual form $g^{\mu\nu} \nabla_\nu \nabla_\mu \phi = 0$.

As it was found in [36], the introduction of a perfect fluid in the gravitational model leads to the following set of gravitational field equations [36]

$$\tilde{G}_{\mu\nu} + \tilde{\nabla}_\nu (\tilde{\nabla}_\mu \phi) - (2\xi - 1) (\tilde{\nabla}_\mu \phi) (\tilde{\nabla}_\nu \phi) + \xi g_{\mu\nu} g^{\kappa\lambda} (\tilde{\nabla}_\kappa \phi) (\tilde{\nabla}_\lambda \phi) - \Lambda g_{\mu\nu} = e^{-\frac{\phi}{2}} T_{\mu\nu}^{(m)}, \quad (9)$$

that is,

$$G_{\mu\nu} - \lambda \left(\phi_{,\mu} \phi_{,\nu} - \frac{1}{2} g_{\mu\nu} \phi^{,\kappa} \phi_{,\kappa} \right) - \Lambda g_{\mu\nu} = e^{-\frac{\phi}{2}} T_{\mu\nu}^{(m)}, \quad (10)$$

where $T_{\mu\nu}^{(m)} = (\rho_m + p_m) u_\mu u_\nu + p_m g_{\mu\nu}$.

Moreover, the modified Klein-Gordon equation follows [36]

$$-g^{\mu\nu} \nabla_\nu \nabla_\mu \phi = \frac{1}{2\lambda} e^{-\frac{\phi}{2}} \rho_m. \quad (11)$$

Equation (11) follows from the identity $G_{;\nu}^{\mu\nu} = 0$, which provides the conserve of the effective energy-momentum tensor.

2.1. FLRW spacetime

Following the cosmological principle, in very large scales the universe is considered to be isotropic and homogeneous. Hence, the physical space is described by the FLRW spacetime, where the three-dimensional surface is a maximally symmetric space and admits six isometries. However, from cosmological observations the spatial curvature is very small, which means that we can consider as background space the spatially flat FLRW metric

$$ds^2 = -dt^2 + a^2(t) (dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2)). \quad (12)$$

Moreover, we assume the comoving observer $u_\mu = \delta_\mu^t$, with expansion rate $\theta = 3\frac{\dot{a}}{a}$, for the line element (12) and for a scalar field $\phi = \phi(t)$, the gravitational field equations are

$$\frac{\theta^2}{3} - \frac{\lambda}{2} \dot{\phi}^2 - \Lambda - e^{-\frac{\phi}{2}} \rho_m = 0, \quad (13)$$

$$\dot{\theta} + \frac{1}{3}\theta^2 + \frac{1}{2}e^{-\frac{\phi}{2}}(\rho_m + 3p_m) + \lambda\dot{\phi}^2 - \Lambda = 0, \quad (14)$$

$$\ddot{\phi} + \theta\dot{\phi} + \frac{1}{2\lambda}e^{-\frac{\phi}{2}}\rho_m = 0 \quad (15)$$

and

$$\dot{\rho}_m + \theta(\rho_m + p_m) - \rho_m \dot{\phi} = 0. \quad (16)$$

From the modified Friedmann equations we observe the existence of a nonzero interacting term for scalar field ϕ and the matter component ρ_m . When $\lambda > 0$, energy decays from scalar field to the ρ_m , while for $\lambda < 0$ energy decays from ρ_m to the field ϕ . Furthermore, the effective equation of state parameter for the effective cosmological matter is defined as $w_{eff} = -1 - 2\frac{\dot{\theta}}{\theta^2}$.

Finally, for the nature of the matter source ρ_m in the following we consider that ρ_m is an ideal gas, or a Chaplygin gas.

3. COSMOLOGICAL DYNAMICS

We continue our analysis with the investigation of the stationary points for the cosmological field equations. In order to proceed with the study we define the new dimensionless variables in the context of θ -normalization

$$x = \sqrt{\frac{3}{2}} \frac{\dot{\phi}}{\theta}, \quad \Omega_\Lambda = \frac{3\Lambda}{\theta^2}, \quad \Omega_m = \frac{3\rho_m}{\theta^2} e^{-\frac{\phi}{2}} \quad (17)$$

where for the equation of state parameter for the matter source we consider (i) ideal gas $p_m = (\gamma - 1)\rho_m$, $0 \leq \gamma < 2$, and (ii) Chaplygin gas $p_m = \frac{A_0}{\rho_m^\alpha}$, $\alpha \geq 1$. Moreover, we define the new independent parameter to be $\tau = \ln(a)$, such that $x' = \frac{dx}{d\tau}$.

At the stationary points the effective equation of the state parameter is defined as $w_{eff} = w_{eff}(x, \Omega_\Lambda, \Omega_m)$, so that the asymptotic solution is described by the scale factor $a(t) = a_0 t^{\frac{2}{3(1+w_{eff})}}$, $w_{eff} \neq -1$ and $a(t) = a_0 e^{H_0 t}$, when $w_{eff} = -1$.

3.1. Ideal gas with $\Lambda = 0$

Assume the equation of state of an ideal gas $p_m = (\gamma - 1)\rho_m$, without the cosmological constant term. Then in the new dimensionless variables (17) the field equations are

$$\Omega_m = 1 - \lambda x^2, \quad (18)$$

$$x' = -\frac{(1 - \lambda x^2)(\sqrt{6} - 6(\gamma - 2)\lambda x)}{12\lambda}. \quad (19)$$

Moreover, Ω_m is bounded as $0 \leq \Omega_m \leq 1$, such that the solution is physically acceptable, that is, from (18) it follows that there are physical stationary points only when $\lambda > 0$.

The stationary points of equation (19) are

$$A_1^\pm : x_1^\pm = \frac{1}{\sqrt{\lambda}}, \quad A_2 : x_2 = \frac{1}{\sqrt{6}(\gamma-2)\lambda}. \quad (20)$$

Points x_1^\pm describe asymptotic solutions where only the scalar field contributes to the cosmological fluid. The effective equation of state parameter is derived to be $w_{eff}(x_1^\pm) = 1$, from which we infer that the solution is that of a stiff fluid. On the other hand, the point x_2 is physically acceptable when $\lambda \geq \frac{1}{6(\gamma-2)^2}$, and the point describes a scaling solution with $w_{eff}(x_2) = -1 + \gamma + \frac{1}{6\lambda(\gamma-2)}$. For $\gamma < \frac{2}{3}$, $\lambda > \frac{1}{8(1-2\gamma)+6\gamma^2}$ it follows that $w_{eff}(x_2) < -\frac{1}{3}$ which means that the asymptotic solution describes an accelerated universe, where in the limit $\lambda = \frac{1}{8(1-2\gamma)+6\gamma^2}$, the asymptotic solution is that of the de Sitter universe.

We proceed with the investigation of the stability properties for the stationary points. We linearize equation (19) and we find the eigenvalues $e_1(x_1^\pm) = 2 - \gamma \mp \frac{1}{\sqrt{6\lambda}}$, $e_1(x_2) = -1 + \frac{\gamma}{2} + \frac{1}{12\lambda(2-\gamma)}$. Thus, point x_1^- is always a source, x_1^+ is an attractor when $\lambda < \frac{1}{6(\gamma-2)^2}$, while x_2 is the unique attractor when it exists.

3.2. Ideal gas with $\Lambda \neq 0$

In the presence of the cosmological constant, that is, $\Lambda \neq 0$, and when the matter term is that of the ideal gas, the field equations are written as follows

$$\Omega_m = 1 - \lambda x^2 - \Omega_\Lambda, \quad (21)$$

$$\Omega'_\Lambda = -\Omega_\Lambda ((\gamma-2)\lambda x^2 + \gamma(\Omega_\Lambda - 1)) \quad (22)$$

and

$$x' = \frac{1}{12\lambda} \left((\lambda x^2 - 1) \left(\sqrt{6} - 6(\gamma-2)\lambda x \right) + \left(\sqrt{6} - 6\gamma\lambda x \right) \Omega_\Lambda \right). \quad (23)$$

Furthermore, we assume that $|\Omega_\Lambda| \leq 1$, from which we infer that x is also bounded, and we do not have to study the dynamical system for the existence of stationary points at infinity.

The stationary points of the dynamics system (22), (23) are defined in the plane $\{x, \Omega_\Lambda\}$, that is $B = (x(B), \Omega_\Lambda(B))$. The points are

$$B_1^\pm = \left(\pm \frac{1}{\sqrt{\lambda}}, 0 \right), \quad B_2 = \left(\frac{1}{\sqrt{6}(\gamma-2)\lambda}, 0 \right), \quad (24)$$

$$B_3 = (0, 1) , B_4 = \left(\sqrt{6}\gamma, 1 + 6(2 - \gamma)\gamma\lambda \right) . \quad (25)$$

Points B_1^\pm , B_2 are actually the stationary points A_1^\pm and A_2 respectively for which the cosmological constant component is zero. The physical properties are the same as before. However, we should investigate the stability analysis.

For point B_3 we derive $w_{eff}(B_3) = -1$, $\Omega_m(B_3) = 0$. Thus point B_3 describes a de Sitter universe.

Furthermore, point B_4 provides $\Omega_m(B_4) = -12\gamma\lambda$, $w_{eff}(B_4) = -1$. The point is physically acceptable when $-\frac{1}{24} \leq \lambda < 0$, or $\lambda < -\frac{1}{24}$ with $\gamma \leq -\frac{1}{12\lambda}$ or $\gamma = 0$. The stationary point describes the de Sitter universe in which all the fluid components contribute in the cosmological solution.

We linearize the dynamical system (22), (23) around the stationary points and we derive the eigenvalues. For points B_1^\pm the eigenvalues are $e_1(B_1^\pm) = 2 - \gamma \mp \frac{1}{\sqrt{6\lambda}}$, $e_2(B_1^\pm) = 2$ from which we infer that B_1^- is always a source, while B_1^+ is a saddle point when $\lambda < \frac{1}{6(\gamma-2)^2}$. Otherwise it is a source.

For point B_2 the two eigenvalues are $e_1(B_2) = -1 + \frac{\gamma}{2} + \frac{1}{12\lambda(2-\gamma)}$, $e_2(B_2) = \gamma + \frac{1}{6\lambda(2-\gamma)}$. Thus, point is always a saddle point when it is physically acceptable because $e_1(B_2)$ is always negative while $e_2(B_2)$ is always positive.

The eigenvalues of the linearized system around the de Sitter point B_3 are calculated to be $e_1(B_3) = -1$ and $e_2(B_3) = -\gamma$, from which we infer that the point is always an attractor. Finally, for point B_4 we find the eigenvalues $e^\pm(B_4) = -\frac{1}{2} \pm \sqrt{1 + 4\gamma(1 + 6(2 - \gamma)\lambda)}$. Consequently, point B_4 is always a saddle point.

3.3. Chaplygin gas with $\Lambda = 0$

Consider now that the matter source satisfies the equation of the state parameter of a Chaplygin gas, $p_m = \frac{A_0}{\rho_m^\alpha}$, for which $\alpha \geq 1$, $A_0 = (-1)^\alpha 3^{-(1+\alpha)} A$ and $\rho_m \neq 0$. The field equations are written as follows

$$\Omega_m = 1 - \lambda x^2 , \quad (26)$$

$$x' = \frac{1}{12} \left(\frac{(\sqrt{6} + 6\lambda x)(\lambda x^2 - 1)}{\lambda} + 6x(\lambda x^2 - 1)^{-\alpha} Y \right) \quad (27)$$

and

$$Y' = \frac{1 + \alpha}{6} Y \left(6 - \sqrt{6}x + 6\lambda x^2 + 6(\lambda x^2 - 1)^{-\alpha} Y \right) , \quad (28)$$

where the new variable Y is defined as $Y = Ae^{-\frac{1}{2}(1+\alpha)\phi}\theta^{-(2+\alpha)}$.

The stationary points $C = (x(C), Y(C))$ of the dynamical system (27), (28), with $\Omega_m > 0$ are

$$C_1 = \left(-\frac{1}{\sqrt{6\lambda}}, 0 \right), \quad (29)$$

$$C_2 = \left(\sqrt{\frac{3}{2}} - \frac{\sqrt{\lambda(1+3\lambda)}}{\sqrt{2\lambda}}, \frac{\left(\sqrt{3\lambda(1+3\lambda)} + 6\lambda \left(1 + 3\lambda - \sqrt{3\lambda(1+3\lambda)} \right) \right) \left(-\frac{1}{2} - 3\lambda\sqrt{3\lambda(1+3\lambda)} \right)^\alpha}{6\lambda} \right) \quad (30)$$

and

$$C_3 = \left(\sqrt{\frac{3}{2}} + \frac{\sqrt{\lambda(1+3\lambda)}}{\sqrt{2\lambda}}, \frac{\left(\sqrt{3\lambda(1+3\lambda)} + 6\lambda \left(1 + 3\lambda + \sqrt{3\lambda(1+3\lambda)} \right) \right) \left(-\frac{1}{2} + 3\lambda\sqrt{3\lambda(1+3\lambda)} \right)^\alpha}{6\lambda} \right) \quad (31)$$

For point C_1 we derive $\Omega(C_1) = 1 - \frac{6}{\lambda}$, $w_{eff}(C_1) = \frac{1}{6\lambda}$. The point is physically acceptable when $\lambda \geq \frac{1}{6}$ while it always describes a universe without acceleration. For $\lambda = \frac{1}{6}$, the asymptotic solution is that of dust, while for $\lambda = \frac{1}{2}$ the asymptotic solution is that of radiation. The eigenvalues of the linearized system around the stationary point are calculated $e_1(C_1) = \frac{(1+\alpha)(1+3\lambda)}{3\lambda}$, $e_2(C_1) = \frac{1-6\lambda}{12\lambda}$, from which we can easily conclude that the stationary point is always a saddle point.

Point C_2 describes a universe for which $\Omega_m(C_2) = \frac{1}{2} - 3\lambda + \sqrt{3\lambda(1+3\lambda)}$ and $w_{eff}(C_2) = \lambda(x(C_2))^2 + (\lambda(x(C_2))^2 - 1)^{-\alpha} Y(C_2)$. The point is well defined when $\lambda > 0$, while for large values of λ it follows that $w_{eff}(C_2; \lambda \gg 1) \simeq -1$, which means that point C_2 can describe a solution near to the de Sitter point. On the other hand, point C_3 is physical acceptable for $0 < \lambda \leq \frac{1}{24}$, while we derive $\Omega_m = -3\lambda + \sqrt{3\lambda(1+3\lambda)}$ and $w_{eff}(C_3) = \lambda(x(C_3))^2 + (\lambda(x(C_3))^2 - 1)^{-\alpha} Y(C_3)$ in which $w_{eff}(C_3; \lambda = \frac{1}{24}) = 1$. Thus point C_3 does not describe any acceleration.

The eigenvalues of the linearized system near to the stationary points C_2 and C_3 are determined. Numerically we find that $e_1(C_2)$, $e_2(C_2)$ have always negative real parts for $\lambda > 0$ and $\alpha \geq 1$; on the other hand $\text{Re}(e_1(C_3)) > 0$, $\text{Re}(e_2(C_3)) > 0$ for $\alpha \geq 1$, $0 < \lambda \leq \frac{1}{24}$. Hence, point C_2 is always an attractor while point C_3 is always a source.

3.4. Chaplygin gas with $\Lambda \neq 0$

In the presence of a nonzero cosmological constant term, the field equations are reduced to the following dynamical system

$$\Omega_m = 1 - \lambda x^2 - \Omega_\Lambda , \quad (32)$$

$$\Omega'_\Lambda = \Omega_\Lambda \left(1 + \lambda x^2 - \Omega_\Lambda + Y (\lambda x^2 + \Omega_\Lambda - 1)^{-\alpha} \right) , \quad (33)$$

$$x' = \frac{1}{12} \left(x^2 (\sqrt{6} + 6\lambda) + \frac{\sqrt{6}}{\lambda} (\Omega_\Lambda - 1) + 6x \left(Y (\lambda x^2 + \Omega_\Lambda - 1)^{-\alpha} - 1 - \Omega_\Lambda \right) \right) , \quad (34)$$

$$Y' = \frac{1+\alpha}{6} Y \left(6 \left(1 + \lambda x^2 + \Omega_\Lambda + Y (\lambda x^2 + \Omega_\Lambda - 1)^{-\alpha} \right) - \sqrt{6}\lambda x \right) . \quad (35)$$

The physically acceptable stationary points $D = (x(D), Y(D), \Omega_\Lambda(D))$ are

$$D_1 = (x(C_1), Y(C_1), 0) , \quad D_2 = (x(C_2), Y(C_2), 0) , \quad (36)$$

$$D_3 = (x(C_3), Y(C_3), 0) , \quad D_4 = \left(\sqrt{6}, 1 + 6\lambda, 0 \right) , \quad (37)$$

where D_1 , D_2 and D_3 have the same physical properties as points C_1 , C_2 and C_3 respectively.

For the point D_4 we find $\Omega_m(D_4) = -12\lambda$ and $w_{eff}(D_4) = -1$, which means that the asymptotic solution is physically acceptable when $-\frac{1}{12} \leq \lambda < 0$, while the asymptotic solution is that of the de Sitter universe.

The eigenvalues of the linearized system near D_1 are $e_1(D_1) = \frac{(1+\alpha)(1+3\lambda)}{3\lambda}$, $e_2(D_1) = \frac{1-6\lambda}{12\lambda}$ and $e_3(D_1) = \frac{1+6\lambda}{6\lambda}$, which means that point D_1 is always a saddle point. For the points D_2 and D_3 we find numerically that D_2 is always an attractor while D_3 is always a source. Finally, for the point D_4 we calculate $e_1(D_4) = -(1+\alpha)$, $e_2^\pm = \frac{1}{2} (-1 \pm \sqrt{5+24\lambda})$, from which it follows that the stationary point is always a saddle point.

4. MINISUPERSPACE DESCRIPTION AND CONSERVATION LAWS

For an ideal gas $p_m = (\gamma - 1) \rho_m$, from equation (16) it follows $\rho_m(t) = \rho_{m0} a^{-3\gamma} e^\phi$ in which ρ_{m0} is a constant of integration.

We substitute this into the rest of the field equations and we end with the following dynamical system

$$\frac{\theta^2}{3} - \frac{\lambda}{2} \phi^2 - \Lambda - \rho_{m0} e^{\frac{\phi}{2}} a^{-3\gamma} = 0, \quad (38)$$

$$\dot{\theta} + \frac{1}{3}\theta^2 + \frac{(3\gamma - 2)}{2}\rho_{m0}e^{\frac{\phi}{2}}a^{-3\gamma} + \lambda\dot{\phi}^2 - \Lambda = 0, \quad (39)$$

$$\ddot{\phi} + \theta\dot{\phi} + \frac{\rho_{m0}}{2\lambda}e^{\frac{\phi}{2}}a^{-3\gamma} = 0. \quad (40)$$

For the second-order differential equations (39), (40) in the space of variables $\{a, \phi\}$, the inverse problem for the determination of a Lagrangian function, provides that the function

$$L(a, \dot{a}, \phi, \dot{\phi}) = -3a\dot{a}^2 + \frac{\lambda}{2}a^3\dot{\phi}^2 - a^3\Lambda - \rho_{m0}e^{\frac{\phi}{2}}a^{3-3\gamma} \quad (41)$$

is an autonomous Lagrangian function for the field equations, while equation (38) is conservation law of “energy”, i.e. the Hamiltonian \mathcal{H} , constraint $\mathcal{H} = 0$.

In general, the field equations for the cosmological model in WIG theory with an ideal gas, for the metric

$$ds^2 = -N^2(t) + a^2(t)(dx^2 + dy^2 + dz^2), \quad (42)$$

follow from the singular point-like Lagrangian

$$\mathcal{L}(a, \dot{a}, \phi, \dot{\phi}) = \frac{1}{N} \left(-3a\dot{a}^2 + \frac{\lambda}{2}a^3\dot{\phi}^2 \right) - N \left(a^3\Lambda + \rho_{m0}e^{\frac{\phi}{2}}a^{3-3\gamma} \right). \quad (43)$$

4.1. Integrability property and analytic solution

Because the field equations admit a point-like Lagrangian various techniques inspired by analytic mechanics be applied for the study of the dynamical system. Indeed, variational symmetries and conservation laws can be determined by using Noether’s theorems [37]. That approach has been widely used in various gravitational systems. New integrable cosmological models as also new analytic and exact solutions were found through the use of variational symmetries, see for instance [38].

We investigate for variational symmetries which have point transformations as generators and provide conservation laws linear in the velocities. Hence, for the Lagrangian function (41) and for $\rho_{m0} \neq 0$, we find that the variational symmetry $X = \frac{2}{3}a\partial_a + 4(\gamma - 2)\partial_\phi$ exists for $\Lambda = 0$, and the corresponding conservation law is

$$F(a, \dot{a}, \phi, \dot{\phi}) = 4a^2\dot{a} - 4(\gamma - 2)\lambda a^3\dot{\phi} - F_0. \quad (44)$$

Function $F(a, \dot{a}, \phi, \dot{\phi})$, $\frac{dF}{dt} = 0$, is the second-conservation law for the dynamical system, which means that the field equations form an integrable dynamical system.

In order to reduce the field equations and determine exact solutions, we apply the Hamilton-Jacobi approach. We define the momentum $p_a = -6a\dot{a}$, and $p_\phi = \lambda a^3 \dot{\phi}$, thus the Hamiltonian function $\mathcal{H}(a, \phi, p_a, p_\phi) = 0$, reads

$$-\frac{p_a^2}{6a} + \frac{p_\phi^2}{\lambda a^3} + 2 \left(a^3 \Lambda + \rho_{m0} e^{\frac{\phi}{2}} a^{3-3\gamma} \right) = 0 \quad (45)$$

while the Hamilton-Jacobi equation is written in the following form

$$-\frac{1}{6a} \left(\frac{\partial}{\partial a} S(a, \phi) \right)^2 + \frac{1}{\lambda a^3} \left(\frac{\partial}{\partial \phi} S(a, \phi) \right)^2 + 2\rho_{m0} e^{\frac{\phi}{2}} a^{3-3\gamma} = 0, \quad (46)$$

where now $p_a = \frac{\partial S}{\partial a}$ and $p_\phi = \frac{\partial S}{\partial \phi}$.

Moreover, the conservation law (44) provides the constraint equation for the Action $S(a, \phi)$

$$\frac{2a}{3} \left(\frac{\partial}{\partial a} S(a, \phi) \right) + 4(\gamma - 2) \frac{\partial}{\partial \phi} (S(a, \phi)) - F_0 = 0. \quad (47)$$

We define the new variable $\phi = 6(\gamma - 2) \ln a + \Phi$, such that the constraint equation becomes

$$\frac{2}{3} a \frac{\partial}{\partial a} (S(a, \Phi)) - F_0 = 0. \quad (48)$$

This new set of variables $\{a, \Phi\}$ are the normal coordinates for the dynamical system.

Consequently, in the normal variables the analytic expression for the Action as provided by the Hamilton-Jacobi equation is

$$S(a, \Phi) = \frac{3}{2} F_0 \ln a + \int \frac{\sqrt{2\lambda} \sqrt{16\rho_{m0} e^{\frac{\Phi}{2}} (6\lambda(\gamma - 2)^2 - 1) + 3F_0 + 6\lambda(\gamma - 2) F_0}}{4(6\lambda(\gamma - 2)^2 - 1)} d\Phi \quad (49)$$

for $(6\lambda(\gamma - 2)^2 - 1) \neq 0$, or

$$S(a, \Phi) = \frac{3}{2} F_0 \ln a + \frac{3F_0^2 \Phi - 32\rho_0 e^{\frac{\Phi}{2}}}{24F_0(\gamma - 2)}, \quad (50)$$

when $(6\lambda(\gamma - 2)^2 - 1) = 0$.

However, in the new coordinates the momentum are defined as

$$p_a = -6a \left((6\lambda(\gamma - 2)^2 - 1) \dot{a} + (\gamma - 2) \lambda a \dot{\Phi} \right), \quad (51)$$

$$p_\Phi = -\lambda a \left(6(\gamma - 2) \dot{a} + a \dot{\Phi} \right), \quad (52)$$

which give the following expressions for the scale factor and the scalar field

$$6a^2 \dot{a} = a p_a - 6(\gamma - 2) p_\Phi, \quad (53)$$

$$\lambda a^3 \dot{\Phi} = -p_{\Phi} - \lambda(\gamma - 2)(Ap_A + 6(\gamma - 2)p_{\Phi}). \quad (54)$$

Hence, by using the Action (49) and expressions (53), (54), the cosmological field equations can be written into an equivalent system. We summarize the results in the following proposition.

Proposition 1: *The field equations in WIG for a FLRW background space with zero spatial curvature and an ideal gas form a Liouville integrable system when there is no cosmological constant term. The analytic solution for the Hamilton-Jacobi equation provides the Action (49), while the field equations can be written into an equivalent set of two first-order ordinary differential equations (53), (54).*

Assume now the simple case for which $\gamma = 1$ and $F_0 = 0$. Moreover, we define the new variable $T = T(t)$, such that $dT = \frac{\sqrt{(6\lambda-1)}}{A^3} dt$ and $\lambda \neq \frac{1}{6}$.

Thus, the field equations are

$$\begin{aligned} \frac{\dot{a}}{a} - \sqrt{2\lambda\rho_{m0}}e^{\frac{\Phi}{4}} &= 0, \\ \dot{\Phi} - \sqrt{\frac{2}{\lambda}\rho_{m0}(6\lambda+1)}e^{\frac{\Phi}{4}} &= 0, \end{aligned} \quad (55)$$

with exact solution

$$a(t) = a_0 t^{\frac{4\lambda}{1+6\lambda}}, \quad \Phi(t) = -2 \ln \left(\frac{(6\lambda+1)\rho_{m0}}{8\lambda} t^2 \right). \quad (56)$$

For this exact solution the background space is

$$ds^2 = -\frac{(6\lambda-1)}{a_0^6} t^{-\frac{24\lambda}{1+6\lambda}} dT^2 + a_0^2 t^{\frac{8\lambda}{1+6\lambda}} (dx^2 + dy^2 + dz^2). \quad (57)$$

The later solution describes a universe dominated by a perfect fluid source with constant equation of state parameter. This specific solution is described by the stationary points A_2 , thus. the results are in agreement with the asymptotic analysis for the dynamics.

5. CONCLUSIONS

In this work we considered WIG to describe the cosmological evolution for the physical parameters in FLRW spacetime with zero spatially curvature. The gravitational field equations in WIG are of second-order and Einstein's theory, with the presence of the of a scalar field, is recovered. Scalar field plays the role for conformal factor which relates the connection of Weyl theory with the Levi-Civita connection of Riemannian geometry. However,

the field equations differ when matter is introduced in the gravitational model. Indeed, in WIG the matter source interacts with the scalar field. The interaction term is introduced naturally from the geometric character of the theory.

In our study we considered the matter source to be described by that of an ideal gas, that is $p_m = (\gamma - 1) \rho_m$, or by the Chaplygin gas $p_m = -\frac{A_0}{\rho_m^\alpha}$. We defined new dimensionless variables based on the Hubble-normalization in order to write the field equations as a system of first-order algebraic differential system. In each model, we determined the stationary points for the latter system and we determined their dynamical properties as also the physical properties of the asymptotic solutions. In our analysis we considered also a nonzero cosmological constant.

For the ideal gas, we found that there exists an attractor with an asymptotic solution that of an ideal gas, but with different parameter for the equation of state. For instance, we can consider the matter source to be that of radiation while the attractor to describe an accelerated universe. In the presence of the cosmological constant, we find two asymptotic solutions which can describe the past acceleration phase of the universe known as inflation, as also the late time acceleration. The future attractor describes the de Sitter universe. When the matter component is that of a Chaplygin gas the stationary points as also the cosmological evolution are similar with the previous case.

Moreover, for the ideal gas case, we solved the inverse problem and determined a Lagrangian function, and a minisuperspace description, which generates the cosmological equations under a variation. We applied Noether's theorems for point transformations in order to construct a nontrivial conservation law when the cosmological constant term is zero. Hence, the cosmological field equations form a Liouville integrable dynamical system. The closed-form expression for the Hamilton-Jacobi equation derived. Finally, for a specific values for the free parameters we were able to construct an exact solution which is in agreement with the asymptotic analysis.

In a subsequent analysis we plan to investigate further the field equations as a Hamilton system and understand how a nonzero cosmological constant affects the integrability

property of the field equations.

- [1] Planck Collaboration: N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters, *A&A* 641, A6 (2020)
- [2] L. Perivolaropoulos and F. Skara, Challenges for Λ CDM: An update, (2021) [2105.05208]
- [3] B. Ratra and P.J.E. Peebles, Cosmological consequences of a rolling homogeneous scalar field, *Phys. Rev. D* 37, 3406 (1988)
- [4] C. Armendariz-Picon, V.F. Mukhanov and P.J. Steinhardt, *Phys. Rev. D* 63, 103510 (2001)
- [5] V. Faraoni, *Cosmology in Scalar-Tensor Gravity*, Kluwer Academic Publishers, Dordrecht, (2004)
- [6] M. C. Bento, O. Bertolami and A. A. Sen, Generalized Chaplygin gas and CMBR constraints, *Phys. Rev. D* 67, 063003 (2003).
- [7] A. Kamenshchik, U. Moschella and V. Pasquier, An alternative to quintessence, *Phys. Lett. B* 511, 265 (2001)
- [8] S. Basilakos, N.E. Mavromatos and J. Solà Peracaula, *Phys. Rev. D* 101, 045001 (2020)
- [9] T. Clifton, P.G. Ferreira, A. Padilla and C. Skordis, Modified gravity and cosmology *Phys. Rept.*, 513, 1, (2012)
- [10] S.I. Nojiri and S.D. Odintsov, Introduction to modified gravity and gravitational alternative for dark energy, *IJGMMP* 4, 115 (2007)
- [11] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D.F. Mota, A.G. Riess and J. Silk, In the Realm of the Hubble tension a Review of Solutions, *Class. Quantum Grav.* 38, 153001 (2021)
- [12] A. P. Billyard and A. A. Coley, Interactions in scalar field cosmology, *Phys Rev D* 61, 083503 (2000)
- [13] S. Pan, G.S. Sharov and W. Yang, Field theoretic interpretations of interacting dark energy scenarios and recent observations, *Phys. Rev. D* 101, 103533 (2020)
- [14] W. Yang, A. Mukherjee, E. Di Valentino and S. Pan, Interacting dark energy with time varying equation of state and the H_0 tension, *Phys. Rev. D* 98, 123527 (2018)
- [15] S. Pan and G.S. Sharov, A model with interaction of dark components and recent observational data, *MNRAS* 472, 4736 (2017)

- [16] E. Di Valentino, A. Melchiorri, O. Mena and S. Vagnozzi, Interacting dark energy in the early 2020s: a promising solution to the H_0 and cosmic shear tensions, *Phys. Dark Univ.* 30, 100666 (2020)
- [17] S. Yu Vernov and E. Pozdeeva, De Sitter Solutions in Einstein–Gauss–Bonnet Gravity, *Universe* 7, 149 (2021)
- [18] A. Yu. Kamenshchik, E.O. Pozdeeva, G. Venturi and S. Yu Vernov, Integrable cosmological models in the Einstein and in the Jordan frames and Bianchi-I cosmology, *Phys. Part. Nucl.* 49, 1 (2018)
- [19] N. Dimakis, A. Paliathanasis, P.A. Terzis and T. Christodoulakis, Cosmological solutions in multiscalar field theory, *EPJC* 79, 618 (2019)
- [20] A. Paliathanasis, De Sitter and scaling solutions in a higher-order modified teleparallel theory, *JCAP* 08, 027 (2017)
- [21] E.J. Copeland, A.R. Liddle and D. Wands, Exponential potentials and cosmological scaling solution, *Phys. Rev. D* 57, 4686 (1998)
- [22] A. Coley and G. Leon, Static Spherically Symmetric Einstein-aether models I: Perfect fluids with a linear equation of state and scalar fields with an exponential self-interacting potential, *Gen. Rel. Grav.* 51, 115 (2019)
- [23] L. Amendola, D. Polarski and S. Tsujikawa, Are $f(R)$ dark energy models cosmologically viable?, *Phys. Rev. Lett.* 98, 131302 (2007)
- [24] L. Amendola, R. Gannouji, D. Polarski and S. Tsujikawa, Conditions for the cosmological viability of $f(R)$ dark energy models, *Phys. Rev. D* 75, 083504 (2007)
- [25] P. Christodoulidis, D. Roest and E.I. Sfakianakis, Scaling attractors in multi-field inflation, *JCAP* 12, 059 (2019)
- [26] C.R. Fadragas, R. Cardenas, M. Rodriguez-Ricard, A. Rivero-Acosta and A. Linares-Rodriguez, Detailed qualitative dynamical analysis of a cosmological Higgs field, *Gen. Rel. Gravit.* 51, 109 (2019)
- [27] T. Gonzalez, G. Leon and I. Quiros, Dynamics of quintessence models of dark energy with exponential coupling to dark matter, *Class. Quantum Grav.* 23, 32165 (2006)
- [28] M. Kerachia, G. Acquaviva and G. Lukes-Gerakopoulos, Dynamics of classes of barotropic fluids in spatially curved FRW spacetimes, *Phys. Rev. D* 101, 043535 (2020)
- [29] J.E. Madriz Aguilar and C. Romero, Inducing the cosmological constant from five-dimensional

- Weyl space, *Found. Phys.* 39, 1205 (2009)
- [30] Y.-X. Liu, K. Yang and Y. Zhong, de Sitter Thick Brane Solution in Weyl Geometry, *JHEP* 10, 069 (2010)
- [31] I.P. Lobo, A.B. Barreto and C. Romero, Space-time singularities in Weyl manifolds, *EPJC* 75, 448 (2015)
- [32] A. Paliathanasis and G. Leon, Integrability and cosmological solutions in Einstein-æther-Weyl theory, *EPJC* 81, 255 (2021)
- [33] M.L. Pucheu, F.A.P. Alves Junior, A.B. Barreto and C. Romero, Cosmological models in Weyl geometrical scalar-tensor theory, *Phys. Rev. D* 94, 064010 (2016)
- [34] J. Miritzis, Isotropic cosmologies in Weyl geometry, *Class. Quantum Grav.* 21, 3043 (2004)
- [35] J.M. Salim and S.L. Sautú, Gravitational collapse in Weyl integrable space-times, *Class. Quantum Grav.* 16, 3281 (1999)
- [36] J.M. Salim and S.L. Sautú, Gravitational theory in Weyl integrable spacetime, *Class. Quantum Grav.* 13, 353 (1996)
- [37] A. Halder, A. Paliathanasis and P.G.L. Leach, Noether's Theorem and Symmetry, *Symmetry* 10, 744 (2018)
- [38] M. Tsampalis and A. Paliathanasis, Symmetries of Differential Equations in Cosmology, *Symmetry* 10, 233 (2018)