

# LP-based Approximations for Disjoint Bilinear and Two-Stage Adjustable Robust Optimization

Omar El Housni

Operations Research and Information Engineering, Cornell University, New York, USA, [oe46@cornell.edu](mailto:oe46@cornell.edu)

Ayoub Foussoul

Industrial Engineering and Operations Research, Columbia University, New York, USA, [af3209@columbia.edu](mailto:af3209@columbia.edu)

Vineet Goyal

Industrial Engineering and Operations Research, Columbia University, New York, USA, [vg2277@columbia.edu](mailto:vg2277@columbia.edu)

## Abstract

We consider the class of disjoint bilinear programs  $\max \{\mathbf{x}^T \mathbf{y} \mid \mathbf{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{Y}\}$  where  $\mathcal{X}$  and  $\mathcal{Y}$  are packing polytopes. We present an  $O(\frac{\log \log m_1}{\log m_1} \frac{\log \log m_2}{\log m_2})$ -approximation algorithm for this problem where  $m_1$  and  $m_2$  are the number of packing constraints in  $\mathcal{X}$  and  $\mathcal{Y}$  respectively. In particular, we show that there exists a near-optimal solution  $(\mathbf{x}, \mathbf{y})$  such that  $\mathbf{x}$  and  $\mathbf{y}$  are “near-integral”. We give an LP relaxation of this problem from which we obtain the near-optimal near-integral solution via randomized rounding. As an application of our techniques, we present a tight approximation for the two-stage adjustable robust problem with covering constraints and right-hand side uncertainty where the separation problem is a bilinear optimization problem. In particular, based on the ideas above, we give an LP restriction of the two-stage problem that is an  $O(\frac{\log n}{\log \log n} \frac{\log L}{\log \log L})$ -approximation where  $L$  is the number of constraints in the uncertainty set. This significantly improves over state-of-the-art approximation bounds known for this problem.

**Keywords:** Disjoint bilinear programming; Two-stage robust optimization; Approximation Algorithms

## 1 Introduction

We consider the following class of disjoint bilinear programs,

$$z_{\text{PDB}} = \max_{\mathbf{x}, \mathbf{y}} \{\mathbf{x}^T \mathbf{y} \mid \mathbf{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{Y}\}, \quad (\text{PDB})$$

where  $\mathcal{X}$  and  $\mathcal{Y}$  are packing polytopes given by an intersection of knapsack constraints. Specifically,

$$\mathcal{X} := \{\mathbf{x} \geq \mathbf{0} \mid \mathbf{P}\mathbf{x} \leq \mathbf{p}\} \quad \text{and} \quad \mathcal{Y} := \{\mathbf{y} \geq \mathbf{0} \mid \mathbf{Q}\mathbf{y} \leq \mathbf{q}\},$$

where  $\mathbf{P} \in \mathbb{R}_+^{m_1 \times n}$ ,  $\mathbf{Q} \in \mathbb{R}_+^{m_2 \times n}$ ,  $\mathbf{p} \in \mathbb{R}_+^{m_1}$  and  $\mathbf{q} \in \mathbb{R}_+^{m_2}$ . We refer to this problem as a packing disjoint bilinear program **PDB**. This is a subclass of the well-studied disjoint bilinear problem:  $\max_{\mathbf{x}, \mathbf{y}} \{\mathbf{x}^T \mathbf{M} \mathbf{y} \mid \mathbf{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{Y}\}$ , where  $\mathbf{M}$  is a general  $n \times n$  matrix.

Disjoint bilinear programming is NP-hard in general (Chen et al. [9]). We show that it is NP-hard to even approximate within any finite factor. Several heuristics have been studied for this problem including cutting-planes algorithms (Konno et al. [24]), polytope generation methods (Vaish et al. [31]), Benders decomposition (Geoffrion [18]), reduction to concave minimization (Thieu [30]) and two-stage robust optimization (Zhen et al. [34]). Algorithms for non-convex quadratic optimization can also be used to solve disjoint bilinear programs.

Many important applications can be formulated as a disjoint bilinear program including fixed charge network flows (Rebennack et al. [27]), concave cost facility location (Soland [29]), bilinear assignment problems (Ćustić et al. [11]), non-convex cutting-stock problems (Harjunkski et al. [23]), multicommodity flow network interdiction problems ([25]), game theory (Mangasarian and Stone [26], Firouzbakht et al. [17]) pooling problems (Gupte et al. [22]).

One important application closely related to disjoint bilinear optimization that we focus on in this paper, is the two-stage adjustable robust optimization. In particular, the separation problem of a two-stage adjustable robust problem can be formulated as a disjoint bilinear optimization problem. More specifically, we consider the following two-stage adjustable robust problem,

$$\begin{aligned} z_{\text{AR}} = \min_{\mathbf{x}, t} \quad & \mathbf{c}^T \mathbf{x} + t \\ & t \geq \mathcal{Q}(\mathbf{x}), \\ & \mathbf{x} \in \mathcal{X}, \end{aligned} \tag{AR}$$

where for all  $\mathbf{x} \in \mathcal{X}$ ,

$$\mathcal{Q}(\mathbf{x}) = \max_{\mathbf{h} \in \mathcal{U}} \min_{\mathbf{y} \geq \mathbf{0}} \{\mathbf{d}^T \mathbf{y} \mid \mathbf{B} \mathbf{y} \geq \mathbf{h} - \mathbf{A} \mathbf{x}\}.$$

Here  $\mathbf{A} \in \mathbb{R}^{m \times n}$ ,  $\mathbf{B} \in \mathbb{R}_+^{m \times n}$ ,  $\mathbf{c} \in \mathbb{R}_+^n$ ,  $\mathbf{d} \in \mathbb{R}_+^n$ ,  $\mathcal{X} \subset \mathbb{R}_+^n$  is a polyhedral cone, and  $\mathcal{U}$  is a polyhedral uncertainty set. The separation problem of **AR** is the following: given a candidate solution  $(\mathbf{x}, t)$ , decide if it is feasible, i.e.,  $\mathbf{x} \in \mathcal{X}$  and  $t \geq \mathcal{Q}(\mathbf{x})$  or give a separating hyperplane. This is equivalent to solving  $\mathcal{Q}(\mathbf{x})$ . We will henceforth refer to  $\mathcal{Q}(\mathbf{x})$  as the separation problem. For ease of notation, we use  $\mathcal{Q}(\mathbf{x})$  to refer to both the problem and its optimal value. In this two-stage problem, the adversary observes the first-stage decision  $\mathbf{x}$  and reveals the worst-case  $\mathbf{h} \in \mathcal{U}$ . Then, the decision maker selects a second-stage recourse decision  $\mathbf{y}$  such that  $\mathbf{A} \mathbf{x} + \mathbf{B} \mathbf{y}$  covers  $\mathbf{h}$ . The goal is to select a first-stage decision such that the total cost in the worst-case is minimized. This model has been widely considered in the literature (Dhamdhare et al. [12], Feige et al. [16], Gupta et al. [21], Bertsimas and Goyal [4], Bertsimas and Bidkhori [5], Bertsimas and de Ruiter [7], Xu et al. [32], Zhen et al. [33], El Housni and Goyal [13], El Housni et al. [15]), and has many applications including set cover, capacity planning and network design problems under uncertain demand.

Several uncertainty sets have been considered in the literature including polyhedral uncertainty sets, ellipsoids and norm balls (see Bertsimas et al. [6]). Some of the most important uncertainty sets are budget of uncertainty sets (Bertsimas and Sim [8], Gupta et al. [20], El Housni and Goyal [14]) and intersections of budget of uncertainty sets such as CLT sets (see Bandi and Bertsimas [1]) and inclusion-constrained budgeted sets (see Gounaris et al. [19]). These have been widely used in practice. Following this motivation, we consider in this paper the following uncertainty set

$$\mathcal{U} := \{\mathbf{h} \geq \mathbf{0} \mid \mathbf{R}\mathbf{h} \leq \mathbf{r}\},$$

where  $\mathbf{R} \in \mathbb{R}_+^{L \times m}$  and  $\mathbf{r} \in \mathbb{R}_+^L$ . This is a generalization of the previously mentioned sets. We refer to this as a packing uncertainty set.

Feige et al. [16] show that **AR** is hard to approximate within any factor better than  $\Omega(\frac{\log n}{\log \log n})$  even in the special case of a single budget of uncertainty set. Feige et al. [16] and Gupta et al. [21] give an  $O(\log n)$ -approximation for the two-stage problem in the special case where  $\mathbf{A} = \mathbf{B} \in \{0, 1\}^{m \times n}$ ,  $\mathbf{d} = \lambda \mathbf{c}$  for some  $\lambda > 0$  and the uncertainty set  $\mathcal{U}$  is a budget of uncertainty set with equal weights, i.e.,  $\mathcal{U} = \{\mathbf{h} \in [0, 1]^m \mid \sum_{i=1}^m h_i \leq k\}$ . Gupta et al. [20] consider a more general uncertainty set, namely, intersection of *p-system* and *q-knapsack* and give an  $O(pq \log n)$ -approximation of the two-stage problem. Bertsimas and Goyal give an  $O(\sqrt{m})$ -approximation in the case where the first-stage matrix  $\mathbf{A}$  is non-negative. Recently, El Housni and Goyal [14] give an  $O(\frac{\log n}{\log \log n})$ -approximation in the case of a single budget of uncertainty set and an  $O(\frac{\log^2 n}{\log \log n})$ -approximation in the case of an intersection of disjoint budgeted sets. In general, they show an  $O(\frac{L \log n}{\log \log n})$ -approximation in the general case of a packing uncertainty set with  $L$  constraints. However, this bound scales linearly with  $L$ .

The goal of this paper is to provide LP-based approximation algorithms with provable guarantees for the packing disjoint bilinear program as well as the two-stage adjustable robust problem **AR** that improve over the approximation bounds known for these problems.

## 1.1 Our contributions

**A polylogarithmic approximation algorithm for PDB.** We present an LP-rounding based randomized approximation algorithm for **PDB**. Our algorithm relies on a new idea that might be of independent interest. In particular, we show the existence of a near-optimal near-integral solution of this problem. That is, a near-optimal solution  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  such that  $\hat{x}_i \in \{0, \max_{\mathbf{x} \in \mathcal{X}} x_i / \zeta_1\}$  and  $\hat{y}_i \in \{0, \max_{\mathbf{y} \in \mathcal{Y}} y_i / \zeta_2\}$  for some logarithmic factors  $\zeta_1$  and  $\zeta_2$ . We give an LP relaxation of **PDB**, i.e., a linear program whose optimal cost is greater than the optimum of **PDB**, from which we obtain such  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  via randomized rounding. More specifically, we have the following theorem,

**Theorem 1.** *There exists an LP-rounding based randomized algorithm that gives an  $O(\frac{\log \log m_1}{\log m_1} \frac{\log \log m_2}{\log m_2})$ -approximation to **PDB**.*

**A polylogarithmic approximation algorithm for AR.** We present an LP-based approximation for AR. The separation problem for AR is a variant of PDB. However, the objective is a difference of a bilinear and a linear term making it challenging to approximate. Our approach approximates AR directly. In particular, using ideas from our approximation of PDB, we give a compact linear restriction of AR, that is, a linear program whose optimal objective is greater than the optimum of AR, and show that it is a polylogarithmic approximation of AR. In particular, we have the following theorem.

**Theorem 2.** *There exists an LP restriction of AR that gives an  $O(\frac{\log n}{\log \log n} \frac{\log L}{\log \log L})$ -approximation to AR.*

Our bound improves significantly over the state-of-the-art approximation bound of  $O(\frac{L \log n}{\log \log n})$  [14] known for this problem.

We also compare the performance of our algorithm to affine policies. This is one of the most widely used approximation techniques in dynamic robust optimization where the second-stage decision  $\mathbf{y}$  is restricted to be an affine function of the uncertain right-hand side  $\mathbf{h}$ . It is known that the optimal affine policy can be found efficiently (Ben-Tal et al. [2]). We show that our algorithm is significantly faster than finding the optimal affine policy while providing good approximate solutions. Specifically, in randomly generated instances with  $n = m = L = 100$ , the cost of our solution is within 30% of the cost of the optimal affine policy in all of the instances we consider. However, our algorithm is significantly faster terminating in less than 0.1s for all instances. In contrast, it takes 1000s or larger on average to compute the optimal affine policy for  $n > 80$ .

## 2 A polylogarithmic approximation for PDB

In this section, we present an  $O(\frac{\log \log m_1}{\log m_1} \frac{\log \log m_2}{\log m_2})$ -approximation for PDB (Theorem 1). To prove this theorem, we show an interesting structural property for near-optimal solutions for PDB. In particular, we show that there exists a near-optimal solution for PDB that is “near-integral”. Let us define for all  $i \in [n]$ ,

$$\theta_i = \max_{\mathbf{x} \in \mathcal{X}} x_i, \quad \gamma_i = \max_{\mathbf{y} \in \mathcal{Y}} y_i, \quad \zeta_1 = \frac{2 \log m_1}{\log \log m_1} + 2 \quad \text{and} \quad \zeta_2 = \frac{2 \log m_2}{\log \log m_2} + 2.$$

We formally state our structural property in the following lemma.

**Lemma 1. (Structural Property).** *There exists a feasible solution  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  of PDB whose objective value is within  $O(\frac{\log \log m_1}{\log m_1} \frac{\log \log m_2}{\log m_2})$  of the optimum and such that  $\hat{x}_i \in \{0, \frac{\theta_i}{\zeta_1}\}$  and  $\hat{y}_i \in \{0, \frac{\gamma_i}{\zeta_2}\}$  for all  $i \in [n]$ .*

We obtain such a solution satisfying the above property using an LP relaxation of PDB via a randomized rounding approach.

**LP relaxation and Rounding.** We consider the following linear program,

$$z_{\text{LP-PDB}} = \max_{\boldsymbol{\omega} \geq \mathbf{0}} \left\{ \sum_{i=1}^n \theta_i \gamma_i \omega_i \mid \begin{array}{l} \sum_{i=1}^n \theta_i \mathbf{P}_i \omega_i \leq \mathbf{p} \\ \sum_{i=1}^n \gamma_i \mathbf{Q}_i \omega_i \leq \mathbf{q} \end{array} \right\}, \quad (\text{LP-PDB})$$

where  $\mathbf{P}_i$  is the  $i$ -th column of  $\mathbf{P}$  and  $\mathbf{Q}_i$  is the  $i$ -th column of  $\mathbf{Q}$ . We first show that **LP-PDB** is a relaxation of **PDB**.

**Lemma 2.**  $z_{\text{PDB}} \leq z_{\text{LP-PDB}}$ .

*Proof.* Let  $(\mathbf{x}^*, \mathbf{y}^*)$  be an optimal solution of **PDB**. Let  $\boldsymbol{\omega}^*$  be such that  $\omega_i^* = \frac{x_i^*}{\theta_i} \cdot \frac{y_i^*}{\gamma_i}$  for all  $i \in [n]$ . By definition, we have  $x_i^* \leq \theta_i$  and  $y_i^* \leq \gamma_i$  for all  $i \in [n]$ . Hence,

$$\sum_{i=1}^n \theta_i \mathbf{P}_i \omega_i^* = \sum_{i=1}^n \theta_i \mathbf{P}_i \frac{x_i^*}{\theta_i} \frac{y_i^*}{\gamma_i} \leq \sum_{i=1}^n \mathbf{P}_i x_i^* \leq \mathbf{p},$$

and

$$\sum_{i=1}^n \gamma_i \mathbf{Q}_i \omega_i^* = \sum_{i=1}^n \gamma_i \mathbf{Q}_i \frac{x_i^*}{\theta_i} \frac{y_i^*}{\gamma_i} \leq \sum_{i=1}^n \mathbf{Q}_i y_i^* \leq \mathbf{q}.$$

Note that we use the fact that  $\mathbf{P}$  and  $\mathbf{Q}$  are non-negative in the above inequalities. Therefore,  $\boldsymbol{\omega}^*$  is feasible for **LP-PDB** with objective value

$$\sum_{i=1}^n \theta_i \gamma_i \omega_i^* = \sum_{i=1}^n x_i^* y_i^* = z_{\text{PDB}},$$

which concludes the proof.  $\square$

Now, to construct our near-optimal near-integral solution, we consider a randomized rounding approach as described in Algorithm 1. Note that by definition of  $\theta_i$ ,

$$\max_{\boldsymbol{\omega}} \{\omega_i \mid \sum_{j=1}^n \theta_j \mathbf{P}_j \omega_j \leq \mathbf{p}, \boldsymbol{\omega} \geq \mathbf{0}\} = 1.$$

Hence,  $\omega_i^* \leq 1$  for all  $i \in [n]$ . This implies the correctness of Algorithm 1.

In our proof of Lemma 1, we use the following variant of Chernoff bounds.

**Lemma 3.** (Chernoff Bounds [10]).

(a) Let  $\chi_1, \dots, \chi_r$  be independent Bernoulli trials. Denote  $\Xi := \sum_{i=1}^r \epsilon_i \chi_i$  where  $\epsilon_1, \dots, \epsilon_r$  are reals in  $[0, 1]$ . Let  $s > 0$  such that  $\mathbb{E}(\Xi) \leq s$ . Then for any  $\delta > 0$  we have,

$$\mathbb{P}(\Xi \geq (1 + \delta)s) \leq \left( \frac{e^\delta}{(1 + \delta)^{1 + \delta}} \right)^s.$$

---

**Algorithm 1**

---

**Input:**  $\epsilon > 0$ .

**Output:** a feasible solution verifying structural property 1 with probability at least  $1 - \epsilon - o(1)$ .

```
1: Let  $\omega^*$  be an optimal solution of LP-PDB and let  $T = 8\lceil\log \frac{1}{\epsilon}\rceil$ .
2: Initialize  $\max = 0$ ,  $\mathbf{x}^* = \mathbf{0}$  and  $\mathbf{y}^* = \mathbf{0}$ .
3: for  $t = 1, \dots, T$  do
4:   let  $\tilde{\omega}_1, \dots, \tilde{\omega}_n$  be i.i.d. Bernoulli variables with  $\mathbb{P}(\tilde{\omega}_i = 1) = \omega_i^*$  for  $i \in [n]$ .
5:   let  $\hat{x}_i = \theta_i \tilde{\omega}_i / \zeta_1$  and  $\hat{y}_i = \gamma_i \tilde{\omega}_i / \zeta_2$  for  $i \in [n]$ .
6:   if  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  is feasible for PDB and  $\hat{\mathbf{x}}^T \hat{\mathbf{y}} \geq \max$  then
7:     set  $x_i^* = \hat{x}_i$  and  $y_i^* = \hat{y}_i$  for  $i \in [n]$ .
8:     set  $\max = \hat{\mathbf{x}}^T \hat{\mathbf{y}}$ .
9:   end if
10: end for
11: return  $(\mathbf{x}^*, \mathbf{y}^*)$ 
```

---

(b) Let  $\chi_1, \dots, \chi_r$  be independent Bernoulli trials. Denote  $\Xi := \sum_{i=1}^r \epsilon_i \chi_i$  where  $\epsilon_1, \dots, \epsilon_r$  are reals in  $(0, 1]$ . Then for any  $0 < \delta < 1$ ,

$$\mathbb{P}(\Xi \leq (1 - \delta)\mathbb{E}(\Xi)) \leq e^{-\frac{1}{2}\delta^2\mathbb{E}(\Xi)}.$$

For the sake of completeness, we present the proof for these bounds in Appendix A.

*Proof of Lemma 1.* It is sufficient to prove that, with constant probability,  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  constructed at each iteration of Algorithm 1 is a feasible solution of PDB verifying the structural property. In particular, let  $\omega^*$  be an optimal solution of LP-PDB as defined in Algorithm 1. Consider some iteration  $t \in [T]$ . Let  $\tilde{\omega}_1, \dots, \tilde{\omega}_n$  and  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  be as defined in iteration  $t$  of the main loop. We show that the following properties hold with a constant probability,

$$\begin{aligned} \sum_{i=1}^n \mathbf{P}_i \hat{x}_i &\leq \mathbf{p}, \\ \sum_{i=1}^n \mathbf{Q}_i \hat{y}_i &\leq \mathbf{q}, \\ \sum_{i=1}^n \hat{x}_i \hat{y}_i &\geq \frac{z_{\text{LP-PDB}}}{2\zeta_1 \zeta_2}. \end{aligned} \tag{1}$$

First, we have,

$$\begin{aligned}
\mathbb{P}\left(\sum_{i=1}^n \mathbf{P}_i \hat{x}_i > \mathbf{p}\right) &= \mathbb{P}\left(\sum_{i=1}^n \theta_i \mathbf{P}_i \frac{\tilde{\omega}_i}{\zeta_1} > \mathbf{p}\right) \\
&\leq \sum_{j=1}^{m_1} \mathbb{P}\left(\sum_{i=1}^n \theta_i P_{ji} \frac{\tilde{\omega}_i}{\zeta_1} > p_j\right) \\
&= \sum_{j \in [m_1]: p_j > 0} \mathbb{P}\left(\sum_{i=1}^n \frac{\theta_i P_{ji}}{p_j} \tilde{\omega}_i > \zeta_1\right) \\
&\leq \sum_{j \in [m_1]: p_j > 0} \left(\frac{e^{\zeta_1 - 1}}{(\zeta_1)^{\zeta_1}}\right) \\
&\leq m_1 \frac{e^{\zeta_1 - 1}}{(\zeta_1)^{\zeta_1}},
\end{aligned}$$

where the first inequality follows from a union bound on  $m_1$  constraints. The second equality holds because for all  $j \in [m_1]$  such that  $p_j = 0$ , we have

$$\mathbb{P}\left(\sum_{i=1}^n \theta_i P_{ji} \frac{\tilde{\omega}_i}{\zeta_1} > p_j\right) = 0.$$

In fact, if  $p_j = 0$ , by feasibility of  $\boldsymbol{\omega}^*$  in **LP-PDB**, we get

$$\sum_{i=1}^n \theta_i P_{ji} \frac{\omega_i^*}{\zeta_1} = 0,$$

hence, we almost surely have

$$\sum_{i=1}^n \theta_i P_{ji} \frac{\tilde{\omega}_i}{\zeta_1} = 0.$$

The second inequality follows from the Chernoff bounds (a) with  $\delta = \zeta_1 - 1$  and  $s = 1$ . In particular,  $\frac{\theta_i P_{ji}}{p_j} \in [0, 1]$  by definition of  $\theta_i$  for all  $i \in [n]$  and  $j \in [m_1]$  such that  $p_j > 0$  and for all  $j \in [m_1]$  such that  $p_j > 0$  we have,

$$\mathbb{E}\left[\sum_{i=1}^n \frac{\theta_i P_{ji}}{p_j} \tilde{\omega}_i\right] = \sum_{i=1}^n \frac{\theta_i P_{ji}}{p_j} \omega_i^* \leq 1,$$

which holds by feasibility of  $\boldsymbol{\omega}^*$ . Next, note that

$$\frac{e^{\zeta_1 - 1}}{(\zeta_1)^{\zeta_1}} = O\left(\frac{1}{m_1^2}\right).$$

Therefore, there exists a constant  $c > 0$  such that,

$$\mathbb{P}\left(\sum_{i=1}^n \mathbf{P}_i \hat{x}_i > \mathbf{p}\right) \leq \frac{c}{m_1}. \quad (2)$$

By a similar argument there exists a constant  $c' > 0$ , such that

$$\mathbb{P}\left(\sum_{i=1}^n \mathbf{Q}_i \hat{y}_i > \mathbf{q}\right) \leq \frac{c'}{m_2}, \quad (3)$$

Finally we have,

$$\begin{aligned} \mathbb{P}\left(\sum_{i=1}^n \hat{x}_i \hat{y}_i < \frac{1}{2\zeta_1 \zeta_2} z_{\text{LP-PDB}}\right) &= \mathbb{P}\left(\sum_{i=1}^n \frac{\theta_i \gamma_i \tilde{\omega}_i^2}{\zeta_1 \zeta_2} < \frac{1}{2\zeta_1 \zeta_2} \sum_{i=1}^n \theta_i \gamma_i \omega_i^*\right) \\ &= \mathbb{P}\left(\sum_{i=1}^n \frac{\theta_i \gamma_i}{\sum_{j=1}^n \theta_j \gamma_j \omega_j^*} \tilde{\omega}_i < \frac{1}{2}\right) \\ &\leq e^{-\frac{1}{8}}, \end{aligned} \quad (4)$$

where the last inequality follows from Chernoff bounds (b) with  $\delta = 1/2$ . In particular, for  $i \in [n]$ ,

$$\frac{\theta_i \gamma_i}{\sum_{j=1}^n \theta_j \gamma_j \omega_j^*} \leq 1.$$

This is because the unit vector  $\mathbf{e}_i$  is feasible for LP-PDB for all  $i \in [n]$  such as

$$\theta_i \gamma_i \leq z_{\text{LP-PDB}} = \sum_{j=1}^n \theta_j \gamma_j \omega_j^*,$$

and we also have,

$$\mathbb{E}\left[\sum_{i=1}^n \frac{\theta_i \gamma_i}{\sum_{j=1}^n \theta_j \gamma_j \omega_j^*} \tilde{\omega}_i\right] = 1.$$

Combining inequalities (2), (3) and (4) we get that  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  verifies the properties (1) with probability at least

$$1 - \frac{c}{m_1} - \frac{c'}{m_2} - e^{-\frac{1}{8}} = 1 - e^{-\frac{1}{8}} - o(1),$$

which is greater than a constant for  $m_1$  and  $m_2$  large enough. This implies in particular that with positive probability,  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  is feasible and has an objective value that is greater than  $\frac{1}{2\zeta_1 \zeta_2} z_{\text{LP-PDB}}$ . From Lemma 2, this is greater than  $\frac{1}{2\zeta_1 \zeta_2} z_{\text{PDB}}$ . Therefore, with constant probability,  $(\hat{\mathbf{x}}, \hat{\mathbf{y}})$  is a feasible solution of PDB that verifies the structural property.  $\square$

*Proof of Theorem 1.* Let  $(\mathbf{x}^*, \mathbf{y}^*)$  be the output solution of Algorithm 1. Then  $(\mathbf{x}^*, \mathbf{y}^*)$  is such that

$$\mathbf{x}^* \in \mathcal{X}, \quad \mathbf{y}^* \in \mathcal{Y}, \quad \mathbf{x}^{*T} \mathbf{y}^* \geq \frac{1}{2\zeta_1 \zeta_2} \cdot z_{\text{PDB}},$$

if and only if

$$\hat{\mathbf{x}} \in \mathcal{X}, \quad \hat{\mathbf{y}} \in \mathcal{Y}, \quad \hat{\mathbf{x}}^T \hat{\mathbf{y}} \geq \frac{1}{2\zeta_1 \zeta_2} \cdot z_{\text{PDB}},$$



for some iteration  $t$  of the main loop. From our proof of lemma 1, this happens with probability at least

$$\begin{aligned} 1 - (e^{-\frac{1}{8}} - o(1))^T &= 1 - e^{-\frac{T}{8}} - o(1) \\ &= 1 - e^{-\lceil \log \frac{1}{\epsilon} \rceil} - o(1) \\ &\geq 1 - \epsilon - o(1). \end{aligned}$$

Therefore, with probability at least  $1 - \epsilon - o(1)$ , Algorithm 1 outputs a feasible solution of PDB whose objective value is within  $O(\frac{\log \log m_1}{\log m_1} \frac{\log \log m_2}{\log m_2})$  of  $z_{\text{PDB}}$ .  $\square$

**Hardness of the general disjoint bilinear program.** Like packing linear programs, the covering linear programs are known to have logarithmic integrality gaps. Hence, a natural question to ask would be whether similar results can be proven for an equivalent covering version of PDB, i.e., a disjoint bilinear program of the form,

$$z_{\text{cdb}} = \min_{\mathbf{x}, \mathbf{y}} \{ \mathbf{x}^T \mathbf{y} \mid \mathbf{P}\mathbf{x} \geq \mathbf{p}, \mathbf{Q}\mathbf{y} \geq \mathbf{q}, \mathbf{x}, \mathbf{y} \geq \mathbf{0} \} \quad (\text{CDB})$$

where  $\mathbf{P} \in \mathbb{R}_+^{m_1 \times n}$ ,  $\mathbf{Q} \in \mathbb{R}_+^{m_2 \times n}$ ,  $\mathbf{p} \in \mathbb{R}_+^{m_1}$  and  $\mathbf{q} \in \mathbb{R}_+^{m_2}$ . However, the previous analysis does not extend to the covering case. In particular, we have the following inapproximability results.

**Theorem 3.** *The covering disjoint bilinear program CDB is NP-hard to approximate within any finite factor.*

The proof of Theorem 3 uses a polynomial time transformation from the *Monotone Not-All-Equal 3-Satisfiability* (MNAE3SAT) NP-complete problem and is given in Appendix B.

### 3 From disjoint bilinear optimization to two-stage adjustable robust optimization

In this section, we present a polylogarithmic approximation algorithm for AR. In particular, we give a compact linear restriction of AR that provides near-optimal first-stage solutions with cost that is within a factor of  $O(\frac{\log n}{\log \log n} \frac{\log L}{\log \log L})$  of  $z_{\text{AR}}$ . Our proof uses ideas from our approximation of PDB applied to the separation problem  $\mathcal{Q}(\mathbf{x})$ . We begin by discussing our linear restriction below.

#### 3.1 A linear restriction of AR

Recall the two-stage adjustable problem AR,

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbf{c}^T \mathbf{x} + \mathcal{Q}(\mathbf{x}),$$

where for all  $\mathbf{x} \in \mathcal{X}$ ,

$$\mathcal{Q}(\mathbf{x}) = \max_{\mathbf{h} \in \mathcal{U}} \min_{\mathbf{y} \geq \mathbf{0}} \{\mathbf{d}^T \mathbf{y} \mid \mathbf{B}\mathbf{y} \geq \mathbf{h} - \mathbf{A}\mathbf{x}\}.$$

Let us write  $\mathcal{Q}(\mathbf{x})$  in its bilinear form. In particular, we take the dual of the inner minimization problem on  $\mathbf{y}$  to get,

$$\mathcal{Q}(\mathbf{x}) = \max_{\mathbf{h}, \mathbf{z} \geq \mathbf{0}} \left\{ \mathbf{h}^T \mathbf{z} - (\mathbf{A}\mathbf{x})^T \mathbf{z} \mid \begin{array}{l} \mathbf{B}^T \mathbf{z} \leq \mathbf{d} \\ \mathbf{R}\mathbf{h} \leq \mathbf{r} \end{array} \right\}.$$

For the special case where  $\mathbf{A} = \mathbf{0}$ , the optimal first-stage solution is  $\mathbf{x} = \mathbf{0}$  and **AR** reduces to an instance of **PDB**. Therefore, our algorithm for **PDB** gives an  $O(\frac{\log n}{\log \log n} \frac{\log L}{\log \log L})$ -approximation algorithm of **AR** in this special case.

In the general case, the separation problem  $\mathcal{Q}(\mathbf{x})$  is the difference of a bilinear and a linear term. This makes it challenging to approximate  $\mathcal{Q}(\mathbf{x})$ . Instead, we attempt to approximate **AR** directly. In particular, for any  $\mathbf{x} \in \mathcal{X}$  and  $\mathbf{y}_0 \geq \mathbf{0}$  such that  $\mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{y}_0 \geq \mathbf{0}$ , we consider the following linear program:

$$\mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0) = \max_{\boldsymbol{\omega} \geq \mathbf{0}} \left\{ \sum_{i=1}^m (\theta_i \gamma_i - \theta_i \mathbf{a}_i^T \mathbf{x} - \theta_i \mathbf{b}_i^T \mathbf{y}_0) \omega_i \mid \begin{array}{l} \sum_{i=1}^m \theta_i \mathbf{b}_i \omega_i \leq \mathbf{d} \\ \sum_{i=1}^m \gamma_i \mathbf{R}_i \omega_i \leq \mathbf{r} \end{array} \right\},$$

where for all  $i \in [m]$ ,

$$\theta_i := \max_{\mathbf{z}} \{z_i \mid \mathbf{B}\mathbf{z} \leq \mathbf{d}, \mathbf{z} \geq \mathbf{0}\}, \quad \gamma_i := \max_{\mathbf{h}} \{h_i \mid \mathbf{R}\mathbf{h} \leq \mathbf{r}, \mathbf{h} \geq \mathbf{0}\},$$

$\mathbf{a}_i$  and  $\mathbf{b}_i$  are the  $i$ -th row of  $\mathbf{A}$  and  $\mathbf{B}$  respectively and  $\mathbf{R}_i$  is the  $i$ -th column of  $\mathbf{R}$ . The role of  $\mathbf{y}_0$  here is to handle the case when some of the entries of  $\mathbf{A}$  are negative. In fact, our approximation relies on the non-negativity of  $\mathbf{A}\mathbf{x}$ . Since this is not the case in general, we add a second-stage covering term  $\mathbf{B}\mathbf{y}_0$  to  $\mathbf{A}\mathbf{x}$  for some static second-stage solution  $\mathbf{y}_0 \geq \mathbf{0}$  such that  $\mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{y}_0 \geq \mathbf{0}$ . For ease of notation, we use  $\mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0)$  to refer to both the problem and its optimal value. Let

$$\eta := \frac{2 \log n}{\log \log n} + 2, \quad \beta := \frac{2 \log L}{\log \log L} + 2.$$

Similar to **PDB**, we show the following structural property of the separation problem.

**Structural property.** For  $\mathbf{x} \in \mathcal{X}$  and  $\mathbf{y}_0 \geq \mathbf{0}$  such that  $\mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{y}_0 \geq \mathbf{0}$ , there exists a near-integral

solution  $(\mathbf{h}, \mathbf{z}) \in \{0, \frac{\gamma_i}{\beta}\}^m \times \{0, \frac{\theta_i}{\eta}\}^m$  of  $\mathcal{Q}(\mathbf{x})$  such that,

$$\begin{aligned} \sum_{i=1}^m \mathbf{b}_i z_i &\leq \mathbf{d}, \\ \sum_{i=1}^m \mathbf{R}_i h_i &\leq \mathbf{r}, \\ \sum_{i=1}^m h_i z_i - (\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0) z_i &\geq \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta\mathbf{x}, \beta\mathbf{y}_0). \end{aligned} \tag{5}$$

*Proof.* We construct such solution following a similar procedure as in Algorithm 1. In particular, let  $\omega^*$  be an optimal solution of  $\mathcal{Q}^{\text{LP}}(\beta\mathbf{x}, \beta\mathbf{y}_0)$ , consider  $\tilde{\omega}_1, \dots, \tilde{\omega}_m$  i.i.d. Bernoulli random variables such that  $\mathbb{P}(\tilde{\omega}_i = 1) = \omega_i^*$  for all  $i \in [m]$  and let  $(\mathbf{h}, \mathbf{z})$  and let  $h_i = \frac{\gamma_i \tilde{\omega}_i}{\beta}$  and  $z_i = \frac{\theta_i \tilde{\omega}_i}{\eta}$  for all  $i \in [m]$ . Such  $(\mathbf{h}, \mathbf{z})$  satisfies the properties (5) with a constant probability. The proof of this fact is similar to the proof of Lemma 1. In particular, we have,

$$\begin{aligned} \mathbb{P}\left(\sum_{i=1}^m \mathbf{b}_i z_i > \mathbf{d}\right) &= \mathbb{P}\left(\sum_{i=1}^m \theta_i \mathbf{b}_i \frac{\tilde{\omega}_i}{\eta} > \mathbf{d}\right) \\ &\leq \sum_{j=1}^n \mathbb{P}\left(\sum_{i=1}^m \theta_i B_{ij} \frac{\tilde{\omega}_i}{\eta} > d_j\right) \\ &= \sum_{j \in [n]: d_j > 0} \mathbb{P}\left(\sum_{i=1}^m \frac{\theta_i B_{ij}}{d_j} \tilde{\omega}_i > \eta\right) \\ &\leq \sum_{j \in [n]: d_j > 0} \left(\frac{e^{\eta-1}}{\eta^\eta}\right) \\ &\leq n \frac{e^{\eta-1}}{\eta^\eta}, \end{aligned}$$

where the first inequality follows from a union bound on  $n$  constraints. The second equality holds because for all  $j \in [n]$  such that  $d_j = 0$  we have

$$\mathbb{P}\left(\sum_{i=1}^m \theta_i B_{ij} \frac{\tilde{\omega}_i}{\eta} > d_j\right) = 0.$$

Note that  $d_j = 0$  implies

$$\sum_{i=1}^m \theta_i B_{ij} \frac{\omega_i^*}{\eta} = 0,$$

by feasibility of  $\omega^*$  in  $\mathcal{Q}^{\text{LP}}(\beta\mathbf{x}, \beta\mathbf{y}_0)$ . Therefore,

$$\sum_{i=1}^m \theta_i B_{ij} \frac{\tilde{\omega}_i}{\eta} = 0,$$

almost surely. The second inequality follows from the Chernoff bounds (a) with  $\delta = \eta - 1$  and

$s = 1$ . In particular,  $\frac{\theta_i B_{ij}}{d_j} \in [0, 1]$  by definition of  $\theta_i$  for all  $i \in [m]$  and  $j \in [n]$  such that  $d_j > 0$  and for all  $j \in [n]$  such that  $d_j > 0$  we have,

$$\mathbb{E} \left[ \sum_{i=1}^m \frac{\theta_i B_{ij}}{d_j} \tilde{\omega}_i \right] = \sum_{i=1}^m \frac{\theta_i B_{ij}}{d_j} \omega_i^* \leq 1,$$

by feasibility of  $\omega^*$ . Next, note that

$$\frac{e^{\eta-1}}{\eta^\eta} = O\left(\frac{1}{n^2}\right).$$

Therefore, there exists a constant  $c > 0$  such that,

$$\mathbb{P}\left(\sum_{i=1}^m \mathbf{b}_i z_i > \mathbf{d}\right) \leq \frac{c}{n}. \quad (6)$$

By a similar argument there exists a constant  $c' > 0$ ,

$$\mathbb{P}\left(\sum_{i=1}^m \mathbf{R}_i h_i > \mathbf{q}\right) \leq \frac{c'}{L}, \quad (7)$$

Finally we have,

$$\begin{aligned} & \mathbb{P}\left(\sum_{i=1}^m h_i z_i - (\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0) z_i < \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)\right) \\ &= \mathbb{P}\left(\sum_{i=1}^m \frac{\theta_i \gamma_i \tilde{\omega}_i^2}{\eta\beta} - \theta_i (\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0) \frac{\tilde{\omega}_i}{\eta} < \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)\right) \\ &= \mathbb{P}\left(\frac{1}{\eta\beta} \sum_{i=1}^m (\theta_i \gamma_i - \theta_i \mathbf{a}_i^T(\beta \mathbf{x}) + \theta_i \mathbf{b}_i^T(\beta \mathbf{y}_0)) \tilde{\omega}_i < \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)\right). \end{aligned}$$

Let  $\mathcal{I}$  denote the subset of indices  $i \in [m]$  such that

$$(\theta_i \gamma_i - \theta_i \mathbf{a}_i^T(\beta \mathbf{x}) + \theta_i \mathbf{b}_i^T(\beta \mathbf{y}_0)) \geq 0.$$

Since  $\omega^*$  is the optimal solution of the maximization problem  $\mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)$  we can suppose without loss of generality that  $\omega^* = 0$  for all  $i \notin \mathcal{I}$ . Hence,  $\tilde{\omega}_i = 0$  almost surely for all  $i \notin \mathcal{I}$  and we have,

$$\begin{aligned} & \mathbb{P}\left(\sum_{i=1}^m h_i z_i - (\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0) z_i < \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)\right) \\ &= \mathbb{P}\left(\frac{1}{\eta\beta} \sum_{i \in \mathcal{I}} (\theta_i \gamma_i - \theta_i \mathbf{a}_i^T(\beta \mathbf{x}) + \theta_i \mathbf{b}_i^T(\beta \mathbf{y}_0)) \tilde{\omega}_i < \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)\right) \quad (8) \\ &= \mathbb{P}\left(\sum_{i \in \mathcal{I}} \frac{(\theta_i \gamma_i - \theta_i \mathbf{a}_i^T(\beta \mathbf{x}) + \theta_i \mathbf{b}_i^T(\beta \mathbf{y}_0))}{\mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)} \tilde{\omega}_i < \frac{1}{2}\right) \leq e^{-\frac{1}{8}}, \end{aligned}$$

where the last inequality follows from Chernoff bounds (b) with  $\delta = 1/2$ . In particular we have for all  $i \in \mathcal{I}$

$$\frac{(\theta_i \gamma_i - \theta_i \mathbf{a}_i^T(\beta \mathbf{x}) + \theta_i \mathbf{b}_i^T(\beta \mathbf{y}_0))}{\mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)} \leq 1.$$

This is because the unit vector  $\mathbf{e}_i$  is feasible for  $\mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)$  for all  $i \in \mathcal{I}$  such as

$$(\theta_i \gamma_i - \theta_i \mathbf{a}_i^T(\beta \mathbf{x}) + \theta_i \mathbf{b}_i^T(\beta \mathbf{y}_0)) \leq \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0).$$

We also have,

$$\mathbb{E} \left[ \sum_{i \in \mathcal{I}} \frac{(\theta_i \gamma_i - \theta_i \mathbf{a}_i^T(\beta \mathbf{x}) + \theta_i \mathbf{b}_i^T(\beta \mathbf{y}_0))}{\mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)} \tilde{\omega}_i \right] = 1.$$

Combining inequalities (6), (7) and (8) we get that  $(\mathbf{h}, \mathbf{z})$  verifies the properties (5) with probability at least

$$1 - \frac{c}{n} - \frac{c'}{L} - e^{-\frac{1}{8}},$$

which is greater than a constant for  $n$  and  $L$  large enough. Which concludes the proof of the structural property. □

Because of the linear term, the solution given by this structural property is not necessarily a near-optimal solution of  $\mathcal{Q}(\mathbf{x})$  anymore. However, the existence of such solution allows us to bound  $\mathcal{Q}(\mathbf{x})$  as follows.

**Lemma 4.** For  $\mathbf{x} \in \mathcal{X}$  and  $\mathbf{y}_0 \geq \mathbf{0}$  such that  $\mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{y}_0 \geq \mathbf{0}$  we have,

$$\frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0) \leq \mathcal{Q}(\mathbf{x}) \leq \mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0) + \mathbf{d}^T \mathbf{y}_0.$$

**Our Linear Restriction.** Before proving Lemma 4, let us discuss how we derive our linear restriction of AR. In particular, consider the following problem where  $\mathcal{Q}(\mathbf{x})$  is replaced by  $\mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0)$  in the expression of AR:

$$z_{\text{LP-AR}} = \min_{\mathbf{x} \in \mathcal{X}, \mathbf{y}_0 \geq \mathbf{0}} \left\{ \mathbf{c}^T \mathbf{x} + \mathbf{d}^T \mathbf{y}_0 + \mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0) \mid \mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{y}_0 \geq \mathbf{0} \right\}. \quad (9)$$

Note for given  $\mathbf{x}, \mathbf{y}_0$ ,  $\mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0)$  is a maximization LP. Taking its dual and substituting in (9), we get the following LP:

$$\begin{aligned}
z_{\text{LP-AR}} &= \min_{\mathbf{x}, \mathbf{y}_0, \mathbf{y}, \boldsymbol{\alpha}} \quad \mathbf{c}^T \mathbf{x} + \mathbf{d}^T \mathbf{y}_0 + \mathbf{d}^T \mathbf{y} + \mathbf{r}^T \boldsymbol{\alpha} \\
&\text{s.t.} \quad \theta_i \mathbf{a}_i^T \mathbf{x} + \theta_i \mathbf{b}_i^T \mathbf{y}_0 + \theta_i \mathbf{b}_i^T \mathbf{y} + \gamma_i \mathbf{R}_i^T \boldsymbol{\alpha} \geq \theta_i \gamma_i \quad \forall i, \\
&\quad \mathbf{A} \mathbf{x} + \mathbf{B} \mathbf{y}_0 \geq \mathbf{0}, \\
&\quad \mathbf{x} \in \mathcal{X}, \mathbf{y}_0, \mathbf{y} \geq \mathbf{0}, \boldsymbol{\alpha} \geq \mathbf{0}.
\end{aligned} \tag{LP-AR}$$

We claim that **LP-AR** is a restriction of **AR** and gives an  $O(\frac{\log L}{\log \log L} \frac{\log n}{\log \log n})$ -approximation for **AR**.

We first give the proof of Lemma 4.

*Proof of Lemma 4.* First, let  $(\mathbf{h}^*, \mathbf{z}^*)$  be an optimal solution of  $\mathcal{Q}(\mathbf{x})$ . Define  $\boldsymbol{\omega}^*$  such that  $\omega_i^* = \frac{z_i^*}{\theta_i} \cdot \frac{h_i^*}{\gamma_i}$  for all  $i \in [m]$ . Then  $\boldsymbol{\omega}^*$  is feasible for  $\mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0)$  with objective value,

$$\begin{aligned}
\sum_{i=1}^m (\theta_i \gamma_i - \theta_i \mathbf{a}_i^T \mathbf{x} - \theta_i \mathbf{b}_i^T \mathbf{y}_0) \omega_i^* &= \sum_{i=1}^m h_i^* z_i^* - (\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0) \frac{h_i^*}{\gamma_i} z_i^* \\
&\geq \sum_{i=1}^m h_i^* z_i^* - (\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0) z_i^* \\
&= \sum_{i=1}^m h_i^* z_i^* - \mathbf{a}_i^T \mathbf{x} z_i^* - \left( \sum_{i=1}^m \mathbf{b}_i z_i^* \right)^T \mathbf{y}_0 \\
&\geq \mathcal{Q}(\mathbf{x}) - \mathbf{d}^T \mathbf{y}_0,
\end{aligned}$$

where the first inequality follows from the fact that  $\frac{h_i}{\gamma_i} \leq 1$  and  $\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0 \geq 0$  for all  $i \in [m]$ , and the last inequality follows from the fact that  $\sum_{i=1}^m \mathbf{b}_i z_i^* \leq \mathbf{d}$ . Hence  $\mathcal{Q}(\mathbf{x}) - \mathbf{d}^T \mathbf{y}_0 \leq \mathcal{Q}^{\text{LP}}(\mathbf{x}, \mathbf{y}_0)$ .

Now, consider  $(\mathbf{h}, \mathbf{z}) \in \{0, \frac{\gamma_i}{\beta}\}^m \times \{0, \frac{\theta_i}{\eta}\}^m$  satisfying properties (5). The first two properties imply that  $(\mathbf{h}, \mathbf{z})$  is a feasible solution for  $\mathcal{Q}(\mathbf{x})$ . The objective value of this solution is given by,

$$\sum_{i=1}^m h_i z_i - \mathbf{a}_i^T \mathbf{x} z_i \geq \sum_{i=1}^m h_i z_i - (\mathbf{a}_i^T \mathbf{x} + \mathbf{b}_i^T \mathbf{y}_0) z_i \geq \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0).$$

The first inequality holds because  $\mathbf{b}_i^T \mathbf{y}_0 \geq 0$  for all  $i \in [m]$ , and the second inequality follows from the properties (5). Hence,  $\mathcal{Q}(\mathbf{x}) \geq \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta \mathbf{x}, \beta \mathbf{y}_0)$ . □

Now, we are ready to prove Theorem 2.

*Proof of Theorem 2.* We prove the following:

$$z_{\text{AR}} \leq z_{\text{LP-AR}} \leq 3\eta\beta z_{\text{AR}}.$$

Let  $\mathbf{x}_{\text{LP}}^*$ ,  $\mathbf{y}_{0,\text{LP}}^*$  denote an optimal solution of (9). We have,

$$\begin{aligned} z_{\text{LP-AR}} &= \mathbf{c}^T \mathbf{x}_{\text{LP}}^* + \mathbf{d}^T \mathbf{y}_{0,\text{LP}}^* + \mathcal{Q}^{\text{LP}}(\mathbf{x}^*, \mathbf{y}_{0,\text{LP}}^*) \\ &\geq \mathbf{c}^T \mathbf{x}_{\text{LP}}^* + \mathcal{Q}(\mathbf{x}_{\text{LP}}^*) \\ &\geq z_{\text{AR}}, \end{aligned}$$

where the first inequality follows from Lemma 4 and the last inequality follows from the feasibility of  $\mathbf{x}_{\text{LP}}^*$  in AR.

To prove the upper bound for  $z_{\text{LP-AR}}$ , let  $\mathbf{x}^*$  denote an optimal first-stage solution of AR and let  $\mathbf{y}_0^* \in \operatorname{argmin}_{\mathbf{y} \geq \mathbf{0}} \{\mathbf{d}^T \mathbf{y} \mid \mathbf{A}\mathbf{x}^* + \mathbf{B}\mathbf{y} \geq \mathbf{0}\}$ . Since  $\mathbf{0} \in \mathcal{U}$  is a feasible second-stage scenario,

$$\mathbf{c}^T \mathbf{x}^* + \mathbf{d}^T \mathbf{y}_0^* \leq z_{\text{AR}}. \quad (10)$$

Now, we have

$$\begin{aligned} z_{\text{AR}} &\geq \mathcal{Q}(\mathbf{x}^*) \\ &\geq \frac{1}{2\eta\beta} \mathcal{Q}^{\text{LP}}(\beta\mathbf{x}^*, \beta\mathbf{y}_0^*) \\ &= \frac{1}{2\eta\beta} (\beta\mathbf{c}^T \mathbf{x}^* + \beta\mathbf{d}^T \mathbf{y}_0^* + \mathcal{Q}^{\text{LP}}(\beta\mathbf{x}^*, \beta\mathbf{y}_0^*)) - \frac{1}{2\eta} (\mathbf{c}^T \mathbf{x}^* + \mathbf{d}^T \mathbf{y}_0^*) \\ &\geq \frac{1}{2\eta\beta} (\beta\mathbf{c}^T \mathbf{x}^* + \beta\mathbf{d}^T \mathbf{y}_0^* + \mathcal{Q}^{\text{LP}}(\beta\mathbf{x}^*, \beta\mathbf{y}_0^*)) - \frac{1}{2} z_{\text{AR}} \\ &\geq \frac{1}{2\eta\beta} z_{\text{LP-AR}} - \frac{1}{2} z_{\text{AR}}, \end{aligned}$$

where the second inequality follows from Lemma 4, the third inequality follows from (10) and the fact that  $\eta \geq 1$ . For the last inequality, note that  $\beta\mathbf{x}^* \in \mathcal{X}$ ,  $\beta\mathbf{y}_0^* \geq \mathbf{0}$  and  $\beta\mathbf{A}\mathbf{x}^* + \beta\mathbf{B}\mathbf{y}_0^* \geq \mathbf{0}$ . Therefore,  $\beta\mathbf{x}^*$ ,  $\beta\mathbf{y}_0^*$  is a feasible solution for (9). This implies that  $z_{\text{LP-AR}} \leq 3\eta\beta z_{\text{AR}}$ .  $\square$

### 3.2 Numerical experiments

In this subsection, we propose to study the performance of our algorithm by comparing it to finding the optimal affine policy, a widely used technique to approximate AR. We show that our algorithm is significantly faster and provides good approximate solutions. The results of the experiment are given in Table 1.

**Experimental setup.** We consider the same instances as in Ben-Tal et al. [3], namely, we consider instances of AR where  $n = m$ ,  $\mathbf{c} = \mathbf{d} = \mathbf{e}$  and  $\mathbf{A} = \mathbf{B} = \mathbf{I}_m + \mathbf{G}$ , where  $\mathbf{I}_m$  is the identity matrix and  $\mathbf{G}$  is a random normalized Gaussian matrix. We consider the case where  $\mathcal{X} = \mathbb{R}_+^m$  and  $\mathcal{U}$

Table 1: Comparison of the optimal value and the running time in seconds between our algorithm and the optimal affine policy, for different values of  $n = m$  and  $L$ .

n	T <sub>Aff</sub>	T <sub>LP</sub>	$\frac{z_{LP-AR}}{z_{Aff}}$
20	0.57	0.07	1.28
30	3.70	0.08	1.25
40	12.5	0.06	1.25
50	33.2	0.10	1.23
60	76.0	0.03	1.23
70	222	0.03	1.22
80	430	0.04	1.21
90	768	0.06	1.21
100	1790	0.17	1.22

(a) L=20

n	T <sub>Aff</sub>	T <sub>LP</sub>	$\frac{z_{LP-AR}}{z_{Aff}}$
20	1.15	0.05	1.32
30	3.03	0.03	1.30
40	8.58	0.03	1.31
50	28.2	0.02	1.28
60	78.7	0.04	1.27
70	193	0.10	1.27
80	508	0.10	1.26
90	1116	0.06	1.25
100	1714	0.04	1.22

(b) L=50

n	T <sub>Aff</sub>	T <sub>LP</sub>	$\frac{z_{LP-AR}}{z_{Aff}}$
20	1.58	0.07	1.38
30	3.62	0.07	1.35
40	11.5	0.05	1.32
50	29.6	0.08	1.33
60	78.7	0.05	1.30
70	175	0.09	1.27
80	386	0.10	1.28
90	657	0.13	1.27
100	1354	0.12	1.27

(c) L=100

is an intersection of  $L$  budget of uncertainty sets of the form  $\mathcal{U} = \{ \mathbf{h} \in [0, 1]^m \mid \boldsymbol{\omega}_l^T \mathbf{h} \leq 1 \ \forall l \in [L] \}$ , where the weight vectors  $\boldsymbol{\omega}_l$  are normalized Gaussian vectors, i.e.,  $\omega_{l,i} = \frac{|G_{l,i}|}{\sqrt{\sum_i (G_{l,i})^2}}$  for  $\{G_{l,i}\}$  i.i.d. standard Gaussian variables.

We compare the running time of our algorithm in seconds denoted by  $T_{LP}(s)$  with the running time needed to compute the optimal affine policy denoted by  $T_{aff}(s)$ , for different values of  $n = m$  and  $L$ . We also compare the ratio between the optimal value of our linear restriction  $z_{LP-AR}$  and the cost of the optimal affine policy denoted by  $z_{Aff}$ . The results are given in Table 1 and were obtained using Gurobi v9.1.2 on a dual-core laptop with 8Go of RAM and 1.8GHz processor.

**Results.** Table 1 shows that solving our restriction is significantly faster than finding the optimal affine policy. For example, when  $n = m = 100$  computing the optimal affine policy is more than 10000 times slower than LP-AR for all considered values of  $L$ . This is due to the small size of our restriction, in fact, while computing the optimal affine policy requires solving a linear program of  $O(m(n + L))$  variables and constraints (Ben-Tal et al. [2]), our linear restriction has only  $O(n+m+L)$  variables and constraints. Furthermore, the cost of LP-AR stays within approximately 30% of the cost of the optimal affine policy. We also observe numerically that this gap gets smaller when we increase the dimension of our problem and therefore our algorithm gets close to the optimal affine policy for large instances, which are usually the computationally challenging instances for the optimal affine policy.



## 4 Conclusion

In this paper, we consider the class of packing disjoint bilinear programs [PDB](#) and present an LP rounding based approximation algorithm for this problem. In particular, we show existence of a near-optimal near-integral solution for [PDB](#). We also give an LP relaxation for the problem and use a randomized rounding approach to round the optimal solution of the LP relaxation to obtain the approximate solution with the desired guarantees. We apply our ideas to the two-stage adjustable problem [AR](#) whose separation problem is a variant of [PDB](#). While a direct application of the approximation algorithm for [PDB](#) does not work for [AR](#), we derive a linear restriction of [AR](#) based on similar ideas and give a polylogarithmic approximation for [AR](#). We compare our algorithm with the widely used affine policies and show that it is significantly faster and provides near-optimal solutions.

## References

- [1] Chaithanya Bandi and Dimitris Bertsimas. “Tractable stochastic analysis in high dimensions via robust optimization”. In: *Mathematical Programming* 134.1 (2012), pp. 23–70.
- [2] A. Ben-Tal et al. “Adjustable Robust Solutions of Uncertain Linear Programs”. In: *Mathematical Programming* 99 (2004), pp. 351–376.
- [3] Aharon Ben-Tal, Omar El Housni, and Vineet Goyal. “A tractable approach for designing piecewise affine policies in two-stage adjustable robust optimization”. In: *Mathematical Programming* 182 (2020), pp. 57–102.
- [4] D. Bertsimas and Vineet Goyal. “On the power and limitations of affine policies in two-stage adaptive optimization”. In: *Mathematical Programming* 134 (2012), pp. 491–531.
- [5] Dimitris Bertsimas and Hoda Bidkhori. “On the performance of affine policies for two-stage adaptive optimization: a geometric perspective”. In: *Mathematical Programming* 153.2 (2015), pp. 577–594.
- [6] Dimitris Bertsimas, David B. Brown, and Constantine Caramanis. “Theory and Applications of Robust Optimization”. In: *SIAM Rev.* 53.3 (2011), pp. 464–501.
- [7] Dimitris Bertsimas and Frans Ruiters. “Duality in Two-Stage Adaptive Linear Optimization: Faster Computation and Stronger Bounds”. In: *INFORMS Journal on Computing* 28 (2016), pp. 500–511.
- [8] Dimitris Bertsimas and Melvyn Sim. “The Price of Robustness”. In: *Operations Research* 52 (2004), pp. 35–53.
- [9] Xi Chen, Xiaotie Deng, and Shang-Hua Teng. “Settling the Complexity of Computing Two-Player Nash Equilibria”. In: *J. ACM* 56.3 (2009), pp. 1–57.

- [10] Herman Chernoff. “A Measure of Asymptotic Efficiency for Tests of a Hypothesis Based on the sum of Observations”. In: *The Annals of Mathematical Statistics* 23.4 (1952), pp. 493–507.
- [11] Ante Čustić et al. “The bilinear assignment problem: complexity and polynomially solvable special cases”. In: *Mathematical Programming* 166.1-2 (2017), pp. 185–205.
- [12] K. Dhamdhere et al. “How to pay, come what may: approximation algorithms for demand-robust covering problems”. In: *46th Annual IEEE Symposium on Foundations of Computer Science (FOCS’05)*. 2005, pp. 367–376.
- [13] Omar El Housni and Vineet Goyal. “Beyond Worst-Case: A Probabilistic Analysis of Affine Policies in Dynamic Optimization”. In: *Proceedings of the 31st International Conference on Neural Information Processing Systems*. 2017, pp. 4759–4767.
- [14] Omar El Housni and Vineet Goyal. “On the Optimality of Affine Policies for Budgeted Uncertainty Sets”. In: *Mathematics of Operations Research* 46.2 (2021), pp. 674–711.
- [15] Omar El Housni, Vineet Goyal, and David Shmoys. “On the power of static assignment policies for robust facility location problems”. In: *International Conference on Integer Programming and Combinatorial Optimization*. 2021, pp. 252–267.
- [16] Uriel Feige et al. “Robust Combinatorial Optimization with Exponential Scenarios”. In: *Integer Programming and Combinatorial Optimization*. Ed. by Matteo Fischetti and David P. Williamson. 2007, pp. 439–453.
- [17] Koorosh Firouzbakht, Guevara Noubir, and Masoud Salehi. “Linearly Constrained Bimatrix Games in Wireless Communications”. In: *IEEE Transactions on Communications* 64.1 (2016), pp. 429–440.
- [18] Arthur Geoffrion. “Generalized Benders Decomposition”. In: *Journal of Optimization Theory and Applications* 10 (1972), pp. 237–260.
- [19] Chrysanthos Gounaris et al. “An Adaptive Memory Programming Framework for the Robust Capacitated Vehicle Routing Problem”. In: *Transportation Science* 50 (2014), pp. 1139–1393.
- [20] Anupam Gupta, Viswanath Nagarajan, and R. Ravi. “Robust and MaxMin Optimization under Matroid and Knapsack Uncertainty Sets”. In: *ACM Trans. Algorithms* 12.1 (2015), pp. 1–21.
- [21] Anupam Gupta, Viswanath Nagarajan, and Ramamoorthi Ravi. “Thresholded covering algorithms for robust and max–min optimization”. In: *Mathematical Programming* 146 (2014), pp. 583–615.
- [22] Akshay Gupte et al. “Relaxations and discretizations for the pooling problem”. In: *Journal of Global Optimization* 67 (2017), pp. 631–669.

- [23] I. Harjunkski et al. “Different transformations for solving non-convex trim-loss problems by MINLP”. In: *European Journal of Operational Research* 105 (1998), pp. 594–603.
- [24] Hiroshi Konno. “A cutting plane algorithm for solving bilinear programs”. In: *Mathematical Programming* 11 (1976), pp. 14–27.
- [25] Churlzu Lim and J. Cole Smith. “Algorithms for discrete and continuous multicommodity flow network interdiction problems”. In: *IIE Transactions* 39.1 (2007), pp. 15–26.
- [26] O.L Mangasarian and H Stone. “Two-person nonzero-sum games and quadratic programming”. In: *Journal of Mathematical Analysis and Applications* 9.3 (1964), pp. 348–355.
- [27] Steffen Rebennack, Artyom Nahapetyan, and Panos Pardalos. “Bilinear modeling solution approach for fixed charge network flow problems”. In: *Optimization Letters* 3 (2009), pp. 347–355.
- [28] Thomas J. Schaefer. “The Complexity of Satisfiability Problems”. In: *Proceedings of the Tenth Annual ACM Symposium on Theory of Computing*. STOC ’78. San Diego, California, USA: Association for Computing Machinery, 1978, pp. 216–226.
- [29] Richard Soland. “Optimal Facility Location with Concave Costs”. In: *Operations Research* 22 (1974), pp. 373–382.
- [30] Tran Vu Thieu. “A note on the solution of bilinear programming problems by reduction to concave minimization”. In: *Math. Program.* 41.1-3 (1988), pp. 249–260.
- [31] Harish Vaish and C. M. Shetty. “The bilinear programming problem”. In: *Naval Research Logistics Quarterly* 23.2 (1976), pp. 303–309.
- [32] Guanglin Xu and Samuel Burer. “A copositive approach for two-stage adjustable robust optimization with uncertain right-hand sides”. In: *Computational Optimization and Applications* 70.1 (2018), pp. 33–59.
- [33] Jianzhe Zhen, Dick den Hertog, and M. Sim. “Adjustable robust optimization via Fourier-Motzkin elimination”. In: *Operations Research* 66.4 (2018), pp. 1086–1100.
- [34] Jianzhe Zhen et al. “Disjoint bilinear programming: a two-stage robust optimization perspective”. In: *Optimization Online* (2018). URL: [www.optimization-online.org/DB\\_HTML/2018/06/6685.html](http://www.optimization-online.org/DB_HTML/2018/06/6685.html).

## Appendix A Chernoff bounds

**Proof of Chernoff bounds (a).** From Markov’s inequality we have for all  $t > 0$ ,

$$\mathbb{P}(\Xi \geq (1 + \delta)s) = \mathbb{P}(e^{t\Xi} \geq e^{t(1+\delta)s}) \leq \frac{\mathbb{E}(e^{t\Xi})}{e^{t(1+\delta)s}}.$$

Denote  $p_i$  the parameter of the Bernoulli  $\chi_i$ . By independence, we have

$$\mathbb{E}(e^{t\Xi}) = \prod_{i=1}^r \mathbb{E}(e^{t\epsilon_i \chi_i}) = \prod_{i=1}^r (p_i e^{t\epsilon_i} + 1 - p_i) \leq \prod_{i=1}^r \exp(p_i(e^{t\epsilon_i} - 1))$$

where the inequality holds because  $1 + x \leq e^x$  for all  $x \in \mathbb{R}$ . By taking  $t = \ln(1 + \delta) > 0$ , the right hand side becomes

$$\prod_{i=1}^r \exp(p_i((1 + \delta)^{\epsilon_i} - 1)) \leq \prod_{i=1}^r \exp(p_i \delta \epsilon_i) = \exp(\delta \cdot \mathbb{E}(\Xi)) \leq e^{\delta s},$$

where the first inequality holds because  $(1 + x)^\epsilon \leq 1 + \alpha x$  for any  $x \geq 0$  and  $\epsilon \in [0, 1]$  and the second one because  $s \geq \mathbb{E}(\Xi) = \sum_{i=1}^r \epsilon_i p_i$ . Hence, we have

$$\mathbb{E}(e^{t\Xi}) \leq e^{\delta s}.$$

On the other hand,

$$e^{t(1+\delta)s} = (1 + \delta)^{(1+\delta)s}.$$

Therefore,

$$\mathbb{P}(\Xi \geq (1 + \delta)s) \leq \left( \frac{e^{\delta s}}{(1 + \delta)^{(1+\delta)s}} \right) = \left( \frac{e^\delta}{(1 + \delta)^{1+\delta}} \right)^s.$$

□

**Proof of Chernoff bounds (b).** We show equivalently that

$$\mathbb{P}(\Xi \leq (1 - \delta)\mathbb{E}(\Xi)) \leq e^{\frac{-\delta^2 \mathbb{E}(\Xi)}{2}}.$$

From Markov's inequality we have for all  $t < 0$ ,

$$\mathbb{P}(\Xi \leq (1 - \delta)\mathbb{E}(\Xi)) = \mathbb{P}(e^{t\Xi} \geq e^{t(1-\delta)\mathbb{E}(\Xi)}) \leq \frac{\mathbb{E}(e^{t\Xi})}{e^{t(1-\delta)\mathbb{E}(\Xi)}}.$$

Denote  $p_i$  the parameter of the Bernoulli  $\chi_i$ . By independence, we have

$$\mathbb{E}(e^{t\Xi}) = \prod_{i=1}^r \mathbb{E}(e^{t\epsilon_i \chi_i}) = \prod_{i=1}^r (p_i e^{t\epsilon_i} + 1 - p_i) \leq \prod_{i=1}^r \exp(p_i(e^{t\epsilon_i} - 1)),$$

where the inequality holds because  $1 + x \leq e^x$  for all  $x \in \mathbb{R}$ . We take  $t = \ln(1 - \delta) < 0$ . We have

$t \leq -\delta$ , hence,

$$\prod_{i=1}^r \exp(p_i(e^{t\epsilon_i} - 1)) = \prod_{i=1}^r \exp(p_i((1 - \delta)^{\epsilon_i} - 1)) \leq \prod_{i=1}^r \exp(-p_i\delta\epsilon_i),$$

where the inequality holds because  $(1 - x)^\epsilon \leq 1 - \epsilon x$  for any  $0 < x < 1$  and  $\epsilon \in [0, 1]$ . Therefore,

$$\mathbb{E}(e^{t\Xi}) \leq \prod_{i=1}^r \exp(-p_i\delta\epsilon_i) = e^{-\delta\mathbb{E}(\Xi)}.$$

On the other hand,

$$e^{t(1-\delta)\mathbb{E}(\Xi)} = (1 - \delta)^{(1-\delta)\mathbb{E}(\Xi)}.$$

Therefore,

$$\mathbb{P}(\Xi \leq (1 - \delta)\mathbb{E}(\Xi)) \leq \left( \frac{e^{-\delta\mathbb{E}(\Xi)}}{(1 - \delta)^{(1-\delta)\mathbb{E}(\Xi)}} \right) = \left( \frac{e^{-\delta}}{(1 - \delta)^{1-\delta}} \right)^{\mathbb{E}(\Xi)}.$$

Finally, we have for any  $0 < \delta < 1$ ,

$$\ln(1 - \delta) \geq -\delta + \frac{\delta^2}{2}$$

which implies

$$(1 - \delta) \cdot \ln(1 - \delta) \geq -\delta + \frac{\delta^2}{2}$$

and consequently

$$\left( \frac{e^{-\delta}}{(1 - \delta)^{1-\delta}} \right)^{\mathbb{E}(\Xi)} \leq e^{\frac{-\delta^2\mathbb{E}(\Xi)}{2}}.$$

□

## Appendix B Proof of Theorem 3

Our proof uses a polynomial time transformation from the *Monotone Not-All-Equal 3-satisfiability* (MNAE3SAT) NP-complete problem (Schaefer [28]). In the (MNAE3SAT), we are given a collection of Boolean variables and a collection of clauses, each of which combines three variables. (MNAE3SAT) is the problem of determining if there exists a truth assignment where each clause has at least one true and one false literal. This is a subclass of the *Not-All-Equal 3-satisfiability* problem where the variables are never negated.

Consider an instance  $\mathcal{I}$  of (MNAE3SAT), let  $\mathcal{V}$  be the set of variables of  $\mathcal{I}$  and let  $\mathcal{C}$  be the set of clauses. Let  $\mathbf{A} \in \{0, 1\}^{|\mathcal{C}| \times |\mathcal{V}|}$  be the variable-clause incidence matrix such that for every variable  $v \in \mathcal{V}$  and clause  $c \in \mathcal{C}$  we have  $A_{cv} = 1$  if and only if the variable  $v$  belongs to the clause  $c$ . We

consider the following instance  $\mathcal{I}'$  of **CDB**,

$$\begin{aligned} \min_{\mathbf{x}, \mathbf{y}} \quad & \sum_{v \in \mathcal{V}} x_v y_v \\ \text{s.t.} \quad & \mathbf{A}\mathbf{x} \geq \mathbf{e}, \quad \mathbf{x} \geq \mathbf{0} \\ & \mathbf{A}\mathbf{y} \geq \mathbf{e}, \quad \mathbf{y} \geq \mathbf{0}. \end{aligned} \tag{\mathcal{I}'}$$

Let  $z_{\mathcal{I}'}$  denote the optimum of  $\mathcal{I}'$ . We show that  $\mathcal{I}$  has a truth assignment where each clause has at least one true and one false literal if and only if  $z_{\mathcal{I}'} = 0$ .

First, suppose  $z_{\mathcal{I}'} = 0$ . Let  $(\mathbf{x}, \mathbf{y})$  be an optimal solution of  $\mathcal{I}'$ . Let  $\tilde{\mathbf{x}}$  such that  $\tilde{x}_v = \mathbf{1}_{\{x_v > 0\}}$  for all  $v \in \mathcal{V}$ . We claim that  $(\tilde{\mathbf{x}}, \mathbf{e} - \tilde{\mathbf{x}})$  is also an optimal solution of  $\mathcal{I}$ . In fact, for all  $c \in \mathcal{C}$ , we have,

$$\sum_{v \in \mathcal{V}} A_{cv} x_v \geq 1,$$

hence,

$$\sum_{v \in \mathcal{V}} A_{cv} \min(x_v, 1) \geq 1,$$

since the entries of  $\mathbf{A}$  are in  $\{0, 1\}$ . Therefore,

$$\sum_{v \in \mathcal{V}} A_{cv} \tilde{x}_v \geq \sum_{v \in \mathcal{V}} A_{cv} \min(x_v, 1) \geq 1.$$

Finally,  $\mathbf{A}\tilde{\mathbf{x}} \geq \mathbf{e}$ . Similarly, for all  $c \in \mathcal{C}$  we have,

$$\sum_{v \in \mathcal{V}} A_{cv} y_v \geq 1,$$

hence,

$$\sum_{v \in \mathcal{V}} A_{cv} \min(y_v, 1) \geq 1,$$

since the entries of  $\mathbf{A}$  are in  $\{0, 1\}$ . Therefore,

$$\sum_{v \in \mathcal{V}} A_{cv} (1 - \tilde{x}_v) \geq \sum_{v \in \mathcal{V}} A_{cv} (1 - \tilde{x}_v) \min(y_v, 1) = \sum_{v \in \mathcal{V}} A_{cv} \min(y_v, 1) \geq 1.$$

where the equality follows from the fact that  $y_v = 0$  for all  $v$  such that  $\tilde{x}_v = 1$ . Note that  $\sum_{v \in \mathcal{V}} x_v y_v = z_{\mathcal{I}'} = 0$ . This implies that  $\mathbf{A}(\mathbf{e} - \tilde{\mathbf{x}}) \geq \mathbf{e}$ . Therefore,  $(\tilde{\mathbf{x}}, \mathbf{e} - \tilde{\mathbf{x}})$  is a feasible solution of  $\mathcal{I}'$  with objective value  $\sum_{v \in \mathcal{V}} \tilde{x}_v (1 - \tilde{x}_v) = 0 = z_{\mathcal{I}'}$ . It is therefore an optimal solution. Consider now the truth assignment where a variable  $v$  is set to be true if and only if  $\tilde{x}_v = 1$ . We show that each clause in such assignment has at least one true and one false literal. In particular, consider a

clause  $c \in \mathcal{C}$ , since,

$$\sum_{v \in \mathcal{V}} A_{cv} \tilde{x}_v \geq 1,$$

there must be a variable  $v$  belonging to the clause  $c$  such that  $\tilde{x}_v = 1$ . Similarly, since

$$\sum_{v \in \mathcal{V}} A_{cv} (1 - \tilde{x}_v) \geq 1,$$

there must be a variable  $v$  belonging to the clause  $c$  such that  $\tilde{x}_v = 0$ . This implies that our assignment is such that the clause  $c$  has at least one true and one false literal.

Conversely, suppose there exists a truth assignment of  $\mathcal{I}$  where each clause has at least one false and one true literal. We show that  $z_{\mathcal{I}'} = 0$ . In particular, define  $\mathbf{x}$  such that for all  $v \in \mathcal{V}$  we have  $x_v = 1$  if and only if  $v$  is assigned true. We have for all  $c \in \mathcal{C}$ ,

$$\sum_{v \in \mathcal{V}} A_{cv} x_v \geq 1,$$

since at least one of the variables of  $c$  is assigned true. And,

$$\sum_{v \in \mathcal{V}} A_{cv} (1 - x_v) \geq 1,$$

since at least one of the variables of  $c$  is assigned false. Hence,  $(\mathbf{x}, \mathbf{e} - \mathbf{x})$  is feasible for  $\mathcal{I}'$  with objective value  $\sum_{v \in \mathcal{V}} x_v (1 - x_v) = 0$ . It is therefore an optimal solution and  $z_{\mathcal{I}'} = 0$ .

If there exists a polynomial time algorithm approximating **CDB** to some finite factor, such an algorithm can be used to decide in polynomial time whether  $z_{\mathcal{I}'} = 0$  or not, which is equivalent to solving  $\mathcal{I}$  in polynomial time; a contradiction.

□