

Individual Representation in Approval-Based Committee Voting*

Markus Brill,¹ Jonas Israel,¹ Evi Micha,² Jannik Peters¹

¹ Research Group Efficient Algorithms, TU Berlin, Germany

² Department of Computer Science, University of Toronto, Canada

Abstract

When selecting multiple candidates based on approval preferences of agents, the proportional representation of agents’ opinions is an important and well-studied desideratum. Existing criteria for evaluating the representativeness of outcomes focus on *groups* of agents and demand that sufficiently large and cohesive groups are “represented” in the sense that candidates approved by *some* group members are selected. Crucially, these criteria say nothing about the representation of *individual* agents, even if these agents are members of groups that deserve representation. In this paper, we formalize the concept of *individual representation (IR)* and explore to which extent, and under which circumstances, it can be achieved. We show that checking whether an IR outcome exists is computationally intractable, and we verify that all common approval-based voting rules may fail to provide IR even in cases where this is possible. We then focus on domain restrictions and establish an interesting contrast between “voter interval” and “candidate interval” preferences. This contrast can also be observed in our experimental results, where we analyze the attainability of IR for realistic preference profiles.

1 Introduction

We consider the problem of selecting a fixed-size subset of candidates (a so-called *committee*) based on the approval preferences of agents. This problem has been extensively studied in recent years (Lackner and Skowron 2021) and has a wide variety of applications, including political elections (Brill et al. 2020), recommender systems (Skowron et al. 2017), medical diagnostic decision-making (Gangl et al. 2019), and participatory budgeting (Peters, Pierczyński, and Skowron 2021).

A central concern in committee voting is the principle of *proportional representation*, which states that the agents’ interests and opinions should be reflected proportionately in the committee. While proportional representation is intuitive to understand in scenarios such as apportioning parliamentary seats based on vote shares (Balinski and Young 1982; Pukelsheim 2014), it is less straightforward to formalize in the context of approval-based committee elections.

Indeed, the literature has defined a number of different concepts aiming to capture proportional representation.

Most (if not all) of these approval-based proportionality notions focus on the representation of *groups* of agents. Specifically, it is usually required that each sufficiently large group of agents is “represented” in the committee,¹ where the interpretation of “representation” differs across different notions. For example, *extended justified representation* (Aziz et al. 2017) prescribes that there exists at least one agent in the group approving a certain number of committee members, whereas *proportional justified representation* (Sánchez-Fernández et al. 2017) demands that there are sufficiently many committee members that are each approved by at least one agent in the group. Notably, neither definition comprises any representation requirements for *individual* agents in a group. Thus, a group may count as “represented” even though some agents in the group do not approve a single committee member.

In this paper, we adopt an *individualistic* point of view: our goal is to provide all members of a voter group equal guarantees. Intuitively, when a population consists of n agents and a committee of k representatives is elected, we expect every cohesive voter group of size $\ell \cdot n/k$ to be represented by ℓ representatives in the committee; thus, each individual group member might reasonably hope that at least ℓ candidates represent her in the committee. This notion, which we call *individual representation*, is aligned with the notion of “individual fairness” that was recently introduced in clustering (and in particular in facility location problems) by Jung, Kannan, and Lutz (2020): there, each individual expects to be served by a facility in distance proportional to the radius of the ball that captures its n/k closest neighbors, where n is the number of individuals and k is the number of facilities.

Individual representation, as defined in this paper, is a strengthening of a notion called *semi-strong justified representation* by Aziz et al. (2017). The latter property requires that all members of a group are represented in the committee *at least once*, given that the group is large and cohesive

*A short version of this paper appears in the proceedings of AAAI-2022 (Brill et al. 2022).

¹Often, there is also a condition on the “cohesiveness” of the group, stating that the approval preferences of group members need to be sufficiently aligned.

enough. Individual representation strengthens this requirement by demanding that all members of cohesive groups are represented multiple times (in proportion to the group size). Aziz et al. (2017) observed that semi-strong JR cannot be provided in all instances; this immediately implies our stronger requirement is not universally attainable either.

Our contribution. In this paper, we systematically study *individual representation (IR)*. Notwithstanding the observation that IR demands cannot always be met, we clarify how IR relates to existing axioms and we show that a large range of common approval-based committee voting rules can fail to provide IR even in cases where IR is achievable. We observe that even committees *approximating* IR may fail to exist. Moreover, we answer a question by Aziz et al. (2017) by showing that it is computationally intractable to decide whether a given instance admits a committee providing semi-strong JR or individual representation. We then turn our attention to *restricted domains* of preferences (Elkind and Lackner 2015; Yang 2019) and demonstrate that positive results can be obtained. Doing so, we uncover a striking difference between the *candidate interval* and *voter interval* domains: whereas the former restriction does not admit any non-trivial approximation of IR, we devise an efficient algorithm for selecting committees approximating IR for the latter. This is surprising insofar as these two domain restrictions often exhibit similar behavior (Pierczyński and Skowron 2021; Terzopoulou, Karpov, and Obraztsova 2021). Finally, we experimentally study how often IR is achievable for a wide variety of generated preference data, and how often established voting rules select IR outcomes.

2 Preliminaries

For $t \in \mathbb{N} = \{1, 2, \dots\}$, let $[t]$ denote the set $\{1, 2, \dots, t\}$.

Let $N = [n]$ be a set of n voters (or agents) and $C = \{c_1, \dots, c_m\}$ be a set of m candidates. Each voter $i \in N$ approves a subset $A_i \subseteq C$ of candidates. An (approval) profile $A = (A_1, \dots, A_n)$ contains the approval set A_i for each $i \in N$.

Given a committee size $k \in [m]$, we want to select a subset $W \subseteq C$ of size $|W| = k$, referred to as a *committee*. We call (A, k) an *approval-based committee (ABC) election*. An ABC voting rule takes as input an ABC election (A, k) and outputs one or more committees of size k .

As is standard in the ABC election literature, we assume that voters only care about the number of approved candidates in the committee, i.e., voter i evaluates a committee W by $|A_i \cap W|$. Given a subset $S \subseteq C$ of candidates, we let $N(S)$ denote the set of voters who approve all candidates in S , i.e., $N(S) = \{i \in N : S \subseteq A_i\}$.

Given an ABC election (A, k) and $\ell \in \mathbb{N}$, we call a group $V \subseteq N$ of voters ℓ -cohesive if

$$|V| \geq \ell \cdot \frac{n}{k} \quad \text{and} \quad \left| \bigcap_{i \in V} A_i \right| \geq \ell.$$

The following representation notions are due to Aziz et al. (2017). A committee $W \subseteq C$ of size k provides

- *justified representation (JR)* if for each 1-cohesive group $V \subseteq N$, there is a voter $i \in V$ with $|W \cap A_i| \geq 1$;
- *extended justified representation (EJR)* if for each $\ell \in \mathbb{N}$ and each ℓ -cohesive group $V \subseteq N$, there is a voter $i \in V$ with $|W \cap A_i| \geq \ell$; and
- *core stability* if for each group $V \subseteq N$ (independent of V being ℓ -cohesive) and $S \subseteq C$ with $|S| \leq |V| \cdot \frac{n}{k}$ there is a voter $i \in V$ with $|W \cap A_i| \geq |S \cap A_i|$.

All of these notions have in common that they consider a group of voters “represented” as long as at least one voter in the group is sufficiently represented. This point of view might be hard to justify in many contexts. In the following section, we present our approach to representation that takes into account every voter in a group individually.

3 Individual Representation

In this section, we define the main concept of this paper: individual representation. This notion builds on the idea of (semi-)strong justified representation defined by Aziz et al. (2017) and the notion of individual fairness in clustering as defined by Jung, Kannan, and Lutz (2020).

Similarly to the proportionality notions defined above, we assume that a voter $i \in N$ deserves some representation in an ABC election if i can find enough other voters who all approve a subset of candidates in common. This follows the rationale that everyone in a group of voters that makes up a sizable part of the electorate and can come to an agreement on how (part of) the committee ought to be filled, should be represented accordingly.

Given an ABC election (A, k) , we determine the number of seats that voter $i \in N$ can justifiably demand as

$$f_i := \max_{S \subseteq A_i} \{|S| : |N(S)| \geq |S| \cdot n/k\}.$$

In words, f_i is the largest value f such that voter i can find enough like-minded voters to form an f -cohesive group.

Definition 1 (Individual Representation). *Given an ABC election (A, k) , a committee $W \subseteq C$ of size at most k provides individual representation (IR) if $|W \cap A_i| \geq f_i$ for all voters $i \in N$.*

When only requiring $|W \cap A_i| \geq 1$ for every voter with $f_i \geq 1$ we get *semi-strong justified representation (semi-strong JR)* as defined by Aziz et al. (2017). The authors of that paper provide an example showing that semi-strong JR committees do not always exist. Since individual representation clearly is a more demanding property, it immediately follows that *IR committees* (i.e., committees providing IR) do not need to exist either.

IR not only implies semi-strong JR, but also EJR. More precisely, every IR committee also provides EJR. To build intuition on how our notion differs from established ones, and on how it leads to the election of committees that are “fair” from an individual voter perspective, consider the following examples, illustrated in Figure 1.

Example 1. *The first part of Figure 1 shows an approval profile with 8 voters and 3 candidates. Assuming $k = 2$,*




Figure 1: Two profiles admitting IR committees that are not identified by common voting rules or other representation axioms. Voters correspond to integers and approve all candidates placed above them.

every voter $i \in N$ has $f_i = 1$. Thus, the only committee providing IR is $W = \{c_1, c_2\}$, which represents every voter once and, moreover, satisfies core stability. However, both $W' = \{c_1, c_3\}$ and $W'' = \{c_2, c_3\}$ are core stable as well (and in fact would be selected when choosing a committee maximizing social welfare). Many common ABC voting rules would select either W' or W'' (see Proposition 4). One can argue that committee W is a much “fairer” or “more representative” choice in this example.

Example 2. The second part of Figure 1 shows an approval profile with 12 voters and 10 candidates. For $k = 6$ we have $f_i = 1$ for $i \in \{1, \dots, 4\}$ and $f_i = 2$ for $i \in \{5, \dots, 12\}$. Here, the only committee providing IR is $W = \{c_1, c_2, c_3, c_4, c_9, c_{10}\}$ representing each of the first four voters once, while representing all other voters at least twice. This committee is not core stable, because the voter group $\{5, \dots, 12\}$ would prefer $\{c_5, c_6, c_7, c_8\}$ to W . In order to appreciate the IR committee W , consider voters 1 to 4 and observe that these voters are completely “symmetric.” Hence, from an “equal treatment of equals” perspective, if one of them is represented by an approved candidate in the committee, the same should hold for the others. In fact, the only core-stable committees that provide this kind of symmetry are $\{c_5, \dots, c_{10}\}$, in which one third of the electorate is not represented at all, or committees containing only two candidates among c_5 to c_8 . In the latter case, by noticing that voters 9 to 12 are “symmetric” as well, we can argue similarly as above that they are not treated equally. Thus, the committee W that uniquely provides IR can be considered the “fairest” choice under an individualistic point of view.

The instance in Example 2 shows that core stability and individual representation are incompatible, in the strong sense that the (nonempty) set of IR committees and the (nonempty) set of core-stable committees are disjoint.




Figure 2: Relationships between individual representation (IR) and the representation notions introduced by Aziz et al. (2017). An arrow from X to Y depicts that X implies Y (if X is possible in the given instance).

Proposition 1. *IR is incompatible with core stability.*

Next, we show a further incompatibility result.

Proposition 2. *Semi-strong JR is incompatible with EJR.*

Proof. Consider an ABC election with $n = 8$, $k = 4$ and the following approval profile: $A_1 = \dots = A_4 = \{c_1, c_2\}$, $A_5 = \{c_3, c_4, c_5\}$, $A_6 = \{c_3\}$, $A_7 = \{c_4\}$, and $A_8 = \{c_5\}$. As $f_6 = f_7 = f_8 = 1$, every committee W that provides semi-strong JR must satisfy $\{c_3, c_4, c_5\} \subseteq W$. But then we have $|W \cap A_i| < 2$ for all $i \in [4]$, even though these four voters form a 2-cohesive group. \square

A graphical representation of these results can be found in Figure 2.

Approximate notions. As we mentioned above, IR is not always attainable. The immediate next question is whether we can guarantee IR in an approximate sense. To study this question, we introduce the notion of (α, β) -individual representation, which uses additive and multiplicative approximation parameters.

Definition 2 ((α, β) -Individual Representation). *Given an ABC election (A, k) , a committee W provides (α, β) -individual representation $((\alpha, \beta)$ -IR) if for every voter $i \in N$, $\alpha \cdot |A_i \cap W| + \beta \geq f_i$, with $\alpha \geq 1$ and $\beta \geq 0$.*

Unfortunately, non-trivial approximation guarantees are impossible to obtain without restricting the set of profiles.

Theorem 3. *There exists an instance that does not admit an (α, β) -IR committee for $\beta < k - 1$, and any $\alpha \geq 1$.*

Proof. Assume that $n = k \cdot (k + 1)$. Then it holds that $n/k = k + 1 > k$. Consider a profile in which for each voter $i \in [n/k]$, $A_i = \{c_{(k-1) \cdot (i-1) + 1}, \dots, c_{(k-1) \cdot i}\}$ and all remaining $n - n/k$ voters approve all candidates. Thus, for every voter $i \in [n/k]$ we get that $|N(A_i)| = 1 + n - n/k = 1 + (k - 1)(k + 1)$. Since $n/k = k + 1$, this implies that $f_i \geq k - 1$. Further, for all distinct voters $i, i' \in [n/k]$ it holds that $|A_i \cap A_{i'}| = 0$. However, since $n/k > k$, for each $W \subseteq C$ with $|W| = k$ there is a voter $i \in [n/k]$ with $|A_i \cap W| = 0$. Thus, for any $\alpha \geq 1$ and $\beta < k - 1$ this instance does not admit a (α, β) -IR committee. \square

To see that this bound is worst-possible, note that if $f_i = k$ for some voter i , this means that all voters have a set of at least k jointly approved candidates (and a committee consisting of such candidates would provide IR). On the other hand, every committee trivially provides $(1, k - 1)$ -IR whenever $f_i < k$ for all $i \in N$.

3.1 ABC Rules Violating IR

While the inapproximability result in the previous section is quite discouraging, a natural follow-up question is whether we can at least find IR committees *whenever they exist*. This question was already raised by Aziz et al. (2017) in the context of semi-strong JR, but remained open. In other words, we look for rules that are “consistent” with individual representation.

Definition 3 (IR-consistency). *An ABC rule is consistent with individual representation, or short IR-consistent, if it outputs at least one IR committee for every ABC election that admits one.*

Consistency with semi-strong JR can be defined analogously. We show that all common ABC voting rules fail consistency with respect to both IR and semi-strong JR.²

Example 1 already rules out any rule that always selects one of the candidates with the highest numbers of approvals, so-called *approval winners*. In particular, this class of rules includes all common committee-monotonic ABC rules as well as other “sequential” rules like Rule X, as these rules select one of the approval winners in the very first round.³

Proposition 4. *No ABC voting rule that always selects one of the approval winners into the winning committee is IR-consistent.*

Moreover, the rules PAV, Satisfaction-AV, and reverse-seqPAV select only committees including c_3 in Example 1, and thus fail IR-consistency. In Appendix B, we provide additional examples showing that all remaining ABC rules mentioned in Table 1 of the survey by Lackner and Skowron (2021) fail IR-consistency as well.

3.2 Computational Complexity

Another open problem stated by Aziz et al. (2017) concerns the computational complexity of deciding whether a given ABC election admits a committee providing semi-strong JR. We settle this question and the analogous one for individual representation by showing that both problems are NP-hard.

Theorem 5. *It is NP-hard to decide whether an ABC election admits an IR committee or a semi-strong JR committee.*

Proof. We reduce from *exact cover by 3-sets*. Here, we are given a set of elements $X = \{x_1, \dots, x_{3\ell}\}$ and a family of sets $\mathcal{T} \subseteq 2^X$ such that $|T| = 3$ for all $T \in \mathcal{T}$. The goal is to find a partition of X into sets from \mathcal{T} . The problem is NP-hard even if each element appears in exactly three sets (Garey and Johnson 1979). We construct an ABC instance by setting $N = X$ and $C = \mathcal{T}$. Further, each set $T \in \mathcal{T} = \{x_i, x_j, x_l\}$ is approved exactly by voters x_i, x_j , and x_l .

²Since neither IR nor semi-strong JR is always achievable (and semi-strong JR may be achievable in instances where IR is not) we can, in general, not deduce consistency regarding one of the notions from consistency regarding the other. However, all our examples in this section satisfy $f_i \leq 1$ for all voters i , such that semi-strong JR and IR coincide.

³For the common rules defined in this paper, we refer the reader to the survey (Lackner and Skowron 2021).

We set $k = \ell$. Hence, only groups of 3 voters corresponding to sets in \mathcal{T} are 1-cohesive, and we get $f_{x_i} = 1$ for each $x_i \in X$.

Every exact cover by 3-sets corresponds to a committee of size k where every voter is represented exactly once and thus provides IR in this instance. Conversely, every IR committee of the constructed ABC instance corresponds to a selection of sets from \mathcal{T} such that every element in X is covered exactly once. Since $f_i = 1$ for every voter, the same argument holds for semi-strong JR as well. \square

Moreover, it is hard to decide whether a given voter’s f_i -value exceeds a given value.

Theorem 6. *Given an ABC instance, a voter i , and $j \in \mathbb{N}$, it is NP-complete to decide whether $f_i \geq j$.*

Proof. It is easy to see that this problem is in NP since any subset of voters including voter i of size $j \cdot \frac{|N|}{k}$ and any subset of candidates of size j approved by all selected voters serves as a witness.

We reduce from *balanced complete bipartite subgraph*. Here, we are given a bipartite graph $G = (V_1 \cup V_2, E)$ and an integer j and the goal is to decide whether G has $K_{j,j}$ as a subgraph, i.e., a subgraph consisting of j vertices from V_1 and j vertices from V_2 forming a bipartite clique. The problem is known to be NP-hard (Garey and Johnson 1979). We construct an ABC instance by setting $N = V_1 \cup \{x\}$, $C = V_2 \cup \{y\}$ and $k = |V_1| + 1$. Thus, $\frac{n}{k} = 1$. Each $v \in V_1$ approves exactly its neighbors in G , as well as y , while x approves all candidates. It follows that $f_x \geq j + 1$ if and only if there is a set of j voters different from x approving at least a common set of $j + 1$ candidates. Since all voters approve y , this is equivalent to these j voters all approving j candidates different from y and therefore by definition all being connected to these j vertices in V_2 . Thus, they form a $K_{j,j}$ if and only if $f_x \geq j + 1$. \square

4 Domain Restrictions

We have seen in Theorem 3 that non-trivial approximations of individual representation are impossible to obtain in general. In this section, we explore whether this negative result can be circumvented by considering restricted domains of preferences. Domain restrictions for dichotomous (i.e., approval) preferences have been studied by Elkind and Lackner (2015) and Yang (2019).

Restricting attention to a well-structured domain often allows for axiomatic and algorithmic results that are not achievable otherwise (Elkind, Lackner, and Peters 2017). In the ABC setting, for example, it has recently been shown that a core-stable committee always exists in certain restricted domains (Pierczyński and Skowron 2021), whereas the existence of such committees is an open problem for the unrestricted domain.

We start by recalling the definitions of two classic restricted domains of dichotomous preferences: *candidate interval* and *voter interval* (Elkind and Lackner 2015).

Definition 4 (Candidate Interval). *An approval profile A satisfies candidate interval (CI) if there is a linear order over the candidates C such that for every voter $i \in N$, A_i consists of an interval of that order.*

Definition 5 (Voter Interval). *An approval profile A satisfies voter interval (VI) if there is a linear order over the voters N such that for every candidate $c_j \in C$, the voters that approve c_j build an interval of that order.*

The profile in Example 1 satisfies both candidate interval and voter interval. In fact, a voter order witnessing VI is given in Figure 1. To see that the profile satisfies CI as well, consider the order (c_1, c_3, c_2) . The profile in Example 2, on the other hand, satisfies neither CI nor VI.

Elkind and Lackner (2015) have shown that it can be checked in polynomial time whether a profile satisfies CI or VI. (If the answer is yes, a linear order over candidates/voters can be found efficiently as well.)

Our first observation is that the *candidate interval* domain is not helpful for our purposes: Indeed, the approval profile used to establish Theorem 3 can easily be seen to satisfy CI. Thus, restricting preferences in this way does not yield any improved bounds.

Corollary 7. *There exists a CI profile that does not admit an (α, β) -IR committee with $\beta < k - 1$ and any $\alpha \geq 1$.*

Now, we turn our attention to the *voter interval* domain. Due to the similarity between VI and CI, one might expect a similar result here. Surprisingly, however, we can prove a positive result for VI: We provide an algorithm that finds a $(2, 4)$ -IR committee in polynomial time for any VI profile. In Appendix C, we show that such a guarantee cannot be provided by common ABC rules.

Before describing the high level idea of our algorithm we state a useful property of VI profiles. Without loss of generality, we assume that the linear order witnessing VI is given by $(1, \dots, n)$. Moreover, for $a, b \in \mathbb{Z}$ with $a \leq b$, we let $[a, b]$ denote the integer interval $\{a, a + 1, \dots, b\}$.

Observation 8. *Let $i_1, i_2, i_3 \in [n]$ such that $i_1 < i_2 < i_3$. For any $S \subseteq C$, if $i_1 \in N(S)$ and $i_3 \in N(S)$, then $i_2 \in N(S)$.*

Let $S_i^* := \arg \max_{S \subseteq A_i} \{|S| : |N(S)| \geq |S| \cdot n/k\}$, i.e., a largest subset of A_i approved by sufficiently many voters to validate the f_i -value. (If multiple such sets exist, ties are broken arbitrarily.) From Observation 8 we know that if $i_1, i_2, i \in [n]$ such that $i_1 < i_2 < i$ or $i < i_2 < i_1$, and if $i_1 \in N(S_i^*)$, then $i_2 \in N(S_i^*)$, i.e., $N(S_i^*)$ forms an interval of the order of voters including i .

Further, let $N_{<i} := \{i' \in N(S_i^*) : i' \leq i - 1\}$ denote the set of voters in $N(S_i^*)$ that are ordered *before* i and let $N_{\geq i} := \{i' \in N(S_i^*) : i' \geq i\}$ denote the set of voters in $N(S_i^*)$ that are ordered *after* i (including i itself).

Observation 9. *For each voter i , there exist $a_i \in [1, i]$ and $b_i \in [i, n]$ such that $N_{<i} = [a_i, i - 1]$ and $N_{\geq i} = [i, b_i]$.*

Using this observation and the fact that $f_i = |S_i^*| \leq (|N_{<i}| + |N_{\geq i}|) \cdot k/n$, Algorithm 1 returns a $(2, 4)$ -IR committee W for any VI profile as follows. In the first

Algorithm 1: $(2, 4)$ -IR for Voter Interval Profiles

```

1:  $W_0 \leftarrow \emptyset$  // Round 1
2: for  $i = 1$  to  $n$  do
3:   if  $|W_{i-1} \cap A_i| < \lfloor |N_{\geq i}| \cdot k/(2n) \rfloor$  then
4:     Let  $S_i$  be an arbitrary subset of  $S_i^*$  of size equal to
        $\lfloor |N_{\geq i}| \cdot k/(2n) \rfloor - |W_{i-1} \cap A_i|$  such that  $|S_i \cap$ 
        $W_{i-1}|$  is minimized
5:      $W_i \leftarrow W_{i-1} \cup S_i$ 
6:  $\hat{W}_1 \leftarrow \emptyset$  // Round 2
7: for  $i = n$  to  $1$  do
8:   if  $|\hat{W}_{n-i} \cap A_i| < \lfloor |N_{<i}| \cdot k/(2n) \rfloor$  then
9:     Let  $S_i$  be an arbitrary subset of  $S_i^*$  of size equal to
        $\lfloor |N_{<i}| \cdot k/(2n) \rfloor - |\hat{W}_{n-i} \cap A_i|$  such that  $|S_i \cap$ 
        $(\hat{W}_{n-i} \cup W_n)|$  is minimized
10:     $\hat{W}_{n-i+1} \leftarrow \hat{W}_{n-i} \cup S_i$ 
11:  $S \leftarrow$  an arbitrary subset of  $C$  with  $S \cap (W_n \cup \hat{W}_n) = \emptyset$ 
       and  $|S| = k - |W_n| - |\hat{W}_n|$ 
12: return  $W_n \cup \hat{W}_n \cup S$ 

```

round, iterating from voter $i = 1$ to n , it selects at least $\lfloor |N_{\geq i}| \cdot k/(2n) \rfloor$ candidates that are approved by voter i . In the second round, iterating from voter $i = n$ to 1 , it selects at least $\lfloor |N_{<i}| \cdot k/(2n) \rfloor$ candidates that are approved by voter i (excluding the candidates that are selected in the first round). Together, this ensures $|W \cap A_i| \geq f_i/2 - 2$, where W is the set of selected candidates.

Theorem 10. *For every approval profile satisfying voter interval, Algorithm 1 returns a $(2, 4)$ -IR committee in polynomial time.*

Proof. Let $W = W_n \cup \hat{W}_n \cup S$ be the committee returned by Algorithm 1, and let $f_{\geq i} = |N_{\geq i}| \cdot k/n$ and $f_{<i} = |N_{<i}| \cdot k/n$. In the first round we ensure that $|W_n \cap A_i| \geq \lfloor f_{\geq i}/2 \rfloor$, as at iteration i if $|W_{i-1} \cap A_i| < \lfloor f_{\geq i}/2 \rfloor$, we include $\lfloor f_{\geq i}/2 \rfloor - |W_{i-1} \cap A_i|$ candidates into W_i that are not already included. Similarly, in the second round we ensure that $|\hat{W}_n \cap A_i| \geq \lfloor f_{<i}/2 \rfloor$ as at iteration $n - i + 1$ if $|\hat{W}_{n-i} \cap A_i| < \lfloor f_{\geq i}/2 \rfloor$, we include $\lfloor f_{\geq i}/2 \rfloor - |\hat{W}_{n-i} \cap A_i|$ candidates that are not already included. As $f_i \leq f_{\geq i} + f_{<i}$, for each $i \in N$ we have that

$$|W \cap A_i| \geq \lfloor f_{\geq i}/2 \rfloor + \lfloor f_{<i}/2 \rfloor \geq f_i/2 - 2,$$

and therefore $2 \cdot |W \cap A_i| + 4 \geq f_i$. Thus, we conclude that W provides $(2, 4)$ -IR.

Now we show that $|W_n| \leq k/2$ and $|\hat{W}_n| < k/2$. We first consider W_n .

Lemma 11. $|W_i| \leq \frac{((i-1)+|N_{>i}|) \cdot k}{2n}$ for all $i \in [n]$.

Proof. We prove the lemma using induction. For $i = 1$, $W_1 = \lfloor f_{\geq 1}/2 \rfloor \leq \frac{|N_{\geq 1}| \cdot k}{2n}$ and the statement holds. Assume that for all $t' < t$, we have $|W_{t'}| \leq \frac{((t'-1)+|N_{\geq t'}|) \cdot k}{2n}$.

We show that the statement holds for W_t . Note that

$$|W_t| = |W_{t-1}| + \lfloor f_{\geq t}/2 \rfloor - |W_{t-1} \cap A_t| \quad (1)$$

as at iteration t , the algorithm adds $\lfloor f_{\geq t/2} \rfloor - |W_{t-1} \cap A_t|$ candidates to W_t . Let $t^* = \max\{r \in [t-1] : t \in N(S_{t-r}^*)\}$. If $t^* = 0$, from Equation (1) we get that

$$\begin{aligned} |W_t| &\leq \frac{((t-2) + |N_{\geq t-1}|) \cdot k}{2n} \\ &\quad + \lfloor f_{\geq t/2} \rfloor - |W_{t-1} \cap A_t| \\ &\leq \frac{((t-2) + 1) \cdot k}{2n} + \lfloor f_{\geq t/2} \rfloor - |W_{t-1} \cap A_t| \\ &\leq \frac{(t-1 + |N_{\geq t}|) \cdot k}{2n}, \end{aligned}$$

where the last inequality follows from the fact that $|N_{\geq t-1}| = 1$, as $t \notin N(S_{t-1}^*)$. Now assume that $t^* > 0$. First, using induction, we show that

$$|W_t| = |W_{t-r}| + \lfloor f_{\geq t/2} \rfloor - |W_{t-r} \cap A_t|$$

for any $r \in [1, t^* + 1]$.

For $r = 1$, the claim immediately follows from Equation (1). Assume that for all $q' < q$ it holds that $|W_{t-q'}| = |W_{t-q'}| + \lfloor f_{\geq t/2} \rfloor - |W_{t-q'} \cap A_t|$. Now, we have

$$\begin{aligned} |W_{t-(q-1)} \cap A_t| &= \lfloor f_{\geq t-(q-1)/2} \rfloor - |W_{t-q} \cap A_{t-(q-1)}| \\ &\quad + |W_{t-q} \cap A_t| \end{aligned} \quad (2)$$

as at iteration $t - (q - 1)$, $\lfloor f_{\geq t-(q-1)/2} \rfloor - |W_{t-q} \cap A_{t-(q-1)}|$ candidates from $S_{t-(q-1)}^*$ are added to $W_{t-(q-1)}$, and as $t \in N(S_{t-(q-1)}^*)$, these candidates are approved by t , too. Then,

$$\begin{aligned} |W_t| &= |W_{t-(q-1)}| + \lfloor f_{\geq t/2} \rfloor - |W_{t-(q-1)} \cap A_t| \\ &= |W_{t-q}| + \lfloor f_{\geq t-(q-1)/2} \rfloor - |W_{t-q} \cap A_{t-(q-1)}| \\ &\quad + \lfloor f_{\geq t/2} \rfloor - |W_{t-(q-1)} \cap A_t| \\ &= |W_{t-q}| + \lfloor f_{\geq t/2} \rfloor - |W_{t-q} \cap A_t|, \end{aligned}$$

where the second transition follows since at iteration $t - (q - 1)$, $\lfloor f_{\geq t-(q-1)/2} \rfloor - |W_{t-q} \cap A_{t-(q-1)}|$ candidates are added to $W_{t-(q-1)}$, and the third transition follows from Equation (2).

Now, we distinguish two cases.

Case 1: $t^* = t - 1$. Here, we have

$$\begin{aligned} |W_t| &= |W_{t-(t^*+1)}| + \lfloor f_{\geq t/2} \rfloor - |W_{t-(t^*+1)} \cap A_t| \\ &= |W_0| + \lfloor f_{\geq t/2} \rfloor - |W_0 \cap A_t| = \frac{|N_{\geq t}| \cdot k}{2n}. \end{aligned}$$

Case 2: $t^* > 1$. Here, we have

$$\begin{aligned} |W_t| &= |W_{t-(t^*+1)}| + \lfloor f_{\geq t/2} \rfloor - |W_{t-(t^*+1)} \cap A_t| \\ &\leq \frac{(t - (t^* + 1) - 1 + |N_{\geq t-(t^*+1)}|) \cdot k}{2n} \\ &\quad + \frac{|N_{\geq t}| \cdot k}{2n} \\ &\leq \frac{(t - 1 + |N_{\geq t}|) \cdot k}{2n}, \end{aligned}$$

where the third inequality follows from the fact that $|N_{\geq t-(t^*+1)}| \leq t - (t - (t^* + 1))$, as t is not in $N(S_{t-(t^*+1)}^*)$. \square

As Rounds 1 and 2 of Algorithm 1 are symmetric, with similar arguments, we can show the following lemma.

Lemma 12. $|\hat{W}_i| \leq \frac{((i-1) + |N_{<n-i+1}|) \cdot k}{2n}$ for all $i \in [n]$.

From Lemma 11, for $i = n$, we have $|N_{\geq n}| \leq 1$, and hence

$$|W_n| \leq \frac{((n-1) + 1) \cdot k}{2n} \leq k/2.$$

From Lemma 12, for $i = n$, we have that $|N_{<1}| = 0$, and hence

$$|W_n| \leq \frac{((n-1) + 0) \cdot k}{2n} < k/2.$$

Thus, $|W| = |W_n| + |\hat{W}_n| < k$.

Lastly, we show that S_i^* , and thus f_i , can be computed in polynomial time. For this, we employ Observation 9, i.e., the fact that $N(S_i^*)$ forms an interval of voters that includes i . We consider all such intervals and for each of them calculate the maximum subset of candidates that the voters in this interval deserve due to their size.

Lemma 13. For any voter i , f_i and S_i^* can be computed in polynomial time.

This concludes the proof of Theorem 10. \square

Further, we can show that the bound provided by Theorem 10 is almost tight up to the additive part of 4.

Theorem 14. There exists a VI approval profile that does not admit an $(\alpha, 0)$ -IR committee with $\alpha < 2 - 2/k$.

Proof. Consider the following instance with $k > 2$ and $m = 2(k - 1)$. All the voters $i \in [2, n - 1]$ approve all the candidates, while $A_1 = \{c_1, \dots, c_{k-1}\}$ and $A_n = \{c_{m-k+2}, \dots, c_m\}$. Notice that this profile is VI. Indeed, if we order the voters as $1, 2, \dots, n$, then the voters that approve each candidate form an interval of the ordering. Now, we can see that $f_1 = f_n = k - 1$, but for each $W \subseteq C$, with $|W| = k$, either $|A_1 \cap W| \leq k/2$ or $|A_n \cap W| \leq k/2$. \square

Beyond VI and CI, many other domain restrictions have been studied in the literature. In Appendix C, we provide lower and upper bounds for (α, β) -IR for all domain restrictions introduced by Elkind and Lackner (2015) and Yang (2019). Any domain that is more restrictive than VI inherits the guarantee of a $(2, 4)$ -IR committee from VI—but we show that in some cases we can achieve better approximation guarantees. On the other hand, any domain that is more general than CI inherits the inapproximability from CI. In fact, we show that the same lower bound applies even in a slightly more restricted domain introduced by Yang (2019). We also determined for which of the considered domain restrictions a semi-strong JR committee is guaranteed to exist.




Figure 3: The figure on the left shows the ratio of generated profiles that admit an IR committee. In the figure on the right, for each model and each voting rule, the bold colored part of the bar indicates the ratio of instances the rule returned an IR committee, while the pale-colored part indicates the same ratio for semi-strong JR, averaged over all values k with $2 \leq k \leq 20$. For each model, the black line indicates the fraction of instances admitting an IR committee, while the gray dashed line indicates the ratio of instances admitting a semi-strong JR committee.

5 Experimental Results

To complement our theoretical results, we performed experiments on generated approval profiles.

Setup. Inspired by Peters et al. (2021), we used six models to generate approval profiles: a voter-interval model (VI), a candidate-interval model (CI), a two-dimensional Euclidean model (2D), an impartial culture model (IC), an urn model (Urn), and a Mallows model (Mallows). For detailed descriptions of these models, we refer to Appendix D. We considered the ABC rules AV, PAV, seq-PAV, greedy Monroe, Rule X, seq-Phragmén, and sequential Chamberlin–Courant (seq-CC). For all conducted experiments, we have 100 voters and 50 candidates and generated 500 instances for each model and for each $k \in [50]$.

Results. First, we studied how often the generated approval profiles admit an IR committee. The results are shown in the left graph of Figure 3. We found that IR committees exist quite often, especially for larger values of k . In particular, profiles generated by the VI model or by the urn model almost always admit IR committees. On the other hand, profiles generated by the CI model almost never admit IR committees (except for $k \geq 35$). This striking contrast between VI and CI, which is reminiscent of our theoretical results in Section 4, can be explained with a feature of the preference generation model: Due to the way we generate CI preferences, almost 20% of voters have an approval set spanning at least 20% of the candidates. These voters approving many candidates are then part of multiple cohesive groups, not all of which can be represented in an IR manner. (A similar situation can be observed in the profile that proves Theorem 3.)

Second, we studied how often different ABC rules select a committee providing IR (or semi-strong JR). In order not to dilute our results, we restricted k to the “interesting” range between 2 and 20. The results are shown in the right

graph of Figure 3. Of course, the fraction of profiles for which a rule selects an IR (or semi-strong JR) committee is upper-bounded by the fraction of profiles that admit such a committee. For each model, the latter fraction is depicted in the graph as a solid black line for IR, and a dashed gray line for semi-strong JR. While no rule manages to find an IR committee every time one exists, the rules PAV, sequential PAV, sequential Phragmén, and Rule X select IR committees often. For the small fraction of CI profiles that admit an IR committee, all considered rules do a good job in finding one. Since seq-CC greedily optimizes the amount of voters that are represented at least once, it finds a committee providing semi-strong JR in almost all profiles that admit one. But as the rule does not aim at representing voters more than once, it almost never produces IR committees. In the profiles generated by the IC model, IR often coincides with semi-strong JR (for $k \leq 20$) because almost all non-zero f_i -values are 1. This is in line with the effect noticed by Brederbeck et al. (2019), whose experiments showed that EJR and JR are very likely to coincide under IC.

6 Discussion

Based on the observations that common axioms in approval-based committee voting do not address the representation of individual voters, and that common voting rules sometimes unfairly distinguish between such voters, we formalize *individual representation* (IR) as a requirement for committees. We find that all common voting rules fail to provide IR committees, even when these exist. Nevertheless, for some restricted domains—most prominently, voter interval preferences—we provide polynomial-time algorithms for finding committees that provide a good approximation to IR. Our experimental results suggest that IR is achievable in many instances that follow somewhat realistic preferences. It remains an open problem to find intuitive voting rules that provide (approximate) IR whenever possible.

Acknowledgments

We thank Nisarg Shah and the anonymous reviewers of AAAI 2022 for their helpful comments and discussion. This work was supported by the Deutsche Forschungsgemeinschaft (DFG) under grant BR 4744/2-1 and the Graduiertenkolleg “Facets of Complexity” (GRK 2434).

References

- Aziz, H.; Brill, M.; Conitzer, V.; Elkind, E.; Freeman, R.; and Walsh, T. 2017. Justified Representation in Approval-Based Committee Voting. *Social Choice and Welfare*, 48(2): 461–485.
- Balinski, M.; and Young, H. P. 1982. *Fair Representation: Meeting the Ideal of One Man, One Vote*. Yale University Press.
- Bredereck, R.; Faliszewski, P.; Kaczmarczyk, A.; and Niedermeier, R. 2019. An Experimental View on Committees Providing Justified Representation. In *Proceedings of the 28th International Joint Conference on Artificial Intelligence (IJCAI)*, 109–115.
- Brill, M.; Gözl, P.; Peters, D.; Schmidt-Kraepelin, U.; and Wilker, K. 2020. Approval-Based Apportionment. In *Proceedings of the 34th AAAI Conference on Artificial Intelligence (AAAI)*, 1854–1861. AAAI Press.
- Brill, M.; Israel, J.; Micha, E.; and Peters, J. 2022. Individual Representation in Approval-Based Committee Voting. In *Proceedings of the 36th AAAI Conference on Artificial Intelligence (AAAI)*. AAAI Press. Forthcoming.
- Elkind, E.; and Lackner, M. 2015. Structure in Dichotomous Preferences. In *Proceedings of the 24th International Joint Conference on Artificial Intelligence (IJCAI)*, 2019–2025. AAAI Press.
- Elkind, E.; Lackner, M.; and Peters, D. 2017. Structured Preferences. In Endriss, U., ed., *Trends in Computational Social Choice*, chapter 10.
- Gangl, C.; Maly, J.; Lackner, M.; and Woltran, S. 2019. Aggregating Expert Opinions in Support of Medical Diagnostic Decision-Making. In *Proceedings of the 11th International Workshop on Knowledge Representation for Health Care (KR4HC-2019)*.
- Garey, M. R.; and Johnson, D. S. 1979. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman.
- Jung, C.; Kannan, S.; and Lutz, N. 2020. A Center in your Neighborhood: Fairness in Facility Location. In *Proceedings of the Symposium on Foundations of Responsible Computing (FORC)*, 5:1–5:15.
- Lackner, M.; Regner, P.; Krenn, B.; and Forster, S. S. 2021. abc voting: A Python Library of Approval-Based Committee Voting Rules. Current version: <https://github.com/martinlackner/abcvoting>.
- Lackner, M.; and Skowron, P. 2021. Approval-Based Committee Voting: Axioms, Algorithms, and Applications. Technical report, arXiv:2007.01795 [cs.GT].
- Peters, D.; Pierczyński, G.; Shah, N.; and Skowron, P. 2021. Market-Based Explanations of Collective Decisions. In *Proceedings of the 35th AAAI Conference on Artificial Intelligence (AAAI)*, 5656–5663. AAAI Press.
- Peters, D.; Pierczyński, G.; and Skowron, P. 2021. Proportional Participatory Budgeting with Additive Utilities. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 34. Forthcoming.
- Pierczyński, G.; and Skowron, P. 2021. Core-Stable Committees under Restricted Domains. Technical report, arXiv:2108.01987 [cs.GT].
- Pukelsheim, F. 2014. *Proportional Representation: Apportionment Methods and Their Applications*. Springer.
- Sánchez-Fernández, L.; Elkind, E.; Lackner, M.; Fernández, N.; Fisteus, J. A.; Basanta Val, P.; and Skowron, P. 2017. Proportional Justified Representation. In *Proceedings of the 31st AAAI Conference on Artificial Intelligence (AAAI)*, 670–676. AAAI Press.
- Skowron, P.; Lackner, M.; Brill, M.; Peters, D.; and Elkind, E. 2017. Proportional Rankings. In *Proceedings of the 26th International Joint Conference on Artificial Intelligence (IJCAI)*, 409–415. IJCAI.
- Terzopoulou, Z.; Karpov, A.; and Obraztsova, S. 2021. Restricted Domains of Dichotomous Preferences with Possibly Incomplete Information. In *Proceedings of the 35th AAAI Conference on Artificial Intelligence (AAAI)*, 5726–5733. AAAI Press.
- Yang, Y. 2019. On the Tree Representations of Dichotomous Preferences. In *Proceedings of the 28th International Joint Conference on Artificial Intelligence (IJCAI)*, 644–650.

A Omitted Proofs

A.1 Proof of Lemma 12

Lemma 12. $|\hat{W}_i| \leq \frac{((i-1)+|N_{<n-i+1}|) \cdot k}{2n}$ for all $i \in [n]$.

Proof. We prove the lemma using induction. For $i = 1$, $\hat{W}_1 = \lfloor f_{<n}/2 \rfloor \leq \frac{|N_{<n}| \cdot k}{2n}$ and the statement holds. Assume that for all $t' < t$, we have $|\hat{W}_{t'}| \leq \frac{((t'-1)+|N_{<n-t'+1}|) \cdot k}{2n}$.

We show that the statement holds for \hat{W}_t . Note that

$$|\hat{W}_t| = |\hat{W}_{t-1}| + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-1} \cap A_{n-t+1}| \quad (3)$$

as at iteration t , the algorithm adds $\lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-1} \cap A_{n-t+1}|$ candidates to \hat{W}_t . Let $t^* = \max\{r \in [t-1] : n-t+1 \in N(S_{n-t+1+r}^*)\}$. If $t^* = 0$, from Equation (3) we get that

$$\begin{aligned} |\hat{W}_t| &\leq \frac{((t-2) + |N_{<n-t+2}|) \cdot k}{2n} \\ &\quad + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-1} \cap A_{n-t+1}| \\ &\leq \frac{((t-2) + 0) \cdot k}{2n} \\ &\quad + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-1} \cap A_{n-t+1}| \\ &\leq \frac{((t-2) + |N_{<n-t+1}|) \cdot k}{2n}, \end{aligned}$$

where the last inequality follows from the fact that $|N_{<n-t+2}| = 0$, as $n-t+1 \notin N(S_{n-t+2}^*)$. Now assume that $t^* > 0$. First, using induction, we show that

$$|\hat{W}_t| = |\hat{W}_{t-r}| + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-r} \cap A_{n-t+1}|$$

for any $r \in [1, t^* + 1]$.

For $r = 1$, the claim immediately follows from Equation (3). Assume that for all $q' < q$ it holds that $|\hat{W}_i| = |\hat{W}_{t-q'}| + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-q'} \cap A_{n-t+1}|$. Now, we have

$$\begin{aligned} |\hat{W}_{t-(q-1)} \cap A_{n-t+1}| &= \lfloor f_{<n-t+(q-1)+1}/2 \rfloor \\ &\quad - |\hat{W}_{t-q} \cap A_{n-t+(q-1)+1}| \\ &\quad + |\hat{W}_{t-q} \cap A_{n-t+1}| \quad (4) \end{aligned}$$

as at iteration $t - (q - 1)$, $\lfloor f_{<n-t+(q-1)+1}/2 \rfloor - |\hat{W}_{t-q} \cap A_{n-t+(q-1)+1}|$ candidates from $S_{n-t+(q-1)+1}^*$ are added to $\hat{W}_{t-(q-1)}$, and as $n - t + 1 \in N(S_{n-t+(q-1)+1}^*)$, these candidates are approved by $n - t + 1$, too. Then,

$$\begin{aligned} |\hat{W}_t| &= |\hat{W}_{t-(q-1)}| + \lfloor f_{<n-t+1}/2 \rfloor \\ &\quad - |\hat{W}_{t-(q-1)} \cap A_{n-t+1}| \\ &= |\hat{W}_{t-q}| + \lfloor f_{<n-t+(q-1)+1}/2 \rfloor \\ &\quad - |\hat{W}_{t-q} \cap A_{n-t+(q-1)+1}| \\ &\quad + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-(q-1)} \cap A_{n-t+1}| \\ &= |\hat{W}_{t-q}| + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_{t-q} \cap A_{n-t+1}|, \end{aligned}$$

Algorithm 2: Finding f_i and S_i^*

```

1:  $f_i \leftarrow 0$ 
2:  $S_i^* \leftarrow \emptyset$ 
3: for  $a_i = 1$  to  $i$  do
4:   for  $b_i = i$  to  $n$  do
5:      $S \leftarrow \emptyset$ 
6:     for  $j = 1$  to  $m$  do
7:       if  $\{a_i, \dots, b_i\} \subseteq N(\{c_j\})$  then
8:          $S \leftarrow S \cup \{c_j\}$ 
9:          $\ell^* \leftarrow \arg \max_{\ell \in \mathbb{N}} \{\ell : b_i - a_i + 1 \geq \ell \cdot n/k\}$ 
10:        if  $\min\{\ell^*, |S|\} > f_i$  then
11:           $S_i^* \leftarrow$  an arbitrary subset of  $S$  of size as close
            as possible to  $\ell^*$ 
12:           $f_i \leftarrow |S_i^*|$ 
13: return  $f_i, S_i^*$ 

```

where the second transition follows since at iteration $t - (q - 1)$, $\lfloor f_{<n-t+(q-1)+1}/2 \rfloor - |\hat{W}_{t-q} \cap A_{n-t+(q-1)+1}|$ candidates are added to $\hat{W}_{t-(q-1)}$, and the third transition follows from Equation (4).

Now, we distinguish two cases.

Case 1: $t^* = t - 1$. Here, we have

$$\begin{aligned} |\hat{W}_t| &= |\hat{W}_{t-(t^*+1)}| + \lfloor f_{<n-t+1}/2 \rfloor \\ &\quad - |\hat{W}_{t-(t^*+1)} \cap A_{n-t+1}| \\ &= |\hat{W}_0| + \lfloor f_{<n-t+1}/2 \rfloor - |\hat{W}_0 \cap A_{n-t+1}| \\ &= \frac{|N_{<n-t+1}| \cdot k}{2n}. \end{aligned}$$

Case 2: $t^* > 1$. Here, we have

$$\begin{aligned} |\hat{W}_t| &= |\hat{W}_{t-(t^*+1)}| + \lfloor f_{<n-t+1}/2 \rfloor \\ &\quad - |\hat{W}_{t-(t^*+1)} \cap A_{n-t+1}| \\ &\leq \frac{(t - (t^* + 1) - 1 + |N_{<n-t+(t^*+1)+1}|) \cdot k}{2n} \\ &\quad + \frac{|N_{<n-t+1}| \cdot k}{2n} \\ &\leq \frac{(t - 2 + |N_{<n-t+1}|) \cdot k}{2n}, \end{aligned}$$

where the third inequality follows from the fact that $|N_{<n-t+(t^*+1)+1}| \leq t^*$, as $n - t + 1$ is not in $N(S_{n-t+(t^*+1)+1}^*)$. \square

A.2 Proof Lemma 13

Lemma 13. For any voter i , f_i and S_i^* can be computed in polynomial time.

Proof. We show that Algorithm 2 correctly computes both f_i and S_i^* . For each interval $[a_i, b_i]$ where $a_i \leq i$ and $i \leq b_i$, the algorithm finds the maximum number of candidates that this interval is eligible to elect, denoted by ℓ^* . Moreover, $S \subseteq C$ denotes the set of candidates that is approved by




Figure 4: Relationships between different notions of representation. An arrow from X to Y shows that X implies Y (if X is possible in the given instance). Note that a committee providing the shaded notions does not always exist (the case for core stability is an open problem).

all the voters in the interval. The algorithm calculates the maximum subset of S that can be elected by the voters in the interval as $\min\{\ell^*, |S|\}$. Then, S_i^* and f_i are updated properly by assigning them the biggest subset and the size of it, respectively, that an interval of voters can elect.

Assume for contradiction that the algorithm returns a subset S' such that $|S'| < |S_i^*|$. This means that there is an interval of voters $[a_i, b_i]$ that can elect S_i^* . Thus, when the algorithm considers this interval, it would return a subset of size at least $|S_i^*|$, which concludes the proof. \square

B Additional Details for Section 3

Here we provide additional information on the investigation of the relation of individual representation with representation notions from the literature. First we study the relation of semi-strong JR and IR with PJR and FJR. In the second subsection we investigate the relation of our notion to perfect representation introduced by Sánchez-Fernández et al. (2017). These results are graphically represented in Figure 4. Lastly, we provide the omitted examples showing that basically all common ABC voting rules violate IR-consistency.

B.1 Relation to PJR and FJR

Proportional justified representation was introduced by Sánchez-Fernández et al. (2017). Recall that given an ABC election (A, k) we call a group of voters $V \subseteq N$ ℓ -cohesive if $|V| \geq \ell \cdot n/k$ and $|\bigcap_{i \in V} A_i| \geq \ell$. A committee $W \subseteq C$ of size k provides *proportional justified representation* (PJR) if for each ℓ -cohesive group V it holds that $|W \cap (\bigcup_{i \in V} A_i)| \geq \ell$. It is well-known that PJR is more demanding than JR but implied by EJR.

A strengthening of EJR was recently introduced by Peters, Pierczyński, and Skowron (2021) under the name fully justified representation. We call $V \subseteq N$ *weakly* (β, S) -cohesive for $\beta \in \mathbb{N}$ and $S \subseteq C$ if $|V| \geq |S| \cdot n/k$ and $|S \cap A_i| \geq \beta$ for all voters $i \in V$. A committee $W \subseteq C$ of size k provides *fully justified representation* (FJR) if for each weakly (β, S) -cohesive V there is a voter $i \in V$ with $|W \cap A_i| \geq \beta$.

PJR is implied by EJR and thus also implied by IR (if it exists). On the other hand, there exist profiles where semi-strong JR committees exist that do not provide PJR.

Theorem 15. *Semi-strong JR is incompatible with PJR.*

Proof. Consider the same instance as in the proof of Proposition 2 where $n = 8$ and $k = 4$ with the following approval profile.

$$\begin{aligned}
 A_1 = \dots = A_4 &= \{c_1, c_2\} \\
 A_5 &= \{c_3, c_4, c_5\} \\
 A_6 &= \{c_3\} \\
 A_7 &= \{c_4\} \\
 A_8 &= \{c_5\}
 \end{aligned}$$

As $f_6 = f_7 = f_8 = 1$, for any committee W that provides semi-strong JR, it should hold that $\{c_3, c_4, c_5\} \subseteq W$. But then, while the first four voters form a 2-cohesive group, we have that $|W \cap (\bigcup_{i \in [4]} A_i)| = 1$. \square

Since PJR is implied by EJR and FJR and these are known to always exist, it follows directly that semi-strong JR is also incompatible with EJR and FJR. Since in the example above the core is nonempty (as, e.g., the committee $\{c_1, c_2, c_3, c_4\}$ is core-stable) we can also deduce that semi-strong JR is incompatible with core stability.

Note that in the proof of Proposition 1 the group of voters $1, \dots, 8$ forms a $(3, \{c_1, \dots, c_4\})$ -cohesive group. None of these voters has 3 representatives in the unique IR committee. Further, the committee $\{c_1, \dots, c_5, c_7\}$ providing core stability also by definition provides FJR. Thus, we can conclude the following.

Corollary 16. *IR is incompatible with FJR.*

B.2 Relation to Perfect Representation

Consider an approval profile A and a committee size k such that k divides the number of voters n . We say that a committee W of size k provides *perfect representation* if it is possible to partition the electorate N into k pairwise disjoint subsets N_1, \dots, N_k of size $\frac{n}{k}$ each, and assign a distinct candidate from W to each of the subsets in such a way that for each subset all the voters in the subset approve of the assigned candidate.

It is known that not all ABC voting instances where k divides n allow for a committee providing PR (Sánchez-Fernández et al. 2017). Thus, in the following, we will call an ABC voting instance a *PR-instance* if it allows for a perfect representation committee.

Proposition 17. *On PR-instances, every committee providing perfect representation also provides semi-strong JR, but not the other way around.*

Proof. Assume a PR-instance is given together with a committee W that provides perfect representation. By definition, every voter $i \in N$ witnesses $|A_i \cap W| \geq 1$. Thus, W also provide semi-strong JR. As a counterexample for the other direction, consider the following instance where $n = 6$ and $k = 3$.

$$\begin{aligned}
 A_1 = A_2 &= \{c_1, c_4\} & A_4 &= \{c_2, c_5\} \\
 A_3 &= \{c_2, c_4, c_6\} & A_5 = A_6 &= \{c_3, c_5\}.
 \end{aligned}$$

The committee $W = \{c_4, c_5, c_6\}$ provides semi-strong JR but not perfect representation (while $\{c_1, c_2, c_3\}$ provides both). \square

Proposition 18. *There are PR-instances that do not allow for an IR committee. Additionally, there are PR-instances where an IR committee exists but does not provide perfect representation.*

Proof. Regarding the first claim consider the following instance with $n = 8$ and $k = 4$.

$$\begin{aligned} A_1 &= \{c_1\} \\ A_2 &= \{c_2\} \\ A_3 &= \{c_3\} \\ A_4 &= \{c_4\} \\ A_5 &= \{c_1, c_5, c_6\} \\ A_6 &= \{c_2, c_5, c_6\} \\ A_7 &= \{c_3, c_5, c_6\} \\ A_8 &= \{c_4, c_5, c_6\} \end{aligned}$$

Here $\{c_1, \dots, c_4\}$ provides perfect representation (and semi-strong JR) but not IR (which is not achievable in this instance). Regarding the second claim again consider the instance from the proof of Proposition 17 with $n = 6$ and $k = 3$. Here, the committee $\{c_4, c_5, c_6\}$ provides IR, but not perfect representation. \square

B.3 Counterexamples for IR-Consistency

Here we provide further examples showing that all ABC rules mentioned in Table 1 of the recent survey by Lackner and Skowron (2021) violate IR-consistency. Note that all committee monotone rules (including AV, SAV, sequential PAV, and sequential Phragmén) as well as Rule X and PAV were already ruled out to satisfy IR-consistency in Section 3.1.

Example 3. *Consider the following profile with $n = 16$ voters and assume $k = 4$:*

$$\begin{aligned} A_1 = A_2 = A_3 &= \{c_1\} & A_4 &= \{c_1, c_5\} \\ A_5 = A_6 = A_7 &= \{c_2\} & A_8 &= \{c_2, c_5\} \\ A_9 = A_{10} = A_{11} &= \{c_3\} & A_{12} &= \{c_3, c_5\} \\ A_{13} = A_{14} = A_{15} &= \{c_4\} & A_{16} &= \{c_5\}. \end{aligned}$$

Here we have $f_i = 1$ for all voters $i \in N$ and thus the only committee providing individual representation is $W = \{c_1, c_2, c_3, c_5\}$. Chamberlin-Courant-AV (CCAV), MonroeAV and PAV with the weight-vector $(1, \frac{1}{n}, \frac{1}{n^2}, \dots)$, which provide an IR committee in the example of Proposition 4, choose $\{c_1, c_2, c_3, c_4\}$ and thus fail individual representation.

The only two remaining rules from Table 1 in (Lackner and Skowron 2021) are maxPhragmén and MinimaxAV.

Example 4. *Consider the following profile with $n = 6$ voters and assume $k = 3$:*

$$\begin{aligned} A_1 &= \{c_1\} & A_3 &= \{c_1, \dots, c_5\} & A_5 &= \{c_4, c_5, c_6\} \\ A_2 &= \{c_2\} & A_4 &= \{c_3, \dots, c_6\} & A_6 &= \{c_5, c_6\}. \end{aligned}$$

Here we have $f_i = 1$ for all voters $i \in N$ and thus $\{c_1, c_2, c_5\}$ and $\{c_1, c_2, c_6\}$ are the only committees providing individual representation (or semi-strong JR). It turns out that maxPhragmén does not select any of these two. It is easy to see that the load distribution with the minimum maximal load for either of the two IR committees is $(\frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3})$. The committee $\{c_1, c_4, c_5\}$, however, induces a load distribution of $(\frac{3}{5}, 0, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}, \frac{3}{5})$ which is lower both in terms of the maximum as well as lexicographic ordering.

Example 5. *The final example uses the following approval profile:*

$$99 \times \{c_1, c_2\} \quad 1 \times \{c_3, \dots, c_8\}.$$

Here for $k = 2$, MinimaxAV (which minimises the maximum Hamming-distance among all voters to the winning committee) selects any two candidates from $\{c_3, \dots, c_8\}$ and none that is supported by the 99 voters. This clearly violates individual representation and semi-strong JR.

C Additional Details for Section 4

C.1 Known Voting Rules do not Guarantee (2, 4)-IR for VI Preferences

The two examples below show that many known voting rules are not guaranteed to return a committee that is (2, 4)-IR for VI preferences.

Example 6. *Consider the following profile with $n = 28$ voters and assume $k = 14$:*

$$\begin{aligned} A_1 &= \{c_1, \dots, c_7\} \\ A_2 = \dots = A_{14} &= \{c_1, \dots, c_{20}\} \\ A_{15} = \dots = A_{28} &= \{c_8, \dots, c_{20}\}. \end{aligned}$$

Notice that $f_1 = 7$ and $f_i = 13$ for $i > 1$. Here, Rule X, AV, PAV, seq-PAV, rev-seq-PAV, seq-Phragmén, SAV, and Greedy Monroe (depending on the tie breaking) all return committees W where $\{c_8, \dots, c_{20}\} \subset W$ and thus only one of the candidates among $\{c_1, \dots, c_7\}$ is included. Hence, they all fail to provide a (2, 4)-IR committee.

Example 7. *Now consider the following profile with, again, $n = 28$ voters and assume $k = 14$:*

$$\begin{aligned} A_1 = \dots = A_{14} &= \{c_1, \dots, c_7\} \\ A_{10+i} &= \{c_{5+i}\}, \quad \forall i \in \{1, \dots, 14\}. \end{aligned}$$

Again, we have $f_1 = 7$. CC and seq-CC return committees W where only one candidate among $\{c_1, \dots, c_7\}$ is included. Thus, they fail to provide a (2, 4)-IR committee.

A simple adaption of Example 5 shows that MinimaxAV fails to provide any good approximation for IR even on t -PART instances. From Table 1 of the recent survey by

	Individual Representation		ss-JR
	Lower Bound	Upper Bound	
PART	(1, 0)	(1, 0)	Yes
α -TR	(1, 0)	(1, 0)	Yes
VEI	$(2 - 2/k, 0)$	(2, 0)	Yes
CEI	$(2 - 2/k, 0)$	(2, 0)	Yes
DUE	$(2 - 2/k, 0)$	(2, 4)	No
VI	$(2 - 2/k, 0)$	(2, 4)	No
CI	(1, $k - 1$)	(1, $k - 1$)	No

Table 1: Individual representation guarantees in structured profiles. The column Lower Bound indicates that, while the column Upper Bound gives us values (α, β) such that an (α, β) individually representative committee always exists (and can be computed efficiently). Finally, the column ss-JR refers to the existence of semi-strong JR committees in every instance.

Lackner and Skowron (2021), the only rules missing are maxPhragmén and MonroeAV. In the supplementary material we provide an example profile that is VI with $n = 30$ voters and $m = 300$ candidates where these two rules also fail to provide a $(2, 4)$ -IR committee. Thus, we can conclude that Algorithm 1 outperforms all common ABC voting rules in terms of approximating IR on VI profiles.

C.2 Further Domain Restrictions

Here, we consider all the restricted domains that are mentioned in (Elkind and Lackner 2015; Yang 2019). An overview of the relation of these restrictions (which is adapted from (Yang 2019)) can be found in Figure 5. An overview of the results of this section can be found in Table 1.

t -PART

Definition 6 (t -partition (t -PART)). *An approval profile A satisfies t -partition (t -PART) if there is a partition (C_1, \dots, C_t) of C such that for every voter $i \in N$ there exists C_j such that $A_i = C_j$.*

Theorem 19. *Under t -PART approval profiles an IR committee always exists.*

Proof. Note that for each voter $i \in N$ with $A_i = C_j$ for some $j \in [t]$, $f_i = \lfloor N(C_j) \cdot k/n \rfloor$. Now consider the committee that contains $\lfloor N(C_j) \cdot k/n \rfloor$ candidates from each C_j . Clearly, W provides IR and also

$$|W| = \sum_{j \in T} \left\lfloor N(C_j) \cdot \frac{k}{n} \right\rfloor \leq n \cdot \frac{k}{n} = k,$$

which completes the proof. \square

CEI, VEI and WSC

Definition 7 (Candidate Extremal Interval (CEI)). *An approval profile A satisfies Candidate Extremal Interval (CEI) if there is a linear order of C such that for every voter $i \in N$, A_i consist a prefix or a suffix of the order.*




Figure 5: Upper (green) and lower (red) bounds of approximate IR and guaranteed existence (white) or not (gray) of semi-strong JR committees under the respective domain. An arrow from X to Y shows that X implies Y. An upper (resp. lower) bound for each domain is the best (resp. worst) upper (lower) bound of any of its descendants (resp. ancestors), including itself.

Definition 8 (Voter Extremal Interval (VEI)). *An approval profile A satisfies Voter Extremal Interval (VEI) if there is a linear order of N such that for every candidate $c_j \in C$, the voters that approve c_j consist a prefix or a suffix of the order.*

Definition 9 (Weakly Single-Crossing (WSC)). *An approval profile A satisfies Weakly Single-Crossing (WSC) if there is a linear order of N such that for each pair of candidates c, c' in C it holds that each of the vote sets $N_1 = \{i : c \in A_i, c' \notin A_i\}$, $N_2 = \{i : c \notin A_i, c' \in A_i\}$ and $N_3 = N \setminus (N_1 \cup N_2)$ forms an interval of this ordering, and N_3 appears between N_1 and N_2 .*

Theorem 20. *For any CEI and VEI profile, there exists a $(2, 0)$ -IR and a semi-strong JR committee, and both can be found in polynomial time.*

Proof. Assume we are given a CEI profile and the order c_1, \dots, c_m over the candidates such that each A_i is either a prefix or a suffix of that order. If $f_i = k$ for some voter, then this means that there are at least k candidates that are approved by all the voters and these candidates can form the winning committee which clearly is IR and semi-strong JR. Otherwise, if $f_i \leq k - 1$ for all voters $i \in [n]$, we set

$$W = \{c_1, \dots, c_{\lfloor k/2 \rfloor}, c_{m - \lfloor k/2 \rfloor + 1}, \dots, c_m\}$$

, i.e., W consists of the first $\lfloor k/2 \rfloor$ candidates and the last $\lfloor k/2 \rfloor$ candidates of the order. Hence, for each $i \in [n]$, it holds that $|A_i \cap W| \geq \min\{\lfloor k/2 \rfloor - 1, |A_i|\}$, and as $f_i \leq k - 1$, we get that W is $(2, 0)$ -individually representative. Now notice that as W consists of the first and the last candidate in the order. Hence, W provides semi-strong-JR as each voter approves one of these two candidates.

Now, assume we are given a VEI profile and the order over the voters. We order the candidates as follows. Let C_p

and C_s be the set of candidates such that the voters that approve a candidate in C_p and C_s form a prefix and a suffix of N , respectively. Without loss of generality, a candidate c such that $N(c) = N$ is assigned to C_p . A candidate c in C_p is ordered before c' in C_p if the last voter that approves c is ordered after the last voter that approves c' (break ties arbitrarily). A candidate c in C_s is ordered before c' in C_s if the first voter that approves c is ordered after the first voter that approves c' (break ties arbitrary). All the candidates in C_p are ordered before the candidates in C_s . As above, if $f_i = k$ for some voter, then this means that there are at least k candidates that are approved by all the voters and these candidates form the winning committee which clearly is IR and semi-strong JR. Otherwise, if W consists of the first $\lfloor k/2 \rfloor$ candidates and the last $\lceil k/2 \rceil$ candidates of the order that we describe above, then for each $i \in [n]$, it holds that $|A_i \cap W| \geq \min\{k/2 - 1, |A_i|\}$, and as $f_i \leq k - 1$, we again get that W is $(2, 0)$ -IR. Now similarly as above, as W consists of the first and the last candidate of the order, W is semi-strong-JR as if $f_i \geq 1$ for some $i \in N$, then i approves some candidate between these two candidates. \square

Theorem 21. *For any WSC profile, there exists a semi-strong JR committee and it can be found in polynomial time.*

Proof. Consider the order $1, \dots, n$ of the voters. Without loss of generality assume that for each $i \in N$, $|A_i| > 1$ (otherwise we can just exclude them) and that $k \geq 2$. Now, let c^* be a candidate that is approved by voter 1 and let $i > 1$ be the first voter in the order that does not approve c^* . This means that i approves a candidate c' different from c^* , and the profile is WSC if all the voters from i to n approve c' . Hence, any committee $W \supseteq \{c^*, c'\}$ provides semi-strong JR as all the voters are represented by at least one candidate. \square

Proposition 22. *There exists an instance that is VEI, CEI and WSC, and does not admit a $(2 - 2/k, 0)$ -IR committee.*

Proof. Consider the instance defined in the proof of Theorem 14. We now show that the profile is CEI, VEI and WSC, from which the lemma follows. Indeed, for VEI, if we order the candidates as c_1, c_2, \dots, c_m , then each A_i forms a prefix or a suffix of the ordering. For CEI, if we order the voters as $1, 2, \dots, n$, then the voters that approve each candidate form a prefix or a suffix of the ordering. Under the same ordering of the voters, note that for each candidate $c \in C$ we have $c \in A_i$ for either all $i \in [n - 1]$ or for all $i \in [n] \setminus \{1\}$. Thus, the profile is also WSC. \square

DUE

Definition 10 (Dichotomous Uniformly Euclidean (DUE)). *An approval profile A satisfies Dichotomous uniformly Euclidean (DUE) if there is a mapping of voters and candidates into the real line and a radius r such that every voter*

$i \in N$ approves the candidates that are at most r far from her.

Proposition 23. *There exists a DUE profile which does not admit a semi-strong JR committee.*

Proof. Consider the following instance with $n = 6$ voters and assume $k = 3$:

$$\begin{aligned} A_1 &= \{c_1\} \\ A_2 &= \{c_1, c_2\} \\ A_3 &= \{c_2\} \\ A_4 &= \{c_3\} \\ A_5 &= \{c_3, c_4\} \\ A_6 &= \{c_4\}. \end{aligned}$$

To see that this is a DUE profile, consider the following mapping of the instance onto the real line. Each voter $i \in [6]$ is mapped to the point i and candidates c_1 to c_4 are mapped to 1.5, 2.5, 4.5 and 5.5, respectively. From this mapping we obtain the above profile by using an approval radius of 0.5. We observe $f_i = 1$ for all $i \in [6]$, but for any committee W of size 3 it holds that $|A_i \cap W| = 0$ for some $i \in \{1, 3, 4, 6\}$. \square

α -Tree Representation

Definition 11 (α -tree representation (α -TR)). *An approval profile A satisfies α -tree representation (α -TR) if there exists a rooted tree T with vertices $C \cup \{x\}$ and root $x \notin C$, such that for every voter $i \in N$ there is a candidate $c \in C$ such that A_i equals the set of vertices on the path from x to c (without x but including c).*

Under α -TR preferences, the following algorithm always selects a committee that provide individual representation. Here we use $\text{dist}(c, x)$ to denote the (edge-)distance between the root x and a candidate $c \in C$.

Algorithm 3: Individual representation for α -TR

```

1:  $W \leftarrow \emptyset$ 
2: traverse the tree in any order
3: for  $c \in T$  do
4:   if  $|N_c| \geq \frac{n}{k} \cdot \text{dist}(c, x)$  then
5:      $W \leftarrow W \cup \{c\}$ 
6: if  $|W| < k$  then
7:   fill up  $W$  arbitrarily
8: return  $W$ 

```

Note that this algorithm basically resembles HareAV as defined by Aziz et al. (2017) with a specific tie-breaking mechanism that depends on the tree representation. Due to this tie-breaking, it is clearly polynomial time computable.

Theorem 24. *Under α -TR preferences, a committee providing individual representation always exists and can be found in polynomial time.*

Proof. First, we show that the committee output by Algorithm 3 is of size k and secondly we show that it indeed provides individual representation.

Let $T[W]$ be the subgraph of T induced by the candidate set W chosen by Algorithm 3. By construction, $T[W]$ is a subtree of T with root x . Further, in accordance with the above algorithm we can assign the edges in $T[W]$ pairwise disjoint sets of $\frac{n}{k}$ voters (we might assign a voter fractionally to more than one edge if $\frac{n}{k}$ is not an integer). We assign those voters that were necessary to also pick the second endpoint of the edge in the loop of the algorithm (Lines 4 and 5). If for a candidate $c \in T$ we had $\frac{1}{n} \cdot |N_c| \geq \ell \cdot \frac{\text{dist}(c,x)}{k}$ for some $\ell \geq 2$, then we only pick $\frac{n}{k}$ voters out of N_c so that for a possible neighbor c' of c further away from the root with $\frac{1}{n} \cdot |N_{c'}| \geq \frac{\text{dist}(c',x)}{k}$ we still have at least $\frac{n}{k}$ voters to assign. Thus $T[W]$ can have at most k edges and in turn can have at most k nodes besides the root x . Thus, $|W| = k$ when the algorithm terminates.

To show that Algorithm 3 provides IR committees, consider a group of voters $V \subseteq N$ such that $|V| \geq \ell \cdot \frac{n}{k}$ and $|\bigcap_{i \in V} A_i| \geq \ell$ for some $\ell \in \mathbb{N}$. By the definition of α -TR there is a path of (edge-)length ℓ from x to some $c \in C$ such that the candidates on that path (including c) are a subset of $\bigcap_{i \in V} A_i$ of size ℓ . We call the set of these candidates $A_V \subseteq \bigcap_{i \in V} A_i$. Every candidate in A_V is approved by all of V , i.e., by at least $\ell \cdot \frac{n}{k}$ voters, and is at distance $\leq \ell$ from the root of T . Thus Algorithm 3 chooses all candidates in A_V and therefor for the set of chosen candidates W it holds that $|A_i \cap W| \geq \ell$ for each voter $i \in V$. Thus, a committee providing IR always exists for α -TR. \square

α -VTR

Definition 12 (α -vertex tree representation (α -VTR)). *An approval profile A satisfies α -vertex tree representation (α -VTR) if there exists a rooted tree T with vertices $V \cup \{x\}$ and root $x \notin V$, such that for every candidate $c_j \in C$ there exists a voter approving c_j such that the set of voters $N(\{c_j\})$ form a path from x to that voter (without x but including the voter).*

Proposition 25. *There exists an α -VTR instance that does not admit an (α, β) -IR committee for $\beta < k - 1$ and any $\alpha \geq 1$.*

Proof. We note that the instance in Theorem 3 is also a α -VTR instance, from which the statement follows. To see this, consider a rooted tree with root x and vertices $v \in V$ as follows: the voters $v_{\frac{n}{k}+1}, \dots, v_n$ form a simple path starting from x . All the remaining voters $v_1, \dots, v_{\frac{n}{k}}$ are incident to the vertex farthest away from x and make up the leaves of the tree. The set of voters approving a common candidate in the above instance now form a path from x to one of the leaves $v_1, \dots, v_{\frac{n}{k}}$ and thus this is an α -VTR representation of that instance. \square

Note that the above α -VTR instance also does not admit a committee providing semi-strong JR.

D Additional Details for Section 5

Here we describe the different models we used to generate approval profiles.

Voter Interval Euclidean Our first model is the *voter interval Euclidean model* (VI). Here, we choose a location in $[0, 1]$ uniformly at random for each voter and candidate. Further, for each candidate we choose a radius according to $|\mathcal{N}(0, 0.15)|$. A candidate is approved by all voters in its radius.

Candidate Interval Euclidean Our second model is the *candidate interval Euclidean model* (CI). Here, we again choose a location uniformly at random for each voter and candidate as well as a radius according to $|\mathcal{N}(0, 0.15)|$ for each voter. A voter approves all candidates in its radius.

Two-dimensional Euclidean Here we follow the approach of Peters et al. (2021) for the *two-dimensional Euclidean model* (2D). We uniformly at random generate between one and five centers (with the number of centers being drawn uniformly at random as well) in $[0, 1] \times [0, 1]$ and assign each voter and candidate to one of these centers, also uniformly at random. We then assign each voter and each candidate a position sampled from the normal distribution around their respective center points with standard deviation 0.2. Finally, we assign each candidate a radius sampled from $|\mathcal{N}(0, 0.5)|$ and let each voter approve all candidates within their radius.

Impartial Culture For the *impartial culture model* (IC) each candidate is approved by some voter with probability 0.15.

Urn Models For the *urn model* we again follow the approach of Peters et al. (2021) and use the urn model profile generation of the *abc voting* package (Lackner et al. 2021) with approval size uniformly at random from 1 to $\frac{m}{4}$ and replacement value uniformly at random from 1 to $\frac{m}{4}$.

Mallows Model Following the approach of Peters et al. (2021) again, we use the *abc voting* package (Lackner et al. 2021) to first generate a ranking for each voter according to the mixture of three mallows models generated from three uniform at random base-rankings with each ϕ value being chosen uniformly at random in $[0, 1]$ as well. Finally, each voter approves a prefix of size chosen between $[1, \frac{m}{4}]$, again uniformly at random.