

IDENTIFICATION OF A REVERSIBLE CATENARY SYSTEM BY THE MEASURE OF ONE COMPARTMENT

B. FADIS¹, B. HEBRI², AND S.KHELIFA³

ABSTRACT. The known results on compartment system's identification requires observations and measures on some compartments. However, in many situations some compartments can not be measured, so the minimization of the compartment's number to be measured is a very important problem to solve. In this work, this problem had been solved for a class of compartmental systems. It is proved that all reversible catenary system is identifiable by the measure of one compartment : the principal compartment.

KEYWORDS AND PHRASES. Differential systems; Compartmental analysis; Catenary systems; Will problems; Parameters identification.

1. INTRODUCTION


Catenary systems represents an important class in compartment systems. They are in particular used for modeling transport process through successive biological structures like, for example, the diffusion of some drugs in human body when they are injected and diffused in the blood and finally eliminated by urinary tract. In homogenous media, modeling by using catenary systems can be very convenient to give a discrete representation of some process. Catenary systems can constitute a simplification of a complex process to estimate cellular kinetic parameters (see Golgi [6]).

Catenary systems are also used in ecology for the representation of an age structured population. Some equivalents models had been proposed in epidemiology for the Bilharzia (see Cohen & Rochet [5]). All known identification's methods for this type of models are essentially based on empiric algorithms requiring measurements on several compartments see [1],[2],[3],[4], to the different posed problems in compartmental analysis and their best comprehension.

2. PRELIMINARY RESULTS

2.1. Some generalities on catenary systems.

Let n be an integer greater than 3. A catenary system is a compartmental system which can be drafted by :



The exchange coefficients k_{ij} for this kind of systems are characterized by:

$$\begin{cases} k_{ij} = 0, \forall (i, j) \in \{1, 2, \dots, n\}^2, |i - j| \geq 2, \\ k_{i(i+1)} \neq 0; \forall i \in \{1, 2, \dots, n - 1\}. \end{cases}$$

Catenary systems can be classified into two classes:

- the reversible systems characterized by:

$$k_{(j+1)j} \neq 0, \forall j \in \{1, 2, \dots, n - 1\},$$

- the irreversible systems characterized by:

$$\exists j_0 \in \{1, 2, \dots, n - 1\} \text{ such that } k_{(j_0+1)j_0} = 0.$$

These systems are governed by the following differential system noted (S1), given by:

$$(S1) \quad \begin{cases} x'_1(t) = k_{11}x_1(t) + k_{21}x_2(t), \\ x'_j(t) = k_{(j-1)j}x_{j-1}(t) + k_{jj}x_j(t) + k_{(j+1)j}x_{j+1}(t), \forall j \in \{2, 3, \dots, n - 1\}, \\ x'_n(t) = k_{(n-1)n}x_{n-1}(t) + k_{nn}x_n(t), \end{cases}$$

where

$$\begin{cases} k_{11} = -(k_{12} + k_{1e}), \\ k_{jj} = -(k_{j(j-1)} + k_{j(j+1)}), \forall j \in \{2, 3, \dots, n - 1\}, \\ k_{nn} = -k_{n(n-1)}. \end{cases}$$

We associate the matrix A , $A = (k_{ij})_{1 \leq i, j \leq n}$ to this open n -compartmental system. We can remark that A is a tridiagonal and compartmental matrix so its eigenvalues noted λ_i for $i \in \{1, 2, \dots, n\}$ are reals, strictly negatives and different (see [8]). The general solution of the system (S1) can be written under the form:

$$x_j(t) = \sum_{i=1}^n \beta_i^j \exp(\lambda_i t), \forall j \in \{1, 2, \dots, n\},$$

where $(\beta_i^j)_{1 \leq i \leq n}$ is the column j of the matrix B of elementaries masses associated to the compartment j (see [8]).

Measuring the primary compartment (the catalyser of the reaction) we can determine the coefficients $\beta_1^1, \beta_2^1, \dots, \beta_n^1, \lambda_1, \lambda_2, \dots, \lambda_n$, noted by $\beta_1^{1*}, \beta_2^{1*}, \dots, \beta_n^{1*}, \lambda_1^*, \lambda_2^*, \dots, \lambda_n^*$ after minimization of the functional J introduced by Y. Cherruault (see [3]):

$$J(\beta_1^1, \beta_2^1, \dots, \beta_n^1, \lambda_1, \lambda_2, \dots, \lambda_n) = \sum_{j=1}^m \sum_{i=1}^n (\beta_i^1 \exp(\lambda_i t_j) - \bar{x}_1(t_j))^2,$$

where $\bar{x}_1(t_j)$ is the measure of the primary compartment at time t_j for j in $\{1, 2, 3, \dots, m\}$. These coefficients will be noted : $\beta_1^{1*}, \beta_2^{1*}, \dots, \beta_n^{1*}, \lambda_1^*, \lambda_2^*, \dots, \lambda_n^*$.

Thus, the general solution of the system (S1) can be written:

$$\begin{aligned} x_1(t) &= \sum_{i=1}^n \beta_i^{1*} \exp(\lambda_i^* t), \\ x_j(t) &= \sum_{i=1}^n \beta_i^j \exp(\lambda_i^* t), \quad \forall j \in \{2, 3, \dots, n\}, \end{aligned}$$

where the coefficients β_i^j for $j \in \{2, 3, \dots, n\}$ and $i \in \{1, 2, \dots, n\}$ remain unknowns.

In order to identify the all compartmental system, an injection of substance is made in the primary compartment. So the system (S1) is completed by the initial conditions:

$$x_1(0) = a \quad \text{and} \quad x_j(0) = 0, \quad \forall j \in \{2, 3, \dots, n\}.$$

Hence

$$(2.1) \quad \begin{cases} \sum_{i=1}^n \beta_i^{1*} = x_1(0) = a, \\ \sum_{i=1}^n \beta_i^j = x_j(0) = 0, \quad \forall j \in \{2, 3, \dots, n\}. \end{cases}$$

The linear independency of the functions φ_i , $i \in \{1, 2, \dots, n\}$ defined by:

$$\varphi_i(t) = \exp(\lambda_i^* t), \quad \forall t \geq 0, \quad \forall i \in \{1, 2, \dots, n\},$$

gives, for all integer i in $\{1, 2, \dots, n\}$:

$$(2.2) \quad \begin{cases} \lambda_i^* \beta_i^{1*} = k_{11} \beta_i^{1*} + k_{21} \beta_i^2, \\ \lambda_i^* \beta_i^j = k_{(j-1)j} \beta_i^{j-1} + k_{jj} \beta_i^j + k_{(j+1)j} \beta_i^{j+1}, \quad \forall j \in \{2, 3, \dots, n-1\}, \\ \lambda_i^* \beta_i^n = k_{(n-1)n} \beta_i^{n-1} + k_{nn} \beta_i^n. \end{cases}$$

In the following we need to introduce the following notation, for integer j in $\{1, 2, \dots, n\}$ and integer l in $\{0, 1, \dots, n\}$, let Δ_j^l be:

$$\Delta_j^l = \sum_{i=1}^n (\lambda_i^*)^l \beta_i^j.$$

2.2. Preliminary results.

With the previous notations, we have the following results:

Lemma 2.1. *The following relationships hold:*

$$(2.3) \quad \begin{aligned} &i) \\ &\Delta_1^0 = a \quad \text{and} \quad \Delta_j^0 = 0, \quad \forall j \in \{2, 3, \dots, n\}. \end{aligned}$$

$$(2.4) \quad \begin{aligned} &ii) \\ &\Delta_1^1 = k_{11} a. \end{aligned}$$

iii) $\forall j \in \{2, 3, \dots, n-1\}$,

$$(2.5) \quad \Delta_j^l = k_{(j-1)j} \Delta_{j-1}^{l-1} + k_{jj} \Delta_j^{l-1} + k_{(j+1)j} \Delta_{j+1}^{l-1}, \quad \forall l \in \{1, 2, \dots, j\}.$$

iv)

$$(2.6) \quad \Delta_n^l = k_{(n-1)n} \Delta_{n-1}^{l-1} + k_{nn} \Delta_n^{l-1}, \quad \forall l \in \{1, 2, \dots, n\}.$$

Proof.

i) Δ_j^0 represent initial conditions then the results hold.

ii) We recall that:

$$\Delta_1^1 = \sum_{i=1}^n (\lambda_i^*)^1 \beta_i^1 = \sum_{i=1}^n \lambda_i^* \beta_i^{1*}.$$

Then due to the relationship (2.2) we can write that:

$$\Delta_1^1 = \sum_{i=1}^n (k_{11} \beta_i^{1*} + k_{21} \beta_i^2).$$

Or

$$\Delta_1^1 = k_{11} \sum_{i=1}^n \beta_i^{1*} + k_{21} \sum_{i=1}^n \beta_i^2.$$

Consequently :

$$\Delta_1^1 = k_{11} a.$$

iii) Let j and l be two integers such that $j \in \{2, 3, \dots, n-1\}$ and $l \in \{1, 2, \dots, j\}$, we have:

$$\Delta_j^l = \sum_{i=1}^n (\lambda_i^*)^l \beta_i^j = \sum_{i=1}^n (\lambda_i^*)^{l-1} \lambda_i^* \beta_i^j,$$

then, the relationship (2.2) gives:

$$\Delta_j^l = \sum_{i=1}^n (\lambda_i^*)^{l-1} \left(k_{(j-1)j} \beta_i^{j-1} + k_{jj} \beta_i^j + k_{(j+1)j} \beta_i^{j+1} \right),$$

or

$$\Delta_j^l = k_{(j-1)j} \sum_{i=1}^n (\lambda_i^*)^{l-1} \beta_i^{j-1} + k_{jj} \sum_{i=1}^n (\lambda_i^*)^{l-1} \beta_i^j + k_{(j+1)j} \sum_{i=1}^n (\lambda_i^*)^{l-1} \beta_i^{j+1}.$$

The definition of Δ_q^p let us conclude that:

$$\Delta_j^l = k_{(j-1)j} \Delta_{j-1}^{l-1} + k_{jj} \Delta_j^{l-1} + k_{(j+1)j} \Delta_{j+1}^{l-1}.$$

iv) Let l be an integer in $\{1, 2, \dots, n\}$, we have:

$$\Delta_n^l = \sum_{i=1}^n (\lambda_i^*)^l \beta_i^n = \sum_{i=1}^n (\lambda_i^*)^{l-1} \lambda_i^* \beta_i^n.$$

Then, the relationship (2.2) gives:

$$\Delta_n^l = \sum_{i=1}^n (\lambda_i^*)^{l-1} (k_{(n-1)n} \beta_i^{n-1} + k_{nn} \beta_i^n),$$

or

$$\Delta_n^l = k_{(n-1)n} \sum_{i=1}^n (\lambda_i^*)^{l-1} \beta_i^{n-1} + k_{nn} \sum_{i=1}^n (\lambda_i^*)^{l-1} \beta_i^n.$$

The definition of Δ_q^p let us conclude that:

$$\Delta_n^l = k_{(n-1)n}\Delta_{n-1}^{l-1} + k_{nn}\Delta_n^{l-1}.$$

□

Lemma 2.2. *Let n be an integer strictly greater than 4. Then:*

$$\Delta_j^l = 0, \quad \forall j \in \{3, 4, \dots, n\}, \quad \forall l \in \{1, 2, \dots, n-2\}, \quad j-l \geq 2.$$

Proof. The proof of this lemma is based on a recurrence process.

i) In the first step we set $l = 1$:

a- let j be an integer in $\{3, 4, \dots, n-1\}$. Owing to the relationship (2.5), we have:

$$\Delta_j^1 = k_{(j-1)j}\Delta_{j-1}^0 + k_{jj}\Delta_j^0 + k_{(j+1)j}\Delta_{j+1}^0,$$

the relationships (2.3) let us conclude that:

$$\Delta_j^1 = 0.$$

b- when $j = n$, the relationship (2.6) gives:

$$\Delta_n^1 = k_{(n-1)n}\Delta_{n-1}^0 + k_{nn}\Delta_n^0,$$

so the relationships(2.3) let us conclude that:

$$\Delta_n^1 = 0.$$

ii) In the last step we assume that for all integer l in $\{2, \dots, n-3\}$ and for all integer j in $\{l+2, \dots, n\}$ we have :

$$\Delta_j^l = 0,$$

a- for any integer j in $\{l+3, \dots, n-1\}$, owing to the relationship (2.5), we have:

$$\Delta_j^{l+1} = k_{(j-1)j}\Delta_{j-1}^l + k_{jj}\Delta_j^l + k_{(j+1)j}\Delta_{j+1}^l,$$

then the recurrence hypothesis permits to affirm that:

$$\Delta_{j-1}^l = \Delta_j^l = \Delta_{j+1}^l = 0,$$

because $j-1-l \geq 2$, $j-l \geq 3$ and $j+1-l \geq 4$, so we can conclude that:

$$\Delta_j^{l+1} = 0.$$

b- for $j = n$, the same process gives:

$$\Delta_n^{l+1} = 0.$$

□

Lemma 2.3. *For all integer l in $\{1, 2, \dots, n-1\}$ the following relationship holds:*

$$\Delta_{l+1}^l = a \left(\prod_{i=1}^l k_{i(i+1)} \right).$$

Proof. To show this result we use a recurrence process like in the previous lemma.

1° For $l = 1$, we have:

$$\Delta_2^1 = \sum_{i=1}^n \lambda_i^* \beta_i^2,$$

due to the relationship (2.5), we have:

$$\Delta_2^1 = k_{12}\Delta_1^0 + k_{22}\Delta_2^0 + k_{32}\Delta_3^0.$$

Using relationship (2.3) we can conclude:

$$\Delta_2^1 = a.k_{12} + 0.k_{22} + 0.k_{32} = a.k_{12}.$$

2° Assume that for all integer l in $\{2, \dots, (p-1)\}$ with $p \leq (n-2)$ we have:

$$\Delta_{l+1}^l = a \left(\prod_{i=1}^l k_{i(i+1)} \right).$$

By definition, we know that:

$$\Delta_{p+1}^p = \sum_{i=1}^n (\lambda_i^*)^p \beta_i^{p+1},$$

relationship (2.5), gives:

$$\Delta_{p+1}^p = k_{p(p+1)}\Delta_p^{p-1} + k_{(p+1)(p+1)}\Delta_{p+1}^{p-1} + k_{(p+2)(p+1)}\Delta_{p+2}^{p-1}.$$

As Lemma 2.2:

$$\Delta_{p+1}^{p-1} = \Delta_{p+2}^{p-1} = 0,$$

so we can conclude that:

$$\Delta_{p+1}^p = a \left(\prod_{i=1}^p k_{i(i+1)} \right).$$

3° When $l = n-1$, using relationship (2.6), Lemma 2.2 and the recurrence hypothesis, we obtain:

$$\Delta_n^{n-1} = a.k_{(n-1)n} \cdot \left(\prod_{i=1}^{n-2} k_{i(i+1)} \right) + 0.k_{nn},$$

or else:

$$\Delta_n^{n-1} = a \left(\prod_{i=1}^{n-1} k_{i(i+1)} \right).$$

□

Remark 2.4. Knowing that $a \neq 0$, from Lemma 2.3 it becomes:

$$\Delta_{l+1}^l \neq 0, \forall l \in \{1, 2, \dots, n-1\}.$$

3. MAIN RESULT

3.1. General relationships.

The main result can be obtained as a consequence of the two following propositions. The compartmental matrix A and the matrix B of elementaries masses (see [8]) will be alternately identified, respectively row after row and column after column.

Proposition 3.1. *If the catenary system (S1) is reversible, then the following relationships hold:*

i)

$$k_{11} = \frac{\sum_{i=1}^n \lambda_i^* \beta_i^{1*}}{a}.$$

ii)

$$k_{21} = \frac{\Delta_1^2 - k_{11} \Delta_1^1}{a k_{12}}.$$

iii)

$$\beta_i^2 = \frac{\lambda_i^* \beta_i^{1*} - k_{11} \beta_i^{1*}}{k_{21}}, \quad \forall i \in \{1, 2, \dots, n\}.$$

Proof.

i) Knowing that :

$$\Delta_1^1 = \sum_{i=1}^n \lambda_i^* \beta_i^{1*},$$

relationship(2.4) gives directly the result.

ii) The relationship (2.5) used for $l = 2$, $j = 1$ gives:

$$\Delta_1^2 = k_{11} \Delta_1^1 + k_{21} \Delta_2^1,$$

and

$$k_{21} = \frac{\Delta_1^2 - k_{11} \Delta_1^1}{\Delta_2^1} = \frac{\Delta_1^2 - k_{11} \Delta_1^1}{a k_{12}}.$$

iii) Recall that the relationship (2.2) is given, after minimization of J , by:

$$\lambda_i^* \beta_i^{1*} = k_{11} \beta_i^{1*} + k_{21} \beta_i^2, \quad \forall i \in \{1, 2, \dots, n\},$$

with a simple manipulation , we get :

$$\beta_i^2 = \frac{\lambda_i^* \beta_i^{1*} - k_{11} \beta_i^{1*}}{k_{21}}, \quad \forall i \in \{1, 2, \dots, n\}.$$

□

Remark 3.2. *This result says that the second column of the matrix B is identified.*

Proposition 3.3. *If the catenary system (S1) is reversible, then the following relationships hold:*

i)

$$k_{jj} = \frac{\Delta_j^j - k_{(j-1)j} \Delta_{j-1}^{j-1}}{\Delta_j^{j-1}}, \quad \forall j \in \{2, 3, \dots, n-1\}.$$

ii)

$$k_{(j+1)j} = \frac{\Delta_j^{j+1} - k_{(j-1)j}\Delta_{j-1}^j - k_{jj}\Delta_j^j}{\Delta_{j+1}^j}, \quad \forall j \in \{2, 3, \dots, n-1\}.$$

iii)

$$\beta_i^{j+1} = \frac{\lambda_i^* \beta_i^j - k_{(j-1)j}\beta_i^{j-1} - k_{jj}\beta_i^j}{k_{(j+1)j}}, \quad \forall i \in \{1, 2, \dots, n\}.$$

Proof. We recall that:

$$\Delta_{p+1}^p \neq 0, \quad \forall p \in \{1, 2, \dots, n-1\}.$$

Let j be an integer in $\{2, 3, \dots, n-1\}$.

i) The relationship (2.5) with $l = j$, gives:

$$\Delta_j^j = k_{(j-1)j}\Delta_{j-1}^{j-1} + k_{jj}\Delta_j^{j-1} + k_{(j+1)j}\Delta_{j+1}^{j-1}.$$

Lemma 2.2 indicates that:

$$\Delta_{j+1}^{j-1} = 0,$$

so

$$k_{jj} = \frac{\Delta_j^j - k_{(j-1)j}\Delta_{j-1}^{j-1}}{\Delta_j^{j-1}}.$$

ii) Relationship (2.5) when $l = j+1$, gives:

$$\Delta_j^{j+1} = k_{(j-1)j}\Delta_{j-1}^j + k_{jj}\Delta_j^j + k_{(j+1)j}\Delta_{j+1}^j.$$

By a simple manipulation of this equality, we obtain:

$$k_{(j+1)j} = \frac{\Delta_j^{j+1} - k_{(j-1)j}\Delta_{j-1}^j - k_{jj}\Delta_j^j}{\Delta_{j+1}^j}.$$

iii) The relationship (2.2) for all integer $i \in \{1, 2, \dots, n\}$ being:

$$\lambda_i^* \beta_i^j = k_{(j-1)j} \cdot \beta_i^{j-1} + k_{jj} \beta_i^j + k_{(j+1)j} \beta_i^{j+1},$$

with a simple manipulation of this equality we get:

$$\beta_i^{j+1} = \frac{\lambda_i^* \beta_i^j - k_{(j-1)j} \beta_i^{j-1} - k_{jj} \beta_i^j}{k_{(j+1)j}}, \quad \forall i \in \{1, 2, \dots, n\}.$$

$k_{(j+1)j} \neq 0$ due to the reversibility of the system. □

3.2. Identification theorem.

Generally in compartmental analysis the identifiability of a n -compartmental system requires the measurement of $(n - 1)$ compartments or $(n - 2)$ compartments under some conditions (see [8]). In catenary systems, we obtain the following optimal result :

Theorem 3.4. *Let $(S1)$ be a reversible catenary compartmental system. If the primary compartment is measurable then $(S1)$ is identifiable.*

Proof. This theorem is a consequence of Proposition 3.1 and Proposition 3.3. In fact, owing to Proposition 3.1, we identify $k_{11}, k_{12}, k_{21}, \beta_i^2, \forall i \in \{1, 2, \dots, n\}$ in this order as follow:

- firstly:

$$k_{11} = \frac{\sum_{i=1}^n \lambda_i^* \beta_i^{1*}}{a}.$$

Since,

$$k_{12} = -(k_{1e} + k_{11}),$$

and knowing that (see [10] & [11]):

$$k_{1e} = \frac{-a}{\sum_{i=1}^n \frac{\beta_i^{1*}}{\lambda_i^*}},$$

we can get the value of k_{12} . Consequently, k_{21} can be determined:

$$k_{21} = \frac{\Delta_1^2 - k_{11} \Delta_1^1}{a k_{12}},$$

as Δ_1^2, Δ_1^1 are known because the first column of B is known.

- secondly:

$$\beta_i^2 = \frac{\lambda_i^* \beta_i^{1*} - k_{11} \beta_i^{1*}}{k_{21}}, \quad \forall i \in \{1, 2, \dots, n\}.$$

This means that the first column thus the first row of the matrix A are identified and finally the second column of the matrix B is identified.

We suppose that at the step j , $j \in \{1, 2, \dots, n - 1\}$, the coefficients $k_{pp}, k_{p(p+1)}, k_{(p+1)p}, \beta_i^{(p+1)}$ $\forall i \in \{1, 2, \dots, n\}$, $\forall p \in \{1, 2, \dots, j - 1\}$ had been identified. Owing to Proposition 3.3, we can identify $k_{jj}, k_{j(j+1)}, k_{(j+1)j}, \beta_i^{j+1}$, $\forall i \in \{1, 2, \dots, n\}$ in this order as follows:

$$k_{jj} = \frac{\Delta_j^j - k_{(j-1)j} \Delta_{j-1}^{j-1}}{\Delta_j^{j-1}},$$

and $\Delta_j^j, \Delta_{j-1}^{j-1}, \Delta_j^{j-1}$ are known because the columns $(j - 1)$ and j of the matrix B had been identified above (recurrence hypothesis). Then we can identify $k_{j(j+1)}$ because:

$$k_{j(j+1)} = -(k_{j(j-1)} + k_{jj}),$$

and $k_{(j+1)j}$ is known by recurrence hypothesis. Consequently, $k_{(j+1)j}$ can be determined since:

$$k_{(j+1)j} = \frac{\Delta_j^{j+1} - k_{(j-1)j}\Delta_{j-1}^j - k_{jj}\Delta_j^j}{\Delta_{j+1}^j} = \frac{\Delta_j^{j+1} - k_{(j-1)j}\Delta_{j-1}^j - k_{jj}\Delta_j^j}{k_{12}k_{23} \cdots k_{(j-1)j}},$$

and Δ_j^{j+1} , Δ_{j-1}^j , Δ_j^j are known because the columns $(j-1)$ and j of B had been identified above (recurrence hypothesis).

We have :

$$\beta_i^{j+1} = \frac{\lambda_i^* \beta_i^j - k_{(j-1)j} \beta_i^{j-1} - k_{jj} \beta_i^j}{k_{(j+1)j}}, \quad \forall i \in \{1, 2, \dots, n\},$$

and all the parameters of β_i^{j+1} expression are known, so it can be identified. This means that the column j then the row j of the matrix A are identified and finally the column $j+1$ of the matrix B is identified for all $j \in \{1, 2, \dots, n-1\}$.

To identify k_{nn} , it suffices to remark that:

$$k_{nn} = -k_{n(n-1)},$$

and the coefficient $k_{n(n-1)}$ had been identified above. □

3.3. Identification's algorithm.

The results of the Proposition 3.1 can be used for the algorithm's initialization and the results of the Proposition 3.3 can be used to define iterations which permits to identify at each step the corresponding column and row of the tridiagonal compartmental matrix and the corresponding column of the partial measures matrix.

4. CONCLUSION

The problem of the minimal number of compartment to measure for identify a compartmental system had been totaly solved for reversibles catenarays systems. Moreover, with the obtained results, we can construct an identification algorithm for reversibles catenarays systems with just the measure of the first compartment.

REFERENCES

- [1] Bischi, G.I. . "Compartmental analysis of economic systems with heterogeneous agents: an introduction" in Beyond the Representative Agent (A. Kirman and M. Gallegati Eds.) Elgar Pub. Co, 1998, pp.181-214.
- [2] Cherruault, Y. (1983), *Biomathématiques*, Collection "Que sais-je", 2052, Presses Universitaires de France, (P.U.F) Paris.
- [3] Cherruault, Y. (1986), *Mathematical Modelling in Biomedicine. Optimal control of Biomedical Systems*, Kluwer (Reidel), Dordrecht.
- [4] Cherruault, Y. (1998), *Modèles et méthodes mathématiques pour les sciences du vivant*, Presses Universitaires de France, (P.U.F) Paris.
- [5] Cohen et Marie Joelle Rochet: "pour la bilharziose"
- [6] Golgi : temps de synthèse et durée de transport dans le système de Golgi de cellules de la glande séricigène pour une protéine sécrétée la fibroéine].
- [7] B. Hebri, & Y.Cherruault, , "Contribution to identification of linear compartmental systems.", *Kybernetes*, Vol. 33 No.8, pp. 1277-1291, 2004.

- [8] B.Hebri, & Y.Cherruault, “ Direct identification of general linear compartmental systems by means of $(n-2)$ compartments measures.”, *Kybernetes*, Vol. 34 No7/8, pp. 969-982, 2005.
- [9] B.Hebri, & Y.Cherruault, “New results about identifiability of linear open bicompartamental homogeneous system and identification of open Michaelis-Menten system by a linear approach”, *Kybernetes*, Vol. 34 No7/8, pp. 1159-1186, 2005.
- [10] B.Hebri, & Y.Cherruault, ”Identification of closed Michaelis-Menten system by a linear approach”, *Kybernetes*, Vol. 34 No7/8, pp. 101-1026, 2005.
- [11] J.A. Jacquez, The Inverse Problem for Compartmental Models. Mathematics and Computer in Simulation XXIV, 42-459 (1982).
- [12] Lennart, L. & Torkel, G.L. (1994), *Modelling of dynamic systems*, Prentice Hall.
- [13] Mazumdar, J. (1989), *An introduction to Mathematical Physiology and Biology*, Cambridge Univ. Press.
- [14] Mohammedi, A. (1990), *Systèmes à compartiments déterministes*. Monographie. Ecole Polytechnique Fédérale de Lausanne
- [15] Wagner, J. (1975), *Fundamentals of clinical pharmacokinetics*. Drug Intelligence.

¹HIGHER SCHOOL OF APPLIED SCIENCES OF ALGIERS, ALGERIA.

Email address: ¹badreddine.f@gmail.com

^{2,3}UNIVERSITY OF SCIENCES AND TECHNOLOGY HOUARI BOUMEDIENE (USTHB),
FACULTY OF MATHEMATICS, DEPARTMENT OF CALCULUS, P.B. 32 EL-ALIA, 16111, BAB EZZOUAR, ALGIERS,
ALGERIA.

Email address: ²bhebri@usthb.dz, ³ksices@yahoo.fr