

Superspace BRST/BV operators of superfield gauge theories

I. L. Buchbinder^{1a,b}, S. James Gates Jr.^{2c,d}, K. Koutrolikos^{3c,d}

^a*Center of Theoretical Physics, Tomsk State Pedagogical University,
5 Tomsk 634041, Russia*

^b*Laboratory for Theoretical Cosmology, International Center for Gravity and Cosmos,
Tomsk State University of Control Systems and Radioelectronics (TUSUR),
634050, Tomsk, Russia*

^c*Brown Theoretical Physics Center,
Box S, 340 Brook Street, Barus Hall, Providence, RI 02912, USA*

^d*Department of Physics, Brown University,
Box 1843, 182 Hope Street, Barus & Holley 545, Providence, RI 02912, USA*

ABSTRACT

We consider the superspace BRST and BV description of $4D$, $\mathcal{N} = 1$ Super Maxwell theory and its non-abelian generalization Super Yang-Mills. By fermionizing the superspace gauge transformation of the gauge superfields we define the nilpotent superspace BRST symmetry transformation (\mathfrak{J}). After introducing an appropriate set of anti-superfields and define the superspace anti-bracket, we use it to construct the BV-BRST nilpotent differential operator (\mathfrak{S}) in terms of superspace covariant derivatives. The anti-superfield independent terms of \mathfrak{S} provide a superspace generalization of the Koszul-Tate resolution (\mathfrak{D}). In the linearized limit, the set of superspace differential operators that appear in \mathfrak{S} satisfy a nonlinear algebra which can be used to construct a BRST charge Q without requiring pure spinor variables. Q acts on the Hilbert space of superfield states and its cohomology generates the expected superspace equations of motion.

¹joseph@tspu.edu.ru

²sylvester_gates@brown.edu

³konstantinos_koutrolikos@brown.edu

1 Introduction

The Becchi-Rouet-Stora-Tutin (BRST) formalism offers a modern approach of describing gauge theories. The power of this technique has been demonstrated repeatedly from the formulation of field theories that correspond to given first quantized systems all the way to string field theory and the study of string interactions.

Initially, BRST symmetry [1–4] was introduced as a method of quantizing gauge field theories. In general, the quantization procedure of gauge theories is not straightforward and requires the involvement of ghost fields. The two basic characteristics of gauge theories that lead to the introduction of ghosts are (a) gauge redundancy and (b) the tensorial nature of gauge fields. Specifically, one must eliminate the non-dynamical degrees of freedom of gauge fields and also eliminate the states with non-positive definite norm which correspond to the time components⁴ of gauge fields. The first was completely understood by the Fadeev-Popov procedure from the view point of a path-integral integration measure effect. Both these issues are addressed in the BRST quantization procedure which systematically generates not only the Fadeev-Popov ghosts—which give rise to appropriate gauge fixing conditions—but also additional propagating ghosts (non-minimal sector) that cancel the negative norm states.

This approach was later extended to the Batalin-Vilkovisky (BV) [5–7] formalism which promotes the above BRST symmetry to a fundamental principal of the theory which is automatically incorporated by the introduction of additional fields, the antifields. The antifields have the interpretation of sources of BRST transformations and were initially introduced as a technique to capture the renormalization of the composite operators emerged in the BRST transformations of interacting gauge theories. However, antifields play a crucial role whether one studies quantum or classical aspects of the theory. They allow the definition of a symplectic structure in the space of fields and antifields, the antibracket. The antibracket can be understood as a covariant analog of the Hamiltonian Poisson bracket and as a consequence various Hamiltonian concepts (e.g. canonical transformations) can be introduced and applied [8–10]. This makes the field-antifield formalism a very powerful tool which can be used in cases where the usual Fadeev-Popov method may fail. An important class of such theories are the ones with open symmetry algebras, meaning that the commutator of two symmetries generates trivial symmetries. In general, supersymmetric theories fall in this class, unless one considers theories with off-shell supersymmetry, like the ones that have a superspace description⁵.

Moreover, the antibracket formalism gained popularity among string theorists when it was applied to the bosonic open string field theory [12–15] and later to closed string field theory [16]. This proved to be very useful for the study of string interactions and various string vacua containing non-perturbative objects like D-branes [17]. The BRST approach to string field theory originated in the work of Siegel [18] where a nilpotent, fermionic BRST operator is constructed which commutes with observables, acts on the state space of the theory and its cohomology defines the physical spectrum. This state space BRST operator is a reflection of the BRST symmetry of the target space fields as generated from the antibracket in the Lagrangian path integral formulation. Interestingly, similar state space BRST operators have been constructed to describe higher spin gauge fields [19–34]. This

⁴We use the mostly plus convention.

⁵See the relevant discussion in [11].

is another indication of the close relation between higher spins and string theory.

Finally, for maximally supersymmetric theories, it has been shown that a covariant quantization can be achieved by using pure spinor variables [35] and the spectrum of these theories is captured by the cohomology of a pure spinor BRST operator [36]. The free action was constructed [37] in terms of a pure spinor superfield⁶ $\Psi(x, \theta, \lambda)$ and the pure spinor BRST operator $Q = \lambda^\alpha D_\alpha$. However, these descriptions were not manifestly maximally supersymmetric and only worked for linearized theories. Later, both issues were addressed by the introduction of additional –non-minimal– pure spinors that modified the BRST charge operator [38] and by adopting the BV formalism and solving the master equation [39, 40].

The purpose of this paper is to explore the BRST symmetry, BV and BFV (Batalin-Fradkin-Vilkovisky) formulations of supersymmetric gauge theories⁷ from the superspace point of view without the use of pure spinor variables. Similarly to string theory applications, we construct a superspace BRST operator acting on the (target) space of superfields and a corresponding BRST operator that acts on a Hilbert space of superfield states.

Following the BRST procedure, for a given supersymmetric gauge theory with a superspace description we introduce a nilpotent, BRST symmetry operator \mathfrak{J} ($\mathfrak{J}^2 = 0$) by fermionizing the superspace gauge transformations of the theory via the replacement of all gauge parameter superfields with corresponding ghost superfields of opposite statistics. The superspace Lagrangian is deformed by the addition of an appropriate \mathfrak{J} -exact term which includes additional (non-minimal sector) ghost superfields like the Nakanishi-Lautrup ghost superfield and its BRST-doublet partner.

In the BV description, for every superfield and ghost superfield we introduce a conjugate variable, the anti-superfield. In the superfield/anti-superfield space we define the superspace antibracket and use it to construct a nilpotent, superspace differential operator \mathfrak{S} ($\mathfrak{S}^2 = 0$). This is a superspace BRST differential operator which action on (anti-) superfields can be decomposed into two pieces: $\mathfrak{S} = \gamma + \delta$. The γ part is the now renamed superspace BRST symmetry and its action on superfields coincides with \mathfrak{J} . The second part δ is the superspace generalization of the Koszul-Tate resolution [43] which implements the superspace equations of motion. As expected both γ and δ are nilpotent and anticommute.

Finally, getting inspiration from string field theory, we define a Hilbert space of states where the arbitrary state vector $|\Psi\rangle$ can be expanded in some basis $|\phi_s\rangle$ with superfield coefficients χ_s ($|\Psi\rangle = \sum_s |\phi_s\rangle \chi_s$). In this Hilbert space, we define a nilpotent BRST operator Q ($Q^2 = 0$), such that its action on the Hilbert space vectors coincides with the action of \mathfrak{S} on the superfield coefficients. This is done in two steps: (1) we identify the set of differential operators that appear in \mathfrak{S} and (2) find their algebraic properties in order to construct a nilpotent Hilbert space BRST operator via the Fradkin-Fradkina algorithm [44]. The cohomology of Q must correctly generate the superspace equations of motion and gauge transformations for the superfields of the theory and thus produce the expected physical spectrum.

We would like to emphasize that although the existence of such BRST charge is proven for general gauge theories [45–48], the explicit construction of it can be a nontrivial task. For supersymmetric theories with a superspace description the only available methodology for constructing

⁶A superfield with an additional dependence on a pure spinor variable λ

⁷See [41, 42] for earlier considerations.

the BRST charge is based on pure spinors. In the pure spinor approach, the nilpotence of the BRST charge is based on the pure spinor constraint and not on the algebraic properties of the characteristic set of differential operators that participate in the description of the theory. Our methodology can offer an alternative method of constructing such nilpotent charges. This could be beneficial for the construction of a manifestly supersymmetric BRST charge that describes higher spin supermultiplets.

The proposed procedure can be applied to any gauge theory in superspace. In this paper, we demonstrate it for $4D$, $\mathcal{N} = 1$ Super Yang-Mills (SYM) and its linearized limit, the vector supermultiplet. We explicitly construct the superspace BRST symmetry operator \mathfrak{s} and extract the superspace Koszul-Tate complex δ . For the linearized theory, the set of superspace differential operators that appear in \mathfrak{s} satisfy a nonlinear algebra. Nevertheless, using known corrections [49, 50] to the Fradkin-Fradkina algorithm we find an explicit expression for Q in terms of supersymmetric covariant derivatives without the need to introduce pure spinors. The cohomology of this superspace BRST charge correctly generates the gauge symmetry and superspace equations of motion for the vector supermultiplet.

Before discussing all the above, we first remind to the reader the BRST, BV and BFV descriptions in the context of a simple example: Maxwell theory. This corresponds to the bosonic sector of the linearized SYM and it is the simplest example where the non-supersymmetric shadows ($\mathcal{J}, \mathfrak{s}, \gamma, \delta, Q$) of the quantities discussed above can be introduced and easily constructed. Moreover, it is explicitly shown that there is a Hilbert space of field states and a nilpotent charge Q such that the action of the BRST operator \mathfrak{s} , generated by the antibracket, on fields and antifields components of the field states, is equal to the action of Q on the basis vectors of the Hilbert state $\{|e_i\rangle\}$: $\mathfrak{s}|\psi\rangle = Q|\psi\rangle$. The BRST charge Q constructed this way is identical to the Batalin-Fradkin-Vilkovisky (BFV) nilpotent charge constructed from the algebra of the physical constraints that define the on-shell propagating degrees of freedom of the theory.

2 Review of BRST, BV and BFV descriptions of Maxwell theory

In the Lagrangian path integral approach, the BRST method is based on the fact that if the theory under consideration is a gauge theory, then there is a nilpotent, odd BRST symmetry (\mathcal{J}) and one can deform the Lagrangian of the theory by the addition of an \mathcal{J} -exact term without affecting the path integral

$$\mathcal{L} \rightarrow \mathcal{L} + \mathcal{J}\Omega . \tag{1}$$

The deformed action remains invariant

$$\mathcal{J}\mathcal{L} + \mathcal{J}^2\Omega = 0 \tag{2}$$

because both terms independently vanish. The second term vanishes because \mathcal{J} must be nilpotent ($\mathcal{J}^2 = 0$) and the first term vanishes because we choose the BRST transformation of the fields that appear in the starting Lagrangian \mathcal{L} to have exactly the same form as their corresponding gauge transformations. Therefore, the Bianchi identities that guarantee the gauge invariance of \mathcal{L} will also guarantee \mathcal{J} -invariance. The tension between \mathcal{J} being an odd symmetry and also having the same structure as the gauge transformations is resolved by the replacement of every gauge

parameter by a ghost field of opposite statistics. This way one can immediately define the BRST symmetry transformation of any gauge field. This is usually referred to as fermionization of the gauge transformation. For the case of Maxwell's theory, we get:

$$\delta A_m = \partial_m \lambda \rightarrow \mathcal{J} A_m = \partial_m c \quad (3)$$

where c is a ghost field. Moreover, because of the nilpotence of this BRST transformation, one determines the BRST transformation of the ghost field:

$$\mathcal{J}^2 A_m = \partial_m (\mathcal{J} c) = 0 \Rightarrow \mathcal{J} c = 0 . \quad (4)$$

Because the deformation (1) of the Lagrangian is \mathcal{J} -exact, the theory is independent of any particular choice of Ω . In the context of quantizing the theory one selects this deformation such that it generates an appropriate gauge fixing condition by integrating out a ghost field, as expected from the Fadeev-Popov procedure. For Maxwell's theory usually one chooses the Landau-Fermi gauge or more generally the Feynman gauge:

$$\mathcal{L} \rightarrow \mathcal{L} + \rho (\partial^m A_m + \frac{\xi}{2} \rho) \quad (5)$$

where ρ is a bosonic ghost field and ξ is the usual Feynman gauge parameter. However, this deformation fails to satisfy the consistency condition of being \mathcal{J} -closed:

$$\mathcal{J} [\rho (\partial^m A_m + \frac{\xi}{2} \rho)] = \mathcal{J}(\rho) (\partial^m A_m + \xi \rho) + \rho \square c \neq 0 . \quad (6)$$

Even if we choose $\mathcal{J}\rho = 0$, there is no way to cancel the second term, hence in order for the deformation to be consistent we must add additional terms. It is straightforward to check that the complete answer is:

$$\mathcal{J}\Omega = \rho (\partial^m A_m + \frac{\xi}{2} \rho) - \beta \square c \quad (7)$$

where β is an additional fermionic ghost field. By assigning to ρ and β the following \mathcal{J} transformations

$$\mathcal{J}\rho = 0 , \quad \mathcal{J}\beta = \rho \quad (8)$$

one can check that the right hand side not only is \mathcal{J} -closed but also \mathcal{J} -exact like the left hand side, with Ω taking the form

$$\Omega = \beta (\partial^m A_m + \frac{\xi}{2} \rho) . \quad (9)$$

The fermionic ghost fields β and c correspond to the Fadeev-Popov ghosts and the bosonic ghost field ρ is the Nakanishi-Lautrup field. Moreover, the (β, ρ) pair defines the *non-minimal sector* of the theory, which must have two properties. First, it must allow the construction of an action for the fermionic ghost c (accomplished by β). Second, β can not be BRST isolated (\mathcal{J} -closed) because it may introduce new gauge invariant quantities and affect the physical sector of the theory (accomplished by ρ). Transformations (8) have a special structure which identifies the non-minimal sector ghosts β and ρ as a *BRST-doublet*. BRST-doublets play a very crucial role in BRST constructions because they do not contribute non-trivial pieces in the cohomology of the theory. In the case or

string theory or the (super)particle similar BRST-doublets are introduced.

In the BV formalism, one goes a step further and for every field / ghost introduce a “*conjugate*” variable, the antifield. For the moment, antifields are considered as classical sources that generate \mathcal{J} transformations of the corresponding fields or ghosts and thus appear in the action in the following manner: Antifield $\mathcal{J}(\text{Field})$. Therefore the antifield of a specific field / ghost carries appropriate quantum numbers / indices in order for these terms to be well defined additions to the deformed Lagrangian. For Maxwell theory we get

$$\mathcal{L}_{BV} = \frac{1}{2}A^m(\square A_m - \partial_m \partial^n A_n) + \rho(\partial^m A_m + \frac{\xi}{2}\rho) - \beta \square c + A^{*m} \partial_m c + \beta^* \rho \quad (10)$$

where $A^{*m}, c^*, \rho^*, \beta^*$ are the antifields of A_m, c, ρ, β respectively. In this case, c^* and ρ^* drop out of the Lagrangian because their corresponding ghost fields are \mathcal{J} -closed. Furthermore, in order for these additional terms to be BRST invariant:

$$\mathcal{J}[\text{Antifield } \mathcal{J}(\text{Field})] = \mathcal{J}(\text{Antifield}) \mathcal{J}(\text{Field}) = 0 \Rightarrow \mathcal{J}\text{Antifield} = 0 \quad (11)$$

we choose all antifields to be \mathcal{J} -closed.

For a general gauge theory, the variation of the BV action (S) under the \mathcal{J} -BRST transformations takes the form:

$$\delta S = \int \left\{ \mathcal{J}\Phi^i \frac{\delta S}{\delta \Phi^i} + \mathcal{J}\Phi_i^* \frac{\delta S}{\delta \Phi_i^*} \right\} \quad (12)$$

where Φ^i is symbolically the set of gauge fields and ghosts that participate in the BRST description of the gauge theory as discussed above and Φ_i^* are their corresponding antifields. However, because the antifields are \mathcal{J} -closed and also appear in the action such that $\frac{\delta S}{\delta \Phi_i^*} = \mathcal{J}\Phi^i$ one arrives to the expression:

$$\delta S = \int \frac{\delta S}{\delta \Phi_i^*} \frac{\delta S}{\delta \Phi^i} . \quad (13)$$

This motivates the definition of a binary bracket, called the antibracket

$$(F, G) = \int \left\{ \frac{\delta F}{\delta \Phi_i^*} \frac{\delta G}{\delta \Phi^i} + \frac{\delta G}{\delta \Phi_i^*} \frac{\delta F}{\delta \Phi^i} \right\} . \quad (14)$$

Using the antibracket the \mathcal{J} -invariance of the action S takes the simple form

$$(S, S) = 0 \quad (15)$$

which is known as the *classical master equation*. It is easy to check that the fields and antifields are conjugate variables with respect to the antibracket but most importantly because the action obeys the classical master equation one can define a nilpotent differential operator \mathfrak{s} ($\mathfrak{s}^2 = 0$) that acts in the space of fields and antifields

$$\mathfrak{s}F \equiv (S, F) \quad (16)$$

for any field or antifield F . The \mathfrak{s} transformations of the fields and antifields of Maxwell theory are:

$$\mathfrak{s}A_m = \partial_m c, \quad \mathfrak{s}A^{*m} = \square A^m - \partial^m \partial_n A^n - \partial^m \rho \quad (17a)$$

$$\mathbf{s}\rho = 0 , \quad \mathbf{s}\rho^* = \partial^m A_m + \rho + \beta^* \quad (17b)$$

$$\mathbf{s}\beta = \rho , \quad \mathbf{s}\beta^* = -\square c \quad (17c)$$

$$\mathbf{s}c = 0 , \quad \mathbf{s}c^* = \square\beta + \partial_m A^{*m} \quad (17d)$$

These transformations are nilpotent as expected and they have very interesting structure. The \mathbf{s} -transformation of any field matches exactly their \mathcal{J} transformation, whereas the \mathbf{s} -transformation of the antifields knows about the equations of motion of the corresponding fields.

In general, the differential operator \mathbf{s} can be decomposed into two operators: $\mathbf{s} = \gamma + \delta$. Operator γ is the newly renamed BRST symmetry transformation and is defined as follows:

$$\gamma\Phi^i \equiv \frac{\delta S}{\delta\Phi_i^*} = \mathcal{J}\Phi^i , \quad \gamma\Phi_i^* \equiv \frac{\delta}{\delta\Phi^i} \left[\Phi_j^* \mathcal{J}(\Phi^j) \right] . \quad (18)$$

For fields, γ coincides with the \mathcal{J} transformation but for antifields it has additional terms originating from the variation of the antifield terms in the action. The δ part is called the Koszul-Tate resolution⁸ and it implements the equations of motion. It is defined as follows:

$$\delta\Phi^i \equiv 0 , \quad \delta\Phi_i^* \equiv \frac{\delta}{\delta\Phi^i} \left[S - \Phi_j^* \mathcal{J}(\Phi^j) \right] . \quad (19)$$

Adding equations (18) and (19) automatically gives \mathbf{s} as defined in (16). Additionally, γ and δ are nilpotent and anticommute: $\gamma^2 = 0$, $\delta^2 = 0$, $\gamma\delta + \delta\gamma = 0$.

For Maxwell's theory the action of γ and δ operators on the fields and antifields is:

$$\gamma A_m = \partial_m c , \quad \gamma A^{*m} = 0 , \quad \delta A_m = 0 , \quad \delta A^{*m} = \square A^m - \partial^m \partial_n A^n - \partial^m \rho \quad (20a)$$

$$\gamma\rho = 0 , \quad \gamma\rho^* = \beta^* , \quad \delta\rho = 0 , \quad \delta\rho^* = \partial^m A_m + \rho \quad (20b)$$

$$\gamma\beta = \rho , \quad \gamma\beta^* = 0 , \quad \delta\beta = 0 , \quad \delta\beta^* = -\square c \quad (20c)$$

$$\gamma c = 0 , \quad \gamma c^* = \partial_m A^{*m} , \quad \delta c = 0 , \quad \delta c^* = \square\beta \quad (20d)$$

Having constructed the BRST operator \mathbf{s} on the space of fields and antifields, we can define a corresponding BRST charge operator Q which acts on an appropriately defined Hilbert space of states. For every pair of field Φ^i and antifield Φ_i^* we introduce a corresponding pair of state vectors $|e^{(\Phi^i)}\rangle$ and $|e^{(\Phi_i^*)}\rangle$ and we consider the vector space which is spanned by the states $\{|e^{(\Phi^i)}\rangle, |e^{(\Phi_i^*)}\rangle\}$. The most general state is written as

$$|\psi\rangle = \sum_i \chi_i |e^{(\Phi^i)}\rangle + \sum_i v^i |e^{(\Phi_i^*)}\rangle \quad (21)$$

where χ_i and v^i are field components of $|\psi\rangle$. Furthermore, for each of the basis vectors we assign a ghost number value which is defined to be opposite to the ghost number of the corresponding field or antifield. Specifically, if the field theoretic ghost number value of field Φ^i is $Gh(\Phi^i) = g^i$ then we

⁸See [43, 51]

define the vector space ghost number value of the corresponding vectors to be

$$gh(|e^{(\Phi^i)}\rangle) \equiv -g^i, \quad gh(|e^{(\Phi_i^*)}\rangle) \equiv 1 + g^i. \quad (22)$$

For Maxwell theory we have the following set of states:

$$\begin{aligned} gh\# = 0 &: \left\{ |e^{(A)}\rangle, |e^{(\rho)}\rangle, |e^{(\beta^*)}\rangle \right\} \\ gh\# = 1 &: \left\{ |e^{(A^*)}\rangle, |e^{(\rho^*)}\rangle, |e^{(\beta)}\rangle \right\} \\ gh\# = -1 &: \left\{ |e^{(c)}\rangle \right\} \\ gh\# = 2 &: \left\{ |e^{(c^*)}\rangle \right\} \end{aligned}$$

This structure of states can be captured by a Hilbert space with a vacuum state $|\omega\rangle$ and three fermionic creation operators η, ζ, π_ξ with ghost values $gh(\eta) = gh(\zeta) = -gh(\pi_\xi) = 1$ and $gh(|\omega\rangle) = 0$

$$\begin{aligned} gh\# = 0 &: \left\{ |\omega\rangle, \eta\pi_\xi|\omega\rangle, \zeta\pi_\xi|\omega\rangle \right\} \\ gh\# = 1 &: \left\{ \eta|\omega\rangle, \zeta|\omega\rangle, \eta\zeta\pi_\xi|\omega\rangle \right\} \\ gh\# = -1 &: \left\{ \pi_\xi|\omega\rangle \right\} \\ gh\# = 2 &: \left\{ \eta\zeta|\omega\rangle \right\} \end{aligned}$$

In this basis, the most general state is a linear combination of the above states

$$|\psi\rangle = |\psi^{(-1)}\rangle + |\psi^{(0)}\rangle + |\psi^{(1)}\rangle + |\psi^{(2)}\rangle \quad (23)$$

where

$$|\psi^{(-1)}\rangle = \pi_\xi|u\rangle, \quad |u\rangle = u|\omega\rangle \quad (24a)$$

$$|\psi^{(0)}\rangle = |w_1\rangle + \eta\pi_\xi|w_2\rangle + \zeta\pi_\xi|w_3\rangle, \quad |w_i\rangle = w_i|\omega\rangle \quad (24b)$$

$$|\psi^{(1)}\rangle = \eta|z_1\rangle + \zeta|z_2\rangle + \eta\zeta\pi_\xi|z_3\rangle, \quad |z_i\rangle = z_i|\omega\rangle \quad (24c)$$

$$|\psi^{(2)}\rangle = \eta\zeta|v\rangle, \quad |v\rangle = v|\omega\rangle \quad (24d)$$

and the coefficients u, w_i, z_i, v are elements of the field-antifield space. Therefore the BRST operator \mathbf{s} can act on the state $|\psi\rangle$ by acting on these coefficients. On the other hand, we can define a corresponding BRST operator Q that acts on the states of this Hilbert space which is equivalent to the action of \mathbf{s} (17)

$$Q|\psi\rangle = \mathbf{s}|\psi\rangle. \quad (25)$$

Q must be odd, have ghost value one and must be constructed out of the oscillators of the Hilbert space η, ζ, π_ξ and their conjugate π_η, π_ζ, ξ : $\{\eta, \pi_\eta\} = 1, \{\zeta, \pi_\zeta\} = 1, \{\xi, \pi_\xi\} = 1$. The most general ansatz for Q is

$$Q = \eta\Lambda^{(0)} + \zeta\Lambda^{(1)} + \xi\Lambda^{(-1)} \quad (26)$$

$$\begin{aligned}
& +\eta\zeta\pi_\eta B_1 + \eta\zeta\pi_\zeta B_2 + \eta\zeta\pi_\xi B_3 \\
& +\eta\xi\pi_\eta\Gamma_1 + \eta\xi\pi_\zeta\Gamma_2 + \eta\xi\pi_\xi\Gamma_3 \\
& +\zeta\xi\pi_\eta\Delta_1 + \zeta\xi\pi_\zeta\Delta_2 + \zeta\xi\pi_\xi\Delta_3 \\
& +\eta\zeta\xi\pi_\eta\pi_\zeta K_1 + \eta\zeta\xi\pi_\eta\pi_\xi K_2 + \eta\zeta\xi\pi_\zeta\pi_\xi K_3
\end{aligned}$$

Equation (25) can be expanded according to (23)

$$Q|\psi^{(-1)}\rangle + Q|\psi^{(0)}\rangle + Q|\psi^{(1)}\rangle + Q|\psi^{(2)}\rangle = \mathbf{s}|\psi^{(-1)}\rangle + \mathbf{s}|\psi^{(0)}\rangle + \mathbf{s}|\psi^{(1)}\rangle + \mathbf{s}|\psi^{(2)}\rangle \quad (27)$$

and by matching the Hilbert space ghost number (and the field-antifield space ghost number) of the two sides we get the following set of equations

$$Q|\psi^{(-1)}\rangle = \mathbf{s}|\psi^{(0)}\rangle, \quad (28a)$$

$$Q|\psi^{(0)}\rangle = \mathbf{s}|\psi^{(1)}\rangle, \quad (28b)$$

$$Q|\psi^{(1)}\rangle = \mathbf{s}|\psi^{(2)}\rangle, \quad (28c)$$

$$Q|\psi^{(2)}\rangle = 0 = \mathbf{s}|\psi^{(-1)}\rangle. \quad (28d)$$

These equations can be solved to fix all the coefficients of $|\psi\rangle$ and determine all the field operators in Q . We find:

$$\begin{aligned}
u = c, \quad |u\rangle = |c\rangle = c|\omega\rangle, \quad v = c^*, \quad |v\rangle = |c^*\rangle = c^*|\omega\rangle, \\
w_1 = A^m a_m^\dagger, \quad |w_1\rangle = |A\rangle = A^m a_m^\dagger |\omega\rangle, \quad z_1 = -A^{*m} a_m^\dagger - \partial^m a_m^\dagger \rho^*, \quad |z_1\rangle = -|A^*\rangle - \Lambda^{(-1)}|\rho^*\rangle, \\
w_2 = -\beta^*, \quad |w_2\rangle = -|\beta^*\rangle = -\beta^*|\omega\rangle, \quad z_2 = -\rho^* + (1 + \partial^m a_m^\dagger)\beta, \quad |z_2\rangle = -|\rho^*\rangle + (1 + \Lambda^{(-1)})|\beta\rangle, \\
w_3 = \rho, \quad |w_3\rangle = |\rho\rangle = \rho|\omega\rangle, \quad z_3 = -\square\beta, \quad |z_3\rangle = -\square|\beta\rangle
\end{aligned}$$

where a_m^\dagger and a_m is a pair of bosonic creation and annihilation operators ($[a_m, a_n^\dagger] = \eta_{mn}$) and

$$Q = \eta \square + \zeta \partial^m a_m + \xi \partial^m a_m^\dagger - \zeta \xi \pi_\eta. \quad (29)$$

This Q operator is nilpotent by construction and it is identical to the BFV-BRST charge constructed from the algebra of operators $\Lambda^{(0)} = \square$, $\Lambda^{(1)} = \partial^m a_m$, $\Lambda^{(-1)} = \partial^m a_m^\dagger$:

$$[\Lambda^{(0)}, \Lambda^{(1)}] = 0, \quad [\Lambda^{(0)}, \Lambda^{(-1)}] = 0, \quad [\Lambda^{(1)}, \Lambda^{(-1)}] = \Lambda^{(0)}. \quad (30)$$

The general physical state (zero ghost number) cohomology of Q is given by:

$$|\phi\rangle = |A\rangle + \eta\pi_\xi|B\rangle + \zeta\pi_\xi|C\rangle \quad (31)$$

with a transformation:

$$\delta|\phi\rangle = Q|L\rangle, \quad |L\rangle = \pi_\xi|\lambda\rangle \Rightarrow \begin{cases} \delta|A\rangle = \Lambda^{(-1)}|\lambda\rangle, \\ \delta|B\rangle = \Lambda^{(0)}|\lambda\rangle, \\ \delta|C\rangle = \Lambda^{(1)}|\lambda\rangle \end{cases} \quad (32)$$

and equations of motion:

$$Q|\phi\rangle = 0 \Rightarrow \begin{cases} \Lambda^{(0)}|A\rangle - \Lambda^{(-1)}\Lambda^{(1)}|A\rangle + \Lambda^{(-1)}\Lambda^{(-1)}|C\rangle = 0, \\ \Lambda^{(1)}|A\rangle - \Lambda^{(-1)}|D\rangle - |B\rangle = 0, \\ \Lambda^{(0)}|D\rangle - \Lambda^{(1)}\Lambda^{(1)}|A\rangle + \Lambda^{(1)}\Lambda^{(-1)}|D\rangle = 0. \end{cases} \quad (33)$$

For the case $|\phi\rangle = |\psi^{(0)}\rangle$ and $|L\rangle = |\psi^{(-1)}\rangle$ one recovers the Maxwell theory equations and transformations.

3 Superspace BRST description of Super Maxwell theory

In this section, we apply the above concepts in superspace aiming towards a BRST-BV/BFV description of the vector supermultiplet which is the supersymmetric extension of Maxwell theory. The superspace description of the theory is given in terms of a real scalar superfield $V(x, \theta, \bar{\theta})$ with the following superspace action principle

$$S = \frac{1}{2} \int d^4x d^2\theta d^2\bar{\theta} V D^\gamma \bar{D}^2 D_\gamma V \quad (34)$$

which is invariant under the gauge transformation $\delta V = \bar{D}^2 L + D^2 \bar{L}$. The operators D_α and $\bar{D}_{\dot{\alpha}}$ are the supersymmetric covariant derivatives and the off-shell spectrum of the theory is a Maxwell spin 1 gauge field, a decoupled auxiliary real scalar field and a spin 1/2 fermion.

Similarly to the discussion in the previous section, we promote the gauge symmetry of the theory to a nilpotent, superspace BRST symmetry \mathfrak{J} . The action of \mathfrak{J} on the gauge superfield V is found by fermionizing the original gauge transformation and replace the gauge parameter superfields with ghost superfields:

$$\mathfrak{J}V = \bar{D}^2 \mathfrak{c} + D^2 \bar{\mathfrak{c}}. \quad (35)$$

Using the nilpotence of \mathfrak{J} we also find that

$$0 = \mathfrak{J}^2 V = \bar{D}^2 (\mathfrak{J}\mathfrak{c}) + D^2 (\mathfrak{J}\bar{\mathfrak{c}}) \Rightarrow \mathfrak{J}\mathfrak{c} = 0, \quad \mathfrak{J}\bar{\mathfrak{c}} = 0 \quad (36)$$

An important comment is that, unlike Maxwell theory, the differential operators that appears in the gauge transformation of the gauge superfield have a non-zero kernel and based on the algebra of the supersymmetric covariant derivatives we can introduce ghosts for ghosts, ghosts for ghosts for ghosts et cetera ad infinitum⁹. For the purpose of this paper these contributions are not required and we will not consider them.

⁹ $\mathfrak{J}\mathfrak{c} = 0 + \bar{D}^{\dot{\alpha}} \bar{\mathfrak{d}}_{\dot{\alpha}}, \quad \mathfrak{J}\bar{\mathfrak{c}} = 0 + \bar{D}^{\dot{\beta}} \bar{\mathfrak{d}}_{\dot{\alpha}\dot{\beta}}, \quad \mathfrak{J}\bar{\mathfrak{d}}_{\dot{\alpha}\dot{\beta}} = 0 + \bar{D}^{\dot{\gamma}} \bar{\mathfrak{d}}_{\dot{\alpha}\dot{\beta}\dot{\gamma}}, \dots$. Similarly for $\bar{\mathfrak{c}}$.

Next, we deform the Lagrangian by adding an appropriate \mathfrak{J} -exact term, $\mathfrak{J}\Omega$. Motivated from the quantization procedure for this theory, this deformation will include a gauge fixing condition accompanied by the corresponding Lagrange multiplier ghost superfield. The usual superspace gauge fixing conditions for gauge superfield V are $D^2V = 0$ and $\bar{D}^2V = 0$, which correspond to the superspace extension of the Landau-Fermi gauge in Maxwell theory. Therefore, we consider the deformation:

$$\mathfrak{J}\Omega = \rho(D^2V + \frac{\xi}{2}\bar{\rho}) + \bar{\rho}(\bar{D}^2V + \frac{\xi}{2}\rho) + \dots \quad (37)$$

where ρ and $\bar{\rho}$ are Nakanishi-Lautrup superfield ghosts, ξ is the Feynman gauge parameter and the dots represent additional terms that we have in order to make the right hand side of the equation \mathfrak{J} -exact. Using the nilpotence of \mathfrak{J} we find the consistence condition

$$0 = (\mathfrak{J}\rho)(D^2V + \xi\bar{\rho}) + (\mathfrak{J}\bar{\rho})(\bar{D}^2V + \xi\rho) + \rho D^2\bar{D}^2c + \bar{\rho}\bar{D}^2D^2\bar{c} + \mathfrak{J}(\dots) \quad (38)$$

which determines the missing terms to be

$$\mathfrak{J}\Omega = \rho(D^2V + \frac{\xi}{2}\bar{\rho}) + \bar{\rho}(\bar{D}^2V + \frac{\xi}{2}\rho) - \beta D^2\bar{D}^2c - \bar{\beta}\bar{D}^2D^2\bar{c}. \quad (39)$$

The remaining \mathfrak{J} transformations are

$$\mathfrak{J}\beta = \rho, \quad \mathfrak{J}\rho = 0, \quad \mathfrak{J}\bar{\beta} = \bar{\rho}, \quad \mathfrak{J}\bar{\rho} = 0 \quad (40)$$

and β is a fermionic ghost superfield which forms a BRST-doublet with ρ . Similarly for $\bar{\beta}$ and $\bar{\rho}$. This is expected because there are two gauge parameter superfields (L, \bar{L}) and therefore by fermionizing them we get two fermionic ghost superfields (c, \bar{c}) and for each one of them we must introduce a BRST-doublet, as discussed in the previous section. The non-minimal sector of this theory is the two BRST-doublets (β, ρ) and $(\bar{\beta}, \bar{\rho})$. Also, using the above \mathfrak{J} transformations one can check that the right hand side of (39) is \mathfrak{J} -exact and solve for Ω

$$\Omega = \beta(D^2V + \frac{\xi}{2}\bar{\rho}) + \bar{\beta}(\bar{D}^2V + \frac{\xi}{2}\rho). \quad (41)$$

This deformation is further extended by the BV anti-superfields. For each (ghost) superfield we introduce an anti-superfield with opposite statistics and appropriate superspace ghost number and add to the action a [Anti-superfield \mathfrak{J} (Superfield)] term. The superspace BV Lagrangian takes the form:

$$\begin{aligned} \mathcal{L}_{BV} = & \frac{1}{2}VD^\gamma\bar{D}^2D_\gamma V + \rho(D^2V + \frac{\xi}{2}\bar{\rho}) + \bar{\rho}(\bar{D}^2V + \frac{\xi}{2}\rho) - \beta D^2\bar{D}^2c - \bar{\beta}\bar{D}^2D^2\bar{c} \\ & + V^*(\bar{D}^2c + D^2\bar{c}) + \beta^*\rho + \bar{\beta}^*\bar{\rho} \end{aligned}$$

where $V^*, \beta^*, \bar{\beta}^*, \rho^*, \bar{\rho}^*, c^*, \bar{c}^*$ are the anti-superfields corresponding to $V, \beta, \bar{\beta}, \rho, \bar{\rho}, c, \bar{c}$ respectively. The ghosts $\rho, \bar{\rho}, c, \bar{c}$ are \mathfrak{J} -closed, thus their conjugate anti-superfields drop out of the BV superspace action. In order to maintain the \mathfrak{J} BRST symmetry of the BV action we assign

to all the anti-superfields trivial \mathfrak{J} transformations (\mathfrak{J} -closed):

$$\mathfrak{J}V^* = \mathfrak{J}\beta^* = \mathfrak{J}\bar{\beta}^* = \mathfrak{J}\rho^* = \mathfrak{J}\bar{\rho}^* = \mathfrak{J}\mathbf{c}^* = \mathfrak{J}\bar{\mathbf{c}}^* = 0 . \quad (42)$$

Now we can define a full superspace antibracket (\cdot, \cdot)

$$\begin{aligned} (\mathbf{F}, \mathbf{G}) &\equiv \int d^4x d^2\theta d^2\bar{\theta} \left\{ \frac{\delta\mathbf{F}}{\delta\Phi_i^*} \frac{\delta\mathbf{G}}{\delta\Phi^i} + \frac{\delta\mathbf{G}}{\delta\Phi_i^*} \frac{\delta\mathbf{F}}{\delta\Phi^i} \right\} \\ &= \int d^4x d^2\theta d^2\bar{\theta} \left\{ \frac{\delta\mathbf{F}}{\delta V^*} \frac{\delta\mathbf{G}}{\delta V} + \frac{\delta\mathbf{G}}{\delta V^*} \frac{\delta\mathbf{F}}{\delta V} \right. \\ &\quad + \frac{\delta\mathbf{F}}{\delta\beta^*} \frac{\delta\mathbf{G}}{\delta\beta} + \frac{\delta\mathbf{G}}{\delta\beta^*} \frac{\delta\mathbf{F}}{\delta\beta} + \frac{\delta\mathbf{F}}{\delta\bar{\beta}^*} \frac{\delta\mathbf{G}}{\delta\bar{\beta}} + \frac{\delta\mathbf{G}}{\delta\bar{\beta}^*} \frac{\delta\mathbf{F}}{\delta\bar{\beta}} \\ &\quad + \frac{\delta\mathbf{F}}{\delta\rho^*} \frac{\delta\mathbf{G}}{\delta\rho} + \frac{\delta\mathbf{G}}{\delta\rho^*} \frac{\delta\mathbf{F}}{\delta\rho} + \frac{\delta\mathbf{F}}{\delta\bar{\rho}^*} \frac{\delta\mathbf{G}}{\delta\bar{\rho}} + \frac{\delta\mathbf{G}}{\delta\bar{\rho}^*} \frac{\delta\mathbf{F}}{\delta\bar{\rho}} \\ &\quad \left. + \frac{\delta\mathbf{F}}{\delta\mathbf{c}^*} \frac{\delta\mathbf{G}}{\delta\mathbf{c}} + \frac{\delta\mathbf{G}}{\delta\mathbf{c}^*} \frac{\delta\mathbf{F}}{\delta\mathbf{c}} + \frac{\delta\mathbf{F}}{\delta\bar{\mathbf{c}}^*} \frac{\delta\mathbf{G}}{\delta\bar{\mathbf{c}}} + \frac{\delta\mathbf{G}}{\delta\bar{\mathbf{c}}^*} \frac{\delta\mathbf{F}}{\delta\bar{\mathbf{c}}} \right\} \end{aligned} \quad (43)$$

Using this bracket, the \mathfrak{J} invariance of the BV superspace action takes the form of a superspace classical master equation

$$(S, S) = 2\mathfrak{J}S = 0 . \quad (44)$$

As discussed previously, by populating the first slot of the antibracket with the superspace BV action S , we define a nilpotent superspace BRST differential operator \mathfrak{s} in the space of superfields and anti-superfields

$$\mathfrak{s}\mathbf{F} \equiv (S, \mathbf{F}) . \quad (45)$$

The action of \mathfrak{s} on the superfields and anti-superfields of the vector supermultiplet is:

$$\mathfrak{s}V = \bar{D}^2\mathbf{c} + D^2\bar{\mathbf{c}} , \quad \mathfrak{s}V^* = D^\gamma\bar{D}^2D_\gamma V + D^2\rho + \bar{D}^2\bar{\rho} , \quad (46a)$$

$$\mathfrak{s}\rho = 0 , \quad \mathfrak{s}\rho^* = \beta^* + D^2V + \xi\bar{\rho} , \quad (46b)$$

$$\mathfrak{s}\bar{\rho} = 0 , \quad \mathfrak{s}\bar{\rho}^* = \bar{\beta}^* + \bar{D}^2V + \xi\rho , \quad (46c)$$

$$\mathfrak{s}\beta = \rho , \quad \mathfrak{s}\beta^* = -D^2\bar{D}^2\mathbf{c} , \quad (46d)$$

$$\mathfrak{s}\bar{\beta} = \bar{\rho} , \quad \mathfrak{s}\bar{\beta}^* = -\bar{D}^2D^2\bar{\mathbf{c}} , \quad (46e)$$

$$\mathfrak{s}\mathbf{c} = 0 , \quad \mathfrak{s}\mathbf{c}^* = -\bar{D}^2V^* + \bar{D}^2D^2\beta , \quad (46f)$$

$$\mathfrak{s}\bar{\mathbf{c}} = 0 , \quad \mathfrak{s}\bar{\mathbf{c}}^* = -D^2V^* + D^2\bar{D}^2\bar{\beta} . \quad (46g)$$

As expected the action of \mathfrak{s} on superfields is identical to their \mathfrak{J} transformations, while \mathfrak{s} on anti-superfields gives the equations of motion for the corresponding superfields. By splitting these equations of motion into two pieces, the piece coming from the anti-superfield terms of the action and the rest we can decompose \mathfrak{s} into two nilpotent and anticommuting operators γ and δ : $\mathfrak{s} = \gamma + \delta$. The γ part is the superspace BRST symmetry operator and δ is the superspace Koszul-Tate resolution differential:

$$\gamma V = \bar{D}^2\mathbf{c} + D^2\bar{\mathbf{c}} , \quad \gamma V^* = 0 , \quad \delta V = 0 , \quad \delta V^* = D^\gamma\bar{D}^2D_\gamma V + D^2\rho + \bar{D}^2\bar{\rho} , \quad (47a)$$

$$\gamma\rho = 0, \quad \gamma\rho^* = \beta^*, \quad \delta\rho = 0, \quad \delta\rho^* = D^2V + \xi\bar{\rho}, \quad (47b)$$

$$\gamma\bar{\rho} = 0, \quad \gamma\bar{\rho}^* = \bar{\beta}^*, \quad \delta\bar{\rho} = 0, \quad \delta\bar{\rho}^* = \bar{D}^2V + \xi\rho, \quad (47c)$$

$$\gamma\beta = \rho, \quad \gamma\beta^* = 0, \quad \delta\beta = 0, \quad \delta\beta^* = -D^2\bar{D}^2c, \quad (47d)$$

$$\gamma\bar{\beta} = \bar{\rho}, \quad \gamma\bar{\beta}^* = 0, \quad \delta\bar{\beta} = 0, \quad \delta\bar{\beta}^* = -\bar{D}^2D^2\bar{c}, \quad (47e)$$

$$\gamma c = 0, \quad \gamma c^* = -\bar{D}^2V^*, \quad \delta c = 0, \quad \delta c^* = \bar{D}^2D^2\beta, \quad (47f)$$

$$\gamma\bar{c} = 0, \quad \gamma\bar{c}^* = -D^2V^*, \quad \delta\bar{c} = 0, \quad \delta\bar{c}^* = D^2\bar{D}^2\bar{\beta}. \quad (47g)$$

Using the above it is straightforward to verify that $\gamma^2 = 0$, $\delta^2 = 0$, $\{\gamma, \delta\} = 0$.

The last step, is to use the \mathfrak{S} transformations in order to define a nilpotent BRST charge operator in a Hilbert space of states. In the previous section, we demonstrated explicitly how this can be done for the Maxwell theory. The result of that procedure can be summarized in four steps. First, we take as an input the list of operators that appear in the nilpotent differential operator defined by the antibracket. Second, if some of these operators carry free indices then we introduce pairs of creation and annihilation oscillators with appropriate statistics to dress them and absorb these free indices. Third, we calculate the algebra of all these dressed operators and apply the Fradkin and Fradkina algorithm [44] for the construction of a corresponding Hilbert space, nilpotent BRST charge operator. Finally, we select an appropriate vacuum state in order to define the cohomology of the BRST charge in this Hilbert space.

In this case, according to equations (46) we must consider the following list of superspace operators

$$\left\{ D^\gamma\bar{D}^2D_\gamma, D^2, \bar{D}^2, D^2\bar{D}^2, \bar{D}^2D^2 \right\} \quad (48)$$

None of these operators carry free indices, therefore we do not require the introduction of any additional oscillators besides the ghost oscillators of the Fradkin-Fradkina process. The algebra of these operators is as follows:

$$[D^\gamma\bar{D}^2D_\gamma, D^2] = 0, \quad [D^\gamma\bar{D}^2D_\gamma, \bar{D}^2] = 0, \quad [D^\gamma\bar{D}^2D_\gamma, D^2\bar{D}^2] = 0, \quad [D^\gamma\bar{D}^2D_\gamma, \bar{D}^2D^2] = 0, \quad (49a)$$

$$[D^2, \bar{D}^2] = D^2\bar{D}^2 - \bar{D}^2D^2, \quad [D^2, D^2\bar{D}^2] = -\square D^2, \quad [D^2, \bar{D}^2D^2] = \square D^2, \quad (49b)$$

$$[\bar{D}^2, D^2\bar{D}^2] = \square \bar{D}^2, \quad [\bar{D}^2, \bar{D}^2D^2] = -\square \bar{D}^2, \quad (49c)$$

$$[D^2\bar{D}^2, \bar{D}^2D^2] = 0. \quad (49d)$$

Notice that the d'Alembertian operator emerges in some of the commutators. In superspace, this is not an independent operator because it can be expressed as a linear combination of other operators ¹⁰:

$$\square = D^2\bar{D}^2 + \bar{D}^2D^2 - D^\gamma\bar{D}^2D_\gamma. \quad (50)$$

This makes the above algebra non-linear. The construction of BRST operators of non-linear algebras has been found in [49, 50]. Using these results, we find that the most general BRST charge we can write is:

$$\mathbf{Q} = \eta D^\gamma\bar{D}^2D_\gamma + \bar{\zeta} D^2 + \zeta \bar{D}^2 + \xi D^2\bar{D}^2 + \bar{\xi} \bar{D}^2D^2 + \kappa \square - \bar{\zeta}\zeta\pi_\xi + \bar{\zeta}\zeta\pi_{\bar{\xi}} \quad (51)$$

¹⁰We follow the conventions of *Superspace*

where $\kappa = -\xi\bar{\zeta}\pi_{\bar{\zeta}} - \bar{\zeta}\bar{\xi}\pi_{\bar{\zeta}} - \zeta\xi\pi_{\zeta} - \bar{\xi}\zeta\pi_{\zeta}$ and $\eta, \zeta, \bar{\zeta}, \xi, \bar{\xi}$ are fermionic ghost oscillators with conjugate oscillators $\pi_{\eta}, \pi_{\zeta}, \pi_{\bar{\zeta}}, \pi_{\xi}, \pi_{\bar{\xi}}$. The $D^2\bar{D}^2$ and \bar{D}^2D^2 terms in the above expression can be decomposed into an anticommutator and a commutator

$$D^2\bar{D}^2 = \frac{1}{2}\{D^2, \bar{D}^2\} + \frac{1}{2}[D^2, \bar{D}^2], \quad \bar{D}^2D^2 = \frac{1}{2}\{D^2, \bar{D}^2\} - \frac{1}{2}[D^2, \bar{D}^2] \quad (52)$$

and the BRST charge takes the form

$$\mathbf{Q} = \eta D^\gamma \bar{D}^2 D_\gamma + \bar{\zeta} D^2 + \zeta \bar{D}^2 + \rho [D^2, \bar{D}^2] + 2(\bar{\zeta}\rho\pi_{\bar{\zeta}} - \zeta\rho\pi_{\zeta}) \square - \bar{\zeta}\zeta\pi_{\rho} + \beta \{D^2, \bar{D}^2\} \quad (53)$$

where $\beta = \frac{1}{2}(\xi + \bar{\xi})$, $\rho = \frac{1}{2}(\xi - \bar{\xi})$ and $\pi_{\rho} = \pi_{\xi} - \pi_{\bar{\xi}}$ is ρ 's conjugate oscillator $\{\rho, \pi_{\rho}\} = 1$. Because of (50), the $\{D^2, \bar{D}^2\}$ term is not an independent generator and its effect is already captured by the $D^\gamma \bar{D}^2 D_\gamma$ and \square terms. Therefore, we can ignore this term (choose $\beta = 0$) and focus on the cohomology of the following BRST charge

$$\mathbf{Q} = \eta D^\gamma \bar{D}^2 D_\gamma + \bar{\zeta} D^2 + \zeta \bar{D}^2 + \rho [D^2, \bar{D}^2] + 2(\bar{\zeta}\rho\pi_{\bar{\zeta}} - \zeta\rho\pi_{\zeta}) \square - \bar{\zeta}\zeta\pi_{\rho} \quad (54)$$

acting in the reduced Hilbert space generated out of the fermionic oscillators $\eta, \zeta, \bar{\zeta}, \rho$ and their conjugates. In this Hilbert space, we select the vacuum $|0\rangle$ such that it is annihilated by the oscillators $\pi_{\eta}, \bar{\zeta}, \zeta, \rho$

$$\pi_{\eta}|0\rangle = \bar{\zeta}|0\rangle = \zeta|0\rangle = \rho|0\rangle = 0. \quad (55)$$

Thus the most general state of this Hilbert space takes the form

$$|\Phi\rangle = \sum_{a,b,c,d} \eta^a \pi_{\rho}^b \pi_{\bar{\zeta}}^c \pi_{\zeta}^d |\Phi_{a,b,c,d}\rangle \quad (56)$$

where $a, b, c, d = 0, 1$, $|\Phi_{a,b,c,d}\rangle = \Phi_{a,b,c,d}|0\rangle$ and $\Phi_{a,b,c,d}$ is a superfield coefficient. The Hilbert space ghost number of this state is:

$$gh(|\Phi\rangle) = a - b - c - d. \quad (57)$$

The zero ghost state (physical state) is

$$|\Psi\rangle = |\mathbf{V}\rangle + \eta\pi_{\rho}|\mathbf{A}\rangle + \eta\pi_{\bar{\zeta}}|\mathbf{B}\rangle + \eta\pi_{\zeta}|\bar{\mathbf{B}}\rangle \quad (58)$$

and we can also construct three gauge parameter states with ghost values -1 ($|\mathbf{\Lambda}_{-1}\rangle$), -2 ($|\mathbf{\Lambda}_{-2}\rangle$) and -3 ($|\mathbf{\Lambda}_{-3}\rangle$):

$$|\mathbf{\Lambda}_{-1}\rangle = \pi_{\rho}|\boldsymbol{\lambda}\rangle + \pi_{\bar{\zeta}}|\mathbf{L}\rangle + \pi_{\zeta}|\bar{\mathbf{L}}\rangle + \eta\pi_{\rho}\pi_{\bar{\zeta}}|\boldsymbol{\omega}_1\rangle + \eta\pi_{\rho}\pi_{\zeta}|\boldsymbol{\omega}_2\rangle + \eta\pi_{\bar{\zeta}}\pi_{\zeta}|\boldsymbol{\omega}_3\rangle, \quad (59a)$$

$$|\mathbf{\Lambda}_{-2}\rangle = \pi_{\rho}\pi_{\bar{\zeta}}|\boldsymbol{w}_1\rangle + \pi_{\rho}\pi_{\zeta}|\boldsymbol{w}_2\rangle + \pi_{\bar{\zeta}}\pi_{\zeta}|\boldsymbol{w}_3\rangle + \eta\pi_{\rho}\pi_{\bar{\zeta}}\pi_{\zeta}|\boldsymbol{w}_4\rangle, \quad (59b)$$

$$|\mathbf{\Lambda}_{-3}\rangle = \pi_{\rho}\pi_{\bar{\zeta}}\pi_{\zeta}|\boldsymbol{z}\rangle. \quad (59c)$$

The transformation laws of the superfield coefficients at each ghost level are derived by the action

of the BRST charge (54) on the previous level state. Hence we get:

$$\delta_Q|\Lambda_{-2}\rangle = Q|\Lambda_{-3}\rangle \Rightarrow \begin{cases} \delta|\mathbf{w}_1\rangle = \bar{D}^2|z\rangle, \\ \delta|\mathbf{w}_2\rangle = -D^2|z\rangle, \\ \delta|\mathbf{w}_3\rangle = [D^2, \bar{D}^2]|z\rangle, \\ \delta|\mathbf{w}_4\rangle = D^\gamma \bar{D}^2 D_\gamma |z\rangle \end{cases} \quad (60)$$

$$\delta_Q|\Lambda_{-1}\rangle = Q|\Lambda_{-2}\rangle \Rightarrow \begin{cases} \delta|\boldsymbol{\lambda}\rangle = -D^2|\mathbf{w}_1\rangle - \bar{D}^2|\mathbf{w}_2\rangle + |\mathbf{w}_3\rangle, \\ \delta|\mathbf{L}\rangle = -\bar{D}^2|\mathbf{w}_3\rangle + [D^2, \bar{D}^2]|\mathbf{w}_1\rangle + 2\Box|\mathbf{w}_1\rangle, \\ \delta|\bar{\mathbf{L}}\rangle = D^2|\mathbf{w}_3\rangle + [D^2, \bar{D}^2]|\mathbf{w}_2\rangle - 2\Box|\mathbf{w}_2\rangle, \\ \delta|\boldsymbol{\omega}_1\rangle = D^\gamma \bar{D}^2 D_\gamma |\mathbf{w}_1\rangle - \bar{D}^2|\mathbf{w}_4\rangle, \\ \delta|\boldsymbol{\omega}_2\rangle = D^\gamma \bar{D}^2 D_\gamma |\mathbf{w}_2\rangle + D^2|\mathbf{w}_4\rangle, \\ \delta|\boldsymbol{\omega}_3\rangle = D^\gamma \bar{D}^2 D_\gamma |\mathbf{w}_3\rangle - [D^2, \bar{D}^2]|\mathbf{w}_4\rangle \end{cases} \quad (61)$$

$$\delta_Q|\Psi\rangle = Q|\Lambda_{-1}\rangle \Rightarrow \begin{cases} \delta|\mathbf{V}\rangle = D^2|\mathbf{L}\rangle + \bar{D}^2|\bar{\mathbf{L}}\rangle + [D^2, \bar{D}^2]|\boldsymbol{\lambda}\rangle, \\ \delta|\mathbf{A}\rangle = D^\gamma \bar{D}^2 D_\gamma |\boldsymbol{\lambda}\rangle + D^2|\boldsymbol{\omega}_1\rangle + \bar{D}^2|\boldsymbol{\omega}_2\rangle - |\boldsymbol{\omega}_3\rangle, \\ \delta|\mathbf{B}\rangle = D^\gamma \bar{D}^2 D_\gamma |\mathbf{L}\rangle + \bar{D}^2|\boldsymbol{\omega}_3\rangle - [D^2, \bar{D}^2]|\boldsymbol{\omega}_1\rangle - 2\Box|\boldsymbol{\omega}_1\rangle, \\ \delta|\bar{\mathbf{B}}\rangle = D^\gamma \bar{D}^2 D_\gamma |\bar{\mathbf{L}}\rangle - D^2|\boldsymbol{\omega}_3\rangle - [D^2, \bar{D}^2]|\boldsymbol{\omega}_2\rangle + 2\Box|\boldsymbol{\omega}_2\rangle \end{cases} \quad (62)$$

The equation of motion for the physical state is:

$$Q|\Psi\rangle = 0 \Rightarrow D^\gamma \bar{D}^2 D_\gamma |\mathbf{V}\rangle - D^2|\mathbf{B}\rangle - \bar{D}^2|\bar{\mathbf{B}}\rangle - [D^2, \bar{D}^2]|\mathbf{A}\rangle = 0. \quad (63)$$

Using the gauge freedom, the states $|\mathbf{A}\rangle$ and $|\boldsymbol{\lambda}\rangle$ can be eliminate and we get

$$D^\gamma \bar{D}^2 D_\gamma |\mathbf{V}\rangle = |\Phi\rangle + |\bar{\Phi}\rangle \quad (64)$$

where $|\Phi\rangle$ is the chiral state $|\Phi\rangle = \bar{D}^2|\bar{\mathbf{B}}\rangle$, $\bar{D}_\alpha|\Phi\rangle = 0$ and $|\bar{\Phi}\rangle$ is the antichiral state $|\bar{\Phi}\rangle = D^2|\mathbf{B}\rangle$, $D_\alpha|\bar{\Phi}\rangle = 0$. The transformation of the states $|\mathbf{V}\rangle$, $|\Phi\rangle$, $|\bar{\Phi}\rangle$ are

$$\delta|\mathbf{V}\rangle = D^2|\bar{\mathbf{L}}\rangle + \bar{D}^2|\mathbf{L}\rangle, \quad \delta|\Phi\rangle = 0, \quad \delta|\bar{\Phi}\rangle = 0 \quad (65)$$

and as an integrability condition to (64), the chiral and antichiral states satisfy their expected equations of motion

$$D^2|\Phi\rangle = 0, \quad \bar{D}^2|\bar{\Phi}\rangle = 0. \quad (66)$$

This includes a consistent sector of the theory where the chiral and antichiral state vanish ($|\Phi\rangle = 0 = |\bar{\Phi}\rangle$). This sector correspond to the free, massless vector supermultiplet given by the state $|\mathbf{V}\rangle$ which satisfies the equation of motion

$$D^\gamma \bar{D}^2 D_\gamma |\mathbf{V}\rangle = 0, \quad \delta|\mathbf{V}\rangle = D^2|\bar{\mathbf{L}}\rangle + \bar{D}^2|\mathbf{L}\rangle. \quad (67)$$

as expected from the superspace action (34).

4 Superspace BRST description of Super Yang Mills

The full $4D$, $\mathcal{N} = 1$ SYM theory, which is the non-abelian extension of the vector multiplet, is known to be described by the superspace action

$$S = -\frac{1}{4} Tr \int d^4x d^4\theta (e^{-V} D^\gamma e^V) \bar{D}^2 (e^{-V} D_\gamma e^V) + h.c. \quad (68)$$

where $V = V^I T_I$ and T_I are the generators of an internal symmetry group. The action is invariant under the gauge transformation

$$\begin{aligned} e^{V'} = e^{i\bar{\Lambda}} e^V e^{-i\Lambda} &\Rightarrow \delta V = -\frac{i}{2} L_V [\Lambda + \bar{\Lambda} + \coth(\frac{1}{2} L_V) (\Lambda - \bar{\Lambda})] \\ &= D^2 \bar{L} + \bar{D}^2 L - \frac{1}{2} [V, D^2 \bar{L} - \bar{D}^2 L] + \mathcal{O}(V^2) \end{aligned} \quad (69)$$

where $i\bar{\Lambda} = D^2 \bar{L}$, $i\Lambda = -\bar{D}^2 L$ and $L = L^I T_I$ and $\bar{L} = \bar{L}^I T_I$.

The procedure discussed in the previous section can also be applied to this theory. The superspace \mathfrak{J} BRST transformation is obtained by fermionizing the above gauge transformation

$$\mathfrak{J}V = D^2 \bar{c} + \bar{D}^2 c - \frac{1}{2} [V, D^2 \bar{c} - \bar{D}^2 c] + \dots \quad (70)$$

where $c = c^I T_I$ and $\bar{c} = \bar{c}^I T_I$. The nilpotence of \mathfrak{J} on V fixes the action of \mathfrak{J} on the ghosts c and \bar{c} :

$$\mathfrak{J}^2 V = 0 \Rightarrow \mathfrak{J}c = -\frac{1}{2} \{c, \bar{D}^2 c\}, \quad \mathfrak{J}\bar{c} = \frac{1}{2} \{\bar{c}, D^2 \bar{c}\} \quad (71)$$

Using these transformations, the \mathfrak{J} -exact deformation of the SYM Lagrangian is

$$\begin{aligned} \mathfrak{J}\Omega = Tr \left\{ \rho (D^2 V + \frac{\xi}{2} \bar{\rho}) + \bar{\rho} (\bar{D}^2 V + \frac{\xi}{2} \rho) \right. \\ \left. -\beta D^2 \left\{ -\frac{i}{2} L_V [i \bar{D}^2 c - i D^2 \bar{c} + \coth(\frac{1}{2} L_V) (i \bar{D}^2 c + i D^2 \bar{c})] \right\} \right. \\ \left. -\bar{\beta} \bar{D}^2 \left\{ -\frac{i}{2} L_V [i \bar{D}^2 c - i D^2 \bar{c} + \coth(\frac{1}{2} L_V) (i \bar{D}^2 c + i D^2 \bar{c})] \right\} \right\} \end{aligned} \quad (72)$$

where (ρ, β) and $(\bar{\rho}, \bar{\beta})$ are the two algebra valued BRST doublets: $\mathfrak{J}\beta = \rho$, $\mathfrak{J}\rho = 0$, $\mathfrak{J}\bar{\beta} = \bar{\rho}$, $\mathfrak{J}\bar{\rho} = 0$.

By adding the corresponding anti-superfields, the superspace BV Lagrangian for SYM takes the form

$$\begin{aligned} \mathcal{L}_{BV} = Tr \left\{ -\frac{1}{4} (e^{-V} D^\gamma e^V) \bar{D}^2 (e^{-V} D_\gamma e^V) - \frac{1}{4} (e^{-V} \bar{D}^\dot{\gamma} e^V) D^2 (e^{-V} \bar{D}_\dot{\gamma} e^V) \right. \\ \left. + \rho (D^2 V + \frac{\xi}{2} \bar{\rho}) + \bar{\rho} (\bar{D}^2 V + \frac{\xi}{2} \rho) \right. \\ \left. -\beta D^2 \left\{ -\frac{i}{2} L_V [i \bar{D}^2 c - i D^2 \bar{c} + \coth(\frac{1}{2} L_V) (i \bar{D}^2 c + i D^2 \bar{c})] \right\} \right\} \end{aligned} \quad (73)$$

$$\begin{aligned}
& -\bar{\beta} \bar{D}^2 \left\{ -\frac{i}{2} L_V [i \bar{D}^2 \mathbf{c} - i D^2 \bar{\mathbf{c}} + \coth(\frac{1}{2} L_V) (i \bar{D}^2 \mathbf{c} + i D^2 \bar{\mathbf{c}})] \right\} \\
& + V^* \left\{ -\frac{i}{2} L_V [i \bar{D}^2 \mathbf{c} - i D^2 \bar{\mathbf{c}} + \coth(\frac{1}{2} L_V) (i \bar{D}^2 \mathbf{c} + i D^2 \bar{\mathbf{c}})] \right\} \\
& -\frac{1}{2} \mathbf{c}^* \{ \mathbf{c}, \bar{D}^2 \mathbf{c} \} + \frac{1}{2} \bar{\mathbf{c}}^* \{ \bar{\mathbf{c}}, \bar{D}^2 \bar{\mathbf{c}} \} + \beta^* \boldsymbol{\rho} + \bar{\beta}^* \bar{\boldsymbol{\rho}} \left. \right\}.
\end{aligned}$$

After expanding around the free theory of the previous section and taking the trace the BV Lagrangian becomes

$$\begin{aligned}
\mathcal{L}_{BV} = & \frac{1}{2} V^I D^\gamma \bar{D}^2 D_\gamma V_I + \frac{1}{4} V^I D^\gamma \bar{D}^2 (D_\gamma V^K V^\Lambda) f_{K\Lambda I} + \frac{1}{4} V^I \bar{D}^\gamma D^2 (\bar{D}_\gamma V^K V^\Lambda) f_{K\Lambda I} \\
& + \boldsymbol{\rho}^I (D^2 V_I + \frac{\xi}{2} \bar{\boldsymbol{\rho}}_I) + \bar{\boldsymbol{\rho}}^I (\bar{D}^2 V_I + \frac{\xi}{2} \boldsymbol{\rho}_I) \\
& - \beta^I D^2 \bar{D}^2 \mathbf{c}_I + \frac{1}{2} \beta^I D^2 (V^K (D^2 \bar{\mathbf{c}}^\Lambda - \bar{D}^2 \mathbf{c}^\Lambda)) f_{K\Lambda I} \\
& - \bar{\beta}^I \bar{D}^2 D^2 \bar{\mathbf{c}}_I + \frac{1}{2} \bar{\beta}^I \bar{D}^2 (V^K (D^2 \bar{\mathbf{c}}^\Lambda - \bar{D}^2 \mathbf{c}^\Lambda)) f_{K\Lambda I} \\
& + V^{*I} \bar{D}^2 \mathbf{c}_I + V^{*I} D^2 \bar{\mathbf{c}}_I - \frac{1}{2} V^{*I} V^K (D^2 \bar{\mathbf{c}}^\Lambda - \bar{D}^2 \mathbf{c}^\Lambda) f_{K\Lambda I} \\
& - \frac{1}{2} \mathbf{c}^{*I} \mathbf{c}^K \bar{D}^2 \mathbf{c}^\Lambda f_{K\Lambda I} + \frac{1}{2} \bar{\mathbf{c}}^{*I} \bar{\mathbf{c}}^K \bar{D}^2 \bar{\mathbf{c}}^\Lambda f_{K\Lambda I} + \beta^{*I} \boldsymbol{\rho}_I + \bar{\beta}^{*I} \bar{\boldsymbol{\rho}}_I \\
& + \dots
\end{aligned}$$

where $f_{IJ}{}^K$ are the structure constants of the internal Lie algebra $[T_I, T_J] = f_{IJ}{}^K T_K$. The algebra indices are lowered using the Cartan-Killing metric $g_{IJ} = Tr(T_I T_J)$. The constants $f_{IJK} = f_{IJ}{}^\Lambda g_{\Lambda K}$ have the properties $f_{IJK} = -f_{JIK} = -f_{IKJ}$.

Using the superspace antibracket, we find the action of \mathfrak{s} on the superfields and anti-superfields of SYM:

$$\mathfrak{s} V^I = \bar{D}^2 \mathbf{c}^I + D^2 \bar{\mathbf{c}}^I - \frac{1}{2} V^K (D^2 \bar{\mathbf{c}}^\Lambda - \bar{D}^2 \mathbf{c}^\Lambda) f_{K\Lambda}{}^I, \quad (74a)$$

$$\begin{aligned}
\mathfrak{s} V^*{}_I = & D^\gamma \bar{D}^2 D_\gamma V_I + \frac{1}{4} \left[D^\gamma \bar{D}^2 (D_\gamma V^K V^\Lambda) + D^\gamma (\bar{D}^2 D_\gamma V^K V^\Lambda) - \bar{D}^2 D^\gamma V^K D_\gamma V^\Lambda \right] f_{K\Lambda I} \quad (74b) \\
& + \frac{1}{4} \left[\bar{D}^\gamma D^2 (\bar{D}_\gamma V^K V^\Lambda) + \bar{D}^\gamma (D^2 \bar{D}_\gamma V^K V^\Lambda) - D^2 \bar{D}^\gamma V^K \bar{D}_\gamma V^\Lambda \right] f_{K\Lambda I} + D^2 \boldsymbol{\rho}_I + \bar{D}^2 \bar{\boldsymbol{\rho}}_I \\
& + \frac{1}{2} \left[D^2 \beta^K + \bar{D}^2 \bar{\beta}^K - V^{*K} \right] (D^2 \bar{\mathbf{c}}^\Lambda - \bar{D}^2 \mathbf{c}^\Lambda) f_{I\Lambda K},
\end{aligned}$$

$$\mathfrak{s} \boldsymbol{\rho}^I = 0, \quad \mathfrak{s} \boldsymbol{\rho}^*{}_I = \beta^*{}_I + D^2 V_I + \xi \bar{\boldsymbol{\rho}}_I, \quad (74c)$$

$$\mathfrak{s} \bar{\boldsymbol{\rho}}^I = 0, \quad \mathfrak{s} \bar{\boldsymbol{\rho}}^*{}_I = \bar{\beta}^*{}_I + \bar{D}^2 V_I + \xi \boldsymbol{\rho}_I, \quad (74d)$$

$$\mathfrak{s} \beta^I = \boldsymbol{\rho}^I, \quad \mathfrak{s} \beta^*{}_I = -D^2 \bar{D}^2 \mathbf{c}_I + \frac{1}{2} D^2 (V^K (D^2 \bar{\mathbf{c}}^\Lambda - \bar{D}^2 \mathbf{c}^\Lambda)) f_{K\Lambda I}, \quad (74e)$$

$$\mathfrak{s} \bar{\beta}^I = \bar{\boldsymbol{\rho}}^I, \quad \mathfrak{s} \bar{\beta}^*{}_I = -\bar{D}^2 D^2 \bar{\mathbf{c}}_I + \frac{1}{2} \bar{D}^2 (V^K (D^2 \bar{\mathbf{c}}^\Lambda - \bar{D}^2 \mathbf{c}^\Lambda)) f_{K\Lambda I}, \quad (74f)$$

$$\mathfrak{s} \mathbf{c}^I = -\frac{1}{2} \mathbf{c}^K \bar{D}^2 \mathbf{c}^\Lambda f_{K\Lambda}{}^I, \quad (74g)$$

$$\begin{aligned}
\mathfrak{s} \mathbf{c}^*{}_I = & \bar{D}^2 D^2 \boldsymbol{\beta}_I - \bar{D}^2 V^*{}_I + \frac{1}{2} \bar{D}^2 \left[(D^2 \beta^K + \bar{D}^2 \bar{\beta}^K - V^{*K}) V^\Lambda \right] f_{\Lambda I K} \quad (74h) \\
& - \frac{1}{2} \left(\mathbf{c}^{*K} \bar{D}^2 \mathbf{c}^\Lambda + \bar{D}^2 (\mathbf{c}^{*K} \mathbf{c}^\Lambda) \right) f_{I\Lambda K},
\end{aligned}$$

$$\mathfrak{s}\bar{\mathbf{c}}^I = \frac{1}{2} \bar{\mathbf{c}}^K \mathbb{D}^2 \bar{\mathbf{c}}^\Lambda f_{K\Lambda}{}^I, \quad (74i)$$

$$\begin{aligned} \mathfrak{s}\bar{\mathbf{c}}^*{}_I &= \mathbb{D}^2 \bar{\mathbb{D}}^2 \bar{\beta}_I - \mathbb{D}^2 \mathbf{V}_I^* - \frac{1}{2} \mathbb{D}^2 \left[(\mathbb{D}^2 \beta^K + \bar{\mathbb{D}}^2 \bar{\beta}^K - \mathbf{V}^{*K}) V^\Lambda \right] f_{\Lambda IK} \\ &\quad + \frac{1}{2} \left(\bar{\mathbf{c}}^{*K} \bar{\mathbb{D}}^2 \bar{\mathbf{c}}^\Lambda + \mathbb{D}^2 (\bar{\mathbf{c}}^{*K} \bar{\mathbf{c}}^\Lambda) \right) f_{I\Lambda K}. \end{aligned} \quad (74j)$$

It is straight forward to decompose these transformations to their γ and δ components as defined in (18) and (19). The action of the γ -BRST symmetry transformation on the SYM superfields and anti-superfields is:

$$\gamma V^I = \bar{\mathbb{D}}^2 \mathbf{c}^I + \mathbb{D}^2 \bar{\mathbf{c}}^I - \frac{1}{2} V^K (\mathbb{D}^2 \bar{\mathbf{c}}^\Lambda - \bar{\mathbb{D}}^2 \mathbf{c}^\Lambda) f_{K\Lambda}{}^I, \quad (75a)$$

$$\gamma \mathbf{V}^*{}_I = -\frac{1}{2} \mathbf{V}^{*K} (\mathbb{D}^2 \bar{\mathbf{c}}^\Lambda - \bar{\mathbb{D}}^2 \mathbf{c}^\Lambda) f_{I\Lambda K}, \quad (75b)$$

$$\gamma \rho^I = 0, \quad \gamma \rho^*{}_I = \beta^*{}_I, \quad (75c)$$

$$\gamma \bar{\rho}^I = 0, \quad \gamma \bar{\rho}^*{}_I = \bar{\beta}^*{}_I, \quad (75d)$$

$$\gamma \beta^I = \rho^I, \quad \gamma \beta^*{}_I = 0, \quad (75e)$$

$$\gamma \bar{\beta}^I = \bar{\rho}^I, \quad \gamma \bar{\beta}^*{}_I = 0, \quad (75f)$$

$$\gamma \mathbf{c}^I = -\frac{1}{2} \mathbf{c}^K \bar{\mathbb{D}}^2 \mathbf{c}^\Lambda f_{K\Lambda}{}^I, \quad (75g)$$

$$\gamma \mathbf{c}^*{}_I = -\bar{\mathbb{D}}^2 \mathbf{V}^*{}_I - \frac{1}{2} \bar{\mathbb{D}}^2 \left[\mathbf{V}^{*K} V^\Lambda \right] f_{\Lambda IK} - \frac{1}{2} \left(\bar{\mathbf{c}}^{*K} \bar{\mathbb{D}}^2 \mathbf{c}^\Lambda + \bar{\mathbb{D}}^2 (\bar{\mathbf{c}}^{*K} \mathbf{c}^\Lambda) \right) f_{I\Lambda K}, \quad (75h)$$

$$\gamma \bar{\mathbf{c}}^I = \frac{1}{2} \bar{\mathbf{c}}^K \mathbb{D}^2 \bar{\mathbf{c}}^\Lambda f_{K\Lambda}{}^I, \quad (75i)$$

$$\gamma \bar{\mathbf{c}}^*{}_I = -\mathbb{D}^2 \mathbf{V}_I^* + \frac{1}{2} \mathbb{D}^2 \left[\mathbf{V}^{*K} V^\Lambda \right] f_{\Lambda IK} + \frac{1}{2} \left(\bar{\mathbf{c}}^{*K} \bar{\mathbb{D}}^2 \bar{\mathbf{c}}^\Lambda + \mathbb{D}^2 (\bar{\mathbf{c}}^{*K} \bar{\mathbf{c}}^\Lambda) \right) f_{I\Lambda K} \quad (75j)$$

and the action of the Koszul-Tate complex δ on the SYM anti-superfields is:

$$\begin{aligned} \delta \mathbf{V}^*{}_I &= \mathbb{D}^\gamma \bar{\mathbb{D}}^2 \mathbb{D}_\gamma V_I + \frac{1}{4} \left[\mathbb{D}^\gamma \bar{\mathbb{D}}^2 (\mathbb{D}_\gamma V^K V^\Lambda) + \mathbb{D}^\gamma (\bar{\mathbb{D}}^2 \mathbb{D}_\gamma V^K V^\Lambda) - \bar{\mathbb{D}}^2 \mathbb{D}^\gamma V^K \mathbb{D}_\gamma V^\Lambda \right] f_{K\Lambda I} \\ &\quad + \frac{1}{4} \left[\bar{\mathbb{D}}^\gamma \mathbb{D}^2 (\bar{\mathbb{D}}_\gamma V^K V^\Lambda) + \bar{\mathbb{D}}^\gamma (\mathbb{D}^2 \bar{\mathbb{D}}_\gamma V^K V^\Lambda) - \mathbb{D}^2 \bar{\mathbb{D}}^\gamma V^K \bar{\mathbb{D}}_\gamma V^\Lambda \right] f_{K\Lambda I} + \mathbb{D}^2 \rho_I + \bar{\mathbb{D}}^2 \bar{\rho}_I \\ &\quad + \frac{1}{2} \left[\mathbb{D}^2 \beta^K + \bar{\mathbb{D}}^2 \bar{\beta}^K \right] (\mathbb{D}^2 \bar{\mathbf{c}}^\Lambda - \bar{\mathbb{D}}^2 \mathbf{c}^\Lambda) f_{I\Lambda K}, \end{aligned} \quad (76a)$$

$$\delta \rho^*{}_I = \mathbb{D}^2 V_I + \xi \bar{\rho}_I, \quad \delta \bar{\rho}^*{}_I = \bar{\mathbb{D}}^2 V_I + \xi \rho_I, \quad (76b)$$

$$\delta \beta^*{}_I = -\mathbb{D}^2 \bar{\mathbb{D}}^2 \mathbf{c}_I + \frac{1}{2} \mathbb{D}^2 \left(V^K (\mathbb{D}^2 \bar{\mathbf{c}}^\Lambda - \bar{\mathbb{D}}^2 \mathbf{c}^\Lambda) \right) f_{K\Lambda I}, \quad (76c)$$

$$\delta \bar{\beta}^*{}_I = -\bar{\mathbb{D}}^2 \mathbb{D}^2 \bar{\mathbf{c}}_I + \frac{1}{2} \bar{\mathbb{D}}^2 \left(V^K (\mathbb{D}^2 \bar{\mathbf{c}}^\Lambda - \bar{\mathbb{D}}^2 \mathbf{c}^\Lambda) \right) f_{K\Lambda I}, \quad (76d)$$

$$\delta \mathbf{c}^*{}_I = \bar{\mathbb{D}}^2 \mathbb{D}^2 \beta_I + \frac{1}{2} \bar{\mathbb{D}}^2 \left[(\mathbb{D}^2 \beta^K + \bar{\mathbb{D}}^2 \bar{\beta}^K) V^\Lambda \right] f_{\Lambda IK}, \quad (76e)$$

$$\delta \bar{\mathbf{c}}^*{}_I = \mathbb{D}^2 \bar{\mathbb{D}}^2 \bar{\beta}_I - \frac{1}{2} \mathbb{D}^2 \left[(\mathbb{D}^2 \beta^K + \bar{\mathbb{D}}^2 \bar{\beta}^K) V^\Lambda \right] f_{\Lambda IK}$$

Unlike the free theory example of section 3, the Hilbert space description of SYM can not be captured by a sole BRST charge \mathbf{Q} . The nonlinear terms require the introduction of a new kind

of product “ $*$ ” in the superfield state space. This product will assign to a pair of superfield states $(|A\rangle, |B\rangle)$ a new state $|A\rangle * |B\rangle$. Hence, for interacting theories, the Lagrangian BRST symmetry operator \mathfrak{s} corresponds to the doublet $(Q, *)$ acting in an appropriately defined Hilbert space of superfield states $\mathfrak{s}|\Psi\rangle = Q|\Psi\rangle + |\Psi\rangle * |\Psi\rangle$. This is similar to string field theory, where there is a $*$ -product and the physical spectrum of the theory is described by the equation $Q\Psi + \Psi * \Psi = 0$.

5 Summary

It has been demonstrated repeatedly that knowing the BRST description of a gauge theory is a very powerful tool. In this paper we explore the BRST description of supersymmetric gauge theories while maintaining supersymmetry manifest. For a given superspace formulated gauge theory, we follow the usual BRST procedure and construct a nilpotent BRST symmetry operator (\mathfrak{J}) by fermionizing the original gauge symmetry. For every gauge parameter superfield we introduce a Fadeev-Popov ghost superfield and two additional ghost superfields which play the role of a BRST doublet. The superspace action is deformed by an appropriate \mathfrak{J} -exact term which generates gauge fixing conditions and remove negative norm states from the physical spectrum of the theory. This deformation is further extended by the addition of anti-superfield terms according to the BV procedure. Using a superspace anti-bracket we define the nilpotent BRST-BV differential operator (\mathfrak{S}) in superspace and identify its decomposition to the superspace BRST symmetry transformation (γ) and the superspace Koszul-Tate complex (δ) . We apply this procedure to $4D$, $\mathcal{N} = 1$ super Maxwell theory and its non-abelian extension super Yang-Mills. For both theories, we derive explicit expressions for all these nilpotent operators in terms of the superspace covariant derivatives.

Moreover, for the linearized theory we explore its BRST-BFV description in terms of a nilpotent BRST charge operator (Q) acting on the Hilbert space of superfield states. Superspace BRST charges have been constructed previously for maximally supersymmetric theories by introducing pure spinors. The nilpotence of these BRST charges is automatic due to the pure spinor constraint. In this paper, we follow another approach and construct a Hilbert space BRST charge based on the algebra of the appropriate set of superspace differential operators. Specifically we consider the set of linearly independent differential operators that appear in \mathfrak{S} . Due to the algebraic properties of the supersymmetric covariant derivative, the algebra of these operators is nonlinear. Nevertheless, the Fradkin-Fradkina algorithm can be appropriately modified and applied in order to construct a manifestly supersymmetric nilpotent BRST charge without requiring pure spinor variables. By choosing an appropriate vacuum state, we show that its physical state cohomology generates the correct superspace equations. For the interacting theory, the physical spectrum of the theory can not be reproduced solely in terms of a BRST charge but additional structure must be introduced in order to capture the nonlinear terms that correspond to the gluing of superfield states in Hilbert space.

There are several interesting future directions that we want to investigate. First of all, this methodology must be applied to gauge theories with tensorial prepotential superfields such as supergravity and higher spin supermultiplets in order to explore the structure of the nilpotent operators for these theories. It is very interesting that for non-supersymmetric Maxwell theory, the physical state cohomology (33), (32) is also valid for higher spin gauge theories by simply allowing the expansion of the states in terms of bosonic oscillators that carry spacetime indices. This happens

because the gauge invariance of higher spin theories (Bianchi identities) rely on exactly the same algebra as Maxwell theory. However, for supersymmetric higher spin theories this is not true hence, we must apply the same procedure to theories described by higher rank gauge superfields. Secondly, we must explore the properties of the additional product rule ($*$) structure between superfield states required by the BRST-BFV description of interacting gauge theories. String interactions in string field theory offers an example of such structure.

Acknowledgments

The work of I. L. B was partially supported by Ministry of Education of Russian Federation, project FEWF-2020-0003. The work of K. K. and S. J. G. is supported in part by the endowment of the Ford Foundation Professorship of Physics at Brown University. Also K. K. gratefully acknowledges the support of the Brown Theoretical Physics Center.

References

- [1] C. Becchi, A. Rouet and R. Stora, “The Abelian Higgs-Kibble Model. Unitarity of the S Operator”, *Phys. Lett. B* **52** (1974) 344.
- [2] C. Becchi, A. Rouet and R. Stora, “Renormalization of the Abelian Higgs-Kibble Model”, *Commun. Math. Phys.* **42** (1975) 127.
- [3] C. Becchi, A. Rouet and R. Stora, “Renormalization of Gauge Theories”, *Annals Phys.* **98** (1976) 287.
- [4] I. V. Tyutin, “Gauge Invariance in Field Theory and Statistical Physics in Operator Formalism”, [arXiv:0812.0580](https://arxiv.org/abs/0812.0580) [hep-th].
- [5] I. A. Batalin and G. A. Vilkovisky, “Gauge Algebra and Quantization”, *Phys. Lett. B* **102** (1981) 27.
- [6] I. A. Batalin and G. A. Vilkovisky, “Quantization of Gauge Theories with Linearly Dependent Generators”, *Phys. Rev. D* **28** (1983) 2567, [Erratum: *Phys.Rev.D* 30, 508 (1984)].
- [7] I. A. Batalin and G. A. Vilkovisky, “Existence Theorem for Gauge Algebra”, *J. Math. Phys.* **26** (1985) 172.
- [8] B. L. Voronov, P. M. Lavrov and I. V. Tyutin, “Canonical Transformations and the Gauge Dependence in General Gauge Theories. (IN RUSSIAN)”, *Yad. Fiz.* **36** (1982) 498.
- [9] B. L. Voronov and I. V. Tyutin, “Formulation of Gauge Theories of General Form. I”, *Theor. Math. Phys.* **50** (1982) 218.
- [10] B. L. Voronov and I. v. Tyutin, “Formulation of Gauge Theories of General Form. II. Gauge Invariant Renormalizability and Renormalization Structure”, *Theor. Math. Phys.* **52** (1982) 628.
- [11] I. L. Buchbinder, S. J. Gates and K. Koutrolikos, “Superspace First Order Formalism, Trivial Symmetries and Electromagnetic Interactions of Linearized Supergravity”, [arXiv:2107.06854](https://arxiv.org/abs/2107.06854) [hep-th].
- [12] E. Witten, “Noncommutative Geometry and String Field Theory”, *Nucl. Phys. B* **268** (1986) 253.
- [13] M. Bochicchio, “String Field Theory in the Siegel Gauge”, *Phys. Lett. B* **188** (1987) 330.
- [14] C. B. Thorn, “Perturbation Theory for Quantized String Fields”, *Nucl. Phys. B* **287** (1987) 61.
- [15] C. B. Thorn, “String Field Theory”, *Phys. Rept.* **175** (1989) 1.
- [16] B. Zwiebach, “Closed string field theory: Quantum action and the B-V master equation”, *Nucl. Phys. B* **390** (1993) 33, [arXiv:hep-th/9206084](https://arxiv.org/abs/hep-th/9206084).

- [17] W. Taylor and B. Zwiebach, “D-branes, tachyons, and string field theory”, in Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 2001): Strings, Branes and Extra Dimensions, p. 641–759 2003, [arXiv:hep-th/0311017](#).
- [18] W. Siegel, “Introduction to string field theory”, 1988, [arXiv:hep-th/0107094](#).
- [19] X. Bekaert, I. L. Buchbinder, A. Pashnev and M. Tsulaia, “On higher spin theory: Strings, BRST, dimensional reductions”, *Class. Quant. Grav.* **21** (2004) S1457, [arXiv:hep-th/0312252](#).
- [20] I. L. Buchbinder, V. A. Krykhtin and A. Pashnev, “BRST approach to Lagrangian construction for fermionic massless higher spin fields”, *Nucl. Phys. B* **711** (2005) 367, [arXiv:hep-th/0410215](#).
- [21] I. L. Buchbinder and V. A. Krykhtin, “Gauge invariant Lagrangian construction for massive bosonic higher spin fields in D dimensions”, *Nucl. Phys. B* **727** (2005) 537, [arXiv:hep-th/0505092](#).
- [22] I. L. Buchbinder, A. Fotopoulos, A. C. Petkou and M. Tsulaia, “Constructing the cubic interaction vertex of higher spin gauge fields”, *Phys. Rev. D* **74** (2006) 105018, [arXiv:hep-th/0609082](#).
- [23] X. Bekaert, N. Boulanger, S. Cnockaert and S. Leclercq, “On killing tensors and cubic vertices in higher-spin gauge theories”, *Fortsch. Phys.* **54** (2006) 282, [arXiv:hep-th/0602092](#).
- [24] I. L. Buchbinder, V. A. Krykhtin and A. A. Reshetnyak, “BRST approach to Lagrangian construction for fermionic higher spin fields in (A)dS space”, *Nucl. Phys. B* **787** (2007) 211, [arXiv:hep-th/0703049](#).
- [25] D. Polyakov, “Interactions of Massless Higher Spin Fields From String Theory”, *Phys. Rev. D* **82** (2010) 066005, [arXiv:0910.5338 \[hep-th\]](#).
- [26] M. Henneaux, G. Lucena Gómez and R. Rahman, “Higher-Spin Fermionic Gauge Fields and Their Electromagnetic Coupling”, *JHEP* **08** (2012) 093, [arXiv:1206.1048 \[hep-th\]](#).
- [27] R. Metsaev, “BRST-BV approach to cubic interaction vertices for massive and massless higher-spin fields”, *Phys. Lett. B* **720** (2013) 237, [arXiv:1205.3131 \[hep-th\]](#).
- [28] M. Henneaux, G. Lucena Gómez and R. Rahman, “Gravitational Interactions of Higher-Spin Fermions”, *JHEP* **01** (2014) 087, [arXiv:1310.5152 \[hep-th\]](#).
- [29] I. L. Buchbinder and K. Koutrolikos, “BRST Analysis of the Supersymmetric Higher Spin Field Models”, *JHEP* **12** (2015) 106, [arXiv:1510.06569 \[hep-th\]](#).
- [30] I. L. Buchbinder, V. A. Krykhtin and H. Takata, “BRST approach to Lagrangian construction for bosonic continuous spin field”, *Phys. Lett. B* **785** (2018) 315, [arXiv:1806.01640 \[hep-th\]](#).
- [31] I. L. Buchbinder, V. A. Krykhtin, M. Tsulaia and D. Weissman, “Cubic Vertices for N=1 Supersymmetric Massless Higher Spin Fields in Various Dimensions”, *Nucl. Phys. B* **967** (2021) 115427, [arXiv:2103.08231 \[hep-th\]](#).
- [32] I. L. Buchbinder and P. M. Lavrov, “On a gauge-invariant deformation of a classical gauge-invariant theory”, *JHEP* **06** (2021) 097, [arXiv:2104.11930 \[hep-th\]](#).
- [33] I. L. Buchbinder and P. M. Lavrov, “On classical and quantum deformations of gauge theories”, *Eur. Phys. J. C* **81** (2021) 856, [arXiv:2108.09968 \[hep-th\]](#).
- [34] I. L. Buchbinder and A. A. Reshetnyak, “General cubic interacting vertex for massless integer higher spin fields”, *Phys. Lett. B* **820** (2021) 136470, [arXiv:2105.12030 \[hep-th\]](#).
- [35] N. Berkovits, “Super Poincare covariant quantization of the superstring”, *JHEP* **04** (2000) 018, [arXiv:hep-th/0001035](#).

- [36] M. Cederwall, B. E. W. Nilsson and D. Tsimpis, “Spinorial cohomology and maximally supersymmetric theories”, *JHEP* **02** (2002) 009, [arXiv:hep-th/0110069](#).
- [37] N. Berkovits, “ICTP lectures on covariant quantization of the superstring”, *ICTP Lect. Notes Ser.* **13** (2003) 57, [arXiv:hep-th/0209059](#).
- [38] N. Berkovits, “Pure spinor formalism as an N=2 topological string”, *JHEP* **10** (2005) 089, [arXiv:hep-th/0509120](#).
- [39] M. Cederwall, “D=11 supergravity with manifest supersymmetry”, *Mod. Phys. Lett. A* **25** (2010) 3201, [arXiv:1001.0112 \[hep-th\]](#).
- [40] M. Cederwall, “Pure spinor superfields – an overview”, *Springer Proc. Phys.* **153** (2014) 61, [arXiv:1307.1762 \[hep-th\]](#).
- [41] S. J. Gates, Jr., M. T. Grisaru, U. Lindstrom, M. Rocek, W. Siegel, P. van Nieuwenhuizen and A. E. van de Ven, “Lorentz Covariant Quantization of the Heterotic Superstring”, *Phys. Lett. B* **225** (1989) 44.
- [42] S. J. Gates, “Unconstrained BRST: A New prototype for covariant string field theory?”, *Conf. Proc. C* **9003122** (1990) 244.
- [43] M. Henneaux and C. Teitelboim, “Quantization of gauge systems”, 1992.
- [44] E. S. Fradkin and T. E. Fradkina, “Quantization of Relativistic Systems with Boson and Fermion First and Second Class Constraints”, *Phys. Lett. B* **72** (1978) 343.
- [45] E. S. Fradkin and G. A. Vilkovisky, “Quantization of Relativistic Systems with Constraints”, *Phys. Lett. B* **55** (1975) 224.
- [46] I. A. Batalin and G. A. Vilkovisky, “Relativistic S Matrix of Dynamical Systems with Boson and Fermion Constraints”, *Phys. Lett. B* **69** (1977) 309.
- [47] I. A. Batalin and E. S. Fradkin, “A Generalized Canonical Formalism and Quantization of Reducible Gauge Theories”, *Phys. Lett. B* **122** (1983) 157.
- [48] M. Henneaux, “Hamiltonian Form of the Path Integral for Theories with a Gauge Freedom”, *Phys. Rept.* **126** (1985) 1.
- [49] K. Schoutens, A. Sevrin and P. van Nieuwenhuizen, “Quantum BRST Charge for Quadratically Nonlinear Lie Algebras”, *Commun. Math. Phys.* **124** (1989) 87.
- [50] I. L. Buchbinder and P. M. Lavrov, “Classical BRST charge for nonlinear algebras”, *J. Math. Phys.* **48** (2007) 082306, [arXiv:hep-th/0701243](#).
- [51] M. Henneaux, “Lectures on the Antifield-BRST Formalism for Gauge Theories”, *Nucl. Phys. B Proc. Suppl.* **18** (1990) 47.