

$SU_2(\mathbb{C})$ SYMMETRY IN QUANTUM SPIN CHAIN GROUND STATES AND HALDANE'S CONJECTURE

ANILESH MOHARI

Abstract

In this paper, we prove that any translation and $SU_2(\mathbb{C})$ -invariant pure state of $\mathbb{M} = \otimes_{k \in \mathbb{Z}} M_d^{(k)}(\mathbb{C})$, that is also real, lattice symmetric and reflection positive with a certain twist $r_0 \in U_d(\mathbb{C})$, is finitely correlated and its two-point spatial correlation function decays exponentially whenever d is an odd integer. In particular, the Heisenberg iso-spin anti-ferromagnetic integer spin model admits unique low temperature limiting ground state and its spatial correlation function decays exponentially. The unique low temperature limiting ground state of the Hamiltonian is determined by the unique solution to Clebsch-Gordon inter-twinning isometry between two representations of $SU_2(\mathbb{C})$.

1. Introduction

In this paper, we resume our investigation [26] for various order properties of ground states of translation invariant Hamiltonian models [8, 34] in the two-side infinite quantum spin chain $\mathbb{M} = \otimes_{k \in \mathbb{Z}} M_d^{(k)}(\mathbb{C})$ of the following formal form

$$(1) \quad H = \sum_{n \in \mathbb{Z}} \theta^n(h_0),$$

with $h_0^* = h_0 \in \mathbb{M}_{loc}$, where \mathbb{M}_{loc} is the union of local sub-algebras of \mathbb{M} and θ is the right translation on \mathbb{M} . In particular, our results are aimed to investigate the set of ground states for the Heisenberg anti-ferromagnet iso-spin model H^{XXX} [7] with nearest neighbour interactions

$$(2) \quad h_0^{xxx} = J(\sigma_x^0 \otimes \sigma_x^1 + \sigma_y^0 \otimes \sigma_y^1 + \sigma_z^0 \otimes \sigma_z^1),$$

where σ_x^k, σ_y^k and σ_z^k are Pauli spin matrices located at lattice site $k \in \mathbb{Z}$ and $J > 0$ is a constant. It is well known that any finite volume truncation of H^{XXX} with periodic boundary condition admits a unique ground state [2, 7]. However, no clear picture had emerged in the past literature about the set of ground states for the two sided infinite volume anti-ferromagnet Heisenberg H^{XXX} model. Nevertheless, many interesting results on ground states, known for other specific Heisenberg type of models [21], such as Ghosh-Majumder (GM) model [15] and Affleck-Kenedy-Lieb-Tasaki (AKLT) model [1], gave interesting conjectures on the general behaviour of ground states and its physical implication for anti-ferromagnetic Hamiltonian H^{XXX} model.

Haldane [2] conjecture stated that H^{XXX} has a unique ground state and the ground state admits a mass gap with its two-point spatial correlation function decaying exponentially for integer spin s (odd integer d , where $d = 2s + 1$). Whereas for the even values of d (i.e. s is a $\frac{1}{2}$ odd integer spin, where $d = 2s + 1$),

2020 *Mathematics Subject Classification.* 46L.

Key words and phrases. Uniformly hyperfinite factors. Cuntz algebra, Popescu dilation, Spontaneous symmetry breaking, Heisenberg iso-spin anti-ferromagnetic model, ground states, reflection positive, Haldane's conjecture .

...

the conjecture stated that H^{XXX} has a unique ground state with no mass gap and its two-point spatial correlation function does not decay exponentially. A well known result of Affleck and Lieb [2] says: if H^{XXX} admits a unique ground state for even values of d then the ground state has no mass gap and its two-point spatial correlation function does not decay exponentially. In contrast, if the integer spin H^{XXX} model admits a unique ground state with a mass gap, a recent result [29] says that its two-point spatial correlation function decays exponentially. Thus the uniqueness of the hypothesis on the ground state for H^{XXX} model is a critical issue to settle a part of the conjecture.

For even values of d , we have proved in [26] that ground state of anti-ferromagnetic Heisenberg model H^{XXX} is not unique and any low temperature limiting ground state of H^{XXX} is not pure. Nevertheless, any low temperature limiting ground state admits no mass gap in its spectrum and its spatial correlation function does not decay exponentially provided the limiting ground state is non degenerate. In the present exposition we resume general mathematical set up [25, 26] and address Haldane's conjecture for odd values for d .

In the following text, we will now formulate the problem in the general framework of C^* -dynamical system [8, 9] valid for two-sided one-dimensional quantum spin chain models. The uniformly hyper-finite [32] C^* -algebra $\mathbb{M} = \otimes_{k \in \mathbb{Z}} M_d^{(k)}(\mathbb{C})$ of infinite tensor product of $d \times d$ -square matrices $M_d^{(k)}(\mathbb{C}) \equiv M_d(\mathbb{C})$, levelled by k in the lattice \mathbb{Z} of integers, is the norm closure of the algebraic inductive limit of the net of finite dimensional C^* algebras $\mathbb{M}_\Lambda = \otimes_{k \in \Lambda} M_d^{(k)}(\mathbb{C})$, where $\Lambda \subset \mathbb{Z}$ are finite subsets and an element Q in \mathbb{M}_{Λ_1} is identified with the element $Q \otimes I_{\Lambda_2 \cap \Lambda_1^c}$ in \mathbb{M}_{Λ_2} , i.e. by the inclusion map if $\Lambda_1 \subseteq \Lambda_2$, where Λ^c is the complementary set of Λ in \mathbb{Z} . We will use the symbol \mathbb{M}_{loc} to denote the union of all local algebras $\{\mathbb{M}_\Lambda : \Lambda \subset \mathbb{Z}, |\Lambda| < \infty\}$. Thus \mathbb{M} is a quasi-local C^* -algebra with local algebras $\{\mathbb{M}_\Lambda : |\Lambda| < \infty\}$ and $\mathbb{M}'_\Lambda = \mathbb{M}_{\Lambda^c}$, where \mathbb{M}'_Λ is the commutant of \mathbb{M}_Λ in \mathbb{M} . We refer readers to Chapter 6 of [8] for more details on quasi-local C^* -algebras.

The lattice \mathbb{Z} is a group under addition and for each $n \in \mathbb{Z}$, we have an automorphism θ^n , extending the translation action, which takes $Q^{(k)}$ to $Q^{(k+n)}$ for any $Q \in M_d(\mathbb{C})$ and $k \in \mathbb{Z}$, by the linear and multiplicative properties on \mathbb{M} . A unital positive linear functional ω of \mathbb{M} is called *state*. It is called *translation-invariant* if $\omega = \omega\theta$. A linear automorphism or anti-automorphism β [16] on \mathbb{M} is called *symmetry* for ω if $\omega\beta = \omega$. Our primary objective is to study translation-invariant states and their symmetries that find relevance in Hamiltonian dynamics of quantum spin chain models H [15, 33, 34].

We consider [8, 33, 34] quantum spin chain Hamiltonian in one dimensional lattice \mathbb{M} of the following form

$$(3) \quad H = \sum_{n \in \mathbb{Z}} \theta^n(h_0)$$

for $h_0^* = h_0 \in \mathbb{M}_{loc}$, where the formal sum in (3) gives a group of auto-morphisms $\alpha = (\alpha_t : t \in \mathbb{R})$ by the thermodynamic limit: $\lim_{\Lambda_\eta \uparrow \mathbb{Z}} \|\alpha_t^{\Lambda_\eta}(A) - \alpha_t(A)\| = 0$ for all $A \in \mathbb{M}$ and $t \in \mathbb{R}$ for a net of finite subsets Λ_η of \mathbb{Z} with uniformly bounded surface energy, where automorphisms $\alpha_t^\Lambda(x) = e^{itH_\Lambda} x e^{-itH_\Lambda}$ is determined by the finite subset Λ of \mathbb{Z}^k and $H_\Lambda = \sum_{n \in \Lambda} \theta^n(h_0)$. Furthermore, the limiting automorphism (α_t) does not depend on the net that we choose in the thermodynamic limit $\Lambda_\eta \uparrow \mathbb{Z}$, provided the surface energies of H_{Λ_η} are kept uniformly bounded. The uniquely determined group of automorphisms (α_t) on \mathbb{M} is called *Heisenberg flows* of H .

In particular, we have $\alpha_t \circ \theta^n = \theta^n \circ \alpha_t$ for all $t \in \mathbb{R}$ and $n \in \mathbb{Z}$. Any linear automorphism or anti-automorphism β on \mathbb{M}_{loc} , keeping the formal sum (3) in H invariant, will also commute with (α_t) .

A state ω is called *stationary* for H if $\omega\alpha_t = \omega$ on \mathbb{M} for all $t \in \mathbb{R}$. The set of stationary states of H is a non-empty compact convex set and has been extensively studied in the last few decades within the framework of ergodic theory for C^* -dynamical systems [12, Chapter 4]. However, a stationary state of H need not be always translation-invariant. A stationary state ω of \mathbb{M} for H is called β -KMS state at an inverse positive temperature $\beta > 0$ if there exists a function $z \rightarrow f_{A,B}(z)$, analytic on the open strip $0 < \text{Im}(z) < \beta$, bounded continuous on the closed strip $0 \leq \text{Im}(z) \leq \beta$ with boundary condition

$$f_{A,B}(t) = \omega_\beta(\alpha_t(A)B), \quad f_{A,B}(t + i\beta) = \omega_\beta(\alpha_t(B)A)$$

for all $A, B \in \mathbb{M}$. Using weak* compactness of convex set of states on \mathcal{M} , finite volume Gibbs state $\omega_{\beta,\Lambda}$ is used to prove existence of a KMS state ω_β for (α_t) at inverse positive temperature $\beta > 0$. The set of KMS states of H at a given inverse positive temperature β is singleton set i.e. there is a unique β KMS-state at a given inverse positive temperature $\beta = \frac{1}{kT}$ for H which has a finite range interaction [3,4,16] and thus inherits translation and other symmetry of the Hamiltonian. The unique KMS state of H at a given inverse temperature is ergodic for translation dynamics. This gives a strong motivation to study translation-invariant states in a more general framework of C^* -dynamical systems [8].

A state ω of \mathbb{M} is called *ground state* for H , if the following two conditions are satisfied:

- (a) $\omega(\alpha_t(A)) = \omega(A)$ for all $t \in \mathbb{R}$;
- (b) If we write on the GNS space $(\mathcal{H}_\omega, \pi_\omega, \zeta_\omega)$ of (\mathbb{M}, ω) ,

$$\alpha_t(\pi_\omega(A)) = e^{itH_\omega} \pi_\omega(A) e^{-itH_\omega}$$

for all $A \in \mathbb{M}$ with $H_\omega \zeta_\omega = 0$, then $H_\omega \geq 0$.

Furthermore, we say a ground state ω is *non-degenerate*, if null space of H_ω is spanned by ζ_ω only. We say ω has a *mass gap*, if the spectrum $\sigma(H_\omega)$ of H_ω is a subset of $\{0\} \cup [\delta, \infty)$ for some $\delta > 0$. For a wide class of spin chain models [27], which includes Hamiltonian H with finite range interaction, h_0 being in \mathbb{M}_{loc} , the existence of a non vanishing spectral gap of a ground state ω of H implies exponential decaying two-point spatial correlation functions. Now we present a precise definition for exponential decay of two-point spatial correlation functions of a state ω of \mathbb{M} . We use symbol Λ_m^c for complementary set of the finite volume box $\Lambda_m = \{n : -m \leq n \leq m\}$ for $m \geq 1$.

DEFINITION 1.1. Let ω be a translation-invariant state of \mathbb{M} . We say that the two-point spatial correlation functions of ω *decay exponentially*, if there exists a $\delta > 0$ satisfying the following condition: for any two local elements $Q_1, Q_2 \in \mathbb{M}$ and $\epsilon > 0$, there exists an integer $m \geq 1$ such that

$$(4) \quad e^{\delta|n|} |\omega(Q_1 \theta^n(Q_2)) - \omega(Q_1)\omega(Q_2)| \leq \epsilon$$

for all $n \in \Lambda_m^c$.

By taking low temperature limit of ω_β as $\beta \rightarrow \infty$, one also proves existence of a ground state for H [6, 9, 32, 33]. On the contrary to KMS states, the set of ground states is a convex face in the set of the convex set of (α_t) invariant states of \mathbb{M} and its extreme points are *pure* states of \mathbb{M} i.e. A state is called

pure if it can not be expressed as convex combination of two different states of \mathbb{M} . Thus low temperature limit points of unique β -KMS states give ground states for the Hamiltonian H inheriting translation and other symmetry of the Hamiltonian. In general the set of ground states need not be a singleton set and there could be other states which are not translation invariant but still a ground state for a translation invariant Hamiltonian. Ising model admits non translation invariant ground states known as Néel state [9]. However ground states that appear as low temperature limit of β -KMS states of a translation invariant Hamiltonian, inherit translation and other symmetry of the Hamiltonian. In particular if ground state of a translation invariant Hamiltonian model (3) is unique, then the ground state is a translation invariant pure state.

Let $Q \rightarrow \tilde{Q}$ be the automorphism on \mathbb{M} that maps an element

$$Q = Q_{-l}^{(-l)} \otimes Q_{-l+1}^{(-l+1)} \otimes \dots \otimes Q_{-1}^{(-1)} \otimes Q_0^{(0)} \otimes Q_1^{(1)} \dots \otimes Q_n^{(n)}$$

by reflecting around the point $\frac{1}{2}$ of the lattice \mathbb{Z} to

$$\tilde{Q} = Q_n^{(-n+1)} \dots \otimes Q_1^{(0)} \otimes Q_0^{(1)} \otimes Q_{-1}^{(2)} \otimes \dots \otimes Q_{-l+1}^{(l)} \otimes Q_{-l}^{(l+1)}$$

for all $n, l \geq 1$ and $Q_{-l}, \dots, Q_{-1}, Q_0, Q_1, \dots, Q_n \in M_d(\mathbb{C})$.

For a state ω of \mathbb{M} , we set a state $\tilde{\omega}$ of \mathbb{M} by

$$(5) \quad \tilde{\omega}(Q) = \omega(\tilde{Q})$$

for all $Q \in \mathbb{M}$. Thus $\omega \rightarrow \tilde{\omega}$ is an affine one to one onto map on the convex set of states of \mathbb{M} . The state $\tilde{\omega}$ is translation-invariant if and only if ω is translation-invariant state. We say a state ω is *lattice reflection-symmetric* or in short *lattice symmetric* if $\omega = \tilde{\omega}$.

The group of unitary matrices $u \in U_d(\mathbb{C})$ acts naturally on \mathbb{M} as a group of automorphisms of \mathbb{M} defined by

$$(6) \quad \beta_u(Q) = (\dots \otimes u \otimes u \otimes \dots)Q(\dots \otimes u^* \otimes u^* \otimes u^* \dots)$$

We also set automorphism $\tilde{\beta}_u$ on \mathbb{M} defined by

$$(7) \quad \tilde{\beta}_u(Q) = \beta_u(\tilde{Q})$$

for all $Q \in \mathbb{M}$. So for $u, w \in U_d(\mathbb{C})$, we have

$$\tilde{\beta}_u \tilde{\beta}_w = \beta_{uw}$$

In particular, $\tilde{\alpha}_w^2(Q) = Q$ for all $Q \in \mathbb{M}$ if and only if $w^2 = I_d$. We say a state ω of \mathbb{M} is *lattice symmetric with a twist* $w \in U_d(\mathbb{C})$ if

$$(8) \quad w^2 = I_d, \quad \omega(\tilde{\beta}_w(Q)) = \omega(Q)$$

We fix an orthonormal basis $e = (e_i)$ of \mathbb{C}^d and $Q^t \in M_d(\mathbb{C})$ be the transpose of $Q \in M_d(\mathbb{C})$ with respect to an orthonormal basis (e_i) for \mathbb{C}^d (not complex conjugate). Let $Q \rightarrow Q^t$ be the linear anti-automorphism map on \mathbb{M} that takes an element

$$Q = Q_0^{(l)} \otimes Q_1^{(l+1)} \otimes \dots \otimes Q_m^{(l+m)}$$

to its transpose with respect to the basis $e = (e_i)$ defined by

$$Q^t = Q_0^{t(l)} \otimes Q_1^{t(l+1)} \otimes \dots \otimes Q_m^{t(l+m)},$$

where Q_0, Q_1, \dots, Q_m are arbitrary elements in $M_d(\mathbb{C})$. We also note that Q^t depends on the basis e that we choose and avoided use of a suffix e . We assume that it won't confuse an attentive reader since we have fixed an orthonormal basis (e_i) for our

consideration through out this paper. For more general $Q \in \mathbb{M}_{loc}$, we define Q^t by extending linearly and take the unique bounded linear extension for any $Q \in \mathbb{M}$. For a state ω of \mathbb{M} , we define a state $\bar{\omega}$ on \mathbb{M} by the following prescription

$$(9) \quad \bar{\omega}(Q) = \omega(Q^t)$$

Thus the state $\bar{\omega}$ is translation-invariant if and only if ω is translation-invariant. We say ω is *real*, if $\bar{\omega} = \omega$. The formal Hamiltonian H is called *reflection symmetric with twist w* if $\beta_w(\tilde{H}) = H$ and *real* if $H^t = H$.

We also set a conjugate linear map $Q \rightarrow \bar{Q}$ on \mathbb{M} with respect to the basis (e_i) for \mathbb{C}^d defined by extending the identity action on elements $\dots I_d \otimes |e_{i_0}\rangle\langle e_{j_0}|^{(k)} \otimes |e_{i_1}\rangle\langle e_{j_1}|^{(k+1)} \otimes \dots |e_{i_n}\rangle\langle e_{j_n}|^{(k+n)} \otimes I_d \dots$, $1 \leq i_k, j_k \leq d$, $k \in \mathbb{Z}$, $n \geq 0$ anti-linearly. Thus by our definition we have

$$Q^* = \bar{Q}^t$$

and

$$(\bar{Q})^* = \overline{Q^*}$$

We set the following anti-linear reflection map $\mathcal{J}_w : \mathbb{M} \rightarrow \mathbb{M}$ with twist $w \in U_d(\mathbb{C})$, defined by

$$(10) \quad \mathcal{J}_w(Q) = \overline{\beta_w(\bar{Q})}$$

for all $Q \in \mathbb{M}$.

Following a well known notion [12], a state ω on \mathbb{M} is called *reflection positive with a twist $r_0 \in U_d(\mathbb{C})$* , $r_0^2 = I_d$, if

$$(11) \quad \omega(\mathcal{J}_{r_0}(Q)Q) \geq 0$$

for all $Q \in \mathbb{M}_R$. Thus the notion of reflection positivity also depends explicitly on the underlining fixed orthonormal basis $e = (e_i)$ of \mathbb{C}^d .

Let G be a compact group and $g \rightarrow u(g)$ be a d -dimensional unitary representation of G . By γ_g we denote the product action of G on the infinite tensor product \mathbb{M} induced by $u(g)$,

$$(12) \quad \gamma_g(Q) = (\dots \otimes u(g) \otimes u(g) \otimes u(g) \dots) Q (\dots \otimes u(g)^* \otimes u(g)^* \otimes u(g)^* \dots)$$

for any $Q \in \mathbb{M}$, i.e. $\gamma_g = \beta_{u(g)}$. We say ω is G -invariant, if

$$(13) \quad \omega(\gamma_g(Q)) = \omega(Q)$$

for all $Q \in \mathbb{M}_{loc}$. If $G = U_d(\mathbb{C})$ and $u : U_d(\mathbb{C}) \rightarrow U_d(\mathbb{C})$ is the natural representation $u(g) = g$, then we will identify the notation β_g with γ_g for simplicity. Formal Hamiltonian H given in (3) is called G -gauge invariant if $\gamma_g(H) = H$ for all $g \in G$.

We recall now [20, 30] if H in (3) has the following form

$$(14) \quad -H = B + \mathcal{J}_{r_0}(B) + \sum_i C_i \mathcal{J}_{r_0}(C_i)$$

for some $B, C_i \in \mathbb{M}_R$ then the unique KMS state at inverse positive temperature β is reflection positive with the twist r_0 . We refer to [14] for details, which we will cite frequently while dealing with examples satisfying (12). Since the weak*-limit of a sequence of reflection positive states with the twist r_0 is also a reflection positive state with the twist r_0 , weak*-limit points of the unique β -KMS state of H as $\beta \rightarrow \infty$, are also reflection positive with the twist r_0 . Thus any weak* low temperature limit point ground state of H is reflection positive with a twist r_0 if

H is given by (14). In particular, the unique β -KMS state of anti-ferromagnetic H^{XXX} model is real and reflection symmetric and reflection positive with twist r_0 since H^{XXX} admits the form (12) [14] with $r_0 = \sigma_y$. Furthermore H^{XXX} admits $SU_2(\mathbb{C})$ gauge symmetry with irreducible representation $g \rightarrow u(g)$.

In general, for a translation invariant pure state ω , it is known that $\pi_\omega(\mathbb{M}_R)''$ is either a type-I or a type-III factor [23]. A pure mathematical question that arise here: Do these additional symmetries of ω help to understand what type of factor $\pi_\omega(\mathbb{M}_L)''$ is? In the present exposition, as an application of our main mathematical results of [26], we will prove the following theorem in the forth section.

THEOREM 1.2. *Let ω be a translation invariant, real, lattice symmetric and reflection positive with twist $r_0 \in U_d(\mathbb{C})$ state of $\mathbb{M} = \otimes_{k \in \mathbb{Z}} M_d^{(k)}(\mathbb{C})$ and the following two statements be true for odd values of d :*

- (a) ω is pure;
- (b) ω is $SU_2(\mathbb{C})$ -invariant, where $g \rightarrow u(g) \in U_d(\mathbb{C})$ in (13) is an irreducible representation of $SU_2(\mathbb{C})$ satisfying

$$(15) \quad r_0^2 = I_d, \quad r_0 u(g) r_0^* = u(\bar{g})$$

for all $g \in SU_2(\mathbb{C})$, where the matrix conjugation with respect to an orthonormal basis $e = (e_i)$ of \mathbb{C}^d .

Then $\pi_\omega(\mathbb{M}_L)''$ is a type-I factor and two-point spatial correlation function of ω decays exponentially.

As an application of Theorem 1.2, we will prove the following theorems in section 5.

THEOREM 1.3. *Let H be a translation invariant Hamiltonian of the form $H = \sum_{k \in \mathbb{Z}} \theta_k(h_0)$ with $h_0 = h_0^* \in \mathbb{M}_{loc}$. and H be also $SU_2(\mathbb{C})$ invariant with an irreducible representation $g \rightarrow u(g)$ of $SU_2(\mathbb{C})$ and r_0 be the element in $U_d(\mathbb{C})$ satisfying (15). Let H be also real (with respect to the basis $e = (e_i)$), lattice reflection symmetric and unique β -KMS at inverse positive temperature be lattice symmetric and reflection positive with the twist $r_0 \in U_d(\mathbb{C})$.*

Then any low temperature limiting ground state of H admits ergodic decomposition, in the convex set of real, lattice reflection symmetric and translation invariant states, satisfying the following:

- (a) If d is an odd integer, then all extreme points in the ergodic decomposition are pure and $SU_2(\mathbb{C})$ invariant ground states of H ;
- (b) If d is an even integer, then none of its extreme points in the ergodic decomposition are factor states of \mathbb{M} though $SU_2(\mathbb{C})$ invariant ground states of H .

At this point we recall well known results valid for a class of $SU_2(\mathbb{C})$ invariant Hamiltonians investigated in [5, 35] that are not of the form (14) and thus not evident that their unique finite temperature states admit reflection positive property. As an example, we can verify our claim easily that the Hamiltonian investigated in [35] is not reflection positive with twist.

Besides, in the general framework [30], it is also well known that mass gap of such a Hamiltonian in its ground state implies that its spatial correlation function

decays exponentially, though the converse statement in the general framework is not true. For counter examples, we refer to example 2, page 596 in [28].

Nevertheless, the converse statement is likely to be true, for Hamiltonian of the form (14) with the additional discrete and continuous symmetry. We include a proof for the following theorem in support (but not assured) of Haldane's conjecture in section 6.

THEOREM 1.4. *Low temperature limiting ground state of anti-ferromagnet Heisenberg H^{XXX} model is unique and pure for odd values of $d = 3$. Moreover, the state is finitely correlated and its spatial two-point correlation function decays exponentially.*

Thus the important question that remains to be answered whether the limiting ground state $\omega_{\frac{1}{2}}$ for integer spin H^{XXX} model is having a mass gap in its spectrum from its ground state. Also note that Theorem 1.4 does not rule out possible existence of Néel type of ground states.

The paper is organized as follows: In section 2 and 3, we will recall basic mathematical set up required from our earlier paper [26] and explain basic ideas involved in the proof of Theorem 1.2. In section 4, we prove Theorem 1.2. In section 5, we include a proof for Theorem 1.3. In the last section, we will illustrate our results with models of physical interest such as H^{XXX} and H^{AKLT} anti-ferromagnetic models. In particular, we will give proof of Theorem 1.4. One can use similar computation for a possible proof extending Theorem 1.4 for any odd values of d .

2. Amalgamated representation of \mathcal{O}_d and $\tilde{\mathcal{O}}_d$:

A state ψ on a C^* -algebra \mathcal{A} is called *factor*, if the center of the von-Neumann algebra $\pi_\psi(\mathcal{A})''$ is trivial, where $(\mathcal{H}_\psi, \pi_\psi, \zeta_\psi)$ is the Gelfand-Naimark-Segal (GNS) space associated with ψ on \mathcal{A} [8] and $\pi_\psi(\mathcal{A})''$ is the double commutant of $\pi_\psi(\mathcal{A})$ and $\psi(x) = \langle \zeta_\psi, \pi_\psi(x)\zeta_\psi \rangle$. Here we fix our convention that Hilbert spaces that are considered here are always equipped with inner products $\langle \cdot, \cdot \rangle$ which are linear in the second variable and conjugate linear in the first variable. A state ψ on \mathcal{A} is called *pure*, if $\pi_\psi(\mathcal{A})'' = \mathcal{B}(\mathcal{H}_\psi)$, the algebra of all bounded operators on \mathcal{H}_ψ .

We recall that the Cuntz algebra $\mathcal{O}_d (d \in \{2, 3, \dots\})$ [10] is the universal unital C^* -algebra generated by the elements $\{s_1, s_2, \dots, s_d\}$ subjected to the following relations:

$$(16) \quad s_i^* s_j = \delta_j^i I, \quad \sum_{1 \leq i \leq d} s_i s_i^* = I$$

Let $\Omega = \{1, 2, 3, \dots, d\}$ be a set of d elements. \mathcal{I} be the set of finite sequences $I = (i_1, i_2, \dots, i_m)$ of elements, where $i_k \in \Omega$ and $m \geq 1$ and we use notation $|I|$ for the cardinality of I . We also include null set denoted by \emptyset in the collection \mathcal{I} and set $s_\emptyset = s_\emptyset^* = I$ identity of \mathcal{O}_d and $s_I = s_{i_1} \dots s_{i_m} \in \mathcal{O}_d$ and $s_I^* = s_{i_m}^* \dots s_{i_1}^* \in \mathcal{O}_d$.

The group $U_d(\mathbb{C})$ of $d \times d$ unitary matrices acts canonically on \mathcal{O}_d as follows:

$$\beta_u(s_i) = \sum_{1 \leq j \leq d} u_i^j s_j$$

for $u = ((u_j^i) \in U_d(\mathbb{C}))$. In particular, the gauge action is defined by

$$\beta_z(s_i) = z s_i, \quad z \in \mathbb{T} = S^1 = \{z \in \mathbb{C} : |z| = 1\}.$$

The fixed point sub-algebra of \mathcal{O}_d under the gauge action i.e., $\{x \in \mathcal{O}_d : \beta_z(x) = x, z \in S^1\}$ is the closure of the linear span of all Wick ordered monomials of the form

$$(17) \quad s_{i_1} \dots s_{i_k} s_{j_k}^* \dots s_{j_1}^* : I = (i_1, \dots, i_k), J = (j_1, j_2, \dots, j_k)$$

and is isomorphic to the uniformly hyper-finite C^* sub-algebra

$$\mathbb{M}_R = \otimes_{1 \leq k < \infty} M_d^{(k)}(\mathbb{C})$$

of \mathbb{M} , where the isomorphism carries the Wick ordered monomial (20) into the following matrix element

$$(18) \quad |e^{i_1}\rangle \langle e_{j_1}|^{(1)} \otimes |e^{i_2}\rangle \langle e_{j_2}|^{(2)} \otimes \dots \otimes |e^{i_k}\rangle \langle e_{j_k}|^{(k)} \otimes 1 \otimes 1 \dots$$

We use notation UHF_d for the fixed point C^* sub-algebra of \mathcal{O}_d under the gauge group action ($\beta_z : z \in S^1$). The restriction of β_u to UHF_d is then carried into action

$$Ad(u) \otimes Ad(u) \otimes Ad(u) \otimes \dots$$

on \mathbb{M}_R .

We also define the canonical endomorphism λ on \mathcal{O}_d by

$$(19) \quad \lambda(x) = \sum_{1 \leq i \leq d} s_i x s_i^*$$

and the isomorphism carries λ restricted to UHF_d into the one-sided shift

$$y_1 \otimes y_2 \otimes \dots \rightarrow 1 \otimes y_1 \otimes y_2 \dots$$

on \mathbb{M}_R . We note for all $u \in U_d(\mathbb{C})$ that $\lambda\beta_u = \beta_u\lambda$ on \mathcal{O}_d and so in particular, also on UHF_d .

Let ω' be a λ -invariant state on the UHF_d sub-algebra of \mathcal{O}_d . Following [16], section 7] and ω be the inductive limit state ω of $\mathbb{M} \equiv \tilde{\text{UHF}}_d \otimes \text{UHF}_d$. In other word $\omega' = \omega_R$ once we make the identification UHF_d with \mathbb{M}_R . We consider the set

$$K_\omega = \{\psi : \psi \text{ is a state on } \mathcal{O}_d \text{ such that } \psi\lambda = \psi \text{ and } \psi|_{\text{UHF}_d} = \omega_R\}$$

By taking invariant mean on an extension of ω_R to \mathcal{O}_d , we verify that K_ω is non empty and K_ω is clearly convex and compact in the weak topology. In case ω is an ergodic state (extremal state) then, ω_R is as well an extremal state in the set of λ -invariant states of \mathbb{M} . Thus K_ω is a face in the λ invariant states. Now we recall Lemma 7.4 of [9] in the following proposition which quantifies what we can gain by considering a factor state on \mathcal{O}_d instead of its restriction to UHF_d .

Our next two propositions are adapted from results in section 6 and section 7 of [9] as stated in the present form in Proposition 2.5 and Proposition 2.6 in [25].

PROPOSITION 2.1. *Let ω be a translation invariant ergodic state of \mathbb{M} then K_ω is a face in the convex set of λ -invariant states of \mathcal{O}_d . Moreover the following holds:*

(a) *An element $\psi \in K_\omega$ is ergodic if and only if ψ is a factor state. Furthermore, any other extremal point in K_ω is of the form $\psi\beta_z$ for some $z \in S^1$;*

(b) *The close subgroup $H = \{z \in S^1 : \psi\beta_z = \psi\}$ is independent of the extremal point $\psi \in K_\omega$ of our choice;*

PROOF. For the proof for (a) and (b), we refer to Lemma 7.4 in [9]. ■

PROPOSITION 2.2. *Let ψ be a λ invariant ergodic state on \mathcal{O}_d and $(\mathcal{H}_\psi, \pi_\psi, \zeta_\psi)$ be its GNS representation. Then the following holds:*

(a) *The closed subgroup $H = \{z \in S^1 : \psi\beta_z = \psi\}$ is equal to*

$$\{z \in S^1 : \beta_z \text{ extends to an automorphism of } \pi_\psi(\mathcal{O}_d)''\}$$

(b) *Let \mathcal{O}_d^H be the fixed point sub-algebra in \mathcal{O}_d under the gauge group $\{\beta_z : z \in H\}$. Then $\pi_\psi(\mathcal{O}_d^H)'' = \pi_\psi(\text{UHF}_d)''$.*

(c) *Let ω' be a λ -invariant state of UHF_d algebra and $\pi_{\omega'}(\text{UHF}_d)''$ is a type-I factor, then there exists a λ -invariant factor state ψ on \mathcal{O}_d extending ω' such that*

$$\pi_\psi(\text{UHF}_d)'' = \pi_\psi(\mathcal{O}_d)''$$

PROOF. For a proof, we refer to Proposition 2.2 in [26]. ■

PROPOSITION 2.3. *Let $(\mathcal{H}_\psi, \pi_\psi, \zeta_\psi)$ be the GNS representation of a λ invariant state ψ on \mathcal{O}_d and P be the support projection of the normal state $\psi_{\zeta_\psi}(X) = \langle \zeta_\psi, X\zeta_\psi \rangle$ in the von-Neumann algebra $\pi_\psi(\mathcal{O}_d)''$. Then the following holds:*

(a) *P is a sub-harmonic projection for the endomorphism $\Lambda(X) = \sum_k S_k X S_k^*$ on $\pi_\psi(\mathcal{O}_d)''$ i.e. $\Lambda(P) \geq P$ satisfying the following:*

(i) $PS_k^*P = S_k^*P$, $1 \leq k \leq d$;

(ii) *The set $\{S_I f : Pf = f, f \in \mathcal{H}_\psi, |I| < \infty\}$ is total in \mathcal{H}_ψ ;*

(iii) $\Lambda_n(P) \uparrow I$ as $n \uparrow \infty$;

(iv) $\sum_{1 \leq k \leq d} v_k v_k^* = I_{\mathcal{K}}$;

where $S_k = \pi_\psi(s_k)$ and $v_k = PS_kP$ for $1 \leq k \leq d$ are contractive operators on Hilbert subspace \mathcal{K} , the range of the projection P ;

(b) *For any $I = (i_1, i_2, \dots, i_k), J = (j_1, j_2, \dots, j_l)$ with $|I|, |J| < \infty$ we have $\psi(s_I s_J^*) = \langle \zeta_\psi, v_I v_J^* \zeta_\psi \rangle$ and the vectors $\{S_I f : f \in \mathcal{K}, |I| < \infty\}$ are total in \mathcal{H}_ψ ;*

(c) *The von-Neumann algebra $\mathcal{M} = P\pi_\psi(\mathcal{O}_d)''P$, acting on the Hilbert space \mathcal{K} i.e. range of P , is generated by $\{v_k, v_k^* : 1 \leq k \leq d\}''$ and the normal state $\phi(x) = \langle \zeta_\psi, x\zeta_\psi \rangle$ is faithful on the von-Neumann algebra \mathcal{M} .*

(d) *The following statements are equivalent:*

(i) ψ is a factor state of \mathcal{O}_d ;

(ii) \mathcal{M} is a factor;

PROOF. For a proof we refer to Proposition 2.1 in [26]. ■

Let ψ be a λ -invariant state of \mathcal{O}_d as in Proposition 2.2 and $H = \{z \in S^1 : \psi = \psi\beta_z\}$ be the closed subgroup of S^1 . Let $z \rightarrow U_z$ be the unitary representation of H in the GNS space $(\mathcal{H}_\psi, \pi, \zeta_\psi)$ associated with the state ψ of \mathcal{O}_d , defined by

$$(20) \quad U_z \pi_\psi(x) \zeta_\psi = \pi_\psi(\beta_z(x)) \zeta_\psi$$

so that $\pi_\psi(\beta_z(x)) = U_z \pi_\psi(x) U_z^*$ for $x \in \mathcal{O}_d$. We use same notations $(\beta_z : z \in H)$ for its normal extensions as group of automorphisms on $\pi_\psi(\mathcal{O}_d)''$. Furthermore, $\langle \zeta_\psi, P\beta_z(I - P)P\zeta_\psi \rangle = 0$ as $\psi = \psi\beta_z$ for $z \in H$. Since P is the support projection of ψ in $\pi_\psi(\mathcal{O}_d)''$, we have $P\beta_z(I - P)P = 0$ i.e. $\beta_z(P) \geq P$ for all $z \in H$. Since

H is a group, we conclude that $\beta_z(P) = P$. So $P \in \pi_\psi(\text{UHF}_d)''$ by Proposition 2.2 (b).

Since ϕ is a faithful state of \mathcal{M} , ζ_ϕ once identified with $\zeta_\psi \in \mathcal{K}$ is a cyclic and separating vector for \mathcal{M} and the closure of the closable operator $S_0 : a\zeta_\phi \rightarrow a^*\zeta_\phi$, $a \in \mathcal{M}$, S possesses a polar decomposition $S = \mathcal{J}\Delta^{1/2}$, where \mathcal{J} is an anti-unitary and Δ is a non-negative self-adjoint operator on \mathcal{K} . M. Tomita [8] theorem says that $\Delta^{it}\mathcal{M}\Delta^{-it} = \mathcal{M}$, $t \in \mathbb{R}$ and $\mathcal{J}\mathcal{M}\mathcal{J} = \mathcal{M}'$, where \mathcal{M}' is the commutant of \mathcal{M} . We define the modular automorphism group $\sigma = (\sigma_t, t \in \mathbb{T})$ on \mathcal{M} by

$$\sigma_t(a) = \Delta^{it}a\Delta^{-it}$$

which satisfies the modular relation

$$\phi(a\sigma_{-\frac{i}{2}}(b)) = \phi(\sigma_{\frac{i}{2}}(b)a)$$

for any two analytic elements a, b for the group of automorphisms (σ_t) . A more useful modular relation used frequently in this paper is given by

$$(21) \quad \phi(\sigma_{-\frac{i}{2}}(a^*)^*\sigma_{-\frac{i}{2}}(b^*)) = \phi(b^*a)$$

which shows that $\mathcal{J}a\zeta_\phi = \sigma_{-\frac{i}{2}}(a^*)\zeta_\phi$ for an analytic element a for the automorphism group (σ_t) . Anti unitary operator \mathcal{J} and the group of automorphism $\sigma = (\sigma_t, t \in \mathbb{R})$ are called *conjugate operator* and *modular automorphisms* associated with ϕ respectively.

The state $\phi(a) = \langle \zeta_\phi, x\zeta_\phi \rangle$ on \mathcal{M} being faithful and invariant of $\tau : \mathcal{M} \rightarrow \mathcal{M}$, we find a unique unital completely positive map $\tilde{\tau} : \mathcal{M}' \rightarrow \mathcal{M}'$ ([section 8 in [30]]) satisfying the duality relation

$$(22) \quad \langle b\zeta_\phi, \tau(a)\zeta_\phi \rangle = \langle \tilde{\tau}(b)\zeta_\phi, a\zeta_\phi \rangle$$

for all $a \in \mathcal{M}$ and $b \in \mathcal{M}'$. For a proof, we refer to section 8 in the monograph [30] or section 2 in [24].

Since $\tau(a) = \sum_{1 \leq k \leq d} v_k a v_k^*$, $x \in \mathcal{M}$ is an *inner map* i.e. each $v_k \in \mathcal{M}$, we have an explicit formula for $\tilde{\tau}$ as follows: For each $1 \leq k \leq d$, we set contractive operator

$$(23) \quad \tilde{v}_k = \overline{\mathcal{J}\sigma_{\frac{i}{2}}(v_k^*)\mathcal{J}} \in \mathcal{M}'$$

That \tilde{v}_k is indeed well defined as an element in \mathcal{M}' given in section 8 in [9]. By the modular relation (21), we have

$$(24) \quad \sum_k \tilde{v}_k \tilde{v}_k^* = I_{\mathcal{K}} \quad \text{and} \quad \tilde{\tau}(b) = \sum_k \tilde{v}_k b \tilde{v}_k^*, \quad b \in \mathcal{M}'$$

Moreover, if $\tilde{I} = (i_n, \dots, i_2, i_1)$ for $I = (i_1, i_2, \dots, i_n)$, we have

$$\begin{aligned} & \tilde{v}_{\tilde{I}}^* \zeta_\phi \\ &= \mathcal{J}\sigma_{\frac{i}{2}}(v_{\tilde{I}})^* \mathcal{J}\zeta_\phi \\ &= \mathcal{J}\Delta^{\frac{1}{2}} v_{\tilde{I}} \zeta_\phi \\ &= v_{\tilde{I}}^* \zeta_\phi \end{aligned}$$

and

$$(25) \quad \phi(v_I v_J^*) = \phi(\tilde{v}_{\tilde{I}} \tilde{v}_{\tilde{J}}^*), \quad |I|, |J| < \infty$$

We also set $\tilde{\mathcal{M}}$ to be the von-Neumann algebra generated by $\{\tilde{v}_k : 1 \leq k \leq d\}$. Thus $\tilde{\mathcal{M}} \subseteq \mathcal{M}'$.

Following [9] and [24], we consider the amalgamated tensor product $\mathcal{H} \otimes_{\mathcal{K}} \tilde{\mathcal{H}}$ of \mathcal{H} with $\tilde{\mathcal{H}}$ over the joint subspace \mathcal{K} . It is the completion of the quotient of the set

$$\mathbb{C}\bar{I} \otimes \mathbb{C}I \otimes \mathcal{K},$$

where \bar{I}, I both consisting of all finite sequences with elements in $\{1, 2, \dots, d\}$, by the equivalence relation defined by a semi-inner product defined on the set by requiring

$$\begin{aligned} \langle \bar{I} \otimes I \otimes f, \bar{I}\bar{J} \otimes IJ \otimes g \rangle &= \langle f, \tilde{v}_{\bar{J}} v_{Jg} \rangle, \\ \langle \bar{I}\bar{J} \otimes I \otimes f, \bar{I} \otimes IJ \otimes g \rangle &= \langle \tilde{v}_{\bar{J}} f, v_{Jg} \rangle \end{aligned}$$

and all inner product that are not of these form are zero. We also define two commuting representations (S_i) and (\tilde{S}_i) of \mathcal{O}_d on $\mathcal{H} \otimes_{\mathcal{K}} \tilde{\mathcal{H}}$ by the following prescription:

$$\begin{aligned} S_I \lambda(\bar{J} \otimes J \otimes f) &= \lambda(\bar{J} \otimes IJ \otimes f), \\ \tilde{S}_{\bar{I}} \lambda(\bar{J} \otimes J \otimes f) &= \lambda(\bar{J}\bar{I} \otimes J \otimes f), \end{aligned}$$

where λ is the quotient map from the index set to the Hilbert space. Note that the subspace generated by $\lambda(\emptyset \otimes I \otimes \mathcal{K})$ can be identified with \mathcal{H} and earlier S_I can be identified with the restriction of S_I , defined here. Same is valid for $\tilde{S}_{\bar{I}}$. The subspace \mathcal{K} is identified here with $\lambda(\emptyset \otimes \emptyset \otimes \mathcal{K})$. Thus \mathcal{K} is a cyclic subspace for the representation

$$\tilde{s}_j \otimes s_i \rightarrow \tilde{S}_j S_i$$

of $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ in the amalgamated Hilbert space. Let P be the projection on \mathcal{K} . Then we have

$$\begin{aligned} S_i^* P &= P S_i^* P = v_i^* \\ \tilde{S}_i^* P &= P \tilde{S}_i^* P = \tilde{v}_i^* \end{aligned}$$

for all $1 \leq i \leq d$.

We sum up required results in the following proposition.

PROPOSITION 2.4. *Let ψ be an element in K_ω and $(\mathcal{K}, v_k, 1 \leq k \leq d)$ be the elements in the support projection of ψ in $\pi_\psi(\mathcal{O}_d)''$ described in Proposition 2.3 and $(\mathcal{K}, \tilde{v}_k, 1 \leq k \leq d)$ be the dual elements and π be the amalgamated representation of $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$. Then the following holds:*

(a) *For any $1 \leq i, j \leq d$ and $|I|, |J| < \infty$ and $|\bar{I}|, |\bar{J}| < \infty$*

$$\langle \zeta_\psi, \tilde{S}_{\bar{I}} \tilde{S}_{\bar{J}}^* S_i S_I S_J^* S_j^* \zeta_\psi \rangle = \langle \zeta_\psi, \tilde{S}_i \tilde{S}_{\bar{I}} \tilde{S}_{\bar{J}}^* \tilde{S}_j^* S_I S_J^* \zeta_\psi \rangle;$$

(b) *The state $\psi : x \rightarrow \langle \zeta_\psi, x \zeta_\psi \rangle$ defined on $U\tilde{H}F_d \otimes UHF_d$ is equal to ω on \mathbb{M} , where we have identified*

$$\mathbb{M} \equiv \mathbb{M}_{(-\infty, 0]} \otimes \mathbb{M}_{[1, \infty)} \equiv U\tilde{H}F_d \otimes UHF_d;$$

with respect to an orthonormal basis $e = (e_i)$ of \mathbb{C}^d .

If ω is an ergodic state of \mathbb{M} and ψ is an extremal element in K_ω then

(c) $\pi(\mathcal{O}_d^H)'' = \pi(UHF_d)''$ and $\pi(\tilde{\mathcal{O}}_d^H)'' = \pi(UHF_d)''$;

(d) *The following statements are equivalent:*

(i) ω is a factor state of \mathcal{O}_d ;

(ii) $\pi(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d)'' = \mathcal{B}(\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H})$

(iii) $\mathcal{M} \vee \tilde{\mathcal{M}} = \mathcal{B}(\mathcal{K})$.

PROOF. For a proof we refer to Proposition 3.1. in [26]. ■

Let G be a compact group and $g \rightarrow u(g)$ be a d -dimensional unitary representation of G . By γ_g , we denote the product action of G on the infinite tensor product \mathbb{M} induced by $u(g)$,

$$\gamma_g(Q) = (\dots \otimes u(g) \otimes u(g) \otimes u(g) \dots) Q (\dots \otimes u(g)^* \otimes u(g)^* \otimes u(g)^* \dots)$$

for any $Q \in \mathbb{M}$. We recall now that the canonical action of the group $U_d(\mathbb{C})$ of $d \times d$ matrices on \mathcal{O}_d is given by

$$\beta_{u(g)}(s_j) = \sum_{1 \leq i \leq d} s_i u(g)_j^i$$

and thus

$$\beta_{u(g)}(s_j^*) = \sum_{1 \leq i \leq d} u(\bar{g})_j^i s_i^*$$

Note that $u(g)|e_i \rangle \langle e_j| u(g)^* = |u(g)e_i \rangle \langle u(g)e_j| = \sum_{k,l} u(g)_i^l u(\bar{g})_j^k |e_l \rangle \langle e_k|$, where e_1, \dots, e_d are the standard basis for \mathbb{C}^d . Identifying $|e_i \rangle \langle e_j|$ with $s_i s_j^*$, we verify that on \mathbb{M}_R the gauge action $\beta_{u(g)}$ of the Cuntz algebra \mathcal{O}_d and γ_g coincide i.e. $\gamma_g(Q) = \beta_{u(g)}(Q)$ for all $Q \in \mathbb{M}_R$.

PROPOSITION 2.5. *Let ω be a translation invariant ergodic state on \mathbb{M} . Suppose that ω is G -invariant,*

$$\omega(\gamma_g(Q)) = \omega(Q) \text{ for all } g \in G \text{ and any } Q \in \mathbb{M}.$$

Let ψ be an extremal point in K_ω and $(\mathcal{K}, \mathcal{M}, v_k, 1 \leq k \leq d, \phi)$ be the elements associated with $(\mathcal{H}, S_i = \pi(s_i), \zeta_\psi)$, described as in Proposition 2.3. Then we have the following:

(a) *There exists a unitary representation $g \rightarrow \hat{U}(g)$ in $\mathcal{B}(\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H})$ and a representation $g \rightarrow \zeta(g) \in S^1$ so that*

$$(26) \quad \hat{U}(g) S_i \hat{U}(g)^* = \zeta(g) \beta_{u(g)}(S_i), \quad 1 \leq i \leq d$$

and

$$(27) \quad \hat{U}(g) \tilde{S}_i \hat{U}(g)^* = \zeta(g) \beta_{u(g)}(\tilde{S}_i), \quad 1 \leq i \leq d$$

for all $g \in G$.

(b) *There exists a unitary representation $g \rightarrow \hat{u}(g)$ in $\mathcal{B}(\mathcal{K})$ so that $\hat{u}(g) \mathcal{M} \hat{u}(g)^* = \mathcal{M}$ for all $g \in G$ and $\phi(\hat{u}(g)x\hat{u}(g)^*) = \phi(x)$ for all $x \in \mathcal{M}$. Furthermore, the operator $V^* = (v_1^*, \dots, v_d^*)^{tr} : \mathcal{K} \rightarrow \mathbb{C}^d \otimes \mathcal{K}$ is an isometry which intertwines the representation of G ,*

$$(28) \quad (\zeta(g) \hat{u}(g) \otimes u(g)) V^* = V^* \hat{u}(g)$$

for all $g \in G$, where $g \rightarrow \zeta(g)$ is the representation of G in $U(1)$.

(c) $\mathcal{J} \hat{u}(g) \mathcal{J} = \hat{u}(g)$ and $\Delta^{it} \hat{u}(g) \Delta^{-it} = \hat{u}(g)$ for all $g \in G$ and $t \in \mathbb{R}$.

(d) $u_z \hat{u}(g) = \hat{u}(g) u_z$ for all $g \in G$ and $z \in H$.

(e) *If G is simply connected then $\zeta(g) = 1$ for all $g \in G$ and $\psi = \psi \beta_{u(g)}$ and $\psi_0 = \psi_0 \beta_{u(g)}$ for all $g \in G$.*

PROOF. Proof is given in Proposition 2.7 in [26], where we used factor property of ψ but same holds good if ω is an ergodic state of \mathbb{M} once we use Proposition 2.1 (a) instead of factor property of ω . ■

For a given $u \in U_d(\mathbb{C})$, we also extend the map $\mathcal{J}_u : \mathbb{M} \rightarrow \mathbb{M}$ defined in (10) to an anti-linear automorphism on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$, defined by

$$(29) \quad \mathcal{J}_u(\tilde{s}_I \tilde{s}_{J'}^* \otimes s_I s_{J'}^*) = \beta_{\bar{u}}(\tilde{s}_I \tilde{s}_J \otimes s_I s_{J'}^*)$$

for all $|I|, |J|, |I'|, |J'| < \infty$ and then extend anti-linearly for an arbitrary element of $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$. So we have

$$(30) \quad \mathcal{J}_u = \beta_{\bar{u}} \mathcal{J}_{I_d} = \mathcal{J}_{I_d} \beta_u$$

So these maps are defined after fixing the orthonormal basis $e = (e_i)$, which have identified $\text{UHF} \otimes \text{UHF}_d$ with $\mathbb{M}_L \otimes \mathbb{M}_R = \mathbb{M}$ as in Proposition 2.4 (b), where the monomial given (17) is identified with the matrix given in (18).

We make few simple observations in the following for $u, w \in U_d(\mathbb{C})$:

$$(31) \quad \begin{aligned} & \mathcal{J}_u \mathcal{J}_w \\ &= \beta_{\bar{u}} \mathcal{J}_{I_d} \mathcal{J}_{I_d} \beta_w \\ &= \beta_{\bar{u}w}, \end{aligned}$$

and

$$(32) \quad \begin{aligned} & \mathcal{J}_w \beta_u \\ &= \mathcal{J}_{I_d} \beta_w \beta_u \\ &= \mathcal{J}_{I_d} \beta_{wu} \\ &= \mathcal{J}_{wu} \end{aligned}$$

Also

$$(33) \quad \begin{aligned} & \beta_u \mathcal{J}_w \\ &= \beta_u \mathcal{J}_{I_d} \beta_w \\ &= \mathcal{J}_{I_d} \beta_{\bar{u}} \beta_w \\ &= \mathcal{J}_{I_d} \beta_{\bar{u}w} \\ &= \mathcal{J}_{\bar{u}w} \end{aligned}$$

i.e. \mathcal{J}_w commutes with β_u if $wuw^* = \bar{u}$.

By combining relations (32)-(33), we have the following identities

$$(34) \quad \begin{aligned} & \mathcal{J}_w \beta_u \\ &= \mathcal{J}_{wu} \text{ by (52)} \\ &= \mathcal{J}_{\bar{u}w} \text{ (provided } wuw^* = \bar{u}) \\ &= \beta_u \mathcal{J}_w \text{ by (33)} \end{aligned}$$

Let G be the simply connected Lie group $SU_2(\mathbb{C})$ and $g \rightarrow u(g)$ be a d -dimensional unitary irreducible representation of G . Then there exists a $r \in U_d(\mathbb{C})$ such that

$$(35) \quad ru(g)r^* = u(\bar{g})$$

for all $g \in G$. The element $r \in U_d(\mathbb{C})$ is determined uniquely modulo a phase factor in S^1 . In particular any element

$$(36) \quad r_z = zr_0, \quad z \in S^1$$

satisfies (35), where we have fixed a r_0 satisfying (35) with additional condition

$$(37) \quad r_0^2 = I_d$$

In our notions $r_1 = r_0$. In such a case, $r_{-1} = -r_0$ is the only other choice that satisfies (35) and (37) instead of r_0 .

With such a choice for r_0 , for all $g \in SU_2(\mathbb{C})$ we have

$$r_0 u(g) r_0 = u(\bar{g})$$

Taking conjugation on both sides, we have $\bar{r}_0 u(\bar{g}) \bar{r}_0 = u(g)$ i.e.

$$\bar{r}_0^* u(g) \bar{r}_0^* = u(\bar{g})$$

Since $\bar{r}_0^2 = I$, we arrive at

$$\bar{r}_0 u(g) \bar{r}_0 = u(\bar{g})$$

for all $g \in G$. So by the irreducible property of the representation $g \rightarrow u(g)$, we conclude that $\bar{r}_0 = \mu r_0$, where $\mu^2 = 1$ since $r_0^2 = \bar{r}_0^2 = 1$ i.e. μ is either 1 or -1 .

Taking determinants of matrices on both sides of $\bar{r}_0 r_0 = \mu I_d$, we get $\mu^d = \det(r_0) \det(\bar{r}_0) = |\det(r_0)|^2 = 1$. This shows that $\mu = 1$ if d is an odd integer. For even values of d , we make a direct calculation to show $\mu = -1$ as follows:

For $d = 2$, let σ_x, σ_y and σ_z be the Pauli matrices in $M_2(\mathbb{C})$ i.e. the standard (irreducible) representation of Lie algebra $su_2(\mathbb{C})$ in \mathbb{C}^2 :

$$\sigma_x = \begin{pmatrix} 0 & , & 1 \\ 1 & , & 0 \end{pmatrix},$$

$$\sigma_y = \begin{pmatrix} 0 & , & i \\ -i & , & 0 \end{pmatrix},$$

$$\sigma_z = \begin{pmatrix} 1 & , & 0 \\ 0 & , & -1 \end{pmatrix}.$$

A direct commutation shows that r_0 is given by

$$r_0 = \begin{pmatrix} 0 & , & i \\ -i & , & 0 \end{pmatrix}$$

The self-adjoint matrix σ_y is also a unitary i.e. $\sigma_y^2 = I_2$ and

$$\sigma_y \sigma_x \sigma_y = -\sigma_x$$

and

$$\sigma_y \sigma_z \sigma_y = -\sigma_z$$

Since $\sigma_x = \bar{\sigma}_x$ and $\sigma_z = \bar{\sigma}_z$, σ_y inter-twins $e^{it\sigma_x}$ and $e^{it\sigma_z}$ with their conjugate matrices $e^{-it\sigma_x}$ and $e^{-it\sigma_z}$ respectively for all $t \in \mathbb{R}$. In contrast, since $\bar{\sigma}_y = -\sigma_y$, we also get σ_y inter-twins $e^{it\sigma_y}$ with $e^{-it\sigma_y}$ for all $t \in \mathbb{R}$. So we set $r_0 = \sigma_y$ (other choice we can make for r_0 is $-\sigma_y$) and verify directly that $\bar{r}_0 = -r_0$ i.e. $\mu = -1$ if $d = 2$.

We write $i\sigma_y = e^{it_0\sigma_y} \in SU_2(\mathbb{C})$, where $t_0 = \frac{\pi}{2}$ and verify that

$$u(e^{it_0\sigma_y})u(g)u(e^{-it_0\sigma_y})$$

$$\begin{aligned} &= u(i\sigma_y)u(g)u(i\sigma_y)^* \\ &= u((i\sigma_y)g(i\sigma_y)^*) \\ &= u(\bar{g}) \end{aligned}$$

Since $su_2(\mathbb{C})$ is a real Lie algebra that has unique Lie algebra extension to a complex Lie algebra $sl_2(\mathbb{C})$, i.e. Lie algebra over the field of complex numbers, we also have

$$u(\bar{g}) = u(g)$$

for all $g \in SU_2(\mathbb{C})$ (Lie-derivatives of the representations in both sides are equal as element in $sl_2(\mathbb{C})$). So we have

$$(38) \quad u(e^{it_0\sigma_y})u(g)u(e^{-it_0\sigma_y}) = \overline{u(g)}$$

If π_u is the associated Lie-representation of $su_2(\mathbb{C})$, we have

$$u(e^{it_0\sigma_y}) = e^{it_0\pi_u(\sigma_y)}$$

for even integer values of d , whereas

$$u(e^{it_0\sigma_y}) = e^{2it_0\pi_u(\sigma_y)}$$

for odd integer values of d . Thus for an arbitrary even values of d , the unitary matrix $r_0 = e^{it_0\pi_u(\sigma_y)}$ satisfies (35) and (37). In contrast, for an arbitrary odd values of d , the unitary matrix $r_0 = e^{i2t_0\pi_u(\sigma_y)}$ satisfies (35) and (37). In short, $\mu = 1$ if d is an odd integer and -1 if d is an even integer.

We write $\mu = \zeta^2$ and set $r_0 \in U_d(\mathbb{C})$, such that

$$\zeta r_0 = u(e^{it_0\sigma_y}) \in U_d(\mathbb{C}),$$

where $\zeta^2 = \mu$ and so μ is 1 for odd values of d otherwise -1 . In the last section, we will recall standard explicit description of r_0 and $g \rightarrow u(g)$ that satisfies (35) and (37). Note also that $r_\zeta = \zeta r_0$ is a matrix with real entries irrespective of values taken for d .

Now we go back to our main text. So we have

$$(39) \quad \mathcal{J}_{r_z}^2(x) = \beta_{\bar{r}_z r_z}(x) = \beta_{\mu I_d}(x)$$

for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$, where $\mu = 1$ or -1 depending on d odd or even. In any case, by (34) and (35), we also have

$$(40) \quad \mathcal{J}_{r_z} \beta_{u(g)} = \beta_{u(g)} \mathcal{J}_{r_z}$$

for all $g \in SU_2(\mathbb{C})$.

Let ω be a translation invariant factor state of \mathbb{M} and ψ be an extremal element in K_ω . We define a state $\psi_0 : \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d \rightarrow \mathbb{C}$ by extending both $\tilde{\psi} : \tilde{\mathcal{O}}_d \rightarrow \mathbb{C}$ and $\psi : \mathcal{O}_d \rightarrow \mathbb{C}$ by

$$(41) \quad \psi_0(\tilde{s}_{I'} \tilde{s}_{J'}^* \otimes s_I s_J^*) = \langle \zeta_\psi, \tilde{v}_{I'} \tilde{v}_{J'}^*, v_I^* v_J^* \zeta_\psi \rangle$$

for all $|I'|, |J'|, |I|$ and $|J| < \infty$. Proposition 2.4 says that $(\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H}, \pi, \zeta_\psi)$ is the GNS representation $(\mathcal{H}_{\psi_0}, \pi_{\psi_0}, \zeta_{\psi_0})$ of $(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d, \psi_0)$.

PROPOSITION 2.6. *Let ω be a translation and $SU_2(\mathbb{C})$ invariant ergodic state of \mathbb{M} . Then the following statement holds for any extremal element $\psi \in K_\omega$:*

(a) $\psi \beta_{u(g)} = \psi$ on \mathcal{O}_d ;

(b) $\psi_0 \beta_{u(g)} \otimes \beta_{u(g)} = \psi_0$ on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ for all $g \in SU_2(\mathbb{C})$;

(c) $\psi_0 \beta_{r_\zeta} \otimes \beta_{r_\zeta} = \psi_0$ on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$, where $r_\zeta = \zeta r_0 \in u(SU_2(\mathbb{C}))$;

(d) $\psi_0 \beta_{r_0} \otimes \beta_{r_0} = \psi_0$ on $U\tilde{H}F_d \otimes UHF_d$.

PROOF. For (a),(b) and (c) we refer to Proposition 3.1 in [26], where we used factor property of ψ but same holds good for ergodic state as well once we use Proposition 2.1 (a). The last statement is a simple consequence of (c) since $\beta_{r_\zeta} = \beta_{r_0}$ on UHF_d and $U\tilde{H}F_d$ as $r_\zeta = \zeta r_0$, where $\zeta^2 = \mu$ is either 1 or -1 . ■

3. Real, lattice symmetric, $SU_2(\mathbb{C})$ and translation invariant ergodic states

We quickly recall from [25] the following definitions. Given a λ -invariant state of \mathcal{O}_d , we define its lattice symmetric state $\tilde{\psi} : \mathcal{O}_d \rightarrow \mathbb{C}$ by

$$\tilde{\psi}(s_I s_J^*) = \psi(s_{\bar{I}} s_{\bar{J}})$$

for all $|I|, |J| < \infty$ and extend linearly. Also we set state $\bar{\psi} : \mathcal{O}_d \rightarrow \mathbb{C}$ defined by

$$\bar{\psi}(s_I s_J^*) = \psi(s_J s_I)$$

for all $|I|, |J| < \infty$ and extend linearly.

PROPOSITION 3.1. *Let ω be a real, lattice symmetric translation invariant state of \mathbb{M} and extremal in the convex set of translation invariant states of \mathbb{M} . Then there exists an extremal element ψ in K_ω , so that the associated amalgamated representation $\pi_{\psi_0} : \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d \rightarrow \mathcal{B}(\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H})$ determines an anti-automorphism \mathcal{J}_{r_ζ} that takes $\pi_{\psi_0}(\tilde{s}_{I'} \tilde{s}_{J'} s_I s_J^*)$ to $\psi_{\psi_0}(\beta_{r_\zeta}(\tilde{s}_I \tilde{s}_J^* s_I s_J^*))$ extending anti-linearly such that*

$$(42) \quad \mathcal{J}_{r_\zeta}(X) = \mathcal{J}_\zeta X \mathcal{J}_\zeta^*$$

where \mathcal{J}_ζ is an anti-unitary operator on $\mathcal{H} \otimes_{\mathcal{K}} \tilde{\mathcal{H}}$ that takes

$$\pi(s_I s_J^* \tilde{s}_{I'} \tilde{s}_{J'}^*) \zeta_\psi \rightarrow \beta_\zeta(\pi(s_{I'} s_{J'}^* \tilde{s}_I \tilde{s}_J^*)) \zeta_\psi$$

for all $|I|, |J|, |I'|, |J'| < \infty$ and then extending anti-linearly on their linear span. Furthermore, we have

$$\psi_0(\mathcal{J}_{r_\zeta}(X)) = \psi_0(X^*)$$

for all $X \in \pi_{\psi_0}(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d)''$.

Furthermore, if ω is also pure then the Popescu elements $(\mathcal{K}, \mathcal{M}, v_k, 1 \leq k \leq d, \phi)$ of ψ as given in Proposition 2.4 satisfies the following:

(a) There exists a unique unitary operator γ on \mathcal{K} such that $\gamma \zeta_\psi = \zeta_\psi$ and

$$(43) \quad \gamma \left(\sum c_{I', J', I, J} \tilde{v}_{I'} \tilde{v}_{J'}^* v_I v_J^* \right) \gamma^* = \sum c_{I', J', I, J} \mathcal{J} \tilde{v}_I \tilde{v}_J v_{I'} v_{J'}^* \mathcal{J}$$

for all $I', |J'|, |I|$ and $|J| < \infty$, where γ is also self adjoint, commuting with modular elements $\Delta^{\frac{1}{2}}, \mathcal{J}$.

(b) $\gamma u_z = u_{\bar{z}} \gamma$ for all $z \in H$.

(c) The anti-unitary map $\mathcal{J} \gamma : \mathcal{K} \rightarrow \mathcal{K}$ extends to anti-unitary map \mathcal{J}_γ on $\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H}$ so that $\mathcal{J}_\gamma^2 = I$ and

$$\pi(\mathcal{J}_{I_d}(x)) = \mathcal{J}_\gamma \pi(x) \mathcal{J}_\gamma$$

for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$.

Let ω, ψ be as in Proposition 3.1 and ω be also $SU_2(\mathbb{C})$ invariant. By Proposition 2.6, we have $\psi_0\beta_{u(g)} = \psi_0$ for all $g \in SU_2(\mathbb{C})$ on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$. Since $r_\zeta = u(i\sigma_y)$, there exists a unitary operator $\hat{r}_\zeta : \tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H} \rightarrow \tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H}$ such that $\hat{r}_\zeta\zeta_\psi = \zeta_\psi$ and

$$Ad_{\hat{r}_\zeta}(\pi(x)) = \pi(\beta_{r_\zeta}(x))$$

for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$.

We recall for each $z \in H$,

$$Ad_{U_z}(\pi(x)) = \pi(\beta_z(x))$$

for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$.

We consider the anti-automorphism $\hat{\mathcal{J}}_{\gamma_{r_\zeta}}$ on $\mathcal{B}(\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H})$, defined by

$$\hat{\mathcal{J}}_{\gamma_{r_\zeta}}(X) = \mathcal{J}_\gamma \hat{r}_\zeta X \hat{r}_\zeta^* \mathcal{J}_\gamma$$

and verify the following

$$\begin{aligned} & \hat{\mathcal{J}}_{\gamma_{r_\zeta}}(\pi(\tilde{s}_{I'} \tilde{s}_{J'} s_I s_J^*)) \\ &= \mathcal{J}_\gamma \pi(\beta_{r_\zeta}(\tilde{s}_{I'} \tilde{s}_{J'} s_I s_J^*)) \mathcal{J}_\gamma \\ &= \beta_{\bar{r}_\zeta}(\mathcal{J}_\gamma \pi(\tilde{s}_{I'} \tilde{s}_{J'} s_I s_J^*) \mathcal{J}_\gamma^*) \\ &= \beta_{\bar{r}_\zeta}(\pi(\mathcal{J}_{I_d}(\tilde{s}_{I'} \tilde{s}_{J'} s_I s_J^*))) \\ &= \pi(\mathcal{J}_{r_\zeta}(\tilde{s}_{I'} \tilde{s}_{J'} s_I s_J^*)) \end{aligned}$$

PROPOSITION 3.2. *Let ω, ψ be as in Proposition 3.1 and ω be also $SU_2(\mathbb{C})$ invariant. Then the anti-automorphism \mathcal{J}_{r_ζ} on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ induces a well defined anti-automorphism map $\hat{\mathcal{J}}_{r_\zeta}$ on $\pi(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d)''$ by*

$$(44) \quad \hat{\mathcal{J}}_{r_\zeta}(\pi(x)) = \pi(\mathcal{J}_{r_\zeta}(x))$$

for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ and

$$\hat{\mathcal{J}}_{r_\zeta} = \hat{\mathcal{J}}_{\gamma_{r_\zeta}}$$

i.e.

$$(45) \quad \hat{\mathcal{J}}_{r_\zeta}(X) = \mathcal{J}_\gamma \hat{r}_\zeta X \hat{r}_\zeta^* \mathcal{J}_\gamma$$

for all $X \in \pi(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d)''$.

Furthermore, if ω is also pure then $\hat{\mathcal{J}}_{r_\zeta}(P) = P$ and the corner anti-automorphism, defined by

$$\hat{\mathcal{J}}_{r_\zeta}(a) = P \hat{\mathcal{J}}_{r_\zeta}(PaP)P$$

for all $a \in \mathcal{B}(\mathcal{K})$ satisfies the following:

$$(46) \quad \hat{\mathcal{J}}_{r_\zeta}(a) = \mathcal{J}_\gamma \hat{r}_\zeta a \hat{r}_\zeta^* \gamma^* \mathcal{J}$$

for all $a \in \mathcal{B}(\mathcal{K})$. Furthermore, we have the following consequences:

$$(a1) \quad \hat{\mathcal{J}}_{r_\zeta}^2 = \beta_\mu;$$

$$(a2) \quad \beta_{\bar{r}_\zeta}(\tilde{S}_I \tilde{S}_J^* S_{I'} S_{J'}^*) \mathcal{J}_\gamma \hat{r}_\zeta = \mathcal{J}_\gamma \hat{r}_\zeta S_I S_{J'}^* \tilde{S}_I \tilde{S}_J^*, \text{ for all } |I'|, |J'|, |I| \text{ and } |J| < \infty.$$

$$(a3) \quad Ad_{\hat{U}(g)} \hat{\mathcal{J}}_{r_\zeta} = \hat{\mathcal{J}}_{r_\zeta} Ad_{\hat{U}(g)} \text{ for all } g \in SU_2(\mathbb{C});$$

$$(b1) \quad Ad_{\gamma_{r_\zeta}}^2 = \beta_{\zeta^2 I_d}, \text{ where } \gamma_{r_\zeta} = \gamma \hat{r}_\zeta \text{ commutes with modular elements } \mathcal{J} \text{ and } \Delta^{\frac{1}{2}};$$

$$(b2) \quad \beta_{\bar{r}_\zeta}(\tilde{v}_I \tilde{v}_J^* v_{I'} v_{J'}^*) \mathcal{J}_\gamma \hat{r}_\zeta = \mathcal{J}_\gamma \hat{r}_\zeta v_I v_J^* \tilde{v}_{I'} \tilde{v}_{J'}^*, \text{ for all } |I'|, |J'|, |I| \text{ and } |J| < \infty;$$

(b3) γ_{r_ζ} commutes the representation $\{\hat{u}(g) : g \in SU_2(\mathbb{C})\}$;

There exists a unique unitary operator Γ_{r_ζ} and an anti-unitary operator extending \mathcal{J} on $\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H}$ extending unitary $\gamma_{r_\zeta} : \mathcal{K} \rightarrow \mathcal{K}$ and anti-unitary operator $\mathcal{J} : \mathcal{K} \rightarrow \mathcal{K}$ respectively such that

(c1) $Ad_{\Gamma_{r_\zeta}}^2 = \beta_{\zeta^2}$; $Ad_{\Gamma_{r_\zeta}}$ acts on $\pi(\mathcal{O}_d)''$ and $\pi(UHF_d)''$ ($\pi(\tilde{\mathcal{O}}_d)''$ and $\pi(U\tilde{H}F_d)''$) respectively;

(c2) $\beta_{r_\zeta}(\tilde{S}_I \tilde{S}_J^* S_{I'} S_{J'}^*) \mathcal{J} \Gamma_{r_\zeta} = \mathcal{J} \Gamma_{r_\zeta} S_I S_J^* \tilde{S}_{I'} \tilde{S}_{J'}^*$ for all $|I'|, |J'|, |I|$ and $|J| < \infty$;

(c3) Γ_{r_ζ} and \mathcal{J} commutes the representation $\{\hat{U}(g) : g \in SU_2(\mathbb{C})\}$;

PROPOSITION 3.3. *Let ω and ψ be as in Proposition 3.2 and ω is also reflection positive with twist r_0 . Then we have $Ad_{\gamma_{r_\zeta}}(a) = a$ for all $a \in \mathcal{M}_0$, where $\mathcal{M}_0 = \{a \in \mathcal{M} : \beta_z(a) = a \forall z \in H\} = P\pi_\psi(UHF_d)''P$.*

PROOF. We recall that $r_\zeta = \zeta r_0$ and thus Theorem 3.5 (d) in [25] gives a proof as $Ad_{\gamma_{r_\zeta}} = Ad_{\gamma_{r_0}}$ on \mathcal{M}_0 , where Ad_{γ_0} is defined in Theorem 3.5 in [23] using invariance property $\psi_0 \beta_{r_0} \otimes \beta_{r_0} = \psi_0$ on \mathbb{M} which is identified as before with $U\tilde{H}F_d \otimes UHF_d$. Alternatively, we can use the same argument directly with Ad_{r_ζ} using our hypothesis that ω is reflection positive with the twist r_0 to conclude Ad_{r_ζ} on $\pi(UHF_d)''$ is I . \blacksquare

In general ergodic states of (\mathbb{M}, θ) need not be factor states of \mathbb{M} [9] though ergodic states of (\mathcal{O}_d, λ) are factor states of \mathcal{O}_d . One of the central question that arises while dealing with ground states of H , whether additional symmetries of H make extremal decomposition of its ground state ω in the convex set of translation and $SU_2(\mathbb{C})$ invariant states to be a factor decomposition.

PROPOSITION 3.4. *Let ω be an extreme point in the convex set of translation and G -invariant states of \mathbb{M} and $K_{\omega, G}$ be the non-empty compact convex set of λ and G -invariant states ψ so that $\psi|_{UHF_d} = \omega|_{\mathbb{M}_R}$ as in Proposition 2.1. Then the following statements are true:*

(a) *The set $K_{\omega, G}$ is a face in the convex set of translation and G invariant states of \mathcal{O}_d ;*

(b) *For an extremal element $\psi \in K_{\omega, G}$, let $\psi = \int^\oplus \psi'_\alpha d\mu(\alpha)$ be an ergodic decomposition in the convex set of λ -invariant states of \mathcal{O}_d and $\psi_\alpha = \int_G \psi'_\alpha \beta_{u(g)} dg$, where dg is the normalised Haar measure on G . Then μ -almost every where, $\psi = \psi_\alpha$.*

(c) *For two elements g_1 and g_2 , either $\psi'_\alpha \beta_{g_1}$ and $\psi'_\alpha \beta_{g_2}$ are same or orthogonal and*

$$\pi_\psi(x) = \int_{G/G_\alpha}^\oplus \pi_{\psi'_\alpha \beta_{u(g')}}(x) dg'$$

for all $x \in \mathcal{O}_d$, where $G_\alpha = \{g \in G : \psi'_\alpha \beta_{u(g)} = \psi'_\alpha\}$ and dg' is the induced measure on the cosets of G_α . Moreover, $L^\infty(G/G_\alpha, dg') \otimes I \subset \pi_\psi(\mathcal{O}_d)''$.

(d) *If ψ_1 and ψ_2 are two extreme points in $K_{\omega, G}$ then $\psi_2 = \psi_1 \beta_z$ for some $z \in S^1$; The closed set $H = \{z \in S^1 : \psi \beta_z = \psi\}$ is independent of the extreme point $\psi \in K_{\omega, G}$ we choose;*

(e) For an extremal element $\psi \in K_{\omega, G}$, we have $\pi_\psi(\mathcal{O}_d^H)'' = \pi_\psi(\text{UHF}_d)''$;

PROOF. For any two G and λ -invariant states ψ and ψ' and $\lambda \in (0, 1)$, if the state $\psi_\lambda = \lambda\psi_1 + (1-\lambda)\psi_0$ is in $K_{\omega, G}$ then its restriction to UHF_d being an extremal element in the convex set of G and λ invariant states, $\psi_1 = \psi_2 = \omega$ on UHF_d . So ψ_1 and ψ_1 are elements in $K_{\omega, G}$. This proves (a).

The statement (b) follows as $\psi_\alpha \in K_{\omega, G}$,

$$\begin{aligned} & \int \psi_\alpha d\mu_\alpha \\ &= \int \psi'_\alpha \beta_{u(g)} dg d\mu(\alpha) \\ &= \int \psi'_\alpha d\mu(\alpha) \beta_{u(g)} dg \\ &= \int \psi \beta_{u(g)} dg \\ &= \psi \end{aligned}$$

and ψ is an extremal element in $K_{\omega, G}$.

For (c), we use Theorem 4.3.19 in [7] since $\psi' \beta_g$ commutes with λ and centrally ergodic, being an extremal element in the convex set of λ -invariant states. For the last statement $L^\infty(G/G_\alpha, dg') \otimes I \subset \pi_\psi(\mathcal{O}_d)''$, ψ' being an extremal element in the λ invariant states of \mathcal{O}_d , we have

$$\begin{aligned} & \frac{1}{n} \sum_{0 \leq k \leq n-1} \pi_\psi(\lambda^k(x))(g') \\ &= \frac{1}{n} \sum_{0 \leq k \leq n-1} \pi_{\psi'_\alpha}(\lambda^k(\beta_{u(g')}(x)) \rightarrow \psi'_\alpha(\beta_{u(g')}(x)) \end{aligned}$$

So $g' \rightarrow \psi'_\alpha(\beta_{u(g')}(x))$ is in $\pi_\psi(\mathcal{O}_d)''$. Since the collection of functions separates points in G/G_α , we conclude that $L^\infty(G/G_\alpha, dg') \otimes I_{\alpha'} \subset \pi_\psi(\mathcal{O}_d)''$. This shows that ergodic decomposition is a central decomposition.

For (d), we use ergodic decomposition given in (c) for ψ_1 and ψ_2 in the convex set of λ -invariant states and use Lemma 7.4 in [9] to conclude $\psi'_1(\alpha) = \psi'_2(\alpha)\beta_z$ for some $z \in S^1$. Since β_z commutes with $\{\beta_{u(g)} : g \in SU_2(\mathbb{C})\}$, we get $\psi_1 = \psi_2\beta_z$ as well by (b).

Now proof for (e), follows along the same line of Lemma 6.8 in [9], once we use ergodic decomposition given in (b) and (c) since ζ_z commutes with $\{\beta_{u(g)} : g \in SU_2(\mathbb{C})\}$. ■

In Proposition 3.4, we have used the commuting property of $\{\beta_z : z \in S^1\}$ with the group action $SU_2(\mathbb{C})$ and λ on \mathcal{O}_d . Our next proposition, in contrast having two additional \mathbb{Z}_2 actions on \mathcal{O}_d namely $\psi \rightarrow \tilde{\psi}$ and $\bar{\psi}$ of which one of them $\psi \rightarrow \bar{\psi}$ does not commute with $\{\beta_z : z \in S^1\}$ action on \mathcal{O}_d . Nevertheless, next proposition exploits commuting property of the action $\psi \rightarrow \tilde{\psi}\beta_{r_\zeta}$ with $\{\beta_{u(g)} : g \in SU_2(\mathbb{C})\}$ on \mathcal{O}_d .

PROPOSITION 3.5. *Let ω be an extremal element in the convex set of real, lattice reflection symmetric, translation and $SU_2(\mathbb{C})$ -invariant states of \mathbb{M} . Let K_{ω, G_r} be the non-empty convex set of λ and $SU_2(\mathbb{C})$ invariant states of \mathcal{O}_d satisfying $\tilde{\psi} = \bar{\psi} = \psi$. Let ψ be an extremal element in K_{ω, G_r} and $\psi = \int \psi'_\alpha d\mu(\alpha)$ be an ergodic decomposition of ψ in the convex set of λ -invariant states of \mathcal{O}_d with $\psi = \bar{\psi} = \tilde{\psi}$. Then there exists anti-automorphism \mathcal{J}_{r_ζ} on $\pi_{\psi_0}(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d)''$ that takes*

$$\pi_{\psi_0}(\tilde{s}_I, \tilde{s}_J^*, s_I s_J^*) \rightarrow \pi_{\psi_0}(\beta_{r_\zeta}(\tilde{s}_I \tilde{s}_J^* s_I s_J^*))$$

by extending anti-linearly in their linear span. Moreover, the following statements are true:

(a) *There exists an anti-automorphism $\mathcal{J}_{r_\zeta}^\alpha$ on $\pi_{\psi'_\alpha}(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d)''$ satisfying*

$$\mathcal{J}_{r_\zeta} = \int_{G/G_\alpha}^\oplus \mathcal{J}_{r_\zeta}^\alpha \beta_{u(g')} dg'$$

if and only if $\psi'_{\alpha,0} \beta_{\zeta^2 I_d} = \psi'_{\alpha,0}$ on \mathcal{O}_d . In such a case, $\psi'_{\alpha,0} \mathcal{J}_{r_\zeta}(x^*) = \psi'_{\alpha,0}(x)$ for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$.

(b) *There exists an anti-automorphism $\mathcal{J}_{r_\zeta}^\alpha$ on $\pi_{\psi'_\alpha}(U\tilde{H}F_d \otimes UHF_d)''$ satisfying*

$$\mathcal{J}_{r_\zeta} = \int_{G/G_\alpha}^\oplus \mathcal{J}_{r_\zeta}^\alpha \beta_{u(g')} dg'$$

and $\psi'_\alpha(\mathcal{J}_{r_\zeta}(x^*)) = \psi'_\alpha(x)$ for all $x \in UHF_d$.

(c) *If $\psi'_\alpha \beta_{\zeta^2 I_d} = \psi'_\alpha$ on \mathcal{O}_d then $\psi'_\alpha \beta_{u(g)} = \psi'_\alpha$ on \mathcal{O}_d for all $g \in SU_2(\mathbb{C})$; In such a case $\psi = \psi'_\alpha$ μ -almost everywhere, is an ergodic state in the convex set of λ -invariant states of \mathcal{O}_d satisfying $\psi = \bar{\psi} = \tilde{\psi}$.*

(d) *The state ω is as well an extremal element in the convex set of real, lattice reflection symmetric and translation invariant states of \mathbb{M} .*

PROOF. It is routine to check that similar statements of Proposition 3.4 (a),(b) and (c) are valid for any extremal element $\psi \in K_{\omega, G_r}$ as well. That we can choose an extremal element $\psi \in K_{\omega, G_r}$ with $\psi = \psi_{\beta_{r_\zeta}}$ follows trivially as $r_\zeta \in u(SU_2(\mathbb{C}))$ and ψ is $SU_2(\mathbb{C})$ invariant. Thus along the same line of the proof of Proposition 4.2 in [25], we ensure an anti-automorphism \mathcal{J}_{r_ζ} satisfying (42).

In particular, anti-automorphism \mathcal{J}_{r_ζ} also preserves the centre \mathcal{Z}_{ψ_0} of $\pi_{\psi_0}(\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d)''$. By the last Proposition 3.4 (c), \mathcal{Z}_{ψ_0} is isomorphic to $L^\infty(SU_2(\mathbb{C})/G_\alpha, dg')$ and so by the uniqueness of factor decomposition of ψ_0 , with some $h_\zeta \in SU_2(\mathbb{C})/G_\alpha$ we have

$$\psi'_{\alpha,0}(\mathcal{J}_{r_\zeta}(x^*)) = \psi'_{\alpha,0}(\beta_{u(h_\zeta)}(x))$$

for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ as

$$\begin{aligned} & \int_{G/G_\alpha} \psi'_{\alpha}(\mathcal{J}_{r_\zeta}(\beta_{u(g')}(x^*))) dg' \\ &= \int_{G/G_\alpha} \psi'_{\alpha}(\beta_{u(g')}(\mathcal{J}_{r_\zeta}(x^*))) dg' \\ &= \psi(\mathcal{J}_{r_\zeta}(x^*)) \\ &= \psi(x) \end{aligned}$$

We recall that $\mathcal{J}_{r_\zeta}^2 = \beta_{\mu I_d}$, where $\mu = 1$ for odd values of d and -1 for even values of d and compute the following easy steps:

$$\begin{aligned} & \psi'_{\alpha,0} \beta_{\mu I_d}(x) \\ &= \psi'_{\alpha,0} \mathcal{J}_{r_\zeta} \mathcal{J}_{r_\zeta}(x) \\ &= \psi'_{\alpha,0} \beta_{u(h_\zeta)} \mathcal{J}_{r_\zeta}(x^*) \end{aligned}$$

for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$. However \mathcal{J}_{r_ζ} commutes with $\{\beta_{u(g)} : g \in SU_2(\mathbb{C})\}$ and so

$$\begin{aligned} & \psi'_{\alpha,0} \beta_{\mu I_d}(x) \\ &= \psi'_{\alpha,0} \mathcal{J}_{r_\zeta}(\beta_{u(h_\zeta)}(x^*)) \\ &= \psi'_{\alpha,0} \beta_{u(h_\zeta^2)}(x) \end{aligned}$$

So we have

$$\begin{aligned} & \psi'_{\alpha,0} \beta_{u(h_\zeta)} \mathcal{J}_{r_\zeta}(x^*) \\ &= \psi'_{\alpha,0} \mathcal{J}_{r_\zeta} \beta_{u(h_\zeta)}(x^*) \\ &= \psi'_{\alpha,0} \beta_{u(h_\zeta^2)}(x) \\ &= \psi'_{\alpha,0} \beta_{\mu I_d}(x) \end{aligned}$$

We may reset ψ'_α as $\psi'_\alpha \beta_{u(h_\zeta)}$ and verify that $\psi'_{\alpha,0} \mathcal{J}_{r_\zeta}(x^*) = \psi'_{\alpha,0}(x)$ for $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ if and only if $\psi'_{\alpha,0} \beta_{\mu I_d} = \psi'_{\alpha,0}$ on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ for some $\psi'_{\alpha,0}$ in the factor decomposition of ψ . Same is true with $\psi'_{\alpha,0} \beta_{v(g)}$ for all $g \in SU_2(\mathbb{C})$ as $\beta_{\mu I}$ also commutes with $\{\beta_{u(g)} : g \in SU_2(\mathbb{C})\}$ on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$.

In particular,

$$\omega_{\alpha,0} \mathcal{J}_{r_\zeta}(x^*) = \omega_{\alpha,0}(x)$$

for all $x \in \tilde{\text{UHF}}_d \otimes \text{UHF}_d$ and thus the direct integral representation in (b) holds on \mathbb{M} for all integers $d \geq 2$.

For any $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$, we have $\psi_{\alpha,0}(\mathcal{J}_{r_\zeta}^\alpha(x^*)) = \psi_{\alpha,0}(x)$ provided $\psi'_{\alpha,0} \beta_{\mu I_d}(x) = \psi'_{\alpha,0}(x)$.

We claim that $\psi_{\alpha,0}(\mathcal{J}_{I_d}(x^*)) = \psi_{\alpha,0}(x)$ and $\psi_{\alpha,0}(\beta_{r_\zeta}(x)) = \psi_{\alpha,0}(x)$ for all $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$. We may write $\mathcal{J}_{r_\zeta} = \mathcal{J}_{I_d} \circ \beta_{r_\zeta}$ to verify first that $\psi_{\alpha,0} \beta_{r_\zeta}(x) = \psi_{\alpha,0}(x)$ for all $x \in \mathcal{O}_d$.

For $x \in \mathcal{O}_d$, $\mathcal{J}_{r_\zeta}(\mathcal{J}_{I_d}(x)) = \beta_{r_\zeta}(\mathcal{J}_{I_d}(x))$ since $\mathcal{J}_{I_d}^2 = I$. In particular, we have

$$\psi'_{\alpha,0} \beta_{r_\zeta}(\mathcal{J}_{I_d}(x)) = \psi_\alpha(\mathcal{J}_{I_d}(x))$$

for $x \in \mathcal{O}_d$. Now we use polarization identity

$$\mathcal{J}_{I_d}(x)y = \sum_{0 \leq k \leq 3} i^k \mathcal{J}_{I_d}(x + i^k y)(x + i^k y)$$

to deduce

$$\psi_\alpha \beta_{r_\zeta}(\mathcal{J}_{I_d}(x)y) = \psi_\alpha(\mathcal{J}_{I_d}(x)y)$$

for all $x, y \in \mathcal{O}_d$. This shows that

$$\psi_{\alpha,0} \beta_{r_\zeta} = \psi_{\alpha,0}$$

on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ whenever

$$\psi'_\alpha \beta_{\zeta^2 I_d} = \psi'_\alpha$$

on \mathcal{O}_d .

In particular, it also shows that $\psi'_{\alpha,0}\beta_{r_\zeta} = \psi'_{\alpha,0}$ on $\tilde{\text{UHF}}_d \otimes \text{UHF}_d$. By our construction, ψ'_α is an extremal element in $K_{\omega'_\alpha}$, where the state ω'_α is a λ -invariant ergodic state of $\mathbb{M}_R = \text{UHF}_d$. By Proposition 4.2 in [25], $\psi'_{\alpha,0}$ is in fact a pure state of $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$.

Since $\psi'_{\alpha,0}\beta_{u(g)}\mathcal{J}_{r_\zeta}(x^*) = \psi'_{\alpha,0}(\beta_{u(g)}(x))$ for all $g \in SU_2(\mathbb{C})$, we as well have $\psi'_{\alpha,0}\beta_{u(g)}\beta_{r_\zeta} = \psi'_{\alpha,0}\beta_{u(g)}$ for all $g \in SU_2(\mathbb{C})$ on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ provided $\psi'_\alpha(\beta_{\mu I_d}(x)) = \psi'_\alpha(x)$ for $x \in \mathcal{O}_d$. In other words, we have

$$\psi'_\alpha\beta_{u(g)}\beta_{r_\zeta}\beta_{u(g)^*} = \psi'_\alpha$$

on UHF_d for all $g \in SU_2(\mathbb{C})$.

We recall that $r_\zeta = u(h)$, where $h \in SU_2(\mathbb{C})$ is given explicitly in section 2, where $h = e^{\pi i \sigma_y}$ for odd values of d and $h = e^{\frac{\pi}{2} i \sigma_y}$. The normal subgroup generated by $\{ghg^{-1} : g \in SU_2(\mathbb{C})\}$ is not a subgroup of $\{-I, I\}$ since $h \notin \{-I, I\}$. Thus the normal subgroup is equal to $SU_2(\mathbb{C})$ i.e. $\psi'_{\alpha,0}\beta_{u(g)} = \psi'_{\alpha,0}$ on UHF_d for $g \in SU_2(\mathbb{C})$. This completes the proof for (c).

The last statement (d) is simple consequence of (c). ■

In Proposition 3.5, if ψ'_α were an extremal element in $K_{\omega'_\alpha}$, then by Proposition 2.6 we could have $\psi'_\alpha\beta_{u(g)} = \psi'_\alpha$ on \mathcal{O}_d for all $g \in SU_2(\mathbb{C})$. Thus in particular, we could have concluded that $\psi'_\alpha\beta_{\mu I_d} = \psi'_\alpha\beta_{r_\zeta}^2 = \psi'_\alpha$. The following proposition says much more.

PROPOSITION 3.6. *Let ω and ψ be states as in Proposition 3.5. Then ω is an extremal element in the convex set of real, lattice reflection symmetric and translation invariant state of \mathbb{M} . Moreover, the statements following hold for odd integer values for d :*

(a) *If ω is an ergodic state in the convex set of translation invariant states of \mathbb{M} then ω is a factor state of \mathbb{M} .*

(b) *If ω is reflection positive with twist r_0 on \mathbb{M} then ω is a factor state of \mathbb{M} .*

PROOF. We only need to prove statements (a) and (b) in light of Proposition 3.5 (d).

We fix a unitary element w in the centre \mathcal{Z}_{ψ_0} of $\pi_{\psi_0}(\tilde{\text{UHF}}_d \otimes \text{UHF}_d)''$. So $\omega Ad_w = \omega$ on \mathbb{M} and so there exists $z_w \in S^1$ so that $\psi Ad_w = \psi\beta_{z_w}$ on \mathcal{O}_d . We compute the following equalities for $x \in \tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$:

$$\begin{aligned} & \psi_0 Ad_w \mathcal{J}_{r_\zeta} Ad_w(x^*) \\ &= \psi_0 \beta_{z_w} \mathcal{J}_{r_\zeta} Ad_w(x^*) \\ &= \psi_0 \mathcal{J}_{r_\zeta} \beta_{\bar{z}_w} Ad_w(x^*) \\ &= \psi_0 Ad_w \beta_{z_w}(x) \\ &= \psi_0 \beta_{z_w} \beta_{\bar{z}_w}(x) \\ &= \psi_0(x) \\ &= \psi_0 \mathcal{J}_{r_\zeta}(x^*) \end{aligned}$$

Since ψ_0 is a pure state of $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$ by Proposition 4.3 in [25] (we need only extremal property of ω to show $\{x \in \mathcal{B}(\mathcal{K}) : \tau(x) = \tilde{\tau}(x) = x\}$ is $\mathbb{C}I$), we get

$$w\mathcal{J}_\zeta w\zeta_\psi = \mathcal{J}_\zeta \zeta_\psi = \zeta_\psi$$

where we have used notation for anti-unitary operator \mathcal{J}_ζ on $\tilde{\mathcal{H}} \otimes_{\mathcal{K}} \mathcal{H}$ for which $\mathcal{J}_{r_\zeta}(X) = \mathcal{J}_\zeta X \mathcal{J}_\zeta^*$ with $\mathcal{J}_\zeta \zeta_\psi = \zeta_\psi$ i.e. $\mathcal{J}_\zeta = \mathcal{J}_\gamma \hat{r}_\zeta$, notation used in Proposition 3.2.

We compute further that

$$\mathcal{J}_\zeta w \mathcal{J}_\zeta \zeta_\psi = w^* \zeta_\psi$$

Since $\mathcal{J}_\zeta w \mathcal{J}_\zeta^* \in \mathcal{Z}_\psi$, we conclude that

$$\mathcal{J}_{r_\zeta}(w)F_0 = w^*F_0$$

where $F_0 = [\pi_\psi(\text{UHF}_d \otimes \text{UHF}_d)''\zeta_\psi]$. Since the above relation holds for all unitary elements w in the centre \mathcal{Z}_{ψ_0} , by Stone's spectral theorem, we have

$$\mathcal{J}_{r_\zeta}(E)F_0 = EF_0$$

for all projection $E \in \mathcal{Z}_{\psi_0}$.

As Λ takes \mathcal{Z}_ψ to itself, we get

$$\mathcal{J}_{r_\zeta}(\Lambda(E))F_0 = \Lambda(E)F_0$$

Since $\mathcal{J}_{r_\zeta}(\Lambda(w)) = \tilde{\Lambda}(\mathcal{J}_{r_\zeta}(w))$, we arrive at $\tilde{\Lambda}(E)F_0 = \Lambda(E)F_0$. The action $X \rightarrow S_i^* X S_i$ also takes elements in \mathcal{Z}_ψ to itself and so we may take unitary element $S_i^* E S_i$ in place of E for

$$\tilde{\Lambda}(S_i^* E S_i)F_0 = \Lambda(S_i^* E S_i)F_0 = EF_0$$

The map $X \rightarrow \tilde{\Lambda}(S_i^* X S_i)$ is the left shift on $\text{UHF}_d \otimes \text{UHF}_d$ and so we conclude that

$$\theta^{-1}(E)F_0 = EF_0$$

Since the relation holds for all projection in the centre \mathcal{Z}_{ψ_0} , we also have

$$\theta(E)F_0 = EF_0$$

for all projection $E \in \mathcal{Z}_{\psi_0}$.

However $\theta(F_0) = F_0$ and thus

$$\theta(E') = E',$$

where $E' = EF_0$ is a projection in the centre of $\pi_\omega(\mathbb{M})''$. The state ω being an ergodic state pf \mathbb{M} , we conclude E' is either 0 or F_0 . Since there is an isomorphism between the centre of $\pi_\omega(\mathbb{M})''$ with $\mathcal{Z}_{\psi_0}F_0$, we conclude that ω is a factor state of \mathbb{M} . This completes the proof of (a).

We will modify our proof for (a) in order to prove (b). Suppose ω is not itself an extremal element in the convex set of translation invariant states. Since ω is an extremal element in the convex set of real, lattice reflection symmetric and translation invariant state, extremal decomposition of ω in the convex set of translation invariant states of \mathbb{M} is given by

$$\omega = \omega^1 \oplus \omega^1 \beta_{r_\zeta}$$

where ω^1 is a translation invariant ergodic state of \mathbb{M} . We may write $\psi_0 = \psi_0^1 \oplus \psi_0^1 \beta_{r_\zeta}$ for some extremal element $\psi^1 \in K_{\omega^1, G}$,

For any projection E' in the centre of $\pi_\omega(\mathbb{M})''$ we have

$$\mathcal{J}_{r_\zeta}(E' \mathcal{J}_{r_\zeta}(E')) = E' \mathcal{J}_{r_\zeta}(E')$$

since $\mathcal{J}_{r_\zeta}^2 = I$ and \mathcal{J}_{r_ζ} keeps $\pi_\omega(\mathbb{M})''$ invariant. By the purity property of the state ψ_0^1 on $\tilde{\mathcal{O}}_d \otimes \mathcal{O}_d$, as in the proof for (a), we arrive at

$$\theta(E' \mathcal{J}_{r_\zeta}(E')) = E' \mathcal{J}_{r_\zeta}(E')$$

Now by extremal property of ω in the convex set of real, lattice reflection symmetric and translation invariant states, we get

$$\psi(X E' \mathcal{J}_{r_\zeta}(E')) = \psi(X) \psi(E' \mathcal{J}_{r_\zeta}(E'))$$

for all $X \in \pi_\psi(\tilde{\text{UHF}}_d \otimes \text{UHF}_d)''$ and so $\psi(E' \mathcal{J}_{r_\zeta}(E'))$ is equal to 0 or 1.

The state ω being reflection positive with twist r_0 on \mathbb{M} and $SU_2(\mathbb{C})$, we have

$$\omega(\mathcal{J}_{r_\zeta}(X)X) \geq 0$$

for all $X \in \pi_\psi(\text{UHF}_d)''$. We consider now the map $(X, Y) \rightarrow \omega(\mathcal{J}_{r_\zeta}(X)Y)$ that is linear in right side and conjugate linear in the left side variables on $\pi_\psi(\mathcal{O}_d)''$. This bilinear map is skew-symmetric if and only if d is an odd integer, here we recall that $\zeta^2 = 1$ or -1 in accordance the values for d are odd or even. So for odd integer d , the map

$$(X, Y) \rightarrow \psi(\mathcal{J}_{r_\zeta}(XE)EY) = \psi(\mathcal{J}_{r_\zeta}(E)E)\psi(\mathcal{J}_{r_\zeta}(X)Y)$$

is a pre-inner product

$$\langle\langle X\zeta_\psi, Y\zeta_\psi \rangle\rangle_E = \omega(\mathcal{J}_{r_\zeta}(EX)EY)$$

on the vector space $[\pi_\psi(\text{UHF}_d)''\zeta_\psi]$.

However we computed above that $\omega(\mathcal{J}_{r_\zeta}(X)\mathcal{J}_{r_\zeta}(E)EX) = 0$ by the first part if $\psi(\mathcal{J}_{r_\zeta}(E)E) = 0$. In such case, we have $EX\zeta_\psi = 0$ i.e. $EF_0 = 0$. Similarly, if $\psi(E\mathcal{J}_{r_\zeta}(E)) = 1$, then $\psi(E) = 1 = \psi(\mathcal{J}_{r_\zeta}(E))$ and so by replacing the role of E by $I - E$, we arrive at $EF_0 = F_0$. Thus E' is either I or F_0 . So ω is a factor state of \mathbb{M} , contradicting our starting assumption that ω is itself not an extremal element in the convex set of translation invariant states. This completes the proof for (b). \blacksquare

In the following theorem, we sum up our main result.

THEOREM 3.7. *Let ω be a translation and $SU_2(\mathbb{C})$ -invariant, real, lattice symmetric state of \mathbb{M} and $\omega = \int^\oplus \omega_\alpha d\mu(\alpha)$ be an ergodic decomposition in the convex set of real, lattice symmetric and translation invariant states of \mathbb{M} . Then μ -almost everywhere ω_α are also $SU_2(\mathbb{C})$ invariant. Moreover, if d is an odd integer and ω is reflection positive with twist r_0 , then μ -almost everywhere ω_α are factor states of \mathbb{M} .*

In general the statement in Theorem 3.7 is false for even values of d and counter examples are included in section 5.

PROOF. Let K_{ω, G_r} be the non empty collection of λ and $SU_2(\mathbb{C})$ -invariant states of \mathcal{O}_d such that $\psi = \bar{\psi} = \tilde{\psi}$ with $\psi|_{\text{UHF}_d} = \omega|_{\mathbb{M}_R}$. We choose an extreme point $\psi \in K_{\omega, G_r}$ and write its extremal decomposition $\psi = \int \psi'_\alpha d\mu(\alpha)$ in the convex set of λ -invariant real and lattice symmetric states of \mathcal{O}_d , which need not be a factor state decomposition of \mathcal{O}_d . We set $\omega'_\alpha = \psi'_\alpha|_{\text{UHF}_d}$ and ω_α be its inductive limit translation invariant state of \mathbb{M} . So by our construction $\omega = \int \omega_\alpha d\mu(\alpha)$ is an extremal decomposition in the convex set of real, lattice symmetric translation

invariant states of \mathbb{M} . Now, we can repeat the argument used in Proposition 3.5 (a)-(c) to show

$$\omega_\alpha(\beta_{u(g)}(x)) = \omega_\alpha(x)$$

for all $x \in \mathbb{M}$ for $d \geq 2$.

However,

$$\psi_\alpha(\beta_{u(g)}(x)) = \psi_\alpha(x)$$

for all $x \in \mathcal{O}_d$ if d is an odd integer for which we have $\mu = \zeta^2 = 1$ and also \mathcal{J}_{r_ζ} admits direct integral representation given in Proposition 3.5 (a). Thus we verify reflection positive property of ω_α with the twist r_0 from the same property of ω , where we have used Proposition 3.5 (b) and $SU_2(\mathbb{C})$ -invariance property of ψ'_α . Now factor property of ψ_α follows from Proposition 3.6. ■

Unlike the situation $K_{\omega,G}$, another extremal element in K_{ω,G_r} need not be of the form $\psi\beta_z$ for some $z \in S^1$. Main obstruction comes from the fact that the \mathbb{Z}_2 symmetric action $\psi \rightarrow \bar{\psi}$ on λ -invariant states of \mathcal{O}_d takes $\psi\beta_z \rightarrow \bar{\psi}\beta_z$ on K_{ω,G_r} .

4. Integer spin s and half-odd integer spin s ($2s + 1 = d$):

PROPOSITION 4.1. *Let ω be a translation invariant real lattice symmetric pure state of \mathbb{M} with a twist r_0 as Proposition 3.2 and γ_{r_0} be the self adjoint unitary operator described in Proposition 3.2. If ω is also reflection positive with the twist r_0 then*

$$(47) \quad \langle \Delta^{-\frac{1}{2}} v_i^* \zeta_\psi, \beta_{r_0}(v_j^*) \zeta_\psi \rangle = \langle \Delta^{\frac{1}{2}} v_j^* \zeta_\psi, \beta_{r_0}(v_i^*) \zeta_\psi \rangle$$

$$(48) \quad \langle \Delta^{-\frac{1}{2}} \sigma_t(v_i^*) \zeta_\psi, \beta_{r_0}(\sigma_s(v_j^*)) \zeta_\psi \rangle = \langle \sigma_s(v_j^*) \zeta_\psi, \Delta^{\frac{1}{2}} \beta_{r_0}(\sigma_t(v_i^*)) \zeta_\psi \rangle \forall s, t \in \mathbb{R}$$

$$(49) \quad \langle \Delta^{-\frac{1}{2}} (v_i^{\delta_1})^* \zeta_\psi, \beta_{r_0}((v_j^{\delta_2})^*) \zeta_\psi \rangle = \langle (v_j^{\delta_2})^* \zeta_\psi, \Delta^{\frac{1}{2}} \beta_{r_0}((v_i^{\delta_1})^*) \zeta_\psi \rangle \forall \delta_1, \delta_2 > 0$$

and

$$(50) \quad \langle \zeta_\psi, v_i^{\delta_1} \sigma_{i(y-\frac{1}{2})}(\beta_{r_0}((v_j^{\delta_2})^*)) \zeta_\psi \rangle = \langle \zeta_\psi, v_j^{\delta_2} \sigma_{i(y+\frac{1}{2})}(\beta_{r_0}((v_i^{\delta_1})^*)) \zeta_\psi \rangle$$

for all $1 \leq i, j \leq d$, $t \in \mathbb{R}$ and $s = 1$, where

$$x^\delta = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp^{-\frac{1}{2} \frac{t^2}{\delta^2}} \sigma_t(x) dt$$

for $\delta > 0$ and $z \rightarrow \sigma_z(x)$ is the analytic extension of $t \rightarrow \sigma_t(x)$ for an analytic element $x \in \mathcal{M}$ to \mathbb{C} .

PROOF. We recall, $x\zeta_\psi$ is an element in the domain of $\Delta^{\frac{1}{2}}$ for $x \in \mathcal{M}$ and $y\zeta_\psi$ is an element in the domain of $\Delta^{-\frac{1}{2}}$ for $y \in \mathcal{M}'$ [6]. We also recall that $v_i^* \zeta_\psi = \tilde{v}_i^* \zeta_\psi$ and thus both sides of equalities in (47)-(49) are well defined. We need to establish those equalities.

The element $\sigma_t(v_i) \mathcal{J} \beta_{\bar{r}_\zeta}(\sigma_s(\tilde{v}_j)) \mathcal{J} \in \mathcal{M}_0$, i.e. $(\beta_z : z \in H)$ invariant element in \mathcal{M} as $\beta_z(v_i) = z v_i$ and $\beta_z(\mathcal{J} \beta_{\bar{r}_\zeta}(\tilde{v}_j) \mathcal{J}) = \bar{z} \mathcal{J} \beta_{\bar{r}_\zeta}(\tilde{v}_j) \mathcal{J}$, where we used commuting property of $(\beta_z : z \in H)$ with the modular group (σ_t) on \mathcal{M} and commuting property of $(u_z : z \in H)$ with \mathcal{J} as $\beta_z = Ad_{u_z}$ and $\beta_z(\tilde{v}_j) = z \tilde{v}_j$ for $z \in H$.

By our hypothesis that ω is reflection positive with twist r_0 , any element in \mathcal{M}_0 is $Ad_{\gamma_{r_\zeta}}$ invariant by Theorem 3.5 (d) in [25]. So we have the following equality for any $1 \leq i, j \leq d$:

$$\sigma_t(v_i) \mathcal{J} \beta_{\bar{r}_\zeta}(\sigma_s(\tilde{v}_j)) \mathcal{J}$$

$$\begin{aligned}
&= Ad_{\gamma_{r_\zeta}}(\sigma_t(v_i))Ad_{\gamma_{r_\zeta}}(\mathcal{J}\beta_{r_\zeta}(\sigma_s(\tilde{v}_j))\mathcal{J}) \\
&= \mathcal{J}\beta_{r_\zeta}(\sigma_t(\tilde{v}_i))\mathcal{J}\sigma_s(v_j)
\end{aligned}$$

where we used again modular group commutes with any automorphism that preserves the faithful normal state ϕ on \mathcal{M} . Since $r_\zeta = \zeta r_0$, we get

$$(51) \quad \sigma_t(v_i)\mathcal{J}\beta_{r_0}(\sigma_s(\tilde{v}_j))\mathcal{J} = \mathcal{J}\beta_{r_0}(\sigma_t(\tilde{v}_i))\mathcal{J}\sigma_s(v_j)$$

for all $s, t \in \mathbb{R}$.

Now we use $x^*\zeta_\psi = Sx\zeta_\psi = \mathcal{J}\Delta^{\frac{1}{2}}x\zeta_\psi$ for $x \in \mathcal{M}$ and $y^*\zeta_\psi = \mathcal{J}\Delta^{-\frac{1}{2}}y\zeta_\psi$ for $y \in \mathcal{M}'$ to verify (47) as we can compute the following equalities at ease:

$$\begin{aligned}
&\langle \Delta^{-\frac{1}{2}}\tilde{v}_i^*\zeta_\psi, \beta_{r_0}(v_j^*)\zeta_\psi \rangle \\
&= \langle v_i^*\zeta_\psi, \Delta^{-\frac{1}{2}}\beta_{r_0}(v_j^*)\zeta_\psi \rangle \\
&= \langle v_i^*\zeta_\psi, \Delta^{-\frac{1}{2}}\beta_{r_0}(\tilde{v}_j^*)\zeta_\psi \rangle \\
&= \langle v_i^*\zeta_\psi, \mathcal{J}\mathcal{J}\Delta^{-\frac{1}{2}}\beta_{r_0}(\tilde{v}_j^*)\zeta_\psi \rangle \\
&= \langle \zeta_\psi, v_i\mathcal{J}\beta_{r_0}(\tilde{v}_j)\mathcal{J}\zeta_\psi \rangle \\
&= \langle \zeta_\psi, \mathcal{J}\beta_{r_0}(\tilde{v}_i)\mathcal{J}v_j\zeta_\psi \rangle \\
&= \langle \mathcal{J}\beta_{r_0}(\tilde{v}_i^*)\mathcal{J}\zeta_\psi, v_j\zeta_\psi \rangle \\
&= \langle \mathcal{J}\beta_{r_0}(v_i^*)\zeta_\psi, v_j\zeta_\psi \rangle \\
&= \langle \mathcal{J}v_j\zeta_\psi, \beta_{r_0}(v_i^*)\zeta_\psi \rangle
\end{aligned}$$

(now by (51))

(we used conjugate linear property of \mathcal{J})

$$\begin{aligned}
&= \langle \mathcal{J}\mathcal{J}\Delta^{\frac{1}{2}}v_j^*\zeta_\psi, \beta_{r_0}(v_i^*)\zeta_\psi \rangle \\
&= \langle \Delta^{\frac{1}{2}}v_j^*\zeta_\psi, \beta_{r_0}(v_i^*)\Delta^{-\frac{1}{2}}\zeta_\psi \rangle
\end{aligned}$$

We can verify (48) along the same line. The equality (49) is a simple consequence of (48). For (50), we recall that x^δ is an analytic element for the modular group (σ_t) for any $x \in \mathcal{M}$ or \mathcal{M}' and $\delta > 0$. \blacksquare

THEOREM 4.2. *Let $g \rightarrow v_j^i(g)$ be an irreducible representation of $SU_2(\mathbb{C})$ and the state ω in Proposition 4.1 be also $SU_2(\mathbb{C})$ invariant. Then the following holds:*

- (a) d is an odd integer;
- (b) $\Delta = I$ and $\mathcal{M} = \mathcal{M}_0$ is a finite type-I factor and ϕ is the normalised trace on \mathcal{M} ;
- (c) H is the trivial subgroup of S^1 and $Ad_{\gamma_{r_\zeta}} = I$;
- (d) $v_i^* = \beta_{r_\zeta}(v_i)$;
- (e) There exists an irreducible representation $g \rightarrow \hat{u}(g) \in \mathcal{M}$ such that

$$(52) \quad \hat{u}(g)v_i^*\hat{u}(g)^* = \beta_{u(g)}(v_i^*)$$

and representation $g \rightarrow \hat{u}(g)$ is an odd integer or even integer representation of $SU_2(\mathbb{C})$.

PROOF. By Clebsch-Gordan theory valid for irreducible representation $g \rightarrow u(g)$ of the group $SU_2(\mathbb{C})$, the representation $g \rightarrow u(g) \otimes u(\bar{g})$ in $\mathbb{C}^d \otimes \mathbb{C}^d$ admits a unique invariant subspace. The state ω being $SU_2(\mathbb{C})$ -invariant, the vectors $\langle \zeta_\psi, S_i^* S_j \zeta_\psi \rangle$ and $\langle \zeta_\psi, S_j S_i^* \zeta_\psi \rangle$ are $g \rightarrow u(\bar{g}) \otimes u(g)$ invariant by Proposition 2.5 (a) and thus

$$\langle \zeta_\psi, S_i S_j^* \zeta_\psi \rangle = \frac{\delta_j^i}{d}$$

The state ω being $SU_2(\mathbb{C})$ -invariant, the vector $\langle \zeta_\psi, v_i \Delta^s v_j^* \zeta_\psi \rangle$ is also $g \rightarrow u(g) \otimes u(\bar{g})$ invariant for any real s since Δ commutes with $u(\hat{g})$ for all $g \in SU_2(\mathbb{C})$ by Proposition 2.5 (c).

For the time being, we fix $\delta > 0$ and simplify notation v_i^δ for v_i and compute that

$$(53) \quad \langle \zeta_\psi, v_i \Delta^y v_j^* \zeta_\psi \rangle = \delta_j^i c_y$$

for some positive constant c_y independent of i, j .

Now we use (53) in the equality (50) for $y = \frac{1}{2}$ to conclude that

$$(r_0)_i^j \langle \zeta_\psi v_i v_i^* \zeta_\psi \rangle = (r_0)_j^i \langle \zeta_\psi v_j \Delta v_j^* \zeta_\psi \rangle$$

for all $1 \leq i, j \leq d$. Since $r_0^2 = I$, i.e. $r_0 = r_0^*$ and each row or column vector is non zero, we conclude that

$$(54) \quad \langle \zeta_\psi v_i v_i^* \zeta_\psi \rangle = \langle \zeta_\psi v_j \Delta v_j^* \zeta_\psi \rangle$$

for some i, j and hence for all $1 \leq i, j \leq d$ by (53). Similarly, we also use (53) in the equality (50) for $y = -\frac{1}{2}$ to conclude that

$$(55) \quad \langle \zeta_\psi v_i \Delta^{-1} v_i^* \zeta_\psi \rangle = \langle \zeta_\psi v_j v_j^* \zeta_\psi \rangle$$

for some i, j and hence for all $1 \leq i, j \leq d$ by (53).

So we have by (54) and (55)

$$\begin{aligned} & \|[\Delta^{\frac{1}{2}} v_i^* - \Delta^{-\frac{1}{2}} v_i^*] \zeta_\psi\|^2 \\ &= \|\Delta^{\frac{1}{2}} v_i^* \zeta_\psi\|^2 + \|\Delta^{-\frac{1}{2}} v_i^* \zeta_\psi\|^2 - 2\|v_i^* \zeta_\psi\|^2 \\ &= 0 \end{aligned}$$

By separating property for ζ_ψ for \mathcal{M} , we get

$$\Delta v_i^* \Delta^{-1} = v_i^*$$

i.e. Δ commutes with each v_i^* . Δ being self-adjoint, Δ also commutes with each v_i i.e. Δ commutes with each v_i^δ for any $\delta > 0$ once we remove simplified notation. Thus Δ commutes with each v_i and so $\Delta \in \mathcal{M}'$. Since $\mathcal{J} \Delta \mathcal{J} = \Delta^{-1}$, we also conclude $\Delta \in \mathcal{M}$. The von Neumann algebra \mathcal{M} being a factor and $\Delta \zeta_\psi = \zeta_\psi$, we conclude $\Delta = I$.

We claim that \mathcal{M} is a finite type-I factor rather than a type-II₁ finite factor. Suppose not. Then \mathcal{M}_0 is also a type-II₁ finite factor. The von-Neumann factor \mathcal{M}_0 being the corner of $\pi_\omega(\text{UHF}_d)''$ by $P \in \pi_\psi(\text{UHF}_d)''$, $\pi_\omega(\text{UHF}_d)''$ is also a type-II von-Neumann factor.

We consider the GNS space $(\mathcal{H}_{\omega_R}, \pi_{\omega_R}, \zeta_{\omega_R})$ associated with (\mathbb{M}_R, ω_R) . So $\pi_{\omega_R}(\mathbb{M}_R)''$ is a type-II factor and ζ_{ω_R} is cyclic for $\pi_{\omega_R}(\mathbb{M}_R)''$ in \mathcal{H}_{ω_R} . We will

rule out the following two possible cases: As in (17), we identify \mathbb{M}_R with UHF_d with respect to a orthonormal basis (e_i) for \mathbb{C}^d .

(i) $\pi_{\omega_R}(\text{UHF}_d)''$ is a type-II₁ factor.

In such a case $\pi_{\omega_R}(\text{UHF}_d)''$ admits a unique tracial state say ω_0 [11]. Since $\omega_0\Lambda$ is also a tracial state on $\pi_{\omega_R}(\text{UHF}_d)''$, we get by uniqueness of tracial state, $\omega_0 = \omega_0\Lambda$. But $\omega\Lambda = \omega$ on $\pi_{\omega_R}(\text{UHF}_d)''$ and ω is a factor state, in particular, an ergodic state i.e. unique invariant state of right translation dynamics (UHF_d, θ) . Thus $\omega = \omega_0$ on UHF_d . So ω is the unique trace on \mathbb{M} , contradicting our hypothesis that ω is pure.

(ii) $\pi_{\omega_R}(\text{UHF}_d)''$ is a type-II_∞ factor.

In this case, P is a finite projection in $\pi_{\omega_R}(\text{UHF}_d)''$ and $\mathcal{M}_0 = P\pi_{\omega_R}(\text{UHF}_d)''P$ is type-II₁ factor and $\pi_{\omega}(\text{UHF}_d)''$ is isomorphic to $\mathcal{M}_0 \otimes \mathcal{B}(\mathcal{H})$, where \mathcal{M}_0 is type-II₁ factor acting on \mathcal{K} and \mathcal{H} is an infinite dimensional Hilbert space [11]. Note that \mathcal{M}'_0 is also a type-II₁ factor and $\pi_{\omega_R}(\text{UHF}_d)'$ is isomorphic to \mathcal{M}'_0 .

More generally, we claim that the commutant of $\Lambda^n(\pi_{\omega_R}(\text{UHF}_d)''$ is also a type-II₁ factor isomorphic to $\mathcal{M}'_0 \otimes \{S_I S_J^* : |I| = |J| = n\}''$. That the type-II₁ factor $\Lambda^n(\pi_{\omega_R}(\text{UHF}_d)')$ contains $\mathcal{M}'_0 \otimes \{S_I S_J^* : |I| = |J|\}''$ is obvious. The factor being a hyperfinite type-II₁ factor, we may write $\pi_{\omega_R}(\Lambda^n(\text{UHF}_d)') = \mathcal{N}_0 \otimes \{S_I S_J^* : |I| = |J| = n\}''$ for some type-II₁ factor \mathcal{N}_0 and $\mathcal{M}'_0 \subseteq \mathcal{N}_0$. For the reverse inclusion, if $X \in \mathcal{N}_0$ then $X \in \pi_{\omega_R}(\text{UHF}_d)'$ and so $X \in \mathcal{M}'_0$.

So $\mathcal{M}'_0 \otimes \pi_{\omega}(\text{UHF}_d)''$ admits a tracial state and it is a type-II₁ factor. However, Cuntz relation (16) gives

$$\bigcap_{n \geq 1} \Lambda^n(\pi_{\omega_R}(\text{UHF}_d)'' \subseteq \pi_{\omega_R}(\text{UHF}_d)'' \bigcap \pi_{\omega}(\text{UHF}_d)'$$

So by the factor property of ω_R , we also have $\mathcal{M}'_0 \otimes \pi_{\omega}(\text{UHF}_d)''$ is the algebra of all bounded operators on \mathcal{H}_{ω_R} . This brings a contradiction.

That, d can not be an even integer, is given in [22] since $\omega_R(\text{UHF}_d)''$ is a type-I factor. It also follows by a more general result [26], where we could drop additional assumption that ω is reflection positive with the twist r_0 but here (a) is valid for reflection positive with twist case.

Thus d is an odd integer and ω_R is a type-I factor state of UHF_d with its corner $P\pi_{\omega}(\text{UHF}_d)''P$ equal to a finite type-I factor \mathcal{M} and by Proposition 2.2 (e) in [26] we have

$$\pi_{\psi}(\mathcal{O}_d)'' = \pi_{\psi}(\text{UHF}_d)''$$

Since $\beta_z(S_i) = zS_i$ for any $z \in H$ but $\beta_z(X) = X$ for all $X \in \pi_{\psi}(\text{UHF}_d)'' = \pi_{\psi}(\mathcal{O}_d)''$. So we have $S_i = zS_i$ each $1 \leq i \leq d$ and $z \in H$. This shows $z = 1$ as $S_i^* S_i = I$ for each $1 \leq i \leq d$. Thus $H = \{1\}$.

This also shows that $\mathcal{M}_0 = \mathcal{M}$ since $\mathcal{M}_0 = P\pi_{\psi}(\text{UHF}_d)''P$ and $\mathcal{M} = P\pi_{\psi}(\mathcal{O}_d)''P$. Thus $Ad_{\gamma_{r_\zeta}} = I$ on \mathcal{M} as well and so

$$\begin{aligned} v_i^* &= Ad_{\gamma_{r_\zeta}}(v_i^*) \\ &= \mathcal{J}\beta_{\tilde{r}_\zeta}(\tilde{v}_i^*)\mathcal{J} \\ &= \beta_{r_\zeta}(v_i) \end{aligned}$$

since $\Delta = I$ and so $\tilde{v}_i^* = \mathcal{J}v_i\mathcal{J}$ for $1 \leq i \leq d$

We are left to prove the last statement (e). The factor \mathcal{M} being type-I and $SU_2(\mathbb{C})$ being simply connected, first part follows by a standard result in representation theory [20].

We will prove now that the group action $\alpha_g : x \rightarrow \hat{u}(g)x\hat{u}(g)^*$ on \mathcal{M} is ergodic i.e. there exists no non-trivial invariant element for the group action. Let $(p_i : 1 \leq i \leq m)$ be a maximal set of orthogonal minimal projections in $\mathcal{M}_G = \{x \in \mathcal{M} : \alpha_g(x) = x, \forall g \in SU_2(\mathbb{C})\}$ i.e. elements in \mathcal{M} that are invariant for the group action ($\alpha_g : g \in SU_2(\mathbb{C})$) and u be a unitary element in \mathcal{M} invariant for the group action α_g as well.

So V as well as $V_u = (uv_i u^*)$ satisfies the inter-twinning relation (28). By the uniqueness of Clebs-Gordon coefficients, we get

$$(56) \quad p_i u v_k u^* p_j = c_j^i(u) p_i v_k p_j$$

for all $1 \leq k \leq d$ with some scalars $c_j^i(u) \in \mathbb{C}$.

We compute now the following

$$(57) \quad \sum_{1 \leq k \leq d} p_i u v_k u^* p_j u v_k^* u^* p_i = |c_j^i(u)|^2 \sum_{1 \leq k \leq d} p_i v_k p_j v_k^* p_i$$

Since $\sum_k v_k p_j v_k^*$ is also α_g -invariant, the left hand side of (57) is independent of u , is in the centre of \mathcal{M}_G and so $|c_j^i(u)| = 1$. By (d), we also have $v_k^* = \beta_{r_0}(v_k)$ and the family of vectors $\{p_i v_k p_j v_l^* : 1 \leq k \leq d\}$ are mutually orthogonal for each fix $1 \leq i, j \leq d$ (as $(\phi(p_i v_k p_j v_l^* p_i))$ is a invariant vector for the representation $g \rightarrow u(g) \otimes \overline{u(g)}$ of $SU_2(\mathbb{C})$), $c_j^i(u)$ is a real number and so either equal to 1 or -1 . The set of invariant unitary elements in the centre of \mathcal{M}_G is a connected set and the map $u \rightarrow c_j^i(u)$ is continuous. Thus $c_j^i(u) = 1$. So we get $u v_k u^* = v_k$ for all k i.e. $u \in \mathcal{M}'$. Since \mathcal{M} is a factor, we conclude that u is a scalar multiple of identity element of \mathcal{M} . This shows that \mathcal{M}_G is a subfactor of finite type-I factor \mathcal{M} .

Without loss of generality we write $\mathcal{M} = \mathcal{M}'_G \otimes \mathcal{M}_G$ and $\hat{u}(g) = \hat{u}(g) \otimes I_{\mathcal{M}_G}$ for all $g \in SU_2(\mathbb{C})$ and $\mathcal{M}'_G = \{\hat{u}(g) : g \in SU_2(\mathbb{C})\}''$.

If $u \in \mathcal{M}_G$ in (57), is only an element that commutes with each (p_i) , then the left hand side of (57) is also independent of u since G -invariant element $p_i \tau(p_j) p_i$ is a scalar multiple of p_i as each p_i is a minimal projection in \mathcal{M}_G . Now we follow the same argument used above to conclude that u is a scalar multiple of identity operator. Thus \mathcal{M}_G is trivial. This completes the proof for irreducibility of the representation $g \rightarrow \hat{u}(g)$ of $SU_2(\mathbb{C})$.

That the dimension of the representation $g \rightarrow \hat{u}(g)$ could be an odd integer or even integer follows once we appeal to Clebsch-Gordon theorem for

$$u_s(g) \otimes u_t(g) \equiv u_{|t-s|} \oplus u_{|t-s|+1} \oplus \dots \oplus u_{s+t}(g)$$

to verify that for any integer value of s , $u_t(g)$ is present in the decomposition of $u_s(g) \otimes u_t(g)$ irrespective of the value t that could be either an integer spin or $\frac{1}{2}$ -odd integer spin representation. ■

Now we sum up our main result of this section in the following theorem with a natural generalisation.

THEOREM 4.3. *Let G be a simply connected group and $g \rightarrow u_j^i(g)$ is a d -dimensional irreducible representation of G such that $g \rightarrow u(g) \otimes u(g)$ admits a unique one dimensional invariant subspace in $\mathbb{C}^d \otimes \mathbb{C}^d$ and ω be a real, lattice reflection symmetric with a twist r_0 , translation invariant pure state of \mathbb{M} . If ω is also G -invariant and reflection positive with the twist r_0 then there exists an extremal element $\psi \in K_\omega$ so that its associated elements in Proposition 2.5 ($\mathcal{K}, \mathcal{M}, v_k : 1 \leq k \leq d$) satisfies the following:*

- (a) $\Delta = I$ and $\mathcal{M} = \mathcal{M}_0$ is a finite type-I factor;
- (b) H is the trivial subgroup of S^1 and $Ad_{\gamma_{r_\zeta}} = I$;
- (c) For each $1 \leq i \leq d$, we have $v_i^* = \beta_{r_\zeta}(v_i)$;
- (d) Two-point spatial correlation functions of ω decay exponentially.

PROOF. First part of the statement is a simple generalisation of Theorem 4.2 and its proof follows by simple inspection of the proof where we have used those properties of the representation rather than explicit use of it.

Now we consider the contractive operator $Ta\zeta_\psi = \tau(a)\zeta_\psi$, $a \in \mathcal{M}$, where $\tau(a) = \sum_i v_i a v_i^*$, $x \in \mathcal{M}$ and the tracial state $a \rightarrow \langle \zeta_\psi, a\zeta_\psi \rangle$ on the finite matrix algebra \mathcal{M} is invariant for τ . The equality in (c) in particular says that T is also self adjoint and so T^2 is positive.

Thus the exponentially decaying property of two point spatial correlation would be related with the mass gap in the spectrum of T^2 from 1 once we show that any invariant vector of T^2 is a scalar multiple of ζ_ψ .

Let f be an invariant vector for T^2 . Then we get

$$\begin{aligned} & \langle f, a\zeta_\psi \rangle \\ &= \langle T^{2n} f, a\zeta_\psi \rangle \\ &= \langle f, \tau^{2n}(a)\zeta_\psi \rangle \end{aligned}$$

for all $n \geq 1$ and $a \in \mathcal{M}$. Taking $n \rightarrow \infty$, we conclude that

$$\langle f, a\zeta_\psi \rangle = \langle \zeta_\psi, a\zeta_\psi \rangle \langle f, \zeta_\psi \rangle$$

for all $a \in \mathcal{M}$ i.e. $f = 0$ if f is orthogonal to ζ_ψ .

Let $0 \leq \delta < 1$ and δ^2 be the highest eigen value of $T^2 - |\zeta_\psi\rangle\langle\zeta_\psi|$ and $\beta > 0$ so that $e^\beta \delta < 1$. So we have $\|T - |\zeta_\psi\rangle\langle\zeta_\psi|\| \leq \delta I$ and for any $A, B \in \mathbb{M}$

$$\begin{aligned} & e^{\beta n} |\omega(A\theta^n(B)) - \omega(A)\omega(B)| \\ &= e^{\beta n} |\langle a^* \zeta_\psi, [T - |\zeta_\psi\rangle\langle\zeta_\psi|]^n b \zeta_\psi \rangle| \\ &\leq (e^\beta \delta)^n \|a\| \|b\| \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, where $a = P\pi_\psi(A)P$ and $b = P\pi_\psi(B)P$ are elements in \mathcal{M} and $e^\beta \delta < 1$. ■

5. Ground states of Hamiltonian in quantum spin chain

We are left to discuss few motivating examples for this abstract framework, developed so far to study symmetries of Hamiltonian H that satisfies (3) and (14). Before we take few specific examples, we recall some well known results in the following proposition for our reference and its consequences in light of results proved in section 4 and 5.

PROPOSITION 5.1. *Let H be a Hamiltonian in quantum spin chain $\mathbb{M} = \otimes_{\mathbb{Z}} M_d(\mathbb{C})$ that satisfies relation (3) with $h_0 \in \mathbb{M}_{loc}$. Then the following statements are true:*

(a) *There exists a unique KMS state ω_β for (α_t) at each inverse positive temperature $\beta = \frac{1}{kT} > 0$ and ω_β is a translation invariant factor state of \mathbb{M} .*

(b) *If H also satisfies relation (14) with $J > 0$ and $r_0 \in U_d(\mathbb{C})$, then the unique KMS state ω_β is reflection positive with twist r_0 . Furthermore, any weak* limit point of ω_β as $\beta \rightarrow \infty$ is also reflection positive with twist r_0 ;*

(c) *If H is also $SU_2(\mathbb{C})$ -invariant i.e. $\beta_{u(g)}(h_0) = h_0$ and ω be a low temperature limit point ground state for H described in (b) and $\omega = \int_X^{\oplus} \omega^\alpha d\mu(\alpha)$ be its extremal decomposition in the convex set of real, lattice reflection symmetric and translation invariant states of \mathbb{M} . Then the following hold:*

(i) *For any odd integer $d \geq 3$, μ -almost everywhere, $\omega^\alpha = \omega^\alpha \beta_{u(g)}$ for all $g \in SU_2(\mathbb{C})$ and ω^α are pure ground states of H .*

(ii) *For any even integer $d \geq 2$, μ -almost everywhere $\omega^\alpha = \omega^\alpha \beta_{u(g)}$ for all $g \in SU_2(\mathbb{C})$ but ω^α are not even an extremal element in the convex set of translation invariant states of \mathbb{M} .*

PROOF. For (a), we refer to H. Araki's work [4] and also [19]. For the first statement in (b), we refer to [14]. Last part of (b) is trivial as reflection positive property (11) is closed under weak* limit.

For the statement of (c), we use Theorem 3.7. In particular, $SU_2(\mathbb{C})$ -invariance ensures that μ -almost everywhere ω^α are stationary states for the Hamiltonian dynamics H for which $\beta_{u(g)}(h_0) = h_0$ for all $g \in SU_2(\mathbb{C})$ and so μ -almost everywhere ω^α are as well ground states of H , since the set of ground states is a face in the convex set of stationary states [8].

We use now a standard fact that factor decomposition coincides with extremal decomposition for a ground state of a Hamiltonian H [8] and so extreme points are pure states of \mathbb{M} for odd values of d . For even values of d , these extreme points are not extremal elements in the convex set of translation invariant states of \mathbb{M} as otherwise these states would have been factor states of \mathbb{M} by Proposition 3.6 (a) and so would have been pure states of \mathbb{M} , contradicting the fact that there exists no pure state that is real, lattice symmetric and $SU_2(\mathbb{C})$ -invariant [26]. ■

For an even integer, such a extremal element ω_α in the decomposition given in Proposition 5.1 (c) is an $SU_2(\mathbb{C})$ -invariant ground state of H but not far from being extremal in the convex set of translation invariant states of \mathbb{M} . Nevertheless, we have

$$\omega_\alpha = \omega_\alpha^1 \oplus \omega_\alpha^1 \beta_{r_\zeta}$$

for some translation invariant ergodic states ω_α^1 of \mathbb{M} . In particular, this shows that translation invariant ergodic states ω_α^1 are $SU_2(\mathbb{C})$ invariant ground states of H for μ -almost everywhere but fails to be real and lattice symmetric. In other words, spontaneous \mathbb{Z}_2 symmetry $\omega \rightarrow \bar{\omega}$ breaks down rather than $SU_2(\mathbb{C})$ symmetry [26] in the decomposition given in Proposition 5.1 (c) for even values of $d \geq 2$.

PROOF. (Theorem 1.3) Proof for (a) is given Proposition 5.1 (c) and for (b) we recall that there is no real, lattice symmetric, $SU_2(\mathbb{C})$ and translation invariant pure state of \mathbb{M} for even values of d [26]. ■

6. Haldane's conjecture:

If the ground state of anti-ferromagnetic model H^{XXX} admits unique ground state then the ground state ω_{XXX} is pure, translation invariant, $SU_2(\mathbb{C})$ -invariant, reflection symmetric with twist and positive. The main result, Theorem 4.2 says that such a state is also finitely correlated and its spatial correlation functions decay exponentially. Moreover, there exists an automorphism α on \mathbb{M} commuting with θ such that $\omega\alpha$ is an infinite tensor product state of \mathbb{M} [27]. Furthermore, mean entropy $s(\omega_{XXX})$. Thus by upper-semicontinuity property of mean entropy, $s(\omega_\beta) \rightarrow 0$ as $\beta \rightarrow \infty$, if ω_β is the unique thermal equilibrium factor state at inverse temperature $\beta = \frac{1}{kT}$ of H^{XXX} model. This gives an important criteria for possible purity property of ω_{XXX} by studying its asymptotic behaviour for factor or cluster property [32] of ω_β as described in section 6 of [27]. The following statement is an easy consequence of standard results [9, 21].

THEOREM 6.1. *Let ω_β be the unique thermal equilibrium or KMS factor state at inverse temperature β for anti-ferromagnetic H^{XXX} model with odd integer $d = 2s + 1 \geq 3$ (s is an integer greater than 1) and ω be a limit point of ω_β as $\beta \rightarrow \infty$. Then following holds:*

(a) Then $\omega = \int \omega_r d\mu(r)$, where ω_r is the state defined by

$$\omega_r(e_{j_1}^{i_1} \otimes \dots \otimes e_{j_n}^{i_n}) = \phi(v_I v_J^*)$$

and $v = (v_i)$ is the unique solution to Clebsch-Gordon equation (52) as described in Theorem 4.2 satisfying (d) and (e) with irreducible representations $g \rightarrow u(g) = u_s(g)$ and $g \rightarrow \hat{u}(g) = u_r(g)$ of $SU_2(\mathbb{C})$ with finite I_{2r+1} factor \mathcal{M} for an integer $r \geq 1$ or half-odd integer.

(b) In (a) the dimension of \mathcal{M} i.e. irreducible representation $g \rightarrow \hat{u}_r(g)$ in Theorem 4.2 with dimension $2r + 1$ with half odd-integer r or integer r is determined by minimising mean energy of H^{XXX} over all possible solutions to (52) with irreducible representations $g \rightarrow \hat{u}_l(g)$ of dimension $2l + 1$ with half odd-integer or integer (each ω_l is an invariant state for Hamiltonian flow $\hat{\sigma}_t$ of H^{XXX}) i.e.

$$(58) \quad \omega_r(h_0) = \min_{l=\frac{1}{2}, 1, \frac{3}{2}, \dots} \omega_l(h_0)$$

(c) If there exist unique r for which $\omega_l(h_0^{xxx})$ attains its minimum then $\omega = \omega_r$ i.e. the low temperature limit of ω_β as $\beta \rightarrow \infty$ is unique and its limiting value is ω_r .

We illustrate our results for $d = 2s + 1 = 3$ in the following text for possible further investigation.

Now we briefly discuss the situation when $d = 3$ i.e. $s = 1$. In such a case Pauli spin matrices are given by

$$\sigma_x = 2^{-\frac{1}{2}} \begin{pmatrix} 0 & , & 1, & 0 \\ 1 & , & 0, & 1 \\ 0 & , & 1, & 0 \end{pmatrix},$$

$$\sigma_y = 2^{-\frac{1}{2}} \begin{pmatrix} 0 & , & -i, & 0 \\ i & , & 0, & -i \\ 0 & , & i, & 0 \end{pmatrix},$$

$$\sigma_z = \begin{pmatrix} 1 & , & 0, & 0 \\ 0 & , & 0, & 0 \\ 0 & , & 0, & -1 \end{pmatrix}.$$

and $i\sigma_x, i\sigma_y, i\sigma_z$ are basis for Lie-algebra $su_2(\mathbb{C})$ with

$$[i\sigma_x, i\sigma_y] = -i\sigma_z, \quad [i\sigma_y, i\sigma_z] = -i\sigma_x, \quad [i\sigma_z, i\sigma_x] = -i\sigma_y$$

A direct calculation shows that the inter-twiner $r_\zeta = r_0$ is a matrix with real entries given below

$$r_\zeta = \begin{pmatrix} 0 & , & 0, & -1 \\ 0 & , & 1, & 0 \\ -1 & , & 0, & 0 \end{pmatrix}.$$

and

$$h_0^{xxx} = J(\sigma_x \otimes \sigma_x + \sigma_y \otimes \sigma_y + \sigma_z \otimes \sigma_z)$$

By Clebsch-Gordon decomposition of $SU_2(\mathbb{C})$ representation $g \rightarrow u_1(g) \otimes u_1(g)$, the commutant of $\{u_1(g) \otimes u_1(g) : g \in SU_2(\mathbb{C})\}$ in $\mathbb{M}_3(\mathbb{C}) \otimes \mathbb{M}_3(\mathbb{C})$ is equal to its centre made of orthogonal projections of dimension 1, 3, 5. Since h_0 commutes with $u_1(g) \otimes u_1(g)$, h_0 is in the centre of $\{u_1(g) \otimes u_1(g) : g \in SU_2(\mathbb{C})\}''$ and so $\omega_l(xh_0) = \omega_l(h_0x)$ for any $x \in \mathbb{M}_3(\mathbb{C}) \otimes \mathbb{M}_3(\mathbb{C})$ since ω_l is $SU_2(\mathbb{C})$ -invariant of \mathbb{M} . Thus $\omega_l \sigma_t^{XXX} = \omega_l$ for all $t \in \mathbb{R}$ on local elements of \mathbb{M} and so on \mathbb{M} .

We compute with $J = 1$

$$\begin{aligned} \omega_l(h_0^{xxx}) &= \frac{1}{2}\phi(v_1av_2^* + v_2av_1^* + v_3av_2^* + v_2av_3^*) \\ &\quad - \frac{1}{2}\phi(-v_1bv_2^* + v_2bv_1^* + v_3bv_2^* - v_2bv_3^*) \\ &\quad + \phi(v_1cv_1^* - v_3cv_3^*) \end{aligned}$$

(where $a = v_1v_2^* + v_2v_1^* + v_3v_2^* + v_2v_3^*$, $b = -v_1v_2^* + v_2v_1^* + v_3v_2^* - v_2v_3^*$ and $c = v_1v_1^* - v_3v_3^*$)

$$\begin{aligned} &= \frac{1}{2}\phi((v_2^*v_1 + v_1^*v_2 + v_3^*v_2 + v_2^*v_3)(v_1v_2^* + v_2v_1^* + v_3v_2^* + v_2v_3^*)) \\ &\quad - \frac{1}{2}\phi((-v_2^*v_1 + v_1^*v_2 + v_2^*v_3 - v_3^*v_2)(-v_1v_2^* + v_2v_1^* + v_3v_2^* - v_2v_3^*)) \\ &\quad + \phi((v_1^*v_1 - v_3^*v_3)(v_1v_1^* - v_3v_3^*)) \end{aligned}$$

(where we have tracial state property of ϕ on \mathcal{M})

$$\begin{aligned} &= \phi((v_2^*v_1 + v_3^*v_2)(v_2v_1^* + v_3v_2^*)) \\ &\quad + \phi((v_1^*v_2 + v_2^*v_3)(v_1v_2^* + v_2v_3^*)) \\ &\quad + \phi((v_1^*v_1 - v_3^*v_3)(v_1v_1^* - v_3v_3^*)) \end{aligned}$$

Since $v_1^* = -v_3$, $v_2^* = v_2$ and $v_1v_1^* + v_2v_2^* + v_3v_3^* = I$, we simplify further that

$$\begin{aligned} \omega_l(h_0^{xxx}) &= \phi(\delta_{v_2}(v_1)\delta_{v_2}(v_1^*)) + \phi(\delta_{v_2}(v_3)\delta_{v_2}(v_3^*)) \\ (59) \quad &\quad + \phi(((v_1^*v_1 - v_3^*v_3)(v_1v_1^* - v_3v_3^*))) \end{aligned}$$

where we have used the symbol $\delta_{v_2}(a) = v_2a - av_2$ for $a \in \mathcal{M}$.

Solution to (52) is given by $v_1 = l_+$, $v_3 = l_-$ and $v_2 = il_z$, where

$$\sqrt{l(l+1)}l_+ = \frac{1}{\sqrt{2}}(\bar{\pi}_l(i\sigma_x) + i\bar{\pi}_l(i\sigma_y)),$$

$$\sqrt{l(l+1)}l_- = \frac{1}{\sqrt{2}}(\bar{\pi}_l(i\sigma_x) - i\bar{\pi}_l(i\sigma_y))$$

and

$$\sqrt{l(l+1)}l_z = \bar{\pi}_l(i\sigma_z)$$

where we used notation $\bar{\pi}_l$ for $2l+1$ -dimensional irreducible representation of Lie-algebra $su_2(\mathbb{C})$ of the Lie-group $SU_2(\mathbb{C})$. So

$$\begin{aligned} & l(l+1)\delta_{v_2}(v_1) \\ &= \frac{i}{\sqrt{2}}([\bar{\pi}_l(i\sigma_z), \pi_l(i\sigma_x)] + i[\bar{\pi}_l(i\sigma_z), \bar{\pi}_l(i\sigma_y)]) \\ &= \frac{i}{\sqrt{2}}(\bar{\pi}_l(-i\sigma_y) + i(\bar{\pi}_l(i\sigma_x))) \\ &= \frac{1}{\sqrt{2}}(-i\bar{\pi}_l(i\sigma_y) - \bar{\pi}_l(i\sigma_x)) \\ &= -\sqrt{l(l+1)}v_1 \end{aligned}$$

i.e

$$\sqrt{l(l+1)}\delta_{v_2}(v_1) = -v_1$$

Similarly, we may compute by taking a joint that

$$\sqrt{l(l+1)}\delta_{v_2}(v_3) = v_3$$

We also compute that

$$\begin{aligned} & l(l+1)(v_1v_1^* - v_3v_3^*) \\ &= -l(l+1)(v_1v_3 - v_3v_1) \\ &= -\frac{1}{2}[\pi_l(i\sigma_x) + i\pi_l(i\sigma_y), \pi_l(i\sigma_x) - i\pi_l(i\sigma_y)] \\ &= i[\pi_l(i\sigma_x), \pi_l(i\sigma_y)] \\ &= i\bar{\pi}_l(-i\sigma_z) \\ &= -i\bar{\pi}_l(i\sigma_z) \end{aligned}$$

$$\text{i.e. } \sqrt{l(l+1)}(v_1v_1^* - v_3v_3^*) = -v_2$$

Now from (59), we get

$$\begin{aligned} \omega_l(h_0^{xxx}) &= -\frac{1}{l(l+1)}\phi(v_1v_1^* + v_3v_3^*) - \frac{1}{l(l+1)}\phi(v_2v_2^*) \\ &= -\frac{1}{l(l+1)}\phi(v_1v_1^* + v_2v_2^* + v_3v_3^*) \\ &= -\frac{1}{l(l+1)} \end{aligned}$$

The above computation could have been simplified by using $SU_2(\mathbb{C})$ symmetry of the state ω_l to write

$$\begin{aligned} \omega_l(h_0^{xxx}) &= 3\omega_l(\sigma_z \otimes \sigma_z) \\ &= -\frac{3}{l(l+1)}\omega(v_2v_2^*) \\ &= -\frac{1}{l(l+1)} \end{aligned}$$

The map $l \rightarrow \omega_l(h_0^{xxx})$ increases strictly to zero as $l = \frac{1}{2}, 1, \frac{3}{2}, \dots$ increases to infinity and its minimum value at $l = \frac{1}{2}$ is $-\frac{4}{3}$. This shows that there is a unique limit point in ω_β as $\beta \rightarrow \infty$ and limiting value is $\omega_{\frac{1}{2}}$.

THEOREM 6.2. *Anti-ferromagnetic H^{XXX} model with $d = 3$ admits unique low temperature limiting ground state and the state is given by $\omega_{\frac{1}{2}}$.*

For any arbitrary odd values of d , for uniqueness of low temperature limit points, we need to prove that the mean energy of H^{xxx} i.e. $\omega_l(h_0^{xxx})$ gets minimised by a unique state ω_r for some $\frac{1}{2}$ -odd integer or integer spin r . Uniqueness of low temperature limiting states as well holds for any odd integer $d = 2s + 1$ if the function $l \rightarrow \omega_l(h_0^{xxx})$ is a monotonically increasing function in the variable l . For a possible quick proof, we verify using $SU_2(\mathbb{C})$ symmetry that

$$\begin{aligned} & \frac{1}{3}\omega_l(h_0^{xxx}) \\ &= \omega_l(\sigma_z \otimes \sigma_z) \\ &= \phi_l\left(\sum_{1 \leq k, k' \leq s} kk'((v_{k'}^* v_k - v_{2s+1-k'}^* v_{2s+1-k})(v_k v_k^* - v_{2s+1-k} v_{2s+1-k}^*)),\right) \end{aligned}$$

where ϕ_l is the normalized trace on \mathbb{M}_{2l+1} and $V^* = (v_k^*)$ is the Clebsch-Gordon isometry that intertwins two representations $\pi_l \otimes \pi_s$ and π_l of $SU_2(\mathbb{C})$. Note that there exists a unique intertwiner isometry provided $l \geq |l - s|$ i.e. $l \geq \frac{s}{2}$. So we leave it for future investigation as conjecture that low temperature limiting ground state of H_{XXX} for odd values of $d = 2s + 1$ is $\omega_{\frac{s}{2}}$.

For further illustration of our main result, we consider now well-studied AKLT model H^{AKLT} [1] for which

$$h_0^{aklt} = J\left(\frac{1}{3} + \frac{1}{2}(h_0^{xxx} + \frac{1}{3}(h_0^{xxx})^2)\right)$$

It is well known that $\omega_{\frac{1}{2}}$ is the unique ground state for H^{AKLT} with $\omega_{\frac{1}{2}}(h_0^{aklt}) = 0$.

We may as well compute

$$\begin{aligned} & \omega_l(\sigma_x^2 \otimes \sigma_x^2 + \sigma_y^2 \otimes \sigma_y^2 + \sigma_z^2 \otimes \sigma_z^2) \\ &= \frac{1}{4}\phi((v_1 v_1^* + v_1 v_3^* + 2v_2 v_2^* + v_3 v_1^* + v_3 v_3^*)(v_1^* v_1 + v_3^* v_1 + 2v_2^* v_2 + v_1^* v_3 + v_3^* v_3)) \\ &+ \frac{1}{4}\phi((v_1 v_1^* - v_1 v_3^* + 2v_2 v_2^* - v_3 v_1^* + v_3 v_3^*)(v_1^* v_1 - v_3^* v_1 + 2v_2^* v_2 - v_1^* v_3 + v_3^* v_3)) \\ &+ \phi((v_1 v_1^* + v_3 v_3^*)(v_1^* v_1 + v_3^* v_3)) \\ &= \frac{1}{4}\phi((I + v_1 v_3^* + v_2 v_2^* + v_3 v_1^*)(I + v_3^* v_1 + v_2^* v_2 + v_1^* v_3)) \\ &+ \frac{1}{4}\phi((I - v_1 v_3^* + v_2 v_2^* - v_3 v_1^*)(I - v_3^* v_1 + v_2^* v_2 - v_1^* v_3)) \\ &+ \phi((I - v_2 v_2^*)(I - v_2^* v_2)) \\ &= \frac{1}{2}\phi((I + v_2 v_2^*)(I + v_2^* v_2)) + \frac{1}{2}\phi((v_1 v_3^* + v_3 v_1^*)(v_3^* v_1 + v_1^* v_3)) \\ &+ \phi((I - v_2 v_2^*)(I - v_2^* v_2)) \\ &= \frac{3}{2}(1 + \phi(v_2^4)) - \phi(v_2^2) + \frac{1}{2}\phi((v_1 v_3^* + v_3 v_1^*)(v_3^* v_1 + v_1^* v_3)) \end{aligned}$$

We also compute

$$\begin{aligned} & l(l+1)(v_1 v_3^* + v_3 v_1^*) \\ &= \frac{1}{2}\{(\bar{\pi}_l(i\sigma_x) + i\bar{\pi}_l(i\sigma_y))(-\bar{\pi}_l(i\sigma_x) - i\bar{\pi}_l(i\sigma_y)) + (\bar{\pi}_l(i\sigma_x) - i\bar{\pi}_l(i\sigma_y))(-\bar{\pi}_l(i\sigma_x) + i\bar{\pi}_l(i\sigma_y))\} \\ &= \bar{\pi}_l(i\sigma_y)^2 - \bar{\pi}_l(i\sigma_x)^2 \end{aligned}$$

So

$$\begin{aligned} & l^2(l+1)^2\phi((v_1v_3^* + v_3v_1^*)(v_3^*v_1 + v_1^*v_3)) \\ &= \phi((\bar{\pi}_l(i\sigma_y)^2 - \bar{\pi}_l(i\sigma_x)^2)^2) \\ &= \phi(\bar{\pi}_l(i\sigma_y)^4 + \bar{\pi}_l(i\sigma_x)^4) - 2\phi(\bar{\pi}_l(i\sigma_x)^2\bar{\pi}_l(i\sigma_y)^2) \end{aligned}$$

(ϕ being tracial state on \mathcal{M} , $\phi(\bar{\pi}_l(i\sigma_y)^2\bar{\pi}_l(i\sigma_x)^2) = \phi(\bar{\pi}_l(i\sigma_x)^2\bar{\pi}_l(i\sigma_y)^2)$)

Using symmetry and tracial property of ϕ on \mathcal{M} , we write

$$\alpha_l = \phi(\bar{\pi}_l(i\sigma_x)^4) = \phi(\bar{\pi}_l(i\sigma_y)^4) = \phi(\bar{\pi}_l(i\sigma_z)^4)$$

and

$$\beta_l = \phi(\bar{\pi}_l(i\sigma_x)^2\bar{\pi}_l(i\sigma_y)^2) = \phi(\bar{\pi}_l(i\sigma_y)^2\bar{\pi}_l(i\sigma_z)^2) = \phi(\bar{\pi}_l(i\sigma_z)^2\bar{\pi}_l(i\sigma_x)^2)$$

We use the following identities:

$$\begin{aligned} l^2(l+1)^2 &= \phi((\bar{\pi}_l(i\sigma_x)^2 + \bar{\pi}_l(i\sigma_y)^2 + \bar{\pi}_l(i\sigma_z)^2)^2) \\ &= 3\alpha_l + 6\beta_l \end{aligned}$$

and

$$\alpha_l = \frac{1}{2l+1} \sum_{-l \leq m \leq l} m^4$$

to deduce

$$\begin{aligned} & \omega_l(\sigma_x^2 \otimes \sigma_x^2 + \sigma_y^2 \otimes \sigma_y^2 + \sigma_z^2 \otimes \sigma_z^2) \\ &= \frac{3}{2} - \frac{1}{3} + \frac{1}{l^2(l+1)^2} \left(\frac{3\alpha_l}{2} + \alpha_l - \beta_l \right) \\ &= \frac{7}{6} + \frac{1}{l^2(l+1)^2} \left(\frac{5\alpha_l}{2} - \frac{1}{6}(l^2(l+1)^2 - 3\alpha_l) \right) \\ &= 1 + \frac{3\alpha_l}{l^2(l+1)^2} \end{aligned}$$

We can as well use symmetry and tracial state property of ϕ to compute

$$\begin{aligned} & \omega_l((h_0^{xxx})^2) = 3\omega_l(\sigma_z^2 \otimes \sigma_z^2) + 6\omega_l(\sigma_z\sigma_x \otimes \sigma_z\sigma_x) \\ &= 3\phi((I - v_2^2)(1 - v_2^2)) + 3\phi((v_3v_1^* - v_1v_3^*)(v_1^*v_3 - v_3^*v_1)) \\ &= 1 + 3\phi(v_2^4) + 6\phi(v_2^4) \\ &= 1 + 9\phi(v_2^4) \end{aligned}$$

So

$$\begin{aligned} \omega_l(h_0^{aklt}) &= \frac{1}{3} + \frac{1}{2} \left(\frac{1}{3} + 3\phi(v_2^4) - \frac{1}{l(l+1)} \right) \\ &= \frac{1}{3} + \frac{1}{2} \left(\frac{1}{3} - \frac{1}{l(l+1)} + \frac{3}{l^2(l+1)^2(2l+1)} \sum_{-l \leq m \leq l} m^4 \right) \end{aligned}$$

This clearly shows that the unique ground state $\omega = \omega_{\frac{1}{2}}$ of H^{AKLT} [1] is also the low temperature limiting ground state of H^{XXX} . So we have the following well known result [1].

THEOREM 6.3. *Low temperature limit point as $\beta \rightarrow \infty$ of the unique temperature states ω_β for H^{AKLT} at inverse temperatures $\beta > 0$ is unique and its limiting value is also given by $\omega_{\frac{1}{2}}$.*

7. Disclaimer

Present manuscript being on pure mathematics, to my knowledge, I have no conflict of interest with scientific community.

8. Data Availability

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

REFERENCES

1. Affleck, L., Kennedy, T., Lieb, E.H. and Tasaki, H., "Valence Bond States in Isotropic Quantum Antiferromagnets", *Commun. Math. Phys.* 115, 477-528 (1988).
2. Affleck, L. and Lieb, E.H., "A Proof of Part of Haldane's Conjecture on Spin Chains", *Lett. Math. Phys.* 12, 57-69 (1986).
3. Araki, H., "Gibbs states of a one dimensional quantum lattice", *Comm. Math. Phys.* volume 14, 120-157 (1969).
4. Araki, H., "On uniqueness of KMS-states of one-dimensional quantum lattice systems", *Comm. Maths. Phys.* volume 44, 1-7 (1975).
5. Babujian, H.M.J., "Exact solution of the one-dimensional isotropic Heisenberg chain with arbitrary spins S", *Physics Letters*, 90A (1982) 479-482.
6. Bethe, H., "Zur Theorie der Metalle. I. Eigenwerte und Eigenfunktionen der linearen Atomkette, (On the theory of metals. I. Eigenvalues and eigenfunctions of the linear atom chain)", *Zeitschrift für Physik A*, Vol. 71, 205-226 (1931).
7. Bratteli, Ola, and Robinson, D.W., "Operator algebras and quantum statistical mechanics", I, Springer (1981).
8. Bratteli, Ola, and Robinson, D.W., "Operator algebras and quantum statistical mechanics", II, Springer (1981).
9. Bratteli, Ola, Jorgensen, Palle E.T., Kishimoto, Akitaka and Werner Reinhard F., "Pure states on \mathcal{O}_d ", *J.Operator Theory* volume 43, no-1, 97-143 (200).
10. Cuntz, J., "Simple C^* -algebras generated by isometries", *Comm. Math. Phys.* volume 57, no. 2, 173-185 (1977).
11. Dixmier, J.: *Von Neumann Algebras*, North-Holland, 1981.
12. Dyson, Freeman J.; Lieb, Elliott H. and Simon, B., "Barry Phase transitions in quantum spin systems with isotropic and non-isotropic interactions". *J. Statistical. Phys.* volume 18, no. 4, 335-383 (1978).
13. Fannes, M., Nachtergaele, B. and Werner, R., "Finitely correlated states on quantum spin chains" *Commun. Math. Phys.* volume 144, 443-490 (1992).
14. Fröhlich, J., Israel, R., Lieb, E.H., Simon, B., "Phase Transitions and Reflection Positivity-I general theory and long range lattice models", *Comm. Math. Phys.* 62, 1-34 (1978)
15. Ghosh, D. and Majumdar, C.K., "On Next - Nearest - Neighbour Interaction in Linear Chain". *J. Math. Phys.* 10, 1388 (1969).
16. Hall, B.C., "Lie algebras and representations: an elementary introduction", Springer (2003).
17. Høegh-Krohn, R., Landstad, M.B., Størmer, E., "Compact ergodic groups of automorphisms", *Ann. of Math. (2)* 114 (1981), no. 1, 75-86.
18. Kadison, Richard V., "Isometries of operator algebras", *Annals of Mathematics (2)*. volume 54, 325-338 (1951).
19. Kishimoto, A., "On uniqueness of KMS-states of one-dimensional quantum lattice systems", *Comm. Maths. Phys.* 47, 167-170 (1976).
20. Kirillov, A. A., "Elements of the theory of representations", Springer-Verlag, Berlin-New York, (1976).
21. Lieb, L.; Schultz, T.; Mattis, D., "Two soluble models of an anti-ferromagnetic chain", *Ann. Phys. (N.Y.)* 16, 407-466 (1961).
22. Matsui, T., "The split property and the symmetry breaking of the quantum spin chain", *Comm. Maths. Phys* volume-218, 293-416 (2001)
23. Mohari, A., "Pure inductive limit state and Kolmogorov's property. II", *J. Operator Theory* 72, no. 2, 387-404 (2014).
24. Mohari, A., "Translation invariant pure state on $\otimes_{k \in \mathbb{Z}} M_d^{(k)}(\mathbb{C})$ and Haag duality", *Complex Anal. Oper. Theory* 8, no. 3, 745-789 (2014).
25. Mohari, A., "Translation invariant pure state on $\mathcal{B} = \otimes_{k \in \mathbb{Z}} M_d^{(k)}(\mathbb{C})$ and its split property", *J. Math. Phys.* 56, 061701 (2015).
26. Mohari, A., "Spontaneous $SU_2(\mathbb{C})$ symmetry breaking in the ground states of quantum spin chain", *J. Math. Phys.* 59, 111701 (2018).
27. Mohari, A., "Translation invariant state and its mean entropy-I" arXiv 1705.11038.
28. Nachtergaele, B., "The spectral gap for some spin chains with discrete symmetry breaking", *Communications in Mathematical Physics* volume 175, pages 565-606 (1996)

29. Nachtergaele, B. and Sims, R., “Lieb-Robinson bounds and the exponential clustering theorem”, *Comm. Math. Phys.* 265, 119-130 (2006).
30. Ohya, M., Petz, D.: *Quantum entropy and its use*, Text and monograph in physics, Springer-Verlag.
31. Popescu, G., “Isometric dilations for infinite sequences of non-commutating operators”, *Trans. Amer. Math. Soc.* 316 no-2, 523-536 (1989)
32. Powers, Robert T., “Representations of uniformly hyper-finite algebras and their associated von Neumann rings”, *Annals of Mathematics*, 86, 138-171 (1967).
33. Ruelle, D., “*Statistical Mechanics*”, Benjamin, New York-Amsterdam (1969) .
34. Simon, B., “The statistical mechanics of lattice gases”, volume-1, Princeton series in physics (1993).
35. Takhtajan, L.A., The picture of low-lying excitations in the isotropic Heisenberg chain of arbitrary spins, *Phys. Lett.* 87A (1982) 479.

THE INSTITUTE OF MATHEMATICAL SCIENCES,, CIT CAMPUS, TARAMANI, CHENNAI-600113
E-mail: anilesh@imsc.res.in