

Stochastic COVID-19 Model Influenced by Non-Gaussian Noise

Almaz Tesfay^{1,a}, Anwar Zeb^b, Yu-Ming Chu^{c,*}, Ali Ahmadian^d, Soheil Salahshour^e

^aDepartment of Mathematics, Mekelle University, P.O.Box 231, Mekelle, Ethiopia

^bDepartment of Mathematics, COMSATS University Islamabad, Abbottabad Campus, Abbottabad, 22060, Khyber Pakhtunkhwa, Pakistan

^cDepartment of Mathematics, Huzhou University, Huzhou 313000, P. R. China

^dInstitute of IR 4.0, The National University of Malaysia, 43600 UKM, Bangi, Selangor, Malaysia

^eFaculty of Engineering and Natural Sciences, Bahcesehir University, Istanbul, Turkey

Abstract

Natural changes have been played a vital part within the consider of irresistible illnesses uncommonly coronavirus. It is important to think about the effect of irregular unsettling influence on epidemic models. In this work, first we present a SIR coronavirus mathematical model and consider that the contact rate is perturbed by Lévy noise. To protect and control diseases, the Lévy process should be used. Then, we discuss the dynamics of the deterministic model and for the global positive solution of the stochastic model, present the existence and uniqueness. Further, explore a few conditions for the termination and persistence of the infection. Then derive the basic reproduction number that determines the extinction and the persistence of the disease (infection). When the noise is huge or little, the numerical results appear that the COVID19 vanishes from the individuals on the chance that the reproduction number is less than one; though control the epidemic illnesses in the event that reproduction number is greater than one. Finally, for the demonstration of this phenomenon numerical simulations are presented.

Keywords: stochastic COVID-19, Brownian motion, Lévy noise, extinction, persistence.

2020 MSC: -Mathematics Subject Classification: 39A50, 45K05, 65N22.

1. Introduction

Coronaviruses are a huge family of infections that as a rule cause mild to direct upper-respiratory tract ailments. There are numerous kinds of coronaviruses that circulate among creatures such as pigs, cats and bats but now and then these infections can rise from creatures to humans causing infection. few of these infections caused a deadly infection like SARS coronavirus (SARS-COV). As of late, in December 2019, a modern coronavirus COVID-19 showed up within the seafood market of Wuhan, China. By Walk 2020, the World Health Organization (WHO) pronounced that this infection may be a worldwide widespread. COVID-19 spreads through beads of spit or release shape nose when an infected individual hacks or snuffles. In description of infectious diseases mathematical models have an important role both in theory and practice (see, for example, [6–8, 18, 34]). The development and investigation of models of this sort can offer

*Corresponding author

Email addresses: almaz.tesfay@mu.edu.et (Almaz Tesfay), anwar@cuiatd.edu.pk (Anwar Zeb), chuyuming@zjhu.edu (Yu-Ming Chu), ahmadian.hosseini@gmail.com (Ali Ahmadian), soheil.salahshour@eng.bau.edu.tr (Soheil Salahshour)

assistance us get it the instrument of the transmission as well as characteristics of illnesses, and thus, we are able propose successful techniques to foresee, prevent, and control infections, as well as to ensure population well-being. Up to presently, numerous mathematical models for irresistible illnesses defined by differential conditions have been constructed and analyzed to ponder the spreading of infections, for occurrence, [1–4]. As of late, numerical models for COVID-19 plague have pulled in the consideration of many mathematicians, scientists, disease transmission experts, drug specialists, chemists with numerous notable and imperative comes about see, [5–7, 4, 8] and references therein.

But environmental fluctuations have been played an important role in the study of disease specially coronavirus. Therefore, it is important to study the effect of random disturbance on epidemic models. In effect, disease infestation is highly stochastic, and stochastic noise can raise the probability of disease extinction in the early phase of epidemics. By applying an ODE model, we can get only a certain sample solution, whereas by using a stochastic differential equation (SDE) model, we can obtain the stochastic distribution of disease dynamics. Since epidemic models are environmental fluctuations. Recently, dynamic modeling of infectious diseases based on SDE has received considerable attention from researchers and academics. Up to now, many results on stochastic epidemic models have been investigated. Articles [9–17] have studied stochastic epidemic models influenced by Lévy noise. Motivated from the above researches, we assume that the contact rate is perturbed by Lévy noise. Lévy process should be introduced to protect and control diseases. Then the resultant model is given in Eq. (10); see Section 4. We want to derive the basic reproduction number that determines the extinction and the persistence of the disease (infection).

The rest of paper as follows: In section 2, we show the formulation of model. Section 4, is about the qualitative analysis of model presented in section 2. Section 4, is relate the numerical solution of proposed model. In section 7, lastly we give the conclusion of the whole paper.

2. Model formulation

The deterministic COVILD-19 model is

$$\begin{aligned}\frac{dS_t}{dt} &= \theta - \xi S_t I_t - \eta S_t + \rho R_t \\ \frac{dI_t}{dt} &= \xi S_t I_t - (\eta + \gamma) I_t \\ \frac{dR_t}{dt} &= \gamma I_t - (\eta + \rho) R_t\end{aligned}\tag{1}$$

where θ describes the enroll rate to the population that directly join only the susceptible class S_t , ξ stands for the contact rate mainly incidence rate at which susceptible class infectious class I_t , η out going rate of each class may be in the form of natural death, death due to corona or migration rate from each class, γ is the recovered rate and ρ shows the relapse rate at which the recovered class R_t moves to susceptible class.

2.1. Non-negativity of the solutions

This subsection deals about the positive solutions for future time with their respective initial conditions.

Theorem 1 (Positivity). *Let (S_0, I_0, R_0) be any initial data belonging to \mathbb{R}_+^3 and (S_t, I_t, R_t) be the solution corresponding to the initial data. Then, the set \mathbb{R}_+^3 is a positively invariant set of the model (1).*

Proof 1. *By letting $\alpha = \xi I_t$, first equation of (1) implies that*

$$\frac{dS_t}{dt} = \theta - \alpha S_t - \eta S_t + \rho R_t,$$

or

$$\frac{dS_t}{dt} \geq \theta - \alpha S_t - \eta S_t,$$

For all $J \in [0; +\infty[$, where solution of the system (1) exists, the above differential inequality has the solution as

$$\begin{aligned} \frac{dS_t}{dt} + \alpha S_t + \eta S_t &\geq \theta, \\ \frac{dS_t}{dt} + (\alpha + \eta) S_t &\geq \theta, \\ \frac{d}{dt} \left(S_t e^{\eta t + \int_0^t \alpha(s) ds} \right) &\geq \theta e^{\eta t + \int_0^t \alpha(s) ds}, \end{aligned}$$

it follows that

$$\begin{aligned} S_t e^{\eta t + \int_0^t \alpha(s) ds} - S_0 &\geq \int_0^t \theta e^{\eta t + \int_0^w \alpha(u) du} dw, \\ S_t e^{\eta t + \int_0^t \alpha(s) ds} &\geq S_0 + \int_0^t \theta e^{\eta w + \int_0^w \alpha(u) du} dw, \\ S_0 &\geq S_0 e^{-\eta t - \int_0^t \alpha(s) ds} + e^{-\eta t - \int_0^t \alpha(s) ds} \times \int_0^t \theta e^{\eta t + \int_0^w \alpha(u) du} dw > 0. \end{aligned}$$

Hence, S_t is positive for all values of t in the considered interval. As second equation of the system (1) implies that

$$\begin{aligned} \frac{dI_t}{dt} &\geq -(\eta + \gamma) I_t \\ \frac{dI_t}{I_t} &\geq -(\eta + \gamma) dt, \quad (I_t \neq 0). \end{aligned}$$

By integration, we have

$$\begin{aligned} \ln I_t &\geq -(\eta + \gamma)t + C, \\ I_t &\geq e^{-(\eta + \gamma)t + C}, \\ I_t &\geq C_1 e^{-(\eta + \gamma)t}, \end{aligned}$$

at $t = 0$,

$$I_t \geq I_0 e^{-(\eta + \gamma)t} \geq 0.$$

Hence, I_t is positive for all values of t . Also, from last equation of the system (1) we have

$$\frac{dR_t}{dt} = \gamma I_t - (\eta + \rho) R_t,$$

or

$$\frac{dR_t}{dt} \geq -(\eta + \rho) R_t,$$

can be written as

$$\frac{dR_t}{R_t} \geq -(\eta + \rho) dt, \quad (R_t \neq 0).$$

By integrating, we get

$$\begin{aligned}\ln R_t &\geq -(\eta + \rho)t + K, \\ R_t &\geq e^{-(\eta + \rho)t + K}, \\ R_t &\geq K_1 e^{-(\eta + \rho)t},\end{aligned}$$

at $t = 0$,

$$R_t \geq R_0 e^{-(\eta + \rho)0} \geq 0.$$

Hence, R_t also positive in the given interval.

The proof is complete.

3. Dynamics of the deterministic model

3.1. Free virus equilibrium point and reproduction number

Consider the following algebraic system for finding the equilibria of model (1),

$$\begin{aligned}\theta - \xi S_t I_t - \eta S_t + \rho R_t &= 0, \\ \xi S_t I_t - (\eta + \gamma) I_t &= 0, \\ \gamma I_t - (\eta + \rho) R_t &= 0.\end{aligned}\tag{2}$$

By some algebraic manipulations, we obtain two solutions of the system (2), one of them is the virus free equilibrium point is

$$(S_0, I_0, R_0) = \left(\frac{\theta}{\eta}, 0, 0\right).$$

Further, computing the reproduction number of model (1). Let $y = (I, S)$ and rewrite the model (1) for susceptible and infected classes as in the general form

$$\frac{dy}{dt} = \mathcal{F}(y) - \mathcal{V}(y),\tag{3}$$

where

$$\mathcal{F}(y) = \begin{pmatrix} \xi S_t I_t \\ 0 \end{pmatrix} \quad \mathcal{V}(y) = \begin{pmatrix} (\eta + \gamma) I_t \\ -\theta + \xi S_t I_t + \eta S_t - \rho R_t \end{pmatrix}.\tag{4}$$

Now Jacobian of above matrices at E_0 are

$$F(E_0) = \begin{pmatrix} \frac{\xi\theta}{\eta} & 0 \\ 0 & 0 \end{pmatrix} \quad V(E_0) = \begin{pmatrix} \eta + \gamma & 0 \\ \frac{-\xi\theta}{\eta^2(\gamma + \eta)} & \frac{1}{\eta} \end{pmatrix}.$$

Hence, the reproduction number of the model (1) can be determined by

$$\psi_0 = \rho(FV^{-1}) = \frac{\xi\theta}{\eta(\gamma + \eta)}.\tag{5}$$

3.2. Positive endemic equilibrium point

Theorem 2. For system (1), there exists a unique positive endemic equilibrium point E_+ , if $\psi_0 > 1$.

Proof 2. By some algebraic manipulations, the second solution of the system (2) is

$$\begin{aligned} S_+ &= \frac{\eta + \gamma}{\xi} \\ I_+ &= \frac{(\eta + \rho)(\eta + \gamma)}{\xi(\eta + \rho + \gamma)}(\psi_0 - 1) \\ R_+ &= \frac{\gamma(\eta + \gamma)}{\xi(\eta + \rho + \gamma)}(\psi_0 - 1). \end{aligned}$$

So

$$(S_+, I_+, R_+) = \left(\frac{\eta + \gamma}{\xi}, \frac{(\eta + \rho)(\eta + \gamma)}{\xi(\eta + \rho + \gamma)}(\psi_0 - 1), \frac{\gamma(\eta + \gamma)}{\xi(\eta + \rho + \gamma)}(\psi_0 - 1) \right)$$

It is clear from the values of S_+ , I_+ and R_+ that there exists a unique positive endemic equilibrium point E_+ , if $\psi_0 > 1$.

3.3. Stability analysis of the model

Theorem 3. The system (1) is locally stable related to virus free equilibrium point E_0 , if $\psi_0 < 1$ and unstable if $\psi_0 > 1$.

Proof 3. For local stability the Jacobian of system (1) is

$$J = \begin{pmatrix} -\eta - \xi I_t & -\xi S_t & \rho \\ \xi I_t & \xi S_t - (\eta + \gamma) & 0 \\ 0 & \gamma & -\rho - \eta \end{pmatrix}. \quad (6)$$

At E_0 , the Jacobian reaches to

$$J(E_0) = \begin{pmatrix} -\eta & -\frac{\xi\theta}{\eta} & \rho \\ 0 & \frac{\xi\theta}{\eta} - (\eta + \gamma) & 0 \\ 0 & \gamma & -\rho - \eta \end{pmatrix}, \quad (7)$$

which follows that eignvalues

$$\lambda_1 = -\eta < 0,$$

$$\lambda_3 = -\rho - \eta < 0,$$

and

$$\lambda_2 = \frac{\xi\theta}{\eta} - (\eta + \gamma) = (\eta + \gamma) \left(\frac{\xi\theta}{\eta(\eta + \gamma)} - 1 \right),$$

implies that $\lambda_2 < 0$, if $\psi_0 < 1$. So the system (1) is locally stable for $\psi_0 < 1$ and unstable for $\psi_0 > 1$. The proof is complete.

Theorem 4. If $\psi_0 > 1$, then the DFE point of the system (1) is globally stable.

Proof 4. For the proof of this theorem, first we construct the Lyapunov function $L(S_t, I_t)$ as,

$$L(S_t, I_t) = \left(S_t - \beta - \beta \ln \frac{S_t}{\beta} \right) + (I_t - 1 - \ln I_t). \quad (8)$$

Differentiating the equation (8) with respect to time, we have

$$\frac{d}{dt}L(S_t, I_t) = \left(1 - \frac{\beta}{S_t} \right) \frac{dS_t}{dt} + \left(1 - \frac{1}{I_t} \right) \frac{dI_t}{dt}, \quad (9)$$

by substitution the values of S_t , I_t and manipulating along the point E_0 , we have

$$\begin{aligned} \frac{d}{dt}(L(S_t, I_t)) &= \left(1 - \frac{\beta}{S_t} \right) (\theta - \eta S_t) - \xi S_t + (\eta + \gamma) \\ &= \left(\theta - \eta S_t - \frac{\beta\theta}{S_t} + \beta\eta \right) - \xi S_t + (\eta + \gamma) \\ &= \theta - \theta - \beta\eta + \beta\eta - \frac{\xi\theta}{\eta} + (\eta + \gamma) \\ &= -\frac{\xi\theta}{\eta} + (\eta + \gamma), \\ &= -(\eta + \gamma) \left(\frac{\xi\theta}{\eta(\eta + \gamma)} - 1 \right), \\ &= -(\eta + \gamma) (\psi_0 - 1), \\ &\leq 0 \quad \text{for } \psi_0 > 1. \end{aligned}$$

Therefore, if $\psi_0 > 1$, then $\frac{d}{dt}(L(S_t, I_t)) < 0$, which implies that the DFE point of the system (2) is globally stable for $\psi_0 > 1$.

Remark 1. An interesting mathematical problem is the analysis of stability of E_+ , but here, our main focus on the case $\psi_0 < 1$ to find effective strategy to prevent the disease.

4. Formulation and description of stochastic Covid-19 model

Motivated by the works [10, 18], in this paper, we will study the dynamics of stochastic Covid-19 epidemic model. We suppose that environmental fluctuations in the contact rate parameter ξ , mean that $\xi \rightarrow \xi + \int_{\mathbb{Z}} \epsilon(u) \bar{\pi}(dt, du)$ where $\epsilon(u)$ is a bounded function having the value $0; \epsilon(u) + 1$ on the intervals $1 \leq |u|$ or $1 > |u|$. $\bar{\pi}(t, du)$ is the compensated Poisson random measure satisfying $\bar{\pi}(t, du) = \pi(t, du) - \nu(du)dt$, with $\nu(\cdot)$ is a δ -finite measure on a measurable subset \mathbb{Z} of $(0, \infty)$ and $\infty > \nu(\mathbb{Z})$, and $\pi(t, du)$ is the independent Poisson random measure on $\mathbb{Z}^+ \times \mathbb{R} \setminus \{0\}$.

The following nonlinear stochastic Covid-19 system influenced with Non-Gaussian Noise is considered in this study

$$\begin{aligned} dS_t &= (\theta - \xi S_t I_t - \eta S_t + \rho R_t)dt - \int_{\mathbb{Z}} \epsilon(u) S_t^- I_t^- \bar{\pi}(dt, du) \\ dI_t &= (\xi S_t I_t - (\eta + \gamma) I_t)dt + \int_{\mathbb{Z}} \epsilon(u) S_t^- I_t^- \bar{\pi}(dt, du) \\ dR_t &= (\gamma I_t - (\eta + \rho) R_t)dt, \end{aligned} \quad (10)$$

where S_{t^-} and I_{t^-} are the left limit of S_t and I_t , respectively.

The initial conditions $(S(0), I(0), R(0)) \in \mathbb{R}_+^3$ are all non-negative. Since the first two equations of the stochastic covid-19 models (10) do not depend on the last (third) one, then we consider the following equation:

$$\begin{aligned} dS_t &= (\theta - \xi S_t I_t - \eta S_t + \rho R_t)dt - \int_{\mathbb{Z}} \epsilon(u) S_{t^-} I_{t^-} \bar{\pi}(dt, du) \\ dI_t &= (\xi S_t I_t - (\eta + \gamma) I_t)dt + \int_{\mathbb{Z}} \epsilon(u) S_{t^-} I_{t^-} \bar{\pi}(dt, du) \end{aligned} \quad (11)$$

4.1. Existence and uniqueness of the solution

Throughout this paper, we use the following notations and assumption.

$$\begin{aligned} \mathbb{R}_+^2 &:= \{x = (x_1, x_2) \in \mathbb{R}^2 : x_i \geq 0, i = 1, 2\}, \quad \mathbb{R}_+^3 := \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 : x_i \geq 0, i = 1, 2, 3\}, \\ \mathbb{R}_+ &= (0, \infty), \text{ fo all } x > 0, x - 1 - \ln x \geq 0 \text{ is true.} \end{aligned}$$

Assumption 1. $\epsilon(u)$ is a bounded function with $\epsilon(u) > -1 \quad u \in \mathbb{Z}$ of $(0, \infty)$, there exists $K > 0$ such that $\int_{\mathbb{Z}} (\ln(1 + \epsilon(u)))^2 \nu(du) < K$;

Remark 2. The above assumption implies that the intensities of jump noises are fine (not infinite).

Next, we will show that the stochastic Covid-19 model (11) has a unique, positive and global the solution $(S(0), I(0)) \in \mathbb{R}_+^2$.

Theorem 5. If Assumption (1) holds, then for any given initial value $(S(0), I(0)) \in \mathbb{R}_+^2$ and time $t \geq 0$, there is a unique global positive solution $(S(t), I(t)) \in \mathbb{R}_+^2$ of the model (11).

Proof 5. The stochastic covid-19 model (11) has locally Lipschitz continuous coefficient, thus the system has a unique local solution $(S(t), I(t)) \in \mathbb{R}_+^2$ on $t \in [0, \tau_e)$ where τ_e is the time for noise for explosion. In order to have a global solution, we need to show that $\tau_e = \infty$ almost surely. Assume that m_0 is very large positive number ($m_0 > 0$), then $(S(0), I(0)) \in [\frac{1}{m_0}, m_0]$. For every integer $m \geq m_0$, the stopping time is defined as:

$$\tau_e = \inf\{t \in [0, \tau_e) : \min(S(t), I(t), R(t)) \leq \frac{1}{m_0}, \text{ or } \max(S(t), I(t), R(t)) \geq m\}$$

As m goes to ∞ , τ_m increases. Define $\lim_{m \rightarrow \infty} \tau_m = \tau_\infty$ with $\tau_\infty \leq \tau_e$. If we can prove that $\tau_\infty = \infty$ almost surely, then $\tau_e = \infty$. If this is false, then there are two positive constants $T > 0$ and $\delta \in (0, 1)$ such that

$$\mathbb{P}\{\tau_\infty \leq T\} > \delta.$$

Thus there is $m_1 \geq m_0$ that satisfies

$$\mathbb{P}\{\tau_k \leq T\} \geq \delta, \quad m \geq m_1.$$

Construct a C^2 -function $V: \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ by

$$V(S, I) = (S - \beta - \beta \ln \frac{S}{\beta}) + (I - 1 - \ln I) \quad (12)$$

Applying Itô formula to Eq. (12), obtain

$$\begin{aligned}
dV(S, I) &= d\left[\left(S - \beta - \beta \ln \frac{S}{\beta}\right) + (I - 1 - \ln I)\right] \\
&= \left(1 - \frac{\beta}{S}\right)[(\theta - \xi S_t I_t - \eta S_t + \rho R_t)dt] - \left(1 - \frac{\beta}{S}\right) \int_{\mathbb{Z}} \epsilon(u) S_t I_t \bar{\pi}(dt, du) + \frac{(dS)^2}{2S^2} \\
&+ \left(1 - \frac{1}{I}\right)(\xi S_t I_t - (\eta + \gamma) I_t)dt + \left(1 - \frac{1}{I}\right) \int_{\mathbb{Z}} \epsilon(u) S_t I_t \bar{\pi}(dt, du) + \frac{(dI)^2}{2I^2} \\
&= \left(1 - \frac{\beta}{S}\right)[(\theta - \xi S_t I_t - \eta S_t + \rho R_t)dt] - \left(1 - \frac{\beta}{S}\right) \int_{\mathbb{Z}} \epsilon(u) S_t I_t \bar{\pi}(dt, du) - \beta \int_{\mathbb{Z}} [\ln(1 - \epsilon(u) I_t) + \epsilon(u) I_t] \nu((du) \\
&+ \left(1 - \frac{1}{I}\right)(\xi S_t I_t - (\eta + \gamma) I_t)dt + \left(1 - \frac{1}{I}\right) \int_{\mathbb{Z}} \epsilon(u) S_t I_t \bar{\pi}(dt, du) - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) S_t) - \epsilon(u) S_t] \nu((du) \\
&:= LV dt + \phi, \tag{13}
\end{aligned}$$

where

$$\phi = -\beta \int_{\mathbb{Z}} [\ln(1 - \epsilon(u) I_t) + \epsilon(u) S_t I_t] \bar{\pi}(dt, du) + \int_{\mathbb{Z}} [\epsilon(u) S_t I_t - \ln(1 + \epsilon(u) S_t)] \bar{\pi}(dt, du)$$

and L is the operator that acts on a function V . Here, $LV : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ is defined as follows:

$$\begin{aligned}
LV &= \left(1 - \frac{\beta}{S}\right)[(\theta - \xi S_t I_t - \eta S_t + \rho R_t)] - \beta \int_{\mathbb{Z}} [\ln(1 - \epsilon(u) I_t) + \epsilon(u) I_t] \nu((du) \\
&+ \left(1 - \frac{1}{I}\right)(\xi S_t I_t - (\eta + \gamma) I_t) - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) S_t) - \epsilon(u) S_t] \nu((du) \\
&= \theta - \xi S_t I_t - \eta S_t + \rho R_t + \xi S_t I_t - (\eta + \gamma) I_t + \frac{\beta \theta}{S_t} + \beta \xi I_t + \beta \eta + \beta \rho \frac{R_t}{S_t} - \xi S_t \\
&+ (\eta + \gamma) + \frac{1}{2} \int_{\mathbb{Z}} \epsilon^2(u) S_t^2 \nu(du) + \frac{1}{2} \int_{\mathbb{Z}} \epsilon^2(u) I_t^2 \nu(du) \\
&\leq \theta - (\eta + \gamma) I_t + \beta \xi I_t + (\eta + \gamma) + \beta \eta + \frac{1}{2} \int_{\mathbb{Z}} \epsilon^2(u) S_t^2 \nu(du) + \frac{1}{2} \int_{\mathbb{Z}} \epsilon^2(u) I_t^2 \nu(du) \tag{14}
\end{aligned}$$

By assuming $\beta = \frac{\eta + \gamma}{\xi}$, Eq. (14) becomes

$$\begin{aligned}
LV &\leq \theta + (\eta + \gamma) + \beta \eta - \beta \int_{\mathbb{Z}} [\ln(1 - \epsilon(u) I_t) + \epsilon(u) I_t] \nu((du) - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) S_t) - \epsilon(u) S_t] \nu((du) \\
&:= Q > 0, \tag{15}
\end{aligned}$$

we obtain

$$dV(S_t, I_t) \leq Q dt + \phi$$

For more details; see [14], Theorem 2.1, [9], Theorem 2.1, and [18]. The proof is therefore completed. \square

4.2. Extinction and persistence of the Covid-19 disease with Lévy jumps

In this section, we will find sufficient conditions for extinction and persistence of the disease that can give us new insights in understanding the control of Covid-19.

4.2.1. Reproduction number for stochastic covid-19 model

The existence of virus-free equilibrium and endemic equilibrium are computed after we have the basic reproductive number. In this section, we will find the basic reproductive number of stochastic counterpart that helps to determine extinction and persistence of the coronavirus (disease). Consider the infected class of (11), i.e.

$$dI_t = (\xi S_t I_t - (\eta + \gamma) I_t) dt + \int_{\mathbb{Z}} \epsilon(u) S_t I_t \bar{\pi}(dt, du)$$

Choosing a twice differential function Y_t and applying Ito's formula into Y_t , we can obtain stochastic basic reproduction number. Set $Y_t = \ln I_t$.

$$\begin{aligned} dY_t &= d \ln I_t \\ &= \frac{1}{I_t} dI_t - \frac{1}{2I_t^2} (dI_t)^2 \\ &= \frac{1}{I_t} \left[(\xi S_t I_t - (\eta + \gamma) I_t) dt + \int_{\mathbb{Z}} \epsilon(u) S_t I_t \bar{\pi}(dt, du) \right] \\ &\quad - \frac{1}{2I_t^2} \left[(\xi S_t I_t - (\eta + \gamma) I_t) dt + \int_{\mathbb{Z}} \epsilon(u) S_t I_t \bar{\pi}(dt, du) \right]^2 \\ &= (\xi S_t - (\eta + \gamma)) dt + \int_{\mathbb{Z}} \ln(1 + \epsilon(u)) S_t \bar{\pi}(dt, du) - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) S_t) - \epsilon(u) S_t] \nu((du) dt \\ &= \left[\xi S_t - (\eta + \gamma) - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) S_t) - \epsilon(u) S_t] \nu((du) \right] dt + \int_{\mathbb{Z}} \ln(1 + \epsilon(u)) S_t \bar{\pi}(dt, du) \end{aligned} \quad (16)$$

In fact that $I = R = 0$, then $\theta - \eta S = 0$, that gives $S = \frac{\theta}{\eta}$. Set

$A = \xi S_t - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) S_t) - \epsilon(u) S_t] \nu((du) = \xi \frac{\theta}{\eta} - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) \frac{\theta}{\eta}) - \epsilon(u) \frac{\theta}{\eta}] \nu((du) = \xi \frac{\theta}{\eta} - \varphi$ and $B = \eta + \gamma$ where $\varphi = \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) \frac{\theta}{\eta}) - \epsilon(u) \frac{\theta}{\eta}] \nu((du) \leq \int_{\mathbb{Z}} [\ln(1 + \epsilon(u) \frac{\theta}{\eta})] \nu((du) \leq K$ from Assumption 1.

$$AB^{-1} = \left(\xi \frac{\theta}{\eta} - \varphi \right) \frac{1}{\eta + \gamma} = \frac{\xi \theta}{\eta(\eta + \gamma)} - \frac{\varphi}{\eta + \gamma}$$

Therefore, the reproduction number for stochastic Covid-19 model (10) is

$$\psi = \psi_0 - \frac{\varphi}{\eta + \gamma} \quad (17)$$

where $\psi_0 = \frac{\xi \theta}{\eta(\eta + \gamma)}$ is the reproduction number for deterministic Covid-19 model (1).

Remark 3. From Eq. (17), we can observe that $\psi < \psi_0$, which shows the stochastic approach is more realistic than the deterministic approach.

4.2.2. Extinction of the Covid-19 disease

Theorem 6. Assumption 1 and $\psi < 1$ hold. Then for any initial condition $(S_0, I_0) \in \mathbb{R}^2$, the solution $(S_t, I_t) \in \mathbb{R}^2$ of the stochastic COVID-19 model (4) satisfies

$$\lim_{t \rightarrow \infty} \sup \frac{\ln I_t}{t} \leq (\eta + \gamma)[\psi - 1] < 0, \quad a.s.$$




Figure 1: This Figure shows the effect of the parameters on the basic reproductive number of the stochastic Covid-19 model(10) (a) $\theta = 0.0073$, $\xi = 0.003$, $\eta = 0.001$, $\rho = 0.01$, $\gamma = 0.02$, with vary $\epsilon(u)$ (b) $\xi = 0.003$, $\eta = 0.001$, $\rho = 0.01$, $\gamma = 0.02$, $\epsilon(u) = 0.001$ and vary value of θ . (c) $\theta = 0.0073$, $\xi = 0.003$, $\eta = 0.001$, $\rho = 0.01$, $\gamma = 0.02$, , and non-Gaussian noise intensity $\epsilon(u) = 0.001$. (d) $\theta = 0.0073$, $\xi = 0.003$, $\eta = 0.001$, $\rho = 0.01$, $\gamma = 0.02$, $\epsilon(u) = 0.001$ with vary Ψ_0 .

Proof 6. Integrating both sides of Eq. (16) from 0 to t , becomes

$$\ln I_t = \ln I_0 + \int_0^t \left[\xi S_t - (\eta + \gamma) - \int_{\mathbb{Z}} [\ln(1 + \epsilon(u)S_t) - \epsilon(u)S_t] \nu((du)) \right] dt + \int_0^t \int_{\mathbb{Z}} \ln(1 + \epsilon(u)) S_t \bar{\pi}(dt, du) \quad (18)$$

Evaluating Eq. (18) at the disease free equilibrium, we obtain

$$\begin{aligned} \ln I_t &= \ln I_0 + \int_0^t \left[\xi \frac{\theta}{\eta} - (\eta + \gamma) - \int_{\mathbb{Z}} \left[\ln(1 + \epsilon(u) \frac{\theta}{\eta}) - \epsilon(u) \frac{\theta}{\eta} \right] \nu((du)) \right] dt + \int_0^t \int_{\mathbb{Z}} \ln(1 + \epsilon(u)) \frac{\theta}{\eta} \bar{\pi}(dt, du) \\ &\leq \ln I_0 + \int_0^t \xi \frac{\theta}{\eta} - (\eta + \gamma) - \varphi + \int_0^t \int_{\mathbb{Z}} \ln(1 + \epsilon(u)) \frac{\theta}{\eta} \bar{\pi}(dt, du) \end{aligned} \quad (19)$$

Multiplying both side of Eq. (19) by $\frac{1}{t}$

$$\frac{1}{t} \ln I_t \leq \frac{1}{t} \ln I_0 + \frac{1}{t} \int_0^t (\xi \frac{\theta}{\eta} - (\eta + \gamma) - \varphi) ds + \frac{1}{t} \int_0^t \int_{\mathbb{Z}} \ln(1 + \epsilon(u)) \frac{\theta}{\eta} \bar{\pi}(dt, du) ds \quad (20)$$

Using the large number theorem and applying $\lim_{t \rightarrow \infty} \sup$ to Eq. (20) and $\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \int_{\mathbb{Z}} \ln(1 + \epsilon(u)) \frac{\theta}{\eta} \bar{\pi}(dt, du) ds = 0$, get

$$\begin{aligned} \lim_{t \rightarrow \infty} \sup \frac{1}{t} \ln I_t &\leq \left(\xi \frac{\theta}{\eta} - (\eta + \gamma) - \varphi \right) \\ &= (\eta + \gamma) \left[\xi \frac{\theta}{\eta(\eta + \gamma)} - \frac{\varphi}{\eta + \gamma} - 1 \right] \\ &= (\eta + \gamma) [\psi - 1] < 0 \quad \text{iff} \quad \psi < 1. \end{aligned} \quad (21)$$

Thus if $\psi < 1$, then the virus-free equilibrium point is locally asymptotically stable. In other words

$$\lim_{t \rightarrow \infty} \frac{I_t}{t} = 0.$$

The proof is completed. \square

4.3. Persistence of the Covid-19 disease

In this section, we discuss about the persistence of stochastic model

Theorem 7. *Assumption 1 and $\psi > 1$ hold. Then for any initial condition $(S_0, I_0, R_0) \in \mathbb{R}^3$, the solution $(S_t, I_t, R_t) \in \mathbb{R}^3$ of the stochastic COVID-19 model (4) satisfies*

$$\lim_{t \rightarrow \infty} \langle S \rangle_t = S^* > 0, \quad \lim_{t \rightarrow \infty} \langle I \rangle_t = I^* > 0, \quad \text{and} \quad \lim_{t \rightarrow \infty} \langle R \rangle_t = R^* > a.s.$$

where

$$S^* = \frac{\theta}{\eta} - \left(\frac{\eta + \gamma + \rho}{\eta + \rho} \right) (\eta + \gamma) [\psi - 1], \quad I^* = (\eta + \gamma) [\psi - 1] \quad R^* = \frac{\gamma}{\eta + \rho} (\eta + \gamma) [\psi - 1]$$

Proof 7. *Integrating both sides of stochastic Covid-19 model (10), add all equations, and multiply the result by $\frac{1}{t}$, we obtain*

$$\begin{aligned} \frac{S_t - S_0}{t} + \frac{I_t - I_0}{t} + \frac{R_t - R_0}{t} &= \theta - \xi \langle S \rangle_t + \langle I \rangle_t - \eta \langle S \rangle_t + \rho \langle R \rangle_t - \frac{1}{t} \int_0^t \int_{\mathbb{Z}} \epsilon(u) S_r I_r \bar{\pi}(dr, du) dr \\ &\quad + \xi \langle S \rangle_t + \langle I \rangle_t - (\eta + \gamma) \langle I \rangle_t + \frac{1}{t} \int_0^t \int_{\mathbb{Z}} \epsilon(u) S_r I_r \bar{\pi}(dr, du) dr \\ &\quad + \gamma \langle I \rangle_t - (\eta + \rho) \langle R \rangle_t \\ &= \theta - \eta \langle S \rangle_t - \left((\eta + \gamma) - \frac{\rho \gamma}{\eta + \rho} \right) \langle I \rangle_t \\ &= \theta - \eta \langle S \rangle_t - \left(\frac{(\eta + \gamma)(\eta + \rho) - \gamma \rho}{\eta + \rho} \right) \langle I \rangle_t \end{aligned} \quad (22)$$

Now, let us solve $\langle S \rangle_t$ from Eq. (22)

$$\langle S \rangle_t = \frac{1}{\eta} \left[\frac{S_t - S_0}{t} + \frac{I_t - I_0}{t} + \frac{\rho}{\rho + \eta} \frac{R_t - R_0}{t} \right] + \frac{\theta}{\eta} - \frac{1}{\eta} \left(\frac{(\eta + \gamma)(\eta + \rho) - \gamma \rho}{\eta + \rho} \right) \langle I \rangle_t \quad (23)$$

Finding $\lim_{t \rightarrow \infty}$ to Eq. (23)

$$\lim_{t \rightarrow \infty} \langle S \rangle_t = \frac{\theta}{\eta} - \frac{1}{\eta} \left(\frac{(\eta + \gamma)(\eta + \rho) - \gamma \rho}{\eta + \rho} \right) \langle I \rangle_t \quad (24)$$

Substituting Eq. (21) in to Eq. (24), yields

$$\begin{aligned} \lim_{t \rightarrow \infty} \langle S \rangle_t &= \frac{\theta}{\eta} - \frac{1}{\eta} \left(\frac{(\eta + \gamma)(\eta + \rho) - \gamma \rho}{\eta + \rho} \right) (\eta + \gamma) [\psi - 1] \\ &= \frac{\theta}{\eta} - \left(\frac{\eta + \gamma + \rho}{\eta + \rho} \right) (\eta + \gamma) [\psi - 1] \end{aligned} \quad (25)$$

Next apply $\lim_{t \rightarrow \infty}$ in to the following Equation

$$\frac{R_t - R_0}{t} = \gamma \langle I \rangle_t - (\eta + \rho) \langle R \rangle_t \quad (26)$$

Replace $\langle I \rangle_t$ in Eq. (21), gives

$$\lim_{t \rightarrow \infty} \langle R \rangle_t = \frac{\gamma}{\eta + \rho} (\eta + \gamma) [\psi - 1] \quad (27)$$

The proof of the Theorem is completed. \square




Figure 2: The numerical simulation of model (4) (a) The graph of susceptible. (b) The graph of infected people. Parameters $\theta = 0.0073$, $\xi = 0.003$, $\eta = 0.001$, $\rho = 0.01$, $\gamma = 0.02$, $\psi = 0.9994 < 1$, and non-Gaussian noise intensity $\epsilon(u) = 0.001$.

Fact 1. *The biological indication [19] of Theorem 7 is that a very small flow of infectious individuals will not generate an outbreak of the disease unless if $\psi > 1$.*

5. Numerical experiments

Numerical solutions of system are very useful in the study of epidemic models. This section shows the numerical findings of our model that helps to demonstrate how affect the parameters of the deterministic




Figure 3: The numerical simulation of model (4) (a) The graph of susceptible. (b) The graph of infected people. Parameters $\theta = 0.0073$, $\xi = 0.0033$, $\eta = 0.001$, $\rho = 0.01$, $\gamma = 0.02$, $\psi = 1.1042 > 1$, and non-Gaussian noise intensity $\epsilon(u) = 0.001$.

model (1) and the non-Gaussian noise intensity of the stochastic model (10). We give some numerical experiments to illustrate the extinction and persistence of the new-coronavirus called Covid-19 in deterministic model and corresponding stochastic system for comparison.

We fix the values of the parameters as follow: $\theta = 0.0073$, $\xi = 0.003$, $\eta = 0.001$, $\rho = 0.01$, $\gamma = 0.02$, and non-Gaussian noise intensity $\epsilon(u) = 0.001$ in the deterministic model (2) and stochastic model (11). Then the system (1), becomes

$$\begin{aligned} \frac{dS_t}{dt} &= 0.0073 - 0.003 S_t I_t - 0.001 S_t + 0.01 R_t \\ \frac{dI_t}{dt} &= 0.003 S_t I_t - (0.001 + 0.02) I_t \\ \frac{dR_t}{dt} &= 0.02 I_t - (0.001 + 0.01) R_t \end{aligned}$$

Using the simple calculation, we get $\psi_0 = 1.0429 > 1$ and $\psi = 0.9994 < 1$, then according to Theorem 6 model (1), diseases will prevail in a long run, and from Theorem 7 model (11) implies the disease goes to extinction.

From Figure 1, we observed that if the noise intensity is large, the extinction of the covid-19 happen. For $\theta \in (0, 0.0073)$, $\xi \in (0, 0.003)$, and $\psi_0 \in (0, 1.0435)$, the coronavirus-2 (Covid-19) goes to extinction (die out).

The effects of non-Gaussian noise (jumps) to system (11) is plotted on Fig. 2. $\psi_0 > 1$ shows that the disease prevails in the deterministic model case, but in the stochastic model case, $\psi < 1$ that implies the disease die out. This result indicates that the stochastic approach is more realistic than the deterministic approach. In other words, from this result we can see that $\psi_0 > \psi$ because of the effect of the jump noise.

Next, let us keep the parameters the same as shown in Figure 2, but $\xi = 0.0033$ take different value.

Figure 3 is plotted based on Theorem 7, i.e., for $\psi > 1$, then

$$\begin{aligned} \lim_{t \rightarrow \infty} \langle S \rangle_t &= S^* = 7.2977 > 0, \\ \lim_{t \rightarrow \infty} \langle I \rangle_t &= I^* = 0.0022 > 0, \\ \text{and } \lim_{t \rightarrow \infty} \langle R \rangle_t &= R^* = 0.0015 > 0 \quad \text{a.s.} \end{aligned}$$

From this Figure we can see that I_t is persistent.

6. Conclusion

This paper investigates the dynamic behavior of a Covid-19 epidemic model with non-Gaussian noise. We discussed the dynamics of deterministic form of the model and further prove the existence and uniqueness of non-negative global solution of the stochastic system (10). The obtained findings improve some previous study's mentioned in the Introduction which is shown graphically in this manuscript. In this regard, we presented sufficient conditions for the extinction of the disease of Covid-19 (10) if $\psi < 1$, which can help in the control of Covid-19.

References

- [1] G. Bocharov, V. Volpert, B. Ludewig, and A. Meyerhans, *Mathematical immunology of virus infections*. Springer, 2018.
- [2] J. W. F. Brauer, P. van den Driessche, "Mathematical epidemiology," *Springer Berlin Heidelberg*, 2008.
- [3] F. Brauer, "Mathematical epidemiology: Past, present, and future," *Infectious Disease Modelling 2(2017) 113-127*, 2017.
- [4] M. Martcheva, "An introduction to mathematical epidemiology," *Springer Science+Business Media New York.*, 2015.
- [5] S. Çakan, "Dynamic analysis of a mathematical model with health care capacity for covid-19 pandemic," *Chaos, Solitons & Fractals*, vol. 139, p. 110033, 2020.
- [6] M. Higazy, "Novel fractional order sidarthe mathematical model of covid-19 pandemic," *Chaos, Solitons & Fractals Volume 138,110007*, 2020.
- [7] S. Kumar, J. Cao, and M. Abdel-Aty, "A novel mathematical approach of covid-19 with non-singular fractional derivative," *Chaos, Solitons & Fractals*, vol. 139, p. 110048, 2020.
- [8] Y. Li, Y. Chen, and I. Podlubny, "Mittag-leffler stability of fractional order nonlinear dynamic systems," *Automatica*, vol. 45, no. 8, pp. 1965-1969, 2009.
- [9] Y. Liu, Y. Zhang, and Q. Wang, "A stochastic sir epidemic model with lévy jump and media coverage," *Advances in Difference Equations*, vol. 2020, no. 1, 2020.
- [10] A. Tesfay, T. Saeed, A. Zeb, D. Tesfay, A. Khalaf, and J. Brannan, "Dynamics of a stochastic covid-19 epidemic model with jump-diffusion," *Advances in Difference Equations*, vol. 2021, no. 1, pp. 1-19, 2021.
- [11] A. Tesfay, D. Tesfay, A. Khalaf, and J. Brannan, "Mean exit time and escape probability for the stochastic logistic growth model with multiplicative α -stable lévy noise," *Stochastics and Dynamics*, vol. 21, no. 04, p. 2150016, 2021.
- [12] A. D. Khalaf, A. Tesfay, and X. Wang, "Impulsive stochastic volterra integral equations driven by lévy noise," *Bulletin of the Iranian Mathematical Society*, pp. 1-19, 2020.
- [13] R. Cai, Y. Liu, J. Duan, and A. T. Abebe, "State transitions in the morris-lecar model under stable lévy noise," *arXiv preprint arXiv:1906.07613*, 2019.
- [14] Y. Zhou and W. Zhang, "Threshold of a stochastic sir epidemic model with lévy jumps," *Physica A: Statistical Mechanics and Its Applications*, vol. 446, pp. 204-216, 2016.
- [15] A. Tesfay, D. Tesfay, J. Brannan, and J. Duan, "A logistic-harvest model with allee effect under multiplicative noise," *Stochastics and Dynamics*, p. 2150044, 2021.
- [16] A. Zeb, S. Kumar, A. Tesfay, and A. Kumar, "A stability analysis on a smoking model with stochastic perturbation," *International Journal of Numerical Methods for Heat & Fluid Flow*, 2021.
- [17] A. Tesfay, D. Tesfay, S. Yuan, J. Brannan, and J. Duan, "Stochastic bifurcation in single-species model induced by α -stable lévy noise," *Journal of Statistical Mechanics: Theory and Experiment*, vol. 2021, no. 10, p. 103403, 2021.
- [18] Z. Zhang, A. Zeb, S. Hussain, and E. Alzahrani, "Dynamics of covid-19 mathematical model with stochastic perturbation," *Advances in Difference Equations*, vol. 2020, no. 1, pp. 1-12, 2020.
- [19] S. D. A, G. G. K. A, H. A. B, A. A. A. A. C. D, and I. D. B, "Mathematical modeling, analysis and numerical simulation of the covid-19 transmission with mitigation of control strategies used in cameroon," *Chaos, Solitons & Fractals*, vol. 139, 2020.