

# A FIXED POINT THEOREM IN $B(H, \ell_\infty)$

ANDRZEJ WIŚNICKI

ABSTRACT. We show that if  $X$  is a complete metric space with uniform relative normal structure and  $G$  is a subgroup of the isometry group of  $X$  with bounded orbits, then there is a point in  $X$  fixed by every isometry in  $G$ . As a corollary, we obtain a theorem of U. Lang (2013) concerning injective metric spaces. A few applications of this theorem are given to the problems of inner derivations. In particular, we show that if  $L_1(\mu)$  is an essential Banach  $L_1(G)$ -bimodule, then any continuous derivation  $\delta : L_1(G) \rightarrow L_\infty(\mu)$  is inner. This extends a theorem of B. E. Johnson (1991) asserting that the convolution algebra  $L_1(G)$  is weakly amenable if  $G$  is a locally compact group.

## 1. INTRODUCTION

The writing of this paper has been animated by a fixed point theorem of Bader, Gelfand and Monod in [2]—there is a point fixed by every isometry of an  $L$ -embedded Banach space that preserves a given bounded set. One of its consequences is the optimal solution to the following *derivation problem*: if  $G$  is a locally compact group, then any derivation from the convolution algebra  $L_1(G)$  to  $M(G)$  is *inner*. The problem was studied since 1960s and proved for the first time by Losert [24]. An ingenious proof in [2] relies on the Ryll-Nardzewski theorem. Another consequence of this theorem, observed by Uffe Haagerup, is a direct proof of *weak amenability* of all  $C^*$ -algebras which was proved in [12] using the Grothendieck–Haagerup–Pisier inequality.

There is a counterpart of the Bader–Gelfand–Monod theorem in *injective* Banach spaces  $L_\infty$  (see Section 2), and it is natural to ask about the noncommutative analogue of that result. The relevance of such a question is particularly apparent if we analyse the proof of Corollary C in [2]. The central idea behind the proofs of both the derivation problem and weak amenability of  $C^*$ -algebras there, is to construct a group of affine isometries with bounded orbits acting on the predual of an appropriate von Neumann algebra and then applying [2, Theorem A]. We can use a similar strategy to study derivations into injective Banach bimodules and, more general, into Banach bimodules with *uniform relative normal structure* (see Section 2 for the definition), based this time on Theorem 2.3. In particular, we show that if  $L_1(\mu)$  is an essential Banach  $L_1(G)$ -bimodule, then any continuous derivation  $\delta : L_1(G) \rightarrow L_\infty(\mu)$  is inner, where  $\mu$  is a  $\sigma$ -finite measure on  $(E, \Sigma)$  or a Radon measure on a locally compact space. This extends a theorem of

Johnson (see [16, Theorem]) asserting that the convolution algebra  $L_1(G)$  is weakly amenable if  $G$  is a locally compact group.

In Section 3 we apply Theorem 2.3 in the case of  $B(H, \ell_\infty(\Gamma))$ , the Banach space of bounded operators from a Hilbert space  $H$  into  $\ell_\infty(\Gamma)$  for some  $\Gamma$ . It can throw some new light on the *Kadison similarity problem*. One of the equivalent formulations of this problem concerns inner derivations into Banach modules and translates into the existence or non-existence of a fixed point for the appropriate group of affine isometries acting on  $B(H)$ , the von Neumann algebra of bounded linear operators on a Hilbert space  $H$ . Notice that  $B(H)$  is injective in the category of operator spaces and completely contractive maps. Though it is not injective as a Banach space, an embedding of  $B(H)$  into a Banach space  $B(H, \ell_\infty(\Gamma))$  with uniform relative normal structure seems to lead to something new.

Our strategy is as follows. Let  $\mathcal{A}$  be a unital  $C^*$ -algebra,  $M$  a Banach bimodule over  $\mathcal{A}$  and  $\delta : \mathcal{A} \rightarrow M$  a derivation:  $\delta(ab) = \delta(a)b + a\delta(b)$ . If  $G$  is a group of unitary elements of  $\mathcal{A}$ , then  $\alpha : G \times M \rightarrow M$  defined by  $\alpha(g)x = gxg^{-1} + \delta(g)g^{-1}$ ,  $g \in G, x \in M$ , is an affine and isometric action of  $G$  on  $M$ . It follows from Theorem 2.3 that if  $M$  is a Banach bimodule with uniform relative normal structure, then there is  $x \in M$  such that  $\alpha(g)x = x$ , and consequently,  $\delta(g) = xg - gx$ . Since every element of  $\mathcal{A}$  is a linear combination of three unitaries, we deduce that each derivation  $\delta : \mathcal{A} \rightarrow M$  is inner. The argument works also for some group algebras, see Theorem 2.7. The case of the convolution algebra  $L_1(G)$  and  $M = L_\infty(G)$ , where  $G$  is a locally compact group was studied by Johnson [15, 16]. Let us move on now to the Kadison similarity problem, which is equivalent to the derivation problem for derivations  $\delta : B(H) \supset \mathcal{A} \rightarrow B(H)$  (see [20]). Let us embed a Hilbert space  $H$  isometrically in an injective space  $\ell_\infty(\Gamma)$  and extend each unitary  $g : H \rightarrow H$  to an isometry  $\tilde{g} : \ell_\infty(\Gamma) \rightarrow \ell_\infty(\Gamma)$ . Now, consider the group of isometries  $\alpha_g : B(H, \ell_\infty(\Gamma)) \rightarrow B(H, \ell_\infty(\Gamma))$  defined by  $\alpha_g x = \tilde{g}xg^{-1} + \delta(g)g^{-1}$ . It follows from Theorem 3.1 that there is a bounded linear operator  $T : H \rightarrow \ell_\infty(\Gamma)$  such that  $\delta(g) = Tg - \tilde{g}T$ . Note that in fact a similar result holds for derivations from  $\ell_1(G)$  to  $B(H)$  on which  $G$  acts by isometries. However, in the case of  $C^*$ -algebras our factorization appears to be more effective. For example, properties of the group homomorphism  $g \mapsto \tilde{g}$  could be useful in studying complete boundedness of  $\delta$ . Recall that Christensen's theorem [7] asserts that a derivation  $\delta : \mathcal{A} \rightarrow B(H)$  is inner if and only if it is completely bounded whenever  $\mathcal{A}$  is a unital  $C^*$ -algebra. Thus we can reduce the Kadison similarity problem to the problem of complete boundedness of the appropriate homomorphism from  $\mathcal{A}$  into  $B(Y)$  for some space  $Y$  isomorphic to a Hilbert space (see Remark 3.4).

## 2. SPACES WITH UNIFORM RELATIVE NORMAL STRUCTURE

Let  $\mathcal{A}(X)$  denote the family of admissible subsets of a metric space  $X$ , i.e., the intersections of closed balls of  $X$ . The following definition, originated

in the works of Brodskii and Mil'man [4], and Kirk [21], was introduced by Soardi [30] in the case of Banach spaces (see also [19]).

**Definition 2.1.** A metric space  $X$  is said to have uniform relative normal structure if there exists  $c < 1$  such that for each nonempty and bounded set  $M \in \mathcal{A}(X)$  there exists  $z_M \in X$  such that

- $d(x, z_M) \leq c \operatorname{diam} M$  for every  $x \in M$ ;
- if for some  $y \in X$ ,  $M \subset B(y, c \operatorname{diam} M)$ , then

$$d(y, z_M) \leq c \operatorname{diam} M,$$

where  $B(y, r)$  denotes the closed ball centered at  $y$  and of radius  $r$ .

Examples of spaces with uniform relative normal structure include Hilbert spaces, (1-)injective spaces and their closed balls (see [30]). It was proved in [30] that if  $K$  is a nonempty weak\*-closed subset with uniform relative normal structure of a dual Banach space, and  $T : K \rightarrow K$  is nonexpansive and leaves invariant a weak\*-compact subset of  $K$ , then  $K$  contains a fixed point for  $T$  (see also [10, 19] for some generalizations in metric spaces). Lau [23] used Soardi's arguments to show the following fixed point theorem for isometries.

**Theorem 2.2** (A.T.-M. Lau). *Let  $X$  be a Banach space and let  $K$  be a nonempty bounded closed subset of  $X$  with uniform relative normal structure. Then there is  $z \in K$  such that  $Tz = z$  for each isometry  $T$  from  $K$  onto  $K$ .*

A natural question arises whether it is possible to extend the above results for mappings with bounded orbits, and Prus (see [11]) constructed an isometry  $T : \ell_\infty \rightarrow \ell_\infty$  with bounded orbits but without fixed points. However, it turns out that in the case of invertible isometries the situation is different, and the Soardi–Lau method leaves some room for improvement.

**Theorem 2.3.** *Suppose that  $X$  is a complete metric space with uniform relative normal structure. If  $G$  is a group of isometries  $g : X \rightarrow X$  with bounded orbits, then there exists  $x \in X$  such that  $gx = x$  for every  $g \in G$ .*

*Proof.* For any admissible set  $M \subset X$ , define

$$\begin{aligned} A(M) &= \bigcap_{x \in M} B(x, c \operatorname{diam} M), \\ H(M) &= \bigcap_{y \in A(M)} B(y, c \operatorname{diam} M) \cap A(M), \end{aligned}$$

where  $c < 1$  satisfies the conditions of uniform relative normal structure for  $X$ . Notice that  $H(M) \neq \emptyset$  because  $z_M \in H(M)$ .

Fix  $x_0 \in X$  and let  $A_0 = \bigcap_{g \in G} B(gx_0, \delta(x_0))$ , where

$$\delta(x_0) = \sup_{g, h \in G} d(gx_0, hx_0)$$

denotes the diameter of the orbit of  $x_0$ . Note that  $A_0$  is admissible and

$$g(A_0) = \bigcap_{h \in G} B(ghx_0, \delta(x)) = A_0.$$

We take by recursion  $A_1 = H(A_0)$ ,  $A_{n+1} = H(A_n)$ , and pick  $x_n \in A_n$  for  $n = 1, 2, \dots$ . If  $z \in A_{n+1}$ , then  $d(z, y) \leq c \operatorname{diam} A_n$  for each  $y \in A_{n+1} \subset A(A_n)$  and hence

$$\operatorname{diam} A_{n+1} \leq c \operatorname{diam} A_n \leq \dots \leq c^n \operatorname{diam} A_1.$$

Similarly, since  $x_n \in A(A_n)$ ,

$$d(x_{n+1}, x_n) \leq c \operatorname{diam} A_n \leq \dots \leq c^n \operatorname{diam} A_1.$$

Thus  $(x_n)$  is a Cauchy sequence converging to some  $x \in X$ . Moreover,  $g(A_n) = A_n$  for every  $g \in G$ . It follows that

$$d(gx, x) \leq d(gx, gx_n) + d(gx_n, x_n) + d(x_n, x) \longrightarrow 0 \text{ if } n \rightarrow \infty,$$

and hence  $gx = x$  for every  $g \in G$ .  $\square$

In particular, if  $X = K$  is a bounded closed subset of a Banach space we obtain the Lau theorem 2.2, and Bruhat–Tits' theorem [5] (see also [3, Corollary II.2.8]) follows if we consider a closed convex subset of a Hadamard space. But the applications of Theorem 2.3 are even wider. Recall that a metric space is injective (or hyperconvex) if

$$\bigcap_{\alpha \in I} B(x_\alpha, r_\alpha) \neq \emptyset$$

for any collection of closed balls  $\{B(x_\alpha, r_\alpha)\}_{\alpha \in I}$  such that  $d(x_\alpha, x_\beta) \leq r_\alpha + r_\beta$ ,  $\alpha, \beta \in I$ . Notice that injective metric spaces have uniform relative normal structure with  $c = 1/2$ , and thus we obtain the following corollary proved by Lang [22, Proposition 1.2] with completely different methods (see also [31, Theorem 2.3] for a little more general result).

**Corollary 2.4** (U. Lang). *Let  $X$  be an injective metric space. If  $G$  is a group of isometries  $g : X \rightarrow X$  with bounded orbits, then there is a fixed point of  $G$  in  $X$ .*

In particular, Corollary 2.4 applies to a group of isometries with bounded orbits acting on a real  $L_\infty(\mu)$  whenever  $\mu$  is a  $\sigma$ -finite measure on  $(E, \Sigma)$ , a Radon measure on a locally compact space (see [9, Corollary 3.10]), or a counting measure on any  $\Gamma$ . Note that a complex  $L_\infty(\mu)$  need not be hyperconvex, yet it follows from [30, Lemma] that it has uniform relative normal structure with  $c = 1/\sqrt{2}$ .

**Corollary 2.5.** *Let  $\mu$  be a  $\sigma$ -finite measure on  $(E, \Sigma)$ , a Radon measure on a locally compact space, or a counting measure on any  $\Gamma$ . If  $G$  is a subgroup of the isometry group of a real or complex  $L_\infty(\mu)$  with bounded orbits, then there is a fixed point of  $G$  in  $L_\infty(\mu)$ .*

Equipped with Theorem 2.3, the discussion around Corollary C in [2] can be repeated verbatim for the case of Banach bimodules with the uniform relative normal structure.

If  $\mathcal{A}$  is a Banach algebra [with unit  $e$ ], a Banach space  $M$  is said to be a Banach  $\mathcal{A}$ -bimodule if there are bounded bilinear mappings  $(a, x) \mapsto a.x, (x, a) \mapsto x.a : \mathcal{A} \times M \rightarrow M$  such that  $[e.x = x.e = x$  for every  $x \in M]$  and the usual associative law holds for each type of triple product:  $a_1 a_2 x, a_1 x a_2, x a_1 a_2$ . We will assume that

$$(1) \quad \|a.x\| \leq \|a\| \|x\|, \quad \|x.a\| \leq \|x\| \|a\| \quad (a \in \mathcal{A}, x \in M).$$

Note that for any Banach  $\mathcal{A}$ -bimodule  $M$  we can define an equivalent norm which satisfies (1).

By a derivation, from a Banach algebra  $\mathcal{A}$  into a Banach  $\mathcal{A}$ -bimodule  $M$ , we mean a linear mapping  $\delta : \mathcal{A} \rightarrow M$  such that  $\delta(ab) = \delta(a).b + a.\delta(b)$  for  $a, b \in \mathcal{A}$ . A derivation is inner if there exists  $T \in M$  such that  $\delta(a) = T.a - a.T$  for each  $a \in \mathcal{A}$ .

**Theorem 2.6.** *Let  $\mathcal{A}$  be a unital  $C^*$ -algebra, and let  $M$  be a Banach space with uniform relative normal structure. If  $M$  is a Banach  $\mathcal{A}$ -bimodule, then any derivation  $\delta : \mathcal{A} \rightarrow M$  is inner.*

*Proof.* Let  $U(\mathcal{A})$  denotes the group of unitaries of  $\mathcal{A}$ . For any  $g \in U(\mathcal{A})$  and  $x \in M$ , set

$$\alpha(g)x = g.x.g^{-1} + \delta(g).g^{-1},$$

and notice that  $\alpha : U(\mathcal{A}) \times M \rightarrow M$  is an (affine) and isometric action of  $U(\mathcal{A})$  on  $M$ , and thus

$$\begin{aligned} \alpha(gh)x &= \alpha(g)\alpha(h)x, \\ \|\alpha(g)x - \alpha(g)y\| &= \|x - y\| \quad (g, h \in U(\mathcal{A}), x, y \in M). \end{aligned}$$

Furthermore, Ringrose's theorem [29] yields  $\delta : \mathcal{A} \rightarrow M$  is bounded, and hence the orbit

$$A = \{\delta(g).g^{-1} : g \in U(\mathcal{A})\}$$

of  $\alpha$  at 0 is bounded too. It follows from Theorem 2.3 that there is  $T \in M$  such that  $\alpha(g)T = T$ , which gives

$$\delta(g) = T.g - g.T \quad (g \in U(\mathcal{A})).$$

This finishes the proof since any element of  $\mathcal{A}$  is a linear combination of four unitaries (in fact, three [18]).  $\square$

We now turn to the case of group algebras  $L_1(G)$ , where  $G$  is a locally compact group. This case is a little more subtle. If  $E$  is an essential Banach  $L_1(G)$ -bimodule, i.e., the linear hull of  $\{a.t.b : a, b \in L_1(G), t \in E\}$  is dense in  $E$ , then the dual actions

$$\langle t, x.a \rangle = \langle a.t, x \rangle, \quad \langle t, a.x \rangle = \langle t.a, x \rangle \quad (a \in L_1(G), t \in E, x \in M)$$

of  $L_1(G)$  on  $M = E^*$  can be extended to actions of  $M(G)$ , the space of (complex-valued) Radon measures on  $G$ . Let  $\delta : L_1(G) \rightarrow M$  be a continuous derivation. Then there is a unique derivation  $\tilde{\delta} : M(G) \rightarrow M$  extending  $\delta$  such that  $\|\tilde{\delta}\| = \|\delta\|$ , and

$$\langle a.t, \tilde{\delta}\mu \rangle = \langle t, \delta(\mu * a) \rangle - \langle t.\mu, \delta a \rangle \quad (a \in L_1(G), \mu \in M(G), t \in E),$$

where  $*$  denotes the convolution (with respect to a fixed left Haar measure on  $G$ ) and  $t.\mu$  ( $t \in E, \mu \in M(G)$ ) extends the action of  $L_1(G)$  on  $E$  to  $M(G)$  (see [8, Theorem 5.6.34], [14]). For any  $g \in G$  and  $x \in M$ , define an action  $\alpha : G \times M \rightarrow M$  by setting

$$\alpha(g)x = \delta_g.x.\delta_{g^{-1}} + \tilde{\delta}(\delta_g).\delta_{g^{-1}},$$

where  $\delta_g$  denotes the Dirac measure at  $g$ . Since the dual Banach  $L_1(G)$ -bimodule  $M$  satisfies (1),  $\alpha$  is an isometric action of  $G$  on  $M$  with bounded orbits, and thus by Theorem 2.3, there is  $T \in M$  such that  $\alpha(g)T = T$  whenever  $M$  has uniform relative normal structure. It follows that

$$\tilde{\delta}(\delta_g) = T.\delta_g - \delta_g.T,$$

and then Theorem 5.6.39 in [8] shows that  $\delta : L_1(G) \rightarrow M$  is inner. We are thus led to the following theorem.

**Theorem 2.7.** *Let  $G$  be a locally compact group, and let  $E$  be an essential Banach  $L_1(G)$ -bimodule. If the dual  $L_1(G)$ -bimodule  $E^*$  has uniform relative normal structure, then any continuous derivation  $\delta : L_1(G) \rightarrow E^*$  is inner.*

**Corollary 2.8.** *Let  $G$  be a locally compact group, and let  $L_1(\mu)$  be an essential Banach  $L_1(G)$ -bimodule, where  $\mu$  is a  $\sigma$ -finite measure on  $(E, \Sigma)$  or a Radon measure on a locally compact space. Then any continuous derivation  $\delta : L_1(G) \rightarrow L_\infty(\mu)$  (into the dual  $L_1(G)$ -bimodule  $L_\infty(\mu)$ ) is inner.*

In particular, Corollary 2.8 applies to  $L_1(G)$  itself with the convolution product and in this case gives the theorem of Johnson [16].

### 3. A FIXED POINT THEOREM IN $B(H, \ell_\infty)$

In 1955, motivated by work of Dixmier and Day on uniformly bounded group representations, Kadison [17] raised the problem of whether every bounded homomorphism  $u$  from  $C^*$ -algebra  $\mathcal{A}$  into  $B(H)$  is similar to a  $*$ -homomorphism, i.e., does there exist an invertible operator  $S \in B(H)$  such that the map  $\tilde{u}(x) = S^{-1}u(x)S$  satisfies the condition  $\tilde{u}(x^*) = \tilde{u}(x)^*$ ? The Kadison similarity problem is equivalent to a number of other important problems (see [6, 7, 13, 28, 27]), including the derivation problem [20]—given a  $C^*$ -algebra  $\mathcal{A} \subset B(H)$ , is every derivation  $\delta : \mathcal{A} \rightarrow B(H)$  inner?

Our observation concerning derivations from a  $C^*$ -algebra  $\mathcal{A}$  into  $B(H)$  is based on the following consequence of Theorem 2.3 in Section 2.

**Theorem 3.1.** *Let  $B(H, \ell_\infty(\Gamma))$  be a Banach space of bounded operators from a (complex) Hilbert space  $H$  into a (complex)  $\ell_\infty(\Gamma)$  for some  $\Gamma$ . If  $G$  is a group of isometries  $g : B(H, \ell_\infty(\Gamma)) \rightarrow B(H, \ell_\infty(\Gamma))$  with bounded orbits, then there exists  $x \in B(H, \ell_\infty(\Gamma))$  such that  $gx = x$  for every  $g \in G$ .*

*Proof.* Since  $B(H, \ell_\infty(\Gamma))$  is isometric to  $\ell_\infty(\Gamma, H^*) \simeq \ell_\infty(\Gamma, H)$  by setting  $B(H, \ell_\infty(\Gamma)) \ni F \mapsto f \in \ell_\infty(\Gamma, H^*)$ , where

$$\langle h, f(\gamma) \rangle = F(h)(\gamma) \quad (h \in H, \gamma \in \Gamma),$$

it suffices to show, using Theorem 2.3, that  $\ell_\infty(\Gamma, H)$  has uniform relative normal structure. It is quite simple (see also [30, Proposition]). For a bounded set  $M \subset \ell_\infty(\Gamma, H)$ , let

$$M_i = \{x_i \in H : (x_\gamma)_{\gamma \in \Gamma} \in M\} \quad (i \in \Gamma).$$

Then, for each  $i$ , there exists  $z_i \in \overline{\text{co}} M_i$  such that

$$\|x_i - z_i\| \leq \frac{\sqrt{3}}{2} \text{diam } M_i \leq \frac{\sqrt{3}}{2} \text{diam } M \quad (x_i \in M_i, i \in \Gamma).$$

Now suppose that  $M \subset B((y_\gamma), \frac{\sqrt{3}}{2} \text{diam } M)$  for some  $(y_\gamma) \in \ell_\infty(\Gamma, H)$ . Then  $M_i \subset B(y_i, \frac{\sqrt{3}}{2} \text{diam } M)$  for each  $i$ , and hence also  $\overline{\text{co}} M_i \subset B(y_i, \frac{\sqrt{3}}{2} \text{diam } M)$ . It follows that

$$\|z_i - y_i\| \leq \frac{\sqrt{3}}{2} \text{diam } M$$

for every  $i \in \Gamma$ , which completes the proof.  $\square$

**Remark 3.2.** The method of proof carries over to  $\ell_\infty(\Gamma, X)$  when  $X$  is uniformly convex (see [30]), and more generally when  $X$  has uniform normal structure (see [1, Theorem VI.2.2]).

Given a Hilbert space  $H$ , we can embed it (linearly) isometrically into  $\ell_\infty(\Gamma)$ , where  $\Gamma = B_H$  (or any norming subset of  $B_H$ ) by setting

$$H \ni h \mapsto (\langle h, \gamma \rangle)_{\gamma \in \Gamma} \in \ell_\infty(\Gamma).$$

Let  $\tilde{H}$  denote the isometric copy of  $H$  in  $\ell_\infty(\Gamma)$ , and define

$$\kappa(a)(\langle h, \gamma \rangle)_{\gamma \in \Gamma} = (\langle ah, \gamma \rangle)_{\gamma \in \Gamma} \quad (a \in B(H), (\langle h, \gamma \rangle)_{\gamma \in \Gamma} \in \tilde{H}).$$

Then  $\kappa : B(H) \rightarrow B(\tilde{H})$  is a linear, isometric, unital  $*$ -isomorphism, i.e.,  $\kappa(ab) = \kappa(a) \circ \kappa(b)$  and  $\kappa(a^*) = \kappa(a)^*$  for every  $a, b \in B(H)$ . From now on we will regard  $H$  as a subspace of  $\ell_\infty(\Gamma)$ , and  $B(H)$  as a subspace of  $B(H, \ell_\infty(\Gamma))$ —the Banach space of bounded linear operators from  $H$  into  $\ell_\infty(\Gamma)$ .

Now let  $\mathcal{A} \subset B(H) \subset B(H, \ell_\infty(\Gamma))$  be a unital  $C^*$ -algebra and extend each unitary  $g \in U(\mathcal{A})$  to a linear isometry  $\tilde{g} : \ell_\infty(\Gamma) \rightarrow \ell_\infty(\Gamma)$  by setting

$$\ell_\infty(\Gamma) \ni (z_\gamma)_{\gamma \in \Gamma} \mapsto (z_{g^*\gamma})_{\gamma \in \Gamma} \in \ell_\infty(\Gamma).$$

Then

$$(\tilde{g} \circ \tilde{h})(z_\gamma)_{\gamma \in \Gamma} = \tilde{g}((z_{h^*\gamma})_{\gamma \in \Gamma}) = (z_{h^*g^*\gamma})_{\gamma \in \Gamma} = \widetilde{(g \circ h)}(z_\gamma)_{\gamma \in \Gamma}$$

for every  $g, h \in U(\mathcal{A})$ ,  $(z_\gamma)_{\gamma \in \Gamma} \in \ell_\infty(\Gamma)$  and  $\gamma \in \Gamma$ . In other words,  $(g, (z_\gamma)_{\gamma \in \Gamma}) \mapsto \tilde{g}((z_\gamma)_{\gamma \in \Gamma})$  is a left action of  $U(\mathcal{A})$  on  $\ell_\infty(\Gamma)$ .

We are thus led to the following factorization theorem.

**Theorem 3.3.** *Let  $\mathcal{A} \subset B(H) \subset B(H, \ell_\infty(\Gamma))$  be a unital  $C^*$ -algebra, and let  $\delta : \mathcal{A} \rightarrow B(H)$  be a derivation (with the usual product operation). Then there exists a bounded linear operator  $T : H \rightarrow \ell_\infty(\Gamma)$  such that  $\delta(g) = Tg - \tilde{g}T$  for every  $g \in U(\mathcal{A})$ .*

*Proof.* The proof follows the arguments in Theorem 2.6. For any unitary  $g \in U(\mathcal{A})$  and  $x \in X = B(H, \ell_\infty(\Gamma))$ , set

$$\alpha(g)x = \tilde{g}xg^{-1} + \delta(g)g^{-1},$$

and note that  $\alpha : U(\mathcal{A}) \times X \rightarrow X$  is an isometric action of  $U(\mathcal{A})$  on  $X$  with bounded orbits. Hence Theorem 3.1 yields  $\alpha(g)T = T$  for some  $T \in X$ . It follows that

$$\delta(g) = Tg - \tilde{g}T$$

for each unitary element  $g$ , which completes the proof.  $\square$

**Remark 3.4.** Note that a similar result holds for derivations from  $\ell_1(G)$  to  $B(H)$  on which  $G$  acts by isometries, and it is well known that there are non-inner derivations from  $\ell_1(F_2)$  to  $B(\ell_2(F_2))$  (see, e.g., [25]). However, in the case of  $C^*$ -algebras, the Christensen theorem (see [7]) asserts that a derivation  $\delta : \mathcal{A} \rightarrow B(H)$  is inner if and only if it is completely bounded, and we can use Theorem 3.3 to study complete boundedness of  $\delta$ . Let  $T : H \rightarrow \ell_\infty(\Gamma)$  be a bounded linear operator such that  $\delta(g) = Tg - \tilde{g}T$  for every unitary  $g \in U(\mathcal{A})$ . Following [26], define

$$S = \begin{pmatrix} I & T \\ 0 & I \end{pmatrix} : H \oplus_2 H \rightarrow \ell_\infty(\Gamma) \oplus_2 \ell_\infty(\Gamma),$$

where  $X = \ell_\infty(\Gamma) \oplus_2 \ell_\infty(\Gamma)$  denotes a Banach space with the norm  $\|(x_1, x_2)\| = (\|x_1\|^2 + \|x_2\|^2)^{1/2}$ . Clearly,  $H \oplus_2 H$  is a Hilbert subspace of  $X$ , and note that  $S$  is an invertible operator onto  $Y = S(H \oplus_2 H)$  with the inverse

$$S^{-1} = \begin{pmatrix} I & -T \\ 0 & I \end{pmatrix}. \text{ Furthermore, if we define for } g \in U(\mathcal{A}),$$

$$u(g) = \begin{pmatrix} g & -\delta(g) \\ 0 & g \end{pmatrix} : H \oplus_2 H \rightarrow H \oplus_2 H,$$

$$\tilde{G} = \begin{pmatrix} \tilde{g} & 0 \\ 0 & \tilde{g} \end{pmatrix} : X \rightarrow X,$$

we get  $u(g) = S^{-1}\tilde{G}S$  and hence  $\tilde{G}|_Y = Su(g)S^{-1} : Y \rightarrow Y$  for each unitary  $g$ . Thus we obtain a homomorphism

$$\mathcal{A} \ni a \rightarrow Su(a)S^{-1} \in B(Y)$$

from  $\mathcal{A}$  into  $B(Y)$  that extends the group homomorphism

$$U(\mathcal{A}) \ni g \rightarrow \tilde{G}|_Y \in B(Y),$$



and complete boundedness of this homomorphism is equivalent to innerity of the derivation.

## REFERENCES

- [1] J.M. Ayerbe Toledano, T. Domínguez Benavides and G. López Acedo, Measures of noncompactness in metric fixed point theory, Birkhäuser Verlag, Basel, 1997.
- [2] U. Bader, T. Gähler, and N. Monod, A fixed point theorem for  $L_1$  spaces, *Invent. Math.* 189 (2012), 143–148.
- [3] M.R. Bridson and A. Haefliger, Metric spaces of non-positive curvature, Springer-Verlag, Berlin, 1999.
- [4] M.S. Brodskii and D.P. Mil'man, On the center of a convex set, *Doklady Akad. Nauk SSSR* 59 (1948), 837–840 (in Russian).
- [5] F. Bruhat and J. Tits, Groupes réductifs sur un corps local: I. Données radicielles valuées, *Publications Math. IHES* 41 (1972), 5–251.
- [6] J. Cameron et al., A remark on the similarity and perturbation problems, *C. R. Math. Acad. Sci. Soc. R. Can.* 35 (2013), 70–76.
- [7] E. Christensen, Extensions of derivations II, *Math. Scand.* 50 (1982), 111–122.
- [8] H.G. Dales, Banach Algebras and Automatic Continuity, Oxford University Press, New York, 2000.
- [9] A. Defant and K. Floret, Tensor norms and operator ideals, North-Holland Publishing Co., Amsterdam, 1993.
- [10] R. Espínola-García, M. Japón and D. Souza, New examples of subsets of  $c$  with the FPP and stability of the FPP in hyperconvex spaces, *J. Fixed Point Theory Appl.* 23 (2021), Paper No. 45, 19 pp.
- [11] R. Espínola, M. A. Khamsi, Introduction to hyperconvex spaces, in: Handbook of metric fixed point theory, W. A. Kirk, B. Sims (Eds.), Kluwer Academic Publishers, Dordrecht, 2001, pp. 391–435.
- [12] U. Haagerup, All nuclear  $C^*$ -algebras are amenable, *Invent. Math.* 74 (1983), 305–319.
- [13] D. Hadwin and V. Paulsen, Two reformulations of Kadison's similarity problem, *J. Operator Theory* 55 (2006), 3–16.
- [14] B.E. Johnson, Cohomology in Banach algebras, *Mem. Amer. Math. Soc.* 127 (1972), iii+96 pp.
- [15] B.E. Johnson, Weak amenability of group algebras of connected complex semisimple Lie groups, *Proc. Amer. Math. Soc.* 111 (1991), 177–185.
- [16] B.E. Johnson, Weak amenability of group algebras, *Bull. London Math. Soc.* 23 (1991), no. 3, 281–284.
- [17] R.V. Kadison, On the orthogonalization of operator representations, *Amer. J. Math.* 77 (1955), 600–620.
- [18] R.V. Kadison and G.K. Pedersen, Means and convex combinations of unitary operators, *Math. Scand.* 57 (1985), 249–266.
- [19] M.A. Khamsi and W.A. Kirk, An introduction to metric spaces and fixed point theory, Wiley-Interscience, New York, 2001.
- [20] E. Kirchberg, The derivation problem and the similarity problem are equivalent, *J. Operator Theory* 36 (1996), no. 1, 59–62.
- [21] W.A. Kirk, Nonexpansive mappings and the weak closure of sequences of iterates, *Duke Math. J.* 36 (1969), 639–645.
- [22] U. Lang, Injective hulls of certain discrete metric spaces and groups, *J. Topol. Anal.* 5 (2013), 297–331.
- [23] A.T.-M. Lau, Sets with fixed point property for isometric mappings, *Proc. Amer. Math. Soc.* 79 (1980), 388–392.

- [24] V. Losert, The derivation problem for group algebras, *Ann. of Math. (2)* 168 (2008), 221–246.
- [25] V. Losert, On derivations and crossed homomorphisms, in: *Banach algebras 2009*, R.J. Loy, V. Runde and A. Soltysiak (Eds.), Banach Center Publ. 91, Polish Acad. Sci. Inst. Math., Warsaw, 2010, pp. 199–217.
- [26] V. Paulsen, *Completely bounded maps and dilations*. Pitman Research Notes in Math. 146, Longman, Wiley, New York, 1986.
- [27] G. Pisier, The similarity degree of an operator algebra, *Algebra i Analiz* 10 (1998), 132–186; reprinted in *St. Petersburg Math. J.* 10 (1999), 103–146.
- [28] G. Pisier, *Similarity problems and completely bounded maps*, Springer-Verlag, Berlin, 2001.
- [29] J.R. Ringrose, Automatic continuity of derivations of operator algebras, *J. London Math. Soc.* 5 (1972), 432–438.
- [30] P.M. Soardi, Existence of fixed points of nonexpansive mappings in certain Banach lattices, *Proc. Amer. Math. Soc.* 73 (1979), 25–29.
- [31] A. Wiśnicki and J. Wośko, Uniformly Lipschitzian group actions on hyperconvex spaces, *Proc. Amer. Math. Soc.* 144 (2016), 3813–3824.

PEDAGOGICAL UNIVERSITY OF KRAKÓW, CRACOW, POLAND

*Email address:* `andrzej.wisnicki@up.krakow.pl`