# A NOTE ON THE SQUARES OF THE FORM $\prod_{k=1}^{n} (2k^2 + l)$ WITH l ODD

RUSSELLE GUADALUPE

ABSTRACT. Let l be a positive odd integer. Using Cilleruelo's method, we establish an explicit lower bound  $N_l$  depending on l such that for all  $n \ge N_l$ ,  $\prod_{k=1}^n (2k^2 + l)$  is not a square. As an application, we determine all values of n such that  $\prod_{k=1}^n (2k^2 + l)$  is a square for certain values of l.

### 1. INTRODUCTION

For positive integers a, c and n with gcd(a, c) = 1, define  $P_{a,c}(n) := \prod_{k=1}^{n} (ak^2 + c)$ . The question of determining whether the sequence  $\{P_{a,c}(n)\}_{n\geq 1}$  contains infinitely many squares has been extensively studied. In 2008, Amdeberhan, Medina, and Moll [1] conjectured that  $P_{1,1}(n)$  is not a square for  $n \geq 4$  and  $P_{4,1}(n)$  is not a square for  $n \geq 1$ . In addition, they computationally verified that  $P_{1,1}(n)$  is not a square for  $n \leq 10^{3200}$ . Shortly, Cilleruelo [3] proved the conjecture for  $P_{1,1}(n)$  and showed that  $P_{1,1}(n)$  is square only for n = 3. Fang [4] settled the conjecture for  $P_{4,1}(n)$ . Yang, Togbé, and He [9] found all positive integer solutions to the equation  $P_{a,c}(n) = y^l$  for coprime integers  $a, c \in \{1, \ldots, 20\}$  and  $l \geq 2$ . Yin, Tan, and Luo [10] obtained the *p*-adic valuation of  $P_{1,21}(n)$  for all primes *p* and showed that  $P_{1,21}(n)$  is not a square for all  $n \geq 1$ . Chen, Wang and Hu [2] proved that  $P_{1,23}(n)$  is a square only for n = 3.

Ho [6] studied the sequence  $\{P_{1,m^2}(n)\}_{n\geq 1}$  for  $m\geq 1$  and proved that if m has divisors of the form 4q + 1 and  $N = \max\{m, 10^8\}$ , then  $P_{1,m^2}(n)$  is not a square for all  $n\geq N$ . Recently, Zhang and Niu [11] generalized Ho's result by showing that there is a positive integer  $N_q$  depending on q such that  $P_{1,q}(n)$  is not a square for all  $n\geq N_q$ . Motivated by their work, we consider the problem of finding squares of the form  $P_{2,l}(n)$  for a positive odd integer l. In this paper, we apply Cilleruelo's technique to prove the following result.

**Theorem 1.1.** Let l be a positive odd integer. Then there exists a positive integer  $N_l$  depending on l such that for all  $n \ge N_l$ ,  $P_{2,l}(n) = \prod_{k=1}^n (2k^2 + l)$  is not a square.

As an application of Theorem 1.1, we show that for certain values of l, there are finitely many squares of the form  $P_{2,l}(n)$  for some integer  $n \ge 1$ .

Corollary 1.2.  $P_{2,1}(n) = \prod_{k=1}^{n} (2k^2 + 1)$  is not a square for all  $n \ge 1$ . Corollary 1.3.  $P_{2,3}(n) = \prod_{k=1}^{n} (2k^2 + 3)$  is not a square for all  $n \ge 1$ . Corollary 1.4.  $P_{2,7}(n) = \prod_{k=1}^{n} (2k^2 + 7)$  is a square only for n = 1.

We organize the paper as follows. In Section 2, we present several preliminary lemmas needed for the proof of Theorem 1.1. In Sections 3 and 4, we prove Theorem 1.1 by working on two separate cases:  $l \geq 3$  and l = 1. In Sections 5, 6 and 7, we prove Corollaries 1.2, 1.3 and 1.4 using Theorem 1.1.

<sup>2010</sup> Mathematics Subject Classification: Primary 11D09; Secondary 11C08, 11A15.

Key words and phrases: Quadratic polynomials, Diophantine equation, quadratic reciprocity.

#### RUSSELLE GUADALUPE

#### 2. Preliminaries

In this section, we list some preliminary results that are used in the proofs of our main theorem. Throughout this section, we denote a prime by p and a positive odd integer by l.

**Lemma 2.1.** Let  $n > \sqrt{l/2}$  be such that  $P_{2,l}(n)$  is a square and let p be a prime divisor of  $P_{2,l}(n)$ . Then p < 2n.

*Proof.* Since  $P_{2,l}(n)$  is a square and  $p \mid P_{2,l}(n)$ , we have  $p^2 \mid P_{2,l}(n)$ . We consider two cases:

- (1) Suppose  $p^2 \mid 2k^2 + l$  for some  $1 \le k \le n$ . Then  $p \le \sqrt{2k^2 + l} \le \sqrt{2n^2 + l} < 2n$  since  $l < 2n^2$ .
- (2) Suppose  $p \mid 2k^2 + l$  and  $p \mid 2l^2 + 1$  for all  $1 \le k < l \le n$ . Then  $p \mid 2(l^2 k^2) = 2(l-k)(l+k)$ . Since l is odd,  $P_{2,l}(n)$  is odd, so  $p \nmid 2$ . Thus, we see that either  $p \mid l-k$  or  $p \mid l+k$ , which implies that  $p \le \max\{l-k, l+k\} < 2n$ .

The above lemma implies that if  $n > \sqrt{l/2}$ , then  $P_{2,l}(n) = \prod_{p < 2n} p^{\alpha_p}$ , where p is odd and

(1) 
$$\alpha_p = \sum_{j \le \log(2k^2 + l)/\log p} \#\{1 \le k \le n : p^j \mid 2k^2 + l\}.$$

In particular, we have  $\alpha_2 = 0$ . Now, observe that for  $1 \le k \le n$ ,

$$2k^2 + l = k^{\log(2k^2 + l)/\log k} > k^{\log(2n^2 + l)/\log k}$$

where the last inequality follows from the fact that for  $l \ge 1$ , the function  $f_l(x) := \log(2x^2 + l)/\log x$  is decreasing on  $(1, +\infty)$ . Setting  $\lambda = \log(2n^2 + l)/\log n$ , we have  $P_{2,l}(n) > (n!)^{\lambda}$  and writing  $n! = \prod_{p \le n} p^{\beta_p}$ , where

(2) 
$$\beta_p = \sum_{j \le \log n/\log p} \#\{1 \le k \le n : p^j \mid k\} = \sum_{j \le \log n/\log p} \left\lfloor \frac{n}{p^j} \right\rfloor,$$

we deduce that

(3) 
$$\sum_{p \le n} \beta_p \log p \le \frac{1}{\lambda} \sum_{p < 2n} \alpha_p \log p$$

**Lemma 2.2.** Let p be an odd prime with  $p \nmid l$  and let j > 0. Then the congruence  $2x^2 \equiv -l \pmod{p^j}$  has  $1 + \left(\frac{-2l}{p}\right)$  solutions.

*Proof.* This follows from rewriting the congruence as  $(2x)^2 \equiv -2l \pmod{p^j}$  and applying [7, Thm. 5.1].

**Lemma 2.3.** Let p be an odd prime with  $p \nmid l$ .

(1) If 
$$\left(\frac{-2l}{p}\right) = -1$$
, then  $\alpha_p = 0$ .  
(2) If  $\left(\frac{-2l}{p}\right) = 1$ , then  $\frac{\alpha_p}{\lambda} - \beta_p \le \frac{\log(2n^2 + l)}{\log p}$ 

*Proof.* (1) If  $\left(\frac{-2l}{p}\right) = -1$ , then by Lemma 2.2, the congruence  $2x^2 + l \equiv 0 \pmod{p^j}$  has no solutions for j > 0. Thus,  $p^j \nmid 2k^2 + l$  for all integers  $j, k \ge 1$  and  $\alpha_p = 0$ .

(2) If  $\left(\frac{-2l}{p}\right) = 1$ , then by Lemma 2.2, the congruence  $2x^2 + l \equiv 0 \pmod{p^j}$  has at most two solutions in an interval of length  $p^j$  for each j > 0. From (1), we deduce that

(4) 
$$\alpha_p \le \sum_{j \le \log(2k^2 + l)/\log p} 2\left\lceil \frac{n}{p^j} \right\rceil \le \sum_{j \le \log(2k^2 + l)/\log p} \lambda\left\lceil \frac{n}{p^j} \right\rceil$$

Thus, we obtain

$$\frac{\alpha_p}{\lambda} - \beta_p \le \sum_{j \le \log n/\log p} \left( \left\lceil \frac{n}{p^j} \right\rceil - \left\lfloor \frac{n}{p^j} \right\rfloor \right) + \sum_{\log n/\log p < j \le \log(2n^2 + l)/\log p} \left\lceil \frac{n}{p^j} \right\rceil$$
$$\le \sum_{j \le \log n/\log p} 1 + \sum_{\log n/\log p < j \le \log(2n^2 + l)/\log p} 1 = \frac{\log(2n^2 + l)}{\log p}.$$

**Lemma 2.4.** Suppose  $p^s$  is the largest power of an odd prime p dividing  $l \geq 3$ . Then  $\alpha_p \leq \gamma_{l,p}(n)$ , where

$$\begin{split} \gamma_{l,p}(n) &= \frac{n}{2} \left( \frac{3p - 2 - p^{-s}}{(p - 1)^2} - \frac{s + 3}{p^s(p - 1)} + \frac{4}{p^{s/2}(p - 1)} \right) - \frac{2np^{s/2}}{(p - 1)(2n^2 + l)} \\ &+ \frac{s(s + 5)}{4} + 2p^{s/2} \left( \frac{\log(2n^2 + l)}{\log p} - s \right). \end{split}$$

*Proof.* Write  $l = p^s q$ , where  $p \nmid q$ . We consider the number of solutions to the congruence  $2x^2 \equiv -p^s q \pmod{p^j}$  with j > 0, which is equivalent to  $(2x)^2 \equiv -2p^s q \pmod{p^j}$ . We work on two cases:

- (1) Suppose  $s \ge j$ . Then  $2x \equiv p^{\lceil j/2 \rceil}, p^{\lceil j/2 \rceil+1}, \ldots, p^j \pmod{p^j}$ , so there are exactly  $j \lceil j/2 \rceil + 1$  solutions in this case.
- (2) Suppose s < j. Then  $(2x)^2 = mp^j 2p^sq = p^s(mp^{j-s} 2q)$  for some positive integer m. Since p does not divide  $mp^{j-s} 2q$ , we see that s is even. Write  $2x = p^{s/2}r$ , where  $x \in \{0, \ldots, p^j 1\}$  and  $r \in \{0, \ldots, p^{j-s/2} 1\}$ . The congruence now becomes  $r^2 \equiv -2q \pmod{p^{j-s}}$ , and in view of Lemma 2.2, it has at most two solutions contained in each interval of length  $p^{j-s}$ . Thus, the congruence has at most  $2p^{j-s/2}/p^{j-s} = 2p^{s/2}$  solutions contained in each interval of length  $p^j$ .

Combining these cases, we see that

$$\begin{split} \alpha_p &\leq \sum_{j \leq s} \left( j - \left\lceil \frac{j}{2} \right\rceil + 1 \right) \left\lceil \frac{n}{p^j} \right\rceil + \sum_{s < j \leq \log(2n^2 + l)/\log p} 2p^{s/2} \left\lceil \frac{n}{p^j} \right\rceil \\ &\leq \sum_{j \leq s} \left( j - \left\lceil \frac{j}{2} \right\rceil + 1 \right) \left( \frac{n}{p^j} + 1 \right) + \sum_{s < j \leq \log(2n^2 + l)/\log p} 2p^{s/2} \left( \frac{n}{p^j} + 1 \right) \\ &\leq \sum_{j \leq s} \left( \frac{j}{2} + 1 \right) \left( \frac{n}{p^j} + 1 \right) + 2p^{s/2} \left( \frac{n(\frac{1}{p^{s+1}} - \frac{1}{p(2n^2 + l)})}{1 - \frac{1}{p}} + \frac{\log(2n^2 + l)}{\log p} - s \right) \\ &= \frac{n}{2} \sum_{j \leq s} \frac{j + 2}{p^j} + \frac{s(s + 5)}{4} + \frac{2n}{p^{s/2}(p - 1)} - \frac{2np^{s/2}}{(p - 1)(2n^2 + l)} + 2p^{s/2} \left( \frac{\log(2n^2 + l)}{\log p} - s \right) \\ &= \gamma_{l,p}(n). \end{split}$$

**Lemma 2.5.** For all positive integers n, we have  $\sum_{n .$ 

*Proof.* Observe that for primes p with n , <math>p appears exactly once in the prime factorization of  $\binom{2n}{n}$ . Thus, by binomial theorem, we get  $\prod_{n , which is equivalent to the desired inequality.$ 

**Lemma 2.6.** Let  $\pi(n)$  be the number of primes at most n. Then for all positive integers n, we have  $\pi(n) \leq 2 \log 4 \frac{n}{\log n} + \sqrt{n}$ .

*Proof.* From [5, p. 459, (22.4.2)], we have

$$\pi(n) \le n^{1-\lambda} + \frac{1}{(1-\lambda)\log n} \sum_{p \le n} \log p$$

for all  $n \ge 1$  and  $\lambda \in (0, 1)$ . Setting  $\lambda = \frac{1}{2}$  and using  $\prod_{p < n} p \le 4^n$  (see [8, Thm. 1.4]), the desired inequality follows.

## 3. Proof of Theorem 1.1 for odd $l \geq 3$

We now give the proof of Theorem 1.1 for odd  $l \ge 3$  as follows. Suppose  $n > \sqrt{l/2}$  and define the set  $S = \{p : p \text{ is an odd prime with } (-\frac{2l}{p}) = 1\}$ . Then inequality (3) becomes

(5) 
$$\sum_{p \le n} \beta_p \log p \le \sum_{\substack{p \le n \\ p \in \mathcal{S}}} \frac{\alpha_p}{\lambda} \log p + \sum_{\substack{p \le n \\ p \notin \mathcal{S}}} \frac{\alpha_p}{\lambda} \log p + \sum_{\substack{n$$

Write  $l = \prod_{i=1}^{r} p_i^{e_i}$ , where  $p_1, \ldots, p_r$  are odd primes. By Lemma 2.3, we have

(6) 
$$\sum_{\substack{p \le n \\ p \notin \mathcal{S}}} \beta_p \log p \le \sum_{\substack{p \le n \\ p \in \mathcal{S}}} \log(2n^2 + l) + \frac{1}{\lambda} \sum_{i=1}^r \alpha_{p_i} \log p_i + \sum_{n$$

Observe that if p > n, then  $\alpha_p \leq \lambda$  from (4), so applying Lemma 2.5 in (6) yields

(7) 
$$\sum_{\substack{p \le n \\ p \notin S}} \beta_p \log p \le \log(2n^2 + l) \sum_{\substack{p \le n \\ p \in S}} 1 + \frac{1}{\lambda} \sum_{i=1}^r \alpha_{p_i} \log p_i + n \log 4$$

On the other hand, if  $p \leq n$ , then from (2) we have

$$\beta_p = \sum_{j \le \log n/\log p} \left\lfloor \frac{n}{p^j} \right\rfloor \ge \sum_{j \le \log n/\log p} \left( \frac{n}{p^j} - 1 \right) \ge n \sum_{j \le \log n/\log p} \frac{1}{p^j} - \frac{\log n}{\log p}$$
$$\ge n \left( \frac{1 - \frac{1}{n}}{1 - \frac{1}{p}} - 1 \right) - \frac{\log n}{\log p} = \frac{n - p}{p - 1} - \frac{\log n}{\log p}$$
$$\ge \frac{n - 1}{p - 1} - \frac{\log(2n^2 + l)}{\log p}.$$

Thus, from (7) we get

(8) 
$$(n-1)\sum_{\substack{p \le n \\ p \notin S}} \frac{\log p}{p-1} \le \log(2n^2 + l)\pi(n) + \frac{1}{\lambda}\sum_{i=1}^r \alpha_{p_i}\log p_i + n\log 4$$

and applying Lemma 2.6, we obtain

(9) 
$$\sum_{\substack{p \le n \\ p \notin S}} \frac{\log p}{p-1} \le \frac{\log(2n^2+l)}{n-1} \left( 2\log 4 \frac{n}{\log n} + \sqrt{n} \right) + \sum_{i=1}^r \frac{\alpha_{p_i} \log p_i}{\lambda(n-1)} + \frac{n\log 4}{n-1}.$$

Finally, using Lemma 2.4, we arrive at

$$\sum_{\substack{p \le n \\ p \notin \mathcal{S}}} \frac{\log p}{p-1} \le \frac{\log(2n^2+l)}{n-1} \left( 2\log 4 \frac{n}{\log n} + \sqrt{n} \right) + \frac{\log n}{\log(2n^2+l)} \sum_{i=1}^r \frac{\gamma_{l,p_i}(n)\log p_i}{n-1} + \frac{n\log 4}{n-1} + \frac{n\log 4}{n-1}$$

As n grows sufficiently large, the right-hand side of (10) approaches the limit

(11) 
$$10\log 2 + \frac{1}{4}\sum_{i=1}^{r} \left(\frac{3p_i - 2 - p_i^{-e_i}}{(p_i - 1)^2} - \frac{e_i + 3}{p_i^{e_i}(p_i - 1)} + \frac{4}{p_i^{e_i/2}(p_i - 1)}\right)\log p_i,$$

while the left-hand side becomes unbounded since the prime number theorem implies that

$$\sum_{p \le n} \frac{\log p}{p-1} = \log n - \gamma + o(1)$$

where  $\gamma$  is the Euler-Mascheroni constant (see [8]). Thus, there exists a positive integer  $N_l$  such that  $\sum_{p \leq n, p \notin S} (\log p)/(p-1)$  is greater than (11) for all  $n \geq N_l$ . Hence, we conclude that  $P_{2,l}(n)$  is not a square for all  $n \geq N_l$ .

## 4. Proof of Theorem 1.1 for l = 1

We now give the proof of Theorem 1.1 for l = 1 as follows. Suppose  $P_{2,1}(n)$  is a square for some  $n \ge 1$  and let p be a prime divisor of  $P_{2,1}(n)$ . By Lemma 2.1, we have p < 2n. Since p divides  $2k^2 + 1$  for some  $1 \le k \le n$ , we have  $\left(-\frac{2}{p}\right) = 1$ , so that  $p \equiv 1, 3 \pmod{8}$ . Thus, we have

$$P_{2,1}(n) = \prod_{\substack{p < 2n \\ p \equiv 1,3 \pmod{8}}} p^{\alpha_{2}}$$

and in view of  $P_{2,1}(n) > (n!)^{\lambda}$ , we see that

(12) 
$$\sum_{p \le n} \beta_p \log p \le \frac{1}{\lambda} \sum_{\substack{p < 2n \\ p \equiv 1,3 \pmod{8}}} \alpha_p \log p.$$

By Lemma 2.3, we have

(13) 
$$\sum_{\substack{p \le n \\ p \not\equiv 1,3 \pmod{8}}} \beta_p \log p \le \sum_{\substack{p \le n \\ p \equiv 1,3 \pmod{8}}} \log(2n^2 + 1) + \sum_{n$$

Note that if p > n, then  $\alpha_p \leq \lambda$  from (4) and if  $p \leq n$ , then

$$\beta_p \ge \frac{n-1}{p-1} - \frac{\log(2n^2+1)}{\log p}.$$

Using the above bounds for  $\alpha_p$  and  $\beta_p$  and Lemmas 2.5 and 2.6 to (13), we obtain

(14) 
$$\sum_{\substack{p \le n \\ p \not\equiv 1,3 \pmod{8}}} \frac{\log p}{p-1} \le \frac{\log(2n^2+1)}{n-1} \left(2\log 4\frac{n}{\log n} + \sqrt{n}\right) + \frac{n\log 4}{n-1}.$$

As n tends to infinity, the left-hand side of (14) becomes unbounded, while the right-hand side approaches to  $10 \log 2$ . Since the smallest positive integer n for which

$$\sum_{\substack{p \leq n \\ p \not\equiv 1,3 \pmod{8}}} \frac{\log p}{p-1} > 10 \log 2$$

is n = 706310, we take  $N_1 := 706310$  so that  $P_{2,1}(n)$  is not a square for  $n \ge N_1$ .

## 5. Proof of Corollary 1.2

We now apply Theorem 1.1 to prove Corollary 1.2.

*Proof.* We know from the previous section that  $P_{2,1}(n)$  is not a square for  $n \ge N_1 =$  706310. Since  $P_{2,1}(1) = 3$  and  $P_{2,1}(2) = 27$  are not squares, it suffices to prove that  $P_{2,1}(n)$  is not a square for  $3 \le n \le 706309$ . We proceed as follows:

- Since  $2 \cdot 3^2 + 1 = 19$  is a prime and the next value of k > 3 for which 19 divides  $2k^2 + 1$  is k = 19 3 = 16, we see that  $P_{2,1}(n)$  is not a square for  $3 \le n \le 15$ .
- Since  $2 \cdot 6^2 + 1 = 73$  is a prime and the next value of k > 6 for which 73 divides  $2k^2 + 1$  is k = 73 6 = 67, we see that  $P_{2,1}(n)$  is not a square for  $6 \le n \le 66$ .
- Since  $2 \cdot 21^2 + 1 = 883$  is a prime and the next value of k > 21 for which 883 divides  $2k^2 + 1$  is k = 883 21 = 862, we see that  $P_{2,1}(n)$  is not a square for  $21 \le n \le 861$ .
- Since  $2 \cdot 597^2 + 1 = 712819$  is a prime and the next value of k > 597 for which 712819 divides  $2k^2 + 1$  is k = 712819 597 = 712222, we see that  $P_{2,1}(n)$  is not a square for  $597 \le n \le 712221$ .

Hence,  $P_{2,1}(n)$  is not a square for all  $n \ge 1$ .

### 6. Proof of Corollary 1.3

We next prove Corollary 1.3 using Theorem 1.1.

*Proof.* We set l = 3 with  $r = 1, p_1 = 3$  and  $e_1 = 1$ . In this case, we have  $S = \{p \text{ prime } : p \equiv 1, 5, 7, 11 \pmod{24}\}$  and the limit (11) becomes  $10 \log 2 + \frac{1}{4}(1 + \frac{2}{\sqrt{3}}) \log 3 \approx 7.523267$ . Since the smallest positive integer n for which

$$\sum_{\substack{p \le n \\ p \notin S}} \frac{\log p}{p-1} > 10 \log 2 + \frac{1}{4} \left( 1 + \frac{2}{\sqrt{3}} \right) \log 3$$

is  $n = N_3 := 2189634$ , we see that  $P_{2,3}(n)$  is not a square for all  $n \ge N_3$ . Since  $P_{2,3}(1) = 5$  is not a square, it suffices to prove that  $P_{2,3}(n)$  is not a square for  $2 \le n \le 2189633$ . We proceed as follows:

- Since  $2 \cdot 2^2 + 3 = 11$  is a prime and the next value of k > 2 for which 11 divides  $2k^2 + 3$  is k = 11 2 = 9, we see that  $P_{2,3}(n)$  is not a square for  $2 \le n \le 8$ .
- Since  $2 \cdot 8^2 + 3 = 131$  is a prime and the next value of k > 8 for which 131 divides  $2k^2 + 3$  is k = 131 8 = 123, we see that  $P_{2,3}(n)$  is not a square for  $8 \le n \le 122$ .
- Since  $2 \cdot 37^2 + 3 = 2741$  is a prime and the next value of k > 37 for which 2741 divides  $2k^2 + 3$  is k = 2741 37 = 2704, we see that  $P_{2,3}(n)$  is not a square for  $37 \le n \le 2703$ .
- Since  $2 \cdot 1048^2 + 3 = 2196611$  is a prime and the next value of k > 1048 for which 2196611 divides  $2k^2 + 3$  is k = 2196611 1048 = 2195563, we see that  $P_{2,3}(n)$  is not a square for  $1048 \le n \le 2199562$ .

Hence,  $P_{2,3}(n)$  is not a square for all  $n \ge 1$ .

## 7. Proof of Corollary 1.4

We finally prove Corollary 1.4 using Theorem 1.1.

*Proof.* We set l = 7 with  $r = 1, p_1 = 7$  and  $e_1 = 1$ . In this case, we have  $S = \{p \text{ prime} : p \equiv 1, 3, 5, 9, 13, 15, 19, 23, 25, 27, 39, 45 \pmod{56}\}$  and the limit (11) becomes  $10 \log 2 + \frac{1}{4}(\frac{3}{7} + \frac{2}{3\sqrt{7}}) \log 7 \approx 7.262543$ . Since the smallest positive integer n for which

$$\sum_{\substack{p \le n \\ p \notin S}} \frac{\log p}{p-1} > 10 \log 2 + \frac{1}{4} \left(\frac{3}{7} + \frac{2}{3\sqrt{7}}\right) \log 7$$

is  $n = N_7 := 2142500$ , we see that  $P_{2,7}(n)$  is not a square for all  $n \ge N_7$ . By direct calculation, we deduce that the only value of n in  $1 \le n \le 5$  for which  $P_{2,7}(n)$  is a square is n = 1. Thus, it suffices to prove that  $P_{2,7}(n)$  is not a square for  $6 \le n \le 2142499$ . We proceed as follows:

- Since  $2 \cdot 6^2 + 7 = 79$  is a prime and the next value of k > 6 for which 79 divides  $2k^2 + 7$  is k = 79 6 = 73, we see that  $P_{2,7}(n)$  is not a square for  $6 \le n \le 72$ .
- Since  $2 \cdot 15^2 + 7 = 457$  is a prime and the next value of k > 15 for which 457 divides  $2k^2 + 7$  is k = 457 15 = 432, we see that  $P_{2,7}(n)$  is not a square for  $15 \le n \le 431$ .
- Since  $2 \cdot 39^2 + 7 = 3049$  is a prime and the next value of k > 39 for which 3049 divides  $2k^2 + 7$  is k = 3049 39 = 3010, we see that  $P_{2,7}(n)$  is not a square for  $39 \le n \le 3009$ .
- Since  $2 \cdot 1041^2 + 7 = 2167369$  is a prime and the next value of k > 1041 for which 2167369 divides  $2k^2 + 7$  is k = 2167369 1041 = 2166328, we see that  $P_{2,7}(n)$  is not a square for  $1041 \le n \le 2166327$ .

Hence, the only value of n for which  $P_{2,7}(n)$  is a square is n = 1.

#### 

## References

- T. Amdeberhan, L. Medina and V. Moll, Arithmetical properties of a sequence arising from an arctangent sum, J. Number Theory 128 (2008) 1807–1846.
- [2] H. Chen, C. Wang and S. Hu, Squares in (1<sup>2</sup> + 23)(2<sup>2</sup> + 23) · · · (n<sup>2</sup> + 23) (in Chinese), J. Sichuan Univ. Nat. Sci. Ed. 52 (2015) 21−24, doi.org/10.3969/j.issn.0490-6756.2015.01.005.
- [3] J. Cilleruelo, Squares in  $(1^2 + 1) \cdots (n^2 + 1)$ , J. Number Theory **128** (2008) 2488–2491.
- [4] J. Fang, Neither  $\prod_{k=1}^{n} (4k^2 + 1)$  nor  $\prod_{k=1}^{n} (2k(k-1) + 1)$  is a perfect square, Integers 9 (2009) 177–180.
- [5] G. H. Hardy and E. M. Wright, An Introduction to the Theory of Numbers (6th ed.) (Oxford University Press, 2008).
- [6] P. T. Ho, Squares in  $(1^2 + m^2) \cdots (n^2 + m^2)$ , Integers 9 (2009) 711–716.
- [7] L. K. Hua, Introduction to Number Theory (Springer-Verlag, 1982).
- [8] G. Tenenbaum, Introduction to Analytic and Probabilistic Number Theory (3rd ed.) (American Mathematical Society, 2015).
- [9] S. Yang, A. Togbé and B. He, Diophantine equations with products of consecutive values of a quadratic polynomial, J. Number Theory 131 (2009) 1840–1851.
- [10] Q. Yin, Q. Tan and Y. Luo, *p*-adic Valuation of  $(1^2 + 21) \cdots (n^2 + 21)$  and applications, *Southeast Asian Bull. Math.* **39** (2015) 747–754.
- [11] Q. Zhang and C. Niu, Squares in  $(1^2 + Q)(2^2 + Q) \cdots (n^2 + Q)$  (in Chinese), J. Liaocheng Univ. Nat. Sci. 6(2) (2022) 1–6, doi.org/10.19728/j.issn1672-6634.2021060002.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF THE PHILIPPINES-DILIMAN, QUEZON CITY 1101, PHILIPPINES

Email address: rguadalupe@math.upd.edu.ph