

# Fractal dimension and the counting rule of the Goldstone modes

Qian-Qian Shi,<sup>1</sup> Yan-Wei Dai,<sup>1</sup> Huan-Qiang Zhou,<sup>1</sup> and Ian P. McCulloch<sup>2,3,1</sup>

<sup>1</sup>*Centre for Modern Physics, Chongqing University, Chongqing 400044, The People's Republic of China*

<sup>2</sup>*Department of Physics, National Tsing Hua University, Hsinchu 30013, Taiwan*

<sup>3</sup>*School of Mathematics and Physics, The University of Queensland, St. Lucia, QLD 4072, Australia*

It is argued that highly degenerate ground states arising from spontaneous symmetry breaking with a type-B Goldstone mode are scale-invariant, with a salient feature that the entanglement entropy  $S(n)$  scales logarithmically with the block size  $n$  in the thermodynamic limit. As it turns out, the prefactor is half the number of type-B Goldstone modes  $N_B$ . This is achieved by performing an exact singular value decomposition of the degenerate ground states, thus unveiling their self-similarities - the essence of a fractal. Combining with a field-theoretic prediction [O. A. Castro-Alvaredo and B. Doyon, *Phys. Rev. Lett.* **108**, 120401 (2012)], we are led to the identification of the fractal dimension  $d_f$  with the number of type-B Goldstone modes  $N_B$  for quantum many-body systems undergoing spontaneous symmetry breaking.

## I. INTRODUCTION

Spontaneous symmetry breaking (SSB) is a key ingredient in diverse areas of physics, ranging from condensed matter to field theories. In particular, the emergence of a gapless Goldstone mode (GM), when a continuous symmetry group is spontaneously broken, is of paramount importance, due to its relevance to the low-energy physics. As first stated by Goldstone [1], the number of the GMs is equal to the number of broken symmetry generators  $N_{BG}$  for a relativistic system undergoing SSB. However, complications arise for a nonrelativistic system, as far as the connection between the number of broken symmetry generators  $N_{BG}$  and the number of the GMs is concerned. Since an early work by Nielsen and Chadha [2], much attention has been paid to a proper classification of the GMs [3–9], culminating in the introduction of type-A and type-B GMs [7, 8], based on a previous observation made by Nambu [10]. In this classification, the so-called Watanabe-Brauner matrix [6] plays a crucial role. As a result, when the symmetry group  $G$  is spontaneously broken into  $H$ , the counting rule for the GMs may be formulated as follows

$$N_A + 2N_B = N_{BG}, \quad (1)$$

where  $N_A$  and  $N_B$  are, respectively, the numbers of type-A and type-B GMs, and  $N_{BG}$  is equal to the dimension of the coset space  $G/H$ .

One remarkable distinction may be made between type-A and type-B GMs, since SSB with a type-A GM only happens in the thermodynamic limit, in contrast to SSB with a type-B GM, which survives in a finite-size system. Instead, a finite-size precursor to SSB with a type-A GM appears in the guise of the so-called Anderson tower, first developed in spin wave theory for antiferromagnetism [11]. Meanwhile, a significant development has been achieved to describe SSB with a type-A GM from the perspective of quantum entanglement [12, 13]. However, a systematic investigation is still

lacking for SSB with a type-B GM from the perspective of the entanglement entropy, with a few notable exceptions [14–16], in which the entanglement entropy is discussed for the ferromagnetic states. The other distinction between type-A and type-B GMs concerns their instabilities under quantum fluctuations. In fact, SSB with a type-A GM is forbidden in one spatial dimension, as a result of the Mermin-Wagner-Coleman theorem [17], whereas SSB with a type-B GM survives quantum fluctuations even in one spatial dimension. As a consequence, instead of long-range order resulted from SSB with a type-A GM, there exists only quasi-long-range order in one spatial dimension, which may be characterized by means of conformal field theory [18]. Historically, conformal field theory originated from a speculation made by Polyakov [19] that scale invariance implies conformal invariance, which itself has attracted much attention, in an attempt to prove or disprove it [20–22]. In this regard, an intriguing question arises as to whether or not there is any scale-invariant, but not conformally invariant state, if one takes into account SSB with a type-B GM, which survives even in one spatial dimension.

This work attempts to address this question through a thorough investigation of the scaling behavior of the entanglement entropy for one-dimensional quantum many-body systems undergoing SSB with a type-B GM. We demonstrate that highly degenerate ground states arising from SSB with a type-B GM admit an exact singular value decomposition [23], thus unveiling their self-similarities - the essence of a fractal, characterized in term of the fractal dimension  $d_f$ . As a consequence, the degenerate ground states are scale-invariant, which in turn implies that the entanglement entropy  $S(n)$  scales logarithmically with the block size  $n$  in the thermodynamic limit. As it turns out, the prefactor is half the number of type-B GMs  $N_B$ . Combining with a field-theoretic prediction [16] that the prefactor is half the fractal dimension  $d_f$ , we are led to the identification of the fractal dimension  $d_f$  with the number of type-B GMs  $N_B$ . As an illustration, we investigate the  $SU(2)$  spin- $s$  ferromagnetic model, the  $SU(N+1)$  ferromagnetic model and the  $SU(2)$  spin-1 anisotropic biquadratic model, with the

fractal dimension  $d_f$  being 1,  $N$ , and 1, respectively.

## II. THE ENTANGLEMENT ENTROPY FOR SCALE-INVARIANT STATES

Consider a translation-invariant quantum many-body system, described by the Hamiltonian  $\mathcal{H}$ , with the symmetry group  $SU(N+1)$ , on a one-dimensional lattice. Throughout this work, the size  $L$  is assumed to be even. The symmetry group  $SU(N+1)$  has  $(N+1)^2-1$  generators, and the rank is  $N$ . Accordingly, there are  $N$  commuting Cartan generators [24]  $H_\alpha$  ( $\alpha = 1, \dots, N$ ), which are traceless and diagonal. For each  $H_\alpha$ , there exists a conjugate pair of a raising operator  $E_\alpha$  and a lowering operator  $F_\alpha$  such that  $[H_\alpha, E_\beta] = \beta_\alpha E_\beta$ ,  $[E_\alpha, F_\alpha] = (E_\alpha, F_\alpha)H_\alpha$ ,  $[F_\alpha, H_\beta] = \beta_\alpha F_\beta$ ,  $[E_\alpha, E_\beta] = g_{\alpha, \beta} E_\gamma \delta_{\gamma, \alpha+\beta}$ , and  $[F_\alpha, F_\beta] = g_{-\alpha, -\beta} F_\gamma \delta_{\gamma, \alpha+\beta}$ , with  $\beta$  being the root matrix,  $g_{\alpha, \beta}$  depending on the specific form of the Cartan generators, and  $(E_\alpha, F_\alpha)$  being the Killing form of  $E_\alpha$  and  $F_\alpha$ . Here, we stress that, it is convenient to choose the Cartan generators  $H_\alpha$  in such a way that the set of the lowering operators  $F_\alpha$  commute to each other.

Suppose the symmetry group  $SU(N+1)$  is spontaneously broken into  $SU(N) \times U(1)$ . For simplicity, we assume that the translational symmetry under the one-site translation is not spontaneously broken. Otherwise, a unit cell is needed. With this fact in mind, the highest weight state  $|\text{hws}\rangle$ , which itself is an unentangled ground state, takes the form  $|\text{hws}\rangle = |hh\dots h\rangle$ , with a local component  $|h\rangle_j$  being the eigenvector for  $H_{\alpha,j}$ , satisfying  $E_{\alpha,j}|h\rangle_j = 0$ , but  $F_{\alpha,j}|h\rangle_j \neq 0$ . Here,  $H_{\alpha,j}$ ,  $E_{\alpha,j}$ , and  $F_{\alpha,j}$  represent the local counterparts of the Cartan generators  $H_\alpha$ , the raising operators  $E_\alpha$  and the lowering operators  $F_\alpha$  at a lattice site  $j$  on a one-dimensional lattice:  $H_\alpha = \sum_j H_{\alpha,j}$ ,  $E_\alpha = \sum_j E_{\alpha,j}$  and  $F_\alpha = \sum_j F_{\alpha,j}$ . For the symmetry generators

$E_\alpha$  and  $F_\alpha$ , one may choose  $F_{\alpha,j}$  and  $E_{\alpha,j}$  as the interpolating fields [2, 25], respectively. Given  $\langle [E_\alpha, F_{\alpha,j}] \rangle \propto \langle H_{\alpha,j} \rangle$ ,  $\langle [E_{\alpha,j}, F_\alpha] \rangle \propto \langle H_{\alpha,j} \rangle$  and  $\langle H_{\alpha,j} \rangle \neq 0$ , the  $2N$  symmetry generators  $E_\alpha$  and  $F_\alpha$  are spontaneously broken, with  $\langle H_{\alpha,j} \rangle$  being a local order parameter. Here, the expectation value  $\langle O \rangle$  of an operator  $O$  is taken over the highest weight state  $|\text{hws}\rangle$ . Since no type-A GM survives in one spatial dimension [17], the number of type-A GMs  $N_A$  must be 0. Therefore, according to the counting rule (1), the  $N$  type-B GMs emerge. For a later use, we introduce  $q_\alpha$  to denote the power of  $F_{\alpha,j}$  such that  $F_{\alpha,j}^{q_\alpha} |h\rangle_j \neq 0$ , but  $F_{\alpha,j}^{q_\alpha+1} |h\rangle_j = 0$ .

A sequence of degenerate ground states  $|L, M_1, \dots, M_N\rangle$  are generated from the repeated action of the lowering operators  $F_\alpha$  on the highest weight state  $|\text{hws}\rangle$ :

$$|L, M_1, \dots, M_N\rangle = \frac{1}{Z(L, M_1, \dots, M_N)} \prod_{\alpha=1}^N F_\alpha^{M_\alpha} |\text{hws}\rangle, \quad (2)$$

where  $Z(L, M_1, \dots, M_N)$  is introduced to ensure that  $|L, M_1, \dots, M_N\rangle$  is normalized.

In order to understand SSB with a type-B GM from the perspective of quantum entanglement, the system is partitioned into a block  $B$  and its environment  $E$ . Here, the block  $B$  consists of  $n$  lattice sites that are not necessarily contiguous, with the rest  $L-n$  lattice sites constituting the environment  $E$ . As a convention,  $n \leq L/2$ . Note that  $|\text{hws}\rangle$ , as an unentangled product state, is split into  $|\text{hws}\rangle_B$  and  $|\text{hws}\rangle_E$ . With this in mind, we introduce the counterparts of the symmetry group  $SU(N+1)$  in the block  $B$  and the environment  $E$ , respectively. In particular, the counterparts of the lowering operators  $F_\alpha$  are  $F_{\alpha,B}$  and  $F_{\alpha,E}$ , respectively, in the block  $B$  and the environment  $E$ . This in turn allows us to define the basis states  $|n, k_1, \dots, k_N\rangle$  and  $|L-n, M_1-k_1, \dots, M_N-k_N\rangle$  for the block  $B$  and the environment  $E$ , which take the same form as Eq. (2), with  $F_\alpha$  replaced by  $F_{\alpha,B}$  and  $F_{\alpha,E}$ ,  $M_\alpha$  replaced by  $k_\alpha$  and  $M_\alpha - k_\alpha$ , respectively. Meanwhile,  $Z(n, k_1, \dots, k_N)$  and  $Z(L-n, M_1-k_1, \dots, M_N-k_N)$  need to be introduced to ensure that  $|n, k_1, \dots, k_N\rangle$  and  $|L-n, M_1-k_1, \dots, M_N-k_N\rangle$  are normalized.

A remarkable fact is that  $|L, M_1, \dots, M_N\rangle$  admit an exact singular value decomposition:

$$|L, M_1, \dots, M_N\rangle = \prod_{\alpha=1}^N \sum_{k_\alpha=0}^{\min(M_\alpha, q_\alpha n)} \lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) |n, k_1, \dots, k_N\rangle |L-n, M_1-k_1, \dots, M_N-k_N\rangle, \quad (3)$$

where the singular values  $\lambda(L, k_1, \dots, k_N, M_1, \dots, M_N)$  take the form

$$\lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) = \prod_{\alpha=1}^N C_{M_\alpha}^{k_\alpha} \frac{Z(n, k_1, \dots, k_N) Z(L-n, M_1-k_1, \dots, M_N-k_N)}{Z(L, M_1, \dots, M_N)}, \quad (4)$$

where  $C_{M_\alpha}^{k_\alpha}$  is the binomial coefficients:  $C_{M_\alpha}^{k_\alpha} = M_\alpha! / (k_\alpha! (M_\alpha - k_\alpha)!)$ . This decomposition reflects the self-similarities underlying a fractal, which is characterized in terms of the fractal dimension  $d_f$ , since both the block  $B$  and the environment  $E$ , as a subsystem, share the same type of quantum states as the entire system. This explains why the fractal dimension  $d_f$ , as already introduced in a field-theoretic approach to the ferromagnetic states [16], furnishes a proper description for scale-invariant states.

The entanglement entropy  $S_L(n, M_1, \dots, M_N)$  follows from the reduced density matrix  $\rho_L(n, M_1, \dots, M_N)$ :  $S_L(n, M_1, \dots, M_N) = -\text{Tr}[\rho_L(n, M_1, \dots, M_N) \log_2 \rho_L(n, M_1, \dots, M_N)]$ , with

$$\rho_L(n, M_1, \dots, M_N) = \prod_{\alpha=1}^N \sum_{k_\alpha=0}^{\min(M_\alpha, q_\alpha n)} \Lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) |n, k_1, \dots, k_N\rangle \langle n, k_1, \dots, k_N|.$$

Here,  $\Lambda(L, k_1, \dots, k_N, M_1, \dots, M_N)$  are the eigenvalues of the reduced density matrix  $\rho_L(n, M_1, \dots, M_N)$ :  $\Lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) = [\lambda(L, k_1, \dots, k_N, M_1, \dots, M_N)]^2$ . Therefore,  $S_L(n, M_1, \dots, M_N)$  may be rewritten as

$$S_L(n, M_1, \dots, M_N) = - \prod_{\alpha=1}^N \sum_{k_\alpha=0}^{\min(M_\alpha, q_\alpha n)} \Lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) \log_2 \Lambda(L, k_1, \dots, k_N, M_1, \dots, M_N). \quad (5)$$

This makes it possible to perform a systematic analysis of the block entanglement entropy  $S_L(n, M_1, \dots, M_N)$ , depending on a specific realization of the symmetry group  $\text{SU}(N+1)$  for the Hamiltonian  $\mathcal{H}$  under investigation.

Instead, we turn to a scaling analysis of the entanglement entropy  $S(n, M_1, \dots, M_N)$  for a scale-invariant state in the thermodynamic limit. For this purpose, we introduce the fillings  $f_\alpha = M_\alpha/L$  ( $\alpha = 1, \dots, N$ ) to ensure that  $f_\alpha$  are kept constant, when  $L$  tends to infinity. To ease the notations, we denote  $S(n)$  as the block entanglement entropy for a specific choice of the fillings  $f_\alpha$  ( $\alpha = 1, \dots, N$ ). A heuristic physical argument (cf. Sec. A of the Supplemental Material (SM)) implies that  $S(n)$  scales logarithmically with  $n$ ,

$$S(n) = \frac{N_B}{2} \log_2 n + S_0, \quad (6)$$

where  $S_0$  is an additive contribution to the entanglement entropy, which only depends on the fillings  $f_\alpha$  ( $\alpha = 1, \dots, N$ ). Here, we have assumed that the fillings  $f_\alpha$  ( $\alpha = 1, \dots, N$ ) are nonzero. Combining with a field-theoretic prediction that the prefactor is half the fractal dimension  $d_f$  [16], we conclude that the fractal dimension  $d_f$  is identical to the number of type-B GMs  $N_B$ .

### III. THE $\text{SU}(2)$ FERROMAGNETIC STATES: ARBITRARY SPIN $s$

Consider the  $\text{SU}(2)$  spin- $s$  ferromagnetic Heisenberg model with the nearest-neighbor interaction, described by the Hamiltonian

$$\mathcal{H} = - \sum_{j=1}^L \mathbf{S}_j \cdot \mathbf{S}_{j+1}, \quad (7)$$

where  $\mathbf{S}_j = (S_{x,j}, S_{y,j}, S_{z,j})$ , and  $S_{x,j}$ ,  $S_{y,j}$  and  $S_{z,j}$  represent the spin- $s$  operators at the  $j$ -th site. Here, the symmetry group  $\text{SU}(2)$  is generated by  $S_+ = \sum_j S_{+,j}$ ,  $S_- = \sum_j S_{-,j}$  and  $S_z = \sum_j S_{z,j}$ :  $[S_z, S_+] = S_+$ ,  $[S_+, S_-] = S_z$  and  $[S_-, S_z] = S_-$ ,

with  $S_{+,j}$  and  $S_{-,j}$  being defined by  $S_{\pm,j} = (S_{x,j} \pm iS_{y,j})/\sqrt{2}$ . Suppose  $|m\rangle_j$  are the eigenvectors of  $S_{z,j}$ :  $S_{z,j}|m\rangle_j = m|m\rangle_j$ , with  $m = -s, \dots, s$ . Then, the local Hilbert space constitutes a  $2s+1$ -dimensional irreducible representation of  $\text{SU}(2)$  at each lattice site  $j$ . The action of  $S_{-,j}$  and  $S_{+,j}$  on  $|m\rangle_j$  takes the form:  $S_{-,j}|m\rangle_j = \sqrt{(s+m)(s-m+1)/2}|m-1\rangle_j$  and  $S_{+,j}|m\rangle_j = \sqrt{(s-m)(s+m+1)/2}|m+1\rangle_j$ , respectively. Thus, the highest weight state  $|hws\rangle$  is  $|hws\rangle = |ss\dots s\rangle$ . Since  $S_-^{2s}|s\rangle \neq 0$ , but  $S_-^{2s+1}|s\rangle = 0$ , we have  $q = 2s$ . Note that the model is exactly solvable by means of the Bethe ansatz only when  $s = 1/2$ .

The interpolating fields are  $S_{+,j}$  and  $S_{-,j}$ , for the generator  $S_-$  and  $S_+$ , respectively. Thus,  $\langle S_{z,j} \rangle$  is the local order parameter, given  $\langle [S_{+,j}, S_-] \rangle = \langle [S_+, S_{-,j}] \rangle = \langle S_{z,j} \rangle \neq 0$ . Hence, the two generators  $S_-$  and  $S_+$  are spontaneously broken. According to the counting rule (1), the number of type-B GM is one. Therefore, the model is a specific realization of the general scheme:  $H_1 = S_z$ ,  $E_1 = S_+$  and  $F_1 = S_-$ .

The degenerate ground states  $|L, M\rangle$  are generated from the repeated action of the lowering operator  $S_-$  on the highest weight state  $|ss\dots s\rangle$ :

$$|L, M\rangle = \frac{1}{Z(L, M)} S_-^M |ss\dots s\rangle, \quad (8)$$

where

$$Z(L, M) = \frac{M!}{\sqrt{2^M}} \sqrt{\sum'_{N_{-s}, \dots, N_s} \prod_{r=-s}^{s-1} [\varepsilon(s, r)]^{N_r} C_{L-\sum_{m=-s}^s N_m}^{N_r}},$$

with

$$\varepsilon(s, r) = \frac{\prod_{m=r+1}^s (s+m)(s-m+1)}{\prod_{m=r}^{s-1} (s-m)^2}.$$

Here,  $\sum'_{N_{-s}, \dots, N_s}$  is taken over all the possible values of  $N_{-s}, \dots, N_s$ , subject to the constraints:  $\sum_{m=-s}^s N_m = L$  and  $\sum_{m=-s}^s (s-m)N_m = M$ . We remark that  $|L, M\rangle$  ( $M = 0, \dots, 2sL$ ) span a  $2sL+1$ -dimensional irreducible representation of the symmetry group  $\text{SU}(2)$ . A derivation of the concrete expression for  $Z(L, M)$  is presented in Sec. B of the SM.

The degenerate ground states  $|L, M\rangle$  admit an exact singular value decomposition:

$$|L, M\rangle = \sum_{k=0}^{\min(M, 2sn)} \lambda(L, k, M) |n, k\rangle |L-n, M-k\rangle,$$

where the singular values  $\lambda(L, k, M)$  take the form

$$\lambda(L, k, M) = \frac{\mu(L, k, M)}{\nu(L, k, M)},$$

with

$$\mu(L, k, M) = \sqrt{\sum'_{\substack{n_{-s}, \dots, n_s \\ l_{-s}, \dots, l_s}} \prod_{r,t=-s}^{s-1} \varepsilon(s, r)^{n_r} C_{n-\sum_{m=-s}^{s-1} n_m}^{n_r} \varepsilon(s, t)^{l_t} C_{L-n-\sum_{m=-s}^{s-1} l_m}^{l_t}},$$

and

$$\nu(L, k, M) = \sqrt{\sum'_{N_{-s}, \dots, N_s} \prod_{r=-s}^{s-1} \varepsilon(s, r)^{N_r} C_{L-\sum_{m=-s}^{s-1} N_m}^{N_r}}.$$

Here,  $\sum'_{n_{-s}, \dots, n_s}$  is taken over all the possible values of  $n_{-s}, \dots, n_s$ , subject to the constraints:  $\sum_{m=-s}^s n_m = n$  and  $\sum_{m=-s}^s (s-m)n_m = k$ , and  $\sum'_{l_{-s}, \dots, l_s}$  is taken over all the possible values of  $l_{-s}, \dots, l_s$ , subject to the constraints:  $\sum_{m=-s}^s l_m = L-n$  and  $\sum_{m=-s}^s (s-m)l_m = M-k$ . Then, the eigenvalues  $\Lambda(L, M, k)$  of the reduced density matrix  $\rho_L(n, M)$  follows from  $\Lambda(L, M, k) = [\lambda(L, M, k)]^2$ . Note that the same results for spin  $s = 1/2$ , presented in Ref. [14], are reproduced. Hence, the entanglement entropy  $S_L(n, M)$  follows from Eq. (5). In particular, the logarithmic scaling behaviour may be confirmed from an analytical treatment, based on the Stirling's approximation [26], as done in Ref. [14] for spin  $s = 1/2$ .

In order to understand how the logarithmic scaling behaviour emerges in the thermodynamic limit, we plot  $S_L(n, M)$  vs  $\log_2 n$  in Fig. 1: (a) For  $s = 1$ ,  $M = L/4$ , when  $L$  is varied:  $L = 100, 200, 500$  and  $1000$ . A significant deviation from the logarithmic scaling behaviour is observed when  $L$  is relatively small, but tends to vanish, as  $L$  increases. Indeed, the prefactor is close to the exact value  $1/2$ , with an error being less than 1.1%, when  $L = 1000$ :  $S_{1000}(n, 250) = 0.505 \log_2 n + 0.891$ . (b) For  $s = 1$ ,  $L = 1000$ , when  $M$  is varied:  $M = 250, 500, 750$  and  $1000$ . This amounts to varying the filling  $f$ . The prefactor is close to  $1/2$ , regardless of the values of the filling  $f$ , within an error less than 2.3%. That is, the contribution from the filling  $f$  goes to a non-universal additive constant, as anticipated. (c) For  $M = 250$ ,  $L = 1000$ , when  $s$  is varied:  $s = 1/2, 1, 3/2, 2, 5/2$  and  $3$ . The prefactor is close to  $1/2$  for any spin  $s$ , within an error less than 2.2%. Here,  $n$  ranges from 10 to 40.




FIG. 1. The entanglement entropy  $S_L(n, M)$  vs  $\log_2 n$  for the spin- $s$  SU(2) ferromagnetic states: (a) For  $s = 1$ ,  $M = L/4$ , when  $L$  is varied:  $L = 100, 200, 500$  and  $1000$ . A significant deviation from the logarithmic scaling behaviour is observed when  $L$  is relatively small, but tends to vanish, as  $L$  increases. The prefactor is close to the exact value  $1/2$ , with an error being less than 1.1%, when  $L = 1000$ :  $S_{1000}(n, 250) = 0.505 \log_2 n + 0.891$ . (b) For  $s = 1$ ,  $L = 1000$ , when  $M$  is varied:  $M = 250, 500, 750$  and  $1000$ . The prefactor is close to  $1/2$ , within an error less than 2.3%. (c) For  $M = 250$ ,  $L = 1000$ , when  $s$  is varied:  $s = 1/2, 1, 3/2, 2, 5/2$  and  $3$ . The prefactor is close to  $1/2$  for any spin  $s$ , within an error less than 2.2%. Here,  $n$  ranges from 10 to 40.

#### IV. THE SU(N + 1) FERROMAGNETIC STATES: FUNDAMENTAL REPRESENTATION

The SU(N + 1) ferromagnetic model is described by the Hamiltonian

$$\mathcal{H} = - \sum_j P_{j, j+1}. \quad (11)$$

Here,  $P$  is the permutation operator:  $P = \sum_{u,v=1}^{N+1} e_{uv} \otimes e_{vu}$ , where  $e_{uv} = |u\rangle\langle v|$ , with  $|u\rangle$  and  $|v\rangle$  being the  $u$ -th and  $v$ -th




FIG. 2. (a) The entanglement entropy  $S_L(n, M_1, M_2)$  vs  $\log_2 n$  for the SU(3) ferromagnetic states, with  $M_1 = L/2$  and  $M_2 = L/4$ . (b) The entanglement entropy  $S_L(n, M_1, M_2, M_3)$  vs  $\log_2 n$  for the SU(4) ferromagnetic states, with  $M_1 = M_2 = M_3 = L/4$ . (c) The entanglement entropy  $S_L(n, M_1, M_2, M_3, M_4)$  vs  $\log_2 n$  for the SU(5) ferromagnetic states, with  $M_1 = M_2 = M_3 = M_4 = L/5$ . Here,  $L$  is varied:  $L = 100, 200, 500$  and  $1000$ . A significant deviation from the logarithmic scaling behaviour is observed when  $L$  is relatively small, but tends to vanish, as  $L$  increases. The prefactor is close to the exact value  $N_B/2$ , with an error being less than 2.3%, when  $L = 1000$ :  $S_{1000}(n, 500, 250) = 0.999 \log_2 n + 1.542$ ,  $S_{1000}(n, 250, 250, 250) = 1.509 \log_2 n + 2.007$  and  $S_{1000}(n, 200, 200, 200, 200) = 2.045 \log_2 n + 2.023$ , with  $N_B$  being 2, 3 and 4, respectively, for the SU(3), SU(4) and SU(5) ferromagnetic states. Here,  $n$  ranges from 10 to 40.

states in an orthonormal basis. Physically, the permutation

operator  $P$  may be realized in terms of the spin- $s$  operators  $\mathbf{S} = (S_x, S_y, S_z)$ , with  $N = 2s$ :

$$P = \sum_{r=0}^{2s} (-1)^{2s+r} \prod_{m \neq r}^{2s} \frac{2(\mathbf{S} \otimes \mathbf{S}) - m(m+1) + 2s(s+1)}{r(r+1) - m(m+1)}.$$

Note that, when  $N = 2$ , it is the SU(3) ferromagnetic point for the spin-1 bilinear-biquadratic model [27]. The model is exactly solvable by means of the Bethe ansatz [28].

The model possesses the symmetry group  $SU(N+1)$ , with the local Hilbert space being the fundamental representation space of  $SU(N+1)$  at each lattice site  $j$ , thus constituting a specific realization of the general scheme. The Cartan generators  $H_\alpha = \sum_j H_{\alpha,j}$  may be chosen as  $H_{\alpha,j} = e_{11,j} - e_{\alpha+1,\alpha+1,j}$  for  $\alpha = 1, \dots, N$ . For each  $H_\alpha$ , the lowering operator and the raising operator may be chosen as:  $F_\alpha = \sum_j F_{\alpha,j}$ ,  $E_\alpha = \sum_j E_{\alpha,j}$ , with  $F_{\alpha,j} = e_{\alpha+1,1,j}$  and  $E_{\alpha,j} = e_{1,\alpha+1,j}$ , satisfying  $[H_\alpha, E_\alpha] = 2E_\alpha$ ,  $[E_\alpha, F_\alpha] = H_\alpha$  and  $[F_\alpha, H_\alpha] = 2F_\alpha$ . Define  $|\beta\rangle$  as a  $N+1$ -dimensional vector, with the  $\beta$ -th entry being 1 and the others being 0. Then,  $|\beta\rangle_j$  are the eigenvectors of  $H_{\alpha,j}$ :  $H_{\alpha,j}|\beta\rangle_j = (\delta_{1\beta} - \delta_{\alpha+1,\beta})|\beta\rangle_j$ , for  $\beta = 1, \dots, N+1$ . The action of  $F_{\alpha,j}$  and  $E_{\alpha,j}$  on  $|\beta\rangle_j$  takes the form:  $F_{\alpha,j}|\beta\rangle_j = |\alpha+1\rangle_j$  and  $E_{\alpha,j}|\beta\rangle_j = 0$ . Therefore, the highest weight state  $|\text{hws}\rangle$  is  $|\text{hws}\rangle = |11\dots 1\rangle$ . The interpolating fields are  $E_{\alpha,j}$  and  $F_{\alpha,j}$ , for  $F_\alpha$  and  $E_\alpha$ , respectively. Thus,  $\langle H_{\alpha,j} \rangle$  ( $\alpha = 1, \dots, N$ ) are the local order parameters, given  $\langle [E_{\alpha,j}, F_\alpha] \rangle = \langle [E_\alpha, F_{\alpha,j}] \rangle = \langle H_{\alpha,j} \rangle \neq 0$ . Hence, the  $2N$  symmetry generators  $E_\alpha$  and  $F_\alpha$  are spontaneously broken. According to the counting rule (1), the  $2N$  broken generators yield  $N$  type-B GMs. In addition, since  $F_{\alpha,j}|1\rangle_j \neq 0$ , but  $F_{\alpha,j}^2|1\rangle_j = 0$ , we have  $q_\alpha = 1$  ( $\alpha = 1, \dots, N$ ).

The degenerate ground states  $|L, M_1, \dots, M_N\rangle$  are generated from the repeated action of the lowering operators  $F_\alpha$  on the highest weight state  $|11\dots 1\rangle$ :

$$|L, M_1, \dots, M_N\rangle = \frac{1}{Z(L, M_1, \dots, M_N)} \prod_{\alpha=1}^N F_\alpha^{M_\alpha} |11\dots 1\rangle, \quad (12)$$

which span an irreducible representation of the symmetry group  $SU(N+1)$ , with the dimension being  $C_{L+N}^N$ . Here,  $Z(L, M_1, \dots, M_N)$  takes the form,

$$Z(L, M_1, \dots, M_N) = \prod_{\alpha=1}^N M_\alpha! \sqrt{C_{L-\sum_{\beta=1}^N M_\beta}^{M_\alpha}}.$$

A derivation of the concrete expression for  $Z(L, M_1, \dots, M_N)$  is presented in Sec. C of the SM.

The degenerate ground states  $|L, M_1, \dots, M_N\rangle$  admit an exact singular value decomposition:

$$|L, M_1, \dots, M_N\rangle = \prod_{\alpha=1}^N \sum_{k_\alpha=0}^{\min(M_\alpha, n)} \lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) |n, k_1, \dots, k_N\rangle |L-n, M_1-k_1, \dots, M_N-k_N\rangle, \quad (13)$$

where the singular values  $\lambda(L, k_1, \dots, k_N, M_1, \dots, M_N)$  take the form

$$\lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) = \sqrt{\frac{\prod_{\alpha=1}^N C_{n-\sum_{\beta=1}^{\alpha-1} k_{\beta}}^{k_{\alpha}} \prod_{\gamma=1}^N C_{L-n-\sum_{\beta=1}^{\gamma-1} (M_{\beta}-k_{\beta})}^{M_{\gamma}-k_{\gamma}}}{\prod_{\alpha=1}^N C_{L-\sum_{\beta=1}^{\alpha-1} M_{\beta}}^{M_{\alpha}}}}. \quad (14)$$

Therefore, the eigenvalues  $\Lambda(L, k_1, \dots, k_N, M_1, \dots, M_N)$  of the reduced density matrix  $\rho_L(n, M_1, \dots, M_N)$  are  $\Lambda(L, k_1, \dots, k_N, M_1, \dots, M_N) = [\lambda(L, k_1, \dots, k_N, M_1, \dots, M_N)]^2$ . Hence, the entanglement entropy  $S_L(n, M_1, \dots, M_N)$  follows from Eq. (5). Thus, we reproduce the eigenvalues of the reduced density matrix  $\rho_L(n, M_1, \dots, M_N)$  in Ref. [15]. In particular, the logarithmic scaling behaviour for the entanglement entropy  $S_L(n, M_1, \dots, M_N)$  may be confirmed from an analytical treatment [15].

In order to understand how the logarithmic scaling behaviour emerges in the thermodynamic limit, we plot  $S_L(n, M_1, M_2)$  vs  $\log_2 n$  in Fig. 2 (a) for the SU(3) ferromagnetic states, with  $M_1 = L/2$  and  $M_2 = L/4$ ,  $S_L(n, M_1, M_2, M_3)$  vs  $\log_2 n$  in Fig. 2 (b) for the SU(4) ferromagnetic states, with  $M_1 = M_2 = M_3 = L/4$ , and  $S_L(n, M_1, M_2, M_3, M_4)$  vs  $\log_2 n$  in Fig. 2 (c) for the SU(5) ferromagnetic states, with  $M_1 = M_2 = M_3 = M_4 = L/5$ , when  $L$  is varied:  $L = 100, 200, 500$  and  $1000$ . A significant deviation from the logarithmic scaling behaviour is observed when  $L$  is relatively small, but tends to vanish, as  $L$  increases. The prefactor is close to the exact value  $N_B/2$ , with an error being less than 2.3%, when  $L = 1000$ :  $S_{1000}(n, 500, 250) = 0.999 \log_2 n + 1.542$ ,  $S_{1000}(n, 250, 250, 250) = 1.509 \log_2 n + 2.007$  and  $S_{1000}(n, 200, 200, 200, 200) = 2.045 \log_2 n + 2.023$ , with  $N_B$  being 2, 3 and 4, respectively, for the SU(3), SU(4) and SU(5) ferromagnetic states. Here,  $n$  ranges from 10 to 40.

## V. THE COEXISTING FRACTAL STATES: AN EXAMPLE BEYOND SIMPLE FERROMAGNETISM

Consider the SU(2) spin-1 anisotropic biquadratic model [27], described by the Hamiltonian

$$\mathcal{H} = \sum_j (J_x S_{x,j} S_{x,j+1} + J_y S_{y,j} S_{y,j+1} + J_z S_{z,j} S_{z,j+1})^2, \quad (15)$$

where  $S_{x,j}$ ,  $S_{y,j}$ , and  $S_{z,j}$  are the spin-1 operators at a lattice site  $j$ , and  $J_x$  and  $J_z$  are the anisotropic coupling parameters. The model possesses the staggered symmetry group SU(2) generated by  $K_x$ ,  $K_y$ , and  $K_z$ :  $K_x = \sum_j K_{x,j}$ ,  $K_y = \sum_j K_{y,j}$  and  $K_z = \sum_j K_{z,j}$ , with  $K_{x,j} = \sum_j (-1)^j [S_{x,j}^2 - S_{y,j}^2]/2$ ,  $K_{y,j} = \sum_j (-1)^j (S_{x,j} S_{y,j} + S_{y,j} S_{x,j})/2$  and  $K_{z,j} = \sum_j S_{z,j}/2$ . Accordingly, one may define the raising operator  $K_+ = \sum_j K_{+,j}$  and the lowering operator  $K_- = \sum_j K_{-,j}$ , with  $K_{\pm,j} = (K_{x,j} \pm iK_{y,j})/\sqrt{2}$ :  $[K_z, K_+] = K_+$ ,  $[K_+, K_-] = K_z$  and  $[K_-, K_z] = K_-$ . In addition, it enjoys two extra U(1) symmetry groups, generated by  $K_1$  and  $K_2$ :  $K_1 = \sum_j (-1)^j [S_{y,j}^2 - S_{z,j}^2]/2$  and  $K_2 = \sum_j (-1)^j [S_{z,j}^2 - S_{x,j}^2]/2$ , respectively. Since  $K_1 + K_2 + K_x = 0$ ,

we only need to consider one U(1) symmetry group, generated by  $\sum_j (-1)^j S_{z,j}^2$ , due to the constraints:  $S_{x,j}^2 + S_{y,j}^2 + S_{z,j}^2 = 2$ .

For  $J_x > J_z > 0$ , there are two distinct choices for the highest weight state |hws>: (i) |hws> =  $|1_z \dots 1_z\rangle$  and (ii) |hws> =  $|0_x 0_y \dots 0_x 0_y\rangle$ , in the sense that the first choice is invariant under the one-site translation, whereas the second choice is not invariant under the one-site translation. Here,  $|1_z\rangle$  is the eigenvector of  $S_{z,j}$ , with the eigenvalue being 1, and  $|0_x\rangle/|0_y\rangle$  is the eigenvector of  $S_{x,j}/S_{y,j}$ , with the eigenvalue being 0. However, the two choices are unitarily equivalent under a local unitary transformation  $U$ :  $K_{x,j} \rightarrow K_{y,j}$ ,  $K_{y,j} \rightarrow K_{z,j}$  and  $K_{z,j} \rightarrow K_{x,j}$ . As a consequence, the entanglement entropy  $S(n)$  for the degenerate ground states, corresponding to the two choices, must be identical. Therefore, we only need to focus on the first choice for brevity. Note that the action of  $K_{-,j}$  and  $K_{+,j}$  on  $|1_z\rangle_j$  takes the form:  $K_{-,j}|1_z\rangle_j = (-1)^j \sqrt{2}/2 | -1_z\rangle_j$  and  $K_{+,j}|1_z\rangle_j = 0$ .

The interpolating fields are  $K_{+,j}$  and  $K_{-,j}$ , for the generator  $K_-$  and the generator  $K_+$ , respectively. Thus,  $\langle K_{z,j} \rangle$  is the local order parameter, given  $\langle [K_{+,j}, K_-] \rangle = \langle [K_+, K_{-,j}] \rangle = \langle K_{z,j} \rangle \neq 0$ . Therefore, the two symmetry generators  $K_-$  and  $K_+$  are spontaneously broken. According to the counting rule (1), there is one type-B GM. In addition, since  $K_{-,j}|1_z\rangle_j \neq 0$ , but  $K_{-,j}^2|1_z\rangle_j = 0$ , we have  $q = 1$ . This is a specific realization of the general scheme:  $K_+ = E_1$ ,  $K_- = F_1$  and  $K_z = H_1$ .

A sequence of degenerate ground states  $|L, M\rangle$  are generated from the repeated action of the lowering operator  $K_-$  on the highest weight state  $|1_z \dots 1_z\rangle$ :

$$|L, M\rangle = \frac{1}{Z(L, M)} K_-^M |1_z \dots 1_z\rangle, \quad (16)$$

with

$$Z(L, M) = M! \sqrt{\frac{C_L^M}{2^M}}.$$

We remark that  $|L, M\rangle$  ( $M = 0, \dots, L$ ) span a  $L+1$ -dimensional irreducible representation of the symmetry group SU(2). A derivation of the concrete expression for  $Z(L, M)$  is presented in Sec. D of the SM.




FIG. 3. The entanglement entropy  $S_L(n, M)$  vs  $\log_2 n$  for the coexisting fractal states: (a)  $M = L/2$  and (b)  $M = L/4$ , when  $L$  is varied:  $L = 100, 200, 500,$  and  $1000$ . A significant deviation from the logarithmic scaling behaviour is observed when  $L$  is relatively small, but tends to vanish, as  $L$  increases. The prefactor is close to the exact value  $1/2$ , with an error being less than 2%, when  $L = 1000$ :  $S_{1000}(n, 250) = 0.499 \log_2 n + 0.818$  and  $S_{1000}(n, 500) = 0.490 \log_2 n + 1.080$  for  $M = L/2$  and  $M = L/4$ , respectively. Here,  $n$  ranges from 10 to 40.

The degenerate ground states  $|L, M\rangle$  admit an exact singular value decomposition:

$$|L, M\rangle = \sum_{k=0}^{\min(M, n)} \lambda(L, M, k) |n, k\rangle |L - n, M - k\rangle, \quad (17)$$

where the singular values  $\lambda(L, M, k)$  take the form,

$$\lambda(L, M, k) = \sqrt{\frac{C_n^k C_{L-n}^{M-k}}{C_L^M}}. \quad (18)$$

Therefore, the eigenvalues  $\Lambda(L, M, k)$  of the reduced density matrix  $\rho_L(n, M)$  are  $\Lambda(L, M, k) = [\lambda(L, M, k)]^2$ . Hence, the entanglement entropy  $S_L(n, M)$  follows from Eq. (5). An analytical treatment confirms the logarithmic scaling behaviour, as predicted in Eq. (6).

In order to understand how the logarithmic scaling behaviour emerges in the thermodynamic limit, we plot  $S_L(n, M)$  vs  $\log_2 n$  in Fig. 3 for the coexisting fractal states:

(a)  $M = L/2$  and (b)  $M = L/4$ , when  $L$  is varied:  $L = 100, 200, 500,$  and  $1000$ . A significant deviation from the logarithmic scaling behaviour is observed when  $L$  is relatively small, but tends to vanish, as  $L$  increases. The prefactor is close to the exact value  $1/2$ , with an error being less than 2%, when  $L = 1000$ :  $S_{1000}(n, 250) = 0.499 \log_2 n + 0.818$  and  $S_{1000}(n, 500) = 0.490 \log_2 n + 1.080$  for  $M = L/2$  and  $M = L/4$ , respectively. Here,  $n$  ranges from 10 to 40.

## VI. SUMMARY

In this work, a systematic investigation has been performed for highly degenerate ground states arising from SSB with a type-B GM in the context of quantum entanglement. It is found that the degenerate ground states admit an exact singular value decomposition, which unveils their scale invariance. This implies that the entanglement entropy  $S(n)$  scales logarithmically with the block size  $n$  in the thermodynamic limit, with the prefactor being half the number of type-B GMs  $N_B$ . Meanwhile, as follows from a field-theoretic prediction [16], the prefactor is half the fractal dimension  $d_f$ . Therefore, the fractal dimension  $d_f$  is identical to the number of type-B GMs  $N_B$  for the degenerate ground states. Our claim has been tested for the  $SU(2)$  spin- $s$  ferromagnetic model, the  $SU(N+1)$  ferromagnetic model and the  $SU(2)$  spin-1 anisotropic bi-quadratic model, with the fractal dimension  $d_f$  being 1,  $N$ , and 1, respectively. In addition, an extensive numerical analysis has been performed to reveal how the logarithmic scaling behaviour emerges, as the system size  $L$  increases. This lends further support to our claim.

In closing, a few remarks are in order. First, the extension to a symmetry group  $G$  other than  $SU(N+1)$  is possible, although our discussion focuses on  $SU(N+1)$ . Second, it is straightforward to extend to quantum many-body systems in two and higher spatial dimensions, given that the occurrence of SSB with a type-B GM does not depend on the spatial dimensionality. Last but not least, the scaling behaviour of the entanglement entropy remains unclear in the thermodynamic limit when both type-A and type-B GMs are present in a quantum many-body system.

## VII. ACKNOWLEDGEMENTS

We thank Murray Batchelor, Sam Young Cho, and John Fjaerestad for enlightening discussions. The work is supported by the National Natural Science Foundation of China (Grant No. 11805285).

- 
- [1] J. Goldstone, *Nuovo Cimento* **19**, 154 (1961); J. Goldstone, A. Salam, and S. Weinberg, *Phys. Rev.* **127**, 965 (1962); Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122**, 345 (1961).
- [2] H. B. Nielsen and S. Chadha, *Nucl. Phys. B* **105**, 445 (1976).
- [3] T. Schafer, D. T. Son, M. A. Stephanov, D. Toublan, and J. J. M. Verbaarschot, *Phys. Lett. B* **522**, 67 (2001).
- [4] V. A. Miransky and I. A. Shovkovy, *Phys. Rev. Lett.* **88**, 111601 (2002).
- [5] A. Nicolis and F. Piazza, *Phys. Rev. Lett.* **110**, 011602 (2013); H. Watanabe, T. Brauner, and H. Murayama, *Phys. Rev. Lett.* **111**, 021601 (2013).
- [6] H. Watanabe and T. Brauner, *Phys. Rev. D* **84**, 125013 (2011).
- [7] H. Watanabe and H. Murayama, *Phys. Rev. Lett.* **108**, 251602 (2012).
- [8] H. Watanabe and H. Murayama, *Phys. Rev. X* **4**, 031057 (2014).
- [9] Y. Hidaka, *Phys. Rev. Lett.* **110**, 091601 (2013); T. Hayata and Y. Hidaka, *Phys. Rev. D* **91**, 056006 (2015); D. A. Takahashi and M. Nitta, *Ann. Phys.* **354**, 101 (2015).
- [10] Y. Nambu, *J. Stat. Phys.* **115**, 7 (2004).
- [11] P. W. Anderson, *Phys. Rev.* **86**, 694 (1952).
- [12] M. A. Metlitski and T. Grover, arXiv: 1112.5166v2 (2011).
- [13] L. Rademaker, *Phys. Rev. B* **92**, 144419 (2015).
- [14] V. Popkov and M. Salerno, *Phys. Rev. A* **71**, 012301 (2005).
- [15] V. Popkov, M. Salerno, and G. Schütz, *Phys. Rev. A* **72**, 032327 (2005).
- [16] O. A. Castro-Alvaredo and B. Doyon, *Phys. Rev. Lett.* **108**, 120401 (2012).
- [17] N. D. Mermin and H. Wagner, *Phys. Rev. Lett.* **17**, 1133 (1966); S. R. Coleman, *Commun. Math. Phys.* **31**, 259 (1973).
- [18] P. D. Francesco, P. Mathieu, and D. Sénéchal, *Conformal Field Theory* (Springer, Berlin, 1997).
- [19] A. M. Polyakov, *JETP Lett.* **12**, 381 (1970).
- [20] H. Osborn, *Nucl. Phys. B* **363**, 486 (1991); Z. Komargodski and A. Schwimmer, *J. High Energy Phys.* **12**, 099 (2011); J.-F. Fortin, B. Grinstein, and A. Stergiou, *J. High Energy Phys.* **01**, 184 (2013); Y. Nakayama, *Phys. Rep.* **569**, 1 (2015).
- [21] A. LeClair, J.M. Roman, and G. Sierra, *Nucl. Phys. B* **700**, 407 (2004); A. LeClair and G. Sierra, *J. Stat. Mech.* **0408**, P08004 (2004).
- [22] A. Dymarsky, Z. Komargodski, A. Schwimmer, and S. Theisen, *J. High Energy Phys.* **10**, 171 (2015); B. Delamotte, M. Tissier, and N. Wschebor, *Phys. Rev. E* **93**, 012144 (2015); A. Dymarsky, K. Farnsworth, Z. Komargodski, M. A. Luty, and V. Prilepina, *J. High Energy Phys.* **02**, 099 (2016); A. Dymarsky and A. Zhiboedov, *J. Phys. A: Math. Theor.* **48**, 41FT01 (2015); M. Hogervorst, S. Rychkov, and B. C. van Rees, *Phys. Rev. D* **93**, 125025 (2016).
- [23] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [24] G. P. Hochschild, *Basic Theory of Algebraic Groups and Lie Algebras* (Springer, New York, 1981).
- [25] A. J. Beekman, L. Rademaker, and J. van Wezel, *SciPost Phys. Lect. Notes* **11** (2019).
- [26] W. Feller, *An Introduction to Probability Theory and Its Applications* (Wiley, New York, 1968).
- [27] Q.-Q. Shi, Y.-W. Dai, S.-H. Li, and H.-Q. Zhou, arXiv: 2204.05692 (2022).
- [28] B. Sutherland, *Phys. Rev. B* **12**, 3795 (1975).



# Supplementary Material for “Fractal dimension and the counting rule of the Goldstone modes”

Qian-Qian Shi,<sup>1</sup> Yan-Wei Dai,<sup>1</sup> Huan-Qiang Zhou,<sup>1</sup> and Ian P. McCulloch<sup>2,3,1</sup>

<sup>1</sup>Centre for Modern Physics, Chongqing University, Chongqing 400044, The People’s Republic of China

<sup>2</sup>Department of Physics, National Tsing Hua University, Hsinchu 30013, Taiwan

<sup>3</sup>School of Mathematics and Physics, The University of Queensland, St. Lucia, QLD 4072, Australia

## A. A logarithmic scaling relation between the entanglement entropy $S(n)$ and the block size $n$

We present a heuristic argument, aiming to unveil a logarithmic scaling relation between the entanglement entropy  $S(n)$  and the block size  $n$ , for a scale-invariant state, in the thermodynamic limit, when the fillings  $f_1, \dots, f_N$  are kept to be constant. We focus on scale-invariant states in one spatial dimension, with the block  $B$  consisting of  $n$  contiguous lattice sites for simplicity. The partition of the system into the block  $B$  and the environment  $E$  amounts to introducing a length scale  $n$ , thus the system is expected to react. Physically, it is legitimate to consider an effective (continuum) field theory, for a quantum many-body system on a lattice, when the thermodynamic limit is approached.

Suppose the entanglement entropy  $S(n)$  is  $S(n) = f(n)$ , with  $f(n)$  being a function of  $n$  to be determined. The transformation  $n \rightarrow \lambda n$  amounts to introducing two sequences of the values of the block size  $n$ , with a dimensionless ratio  $\lambda$ . The scale invariance implies that the scaling behaviour with  $n$  must remain the same, with an additional contribution from  $\lambda$  being additive. Mathematically, we have

$$f(\lambda n) = f(n) + F(\lambda), \quad (\text{S.1})$$

where  $F(\lambda)$  is a function of  $\lambda$  to be determined. It is easy to see that

$$F(1) = 0. \quad (\text{S.2})$$

Taking the first-order derivative with respect to the parameter  $\lambda$  on both hand sides of Eq.(S.1), we have

$$nf'(\lambda n) = F'(\lambda). \quad (\text{S.3})$$

Setting  $y = \lambda n$ , we are led to

$$yf'(y) = \lambda F'(\lambda) \quad (\text{S.4})$$

The variables are separated, since the left hand side only depends on  $y$ , and the right hand side only depends on  $\lambda$ . This implies that

$$yf'(y) = \kappa, \quad (\text{S.5})$$

and

$$\lambda F'(\lambda) = \kappa, \quad (\text{S.6})$$

with  $\kappa$  being a constant to be determined. This yields

$$f(y) = \kappa \ln y + f(1), \quad (\text{S.7})$$

and

$$F(\lambda) = \kappa \ln \lambda. \quad (\text{S.8})$$

Physically,  $\kappa$  should be proportional to the number of type-B GMs:  $\kappa = \eta N_B$ , with  $\eta$  being a universal constant, since only the low-lying excitations contribute to the scaling behaviour of the entanglement entropy  $S(n)$  with the block size  $n$ . The value of  $\eta$  may be determined from a specific model, as long as its exact scaling relation between the entanglement entropy  $S(n)$  and the block size  $n$ , together with the number of type-B GMs  $N_B$ , are known. For the SU(2) spin-1/2 ferromagnetic states, an analytic treatment yields that  $S(n) = 1/2 \ln n + S_0$  [S1], with the number of type-B GMs  $N_B = 1$  [S2]. As a consequence, we have  $\eta = 1/2$ . That is, for a scale-invariant state,  $S(n)$  scales logarithmically with  $n$ ,

$$S(n) = \frac{N_B}{2} \ln n + S_0, \quad (\text{S.9})$$

where  $S_0$  is an additive contribution to the entanglement entropy, which is non-universal. In addition, a field-theoretic approach [S3] predicts that

$$S(n) = \frac{d_f}{2} \ln n + S_0, \quad (\text{S.10})$$

with  $d_f$  being the fractal dimension. Therefore, we are led to the identification of the fractal dimension  $d_f$  with the number of type-B GMs:  $d_f = N_B$ , for the degenerate ground states arising from SSB with a type-B GM.

Here, we remark that the fillings  $f_1, \dots, f_N$  are assumed to be nonzero when the scaling relation (S.9) is derived. If the fillings  $f_1, \dots, f_N$  are zero, then the ground state is the highest weight state, which is unentangled. Thus, the entanglement entropy  $S(n)$  is simply zero. This implies that the entanglement entropy  $S(n)$  features a singularity, with a discontinuous jump in the prefactor in front of the logarithmic function, when the zero filling limit is approached. Physically, this is due to the fact that the highest weight state, as an unentangled ground state, is special, in the sense that type-B GMs manifest themselves only when the other degenerate ground states with nonzero fillings are involved. Note that the same argument is valid for the lowest weight state, which is an unentangled ground state, when the fillings  $f_1, \dots, f_N$  reach the maximum values.

The argument may be extended to a scale-invariant state in  $D$  spatial dimensions. Then, (S.9) and (S.10) become

$$S(n) = \frac{DN_B}{2} \ln n + S_0, \quad (\text{S.11})$$

and

$$S(n) = \frac{Dd_f}{2} \ln n + S_0, \quad (\text{S.12})$$

respectively. Here, we stress that  $n$  should be understood as the linear size of the  $D$ -dimensional block.

In passing, we remark that our argument for the logarithmic scaling behavior, up to Eq. (S.7) and Eq. (S.8), also works for conformally invariant states in the thermodynamic limit, since scale invariance is part of conformal invariance. In fact, the prefactor  $\kappa$  must be proportional to central charge:  $\kappa = \zeta c$ , with  $\zeta$  being a constant, given that  $c$  counts the number of gapless excitations [S4]. One may also determine the proportionality constant  $\zeta$  to be  $1/3$  from the XY model in a longitudinal magnetic field – an exactly solvable model [S5], with central charge  $c = 1$  at criticality [S6]. This allows us to reproduce the scaling relation of the entanglement entropy  $S(n)$  with the block size  $n$  for conformally invariant states [S7, S8].

## B. A derivation of $Z(L, M)$ for the SU(2) spin- $s$ ferromagnetic states

For the SU(2) spin- $s$  ferromagnetic states  $|L, M\rangle$  in Eq. (9), we need to figure out a way to derive the normalization factor  $Z(L, M)$ . Here, we resort to the permutation invariance of  $|L, M\rangle$  to facilitate the derivation.

For our purpose, it is convenient to introduce a set of permutation invariant states  $|\psi(N_{-s}, \dots, N_s)\rangle$ , defined as follows

$$|\psi(N_{-s}, \dots, N_s)\rangle = \frac{1}{Z_\psi} \sum_P |\underbrace{-s \dots -s}_{N_{-s}} | \dots | \underbrace{s \dots s}_{N_s}\rangle, \quad (\text{S.13})$$

with

$$Z_\psi = \sqrt{\prod_{r=-s}^{s-1} C_{L-\sum_{m=-s}^{r-1} N_m}^{N_r}}. \quad (\text{S.14})$$

Here,  $Z_\psi$  is introduced to ensure that  $|\psi(N_{-s}, \dots, N_s)\rangle$  is normalized, and the sum  $\sum_P$  is taken over all the possible permutations  $P$  for a given partition  $\{N_{-s}, \dots, N_s\}$ .

Now we are ready to expand  $|L, M\rangle$  in terms of the basis states  $|\psi(N_{-s}, \dots, N_s)\rangle$ :

$$|L, M\rangle = \frac{1}{Z(L, M)} \sum'_{N_{-s}, \dots, N_s} c(L, M, N_{-s}, \dots, N_s) |\psi(N_{-s}, \dots, N_s)\rangle, \quad (\text{S.15})$$

where

$$c(L, M, N_{-s}, \dots, N_s) = Z_\psi \frac{M!}{\sqrt{2^M}} \prod_{r=-s}^{s-1} [\sqrt{\varepsilon(s, r)}]^{N_r}, \quad (\text{S.16})$$

with

$$\varepsilon(s, r) = \frac{\prod_{m=r+1}^s (s+m)(s-m+1)}{\prod_{m=r}^{s-1} (s-m)^2}.$$

Here, the sum  $\sum'_{N_{-s}, \dots, N_s}$  is taken over all the possible values of  $N_{-s}, \dots, N_s$ , subject to the constraints:  $\sum_{m=-s}^s N_m = L$  and  $\sum_{m=-s}^s (s-m)N_m = M$ . Indeed,  $Z(L, M)$  takes the form

$$Z(L, M) = \sqrt{\sum'_{N_{-s}, \dots, N_s} c(L, M, N_{-s}, \dots, N_s)^2}. \quad (\text{S.17})$$

Substituting Eq.(S.16) into Eq.(S.17), we have

$$Z(L, M) = \frac{M!}{\sqrt{2^M}} \sqrt{\sum'_{N_{-s}, \dots, N_s} \prod_{r=-s}^{s-1} [\varepsilon(s, r)]^{N_r} C_{L-\sum_{m=-s}^{r-1} N_m}^{N_r}}. \quad (\text{S.18})$$

## C. A derivation of $Z(L, M_1, \dots, M_N)$ for the SU(N + 1) ferromagnetic states

In order to derive the normalization factor  $Z(L, M_1, \dots, M_N)$  for the SU(N + 1) ferromagnetic states  $|L, M_1, \dots, M_N\rangle$  in Eq.(13), we need to take advantage of the permutation invariance of  $|L, M_1, \dots, M_N\rangle$ .

In fact, one may rewrite  $|L, M_1, \dots, M_N\rangle$  as follows

$$|L, M_1, \dots, M_N\rangle = \frac{\prod_{\alpha=1}^N M_\alpha!}{Z(L, M_1, \dots, M_N)} \sum_P |\underbrace{1 \dots 1}_{L-\sum_{\alpha=1}^N M_\alpha} | \underbrace{2 \dots 2}_{M_1} \dots | \underbrace{N+1 \dots N+1}_{M_N}\rangle, \quad (\text{S.19})$$

where  $|\beta\rangle$  ( $\beta = 1, 2, \dots, N + 1$ ) are defined as a  $N + 1$ -dimensional vector, with the  $\beta$ -th entry being 1, and the others being 0, and the sum  $\sum_P$  is taken over all the permutations  $P$  for a given partition  $\{L - \sum_{\alpha=1}^N M_\alpha, M_1, \dots, M_N\}$ . As a consequence,  $Z(L, M_1, \dots, M_N)$  takes the form

$$Z(L, M_1, \dots, M_N) = \prod_{\alpha=1}^N M_\alpha! \sqrt{C_{L - \sum_{\beta=1}^{\alpha-1} M_\beta}^{M_\alpha}}. \quad (\text{S.20})$$

#### D. A derivation of $Z(L, M)$ for the coexisting fractal states

For the coexisting fractal states  $|L, M\rangle$  in Eq. (18), we need to introduce a unit cell consisting of two nearest-neighbor sites, due to the staggered nature of the symmetry group  $SU(2)$ . Therefore, there are four possible configurations:  $|1_z 1_z\rangle$ ,  $|1_z - 1_z\rangle$ ,  $|-1_z 1_z\rangle$ , and  $|-1_z - 1_z\rangle$  in a unit cell. Here,  $|\pm 1_z\rangle$  are the eigenvectors of  $S_{z,j}$ , with the eigenvalues being  $\pm 1$ .

One may rewrite  $|L, M\rangle$  as follows

$$|L, M\rangle = \frac{M!}{\sqrt{2^M} Z(L, M)} \sum'_{N_{mm}, N_{mp}, N_{pm}, N_{pp}} \sum_P (-1)^{N_{mm} + N_{mp}} \underbrace{|-1_z - 1_z \dots - 1_z - 1_z\rangle}_{N_{mm}} \underbrace{|-1_z 1_z \dots - 1_z 1_z\rangle}_{N_{mp}} \underbrace{|1_z - 1_z \dots 1_z - 1_z\rangle}_{N_{pm}} \underbrace{|1_z 1_z \dots 1_z 1_z\rangle}_{N_{pp}}, \quad (\text{S.21})$$

where the sum  $\sum_P$  is taken over all the permutations  $P$  for a given partition  $\{N_{mm}, N_{mp}, N_{pm}, N_{pp}\}$ , and the sum  $\sum'_{N_{mm}, N_{mp}, N_{pm}, N_{pp}}$  is taken over all the possible values of  $N_{mm}, N_{mp}, N_{pm}, N_{pp}$ , subject to the constraints:  $2N_{mm} + N_{mp} + N_{pm} = M$  and  $N_{mm} + N_{mp} + N_{pm} + N_{pp} = L/2$ . Here,  $N_{mm}, N_{mp}, N_{pm}$ , and  $N_{pp}$  denote the numbers of the unit cells in the configurations  $|-1_z - 1_z\rangle$ ,  $|-1_z 1_z\rangle$ ,  $|1_z - 1_z\rangle$ , and  $|1_z 1_z\rangle$ , respectively.

Then,  $Z(L, M)$  takes the form

$$Z(L, M) = \frac{M!}{\sqrt{2^M}} \sqrt{\sum'_{N_{mm}, N_{mp}, N_{pm}, N_{pp}} C_{L/2}^{N_{mm}} C_{L/2 - N_{mm}}^{N_{pp}} C_{L/2 - N_{mm} - N_{pp}}^{N_{pm}}}, \quad (\text{S.22})$$

which may be simplified as

$$Z(L, M) = M! \sqrt{\frac{C_L^M}{2^M}}. \quad (\text{S.23})$$

We thank Murray Batchelor, Sam Young Cho, and John Fjaerestad for enlightening discussions. The work is sup-

ported by the National Natural Science Foundation of China (Grant No. 11805285).

- [S1] V. Popkov and M. Salerno, Phys. Rev. A **71**, 012301 (2005).  
[S2] H. Watanabe and H. Murayama, Phys. Rev. Lett. **108**, 251602 (2012).  
[S3] O. A. Castro-Alvaredo and B. Doyon, Phys. Rev. Lett. **108**, 120401 (2012).  
[S4] P. D. Francesco, P. Mathieu, and D. Sénéchal, Conformal Field

- Theory (Springer, Berlin, 1997).  
[S5] B.-Q. Jin and V. E. Korepin, J. Stat. Phys. **116**, 79 (2004).  
[S6] S. Sachdev, Quantum Phase Transitions (Cambridge University Press, Cambridge, 1999).  
[S7] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, Phys. Rev. Lett. **90**, 227902 (2003).  
[S8] P. Calabrese and J. Cardy, J. Stat. Mech. **2004**, P06002 (2004).