

# Gauge Invariant Perturbations of General Spherically Symmetric Spacetimes

Wentao Liu,<sup>1</sup> Xiongjun Fang,<sup>1,2,\*</sup> Jiliang Jing,<sup>1,†</sup> and Anzhong Wang<sup>2,‡</sup>

<sup>1</sup>*Department of Physics, Key Laboratory of Low Dimensional Quantum Structures and Quantum Control of Ministry of Education, and Synergetic Innovation Center for Quantum Effects and Applications, Hunan Normal University, Changsha, Hunan 410081, P. R. China*

<sup>2</sup>*GCAP-CASPER, Department of Physics, Baylor University, One Bear Place #97316, Waco, Texas 76798-7316, USA*

In this paper, the gauge choices in general spherically symmetric spacetimes have been explored. We construct the gauge invariant variables and the master equations for both the Detweiler easy gauge and the Regge-Wheeler gauge, respectively. The particular cases for  $l = 0, 1$  are also investigated. Our results provide analytical calculations of metric perturbations in general spherically symmetric spacetimes, which can be applied to various cases, including the Effective-One-Body problem. A simple example is presented to show how the metric perturbation components are related to the source perturbation terms.

## I. INTRODUCTION

Metric perturbations of spacetimes is an important issue. The solution of the Einstein field equations (EFEs) for static, vacuum and spherically symmetric spacetime is the Schwarzschild spacetime. And the metric perturbation of the Schwarzschild black hole has been studied for a long time. To begin with, Regge, Wheeler and Vishveshwara et.al studied the odd-parity perturbation [1, 2], while Zerilli and Moncrief investigated the even-parity perturbation [3–5]. The perturbation theory of the Schwarzschild spacetime has been well summarized in Chandrasekhar’s monograph [7]. After decades of research and development, this theory can be applied to a variety of different physical problems. A useful application is the quasi-normal modes of the perturbed black holes, which was initiated by Vishveshwara [8], Chandrasekhar [6] and Mashhoon [9] et.al., and the review articles of this topic can be found in [10–15]. Another application is studying a particle moving around the Schwarzschild black hole. One can treat this point-particle as a perturbation of the Schwarzschild spacetime [16, 17]. In addition, studying the metric perturbation can promote the analysis of the stability of the Schwarzschild spacetime [18–20].

In perturbation theory in general relativity, the redundant coordinate freedom can be eliminated by choosing specific gauges. The most familiar gauge in the Schwarzschild spacetime is the “Regge-Wheeler (RW) gauge”, which was first presented by Regge and Wheeler [1]. And where they also analysed the spherical harmonics and decomposed the general perturbation in the Schwarzschild spacetime into odd-parity sector and even-parity sectors. The RW gauge has the obvious advantage of algebraic simplicity, and it is widely used in the literature. After that, the construction and the physical meaning of the gauge-invariant properties attract people’s attention. Using Lagrangian and Hamiltonian variational principles for the perturbation, Moncrief considered that the metric perturbations can be decomposed into the gauge invariant part and the gauge dependent part [5, 21]. Gerlach and Sengupta discussed the construction of gauge invariant properties in general spherically symmetric spacetimes [22, 23]. Thorne reviewed and summarized the various scalar, vector and tensor spherical harmonics with a uniform notation [24]. Martel and Poisson presented a gauge-invariant and covariant formalism, and they also showed that the energy or angular-momentum radiation can be expressed in terms of gauge-invariant scalar functions [25]. Recently, Sopuerta considered that the master functions are linear combinations of the metric perturbations and their first-order derivatives, and discussed about the master equation for vacuum spherically symmetric spacetimes [26]. Besides the RW gauge, there exist a variety of gauge choices. For example, the light-cone gauge, which presented by Preston and Poisson [27], can provide geometrical meaning to the coordinate in perturbed spacetimes. Another gauge choice is named as “easy (EZ) gauge” [28], which was devised by Detweiler when he considered the gravitational self-force problem in the perturbed Schwarzschild spacetime [29]. In the EZ gauge, the metric perturbation is singular on the black-hole horizon [30].

Generally speaking, for metric perturbation of a spacetime, the standard process is to decouple the even-parity or the odd-parity EFEs, obtain the wave-like master equations and then solve the one dimensional Schrödinger equation

---

\*Corresponding author: [fangxj@hunnu.edu.cn](mailto:fangxj@hunnu.edu.cn)

†[jljing@hunn.edu.cn](mailto:jljing@hunn.edu.cn)

‡[anzhongwang@baylor.edu](mailto:anzhongwang@baylor.edu)

with an effective potential. One of the most important step is how to construct the gauge-invariant variable and get the master equation. The equations that arise in the above process have been thoroughly studied in the Schwarzschild spacetime. However, for example, if one considers a non-vacuum static black hole with hair [31–33], the metric would be no longer the Schwarzschild spacetime. Especially, when studying the Effective-One-Body problem to general relativistic two-body dynamics, peoples used the static and spherically symmetric effective metric [34, 35]. Therefore, a natural question is how to decouple the EFEs and construct the gauge-invariant variables in general spherically symmetric spacetimes, both for even-parity and odd-parity perturbations.

In this paper, we consider the general spherically symmetric background spacetimes and the most general metric perturbations, and the construction of gauge invariant variables. For even parity perturbations, we find that there exist several gauge choices, including the EZ gauge and the RW gauge. Under the EZ gauge, we construct the gauge-invariant variables and obtain a third order master equations. However, the third order equation can be written as a second order equation after the separation of radial and time variables. While under the RW gauge, we also construct the master equation, and find that the master equation is a second order wave-like equation. For the odd-parity perturbations, the master equation remains as the second order wave-like equation.

Considering the general metric perturbations in static spherically symmetric spacetimes, Throne showed how to construct a ten-spherical-harmonic basis [24]. Through this paper, we use the A-K notation [28], which was first presented by Detweiler when he considered the self-force problem. The advantage of using this notation is that one can obviously find the relation between the metric perturbation components and the gauge invariants. In this paper, through the gauge invariants representing different combinations of the metric components, we show that the gauge-invariant variables have the similar structure under the EZ gauge or the RW gauge.

The rest of this paper is organized as follows. In Sect. 2, we discuss the basic framework. First, the ten orthogonal harmonics basis are introduced. Then the decomposition of non-vacuum Einstein equations and the A-K notation are reviewed. After the investigation of gauge freedom, we consider the EZ gauge and the RW gauge. In Sect. 3, we first consider the gauge invariant properties. Then we focus on constructing the master equations, for both even-parity and odd-parity perturbations. We also study the cases for  $l = 0, 1$ . In Sect. 4, an example that a small particle goes around a circular orbit in spherically symmetric spacetimes is investigated. Finally, we summarize our main results with some discussions.

Throughout this paper, we use the A-K notation similarly to [28]. Units will be chosen in which  $c = G = 1$ . In Sect. 3, the subscript, e.g.,  $\psi_0$  and  $\psi_1$ , always represents quantities for  $l = 0$  case and  $l = 1$  case, respectively. And the superscript with Roman letters, i.e.  $\chi^I$  and  $\chi^{II}$ , represent the quantities under the EZ gauge or under the RW gauge, respectively.

## II. BASIC FRAMEWORK

### A. Orthogonal Harmonics Basis

Let us start with the general spherically symmetric spacetimes

$$ds^2 = g_{ab}^{(0)} dx^a dx^b = -e^{2\Phi} dt^2 + e^{2\Lambda} dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2). \quad (1)$$

To decompose tensor fields on the above background, we choose the orthogonal basis composed of scalar spherical harmonics, vector harmonics and tensor harmonics. First, we define two unnormalized and orthogonal co-vectors  $v$  and  $n$

$$v_a = (-1, 0, 0, 0), \quad n_a = (0, 1, 0, 0), \quad (2)$$

and the projection operator onto the sphere surface

$$\Omega_{ab} = g_{ab}^{(0)} + e^{2\Phi} v_a v_b - e^{2\Lambda} n_a n_b = r^2 \text{diag}(0, 0, 1, \sin^2\theta), \quad (3)$$

and the spatial Levi-Civita tensor,  $\epsilon_{abc} \equiv v^d \epsilon_{dabc}$ , where  $\epsilon_{tr\theta\phi} = e^{\Phi+\Lambda} r^2 \sin\theta$ .

In general, the complete basis on the 2-sphere are constructed by 1-scalar spherical harmonic,  $Y^{lm} = Y^{lm}(\theta, \phi)$ , and 3 pure-spin vector harmonics and 6 tensor harmonics. The pure-spin vector harmonics are given by

$$Y_a^{E,lm} = r \nabla_a Y^{lm}, \quad Y_a^{B,lm} = r \epsilon_{ab}^c n^b \nabla_c Y^{lm}, \quad Y_a^{R,lm} = n_a Y^{lm}. \quad (4)$$

And the pure-spin tensor harmonics are given by

$$T_{ab}^{T0,lm} = \Omega_{ab} Y^{lm}, \quad T_{ab}^{L0,lm} = n_a n_b Y^{lm}, \quad (5)$$

$$T_{ab}^{E1,lm} = r n_{(a} \nabla_{b)} Y^{lm}, \quad T_{ab}^{B1,lm} = r n_{(a} \epsilon_{b)c}{}^d n^c \nabla_d Y^{lm}, \quad (6)$$

$$T_{ab}^{E2,lm} = r^2 \left( \Omega_a{}^c \Omega_b{}^d - \frac{1}{2} \Omega_{ab} \Omega^{cd} \right) \nabla_c \nabla_d Y^{lm}, \quad T_{ab}^{B2,lm} = r^2 \Omega_{(a}{}^c \epsilon_{b)e}{}^d n^e \nabla_c \nabla_d Y^{lm}. \quad (7)$$

Note that the vector harmonics are orthogonal to each other

$$\oint Y_a^{A,lm} (Y_{A',l'm'}^a)^* d\Omega = N^{(vec)}(A, r, l) \delta_{AA'} \delta_{ll'} \delta_{mm'}, \quad (8)$$

with  $\{A, A'\} \in \{E, B, R\}$  and  $N^{(vec)}(A, r, l)$  is the specific normalization factor for vector harmonics. The tensor harmonics are also orthogonal to each other

$$\oint T_{ab}^{A,lm} (T_{A',l'm'}^{ab})^* d\Omega = N^{(ten)}(A, r, l) \delta_{AA'} \delta_{ll'} \delta_{mm'}, \quad (9)$$

with  $\{A, A'\} \in \{T0, L0, E1, E2, B1, B2\}$  and  $N^{(ten)}(A, r, l)$  is the specific normalization factor for tensor harmonics. The expression for these normalization functions  $N^{(vec)}$  and  $N^{(ten)}$  are given in Appendix A.

## B. Decomposition of Linearized Einstein Equations

For perturbed spacetimes, we use  $h_{ab}$  to represent the linear perturbation of the background spacetime  $g_{ab}^{(0)}$ , i.e., the metric of perturbed spacetime can be written as

$$g_{ab} = g_{ab}^{(0)} + h_{ab}. \quad (10)$$

The background metric and the perturbed metric satisfied the Einstein Field Equations (EFEs)

$$G_{ab}(g^{(0)}) = 8\pi T_{ab}, \quad (11)$$

$$G_{ab}(g^{(0)} + h) = 8\pi(T_{ab} + \mathcal{T}_{ab}), \quad (12)$$

where  $T_{ab}$  and  $\mathcal{T}_{ab}$  denote the non-vacuum background and the perturbed energy-momentum tensor, respectively. Expanding the EFEs in terms of  $h_{ab}$ , we get

$$G_{ab}(g^{(0)} + h) = G_{ab}(g^{(0)}) - \frac{1}{2} E_{ab}(h), \quad (13)$$

where  $E_{ab}$  is the linearized Einstein operator

$$-16\pi \mathcal{T}_{ab} = E_{ab}(h) = \square h_{ab} + \nabla_a \nabla_b h^c{}_c - 2\nabla_{(a} \nabla^c h_{b)c} + 2R_a{}^c{}^d h_{cd} + g_{ab}(\nabla^c \nabla^d h_{cd} - \square h^d{}_d) - g_{ab} R^{cd} h_{cd} + R h_{ab}. \quad (14)$$

Note that now the Ricci curvature  $R$  and the scalar curvature  $R_{ab}$  of the background do not vanish in general. If  $R$  and  $R_{ab}$  vanish, then the background metric would reduce to the Schwarzschild metric, which is the situation discussed in [28].

Detweiler decomposed the harmonic modes of the perturbed metric  $h_{ab}$  as

$$h_{ab}^{lm} = A v_a v_b Y^{lm} + 2B v_{(a} Y_{b)}^{E,lm} + 2C v_{(a} Y_{b)}^{B,lm} + 2D v_{(a} Y_{b)}^{R,lm} + E T_{ab}^{T0,lm} + F T_{ab}^{E2,lm} + G T_{ab}^{B2,lm} + 2H T_{ab}^{E1,lm} + 2J T_{ab}^{B1,lm} + K T_{ab}^{L0,lm}, \quad (15)$$

where all coefficients A through K are scalar functions of  $(t, r)$ , which were referred to as the A-K coefficients in [28].

For even-parity (polar part) perturbations with  $l \geq 2$ , the perturbed metric can be decomposed as

$$h_{ab}^{\text{even}} = \begin{pmatrix} A Y_{lm} & -D Y_{lm} & -rB \partial_\theta Y_{lm} & -rB \partial_\phi Y_{lm} \\ \text{Sym} & K Y_{lm} & rH \partial_\theta Y_{lm} & rH \partial_\phi Y_{lm} \\ \text{Sym} & \text{Sym} & r^2 [E + F (\partial_\theta^2 + \frac{1}{2}l(l+1))] Y_{lm} & r^2 F [\partial_\theta \partial_\phi - \cot \theta \partial_\phi] Y_{lm} \\ \text{Sym} & \text{Sym} & \text{Sym} & r^2 \sin^2 \theta [E - F (\partial_\theta^2 + \frac{1}{2}l(l+1))] Y_{lm} \end{pmatrix}. \quad (16)$$

For the odd-parity (axial part) perturbations with  $l \geq 2$ , the perturbed metric can be decomposed as

$$h_{ab}^{\text{odd}} = e^{-\Phi-\Lambda} \begin{pmatrix} 0 & 0 & r \csc \theta C \partial_\phi Y_{lm} & -r \sin \theta C \partial_\theta Y_{lm} \\ 0 & 0 & -r \csc \theta J \partial_\phi Y_{lm} & r \sin \theta J \partial_\theta Y_{lm} \\ \text{Sym} & \text{Sym} & -r^2 \csc \theta G [\partial_\theta \partial_\phi - \cot \theta \partial_\phi] Y_{lm} & -\frac{r^2}{2} G [\csc \theta \partial_\phi^2 + \cos \theta \partial_\theta - \sin \theta \partial_\theta^2] Y_{lm} \\ \text{Sym} & \text{Sym} & \text{Sym} & r^2 G [\sin \theta \partial_\theta \partial_\phi - \cos \theta \partial_\phi] Y_{lm} \end{pmatrix}. \quad (17)$$

These A-K notation could be linearly transformed into the notation taken by Detweiler [28]. To get the explicit expression of coefficients A-K, one can project the metric perturbation onto tensor harmonic basis. We present these coefficients in Appendix A.

Now, we can get the A-K components of any rank-2 tensor in the spherically symmetric background. For example, from Eq.(14), one can write

$$-16\pi \mathcal{T}_A = E_A = e^{4\Phi} \oint E_{ab}(h) (v^a v^b Y_{lm}^*) d\Omega, \quad (18)$$

to represent the A-term of  $\mathcal{T}_{ab}$  or  $E_{ab}(h)$ , which along the direction of  $v^a v^b Y_{lm}^*$ . The expression for  $E_A$ - $E_K$  are given in Appendix B.

### C. The Gauge Transformation

Under a gauge transformation,  $\tilde{x}^a = x^a + \xi^a$ , the first-order metric perturbation  $h_{ab}$  would be transformed as

$$\tilde{h}_{ab} = h_{ab} - 2\nabla_{(a} \xi_{b)}. \quad (19)$$

As a vector,  $\xi^a$  can be projected onto the pure-spin harmonic basis as

$$\xi_a = P v_a Y_{lm} + R Y_a^{R,lm} + S Y_a^{E,lm} + Q Y_a^{B,lm}, \quad (20)$$

where P, R, S and Q are scalar functions of  $(t, r)$ . The function P, R and S describe three degrees of gauge freedom for even-parity perturbations, while the function Q describes one degrees of gauge freedom for odd-parity perturbations. Next we use  $\Delta$  to represent the A-K projections of  $2\nabla_{(a} \xi_{b)}$ . For example,

$$\Delta A = A - \tilde{A} = 2e^{4\Phi} \oint \nabla_{(a} \xi_{b)} (v^a v^b Y_{lm}^*) d\Omega, \quad (21)$$

here A and  $\tilde{A}$  correspond to the projections of  $h_{ab}$  and  $\tilde{h}_{ab}$ , respectively. Projecting  $2\nabla_{(a} \xi_{b)}$  on the tensor harmonic basis, one could obtain the A-K components of  $2\nabla_{(a} \xi_{b)}$  as follows:

$$\begin{aligned} \Delta A &= -2 \frac{\partial P}{\partial t} - 2e^{-2\Lambda+2\Phi} \frac{\partial \Phi}{\partial r} R, & \Delta B &= -\frac{\partial S}{\partial t} + \frac{1}{r} P, \\ \Delta C &= -\frac{\partial}{\partial t} Q, & \Delta D &= \left( \frac{\partial}{\partial r} - 2 \frac{\partial \Phi}{\partial r} \right) P - \frac{\partial R}{\partial t}, \\ \Delta E &= 2 \frac{e^{-2\Lambda}}{r} R - \frac{l(l+1)}{r} S, & \Delta F &= \frac{2}{r} S, \\ \Delta G &= \frac{2}{r} Q, & \Delta H &= \frac{1}{r} R + \left( \frac{\partial}{\partial r} - \frac{1}{r} \right) S, \\ \Delta J &= \left( \frac{\partial}{\partial r} - \frac{1}{r} - \frac{\partial \Phi}{\partial r} - \frac{\partial \Lambda}{\partial r} \right) Q, & \Delta K &= \left( 2 \frac{\partial}{\partial r} - 2 \frac{\partial \Lambda}{\partial r} \right) R. \end{aligned} \quad (22)$$

### D. Gauge Choices

Generally speaking,  $\xi^a$  has four independent functions, representing four degrees of freedom in spherically symmetric spacetimes. Hence by properly choosing these four functions we can work with different gauges. For example, under the gauge transformation, the scalar function  $F(t, r)$  would transform as

$$\tilde{F} = F - \Delta F = F - \frac{2}{r} S. \quad (23)$$

By setting  $S = rF/2$ , one degree of the gauge freedom is fixed, and  $\tilde{F} = 0$ . Then substituting this  $S$  back into Eq.(22), one can move on to eliminate the next degrees of freedom. In even-parity perturbations, properly choosing the functions  $P$ ,  $R$  and  $S$  would fix three variables of the metric perturbation. In the odd-parity properly choosing the function  $Q$  would fix one variable of the metric perturbation.

Note that in the odd-parity sector,  $\Delta G$  is parallel to  $Q$ , but there exists some derivative relation between  $\Delta C$ ,  $\Delta J$  and  $Q$ . If we want to eliminate  $\tilde{G}$  under the gauge transformation, just set  $Q = rG/2$  then  $\tilde{G} = 0$  and  $\Delta C$  and  $\Delta J$  are uniquely determined. If we eliminate  $\tilde{C}$  under the gauge transformation rather than  $\tilde{G}$ , then  $Q$  could be an arbitrary function of  $r$  with some integration constant, which could not determine  $\Delta G$  and  $\Delta J$  completely. Such a choice obviously cannot completely fix the gauge freedom. Similarly, in the even-parity sector, in order to fix the gauge completely, one may first fix  $S$  from  $\Delta F$ . Afterward, there are still several choices to fix  $P$  and  $R$  via  $\Delta A$ ,  $\Delta B$ ,  $\Delta E$  or  $\Delta H$ . Below, we would discuss two useful gauge choices.

**Regge-Wheeler Gauge.** Regge and Wheeler first presented the RW gauge by setting certain RW variables to zero [1]. In the even-parity sector of the Schwarzschild spacetime, to eliminate three gauge freedom, RW set

$$h_0^{RW,even} = h_1^{RW,even} = G^{RW} = 0, \quad (24)$$

which corresponds to setting  $\tilde{B} = \tilde{F} = \tilde{H} = 0$  in the A-K notations. And in the odd-parity sector, RW set  $h_2^{RW} = 0$ , which corresponds to setting  $\tilde{G} = 0$  [28]. For general spherically symmetric spacetimes, we set

$$\begin{aligned} S^{RW} &= \frac{r}{2}F, & P^{RW} &= rB + \frac{r^2}{2}\frac{\partial F}{\partial t}, \\ R^{RW} &= rH - \frac{r}{2}\frac{\partial F}{\partial r}, & Q^{RW} &= \frac{r}{2}G, \end{aligned} \quad (25)$$

which means

$$\xi_a^{RW} = \left( rB + \frac{r^2}{2}\frac{\partial F}{\partial t} \right) v_a Y_{lm} + \left( rH - \frac{r}{2}\frac{\partial F}{\partial r} \right) Y_a^{R,lm} + \frac{r}{2}FY_a^{E,lm} + \frac{r}{2}GY_a^{B,lm}. \quad (26)$$

It should note that this gauge choice is exactly the same as the gauge choice with the A-K notations in the Schwarzschild spacetime, see, for example, Eq.(6.7) in [28].

**Easy gauge.** The EZ gauge was first introduced by Detweiler when he considered the self-force problem, in which the following metric components are set to zero,

$$\tilde{B} = \tilde{E} = \tilde{F} = \tilde{G} = 0. \quad (27)$$

To eliminate these metric components, the components of the gauge vector are chosen as

$$\begin{aligned} S^{EZ} &= \frac{r}{2}F, & P^{EZ} &= rB + \frac{r^2}{2}\frac{\partial F}{\partial t}, \\ R^{EZ} &= rl(l+1)e^{2\Lambda}F + \frac{r}{2}e^{2\Lambda}E, & Q^{EZ} &= \frac{r}{2}G, \end{aligned} \quad (28)$$

which means

$$\xi_a^{EZ} = \left( rB + \frac{r^2}{2}\frac{\partial F}{\partial t} \right) v_a Y_{lm} + \left( rl(l+1)e^{2\Lambda}F + \frac{r}{2}e^{2\Lambda}E \right) Y_a^{R,lm} + \frac{r}{2}FY_a^{E,lm} + \frac{r}{2}GY_a^{B,lm}. \quad (29)$$

### III. GAUGE INVARIANTS AND MASTER EQUATIONS

In this section, we would first introduce a general set of gauge invariant quantities in the spherically symmetric backgrounds. With these gauge invariants, we shall investigate how to construct the single master equation under certain gauge choices.

### A. Gauge Invariants

Generally speaking, when a gauge transformation is performed, we know from Eq.(22), the metric perturbations will change, for example,

$$\Delta G = \frac{2}{r}Q, \quad (30)$$

$$\Delta J = \left( \frac{\partial}{\partial r} - \frac{1}{r} - \frac{\partial \Phi}{\partial r} - \frac{\partial \Lambda}{\partial r} \right) Q, \quad (31)$$

from which, we obtain

$$\Delta J + \frac{r}{2} \left( \frac{\partial \Phi}{\partial r} + \frac{\partial \Lambda}{\partial r} - \frac{\partial}{\partial r} \right) \Delta G = 0. \quad (32)$$

The above equation indicates that one can define

$$\alpha = J + \frac{r}{2} \left( \frac{\partial \Phi}{\partial r} + \frac{\partial \Lambda}{\partial r} - \frac{\partial}{\partial r} \right) G, \quad (33)$$

and  $\alpha$  is a gauge invariant quantity. Note that there are seven even-parity metric components and three odd-parity metric components in the metric perturbation, while there are three even-parity components and one odd-parity component in  $\xi^a$ , which tell us that we can construct four independent even-parity gauge invariants and two independent odd-parity gauge invariants for  $l \geq 2$  cases. Following the construction of  $\alpha$ , we find that they are given by

$$\begin{aligned} \alpha &= J + \frac{r}{2} \left( \frac{\partial \Phi}{\partial r} + \frac{\partial \Lambda}{\partial r} - \frac{\partial}{\partial r} \right) G, \\ \beta &= -C - \frac{r}{2} \frac{\partial}{\partial t} G, \\ \chi &= H - \frac{1}{2} e^{2\Lambda} E - \frac{l(l+1)}{4} e^{2\Lambda} F - \frac{r}{2} \frac{\partial}{\partial r} F, \\ \psi &= \frac{1}{2} K - \frac{r}{2} e^{2\Lambda} \frac{\partial \Lambda}{\partial r} E - \frac{1}{2} e^{2\Lambda} E - \frac{r}{2} e^{2\Lambda} \frac{\partial}{\partial r} E - \frac{r}{4} l(l+1) e^{2\Lambda} \frac{\partial \Lambda}{\partial r} F - \frac{1}{4} l(l+1) e^{2\Lambda} F - \frac{r}{4} l(l+1) e^{2\Lambda} \frac{\partial}{\partial r} F, \\ \delta &= D + \frac{r}{2} e^{2\Lambda} \frac{\partial}{\partial t} E + \left( 2r \frac{\partial \Phi}{\partial r} - 1 \right) B - r \frac{\partial}{\partial r} B - \frac{r^2}{2} \frac{\partial^2}{\partial t \partial r} F - \left[ r - r^2 \frac{\partial \Phi}{\partial r} - \frac{r}{4} l(l+1) e^{2\Lambda} \right] \frac{\partial}{\partial t} F, \\ \epsilon &= -\frac{1}{2} A - \frac{r}{2} e^{2\Phi} \frac{\partial \Phi}{\partial r} E - r \frac{\partial}{\partial t} B - \frac{r}{4} l(l+1) e^{2\Phi} \frac{\partial \Phi}{\partial r} F - \frac{r^2}{2} \frac{\partial^2}{\partial t^2} F. \end{aligned} \quad (34)$$

These relations are the same as Eq. (7.5) of [28] when the background is vacuum, degenerates to the Schwarzschild case.

From now on, we shall work in the coordinates  $\tilde{x}^a$ , but for the save of simplicity, we shall drop all the tildes. Note that we use the superscript I or II to denote the quantities or parameters under the EZ gauge or the RW gauge, respectively. Adopting the specific EZ gauge, we have

$$B = E = F = G = 0. \quad (35)$$

Then, the gauge invariants become

$$\begin{aligned} \alpha &= J, & \beta &= -C, \\ \chi^I &= H, & \psi^I &= \frac{1}{2} K, \\ \delta^I &= D, & \epsilon^I &= -\frac{1}{2} A. \end{aligned} \quad (36)$$

Here  $\alpha$  and  $\beta$  are not superscripted because they are the same under the EZ gauge or the RW gauge. Similarly, adopting the specific RW gauge, we have

$$B = F = H = G = 0, \quad (37)$$

the even-parity gauge invariants become

$$\begin{aligned}\chi^{\text{II}} &= -\frac{1}{2}e^{2\Lambda}\text{E}, & \psi^{\text{II}} &= \frac{1}{2}\text{K} - \frac{1}{2}(r\Lambda' + 1)e^{2\Lambda}\text{E} - \frac{r}{2}e^{2\Lambda}\frac{\partial}{\partial r}\text{E}, \\ \delta^{\text{II}} &= \text{D} + \frac{r}{2}e^{2\Lambda}\frac{\partial}{\partial t}\text{E}, & \epsilon^{\text{II}} &= -\frac{1}{2}\text{A} - \frac{1}{2}r\Phi'e^{2\Phi}\text{E}.\end{aligned}\quad (38)$$

Using these gauges, each A-K projection of the linearized Einstein equations, i.e. Eqs.(B1)-(B10), can be rewritten as a combination of the gauge invariants we listed in Eq.(34). The results can be found in Appendix C. It is obvious that under the EZ gauge, the relationship between the gauge invariants and the perturbed metric components seems simpler, hence we first study the master equation under the EZ gauge.

## B. Master Equations for $l \geq 2$

Assuming that the perturbation of the stress-energy  $\mathcal{T}_{ab}$  is known, i.e.  $E_{\text{A}}-E_{\text{K}}$  are known quantities, now we look for the master equations, in terms of gauge invariants.

### 1. Even-parity Perturbations and the EZ gauge

The Bianchi identities indicate that not all seven even-parity projection equations in Appendix C are independent. It has shown that there are four independent gauge invariants. Noting the specific structure of the expressions of  $E_{\text{A}}^{\text{I}}$ ,  $E_{\text{D}}^{\text{I}}$ ,  $E_{\text{F}}^{\text{I}}$ ,  $E_{\text{H}}^{\text{I}}$  and  $E_{\text{K}}^{\text{I}}$ , we find that one can obtain  $\partial\delta^{\text{I}}/\partial t$  from  $\partial E_{\text{D}}^{\text{I}}/\partial t$ , and  $\epsilon^{\text{I}}$  from  $E_{\text{A}}^{\text{I}}$  or  $E_{\text{F}}^{\text{I}}$ . Substituting these relations into  $2E_{\text{H}}^{\text{I}} + E_{\text{K}}^{\text{I}}$ , after a large but tedious calculation, we find that the coupled partial differential equations for  $\chi^{\text{I}}$  and  $\psi^{\text{I}}$  can be constructed as

$$\frac{\partial}{\partial r}\chi^{\text{I}} = \frac{2e^{2\Lambda-2\Phi}r}{\tau^{\text{I}}(2\sigma^{\text{I}} - e^{2\Lambda}\lambda)} \left( \sigma^{\text{I}}\frac{\partial^2}{\partial t^2}\chi^{\text{I}} - 2\frac{\partial^2}{\partial t^2}\psi^{\text{I}} + \frac{1}{2}e^{2\Lambda}r\frac{\partial}{\partial t}E_{\text{D}}^{\text{I}} \right) + \frac{\gamma^{\text{I}}}{r\tau^{\text{I}}}\chi^{\text{I}} + \frac{\rho^{\text{I}}}{r\tau^{\text{I}}}\psi^{\text{I}} - \frac{1}{2}e^{2\Lambda}rE_{\text{F}}^{\text{I}} + \frac{e^{2\Lambda}r}{2\tau^{\text{I}}}(E_{\text{K}}^{\text{I}} + 2E_{\text{H}}^{\text{I}}), \quad (39)$$

$$\begin{aligned}\frac{\partial}{\partial r}\psi^{\text{I}} &= \frac{\kappa^{\text{I}}e^{2\Lambda-2\Phi}r}{\tau^{\text{I}}(2\sigma^{\text{I}} - e^{2\Lambda}\lambda)} \left( \sigma^{\text{I}}\frac{\partial^2}{\partial t^2}\chi^{\text{I}} - 2\frac{\partial^2}{\partial t^2}\psi^{\text{I}} + \frac{1}{2}e^{2\Lambda}r\frac{\partial}{\partial t}E_{\text{D}}^{\text{I}} \right) + \frac{\mu^{\text{I}}}{2r\tau^{\text{I}}}\chi^{\text{I}} - \frac{\nu^{\text{I}}}{r\tau^{\text{I}}}\psi^{\text{I}} \\ &\quad + \frac{\kappa^{\text{I}}e^{2\Lambda}r}{4\tau^{\text{I}}}(E_{\text{K}}^{\text{I}} + 2E_{\text{H}}^{\text{I}}) - \frac{1}{4}e^{4\Lambda-2\Phi}r(E_{\text{A}}^{\text{I}} + e^{2\Phi}\lambda E_{\text{F}}^{\text{I}}),\end{aligned}\quad (40)$$

where  $\lambda = l(l+1)$ , and  $\tau^{\text{I}}$ ,  $\sigma^{\text{I}}$ ,  $\gamma^{\text{I}}$ ,  $\rho^{\text{I}}$ ,  $\mu^{\text{I}}$ ,  $\nu^{\text{I}}$  are functions determined by the background spacetime, which can be find in Appendix D. Note that in these two equations, there is no spatial derivatives of the source term. Now the goal becomes decouple the gauge invariants  $\chi^{\text{I}}$  and  $\psi^{\text{I}}$ , e.g. the Eqs.(39) and (40). Introducing the gauge invariant

$$Z^{\text{I}(+)} = \sigma^{\text{I}}\chi^{\text{I}} - 2\psi^{\text{I}}, \quad (41)$$

we find that  $Z^{\text{I}(+)}$  satisfies the master equation

$$\left[ \frac{2e^{2\Lambda-2\Phi}r}{e^{2\Lambda}\lambda - 2\sigma^{\text{I}}} \left( 2r(\sigma^{\text{I}} - \kappa^{\text{I}})\frac{\partial^3}{\partial t^2\partial r} - \frac{N_{\text{T}}^{\text{I}}}{N^{\text{I}}(e^{2\Lambda}\lambda - 2\sigma^{\text{I}})\tau^{\text{I}}}\frac{\partial^2}{\partial t^2} \right) + 2r\tau^{\text{I}}\frac{\partial^2}{\partial r^2} - \frac{N_{\text{R}}^{\text{I}}}{N^{\text{I}}\sigma^{\text{I}}}\frac{\partial}{\partial r} - \frac{N_{\text{Z}}^{\text{I}}}{rN^{\text{I}}\sigma^{\text{I}}\tau^{\text{I}}} \right] Z^{\text{I}(+)} = S_{\text{even}}^{\text{I}}, \quad (42)$$

with the source term given by

$$\begin{aligned}S_{\text{even}}^{\text{I}} &= e^{4\Lambda-2\Phi}r \left[ \frac{N_{\text{A}}^{\text{I}}}{2N^{\text{I}}}E_{\text{A}}^{\text{I}} + r\tau^{\text{I}}\frac{\partial}{\partial r}E_{\text{A}}^{\text{I}} - \frac{rN_{\text{D}}^{\text{I}}}{N^{\text{I}}(e^{2\Lambda}\lambda - 2\sigma^{\text{I}})^2\tau^{\text{I}}}\frac{\partial}{\partial t}E_{\text{D}}^{\text{I}} - \frac{2r^2(\sigma^{\text{I}} - \kappa^{\text{I}})}{e^{2\Lambda}\lambda - 2\sigma^{\text{I}}}\frac{\partial^2}{\partial t\partial r}E_{\text{D}}^{\text{I}} \right] \\ &\quad + e^{2\Lambda}r \left[ \frac{N_{\text{F}}^{\text{I}}}{2N^{\text{I}}}E_{\text{F}}^{\text{I}} + r(e^{2\Lambda}\lambda - \sigma^{\text{I}})\tau^{\text{I}}\frac{\partial}{\partial r}E_{\text{F}}^{\text{I}} - \frac{N_{\text{HK}}^{\text{I}}}{N^{\text{I}}\tau^{\text{I}}}\left(E_{\text{H}}^{\text{I}} + \frac{1}{2}E_{\text{K}}^{\text{I}}\right) - r(\kappa^{\text{I}} - \sigma^{\text{I}})\left(\frac{\partial}{\partial r}E_{\text{K}}^{\text{I}} + 2\frac{\partial}{\partial r}E_{\text{H}}^{\text{I}}\right) \right].\end{aligned}\quad (43)$$

In the above equation, the parameters  $N^{\text{I}}$ ,  $N_{\text{T}}^{\text{I}}$ ,  $N_{\text{R}}^{\text{I}}$ ,  $N_{\text{Z}}^{\text{I}}$  and  $N_{\text{A}}^{\text{I}}$ ,  $N_{\text{D}}^{\text{I}}$ ,  $N_{\text{F}}^{\text{I}}$ ,  $N_{\text{HK}}^{\text{I}}$  are functions depending only on the background. The explicit expressions of these parameters can be found in Appendix D. Unlike the well-known Zerilli equation, the master equation under the EZ gauge is a third-order equation. However, rewriting Eq.(42) in the form,

$$\left( a\frac{\partial^3}{\partial t^2\partial r} + b\frac{\partial^2}{\partial t^2} + c\frac{\partial^2}{\partial r^2} + d\frac{\partial}{\partial r} + e \right) Z^{\text{I}(+)} = S_{\text{even}}^{\text{I}}, \quad (44)$$

and setting  $Z^{I(+)} = Z^I(r)e^{i\omega t}$  and  $S_{even}^I = S_{even}^I(r)e^{i\omega t}$ , we obtain a second-order differential equation

$$\left[ c \frac{\partial^2}{\partial r^2} + (d - a\omega^2) \frac{\partial}{\partial r} + (e - b\omega^2) \right] Z^{I(+)}(r) = S_{even}^I(r). \quad (45)$$

If the background becomes the Schwarzschild spacetime, the master equation (42) will reduce to the result that given in [28], in other words, the result of Zerilli [3]. The discussion about the degeneration of our result could be found in Appendix E.

Once Eq.(42) is solved, together with the definition of the master variable given by Eq.(41), the gauge invariants  $\chi$  and  $\psi$  can be solved from

$$\begin{aligned} \chi^I = \frac{1}{N^I} & \left[ \frac{4e^{2\Lambda-2\Phi}r^2(\kappa^I - \sigma^I)}{e^{2\Lambda}\lambda - 2\sigma^I} \frac{\partial^2}{\partial t^2} Z^{I(+)} - 2r\tau^I \frac{\partial}{\partial r} Z^{I(+)} - M_1^I Z^{I(+)} + e^{4\Lambda-2\Phi}r^2\tau^I E_A^I \right. \\ & \left. + \frac{2e^{4\Lambda-2\Phi}r^3(\kappa^I - \sigma^I)}{e^{2\Lambda}\lambda - 2\sigma^I} \frac{\partial}{\partial t} E_D^I + e^{2\Lambda}r^2(e^{2\Lambda}\lambda - \sigma^I)\tau^I E_F^I - e^{2\Lambda}r^2(\kappa^I - \sigma^I)(E_K^I + 2E_H^I) \right], \end{aligned} \quad (46)$$

$$\begin{aligned} \psi^I = \frac{\sigma^I}{2N^I} & \left[ \frac{4e^{2\Lambda-2\Phi}r^2(\kappa^I - \sigma^I)}{e^{2\Lambda}\lambda - 2\sigma^I} \frac{\partial^2}{\partial t^2} Z^{I(+)} - 2r\tau^I \frac{\partial}{\partial r} Z^{I(+)} + \frac{2M_2^I}{\sigma^I} Z^{I(+)} + e^{4\Lambda-2\Phi}r^2\tau^I E_A^I \right. \\ & \left. + \frac{2e^{4\Lambda-2\Phi}r^3(\kappa^I - \sigma^I)}{e^{2\Lambda}\lambda - 2\sigma^I} \frac{\partial}{\partial t} E_D^I + e^{2\Lambda}r^2(e^{2\Lambda}\lambda - \sigma^I)\tau^I E_F^I - e^{2\Lambda}r^2(\kappa^I - \sigma^I)(E_K^I + 2E_H^I) \right], \end{aligned} \quad (47)$$

where  $M_1^I$  and  $M_2^I$  are given in Appendix D. And then the remaining two even-parity gauge invariants  $\delta^I$  and  $\epsilon^I$  can be obtained from Eqs.(C3) and (C5), given respectively by

$$\delta^I = -\frac{1}{2 - 2e^{2\Lambda} + e^{2\Lambda}\lambda + 2r\Phi' - 6r\Phi'' + 2r\Phi'^2 + 2r^2\Phi'' - 2r^2\Lambda'\Phi'} \left( e^{2\Lambda}r^2 E_D^I + e^{2\Lambda}\lambda r \frac{\partial}{\partial t} \chi^I - 4r \frac{\partial}{\partial t} \psi^I \right), \quad (48)$$

$$\epsilon^I = \frac{1}{2} e^{-2\Lambda+2\Phi} \left[ 2r \frac{\partial}{\partial r} \chi^I + 2(1 - r\Lambda' + r\Phi')\chi^I - 2\psi^I + e^{2\Lambda}r^2 E_F^I \right]. \quad (49)$$

From Eq.(36), the even-parity metric perturbation components A, D, H, K could be read out directly.

## 2. Even-parity Perturbations and the RW gauge

Similar to the development provided in the previous subsection, using the field equations  $E_A^{\text{II}}$  to  $E_K^{\text{II}}$  in Appendix C, one can eliminate the gauge invariants  $\delta^{\text{II}}$  and  $\epsilon^{\text{II}}$ , and then obtain the following coupled equations,

$$\begin{aligned} \frac{\partial}{\partial r} \chi^{\text{II}} = \frac{2e^{2\Lambda-2\Phi}r}{\tau^{\text{II}}(\sigma^{\text{II}} - 2r\Lambda' - 2r\Phi')} & \left( \sigma^{\text{II}} \frac{\partial^2}{\partial t^2} \chi^{\text{II}} - 2 \frac{\partial^2}{\partial t^2} \psi^{\text{II}} + \frac{1}{2} e^{2\Lambda} r \frac{\partial}{\partial t} E_D^{\text{II}} \right) + \frac{\rho^{\text{II}}}{r\tau^{\text{II}}} \chi^{\text{II}} + \frac{\sigma^{\text{II}} - 2}{r\tau^{\text{II}}} \psi^{\text{II}} \\ & - \frac{e^{2\Lambda}r}{2\tau^{\text{II}}} (e^{2\Lambda}\lambda + 2r\Phi' - 2) E_F^{\text{II}} + \frac{e^{2\Lambda}r}{2\tau^{\text{II}}} (2E_H^{\text{II}} + E_K^{\text{II}}), \end{aligned} \quad (50)$$

$$\begin{aligned} \frac{\partial}{\partial r} \psi^{\text{II}} = \frac{\sigma^{\text{II}} e^{2\Lambda-2\Phi}r}{\tau^{\text{II}}(\sigma^{\text{II}} - 2r\Lambda' - 2r\Phi')} & \left( \sigma^{\text{II}} \frac{\partial^2}{\partial t^2} \chi^{\text{II}} - 2 \frac{\partial^2}{\partial t^2} \psi^{\text{II}} + \frac{1}{2} e^{2\Lambda} r \frac{\partial}{\partial t} E_D^{\text{II}} \right) + \frac{\mu^{\text{II}}}{r\tau^{\text{II}}} \chi^{\text{II}} - \frac{\nu^{\text{II}}}{r\tau^{\text{II}}} \psi^{\text{II}} \\ & + \frac{e^{2\Lambda}r}{4\tau^{\text{II}}} (\kappa^{\text{II}} E_F^{\text{II}} + 2\sigma^{\text{II}} E_H^{\text{II}} + \sigma^{\text{II}} E_K^{\text{II}}) - \frac{1}{4} e^{4\Lambda-2\Phi} r E_A^{\text{II}}, \end{aligned} \quad (51)$$

where the parameters  $\sigma^{\text{II}}$ ,  $\tau^{\text{II}}$ ,  $\kappa^{\text{II}}$ ,  $\rho^{\text{II}}$ ,  $\mu^{\text{II}}$ ,  $\nu^{\text{II}}$  depend only on the background, and the explicit expressions of them can be found in Appendix D.

We find that the master variable can be similarly constructed as,

$$Z^{\text{II}(+)} = \sigma^{\text{II}} \chi^{\text{II}} - 2\psi^{\text{II}}, \quad (52)$$



and then the master equation under the RW gauge is given by

$$\left[ \frac{2r\sigma^{\text{II}}\tau^{\text{II}}N_T^{\text{II}}}{N^{\text{II}}} \left( -\frac{\partial^2}{\partial t^2} + \frac{1}{N_T^{\text{II}}} \frac{\partial^2}{\partial r^2} \right) + \frac{N_R^{\text{II}}}{(N^{\text{II}})^2} \frac{\partial}{\partial r} + \frac{N_Z^{\text{II}}}{(N^{\text{II}})^2 r\tau^{\text{II}}} \right] Z^{\text{II}(+) } = S_{\text{even}}^{\text{II}}, \quad (53)$$

with the source term is given by

$$S_{\text{even}}^{\text{II}} = e^{4\Lambda-2\Phi} r \left[ \frac{N_A^{\text{II}}}{2(N^{\text{II}})^2} E_A^{\text{II}} + \frac{r\sigma^{\text{II}}\tau^{\text{II}}}{N^{\text{II}}} \frac{\partial}{\partial r} E_A^{\text{II}} + \frac{r\sigma^{\text{II}}}{\tau^{\text{II}}(\sigma^{\text{II}} - 2r\Lambda' - 2r\Phi')} \frac{\partial}{\partial t} E_D^{\text{II}} \right] - e^{2\Lambda} r \left\{ \frac{N_F^{\text{II}}}{2(N^{\text{II}})^2 \tau^{\text{II}}} E_F^{\text{II}} + \frac{r\sigma^{\text{II}}[\kappa^{\text{II}} + \sigma^{\text{II}}(e^{2\Lambda}\lambda + 2r\Phi' - 2)]}{N^{\text{II}}} \frac{\partial}{\partial r} E_F^{\text{II}} - \frac{\sigma^{\text{II}}}{2\tau^{\text{II}}} (2E_H^{\text{II}} + E_K^{\text{II}}) \right\}, \quad (54)$$

where  $N^{\text{II}}$ ,  $N_T^{\text{II}}$ ,  $N_R^{\text{II}}$ ,  $N_Z^{\text{II}}$ ,  $N_A^{\text{II}}$ ,  $N_F^{\text{II}}$  all depend on the background, which can be found in Appendix D. Under the RW gauge, the master equation is a wave-like equation. We find that if the metric reduces to the Schwarzschild spacetime, then Eq.(53) will reduce to that given by Eq.(E6).

Once Eq.(53) is solved, one can get  $\chi^{\text{II}}$  and  $\psi^{\text{II}}$  from

$$\chi^{\text{II}} = \frac{1}{N^{\text{II}}} \left\{ 2r\tau^{\text{II}} \frac{\partial}{\partial r} Z^{\text{II}(+) } + M_1^{\text{II}} Z^{\text{II}(+) } - e^{4\Lambda-2\Phi} r^2 \tau^{\text{II}} E_A^{\text{II}} + e^{2\Lambda} r^2 [\kappa^{\text{II}} + \sigma^{\text{II}}(e^{2\Lambda}\lambda + 2r\Phi' - 2)] E_F^{\text{II}} \right\}, \quad (55)$$

$$\psi^{\text{II}} = \frac{1}{N^{\text{II}}} \left\{ r\sigma^{\text{II}}\tau^{\text{II}} \frac{\partial}{\partial r} Z^{\text{II}(+) } + M_2^{\text{II}} Z^{\text{II}(+) } - \frac{1}{2} e^{4\Lambda-2\Phi} r^2 \sigma^{\text{II}} \tau^{\text{II}} E_A^{\text{II}} + \frac{1}{2} e^{2\Lambda} r^2 \sigma^{\text{II}} [\kappa^{\text{II}} + \sigma^{\text{II}}(e^{2\Lambda}\lambda + 2r\Phi' - 2)] E_F^{\text{II}} \right\}, \quad (56)$$

and the remaining two gauge invariants  $\delta^{\text{II}}$  and  $\epsilon^{\text{II}}$  can be obtained from  $E_D^{\text{II}}$  and  $E_F^{\text{II}}$ , are given respectively by

$$\delta^{\text{II}} = -\frac{e^{2\Lambda} r^2 E_D + r(2 - 2e^{2\Lambda} + e^{2\Lambda}\lambda + 6r\Phi' - 2r\Lambda' + 2r^2\Phi'^2 - 2r^2\Lambda'\Phi' + 2r^2\Phi'') \frac{\partial}{\partial t} \chi - 4r \frac{\partial}{\partial t} \psi}{2 - 2e^{2\Lambda} + e^{2\Lambda}\lambda + 2r\Phi' - 6r\Phi' + 2r\Phi'^2 + 2r^2\Phi'' - 2r^2\Lambda'\Phi'}, \quad (57)$$

$$\epsilon^{\text{II}} = \frac{1}{2} e^{-2\Lambda+2\Phi} \left[ 2r \frac{\partial}{\partial r} \chi + 2(1 - r\Lambda' + r\Phi') \chi - 2\psi + e^{2\Lambda} r^2 E_F \right]. \quad (58)$$

When  $\chi^{\text{II}}$ ,  $\psi^{\text{II}}$ ,  $\delta^{\text{II}}$  and  $\epsilon^{\text{II}}$  are solved, the even-parity metric perturbation components A, D, E, K are given by

$$\begin{aligned} A &= 2e^{-2\Lambda+2\Phi} r\Phi' \chi^{\text{II}} - 2\epsilon^{\text{II}}, & D &= \delta^{\text{II}} + r \frac{\partial}{\partial t} \chi^{\text{II}}, \\ E &= -2e^{-2\Lambda} \chi^{\text{II}}, & K &= 2\psi^{\text{II}} + 2(r\Lambda' - 1)\chi^{\text{II}} - 2r \frac{\partial}{\partial r} \chi^{\text{II}}. \end{aligned} \quad (59)$$

Under the RW gauge, the master variable can be written as the combination of the gauge invariants  $\chi^{\text{II}}$  and  $\psi^{\text{II}}$ . In Zerilli's paper [3], the master variable was written as the combination of  $K_{LM}$  and  $R_{LM}$ , which correspond to a combination of  $\chi^{\text{II}}$  and  $\delta^{\text{II}}$  in our paper.

### 3. Odd-parity Perturbations

For the odd parity perturbations, the EZ gauge and the RW gauge are identical. So, in the following we shall not distinguish them. We construct the odd-parity master variable as

$$Z^{(-)} = (r + r^2\Lambda' + r^2\Phi') \beta - r^2 \frac{\partial}{\partial r} \beta + r^2 \frac{\partial}{\partial t} \alpha, \quad (60)$$

then the master equation is given by

$$\left\{ -\frac{\partial^2}{\partial t^2} + e^{-2\Lambda+2\Phi} \frac{X}{Y} \left[ \frac{\partial^2}{\partial r^2} - \left( \frac{Y'}{Y} + 3\Lambda' + 3\Phi' \right) \frac{\partial}{\partial r} + \frac{N_{\text{odd}}}{r^2} \right] \right\} Z^{(-)} = S_{\text{odd}}, \quad (61)$$

with the source term given by

$$S_{odd} = e^{2\Phi} r^2 \frac{X}{Y} \left[ \left( \frac{1}{r} + \Lambda' - \Phi' - \frac{Y'}{Y} \right) E_C + \frac{\partial}{\partial r} E_C + \frac{Y}{X} \frac{\partial}{\partial t} E_J \right], \quad (62)$$

where  $X$ ,  $Y$  and  $N_{odd}$  are parameters only depending on the background, and the explicit expressions of which can be found in Appendix D. Once Eq.(61) is solved, together with the structure of the odd-parity master variable Eq.(60) and the perturbed field equations Eqs.(C15) and (C17), we have

$$\alpha = -\frac{e^{2\Lambda}}{X} \left( r^2 E_J + e^{-2\Phi} \frac{\partial}{\partial t} Z^{(-)} \right), \quad (63)$$

and

$$\beta = \frac{1}{Y} \left[ e^{2\Lambda} r^2 E_C - \left( \frac{1}{r} - 2\Lambda' - 2\Phi' \right) Z^{(-)} - \frac{\partial}{\partial r} Z^{(-)} \right]. \quad (64)$$

Then, the odd-parity metric perturbation components C and J can be read off from these expressions.

### C. Specific Cases for $l = 0, 1$

In the previous subsection, we have discussed the construction of master variables and master equations for even-parity and odd parity perturbations for the  $l \geq 2$  cases. However, the decomposition of metric perturbation would take some other form for the specific cases of  $l = 0, 1$ . In this section, we investigate how to solve the metric perturbation for  $l = 0, 1$ . Note that in this subsection, we use, e.g.  $A_0$  and  $A_1$ , to represent the scalar functions in metric perturbation and gauge vector for  $l = 0$  and  $l = 1$  cases, respectively. And we also use, e.g.  $E_A^{(l=0)}$  and  $E_A^{(l=1)}$  to represent the projection function  $E_A$  for  $l = 0$  and  $l = 1$  cases, respectively.

#### 1. $l = 0$ case

For the special case  $l = 0$ , there remains one scalar spherical harmonic function  $Y^{00} = \frac{1}{2\sqrt{\pi}}$ , which leads the metric perturbation to

$$h_{ab}^{(l=0)} = \frac{1}{2\sqrt{\pi}} (A_0 v_a v_b + 2D_0 v_{(a} n_{b)} + E_0 \sigma_{ab} + K_0 n_a n_b), \quad (65)$$

and the gauge vector takes the form

$$\xi_a^{(l=0)} = \frac{1}{2\sqrt{\pi}} (P_0 v_a + R_0 n_a). \quad (66)$$

In the  $l = 0$  case, we only have four metric perturbation components, and all of them are even-parity. Under the gauge transformation, the metric perturbation will be transformed as

$$\begin{aligned} \Delta A_0 &= -2 \frac{\partial}{\partial t} P_0 - 2e^{-2\Lambda+2\Phi} \Phi' R_0, & \Delta D_0 &= \left( \frac{\partial}{\partial r} - 2\Phi' \right) P_0 - \frac{\partial}{\partial t} R_0, \\ \Delta E_0 &= 2 \frac{e^{-2\Lambda}}{r} R_0, & \Delta K_0 &= \left( 2 \frac{\partial}{\partial r} - 2\Lambda' \right) R_0. \end{aligned} \quad (67)$$

The structure of  $\Delta A_0$ ,  $\Delta D_0$  and  $\Delta K_0$  are the same as the  $l \geq 2$  cases, but  $\Delta E_0$  is different from the expression given in Eq.(22) since S does not exist for  $l = 0$ . From Eq.(67), we can construct two gauge invariants,

$$\begin{aligned} \psi_0 &= \frac{1}{2} K_0 - \frac{r}{2} e^{2\Lambda} \Lambda' E_0 - \frac{1}{2} e^{2\Lambda} E_0 - \frac{r}{2} e^{2\Lambda} \frac{\partial}{\partial r} E_0, \\ \alpha_0 &= \frac{1}{2} \frac{\partial}{\partial r} A_0 - \Phi' A_0 + \frac{\partial}{\partial t} D_0 + \frac{1}{2} e^{2\Phi} (\Phi' + r\Phi'') E_0 + \frac{1}{2} e^{2\Phi} r \Phi' \frac{\partial}{\partial r} E_0 + \frac{1}{2} e^{2\Lambda} r \frac{\partial^2}{\partial t^2} E_0. \end{aligned} \quad (68)$$

Choosing the gauge  $D_0 = E_0 = 0$ , we find that  $E_A^{(l=0)}$ ,  $E_D^{(l=0)}$ ,  $E_E^{(l=0)}$  and  $E_K^{(l=0)}$  are given by

$$\begin{aligned}
E_A^{(l=0)} &= 2r^{-2}e^{-2\Lambda}(-1 + e^{2\Lambda} - 2r\Phi' + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')A_0 \\
&\quad + 2r^{-2}e^{-4\Lambda+2\Phi}(4r\Lambda' - 1)K_0 - 2r^{-1}e^{-4\Lambda+2\Phi}\frac{\partial}{\partial r}K_0, \\
E_D^{(l=0)} &= 2r^{-1}e^{-2\Lambda}\frac{\partial}{\partial t}K_0, \\
E_E^{(l=0)} &= 2r^{-1}e^{-2\Lambda-2\Phi}(r\Lambda'\Phi' - r\Phi'' - \Phi')A_0 - r^{-1}e^{-2\Lambda-2\Phi}(r\Lambda' + 2r\Phi' - 1)\frac{\partial}{\partial r}A_0 + e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial r^2}A_0 \\
&\quad + 2r^{-1}e^{-4\Lambda}(\Phi' + r\Phi'^2 + r\Phi'' - 2r\Lambda'\Phi' - 2\Lambda')K_0 + r^{-1}e^{-4\Lambda}(r\Phi' + 1)\frac{\partial}{\partial r}K_0 + e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial t^2}K_0, \\
E_K^{(l=0)} &= -4r^{-1}e^{-2\Phi}\Phi'A_0 + 2r^{-1}e^{-2\Phi}\frac{\partial}{\partial r}A_0 + 2r^{-2}e^{-2\Lambda}(e^{2\Lambda} + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')K_0. \tag{69}
\end{aligned}$$

And the gauge invariants  $\psi_0$  and  $o_0$  become

$$\psi_0 = \frac{1}{2}K_0, \quad o_0 = \frac{1}{2}\frac{\partial}{\partial r}A_0 - \Phi'A_0. \tag{70}$$

Note that under this specific gauge,  $E_A^{(l=0)}$  is not easy to be expressed as a combination of  $\psi_0$  and  $o_0$ . A more convenient way is to define the quantity  $E_A^{(l=0)}$  as

$$\begin{aligned}
&\frac{\partial}{\partial r}E_A^{(l=0)} - \mu_0E_A^{(l=0)} - 2\Phi'E_A^{(l=0)} \\
&= 4r^{-2}e^{-2\Lambda}(-1 + e^{2\Lambda} + 2r\Lambda' - 2r\Phi' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')o_0 \\
&\quad + 2\nu_0\psi_0 + 2\tau_0\frac{\partial}{\partial r}\psi_0 - 4r^{-1}e^{-4\Lambda+2\Phi}\frac{\partial^2}{\partial r^2}\psi_0, \tag{71}
\end{aligned}$$

where the explicit expressions of  $\mu_0$ ,  $\nu_0$  and  $\tau_0$  can be found in Appendix D. And  $E_D^{(l=0)}$ ,  $E_E^{(l=0)}$  and  $E_K^{(l=0)}$  are given by

$$E_D^{(l=0)} = 4r^{-1}e^{-2\Lambda}\frac{\partial}{\partial t}\psi_0, \tag{72}$$

$$\begin{aligned}
E_E^{(l=0)} &= -2r^{-1}e^{-2\Lambda-2\Phi}(r\Lambda' - 1)o_0 + 2e^{-2\Lambda-2\Phi}\frac{\partial}{\partial r}o_0 + 4r^{-1}e^{-4\Lambda}(\Phi' - 2\Lambda' + r\Phi'^2 + r\Phi'' - 2r\Lambda'\Phi')\psi_0 \\
&\quad + 2r^{-1}e^{-4\Lambda}(r\Phi' + 1)\frac{\partial}{\partial r}\psi_0 + 2e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial t^2}\psi_0, \tag{73}
\end{aligned}$$

$$E_K^{(l=0)} = 4r^{-1}e^{-2\Phi}o_0 + 4r^{-2}e^{-2\Lambda}(e^{2\Lambda} + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')\psi_0. \tag{74}$$

From Eq.(74) we get

$$o_0 = \frac{r}{4}e^{2\Phi}E_K^{(l=0)} - r^{-1}e^{2\Phi-2\Lambda}(e^{2\Lambda} + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')\psi_0. \tag{75}$$

Then substituting  $o_0$  into Eq.(71),  $\psi_0$  will satisfies the relation

$$\begin{aligned}
&\left[ 2\nu_0 - 4r^{-3}e^{-4\Lambda+2\Phi}(e^{2\Lambda} + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')(-1 + e^{2\Lambda} - 2r\Phi' - r^2\Phi'^2 + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'') \right. \\
&\quad \left. + 2\tau_0\frac{\partial}{\partial r} - 4r^{-1}e^{-4\Lambda+2\Phi}\frac{\partial^2}{\partial r^2} \right]\psi_0 = S_1^{(l=0)}, \tag{76}
\end{aligned}$$

where  $S^{(l=0)}$  is the source term given by

$$S_1^{(l=0)} = \frac{\partial}{\partial r}E_A^{(l=0)} - \mu_0E_A^{(l=0)} - 2\Phi'E_A^{(l=0)} - r^{-1}e^{-2\Lambda+2\Phi}(-1 + e^{2\Lambda} + 2r\Lambda' - 2r\Phi' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')E_K^{(l=0)}. \tag{77}$$

Meanwhile, substituting  $o_0$  and (72) into Eq.(73),  $\psi_0$  satisfies

$$\left( \iota_0\frac{\partial}{\partial r} - \sigma_0 \right)\psi_0 = \frac{r}{2}e^{4\Lambda}S_2^{(l=0)}, \tag{78}$$

where  $\iota_0$  and  $\sigma_0$  are parameters which can be found in Appendix D, and  $S_2^{l=0}$  is the source term given by

$$S_2^{(l=0)} = \frac{1}{2}e^{-2\Phi}r\frac{\partial}{\partial t}E_D^{(l=0)} - E_E^{(l=0)} + \frac{1}{2}e^{-2\Lambda}\left[r\frac{\partial}{\partial r}E_K^{(l=0)} + (2 - r\Lambda' + 2r\Phi')E_K^{(l=0)}\right]. \quad (79)$$

One can obtain  $\partial\psi_0/\partial r$  from Eq.(78), and then substitute it to Eq.(76) to get the equation of  $\psi_0$ , which will be an algebraic equation. Then  $o_0$  can also be obtained. Note that in the Schwarzschild spacetime, both sides of Eq.(76) are singular.

## 2. $l = 1$ case

For the case  $l = 1$ , the tensor harmonic basis  $T_{ab}^{E2,1m}$  and  $T_{ab}^{B2,1m}$  vanish, hence the scalar functions  $F_1$  and  $G_1$  would no longer exist. The four components of the gauge vector  $\xi_a$  imply that there are only three gauge invariants for even-parity and one odd gauge invariant for odd-parity. The remaining eight projections of  $2\nabla_{(a}\xi_{b)}$  are the same as for  $l \geq 2$  cases.

First, we investigate the even-parity sector. As we explained for  $l \geq 2$  cases, one should first determine the function  $S$  of the gauge vector  $\xi_a$  to make  $F$  vanish under the gauge transformation. However, when  $l = 1$ , the lack of  $F_1$  prevents us from constructing the gauge invariants as the cases for  $l \geq 2$ . The even-parity  $l = 1$  metric perturbations are related to the linear momentum of the system [4]. Note that if we take the gauge choice  $B_1 = E_1 = H_1 = 0$ , which was introduced by Zerilli in the Schwarzschild spacetime, the relation between  $E_A$  and metric components would no longer be a simple relationship. However, taking the gauge

$$A_1 = E_1 = H_1 = 0, \quad (80)$$

we find that  $E_A$  and the metric perturbed component  $K$  have a simple relation. In particular, we find

$$E_A^{(l=1)} = -2r^{-1}e^{-4\Lambda+2\Phi}\left(\frac{\partial}{\partial r}K_1 + r^{-1}(1 + e^{2\Lambda} - 4r\Lambda')K_1\right), \quad (81)$$

$$E_D^{(l=1)} = 2r^{-2}(1 - 2r\Phi')B_1 + 2r^{-1}\frac{\partial}{\partial r}B_1 + 2r^{-1}e^{-2\Lambda}\frac{\partial}{\partial t}K_1 - 2r^{-2}e^{-2\Lambda}(1 + r\Phi' + r^2\Phi'^2 - 3r\Lambda' - r^2\Lambda'\Phi' + r^2\Phi'')D_1, \quad (82)$$

$$E_K^{(l=1)} + 4E_H^{(l=1)} = 2r^{-2}e^{-2\Lambda}(-2 + e^{2\Lambda} - 2r\Phi' - r^2\Phi'^2 + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'')K_1 + 4r^{-1}e^{-2\Phi}(1 - e^{2\Lambda})\frac{\partial}{\partial t}B_1 - 4e^{-2\Phi}\frac{\partial^2}{\partial t\partial r}B_1. \quad (83)$$

From Eq.(81), the metric perturbation function  $K_1$  can be found. This solution can be used in Eq.(83) to solve  $B_1$ , and then from Eq.(82),  $D_1$  could also be solved. The remaining quantities, such as  $E_B$  or  $E_E$ , can be used to check the consistency of the solutions.

Then we investigate the odd-parity sector. For  $l \geq 2$  cases, one should determine the component function  $Q$  of the gauge vector  $\xi_a$  to make  $G$  vanish under the gauge transformation. For  $l = 1$  case,  $G_1$  no longer exists, which means that we can not construct the gauge invariants  $\alpha$  and  $\beta$  as in the  $l \geq 2$  cases. However, from the components of the projection of  $2\nabla_{(a}\xi_{b)}$ ,

$$\Delta C_1 = -\frac{\partial}{\partial t}Q, \quad (84)$$

$$\Delta J_1 = \left(\frac{\partial}{\partial r} - \frac{1}{r} - \Phi' - \Lambda'\right)Q, \quad (85)$$

we can construct a gauge invariant property  $\alpha_1$  as

$$\alpha_1 = \frac{\partial}{\partial t}J_1 + \frac{\partial}{\partial r}C_1 - \left(\frac{1}{r} + \Phi' + \Lambda'\right)C_1. \quad (86)$$

Then, for  $l = 1$ , the odd-parity of the projection of EFEs are

$$E_C^{(l=1)} = -r^{-2}e^{-2\Lambda} \left( 3 - e^{2\Lambda} - 2r^2\Lambda'^2 + 6r\Phi' + r^2\Phi'^2 - 2r\Lambda' - 7r^2\Lambda'\Phi' + r^2\Lambda'' + 4r^2\Phi'' \right) C_1 \\ - r^{-1}e^{-2\Lambda}(-2 + 3r\Lambda' + 3r\Phi')\frac{\partial}{\partial r}C_1 + e^{-2\Lambda}\frac{\partial^2}{\partial r^2}C_1 - r^{-1}e^{-2\Lambda}(-3 + 2r\Lambda' + 2r\Phi')\frac{\partial}{\partial t}J_1 + e^{-2\Lambda}\frac{\partial^2}{\partial t\partial r}J_1, \quad (87)$$

$$E_J^{(l=1)} = -r^{-2}e^{-2\Lambda} \left( 1 - e^{2\Lambda} - 5r\Lambda' + 3r\Phi' + 3r^2\Phi'^2 + 3r^2\Phi'' - 3r^2\Lambda'\Phi' \right) J_1 \\ - e^{-2\Phi}\frac{\partial^2}{\partial t^2}J_1 + r^{-1}e^{-2\Phi}(1 + r\Lambda' + r\Phi')\frac{\partial}{\partial t}C_1 - e^{-2\Phi}\frac{\partial^2}{\partial t\partial r}C_1. \quad (88)$$

Together with Eq.(86), Eq.(87) and Eq.(88) will be decoupled. And the master equation for the odd-parity perturbation can be written as

$$\left\{ -\frac{\partial^2}{\partial t^2} + e^{-2\Lambda+2\Phi}\frac{Y_1}{X_1} \left[ \frac{\partial^2}{\partial r^2} + \left( \frac{4}{r} - 3\Lambda' - 3\Phi' - \frac{X_1'}{X_1}\frac{\partial}{\partial r} \right) + \frac{N_1}{r^2} \right] \right\} \alpha_1 = S_{odd}^{(l=1)}, \quad (89)$$

where  $S_{odd}^{(l=1)}$  is the source term, which is given by

$$S_{odd}^{(l=1)} = e^{2\Phi}\frac{Y_1}{X_1} \left[ \left( \frac{1}{r} + \Lambda' - \Phi' - \frac{X_1'}{X_1} \right) E_C^{(l=1)} + \frac{\partial}{\partial r}E_C^{(l=1)} + \frac{X_1}{Y_1}\frac{\partial}{\partial t}E_J^{(l=1)} \right]. \quad (90)$$

In the above equations,  $X_1$ ,  $Y_1$  and  $N_1$  are all depend only on background, which can be found in Appendix D. Once Eq.(89) is solved, the metric perturbation components  $C_1$  and  $J_1$  can be determined by

$$C_1 = \frac{r}{X_1} \left[ e^{2\Lambda}rE_C^{(l=1)} + (-3 + 2r\Lambda' + 2r\Phi')\alpha_1 - r\frac{\partial}{\partial r}\alpha_1 \right], \quad (91)$$

$$J_1 = \frac{e^{2\Lambda-2\Phi}}{Y_1}r^2 \left( e^{2\Phi}E_J^{(l=1)} + \frac{\partial}{\partial t}\alpha_1 \right). \quad (92)$$

#### IV. A POINT PARTICLE AS THE SOURCE

In this section, we present a simple example that a small object moves along a circular orbit around the center of a spherically symmetric spacetime. We analyse the solutions for  $l = 0, 1$  in this section. For  $l \geq 2$  cases, we only provide a general approach.

Assuming that the small object moves along the worldline  $z(\tau)$  with mass  $\mu$  and four velocity  $u_a$ , the stress-energy of this point particle takes the form [36]

$$T_{ab} = \int \frac{\mu u_a u_b}{\sqrt{-g^{(0)}}} \delta^4[x - z(\tau)] d\tau, \quad (93)$$

where  $\tau$  is the proper time,  $g^{(0)}$  and  $\delta^4$  are the determinant of the background and the four-dimensional Dirac delta function, respectively. Generalizing the standard analysis from the textbook [37], the time-like geodesics in general spherically symmetric spacetime is

$$-1 = g_{ab}u^a u^b = -e^{2\Phi}\dot{t}^2 + e^{2\Lambda}\dot{r}^2 + r^2\dot{\varphi}^2. \quad (94)$$

Using the static Killing field  $\xi^a = (\partial/\partial t)^a$  and the rotational Killing field  $\psi^a = (\partial/\partial\varphi)^a$ , two conserved quantities  $\mathcal{E}$  and  $L$  can be defined as

$$\mathcal{E} = -g_{ab}\xi^a u^b = e^{2\Phi}\dot{t}, \quad L = g_{ab}\psi^a u^b = r^2\dot{\varphi}. \quad (95)$$

Then the geodesic equation reads

$$\frac{1}{2}\dot{r}^2 + \frac{1}{2}e^{-2\Lambda} \left( \frac{L^2}{r^2} + 1 \right) = \frac{1}{2}e^{-2\Lambda-2\Phi}\mathcal{E}^2. \quad (96)$$

Considering that the particle moves along a circular orbit with radio  $R$ , which is determined by  $\partial V/\partial r = 0$ , the four velocity of the particle can be written as

$$u_a = (-\mathcal{E}, 0, 0, L), \quad (97)$$

and now  $\mathcal{E}$  and  $L$  represent the energy and angular momentum of the particle, given respectively by,

$$\mathcal{E} = \frac{e^{\Phi|_R}}{\sqrt{R\Lambda'|_R + 1}}, \quad L = R\sqrt{\frac{-R\Lambda'|_R}{R\Lambda'|_R + 1}}. \quad (98)$$

where  $|_R$  denotes the corresponding function taking its value at orbit radius  $R$ . The orbital frequency can also be defined as

$$\Omega^2 = \left(\frac{u^\varphi}{u^t}\right)^2 = -\frac{e^{2\Phi}\Lambda'|_R}{R}, \quad (99)$$

which gives two useful relations

$$\mathcal{E} = \frac{e^{2\Phi|_R}}{\Omega R^2} L, \quad \Omega L = -R\Lambda'|_R \cdot \mathcal{E}. \quad (100)$$

Projecting the stress-energy tensor  $T_{ab}$  to the harmonic basis, one can obtain  $E_A$ - $E_K$ . The results reveal that  $E_D$ ,  $E_H$ ,  $E_J$  and  $E_K$  vanish automatically, and the non-vanishing projections of the stress-energy tensor are given as follows. For  $l \geq 0$ ,

$$E_A^{lm} = -16\pi e^{-\Lambda+\Phi} \frac{\mu\mathcal{E}}{r^2} \delta(r-R) Y_{lm}^* \left(\frac{\pi}{2}, \Omega t\right), \quad (101)$$

$$E_E^{lm} = -8\pi e^{-\Lambda-\Phi} \frac{\mu L \Omega R^2}{r^4} \delta(r-R) Y_{lm}^* \left(\frac{\pi}{2}, \Omega t\right), \quad (102)$$

where  $E_A$  and  $E_E$  satisfy the relation  $E_A = -\frac{2e^{2\Phi} r^2}{R^3 \Lambda'} E_E$ . For  $l \geq 1$ ,

$$E_B^{lm} = -\frac{16\pi}{l(l+1)} e^{-\Lambda-\Phi} \frac{\mu\mathcal{E}\Omega R^2}{r^3} \delta(r-R) \left(\frac{\partial}{\partial\varphi} Y_{lm}^*(\theta, \varphi)\right) \Big|_{\theta=\frac{\pi}{2}, \varphi=\Omega t}, \quad (103)$$

$$E_C^{lm} = -\frac{16\pi}{l(l+1)} \frac{\mu\mathcal{E}\Omega R^2}{r^3} \delta(r-R) \left(\frac{\partial}{\partial\theta} Y_{lm}^*(\theta, \varphi)\right) \Big|_{\theta=\frac{\pi}{2}, \varphi=\Omega t}, \quad (104)$$

and for  $l \geq 2$ ,

$$E_F^{lm} = \frac{16\pi(l-2)!}{(l+2)!} e^{-\Lambda-\Phi} \frac{\mu L \Omega R^2}{r^4} \delta(r-R) \left[2\frac{\partial^2}{\partial\theta^2} Y_{lm}^*(\theta, \varphi) + l(l+1)Y_{lm}^*(\theta, \varphi)\right] \Big|_{\theta=\frac{\pi}{2}, \varphi=\Omega t}, \quad (105)$$

$$E_G^{lm} = -\frac{32\pi(l-2)!}{(l+2)!} \frac{\mu L \Omega R^2}{r^4} \delta(r-R) \left(\frac{\partial^2}{\partial\theta\partial\varphi} Y_{l,m}^*(\theta, \varphi)\right) \Big|_{\theta=\frac{\pi}{2}, \varphi=\Omega t}. \quad (106)$$

### A. Perturbations for $l = 0$

For  $l = 0$ , substituting Eqs.(102) into Eq.(78), we have

$$\left(\iota_0 \frac{\partial}{\partial r} - \sigma_0\right) \psi_0 = -\frac{r}{2} e^{4\Lambda} E_E^{00}. \quad (107)$$

Using the standard method to solve the above equation, one can obtain

$$\begin{aligned} \psi_0 &= -\exp\left(\int^r \frac{\sigma_0(r)}{\iota_0(r)} dr\right) \int^r \left[\frac{r' e^{4\Lambda(r')}}{2\iota_0(r')} \exp\left(-\int^{r'} \frac{\sigma_0(r'')}{\iota_0(r'')} dr''\right) E_E^{00}(r')\right] dr' \\ &= -\exp\left(\int^r \frac{\sigma_0}{\iota_0} dr - \int^R \frac{\sigma_0}{\iota_0} dr\right) \frac{R e^{4\Lambda|_R}}{2\iota_0|_R} \bar{E}_E^{00} \Theta(r-R) \\ &= -\exp\left(\int_R^r \frac{\sigma_0}{\iota_0} dr\right) \frac{R e^{4\Lambda|_R}}{2\iota_0|_R} \bar{E}_E^{00} \Theta(r-R), \end{aligned} \quad (108)$$

where the relation  $Y_{00}^*(\theta, \varphi) = (2\sqrt{\pi})^{-1}$  has been used, and  $\bar{E}_E^{00}$  denote the evaluated coefficient given by,

$$\bar{E}_E^{00} = -4\sqrt{\pi}e^{-\Lambda|_R - \Phi|_R} \frac{\mu L \Omega}{R^2} = 4\sqrt{\pi}e^{-\Lambda|_R - \Phi|_R} \frac{\mu \Lambda'|_R \cdot \mathcal{E}}{R}. \quad (109)$$

From Eq.(70), we have

$$K_0 = 2\psi_0 = -4\sqrt{\pi} \frac{\Lambda'|_R \cdot \mu \mathcal{E}}{\iota_0|_R} \exp\left(\int_R^r \frac{\sigma_0}{\iota_0} dr + 3\Lambda|_R - \Phi|_R\right) \Theta(r - R). \quad (110)$$

Using Eq.(75), we can get  $\iota_0$ , and then solve Eq.(70) to obtain  $A_0$ ,

$$A_0 = 4\sqrt{\pi}e^{2\Phi} \frac{\Lambda'|_R \cdot \mu \mathcal{E}}{\iota_0|_R} \cdot \mathcal{A} \cdot \exp\left[-\int^R \frac{\sigma_0}{\iota_0} dr + 3\Lambda|_R - \Phi|_R\right] \Theta(r - R), \quad (111)$$

where

$$\mathcal{A} = \int r^{-1} (e^{2\Lambda} + 2r\Lambda' - r^2\Phi'^2 + r^2\Lambda'\Phi' - r^2\Phi'') \exp\left[\int^r \frac{\sigma_0(r')}{\iota_0(r')} dr' - 2\Lambda\right] dr. \quad (112)$$

With  $A_0$  and  $K_0$  in hand, the metric perturbation can be directly written out. Since both  $A_0$  and  $K_0$  contain step function, it is obvious that inside the orbit, i.e.  $r < R$ , the perturbation would vanishes. While outside the orbit, i.e.  $r > R$ , the perturbed metric can be given by

$$\begin{aligned} h_{ab}^{00} &= \frac{1}{2\sqrt{\pi}} (\Lambda_0 v_a v_b + K_0 n_a n_b) \\ &= 2e^{2\Phi} \frac{\Lambda'|_R \cdot \mu \mathcal{E}}{\iota_0|_R} \cdot \mathcal{A} \cdot \exp\left[-\int^R \frac{\sigma_0}{\iota_0} dr + 3\Lambda|_R - \Phi|_R\right] v_a v_b \\ &\quad - 2 \frac{\Lambda'|_R \cdot \mu \mathcal{E}}{\iota_0|_R} \exp\left[\int^R \frac{\sigma_0}{\iota_0} dr + 3\Lambda|_R - \Phi|_R\right] n_a n_b. \end{aligned} \quad (113)$$

Unlike the Schwarzschild spacetime, there is no clear definition of the mass  $M$  for general spherically symmetric spacetimes. It is obviously that the perturbation for  $l = 0$  only affects  $g_{tt}$  and  $g_{rr}$ . In the Schwarzschild spacetime, this perturbation is equivalent to adding an extra mass  $\delta m$  to the system [4].

## B. Perturbations for $l = 1$

### 1. Even-parity Perturbations

For the even-parity perturbations with  $l = 1$ , we find that metric perturbation components can be set directly to  $A_1 = E_1 = H_1 = 0$ . From Eq.(81), we find that  $K_1$  can be determined by  $E_A$ . Note that  $Y_{1m}^*(\frac{\pi}{2}, \varphi)$  would vanish for  $m = 0$ , hence

$$K_1 = -8\pi R \frac{\mu \mathcal{E}}{r^2} Y_{1,\pm 1}^*\left(\frac{\pi}{2}, \Omega t\right) \exp\left(-\int^r \frac{1 + e^{2\Lambda} - 4r\Lambda'}{r} dr + 3\Lambda|_R - \Phi|_R\right) \Theta(r - R). \quad (114)$$

Substituting  $K_1$  into Eq.(83), together with  $E_H = E_K = 0$ , we have

$$\frac{\partial}{\partial r} \left( \frac{\partial}{\partial t} B_1 \right) - r^{-1} (1 - e^{2\Lambda}) \frac{\partial}{\partial t} B_1 = \mathcal{K} K_1, \quad (115)$$

where

$$\mathcal{K} = \frac{1}{2} r^{-2} e^{2\Phi - 2\Lambda} (-2 + e^{2\Lambda} - 2r\Phi' - r^2\Phi'^2 + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi''). \quad (116)$$

Hence,  $B_1$  can be solved by Eq.(115). Then, considering Eq.(82) with  $E_D = 0$ , in principle, the metric perturbed component  $D_1$  can also be obtained.

## 2. Odd-parity Perturbations

To study the metric perturbation components  $C_1$  and  $J_1$ , we should first solve the master variable  $\alpha_1$  from Eq.(89). Then using Eqs.(91) and (92),  $C_1$  and  $J_1$  can be obtained. However,  $G$  automatically vanish for the  $l = 1$  case, there exists one freedom for the odd parity. If we impose the gauge  $C_1 = 0$ , the master variable becomes

$$\alpha_1 = \frac{\partial}{\partial t} J_1. \quad (117)$$

Eq.(91) becomes

$$\frac{\partial}{\partial r} \alpha_1 + \left( \frac{3}{r} - 2\Lambda' - 2\Phi' \right) \alpha_1 = e^{2\Lambda} E_C. \quad (118)$$

The solution of this equation is

$$\begin{aligned} \alpha_1 &= -\frac{16\pi}{l(l+1)} \cdot \frac{\mu \mathcal{E} \Omega R^2}{r^3} \exp(2\Lambda + 2\Phi - 2\Phi|_R) \cdot \left( \frac{\partial}{\partial \theta} Y_{lm}^*(\theta, \varphi) \right) \Big|_{\theta=\frac{\pi}{2}, \varphi=\Omega t} \\ &= -8\pi \frac{\mu L}{r^3} e^{2\Lambda+2\Phi} \left( \frac{\partial}{\partial \theta} Y_{lm}^*(\theta, \varphi) \right) \Big|_{\theta=\frac{\pi}{2}, \varphi=\Omega t}, \end{aligned} \quad (119)$$

where we have used the relation Eq.(100) and the fact  $l = 1$ . From  $\alpha_1$ , one can solve  $J_1$  with a undetermined function that only depend on  $r$ . However, the metric perturbation component  $J_1$  yields a contribution to  $h_{r\theta}^{1m}$  and  $h_{r\varphi}^{1m}$ .

Next, we impose the gauge  $J_1 = 0$ , and now the gauge invariant  $\alpha_1$  becomes

$$\alpha_1 = \frac{\partial}{\partial r} C_1 - \left( \frac{1}{r} + \Lambda' + \Phi' \right) C_1. \quad (120)$$

Substituting this relation into Eqs.(91) and (92), we know that  $\alpha_1$  should only be a function of  $r$ , and  $C_1$  satisfies the second order differential equation

$$\begin{aligned} \frac{\partial^2}{\partial r^2} C_1 + \left( \frac{2}{r} - 3\Lambda' - 3\Phi' \right) \frac{\partial}{\partial r} C_1 \\ - r^{-2} \left( 3 - e^{2\Lambda} - 2r^2 \Lambda'^2 + 6r\Phi' + r^2 \Phi'^2 - 2r\Lambda' - 7r^2 \Lambda' \Phi' + r^2 \Lambda'' + 4r^2 \Phi'' \right) C_1 = e^{2\Lambda} E_C. \end{aligned} \quad (121)$$

which is Eq.(87) with  $J_1 = 0$ . Considering that  $E_C$  contains the delta function, we predict that the solution of  $C_1$  contain the step function  $\Theta(r-R)$ , with two arbitrary constants of integration. One can be determined by considering that the background spacetime without intrinsic angular momentum, and the other can be determined by the fact that perturbation  $C_1$  would be convergent at infinity [28]. If the background metric degenerates to the Schwarzschild spacetime, our equation would become Eq. (10.21) of [28], which gives a non-vanishing  $h_{t\varphi}^{01}$  outside the circular orbit to describe the adding angular momentum of the system.

## C. Perturbations For $l \geq 2$

Generally speaking, for  $l \geq 2$  the problem is much more mathematically involved, and normally we have to seek help from numerical computations. From the source term, Eqs.(43), (54) or (62), we know that they depend only on Eqs.(101)-(106). All these terms are proportional to  $e^{-im\Omega t}$ , which implies that our master equations, Eqs.(42), (53) and (61), can be written as the second-order ordinary differential equations in  $r$  for each  $l$  and  $m$ . And the distributional sources of these differential equations would be vanish everywhere except at the circular orbit radius  $r = R$ .

An basic outline is as follows. To solve the second-order equations, one can study two regions separately. One region is from the circular orbit radius to infinity, i.e.  $r \in (R, \infty)$ . In this region, a naturally boundary condition is considering an appropriate radiation at spatial infinity. In the other region there are several different situations. For example, if the background spacetime has an event horizon, then one should determine this region from the event horizon to the circular orbit radius, i.e.,  $r \in (r_+, R)$ . Then solve the differential equation with the boundary condition given by a appropriate radiation at the event horizon  $r_+$ . Another example is that the background spacetime is a perfect fluid star without event horizons, then one should determine the other region from the center of the star to the circular orbit radius, i.e.,  $r \in (0, R)$ . Then solve the differential equation with some boundary condition at the center of the star. Finally, matching the solutions obtained in the two separate regions properly across the boundary  $r = R$ , we obtain the perturbations valid over the whole spacetime.



## V. CONCLUSIONS AND DISCUSSIONS

In this paper, we systematically study the gauge invariant perturbations of a general spherically symmetric background. First, we find that, in general spherically symmetric spacetimes, for even-parity, there are several gauge choices. One is the well-known RW gauge, and the other is the EZ gauge. For odd-parity perturbation, only one gauge choice exists, that is, by setting  $G = 0$ . Then, we mainly focus on the construction of master equations for  $l \geq 2$ . Under the EZ gauge, the master equation Eq.(42) for gauge invariants is a third order equation. However, after the separation of variables, the equation becomes a second order equation. Under the RW gauge, the master equation Eq.(53) for gauge invariants is a second order wave-like equation. For odd-parity perturbation, the master equation is also constructed as Eq.(61). Next, the cases for  $l = 0$  and  $l = 1$  are discussed. For the  $l = 0$  case, we present the equations that the gauge invariants satisfies. For even parity perturbations with  $l = 1$ , we find that the metric perturbation components are still determined when the source is specified. And for odd parity perturbations with  $l = 1$ , the master equation Eq.(89) for gauge invariants is still a wave-like equation. Finally, using our general results, we investigate a point particle moving along the circular orbit in general spherically symmetric spacetimes. In particular, the form of the solutions for  $l = 0$  and  $l = 1$  are carefully discussed.

### Acknowledgments

We would like very much to thank Xiaokai He for valuable discussions. This work was partially supported by the National Key Research and Development Program of China Grant No.2020YFC2201503, and the National Natural Science Foundation of China under Grants No. 11705053, No. 11975203 and No. 12035005. The work of Fang was supported in part by China Scholarship Council for the visiting post-doc program to Baylor University.

### Appendix A: Normalization Factors And A-K Decomposition

The normalization factors  $N^{(vec)}(A, r, l)$  and  $N^{(ten)}(A, r, l)$  in Sec.II can be calculated as

$$\begin{aligned} \oint Y_a^{E,lm}(Y_{E,l'm'}^a) d\Omega &= N^{(vec)}(E, r, l) \delta_{ll'} \delta_{mm'} = l(l+1) \delta_{ll'} \delta_{mm'}, \\ \oint Y_a^{B,lm}(Y_{B,l'm'}^a) d\Omega &= N^{(vec)}(B, r, l) \delta_{ll'} \delta_{mm'} = e^{-2\Lambda-2\Phi} l(l+1) \delta_{ll'} \delta_{mm'}, \\ \oint Y_a^{R,lm}(Y_{R,l'm'}^a) d\Omega &= N^{(vec)}(R, r, l) \delta_{ll'} \delta_{mm'} = e^{-2\Lambda} \delta_{ll'} \delta_{mm'}, \end{aligned}$$

and

$$\begin{aligned} \oint T_{ab}^{T0,lm}(T_{T0,l'm'}^{ab})^* d\Omega &= N^{(ten)}(T0, r, l) \delta_{ll'} \delta_{mm'} = 2 \delta_{ll'} \delta_{mm'}, \\ \oint T_{ab}^{L0,lm}(T_{L0,l'm'}^{ab})^* d\Omega &= N^{(ten)}(L0, r, l) \delta_{ll'} \delta_{mm'} = e^{-4\Lambda} \delta_{ll'} \delta_{mm'}, \\ \oint T_{ab}^{E1,lm}(T_{E1,l'm'}^{ab})^* d\Omega &= N^{(ten)}(E1, r, l) \delta_{ll'} \delta_{mm'} = \frac{1}{2} e^{-2\Lambda} l(l+1) \delta_{ll'} \delta_{mm'}, \\ \oint T_{ab}^{B1,lm}(T_{B1,l'm'}^{ab})^* d\Omega &= N^{(ten)}(B1, r, l) \delta_{ll'} \delta_{mm'} = \frac{1}{2} e^{-4\Lambda-2\Phi} l(l+1) \delta_{ll'} \delta_{mm'}, \\ \oint T_{ab}^{E2,lm}(T_{E2,l'm'}^{ab})^* d\Omega &= N^{(ten)}(E2, r, l) \delta_{ll'} \delta_{mm'} = \frac{(l+2)!}{2(l-2)!} \delta_{ll'} \delta_{mm'}, \\ \oint T_{ab}^{B2,lm}(T_{B2,l'm'}^{ab})^* d\Omega &= N^{(ten)}(B2, r, l) \delta_{ll'} \delta_{mm'} = e^{-2\Lambda-2\Phi} \frac{(l+2)!}{2(l-2)!} \delta_{ll'} \delta_{mm'}. \end{aligned}$$

Projecting the perturbed metric  $h_{ab}$  onto these basis, one can obtain the expressions of coefficients A-K:

$$\begin{aligned}
A &= e^{4\Phi} \oint h_{ab}^{lm} (v^a v^b Y_{lm}^*) d\Omega, & B &= -\frac{e^{2\Phi}}{l(l+1)} \oint h_{ab}^{lm} v^a Y_{E,lm}^{b*} d\Omega, \\
C &= -\frac{e^{4\Phi+2\Lambda}}{l(l+1)} \oint h_{ab}^{lm} v^a Y_{B,lm}^{b*} d\Omega, & D &= -e^{2\Phi+2\Lambda} \oint h_{ab}^{lm} v^a Y_{R,lm}^{b*} d\Omega, \\
E &= \frac{1}{2} \oint h_{ab}^{lm} T_{T0,lm}^{ab*} d\Omega, & F &= \frac{2(l-2)!}{(l+2)!} \oint h_{ab}^{lm} T_{E2,lm}^{ab*} d\Omega, \\
G &= \frac{2(l-2)!}{(l+2)!} e^{2\Phi+2\Lambda} \oint h_{ab}^{lm} T_{B2,lm}^{ab*} d\Omega, & H &= \frac{e^{2\Lambda}}{l(l+1)} \oint h_{ab}^{lm} T_{E1,lm}^{ab*} d\Omega, \\
J &= \frac{e^{4\Lambda+2\Phi}}{l(l+1)} \oint h_{ab}^{lm} T_{B1,lm}^{ab*} d\Omega, & K &= e^{4\Lambda} \oint h_{ab}^{lm} T_{L0,lm}^{ab*} d\Omega.
\end{aligned}$$

The above results are only valid for  $l \geq 2$ .

## Appendix B: Decomposition of Linearized Einstein Equations

In this appendix, we show the decomposition of linearized Einstein tensor into the A-K components. Note that these expressions are only valid for  $l \geq 2$ .

### 1. Even parity

$$\begin{aligned}
E_A &= 2r^{-2} e^{-2\Lambda} (-1 + e^{2\Lambda} + 2r\Lambda' - 2r\Phi' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'') A - r^{-2} e^{2\Phi} (l^2 + l - 2) E \\
&\quad - 2r^{-1} e^{-2\Lambda+2\Phi} (r\Lambda' - 3) \frac{\partial}{\partial r} E + 2e^{-2\Lambda+2\Phi} \frac{\partial^2}{\partial r^2} E - \frac{1}{2} r^{-2} e^{2\Phi} l(l+2)(l+1)(l-1) F \\
&\quad - 2r^{-2} e^{-2\Lambda+2\Phi} l(l+1)(r\Lambda' - 2) H + 2r^{-1} e^{-2\Lambda+2\Phi} l(l+1) \frac{\partial}{\partial r} H \\
&\quad - r^{-2} e^{-4\Lambda+2\Phi} [2 + e^{2\Lambda} l(l+1) - 8r\Lambda'] K - 2r^{-1} e^{-4\Lambda+2\Phi} \frac{\partial}{\partial r} K,
\end{aligned} \tag{B1}$$

$$\begin{aligned}
E_B &= r^{-2} e^{-2\Lambda} (-3 + 3e^{2\Lambda} + 4r\Lambda' - 4r\Phi' + 3r^2\Lambda'\Phi' - 3r^2\Phi'^2 - 3r^2\Phi'') B \\
&\quad - r^{-1} e^{-2\Lambda} (-2 + r\Lambda' + r\Phi') \frac{\partial}{\partial r} B + e^{-2\Lambda} \frac{\partial^2}{\partial r^2} B + r^{-1} e^{-2\Lambda} (\Lambda' - \Phi') D - r^{-1} e^{-2\Lambda} \frac{\partial}{\partial r} D \\
&\quad - r^{-1} \frac{\partial}{\partial t} E - \frac{1}{2} r^{-1} (l^2 + l - 2) \frac{\partial}{\partial t} F - r^{-1} e^{-2\Lambda} (-3 + r\Lambda' + r\Phi') \frac{\partial}{\partial t} H + e^{-2\Lambda} \frac{\partial^2}{\partial t \partial r} H \\
&\quad - r^{-1} e^{-2\Lambda} \frac{\partial}{\partial t} K,
\end{aligned} \tag{B2}$$

$$\begin{aligned}
E_D &= r^{-2} l(l+1)(1 - 2r\Phi') B + r^{-1} l(l+1) \frac{\partial}{\partial r} B + 2r^{-1} (-1 + r\Phi') \frac{\partial}{\partial t} E - 2 \frac{\partial}{\partial t \partial r} E \\
&\quad - r^{-2} e^{-2\Lambda} (2 - 2e^{2\Lambda} + e^{2\Lambda} l + e^{2\Lambda} l^2 + 2r\Phi' - 6r\Lambda' + 2r^2\Phi'^2 - 2r^2\Lambda'\Phi' + 2r^2\Phi'') D \\
&\quad - r^{-1} l(l+1) \frac{\partial}{\partial t} H + 2r^{-1} e^{-2\Lambda} \frac{\partial}{\partial t} K,
\end{aligned} \tag{B3}$$

$$\begin{aligned}
E_E = & -\frac{1}{2}r^{-2}e^{-2\Lambda-2\Phi}(e^{2\Lambda}l + e^{2\Lambda}l^2 + 4r\Phi' - 4r^2\Lambda'\Phi' + 4r^2\Phi'')A - r^{-1}e^{-2\Lambda-2\Phi}(-1 + r\Lambda' + 2r\Phi')\frac{\partial}{\partial r}A \\
& + e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial r^2}A - r^{-1}e^{-2\Phi}l(l+1)\frac{\partial}{\partial t}B - 2r^{-1}e^{-2\Lambda-2\Phi}(-1 + r\Lambda')\frac{\partial}{\partial t}D + 2e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial t\partial r}D \\
& + 2r^{-2}e^{-2\Lambda}(-1 + e^{2\Lambda} - 2r\Phi' + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')E + r^{-1}e^{-2\Lambda}(-2 + r\Lambda' - r\Phi')\frac{\partial}{\partial r}E \\
& - e^{-2\Lambda}\frac{\partial^2}{\partial r^2}E + e^{-2\Phi}\frac{\partial^2}{\partial t^2}E + r^{-2}e^{-2\Lambda}l(l+1)(-1 + r\Lambda' - r\Phi')H - r^{-1}e^{-2\Lambda}l(l+1)\frac{\partial}{\partial r}H \\
& + \frac{1}{2}r^{-2}e^{-4\Lambda}(e^{2\Lambda}l + e^{2\Lambda}l^2 + 4r\Phi' - 8r\Lambda' + 4r^2\Phi'^2 - 8r^2\Lambda'\Phi' + 4r^2\Phi'')K + r^{-1}e^{-4\Lambda}(1 + r\Phi')\frac{\partial}{\partial r}K \\
& + e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial t^2}K,
\end{aligned} \tag{B4}$$

$$\begin{aligned}
E_F = & -r^{-2}e^{-2\Phi}A - 2r^{-1}e^{-2\Phi}\frac{\partial}{\partial t}B + 2r^{-2}e^{-2\Lambda}(-1 + e^{2\Lambda} + 2r\Lambda' - 2r\Phi' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')F \\
& - r^{-1}e^{-2\Lambda}(-2 + r\Lambda' - r\Phi')\frac{\partial}{\partial r}F + e^{-2\Lambda}\frac{\partial^2}{\partial r^2}F - e^{-2\Phi}\frac{\partial^2}{\partial t^2}F + 2r^{-2}e^{-2\Lambda}(-1 + r\Lambda' - r\Phi')H \\
& - 2r^{-1}e^{-2\Lambda}\frac{\partial}{\partial r}H + r^{-2}e^{-2\Lambda}K,
\end{aligned} \tag{B5}$$

$$\begin{aligned}
E_H = & r^{-2}e^{-2\Phi}(1 + r\Phi')A - r^{-1}e^{-2\Phi}\frac{\partial}{\partial r}A + r^{-1}e^{-2\Phi}\frac{\partial}{\partial t}B - e^{-2\Phi}\frac{\partial^2}{\partial t\partial r}B - r^{-1}e^{-2\Phi}\frac{\partial}{\partial t}D + r^{-1}\frac{\partial}{\partial r}E \\
& + \frac{1}{2}r^{-1}(l+2)(l-1)\frac{\partial}{\partial r}F + r^{-2}e^{-2\Lambda}(-1 + 3e^{2\Lambda} + 5r\Lambda' - 3r\Phi' + 3r^2\Lambda'\Phi' - 3r^2\Phi'^2 - 3r^2\Phi'')H \\
& - e^{-2\Phi}\frac{\partial^2}{\partial t^2}H - r^{-2}e^{-2\Lambda}(1 + r\Phi')K,
\end{aligned} \tag{B6}$$

$$\begin{aligned}
E_K = & -r^{-2}e^{-2\Phi}(e^{2\Lambda}l + e^{2\Lambda}l^2 + 4r\Phi')A + 2r^{-1}e^{-2\Phi}\frac{\partial}{\partial r}A - 2r^{-1}e^{2\Lambda-2\Phi}l(l+1)\frac{\partial}{\partial t}B + 4r^{-1}e^{-2\Phi}\frac{\partial}{\partial t}D \\
& + r^{-2}e^{2\Lambda}(l^2 + l - 2)E - 2r^{-1}(1 + r\Phi')\frac{\partial}{\partial r}E + 2e^{2\Lambda-2\Phi}\frac{\partial^2}{\partial t^2}E + \frac{1}{2}r^{-2}e^{2\Lambda}l(l+2)(l+1)(l-1)F \\
& - 2r^{-2}l(l+1)(1 + r\Phi')H + 2r^{-2}e^{-2\Lambda}(e^{2\Lambda} + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'')K.
\end{aligned} \tag{B7}$$

## 2. odd parity

$$\begin{aligned}
E_C = & -r^{-2}e^{-2\Lambda}\left(3 - 3e^{2\Lambda} + e^{2\Lambda}l + e^{2\Lambda}l^2 - 2r^2\Lambda'^2 + 6r\Phi' + r^2\Phi'^2 - 2r\Lambda' - 7r^2\Lambda'\Phi' + r^2\Lambda'' + 4r^2\Phi''\right)C \\
& - r^{-1}e^{-2\Lambda}(-2 + 3r\Lambda' + 3r\Phi')\frac{\partial}{\partial r}C + e^{-2\Lambda}\frac{\partial^2}{\partial r^2}C - \frac{1}{2}r^{-1}(l+2)(l-1)\frac{\partial}{\partial t}G \\
& - r^{-1}e^{-2\Lambda}(-3 + 2r\Lambda' + 2r\Phi')\frac{\partial}{\partial t}J + e^{-2\Lambda}\frac{\partial^2}{\partial t\partial r}J,
\end{aligned} \tag{B8}$$

$$\begin{aligned}
E_G = & -2r^{-1}e^{-2\Phi}\frac{\partial}{\partial t}C + r^{-2}e^{-2\Lambda}(-2 + 2e^{2\Lambda} + 2r^2\Lambda'^2 - 6r\Phi' - 2r^2\Phi'^2 + 2r\Lambda' + 4r^2\Lambda'\Phi' - r^2\Lambda'' - 3r^2\Phi'')G \\
& - r^{-1}e^{-2\Lambda}(-2 + 3r\Lambda' + r\Phi')\frac{\partial}{\partial r}G + e^{-2\Lambda}\frac{\partial^2}{\partial r^2}G - e^{-2\Phi}\frac{\partial^2}{\partial t^2}G + 2r^{-2}e^{-2\Lambda}(-1 + 2r\Lambda')J - 2r^{-1}e^{-2\Lambda}\frac{\partial}{\partial r}J,
\end{aligned} \tag{B9}$$

$$\begin{aligned}
E_J = & r^{-1}e^{-2\Phi}(1+r\Lambda'+r\Phi')\frac{\partial}{\partial t}C - e^{-2\Phi}\frac{\partial^2}{\partial t\partial r}C + \frac{1}{2}r^{-1}(l+2)(l-1)(\Lambda'+\Phi')G + \frac{1}{2}r^{-1}(l+2)(l-1)\frac{\partial}{\partial r}G \\
& - r^{-2}e^{-2\Lambda}(1-3e^{2\Lambda}+e^{2\Lambda}l+e^{2\Lambda}l^2-5r\Lambda'+3r\Phi'+3r^2\Phi'^2+3r^2\Phi''-3r^2\Lambda'\Phi')J - e^{-2\Phi}\frac{\partial^2}{\partial t^2}J. \tag{B10}
\end{aligned}$$

### Appendix C: Linearized Einstein Tensor In Terms of Gauge Invariants

In this appendix, we rewrite the linearized Einstein tensor in terms of the gauge invariants under the EZ gauge or the RW gauge. Note that for even-parity perturbations, we use the superscripts to distinguish expressions under the EZ gauge and RW gauge, respectively. These relations are only valid for  $l \geq 2$  case. The relation  $\lambda = l(l+1)$  also has been used in this appendix.

#### 1. Even Parity and the EZ Gauge

$$\begin{aligned}
E_A^I = & -4r^{-2}e^{-2\Lambda}(-1+e^{2\Lambda}+2r\Lambda'-2r\Phi'+r^2\Lambda'\Phi'-r^2\Phi'^2-r^2\Phi'')\epsilon^I - 2r^{-2}e^{-2\Lambda+2\Phi}\lambda(r\Lambda'-2)\chi^I \\
& + 2r^{-1}e^{-2\Lambda+2\Phi}\lambda\frac{\partial}{\partial r}\chi^I - 2r^{-2}e^{-4\Lambda+2\Phi}[2+e^{2\Lambda}\lambda-8r\Lambda']\psi^I - 4r^{-1}e^{-4\Lambda+2\Phi}\frac{\partial}{\partial r}\psi^I, \tag{C1}
\end{aligned}$$

$$E_B^I = r^{-1}e^{-2\Lambda}(\Lambda'-\Phi')\delta^I - r^{-1}e^{-2\Lambda}\frac{\partial}{\partial r}\delta^I - r^{-1}e^{-2\Lambda}(-3+r\Lambda'+r\Phi')\frac{\partial}{\partial t}\chi^I + e^{-2\Lambda}\frac{\partial^2}{\partial t\partial r}\chi^I - 2r^{-1}e^{-2\Lambda}\frac{\partial}{\partial t}\psi^I, \tag{C2}$$

$$\begin{aligned}
E_D^I = & -r^{-2}e^{-2\Lambda}\left[2+e^{2\Lambda}(\lambda-2)+2r\Phi'-6r\Lambda'+2r^2\Phi'^2-2r^2\Lambda'\Phi'+2r^2\Phi''\right]\delta^I \\
& - r^{-1}\lambda\frac{\partial}{\partial t}\chi^I + 4r^{-1}e^{-2\Lambda}\frac{\partial}{\partial t}\psi^I, \tag{C3}
\end{aligned}$$

$$\begin{aligned}
E_E^I = & r^{-2}e^{-2\Lambda-2\Phi}\left(e^{2\Lambda}\lambda+4r\Phi'-4r^2\Lambda'\Phi'+4r^2\Phi''\right)\epsilon^I + 2r^{-1}e^{-2\Lambda-2\Phi}(-1+r\Lambda'+2r\Phi')\frac{\partial}{\partial r}\epsilon^I \\
& - 2e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial r^2}\epsilon^I - 2r^{-1}e^{-2\Lambda-2\Phi}(-1+r\Lambda')\frac{\partial}{\partial t}\delta^I + 2e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial t\partial r}\delta^I + r^{-2}e^{-2\Lambda}\lambda(-1+r\Lambda'-r\Phi')\chi^I \\
& - r^{-1}e^{-2\Lambda}\lambda\frac{\partial}{\partial r}\chi^I + r^{-2}e^{-4\Lambda}\left(e^{2\Lambda}\lambda+4r\Phi'-8r\Lambda'+4r^2\Phi'^2-8r^2\Lambda'\Phi'+4r^2\Phi''\right)\psi^I \\
& + 2r^{-1}e^{-4\Lambda}(1+r\Phi')\frac{\partial}{\partial r}\psi^I + 2e^{-2\Lambda-2\Phi}\frac{\partial^2}{\partial t^2}\psi^I, \tag{C4}
\end{aligned}$$

$$E_F^I = 2r^{-2}e^{-2\Phi}\epsilon^I + 2r^{-2}e^{-2\Lambda}(-1+r\Lambda'-r\Phi')\chi^I - 2r^{-1}e^{-2\Lambda}\frac{\partial}{\partial r}\chi^I + 2r^{-2}e^{-2\Lambda}\psi^I, \tag{C5}$$

$$\begin{aligned}
E_H^I = & -2r^{-2}e^{-2\Phi}(1+r\Phi')\epsilon^I + 2r^{-1}e^{-2\Phi}\frac{\partial}{\partial r}\epsilon^I - r^{-1}e^{-2\Phi}\frac{\partial}{\partial t}\delta^I - 2r^{-2}e^{-2\Lambda}(1+r\Phi')\psi^I \\
& + r^{-2}e^{-2\Lambda}(-1+3e^{2\Lambda}+5r\Lambda'-3r\Phi'+3r^2\Lambda'\Phi'-3r^2\Phi'^2-3r^2\Phi'')\chi^I - e^{-2\Phi}\frac{\partial^2}{\partial t^2}\chi^I, \tag{C6}
\end{aligned}$$

$$\begin{aligned}
E_K^I = & 2r^{-2}e^{-2\Phi}\left(e^{2\Lambda}\lambda+4r\Phi'\right)\epsilon^I - 4r^{-1}e^{-2\Phi}\frac{\partial}{\partial r}\epsilon^I + 4r^{-1}e^{-2\Phi}\frac{\partial}{\partial t}\delta^I \\
& - 2r^{-2}\lambda(1+r\Phi')\chi^I + 4r^{-2}e^{-2\Lambda}(e^{2\Lambda}+2r\Lambda'+r^2\Lambda'\Phi'-r^2\Phi'^2-r^2\Phi'')\psi^I. \tag{C7}
\end{aligned}$$

## 2. Even Parity and the RW Gauge

$$\begin{aligned}
E_A^{\text{II}} = & -4r^{-2}e^{-2\Lambda} \left( -1 + e^{2\Lambda} + 2r\Lambda' - 2r\Phi' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'' \right) \epsilon^{\text{II}} \\
& + r^{-2}e^{-4\Lambda+2\Phi} \left[ 4 - 4e^{2\Lambda} + 4e^{2\Lambda}\lambda - 8r^2\Lambda'^2 - 8r^2\Phi'^2 - 4r^3\Phi'^3 \right. \\
& \left. + 2r\Lambda' \left( -e^{2\Lambda}\lambda + 4r\Phi' + 2r^2\Phi'^2 \right) + 4r^2\Lambda'' - 4r\Phi'(1 - e^{2\Lambda} + 2r^2\Phi'') \right] \chi^{\text{II}} \\
& + 2r^{-1}e^{-2\Lambda+2\Phi} \lambda \frac{\partial}{\partial r} \chi - 2r^{-2}e^{-4\Lambda+2\Phi} (2 + e^{2\Lambda}\lambda - 8r\Lambda') \psi^{\text{II}} - 4r^{-1}e^{-4\Lambda+2\Phi} \frac{\partial}{\partial r} \psi^{\text{II}}, \tag{C8}
\end{aligned}$$

$$\begin{aligned}
E_B^{\text{II}} = & r^{-1}e^{-2\Lambda}(\Lambda' - \Phi')\delta^{\text{II}} - r^{-1}e^{-2\Lambda} \frac{\partial}{\partial r} \delta^{\text{II}} - r^{-1}e^{-2\Lambda}(-3 + r\Lambda' + r\Phi') \frac{\partial}{\partial t} \chi^{\text{II}} + e^{-2\Lambda} \frac{\partial^2}{\partial t \partial r} \chi^{\text{II}} - 2r^{-1}e^{-2\Lambda} \frac{\partial}{\partial t} \psi^{\text{II}}, \tag{C9}
\end{aligned}$$

$$\begin{aligned}
E_D^{\text{II}} = & -r^{-2}e^{-2\Lambda} \left( 2 - 2e^{2\Lambda} + e^{2\Lambda}\lambda + 2r\Phi' - 6r\Lambda' + 2r^2\Phi'^2 - 2r^2\Lambda'\Phi' + 2r^2\Phi'' \right) \delta^{\text{II}} \\
& - r^{-1}e^{-2\Lambda} \left( 2 - 2e^{2\Lambda} + e^{2\Lambda}\lambda + 6r\Phi' + 2r^2\Phi'^2 - 2r\Lambda' - 2r^2\Lambda'\Phi' + 2r^2\Phi'' \right) \frac{\partial}{\partial t} \chi^{\text{II}} + 4r^{-1}e^{-2\Lambda} \frac{\partial}{\partial t} \psi^{\text{II}}, \tag{C10}
\end{aligned}$$

$$\begin{aligned}
E_E^{\text{II}} = & r^{-2}e^{-2\Lambda-2\Phi} \left( e^{2\Lambda}\lambda + 4r\Phi' - 4r^2\Lambda'\Phi' + 4r^2\Phi'' \right) \epsilon^{\text{II}} + 2r^{-1}e^{-2\Lambda-2\Phi}(-1 + r\Lambda' + 2r\Phi') \frac{\partial}{\partial r} \epsilon^{\text{II}} \\
& - 2e^{-2\Lambda-2\Phi} \frac{\partial^2}{\partial r^2} \epsilon^{\text{II}} - 2r^{-1}e^{-2\Lambda-2\Phi}(-1 + r\Lambda') \frac{\partial}{\partial t} \delta^{\text{II}} + 2e^{-2\Lambda-2\Phi} \frac{\partial^2}{\partial t \partial r} \delta^{\text{II}} \\
& + r^{-2}e^{-4\Lambda} \left[ 4 - 4e^{2\Lambda} - e^{2\Lambda}\lambda + 4r^2\Phi'^2 + 4r^2\Lambda^2(1 + r\Phi') - 2r^2\Lambda'' + 6r^2\Phi'' \right. \\
& \left. - r\Phi'(-6 + e^{2\Lambda}\lambda + 2r^2\Lambda'' - 4r^2\Lambda'') - r\Lambda'(6 - e^{2\Lambda}\lambda + 8r\Phi' + 4r^2\Phi'^2 + 6r^2\Phi'') + 2r^3\Phi''' \right] \chi^{\text{II}} \\
& - r^{-1}e^{-2\Lambda} \lambda \frac{\partial}{\partial r} \chi^{\text{II}} + r^{-2}e^{-4\Lambda} \left( e^{2\Lambda}\lambda + 4r\Phi' - 8r\Lambda' + 4r^2\Phi'^2 - 8r^2\Lambda'\Phi' + 4r^2\Phi'' \right) \psi^{\text{II}} \\
& + 2r^{-1}e^{-4\Lambda}(1 + r\Phi') \frac{\partial}{\partial r} \psi^{\text{II}} + 2e^{-2\Lambda-2\Phi} \frac{\partial^2}{\partial t^2} \psi^{\text{II}}, \tag{C11}
\end{aligned}$$

$$\begin{aligned}
E_F^{\text{II}} = & 2r^{-2}e^{-2\Phi} \epsilon^{\text{II}} + 2r^{-2}e^{-2\Lambda}(-1 + r\Lambda' - r\Phi') \chi^{\text{II}} - 2r^{-1}e^{-2\Lambda} \frac{\partial}{\partial r} \chi^{\text{II}} + 2r^{-2}e^{-2\Lambda} \psi^{\text{II}}, \tag{C12}
\end{aligned}$$

$$\begin{aligned}
E_H^{\text{II}} = & -2r^{-2}e^{-2\Phi}(1 + r\Phi')\epsilon^{\text{II}} + 2r^{-1}e^{-2\Phi} \frac{\partial}{\partial r} \epsilon^{\text{II}} - r^{-1}e^{-2\Phi} \frac{\partial}{\partial t} \delta^{\text{II}} - 2r^{-2}e^{-2\Lambda}(1 + r\Phi')\psi^{\text{II}} \\
& + 2r^{-2}e^{-2\Lambda} \left( 1 + r\Phi' + r\Lambda' - r^2\Phi'^2 + r^2\Lambda'\Phi' - r^2\Phi'' \right) \chi^{\text{II}} - e^{-2\Phi} \frac{\partial^2}{\partial t^2} \chi^{\text{II}}, \tag{C13}
\end{aligned}$$

$$\begin{aligned}
E_K^{\text{II}} = & 2r^{-2}e^{-2\Phi}(e^{2\Lambda}\lambda + 4r\Phi')\epsilon^{\text{II}} - 4r^{-1}e^{-2\Phi} \frac{\partial}{\partial r} \epsilon^{\text{II}} + 4r^{-1}e^{-2\Phi} \frac{\partial}{\partial t} \delta^{\text{II}} \\
& - 2r^{-2}e^{-2\Lambda} \left[ e^{2\Lambda}\lambda + r\Phi'(e^{2\Lambda}\lambda - 2) - 2r^2\Phi'^2 - 2r^2\Lambda'^2(r\Phi' + 2) \right. \\
& \left. - 4r^2\Phi'' + 2r\Lambda'(4 - e^{2\Lambda} + 5r\Phi' + r^2\Phi'^2 + r^2\Phi'') \right] \chi^{\text{II}} \\
& - 4r^{-1}e^{-2\Lambda} \left( -1 + e^{2\Lambda} - 2r\Phi' + 2r\Lambda' - r^2\Phi'^2 + r^2\Lambda'\Phi' - r^2\Phi'' \right) \frac{\partial}{\partial r} \chi^{\text{II}} \\
& + 4r^{-2}e^{-2\Lambda} \left( e^{2\Lambda} + 2r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi'^2 - r^2\Phi'' \right) \psi^{\text{II}}. \tag{C14}
\end{aligned}$$

### 3. Odd Parity

$$E_C = r^{-2} e^{-2\Lambda} \left[ 3 + e^{2\Lambda} (\lambda - 3) - 2r^2 \Lambda'^2 + 6r\Phi' + r^2 \Phi'^2 - 2r\Lambda' - 7r^2 \Lambda' \Phi' + r^2 \Lambda'' + 4r^2 \Phi'' \right] \beta \\ + r^{-1} e^{-2\Lambda} (-2 + 3r\Lambda' + 3r\Phi') \frac{\partial}{\partial r} \beta - e^{-2\Lambda} \frac{\partial^2}{\partial r^2} \beta - r^{-1} e^{-2\Lambda} (-3 + 2r\Lambda' + 2r\Phi') \frac{\partial}{\partial t} \alpha + e^{-2\Lambda} \frac{\partial^2}{\partial t \partial r} \alpha, \quad (C15)$$

$$E_G = 2r^{-1} e^{-2\Phi} \frac{\partial}{\partial t} \beta + 2r^{-2} e^{-2\Lambda} (-1 + 2r\Lambda') \alpha - 2r^{-1} e^{-2\Lambda} \frac{\partial}{\partial r} \alpha, \quad (C16)$$

$$E_J = -r^{-2} e^{-2\Lambda} \left[ 1 + e^{2\Lambda} (\lambda - 3) - 5r\Lambda' + 3r\Phi' + 3r^2 \Phi'^2 + 3r^2 \Phi'' - 3r^2 \Lambda' \Phi' \right] \alpha \\ - e^{-2\Phi} \frac{\partial^2}{\partial t^2} \alpha - r^{-1} e^{-2\Phi} (1 + r\Lambda' + r\Phi') \frac{\partial}{\partial t} \beta + e^{-2\Phi} \frac{\partial^2}{\partial t \partial r} \beta. \quad (C17)$$

#### Appendix D: The specific structure of the parameters

##### 1. Even-parity and the EZ gauge for $l \geq 2$

The parameters  $\tau^I, \sigma^I, \kappa^I, \gamma^I, \rho^I, \mu^I, \nu^I$  in Eqs.(39) and (40) have the following form:

$$\tau^I = -2 + e^{2\Lambda} \lambda + 2r\Phi', \quad (D1)$$

$$\sigma^I = 1 - e^{2\Lambda} + e^{2\Lambda} \lambda + r\Phi' - 3r\Lambda' + r^2 \Phi'^2 - r^2 \Lambda' \Phi' + r^2 \Phi'', \quad (D2)$$

$$\kappa^I = 2\sigma^I - e^{2\Lambda} \lambda + 2r\Phi' + 2r\Lambda', \quad (D3)$$

$$\gamma^I = 3 - 3e^{2\Lambda} + 3r\Phi' + r^2 \Phi'^2 - r\Lambda' (7 - e^{2\Lambda} \lambda + r\Phi') + 3r^2 \Phi'', \quad (D4)$$

$$\rho^I = -2e^{2\Lambda} + e^{2\Lambda} \lambda + 4r\Phi' + 2r^2 \Phi'^2 - 2r\Lambda' (2 + r\Phi') + 2r^2 \Phi'', \quad (D5)$$

$$\mu^I = 2 + e^{2\Lambda} (\lambda - 8) + e^{4\Lambda} (6 - 5\lambda + 2\lambda^2) - r [14 + e^{2\Lambda} (9\lambda - 22)] \Lambda' - r [-10 + e^{2\Lambda} (18 - 13\lambda) + 2e^{4\Lambda} \lambda] \Phi' \\ + 20r^2 \Lambda'^2 - r^2 [40 + 3e^{2\Lambda} (3\lambda - 4)] \Lambda' \Phi' + r^2 [20 + e^{2\Lambda} (7\lambda - 12)] \Phi'^2 + r^2 [8 + e^{2\Lambda} (5\lambda - 12)] \Phi'' \\ + 2r^3 [11\Lambda' - (9 + e^{2\Lambda} \lambda) \Phi'] \left( \Lambda' \Phi' - \Phi'^2 - \Phi'' \right) + 6r^4 \left( \Phi'^2 + \Phi'' - \Lambda' \Phi' \right)^2, \quad (D6)$$

$$\nu^I = -4 + 4e^{2\Lambda} + e^{4\Lambda} (\lambda - 2) - 2r [-8 + e^{2\Lambda} (\lambda + 4)] \Lambda' - r [4 + e^{2\Lambda} (\lambda - 6)] \Phi' \\ - r^2 [8\Lambda'^2 - 8\Lambda' \Phi' + 8\Phi'^2 + 4\Phi'' - e^{2\Lambda} (\lambda - 4) \left( \Lambda' \Phi' - \Phi'^2 - \Phi'' \right)] \\ - 2r^3 (4\Lambda' - 3\Phi') \left( \Lambda' \Phi' - \Phi'^2 - \Phi'' \right) - 2r^4 \left( \Phi'^2 + \Phi'' - \Lambda' \Phi' \right)^2. \quad (D7)$$

The parameters in the master equation Eq.(42) and the source term in Eq.(43) are functions of background  $\Psi(r)$  and  $\Lambda(r)$ . To simplify the expression of  $N_R$  and  $N_Z$ , we first introduce two auxiliary parameters  $M^I$  and  $M^{II}$ :

$$M_1^I = 2\nu^I + \rho^I \sigma^I, \quad (D8)$$

$$M_2^I = -\mu^I + \gamma^I \sigma^I + r\tau^I (\sigma^I)', \quad (D9)$$

then we find the parameters in Eq.(42) are given by

$$N^I = 2\mu^I - 2\gamma^I \sigma^I - 2\nu^I \sigma^I - \rho^I (\sigma^I)^2 - 2r\tau^I (\sigma^I)', \quad (D10)$$

$$N_T^I = (N^I)^2 (e^{2\Lambda} \lambda - 2\sigma^I) + 2r (e^{2\Lambda} \lambda - 2\sigma^I) (\sigma^I - \kappa^I) \tau^I (N^I)' \\ + N^I \left\{ 2e^{2\Lambda} \lambda \gamma^I \sigma^I - 4\gamma^I (\sigma^I)^2 + e^{2\Lambda} \lambda \rho^I (\sigma^I)^2 - 2\rho^I (\sigma^I)^3 - 4e^{2\Lambda} \lambda \sigma^I \tau^I + 8(\sigma^I)^2 \tau^I + 2e^{2\Lambda} r \lambda \tau^I (\kappa^I)' \right. \\ \left. - 4r\sigma^I \tau^I (\kappa^I)' + 8r (\sigma^I)^2 \tau^I \Lambda' - 2e^{2\Lambda} r \lambda \tau^I (\sigma^I)' + 4e^{2\Lambda} r \lambda \sigma^I \tau^I \Phi' - 8r (\sigma^I)^2 \tau^I \Phi' \right. \\ \left. + \kappa^I \left[ -\rho^I (e^{2\Lambda} \lambda - 2\sigma^I) \sigma^I + \gamma^I (-2e^{2\Lambda} \lambda^I + 4\sigma^I) - 4\tau^I \left( -r (\sigma^I)' + 2\sigma^I (1 + r\Lambda' - r\Phi') + e^{2\Lambda} \lambda (-1 + r\Phi') \right) \right] \right\}, \quad (D11)$$

$$N_R^I = 2M_2^I N^I + (N^I)^2 + 2r\sigma^I \tau^I (N^I)' + N^I \sigma^I \left[ 2\gamma^I + \rho^I \sigma^I - 2(\tau^I + r(\tau^I)') \right], \quad (D12)$$

$$N_Z^I = 2rN^I \tau^I (M_2^I)' - M_2^I \left( N^I \rho^I \sigma^I + 2r\tau^I (N^I)' \right) + M_1^I N^I \left( \gamma^I \sigma^I + r\tau^I (\sigma^I)' \right), \quad (D13)$$

and the parameters in Eq.(43) are given by

$$N_A^I = -2r\tau^I (N^I)' + N^I \left( -2\gamma^I - \rho^I \sigma^I + 4\tau^I + 8r\tau^I \Lambda' + 2r(\tau^I)' - 4r\tau^I \Phi' \right), \quad (D14)$$

$$N_D^I = -N_T^I + 2\tau^I N^I (e^{2\Lambda} \lambda - 2\sigma^I) (\sigma^I - \kappa^I) (1 + 2r\Lambda'), \quad (D15)$$

$$N_F^I = (N^I)^2 - 2r(e^{2\Lambda} \lambda - \sigma^I) \tau^I (N^I)' + N^I \left[ \rho^I \sigma^I (-e^{2\Lambda} \lambda + \sigma^I) + \gamma^I (-2e^{2\Lambda} \lambda + 2\sigma^I) \right. \\ \left. + 4e^{2\Lambda} \lambda \tau^I - 4\sigma^I \tau^I + 8e^{2\Lambda} r \lambda \tau^I \Lambda' - 4r\sigma^I \tau^I \Lambda' - 2r\tau^I (\sigma^I)' + 2e^{2\Lambda} r \lambda (\tau^I)' - 2r\sigma^I (\tau^I)' \right], \quad (D16)$$

$$N_{HK}^I = (N^I)^2 + 2r\tau^I (\sigma^I - \kappa^I) (N^I)' \\ + N^I \left\{ 2\gamma^I \sigma^I + \rho^I (\sigma^I)^2 - 4\sigma^I \tau^I + 2r\tau^I (\kappa^I)' - 4r\sigma^I \tau^I \Lambda' + \kappa^I [-2\gamma^I - \rho^I \sigma^I + 4\tau^I (1 + r\Lambda')] - 2r\tau^I (\sigma^I)' \right\}. \quad (D17)$$

## 2. Even-parity and the RW gauge for $l \geq 2$

The parameters  $\tau^{II}$ ,  $\sigma^{II}$ ,  $\kappa^{II}$ ,  $\rho^{II}$ ,  $\mu^{II}$  and  $\nu^{II}$  in Eqs.(50) and (51) have the following form:

$$\tau^{II} = \sigma^{II} + 2r\Phi' - 2, \quad (D18)$$

$$\sigma^{II} = 2 - 2e^{2\Lambda} + e^{2\Lambda} \lambda + 4r\Phi' - 4r\Lambda' + 2r^2 \Phi'^2 - 2r^2 \Lambda' \Phi' + 2r^2 \Phi'', \quad (D19)$$

$$\kappa^{II} = -e^{2\Lambda} \lambda (e^{2\Lambda} \lambda + 2r\Phi' - 2) + 4(1 - e^{2\Lambda} + 2r\Phi' - 2r\Lambda' + r^2 \Phi'^2 - r^2 \Lambda' \Phi' + r^2 \Phi'')^2, \quad (D20)$$

$$\rho^{II} = 4r\Lambda' - 4r\Phi' + e^{2\Lambda} (\lambda - 2)r\Lambda' - 2r^2 (2\Lambda'^2 - 5\Lambda' \Phi' + \Phi'^2 + \Phi'') + 2r^3 \Lambda' (\Phi'^2 + \Phi'' - \Lambda' \Phi'), \quad (D21)$$

$$\mu^{II} = e^{2\Lambda} [2 + e^{2\Lambda} (\lambda - 2)] (\lambda - 2) - 4r\Lambda' [1 + e^{2\Lambda} (\lambda - 2)] + 2r\Phi' [4 + 3e^{2\Lambda} (\lambda - 2)] \\ + r^2 \left[ 6\Phi'^2 - 4\Lambda' \Phi' + 2\Phi'' - e^{2\Lambda} (\lambda - 2) (2\Lambda'^2 + \Lambda' \Phi' - 2\Phi'^2 - \Lambda'' - 2\Phi'') \right] \\ + 2r^3 [4\Lambda'^3 - 8\Lambda'^2 \Phi' + \Lambda' (\Phi'^2 - 2\Lambda'') + \Phi' (\Phi'^2 + 3\Lambda'' + \Phi'')] \\ + 2r^4 (2\Lambda'^2 - \Lambda' \Phi' - \Lambda'') (\Lambda' \Phi' - \Phi'^2 - \Phi''), \quad (D22)$$

$$\nu^{II} = e^{2\Lambda} (\lambda - 2) - 4r\Lambda' [1 + e^{2\Lambda} (\lambda - 2)] + r\Phi' [8 + e^{2\Lambda} (\lambda - 2)] \\ + 2r^2 [8\Lambda'^2 - 15\Lambda' \Phi' + 3\Phi'^2 + \Phi''] + 2r^3 (4\Lambda' - \Phi') (\Lambda' \Phi' - \Phi'^2 - \Phi''). \quad (D23)$$

The parameters in the master equation Eq.(53) and the source term Eq.(54) are functions of background  $\Psi(r)$  and  $\Lambda(r)$ . Similarly, to simplify the expressions of  $N_T^{II}$ ,  $N_R^{II}$  and  $N_Z^{II}$ , two auxiliary parameters  $M_1^{II}$  and  $M_2^{II}$  are introduced

$$M_1^{II} = 2\nu^{II} + \sigma^{II} (\sigma^{II} - 2), \quad (D24)$$

$$M_2^{II} = 2\mu^{II} - \sigma^{II} \rho^{II} - r\tau^{II} (\sigma^{II})'. \quad (D25)$$

Then the parameters in Eq.(53) are given by

$$N^{II} = -4\mu^{II} + 2\sigma^{II} \rho^{II} + 2\sigma^{II} \nu^{II} - 2(\sigma^{II})^2 + (\sigma^{II})^3 + 2r\tau^{II} \cdot (\sigma^{II})', \quad (D26)$$

$$N_T^{II} = \frac{e^{2\Lambda - 2\Phi} (\sigma^{II} M_1^{II} - 2M_2^{II})}{(\tau^{II})^2 [\sigma^{II} - 2r(\Lambda' + \Phi')]}, \quad (D27)$$

$$N_R^{II} = - (N^{II})^2 - 2r\tau^{II} \sigma^{II} \cdot (N^{II})' + N^{II} \left[ -4\mu^{II} + M_1^{II} \sigma^{II} + 2(\nu^{II} \sigma^{II} + r\tau^{II} (\sigma^{II})' + \tau^{II} \sigma^{II} + r\sigma^{II} (\tau^{II})') \right], \quad (D28)$$

$$N_Z^{II} = N^{II} \left( 2M_2^{II} \nu^{II} + r\tau^{II} \sigma^{II} \cdot (M_1^{II})' \right) + M_1^{II} \left[ -r\tau^{II} \sigma^{II} (N^{II})' + N^{II} (-2\mu^{II} + r\tau^{II} \cdot (\sigma^{II})') \right]. \quad (D29)$$

and the parameters in Eq.(54) are given by

$$N_A^{\text{II}} = - (N^{\text{II}})^2 - 2r\sigma^{\text{II}}\tau^{\text{II}} \cdot (N^{\text{II}})' + 2N^{\text{II}} \left( -2\mu^{\text{II}} + \nu^{\text{II}}\sigma^{\text{II}} + r\tau^{\text{II}} (\sigma^{\text{II}})' \right) + 2N^{\text{II}}\sigma^{\text{II}} \left( r (\tau^{\text{II}})' + 2\tau^{\text{II}} + 4r\Lambda'\tau^{\text{II}} - 2r\Phi'\tau^{\text{II}} \right), \quad (\text{D30})$$

$$N_F^{\text{II}} = - (N^{\text{II}})^2 \kappa^{\text{II}} + 2N^{\text{II}} [\kappa^{\text{II}} + (e^{2\Lambda}\lambda - 2)\sigma^{\text{II}}] [-2\mu^{\text{II}} + \sigma^{\text{II}} (\nu^{\text{II}} + 2\tau^{\text{II}})] - 2r\kappa^{\text{II}}\sigma^{\text{II}}\tau^{\text{II}} \cdot (N^{\text{II}})' + 2r \left[ N^{\text{II}}\kappa^{\text{II}}\tau^{\text{II}} (\sigma^{\text{II}})' + N^{\text{II}}\sigma^{\text{II}}\tau^{\text{II}} \left( (\kappa^{\text{II}})' + 2\kappa^{\text{II}}\Lambda' + 2e^{2\Lambda}\lambda (\sigma^{\text{II}})' - 4(\sigma^{\text{II}})' + 4r\Phi' (\sigma^{\text{II}})' \right) - 4N^{\text{II}}\sigma^{\text{II}}\mu^{\text{II}}\Phi' \right] + 2r (\sigma^{\text{II}})^2 \left[ 2N^{\text{II}}\nu^{\text{II}}\Phi' - \tau^{\text{II}} (N^{\text{II}})' (e^{2\Lambda}\lambda - 2) + 4\tau^{\text{II}}N^{\text{II}}\Lambda' (e^{2\Lambda}\lambda - 1) + 6\tau^{\text{II}}N^{\text{II}}\Phi' \right] + 4r^2 (\sigma^{\text{II}})^2 \tau^{\text{II}} \left( - (N^{\text{II}})' \Phi' + 2N^{\text{II}}\Lambda'\Phi' + N^{\text{II}}\Phi'' \right). \quad (\text{D31})$$

### 3. Odd-parity for $l \geq 2$

The parameters of  $X$  and  $Y$  in the odd-parity master equation for  $l \geq 2$  are

$$X = 1 - 3e^{2\Lambda} + e^{2\Lambda}\lambda + 3r\Phi' + 3r^2\Phi'^2 - r\Lambda'(5 + 3r\Phi') + 3r^2\Phi'', \quad (\text{D32})$$

$$Y = 1 - 3e^{2\Lambda} + e^{2\Lambda}\lambda + 5r\Phi' + 3r^2\Phi'^2 - 3r\Lambda'(1 + r\Phi') + 3r^2\Phi''. \quad (\text{D33})$$

And the expression of parameters  $N_{\text{odd}}$  is given by

$$N_{\text{odd}} = 2r^2\Lambda'^2 + r\Phi' + 2r^2\Phi'^2 + r\Lambda' + 4r^2\Lambda'\Phi' - 2 - 2r^2\Lambda'' - 2r^2\Phi'' - Y + \frac{rY'}{Y} (-1 + 2r\Lambda' + 2r\Phi'). \quad (\text{D34})$$

### 4. The case for $l = 0$

In this case, the parameters  $\mu_0$ ,  $\nu_0$  and  $\tau_0$  are given by

$$\mu_0 = \frac{2 - 2e^{2\Lambda} - 2r^2\Lambda'^2(2 + r\Phi') + 2r^2\Lambda'' - 2r^2\Phi'' + r^2\Lambda'(4\Phi' + 2r\Phi'^2 + 3r\Phi'') + r\Phi'(2 + r^2\Lambda'' - 2r^2\Phi'') - r^3\Phi'''}{r(-1 + e^{2\Lambda} - 2r\Phi' - r^2\Phi'^2 + r\Lambda'(2 + r\Phi') - r^2\Phi'')}, \quad (\text{D35})$$

$$\nu_0 = 2r^{-2}e^{-4\Lambda+2\Phi} \left\{ \Lambda'(-2 + 8r\Lambda') - 2(-1 + 4r\Lambda')\Phi' + (-1 + 4r\Lambda')(-4\Lambda' + 2\Phi') + 4(\Lambda' + r\Lambda'') + \frac{(-1 + 4r\Lambda') \left[ -2r\Phi'^2 + \Phi'(-2 + r^2\Lambda'' - 2r^2\Phi'') + \Lambda'(2 + 2e^{2\Lambda} + 2r\Phi' + r^2\Phi'') + r(2\Lambda'' - 4\Phi'' - r\Phi''') \right]}{1 - e^{2\Lambda} + 2r\Phi' + r^2\Phi'^2 - r\Lambda'(2 + r\Phi') + r^2\Phi''} \right\}, \quad (\text{D36})$$

$$\tau_0 = \frac{2e^{-4\Lambda+2\Phi} \left[ 1 - e^{2\Lambda} - r^2\Phi'^2 + 2r^2\Lambda'^2(2 + r\Phi') + 2r^2\Lambda'' + r^3\Phi'(\Lambda'' - 2\Phi'') \right]}{r^2(-1 + e^{2\Lambda} - 2r\Phi' - r^2\Phi'^2 + r\Lambda'(2 + r\Phi') - r^2\Phi'')} + \frac{2e^{-4\Lambda+2\Phi} \left[ -3r^2\Phi'' - r\Lambda' \left( 2 - 4e^{2\Lambda} + 3r\Phi' + 2r^2\Phi'^2 + r^2\Phi'' \right) - r^3\Phi'''\right]}{r^2(-1 + e^{2\Lambda} - 2r\Phi' - r^2\Phi'^2 + r\Lambda'(2 + r\Phi') - r^2\Phi'')}, \quad (\text{D37})$$

$$\sigma_0 = 4r\Phi'^2 + 2r^2\Phi'^3 + 3r\Lambda'^2(2 + r\Phi') + \Lambda'(-6 + e^{2\Lambda} - 10r\Phi' - 5r^2\Phi'^2 - 4r^2\Phi'') + \Phi'(2 - 2e^{2\Lambda} - r^2\Lambda'' + 4r^2\Phi'') + r(-2\Lambda'' + 4\Phi'' + r\Phi'''), \quad (\text{D38})$$

$$\iota_0 = -1 + e^{2\Lambda} - r\Phi' - r^2\Phi'^2 + r\Lambda'(2 + r\Phi') - r^2\Phi''. \quad (\text{D39})$$



### 5. Odd-parity for $l = 1$

In Eq.(89), the parameters  $X_1$  and  $Y_1$  are given by

$$X_1 = -1 + e^{2\Lambda} - 5r\Phi' - 3r^2\Phi'^2 + 3r\Lambda'(1 + r\Phi') - 3r^2\Phi'', \quad (\text{D40})$$

$$Y_1 = -1 + e^{2\Lambda} - 3r\Phi' - 3r^2\Phi'^2 + r\Lambda'(5 + 3r\Phi') - 3r^2\Phi'', \quad (\text{D41})$$

and  $N_1$  is given by

$$N_1 = X_1 + \frac{rX_1'}{X_1}(-3 + 2r\Lambda' + 2r\Phi') + r \left[ 2r\Lambda'^2 - 5\Phi' + 2r\Phi'^2 + \Lambda'(-5 + 4r\Phi') - 2r(\Lambda'' + \Phi'') \right]. \quad (\text{D42})$$

### Appendix E: The Degeneration Of Eq.(42)

The coefficient of the third order term in Eq.(42) is given by

$$\frac{2e^{2\Lambda-2\Phi}r}{e^{2\Lambda}\lambda - 2\sigma^{\text{I}}} 2r(\sigma^{\text{I}} - \kappa^{\text{I}}). \quad (\text{E1})$$

This coefficient would vanish when the relation  $\sigma^{\text{I}} - \kappa^{\text{I}} = 0$  is satisfied. From Appendix D we know that

$$\sigma^{\text{I}} - \kappa^{\text{I}} = -1 + e^{2\Lambda} - 3r\Phi' - r^2\Phi'^2 + r\Lambda' + r^2\Lambda'\Phi' - r^2\Phi''. \quad (\text{E2})$$

If the spherically symmetric spacetime Eq.(1) becomes

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (\text{E3})$$

we find that now the coefficient of the third order term of Eq.(42) vanish under the condition that the scala curvature vanish,  $R = 0$ .

Then considering that the metric of the background spacetime degenerates to the Schwarzschild spacetime, i.e.

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (\text{E4})$$

where the Riemann curvature  $R_{ab}$  and the scalar curvature  $R$  would vanish. The master variable and equation Eq.(42) would become

$$Z = \frac{l(l+1)r}{r-2M}\chi^{\text{I}} - 2\psi^{\text{I}}, \quad (\text{E5})$$

$$\begin{aligned} & \frac{2r^3}{(r-2M)(6M+r\tilde{\lambda})} \frac{\partial^2}{\partial t^2} Z - \frac{2r(r-2M)}{6M+r\tilde{\lambda}} \frac{\partial^2}{\partial r^2} Z - \frac{4 \left[ 6M^2 + r^2\tilde{\lambda} + 3(\tilde{\lambda}+4)Mr \right]}{(6M+r\tilde{\lambda})^2} \frac{\partial}{\partial r} Z \\ & - \frac{2[24M^3 + 12(l^2+l-1)M^2r + 2(4-l+2l^2+l^4)Mr^2 + l(2+l-2l^2-l^3)r^3]}{(r-2M)r(6M+r\tilde{\lambda})^2} Z = S_Z, \end{aligned} \quad (\text{E6})$$

with the source term

$$\begin{aligned} S_Z = & \frac{r^4 \left[ 96M^2 + 2(7\tilde{\lambda} - 18)Mr + \tilde{\lambda}(\tilde{\lambda} - 2)r^2 \right]}{2(r-2M)^3(6M+r\tilde{\lambda})^2} E_A - \frac{r^5}{(r-2M)^2(6M+r\tilde{\lambda})} \frac{\partial}{\partial r} E_A \\ & - \frac{r^5}{(r-2M)^2(6M+r\tilde{\lambda})} \frac{\partial}{\partial t} E_D + \frac{l(l+1)r^3}{2(r-2M)^2} E_F - \frac{l(l+1)r^3}{2(r-2M)(6M+r\tilde{\lambda})} (2E_H + E_K), \end{aligned} \quad (\text{E7})$$

where  $\tilde{\lambda} = \lambda - 2 = (l+2)(l-1)$ . It is obvious that our result reduces to that of the Schwarzschild spacetime [28].

---

[1] T. Regge and J. A. Wheeler, *Stability of a Schwarzschild singularity*, Phys. Rev. **108**, 1063 (1957).

- [2] L. A. Edelman and C. V. Vishveshwara, *Differential Equations for Perturbations on the Schwarzschild Metric*, Phys. Rev. D, **1**, 12: 3514 (1970).
- [3] F. J. Zerilli, *Effective potential for even parity Regge-Wheeler gravitational perturbation equations*, Phys. Rev. Lett. **24**, 13: 737 (1970).
- [4] F. J. Zerilli, *Gravitational field of a particle falling in a schwarzschild geometry analyzed in tensor harmonics*, Phys. Rev. D **2**, 10: 2141 p(1970).
- [5] V. Moncrief, *Gravitational perturbations of spherically symmetric systems. I. The exterior problem*, Ann. Phys. **88**, 323 (1974).
- [6] S. Chandrasekhar and S. L. Detweiler, *The quasi-normal modes of the Schwarzschild black hole*, Proc. Roy. Soc. Lond. A **344**, 441 (1975).
- [7] S. Chandrasekhar, *The mathematical theory of black holes* (Oxford University Press, 1992).
- [8] C. V. Vishveshwara, *Scattering of Gravitational Radiation by a Schwarzschild Black-hole*, Nature **227**, 936 (1970).
- [9] V. Ferrari and B. Mashhoon, *Oscillations of a Black Hole* Phys. Rev. Lett. **52**, 16: 1361 (1984).
- [10] K. D. Kokkotas and B. G. Schmidt, *Quasi-Normal Modes of Stars and Black Holes* Living Rev. Rel. **2**, 2 (1999).
- [11] H. P. Nollert, *Class. Quant. Grav. Quasinormal modes: the characteristic ‘sound’ of black holes and neutron stars* **16**, R159 (1999).
- [12] V. Cardoso, *Quasinormal modes and gravitational radiation in black hole spacetimes* arXiv: gr-qc/0404093 (2004).
- [13] E. Berti, V. Cardoso and A. O. Starinets, *Quasinormal modes of black holes and black branes* Class. Quant. Grav. **26**, 163001 (2009).
- [14] P. Pani, *Advanced Methods in Black-hole Perturbation Theory* Int. J. Mod. Phys. A **28**, 22n23, 1340018 (2013).
- [15] R. A. Konoplya and A. Zhidenko, *Quasinormal modes of black holes: From astrophysics to string theory* Rev. Mod. Phys. **83**, 793 (2011).
- [16] M. Davis, R. Ruffini, W. H. Press and R. H. Price, *Gravitational Radiation from a Particle Falling Radially into a Schwarzschild Black Hole* Phys. Rev. Lett. **27**, 21, 1466 (1971).
- [17] K. Martel, *Gravitational waveforms from a point particle orbiting a Schwarzschild black hole* Phys. Rev. D **69**, 044025 (2004).
- [18] R. H. Price, *Nonspherical Perturbations of Relativistic Gravitational Collapse I. Scalar and Gravitational Perturbations* Phys. Rev. D **5**, 10: 2419 (1972).
- [19] M. Dafermos, G. Holzegel and I. Rodnianski, *The linear stability of the Schwarzschild solution to gravitational perturbations* Acta Math. **222**, 1 (2019).
- [20] K. Prabhu and R. M. Wald, *Canonical Energy and Hertz Potentials for Perturbations of Schwarzschild Spacetime* Class. Quant. Grav. **35**, 23, 235004 (2018).
- [21] V. Moncrief, *Gravitational perturbations of spherically symmetric systems. II. Perfect fluid interiors* Ann. Phys. **88**, 343 (1974).
- [22] U. H. Gerlach and U. K. Sengupta, *Gauge Invariant Perturbations On Most General Spherically Symmetric Space-times* Phys. Rev. D **19**, 2268-2272 (1979)
- [23] U. H. Gerlach and U. K. Sengupta, *Gauge Invariant Coupled Gravitational, Acoustical, And Electromagnetic Modes On Most General Spherical Space-times* Phys. Rev. D **22**, 1300-1312 (1980)
- [24] K. S. Thorne, *Multipole Expansions of Gravitational Radiation* Rev. Mod. Phys. **52**, 299 (1980).
- [25] K. Martel and E. Poisson, *Gravitational perturbations of the Schwarzschild spacetime: A Practical covariant and gauge-invariant formalism* Phys. Rev. D **71**, 104003 (2005)
- [26] M. Lenzi and C. F. Sopuerta, *Master functions and equations for perturbations of vacuum spherically symmetric spacetimes* Phys. Rev. D **104**, 084053 (2021)
- [27] B. Preston and E. Poisson, *Light-cone gauge for black-hole perturbation theory* Phys. Rev. D **74**, 064010 (2006)
- [28] J. E. Thompson, B. F. Whiting and H. Chen, *Gauge invariant perturbations of the Schwarzschild spacetime* Class. Quant. Grav. **34**, 174001 (2017).
- [29] S. L. Detweiler, *Consequence of the gravitational self-force for circular orbits of the Schwarzschild geometry* Phys. Rev. D **77**, 124026 (2008).
- [30] E. Corrigan and E. Poisson, *EZ gauge is singular at the event horizon* Class. Quant. Grav. **35**, 137001 (2018).
- [31] D. C. Zou and Y. S. Myung, *Black hole with primary scalar hair in Einstein-Weyl-Maxwell conformal scalar theory* Phys. Rev. D **101**, 084021 (2020).
- [32] L. Heisenberg, R. Kase and S. Tsujikawa, *Odd-parity stability of hairy black holes in  $U(1)$  gauge-invariant scalar-vector-tensor theories* Phys. Rev. D **97**, 124043 (2018).
- [33] K. Tomikawa and T. Kobayashi, *Perturbations and quasi-normal modes of black holes with time-dependent scalar hair in shift-symmetric scalar-tensor theories* Phys. Rev. D **103**, 084041 (2021).
- [34] A. Buonanno and T. Damour, *Effective one-body approach to general relativistic two-body dynamics* Phys. Rev. D **59**, 084006 (1999)
- [35] A. Buonanno and T. Damour, *Transition from inspiral to plunge in binary black hole coalescences* Phys. Rev. D **62**, 064015 (2000)
- [36] E. Poisson, A. Pound and I. Vega, *The Motion of point particles in curved spacetime* Living Rev. Rel. **14**, 7 (2011).
- [37] R. M. Wald, *General relativity* (University of Chicago Press, Chicago, 1984).