

Stable circular orbit of a spinning test particle in rotating boson star

Yu-Peng Zhang^{*}, Yan-Bo Zeng[†], Yong-Qiang Wang[‡], Shao-Wen Wei[§], Yu-Xiao Liu[¶]
*Lanzhou Center for Theoretical Physics, Key Laboratory of Theoretical Physics of Gansu Province,
Institute of Theoretical Physics & Research Center of Gravitation, Lanzhou University, Lanzhou 730000, China
School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

In this paper, we study the circular orbit of the spinning test particle in the background of a rotating boson star. Using the pole-dipole approximation and neglecting the back-reaction of the spinning test particle on the spacetime, the equation of motion of the spinning test particle is described by the Mathisson-Papapetrou-Dixon equation. We solve this equation under the Tulczyjew spin-supplementary condition and obtain the four-momentum and four-velocity of the spinning test particle. Quite different from the spinless particle, the effective potential of the spinning particle with zero orbital angular momentum goes to infinite at the center of the rotating boson star. This will lead to the fact that the spinning particle can not pass through the center of the boson star. However, when the spin angular momentum and orbital angular momentum satisfy $2\bar{s} + \bar{l} = 0$, the effective potential is not divergent anymore and the spinning particle can pass through the center of the rotating boson star. Moreover, the radius and energy of the circular orbit will be decreased or increased by the particle spin. These results will have an important application in testing the gravitational waves in the boson star background.

PACS numbers:

I. INTRODUCTION

Boson stars are formed by the self-gravitating complex scalar fields with everywhere regular localized configurations, and they were firstly introduced in the end of 1960s [1–3]. They can have a larger mass with a small size without any event horizon and singularity. Since boson stars were proposed in 1968, the research of boson stars has received a lot of attention and various boson stars have been proposed [4–23]. As the mimicker of the black hole [24], boson stars are the important objects in the cosmology [25–27]. The possible gravitational waves emitted from boson stars have been still studied extensively [28–31] in terms of the nonlinear evolution. Especially the study in [31] showed that the binary boson stars can also be the degenerate gravitational wave source of the GW190521 [32].

Continued research on boson stars shows that boson stars are very important test beds of the strong gravity system and the corresponding properties should be investigated in detail. The problem of the equations of motion for a test particle in the background of strong gravity system is critical for understanding the motion of various stars in the galaxy. Note that, a boson star is everywhere regular and there is no phenomenon that a test particle plunges into the event horizon of a black hole. Such properties ensure that novel orbits will be obtained in the backgrounds of boson stars [10, 33–36]. For example, a test particle with zero orbital angular mo-

mentum can pass through the center of the boson star. Usually, the test particle approach ignores the inner-structure of a small body. For a test particle with a spin, such as a small body with a dipole inner structure, the spin-curvature force will lead to that the corresponding equation of motion of the test particle will no longer be geodesics. Therefore, there is a natural question: can a test particle pass through the center of the boson star when its spin (dipole inner-structure) can not be neglected? In this paper, we will consider a spinning test particle under the pole-dipole approximation and use the Mathisson-Papapetrou-Dixon (MPD) equations [37–40] to study the properties of the circular orbits in rotating boson stars.

This paper is organized as follows. In Sec. II, we briefly review the construction of a boson star and obtain the four-momentum and four-velocity of a spinning test particle in the equatorial plane by solving the MPD equations. In Sec. III, we derive the radial effective potential of the spinning test particle in the background of the rotating boson star and investigate the properties of the circular orbits of the spinning test particle. Finally, the brief conclusion and discussion are given in Sec. IV.

II. MPD EQUATION SOLUTIONS IN BACKGROUND OF A BOSON STAR

Firstly, we give a brief review about the construction of a boson star solution. A boson star is a strong gravity system formed by the gravitationally bound Bose-Einstein condensate for a light scalar with long de Broglie wavelength. It is described by the following action

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{16\pi G} R - \nabla_\mu \Phi \nabla^\mu \Phi^* - V(\Phi) \right]. \quad (1)$$

^{*}zyp@lzu.edu.cn

[†]zengyb19@lzu.edu.cn

[‡]yqwang@lzu.edu.cn, corresponding author

[§]weishw@lzu.edu.cn

[¶]liuyx@lzu.edu.cn

We take the Planck units and set the Newton gravitational constant G , the speed of light c , and the Planck constant \hbar to be unity ($G = c = \hbar = 1$). We only consider the mini boson star [41] and the potential $V(\Phi)$ is taken as the simplest form

$$V(\Phi) = \frac{\mu^2}{\hbar^2} \Phi \Phi^*, \quad (2)$$

where μ is the mass parameter of the scalar field Φ .

The geometry of a rotating boson star is stationary and axisymmetric, the corresponding ansatz of the complex scalar field Φ is defined as follows

$$\Phi = \phi(r, \theta) \exp [i(\omega t - k\varphi)], \quad (3)$$

where the scalar field is dependent on the coordinates of the spacetime (t, r, θ, φ) and the properties of a rotating boson star are controlled by the frequency ω and angular number k . The stationary and axisymmetry of a rotating boson star demand that the metric of the rotating boson star satisfies the form [42] as follows

$$\begin{aligned} ds^2 &= g_{tt} dt^2 + g_{rr} dr^2 + g_{\theta\theta} d\theta^2 \\ &\quad + 2g_{t\varphi} dt d\varphi + g_{\varphi\varphi} d\varphi^2 \\ &= e^{2F_1} (dr^2 + r^2 d\theta^2) - e^{2F_0} dt^2 \\ &\quad + e^{2F_2} r^2 \sin^2 \theta (d\varphi - W dt)^2, \end{aligned} \quad (4)$$

where the metric functions F_0 , F_1 , F_2 , and W are the functions of the coordinate r and θ . By assuming the proper boundary conditions and using the ansatz of the scalar field (3) and metric (4), one could obtain the numerical solutions of the boson star [42, 43]. In this paper, we consider three solutions of the rotating boson star, we list the properties of these boson star solutions in Tab. I.

TABLE I: Physical properties of the three rotating boson star solutions that we considered.

	$k = 1$	$k = 2$	$k = 3$
$\omega(\mu/\hbar)$	0.8900	0.8900	0.8900
$M(m_p^2/\mu)$	1.1470	1.6388	2.0921

Next we start to solve the equations of motion for a spinning test particle in the background of the rotating boson star. For a spinning test particle, the corresponding equations of motion will no longer be geodesics and they are described by the MPD equations [37–40, 44, 45]

$$\frac{DP^\mu}{D\lambda} = -\frac{1}{2} R^\mu_{\nu\alpha\beta} u^\nu S^{\alpha\beta}, \quad (5)$$

$$\frac{DS^{\mu\nu}}{D\lambda} = P^\mu u^\nu - u^\mu P^\nu, \quad (6)$$

where P^μ , $S^{\mu\nu}$, and u^μ are the four-momentum, spin tensor, and four-velocity of the spinning test particle along

the trajectory, respectively. However, the MPD equations can not uniquely define the motion of the spinning test particle. Hence an additional spin-supplementary condition is necessary [44, 46–48]. Actually, the spin-supplementary condition is related to the center of mass of the spinning test particle with different observers [44, 46–48]. Recently, the works [49–51] showed that the expression of the three-acceleration contains the divergent terms as the three-velocity satisfies $v \rightarrow c$, where c is the speed of light. Such divergence can be avoided by adding a non-minimal spin-gravity interaction through gravimagnetic moment [49]. In this paper, we ignore above behaviors and take the Tulczyjew spin-supplementary condition [52]

$$P_\mu S^{\mu\nu} = 0. \quad (7)$$

The four-momentum P^μ of the spinning test particle keeps timelike along the trajectory and satisfies

$$P^\mu P_\mu = -m^2, \quad (8)$$

where m is the mass of the spinning test particle.

We only consider the equatorial motion of the spinning test particle with spin-aligned or anti-aligned orbits. All the components of the four-momentum and spin tensor should satisfy $P^\theta = 0$ and $S^{\theta\mu} = 0$. Therefore, the non-vanishing independent components of the equatorial orbits are P^t , P^r , P^ϕ , and $S^{r\phi}$. By using the spin-supplementary condition (7), we get [53]

$$S^{rt} = -S^{r\phi} \frac{P_\phi}{P_t}, \quad S^{\phi t} = S^{r\phi} \frac{P_r}{P_t}. \quad (9)$$

The conservation of the spin for the spinning test particle demands that

$$s^2 = \frac{1}{2} S^{\mu\nu} S_{\mu\nu} = S^{\phi r} S_{\phi r} + S^{tr} S_{tr} + S^{t\phi} S_{t\phi}. \quad (10)$$

Combining (8), (9), and (10), we obtain the independent $r - \phi$ component of the spin tensor

$$S^{r\phi} = \frac{s}{m} \frac{P_t}{H}, \quad (11)$$

where the function H is

$$H = \sqrt{g_{rr}(g_{\varphi t}^2 - g_{\varphi\varphi} g_{tt})}. \quad (12)$$

Then the non-vanishing components of the spin tensor in the background of the rotating boson star are

$$\begin{aligned} S^{r\phi} &= -S^{\phi r} = \bar{s} \frac{P_t}{H}, \\ S^{rt} &= -S^{tr} = -S^{r\phi} \frac{P_\phi}{P_t} = -\bar{s} \frac{P_\phi}{H}, \\ S^{\phi t} &= -S^{t\phi} = S^{r\phi} \frac{P_r}{P_t} = \bar{s} \frac{P_r}{H}, \end{aligned} \quad (13)$$

where the parameter $\bar{s} = \frac{s}{m}$ is the per unit mass spin angular momentum of the test particle and the spin direction is perpendicular to the equatorial plane.

The stationary and axisymmetry mean that there are a timelike Killing vector $\xi^\mu = (\partial_t)^\mu$ and a spacelike Killing vector $\eta^\mu = (\partial_\varphi)^\mu$. Due to the spin-curvature force, the related conserved quantities of the spinning test particle will be different from the test point particle. For a Killing vector \mathcal{K}^μ , the conserved quantity is [45, 53]

$$\mathcal{C} = \mathcal{K}^\mu P_\mu - \frac{1}{2} S^{\mu\nu} \nabla_\nu \mathcal{K}_\mu. \quad (14)$$

In the background of a rotating boson star with the metric (4), there are a timelike Killing vector $\xi^\mu = (\partial_t)^\mu$ and a spacelike Killing vector $\eta^\mu = (\partial_\varphi)^\mu$. The two Killing vectors satisfy

$$S^{\mu\nu} \xi_{\mu;\nu} = S^{\mu\nu} \xi^\beta \partial_\nu g_{\beta\mu}, \quad (15)$$

$$S^{\mu\nu} \eta_{\mu;\nu} = S^{\mu\nu} \eta^\beta \partial_\nu g_{\beta\mu}. \quad (16)$$

Then we can get the two conserved quantities [53]

$$\begin{aligned} m\bar{e} &= -\mathcal{C}_t = -\xi^\mu P_\mu + \frac{1}{2} S^{\mu\nu} \xi_{\mu;\nu} \\ &= -P_t - \frac{1}{2} \frac{\bar{s}}{H} P_t \partial_r g_{t\phi} + \frac{1}{2} \frac{\bar{s}}{H} P_\phi \partial_r g_{tt}, \end{aligned} \quad (17)$$

$$\begin{aligned} m\bar{j} &= \mathcal{C}_\phi = \eta^\mu P_\mu - \frac{1}{2} S^{\mu\nu} \eta_{\mu;\nu} \\ &= P_\phi - \frac{1}{2} \frac{\bar{s}}{H} P_\phi \partial_r g_{\phi t} + \frac{1}{2} \frac{\bar{s}}{H} P_t \partial_r g_{\phi\phi}, \end{aligned} \quad (18)$$

where $\bar{e} = \frac{e}{m}$ and $\bar{j} = \frac{j}{m}$. The parameters e , m , and j are the energy, mass, and total angular momentum of the spinning test particle, respectively. We redefine the total angular momentum $\bar{j} = \bar{s} + \bar{l}$, where \bar{l} is the orbital angular momentum.

Solving Eqs. (8), (17), and (18), we get the non-vanishing components of the four-momentum:

$$P_t = \frac{2mH(2\bar{e}H - \bar{e}\bar{s}\partial_r g_{\phi t} - \bar{j}\bar{s}\partial_r g_{tt})}{4g_{rr}H^2 + \bar{s}^2[(\partial_r g_{\phi t})^2 - \partial_r g_{\phi\phi}\partial_r g_{tt}]}, \quad (19)$$

$$P_\phi = \frac{2mH(2\bar{j}H - \bar{e}\bar{s}\partial_r g_{\phi\phi} + \bar{j}\bar{s}\partial_r g_{\phi t})}{4g_{rr}H^2 + \bar{s}^2[(\partial_r g_{\phi t})^2 - \partial_r g_{\phi\phi}\partial_r g_{tt}]}, \quad (20)$$

and

$$(P^r)^2 = -\frac{m^2 + g^{\phi\phi}P_\phi^2 + 2g^{\phi t}P_\phi P_t + g^{tt}P_t^2}{g_{rr}}. \quad (21)$$

Due to the trajectories of the test particle are independent of the affine parameter λ [40, 48], we set the affine parameter λ as the coordinate time and choose $u^t = 1$. We can solve the four-velocity u^μ by using the equations of motion (5) and (6) and the components of $S^{\mu\nu}$ in (13) [54, 55]

$$\frac{DS^{tr}}{D\lambda} = P^t \dot{r} - P^r = -\frac{\bar{s}}{2H} g_{\phi\mu} R_{\nu\alpha\beta}^\mu u^\nu S^{\alpha\beta}, \quad (22)$$

$$\frac{DS^{t\phi}}{D\lambda} = P^t \dot{\phi} - P^\phi = \frac{\bar{s}}{2H} g_{r\mu} R_{\nu\alpha\beta}^\mu u^\nu S^{\alpha\beta}. \quad (23)$$

Finally, we can get the non-vanishing components of the four-velocity as follows

$$\dot{r} = \frac{b_1 c_2 - a_2 c_1}{a_1 a_2 - b_1 b_2}, \quad (24)$$

$$\dot{\phi} = \frac{b_2 c_1 - a_1 c_2}{a_1 a_2 - b_1 b_2}, \quad (25)$$

where the functions a_1 , b_1 , c_1 , a_2 , b_2 , and c_2 are defined as

$$a_1 = P^t + \frac{\bar{s}}{2H} R_{\phi r \mu\nu} S^{\mu\nu}, \quad (26)$$

$$b_1 = \frac{\bar{s}}{2H} R_{\phi\phi\mu\nu} S^{\mu\nu}, \quad (27)$$

$$c_1 = -P^r + \frac{\bar{s}}{2H} R_{\phi t \mu\nu} S^{\mu\nu}, \quad (28)$$

$$a_2 = P^t - \frac{\bar{s}}{2H} R_{r\phi\mu\nu} S^{\mu\nu}, \quad (29)$$

$$b_2 = -\frac{\bar{s}}{2H} R_{rr\mu\nu} S^{\mu\nu}, \quad (30)$$

$$c_2 = -P^\phi - \frac{\bar{s}}{2H} R_{rt\mu\nu} S^{\mu\nu}. \quad (31)$$

Then the corresponding orbital frequency parameter Ω of the test particle is

$$\Omega \equiv \frac{u^\phi}{u^t} = \dot{\phi}. \quad (32)$$

III. EFFECTIVE POTENTIALS AND CIRCULAR ORBITS

Using the radial effective potential is an easy way to determine the radial motion of a test particle. For the spinning test particle in the equatorial plane of the rotating boson star, the radial momentum is proportional to the radial velocity. Thus, we decompose the radial momentum (21) as follows

$$\begin{aligned} (u^r)^2 &= (A\bar{e}^2 + B\bar{e} + C) \\ &\propto \left(\bar{e} - \frac{-B + \sqrt{B^2 - 4AC}}{2A} \right) \\ &\times \left(\bar{e} - \frac{-B - \sqrt{B^2 - 4AC}}{2A} \right), \end{aligned} \quad (33)$$

where the functions A , B , and C are

$$A = -\frac{4m^2 g_{rr}^2}{D} \left\{ -\bar{s} \partial_r g_{\varphi\varphi} \left[2g_{\varphi t} (2H - \bar{s} \partial_r g_{\varphi t}) + \bar{s} \partial_r g_{\varphi\varphi} g_{tt} \right] + 4g_{\varphi\varphi}^2 g_{rr} g_{tt} + g_{\varphi\varphi} \left[\bar{s} \partial_r g_{\varphi t} (4H - \bar{s} \partial_r g_{\varphi t}) - 4g_{\varphi t}^2 g_{rr} \right] \right\}, \quad (34)$$

$$B = \frac{8\bar{j}m^2 g_{rr}^2}{D} \left\{ -g_{\varphi t} \left[\bar{s}^2 (\partial_r g_{\varphi\varphi} \partial_r g_{tt} + \partial_r g_{\varphi t}^2) + 4g_{\varphi\varphi} g_{rr} g_{tt} \right] - \bar{s} \left[g_{\varphi\varphi} \partial_r g_{tt} (2H - \bar{s} \partial_r g_{\varphi t}) - \partial_r g_{\varphi\varphi} g_{tt} (2H + \bar{s} \partial_r g_{\varphi t}) \right] + 4g_{\varphi t}^3 g_{rr} \right\}, \quad (35)$$

$$C = -\frac{m^2 g_{rr}}{D} \left\{ -8g_{\varphi t}^2 g_{rr} \left[\bar{s}^2 (\partial_r g_{\varphi t}^2 - \partial_r g_{\varphi\varphi} \partial_r g_{tt}) + 2g_{rr} g_{tt} (2g_{\varphi\varphi} + \bar{j}^2) \right] + 4g_{\varphi\varphi} g_{rr} \left[4\bar{j}^2 g_{rr} g_{tt}^2 + \bar{s}^2 (2g_{tt} (\partial_r g_{\varphi t}^2 - \partial_r g_{\varphi\varphi} \partial_r g_{tt}) - \bar{j}^2 \partial_r g_{tt}^2) \right] - 8\bar{j}^2 \bar{s} g_{\varphi t} g_{rr} \partial_r g_{tt} (-\bar{s} \partial_r g_{\varphi t} - 2H) + 16g_{\varphi t}^4 g_{rr}^2 + 4\bar{j}^2 \bar{s} \partial_r g_{\varphi t} g_{rr} g_{tt} (-\bar{s} \partial_r g_{\varphi t} - 4H) + 16g_{\varphi\varphi}^2 g_{rr}^2 g_{tt}^2 + \bar{s}^4 (\partial_r g_{\varphi t}^2 - \partial_r g_{\varphi\varphi} \partial_r g_{tt})^2 \right\}. \quad (36)$$

where

$$D = \left[\bar{s}^2 (\partial_r g_{\varphi\varphi} \partial_r g_{tt} - (\partial_r g_{\varphi t})^2) + 4H^2 \right]^2. \quad (37)$$

We can define the effective potential of the spinning test particle with the four-momentum pointing toward future by using the positive square root of Eq. (33) as follows [56]

$$V_{\text{eff}} = \frac{-B + \sqrt{B^2 - 4AC}}{2A}. \quad (38)$$

Now, we have got the four-momentum and four-velocity of the spinning test particle in the equatorial plane of the rotating boson star. Next, we will focus on the properties of the circular orbit $r = r_{\text{co}}$ of the spinning test particle, for which we have

$$\dot{r}|_{r=r_{\text{co}}} = 0, \quad (39)$$

$$\ddot{r}|_{r=r_{\text{co}}} = 0. \quad (40)$$

The above two equations correspond to $\bar{e} = V_{\text{eff}}|_{r=r_{\text{co}}}$ and $\frac{dV_{\text{eff}}}{dr}|_{r=r_{\text{co}}} = 0$. If we further have $\frac{d^2V_{\text{eff}}}{dr^2}|_{r=r_{\text{co}}} > 0$, then we will obtain a stable circular orbit. We have obtained the effective potential (38) of the spinning test

particle with the four-momentum pointing toward future and shown that it is dependent on the spin and angular momenta of the test particle. It can be shown that when the particle spin is zero, the effective potential is the same as that of the point particle [36].

Figure 1 shows the effective potentials for different values of the particle spin \bar{s} , total angular momentum \bar{j} , and angular number k . We can get the radius of the circular orbit by finding the location of the minimum of the effective potential. The corresponding angular frequency and energy can also be obtained by using Eqs. (32) and (39). Figure 2 shows how the circular orbit parameters (the radius r_{co} , energy \bar{e}_{co} , and angular frequency Ω_{co}) depend on the test particle spin.

Checking the effective potentials for spinning test particle, we observe some novel behaviors that are not obtained from the point-like test particle. We list them as follows:

(a) For a spinning particle with zero angular momentum, the spin-curvature force will cause the effective potential at the center of the rotating boson star to be infinite. It means a spinning particle can not pass through the center of the rotating boson star when its orbital angular momentum is zero.

(b) For a spinning particle with fixed orbital angular momentum, when the spin angular momentum changes from an negative value to a positive one, the radius r_{co} and the particle energy \bar{e}_{co} for the circular orbit will decrease first and then increase. With the change of the total angular momentum \bar{j} and spin angular momentum \bar{s} , the angular frequency for the circular orbit will transform from positive to negative, which will make the test particle move in the opposite angular direction. See the details in Fig. 2

(c) The most surprising result is that when the total angular momentum and spin angular momentum of the spinning particle satisfy $\bar{s} + \bar{j} = 0$, the effective potential at the origin is still finite. That is to say, when the spin angular momentum and the orbital angular momentum satisfy $2\bar{s} + \bar{l} = 0$, a spinning particle with suitable energy could pass through the center of the rotating boson star. See the details in Fig. 3.

IV. CONCLUSIONS AND DISCUSSION

In this paper, we investigated the circular orbits of a spinning test particle in the equatorial plane of the rotating boson star with the angular number $k = (1, 2, 3)$ and the frequency $\omega = 0.89$. We solved the four-velocity of the spinning test particle and derived the corresponding radial effective potential. We obtained the four-momentum and four-velocity of the spinning test particle by solving the MPD equations in the rotating boson star background. The radial effective potential was obtained in terms of the radial component of the four-momentum P^r for the spinning test particle with the four-momentum pointing toward future.




FIG. 1: The effective potentials of the spinning test particle for different values of the particle spin \bar{s} , total angular momentum \bar{j} , and angular number k . The angular number k is taken as $k = (1, 2, 3)$ for subfigures (a1-a3), (b1-b3), (c1-c3), respectively. Other parameters are set as $\omega = 0.89$, $\bar{s} = (-0.5, -0.25, 0, 0.25, 0.5)$, and $\bar{j} = (-0.5, 0, 0.5)$.

We found that the particle spin leads to the divergence of the effective potential at the center of the rotating boson star for the test particle with zero orbital angular momentum. We studied the relations between the circular orbit parameters and the particle spin, and showed that the particle spin can increase or decrease the radius and energy of the circular orbit. There is a novel feature for the radial effective potential of the spinning test particle, i.e., when the spin and the total angular momentum of the test particle satisfy $\bar{s} + \bar{j} = 0$, the effective potential at the origin is still finite and the test particle can pass through the center of the rotating boson star. These results will lead to some novel orbits and have an important application in testing the gravitational waves in the boson star background, which will be studied in the near future.

V. ACKNOWLEDGMENTS

This work was supported in part by the National Key Research and Development Program of China (Grant No. 2020YFC2201503), the National Natural Science Foundation of China (Grants No. 12105126, No. 11875151, No. 12075103, and No. 12047501), the China Postdoctoral Science Foundation (Grant No. 2021M701531), the 111 Project under (Grant No. B20063), the Fundamental Research Funds for the Central Universities (Grant No. lzujbky-2021-pd08), “Lanzhou City’s scientific research funding subsidy to Lanzhou University”.

- [1] D. A. Feinblum and W. A. McKinley, Phys. Rev. **168**, 1445 (1968).
 [2] David J. Kaup, Phys. Rev. **172**, 1331 (1968).

- [3] R. Ruffini and S. Bonazzola, Phys. Rev. **187**, 1767 (1969).
 [4] B. W. Lynn, Nucl. Phys. B **321** (1989), 465.




FIG. 2: The relations between the circular orbit parameters and the test particle spin. The parameters ω and k is taken as $\omega = 0.89$ and $k = (1, 2, 3)$ for subfigures (a1-a3), (b1-b3), (c1-c3), respectively.




FIG. 3: The effective potentials of the spinning test particle with $\bar{j} + \bar{s} = 0$. From top line to bottom line, the corresponding spin and orbital angular momentum are taken as $(\bar{s}, \bar{j}) = (-0.5, 0.5), (-0.25, 0.25), (0, 0), (0.25, -0.25), (0.5, -0.5)$.

- [5] F. E. Schunck and E. W. Mielke, Phys. Lett. A **249**, 389 (1998).
 [6] S. Yoshida and Y. Eriguchi, Phys. Rev. D **56**, 762 (1997).
 [7] D. Astefanesei and E. Radu, Nucl. Phys. B **665**, 594 (2003).
 [8] F. E. Schunck and E. W. Mielke, Class. Quant. Grav. **20**, R301 (2003).
 [9] S. L. Liebling and C. Palenzuela, Living Rev. Rel. **15**, 6 (2012).
 [10] P. Grandclément, C. Somé, and E.ourgoulhon, Phys. Rev. D **90**, 024068 (2014).
 [11] C. A. R. Herdeiro, A. M. Pombo, and E. Radu, Phys. Lett. B **773**, 654 (2017).
 [12] M. Alcubierre, J. Barranco, A. Bernal, J. C. Degollado, A. Diez-Tejedor, M. Megevand, D. Nunez, and O. Sarbach, Class. Quant. Grav. **35**, 19 (2018).
 [13] J. F. M. Delgado, C. A. R. Herdeiro, and E. Radu, JCAP **06**, 037 (2020).
 [14] M. Minamitsuji, Phys. Rev. D **97**, 104023 (2018).
 [15] D. Guerra, C. F. B. Macedo, and P. Pani, JCAP **09**, 061 (2019) [erratum: JCAP **06**, E01 (2020)].
 [16] H.-B. Li, Y.-B. Zeng, Y. Song, and Y.-Q. Wang, JHEP

- 04**, 042 (2021).
- [17] C. A. R. Herdeiro, J. Kunz, I. Perapechka, E. Radu, and Y. Shnir, *Phys. Lett. B* **812**, 136027 (2021).
- [18] Y.-B. Zeng, H.-B. Li, S.-X. Sun, S.-Y. Cui, and Y.-Q. Wang, [arXiv:2103.10717 [gr-qc]].
- [19] V. Dzhunushaliev and V. Folomeev, *Phys. Rev. D* **104**, 104024 (2021).
- [20] R. Brito, V. Cardoso, C. A. R. Herdeiro, and E. Radu, *Phys. Lett. B* **752**, 291 (2016).
- [21] E. W. Mielke and R. Scherzer, *Phys. Rev. D* **24**, 2111 (1981).
- [22] Y.-Q. Wang, Y.-X. Liu, and S.-W. Wei, *Phys. Rev. D* **99**, 064036 (2019).
- [23] M. Colpi, S. L. Shapiro, and I. Wasserman, *Phys. Rev. Lett.* **57**, 2485 (1986).
- [24] F. S. Guzman and J. M. Rueda-Becerril, *Phys. Rev. D* **80** (2009), 084023.
- [25] A. R. Liddle and M. S. Madsen, *Int. J. Mod. Phys. D* **1**, 101 (1992).
- [26] T. D. Lee and Y. Pang, *Phys. Rep.* **222**, 251 (1992).
- [27] P. Jetzer, *Phys. Rep.* **220**, 163 (1992).
- [28] J. Balakrishna, R. Bondarescu, G. Daues, F. Siddhartha Guzman, and E. Seidel, *Class. Quant. Grav.* **23**, 2631 (2006).
- [29] C. Palenzuela, I. Olabarrieta, L. Lehner, and S. L. Liebling, *Phys. Rev. D* **75**, 064005 (2007).
- [30] N. Sanchis-Gual, C. Herdeiro, J. A. Font, E. Radu, and F. Di Giovanni, *Phys. Rev. D* **99**, 024017 (2019).
- [31] J. C. Bustillo, N. Sanchis-Gual, A. Torres-Forné, J. A. Font, Avi Vajpeyi, R. Smith, C. A. R. Herdeiro, E. Radu, and S. H. W. Leong, *Phys. Rev. Lett.* **126**, 081101 (2021).
- [32] R. Abbott *et al.* [LIGO Scientific and Virgo], *Phys. Rev. Lett.* **125**, 101102 (2020).
- [33] P. Grandclément, *Phys. Rev. D* **95**, 084011 (2017).
- [34] M. Grould, Z. Meliani, F. H. Vincent, P. Grandclément, and E.ourgoulhon, *Class. Quant. Grav.* **34**, 215007 (2017).
- [35] L. G. Collodel, B. Kleihaus, and J. Kunz, *Phys. Rev. Lett.* **120**, 201103 (2018).
- [36] Y.-P. Zhang, Y.-B. Zeng, Y.-Q. Wang, S.-W. Wei, and Y.-X. Liu, [arXiv:2107.04848 [gr-qc]].
- [37] M. Mathisson, *Acta Phys. Pol.* **6**, 163 (1937).
- [38] A. Papapetrou, *Proc. Roy. Soc. Lond. A* **209**, 248 (1951).
- [39] E. Corinaldesi and A. Papapetrou, *Proc. Roy. Soc. Lond. A* **209**, 259 (1951).
- [40] W. G. Dixon, *Proc. Roy. Soc. Lond. A* **314**, 499 (1970).
- [41] F. E. Schunck and E. W. Mielke, *Class. Quant. Grav.* **20**, R301 (2003).
- [42] C. A. R. Herdeiro and E. Radu, *Class. Quant. Grav.* **32**, 144001 (2015).
- [43] P. Grandclément, *J. Comp. Physics* **229**, 3334 (2010).
- [44] R. M. Wald, *Phys. Rev. D* **6**, 406 (1972).
- [45] S. A. Hojman, PhD thesis, Princeton University (1975) (unpublished).
- [46] G. Lukes-Gerakopoulos, J. Seyrich, and D. Kunst, *Phys. Rev. D* **90**, 104019 (2014).
- [47] L. Filipe O. Costa, G. Lukes-Gerakopoulos, and Oldřich Semerák, *Phys. Rev. D* **97**, 084023 (2018).
- [48] G. Lukes-Gerakopoulos, *Phys. Rev. D* **96**, 104023 (2017).
- [49] A. A. Deriglazov and W. Guzmán Ramírez, *Adv. High Energy Phys.* **2016**, 1376016 (2016).
- [50] A. A. Deriglazov and W. Guzmán Ramírez, *Adv. Math. Phys.* **2017**, 7397159 (2017).
- [51] A. A. Deriglazov and W. Guzmán Ramírez, *Phys. Lett. B* **779**, 210 (2018).
- [52] W. Tulczyjew, *Acta Phys. Pol.* **18**, 393 (1959).
- [53] R. Hojman and S. Hojman, *Phys. Rev. D* **15**, 2724 (1977).
- [54] S. A. Hojman and F. A. Asenjo, *Class. Quant. Grav.* **30**, 025008 (2013).
- [55] Y.-P. Zhang, B.-M. Gu, S.-W. Wei, J. Yang, and Y.-X. Liu, *Phys. Rev. D* **94**, 124017 (2016).
- [56] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (Freeman, New York, 1973), p. 911.