

Cosmological constant problem on the horizon

Hassan Firouzjahi^{1*},

¹ *School of Astronomy, Institute for Research in Fundamental Sciences (IPM)*

P. O. Box 19395-5531, Tehran, Iran

Abstract

We revisit the quantum cosmological constant problem and highlight the important roles played by the dS horizon of zero point energy. We argue that fields which are light enough to have dS horizon of zero point energy comparable to the FLRW Hubble radius are the main contributor to dark energy. On the other hand, the zero point energy of heavy fields develop nonlinearities on sub-Hubble scales and can not contribute to dark energy. Our proposal provides a simple resolution for both the old and new cosmological constant problems by noting that there exists a field, the (lightest) neutrino, which happens to have a mass comparable to the present background photon temperature. The natures of dark energy and dark matter are unified in this proposal in which the zero point energy of light fields are the source of dark energy while dark matter is sourced by the zero point energy of heavy fields. The proposal predicts multiple transient periods of dark energy in early and late expansion history of the universe yielding to a higher value of the current Hubble expansion rate which can resolve the H_0 tension problem.

*firouz@ipm.ir

1 Introduction

The Λ CDM model has emerged as a successful phenomenological model, explaining the dynamics of the evolution of the cosmos with a handful of free parameters [1, 2]. Among the key unknown ingredients are dark matter and dark energy. There are hopes that the former may be addressed one way or another by new physics beyond the Standard Model (SM) of particle physics while the latter has proven more elusive.

There have been numerous attempts to address the nature of dark energy, for a review see [3–5] and references therein. One simple and natural possibility is that the dark energy is simply a cosmological constant term, like what Einstein initially introduced to construct a seemingly static universe. As is well known, the cosmological constant is associated with big problems in theoretical physics, for a classic review see [6]. As we shall review below, it is expected that the cosmological constant receives contributions from the zero point energy of all fields in SM. It is usually argued that the zero point energy density is at the order M^4 in which M is the UV cutoff of theory. The old cosmological constant problem is why it is not so large as expected from the typical energy scale of the high energy physics, like $(\text{TeV})^4$ or M_P^4 in which $M_P \sim 10^{18}\text{GeV}$ is the reduced Planck mass related to the Newton constant G via $M_P^2 = 1/8\pi G$. The new cosmological constant problem is why it is comparable to the energy density of matter at the current epoch in cosmic history, entering to the dynamics of the expansion of the universe at redshift around $z_\Lambda \sim 0.3$.

A number of approaches were proposed to address the cosmological constant problem(s). One approach is to employ the fundamental symmetries, like supersymmetry. In supersymmetric theories there is a one to one relation between the number of bosons and fermions which keep the value of cosmological constant zero. Of course, the Nature is not seen to be supersymmetric on energy scale below TeV so the supersymmetry can not explain the smallness of dark energy. The other approach is based on ideas like quintessence [7, 8] in which the dark energy is dynamically evolving to is attractor value which is zero. However, these kind of models can not address the severe fine-tunings involved as one has to tune the model parameters so carefully to obtain a tiny value of dark energy at the current epoch. Another class of solutions are based on the self-tuning idea in the context of extra dimensions [9–13]. These solutions were not convincing either as the solutions either run into singularities or require fine-tuning of model parameters. A solution of different kind has been proposed based on anthropic principle in which the cosmological constant should choose the range of values to allow the gravitationally bound objects like galaxies to form to host star formation required for the existence of intelligent observer [6, 14].

In this paper we revisit the quantum cosmological constant problem. We emphasize on the important roles played by the scale of dS horizon associated with the zero point energy. In particular we compare the scale of its Hubble radius with the Hubble radius of the FLRW universe and argue that the conventional treatment of the quantum cosmological constant problem misses this important effect.

The rest of the paper is organized as follows. In Section 2 we review the cosmological constant problem and gather some basic formulas. In Section 3 we highlight the important

roles that the horizon scale of the patch of zero point energy play while in Section 4 we show why, unlike in conventional treatment, the heavy fields can not contribute to the cosmological constant. In Section 5 we calculate the zero point energy in dS background while in Section 6 we present various cosmological implications of our proposal. In Sections 7, 8 and 9 we study a toy model which can mimic the dynamics of the patches of zero point energy in a cosmological background followed by the Summary and Discussions in Section 10. The reader who is only interested in the main results of our proposal may skip Sections 7, 8 and 9.

2 The Cosmological Constant Problem

In this section we briefly review the cosmological constant problem. For more extensive review see Refs. [6] and [15].

Numerous observations confirm the detection of dark energy at a level comparable to the matter energy density today [2, 16, 17]. Therefore, the cosmological constant, if not exactly zero by some symmetry considerations, is as small as $\rho_v \sim (10^{-3}\text{eV})^4$. Now the big trouble with the cosmological constant is what mechanism keeps it so small to be consistent with cosmic evolution.

In its simplest form, the cosmological constant is associated with the problem that the zero point energy of the fields has quartic divergence with the UV energy scale. To see it more specifically, consider a real scalar field ϕ with the mass m described by the following simple Lagrangian in flat spacetime,

$$S = \int d^4x \left[-\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{m^2}{2} \phi^2 \right]. \quad (2.1)$$

The energy momentum tensor $T^{\mu\nu}$ is given by

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - \eta_{\mu\nu} \left[\frac{1}{2} g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi + \frac{m^2}{2} \phi^2 \right]. \quad (2.2)$$

In particular, the energy density ρ is given by

$$\rho = T_{00} = \mathcal{H} = \frac{\dot{\phi}^2}{2} + \frac{1}{2} \delta^{ij} \partial_i \phi \partial_j \phi + \frac{m^2}{2} \phi^2, \quad (2.3)$$

in which \mathcal{H} is the Hamiltonian density.

We expand the quantum field in terms of the creation and the annihilation operators in Fourier space,

$$\phi(\mathbf{x}) = \frac{1}{(2\pi)^{3/2}} \int \frac{d^3\mathbf{k}}{\sqrt{2\omega(k)}} \left[e^{ik \cdot x} a_{\mathbf{k}} + e^{-ik \cdot x} a_{\mathbf{k}}^\dagger \right], \quad (2.4)$$

in which $k^\mu = (\omega(k), \mathbf{k})$, $\omega(k) \equiv \sqrt{\mathbf{k}^2 + m^2}$ and $[a_{\mathbf{k}}, a_{\mathbf{q}}^\dagger] = (2\pi)^3 \delta^3(\mathbf{k} - \mathbf{q})$.

The effective energy density contributing to the Einstein field equation is $\langle 0 | \rho | 0 \rangle \equiv \langle \rho \rangle$ in which $\langle 0 | X | 0 \rangle$ means the quantum expectation value of the quantum operator X with respect

to the vacuum. Using the form of the operator ρ from Eq. (2.3) we obtain

$$\langle \rho \rangle = \frac{1}{2} \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \omega(k). \quad (2.5)$$

Performing the same calculation for the pressure p , we obtain [15]

$$\langle p \rangle = \left\langle \frac{1}{3} \perp^{\mu\nu} T_{\mu\nu} \right\rangle = \frac{1}{6} \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \frac{\mathbf{k}^2}{\omega(k)}, \quad (2.6)$$

in which $\perp^{\mu\nu} \equiv \eta^{\mu\nu} + u^\mu u^\nu$ is a projection operator and $u^\mu = (1, \mathbf{0})$ is the comoving four-velocity.

To perform the above integrals, one usually imposes a hard UV momentum cutoff $k \leq M$ and the integral in Eq. (2.5) is estimated as

$$\langle \rho \rangle \simeq \frac{M^4}{16\pi^2}. \quad (2.7)$$

This is the usual representation of the old cosmological constant problem. If one assumes $M \sim M_P$ then the above estimation yields a contribution to $\langle \rho_v \rangle$ larger than the observed value by some factor of 10^{120} . Of course, if one chooses a smaller UV energy scale, say the scale of SM particle physics $M \sim 10^2 \text{ GeV}$, then the discrepancy becomes less severe but still the theoretical estimation is larger by some 10^{56} orders of magnitude from the observed value.

The above analysis can be repeated for other fields, bosonic or fermionic, with the important distinction that for fermionic fields one obtains a negative contribution in $\langle \rho \rangle$. This is because the fermionic fields, unlike the bosonic fields, anti commute with each other.

However, as pointed out in [15, 18–21], the hard momentum cutoff implemented above is problematic as it does not respect the Lorentz invariance of the setup. In other words the hard cutoff only respects the $O(3)$ invariance over the three-dimensional momentum space while breaking the four-dimensional $O(1, 3)$ Lorentz invariance. To see the fatal problem with this naive hard momentum cutoff regularization, note that to leading order we obtain $\langle p \rangle \simeq M^4/(3 \times 16\pi^2)$ which means that $\langle p \rangle/\langle \rho \rangle \simeq 1/3$. On the other hand, the Lorentz invariance of the vacuum requires that

$$\langle T_{\mu\nu} \rangle = -\rho_{\text{vac}} \eta_{\mu\nu}, \quad (2.8)$$

which immediately enforces $\langle p \rangle = -\langle \rho \rangle$. To bypass this problem one has to employ a regularization scheme which respects the underlying Lorentz invariance. A good candidate is the dimensional regularization approach which keeps the four dimensional Lorentz invariance intact.

Performing the integral using dimensional regularization approach (or any Lorentz invariant scheme) for the massive real scalar field one obtains [15, 18–21]

$$\langle \rho \rangle = -\langle p \rangle = \frac{m^4}{64\pi^2} \ln\left(\frac{m^2}{\mu^2}\right), \quad (2.9)$$

in which μ is the renormalization scale.

There are a few comments in order concerning Eq. (2.9). First, the requirement of Lorentz invariance $\langle p \rangle = -\langle \rho \rangle$ is explicit. Second, the massless fields such as gravitons, photons or gluons do not contribute to the energy density of vacuum. This may be understood by noting that for massless particles the equation of state is simply $p = \rho/3$ so the requirement of Lorentz invariance $\langle p \rangle = -\langle \rho \rangle$ can be met only if $\langle \rho \rangle = 0$. Third, depending on the renormalization scale, the vacuum energy density can be either positive (dS spacetime) or negative (AdS spacetime). Again, this is unlike the conclusion based on hard momentum cutoff prescription in which the bosonic (fermionic) fields always yield positive (negative) contributions to $\langle \rho \rangle$. Finally, there is no quartic dependence on the cutoff scale. This can reduce the naive fine-tuning of order 10^{-120} by many orders of magnitudes. As a rough estimate, for the heavy SM fields with mass at the order 10^2 GeV we obtain $|\langle \rho \rangle| \sim 10^{44} \text{ eV}^4$ which is much smaller than the naive estimation $\langle \rho \rangle \sim M_P^4$. However, it is still vastly larger than the observed value of the vacuum energy density. In addition, note that the contribution from the (lightest) neutrino with $m_\nu \sim 10^{-2} \text{ eV}$ is $\langle \rho \rangle \sim m_\nu^4 \sim (10^{-2} \text{ eV})^4$ which is at the same order as the observed value of dark energy. As we shall see this is not an accident and will be part of our solution for the cosmological constant problem!

Combining the contributions from all fields, bosonic and fermionic, the total contribution in vacuum zero point energy is obtained to be [15, 19]

$$\langle \rho \rangle = \sum_i n_i \frac{m_i^4}{64\pi^2} \ln\left(\frac{m_i^2}{\mu^2}\right), \quad (2.10)$$

in which n_i represents the degree of freedom (polarizations) of each field. For example, for a real scalar field $n = 1$, for a massive vector field $n = 3$ while for a Dirac fermion $n = -4$.

3 The Question of dS Horizon

To calculate the vacuum energy density we have assumed a flat background yielding to Eq. (2.9). This is perfectly fine as the energy density is a local quantity and is not sensitive to the large scale properties of the cosmological background. Indeed, the combination of Lorentz invariance and the equivalence principle guarantee that the flat spacetime approximation to calculate the vacuum energy density as given in (2.9) is valid. However, there is a subtle issue that was not taken into account when applying the estimated zero point energy to cosmology. In the presence of gravity, i.e. when the Newton constant G is turned on, associated to each positive vacuum energy there is a horizon radius which controls the causal properties of the corresponding dS spacetime. However, in the usual treatment of cosmological constant problem, it is assumed that the *entire* observable universe with the current Hubble radius H_0^{-1} is *simultaneously* endowed with a vacuum energy density ρ_v given by (2.9). This is equivalent to assuming that the quantum zero point energy fills the entire observable universe in a single patch of size H_0^{-1} . This is too much to ask!

The correct way of thinking about this problem is to look at the Hubble radius associated to Eq. (2.10) for each field. Let us denote $\rho_v(m)$ and $H_{(m)}$ respectively as the vacuum energy density and the Hubble rate of dS spacetime created from the zero point energy of each field with the mass m . Then we have $3M_P^2 H_{(m)}^2 = \rho_v(m) \sim m^4$, yielding

$$H_{(m)} \simeq \frac{m^2}{M_P}. \quad (3.1)$$

This is one important relation to keep in mind. For example, for electron field with $m_e \sim \text{MeV}$, we obtain $H_{(m_e)} \sim 10^{-15} \text{eV}$ yielding the Hubble radius $H_{(m_e)}^{-1} \sim 10^9 m$. Logically, this is the largest dS patch which the zero point energy associated with the electron field can cover coherently. On the other hand the current Hubble radius is $H_0^{-1} \sim \text{Gpc} \sim 10^{26} m$. From the above simple estimation we conclude that the observable FLRW universe encompasses as many as $(H_0^{-1}/H_{(m_e)}^{-1})^3 \sim 10^{51}$ independent patches with the size $H_{(m_e)}^{-1}$. Demanding that as many as 10^{51} dS patches of the electron zero point energy to coherently cover the entire observable universe is too strong a condition to be realistic. Considering fields heavier than electron, the situation becomes even worse, i.e, the ratio $H_0^{-1}/H_{(m)}^{-1}$ becomes much larger.

The picture which emerges from the above discussions is as follows. For heavy fields, the Hubble radius associated with a dS spacetime generated from the zero point energy is much smaller than the Hubble radius of the observable universe, $H_{(m)}^{-1} \ll H_0^{-1}$. Therefore, one needs as many as

$$N_{\text{patches}} \sim \left(\frac{H_{(m)}}{H_0} \right)^3 \sim \left(\frac{m}{10^{-2} \text{eV}} \right)^6 \quad (3.2)$$

tiny dS patches to cover the current observable universe. As these tiny patches are created quantum mechanically they are uncorrelated so they can not provide a coherent energy density to be the origin of the observed cosmological constant (or dark energy). This also provides an interesting resolution for the cosmological constant problem: considering the light fields. Specifically, very light fields have much smaller zero point energy but at the same time they have a very large Hubble radius which can encompass the entire observable universe in a single dS patch. From equation (3.2) we see that the (lightest) neutrino with the mass $m \sim 10^{-2} \text{eV}$ has just the right scale to address the cosmological constant problem in which $H_0^{-1} \sim H_{(m_\nu)}^{-1}$. In this view the entire observable universe currently is within a single dS patch created by the zero point energy of the light neutrino field with the vacuum energy density $(10^{-3} \text{eV})^4$, see the left panel of Fig. 1 for a schematic view. Correspondingly, the observed vacuum energy density today with the fractional energy density [2] $\Omega_\Lambda^{(0)} \sim 0.7$ is sourced by the zero point energy of the lightest neutrino.

In the case of heavy fields, as the huge number of dS patches in Eq. (3.2) are uncorrelated, one expects that the full volume between these regions to be highly inhomogeneous, for a schematic view see the right panel of Fig. 1. Consequently one expects that the variance in the energy density to be large, yielding to a large density contrast $\frac{\delta \rho_v(m)}{\rho_v(m)} \sim 1$, so the spacetime within these regions will collapse to black holes. In the next Section we calculate the variance and the density contrast of the vacuum zero point energy which confirm the




Figure 1: A schematic view comparing the FLRW Hubble radius H_F^{-1} with the dS horizon of size $H_{(m)}^{-1}$ associated with the zero point energy of the field with the mass m . Left: this is the case when the field is light so the FLRW horizon is within a single dS patch of zero point energy. Right: this represents the case when the field is heavy so its dS horizon is much smaller than the FLRW horizon such that a FLRW Hubble patch encompasses many small patches of zero point energy.

above conclusion. In addition, we calculate the correlation length associated with the zero point energy and show that the correlation length is at the order of Compton radius m^{-1} .

However, before calculating the variance of the vacuum energy density, let us check the accumulated energy associated with the modes which are outside the correlation length, i.e. the modes which have crossed the quantum Compton radius with $k \leq m$. Denoting this accumulated energy by $\delta\rho_C(m)$ we obtain

$$\delta\rho_C(m) = \frac{1}{2} \int_0^m \frac{d^3\mathbf{k}}{(2\pi)^3} \sqrt{k^2 + m^2} = \frac{m^4}{32\pi^2} \left[3\sqrt{2} - \ln(1 + \sqrt{2}) \right] \sim \frac{6m^4}{64\pi^2}. \quad (3.3)$$

Now we can compare the above value of the accumulated energy density of the “long modes” with the supposedly background vacuum energy density $\rho_v(m)$ given in Eq. (2.9). Assuming that there is no exponential hierarchy between m and the renormalization scale μ we obtain

$$\frac{\delta\rho_C(m)}{\rho_v(m)} \sim 1. \quad (3.4)$$

We conclude that the accumulated energy density of the modes which are outside the correlation length m^{-1} is comparable to the background vacuum energy density. This rings the bell

that the background space can be highly inhomogeneous. As such, one expects the nearby dS patches to collapse into black holes so they can not provide a coherent background for the large scale universe.

4 Density Contrast and Correlation Length

The quantum average of the vacuum zero point energy density $\langle \rho_v \rangle = \langle \mathcal{H} \rangle$ for the real scalar field is given in Eq. (2.9). Now for the moment suppose we neglect the contribution from the FLRW matter and radiation energy density and assume that ρ_v is the dominant energy density in cosmic expansion. In order for $\langle \rho_v \rangle$ to be viewed as a viable background energy density for the cosmic evolution we have to make sure that the variance in energy density is small, yielding to a small density contrast $\delta \rho_v / \langle \rho_v \rangle \ll 1$. Otherwise, a region with large density contrast $\delta \rho_v / \langle \rho_v \rangle \sim 1$ develops inhomogeneities collapsing to black holes.

In this section we calculate explicitly the variance $\delta \rho_v^2 \equiv \langle \rho_v^2 \rangle - \langle \rho_v \rangle^2$ and the density contrast $\delta \rho_v / \langle \rho_v \rangle$. First we present the analysis for the case of a real scalar field and then go to the case of a Dirac fermion field.

4.1 Real Scalar Field

First we consider the case of a real scalar field with the action (2.1). Let us denote the three different contributions to energy density of the scalar field in Eq. (2.3) by ρ_1, ρ_2 and ρ_3 as follows,

$$\rho_1 \equiv \frac{\dot{\phi}^2}{2}, \quad \rho_2 \equiv \frac{1}{2} \delta^{ij} \partial_i \phi \partial_j \phi, \quad \rho_3 \equiv \frac{m^2}{2} \phi^2. \quad (4.1)$$

To calculate $\langle \rho_i \rangle$ we go to Fourier space and expand the field in terms of the mode functions as in Eq. (2.4). Performing the momentum integrals using the dimensional regularization scheme to regularize the UV divergences (for a sample analysis of dimensional regularization see Section 5) we obtain

$$\langle \rho_1 \rangle = \frac{1}{2} \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{\omega(q)}{2} = \frac{m^4}{128\pi^2} \ln \left(\frac{m^2}{\mu^2} \right), \quad (4.2)$$

$$\langle \rho_2 \rangle = \frac{1}{2} \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{\mathbf{q}^2}{2\omega(q)} = \frac{-3m^4}{128\pi^2} \ln \left(\frac{m^2}{\mu^2} \right), \quad (4.3)$$

$$\langle \rho_3 \rangle = \frac{1}{2} \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{m^2}{2\omega(q)} = \frac{4m^4}{128\pi^2} \ln \left(\frac{m^2}{\mu^2} \right). \quad (4.4)$$

Curiously, note that while all three operators ρ_1, ρ_2 and ρ_3 are positive definite and Hermitian operators but the expectation value of ρ_2 has opposite sign compared to those of ρ_1 and ρ_3 . Also note that $\langle \rho_1 \rangle = \langle \rho_2 \rangle + \langle \rho_3 \rangle$ so

$$\langle \rho \rangle = \langle \rho_1 \rangle + \langle \rho_2 \rangle + \langle \rho_3 \rangle = 2(\langle \rho_2 \rangle + \langle \rho_3 \rangle) = 2\langle \rho_1 \rangle. \quad (4.5)$$

Now to calculate $\delta\rho^2$ we have

$$\begin{aligned}\delta\rho^2 &= \langle\rho_1^2\rangle + \langle\rho_2^2\rangle + \langle\rho_3^2\rangle + \langle\rho_1\rho_2\rangle + \langle\rho_2\rho_1\rangle + \langle\rho_1\rho_3\rangle + \langle\rho_3\rho_1\rangle + \langle\rho_2\rho_3\rangle + \langle\rho_3\rho_2\rangle \\ &- \left(\langle\rho_1\rangle^2 + \langle\rho_2\rangle^2 + \langle\rho_3\rangle^2\right) - 2\langle\rho_1\rangle\langle\rho_2\rangle - 2\langle\rho_1\rangle\langle\rho_3\rangle - 2\langle\rho_2\rangle\langle\rho_3\rangle.\end{aligned}\quad (4.6)$$

Note that since $\dot{\phi}$ and ϕ do not commute we have considered all possible orderings of the operators. However, as we shall see, these orderings do not matter.

First we note that ϕ is a free Gaussian field so using the standard Wick theorem (or by direct calculations) one can show that

$$\langle\rho_i^2\rangle = 3\langle\rho_i\rangle^2, \quad i = 1, 3. \quad (4.7)$$

However, the calculation of $\langle\rho_2^2\rangle$ is somewhat tricky. Performing various contractions we obtain

$$\begin{aligned}\langle\rho_2^2\rangle &= \langle\rho_2\rangle^2 + (2)\frac{1}{4}\int\frac{d^3\mathbf{q}_1}{(2\pi)^3 2\omega(q_1)}\frac{d^3\mathbf{q}_2}{(2\pi)^3 2\omega(q_2)}(\mathbf{q}_1\cdot\mathbf{q}_2)^2 \\ &= \langle\rho_2\rangle^2 + (2\times\frac{1}{3})\langle\rho_2\rangle^2,\end{aligned}\quad (4.8)$$

in which the factor $1/3$ originated from an integral of the form $\int_{-1}^{+1} d(\cos\theta)\cos^2\theta$.

For the mix correlations we calculate one case for illustration and the rest would be similar. For this purpose let us calculate $\langle\rho_1\rho_2\rangle$. To simplify the analysis note that due to translation invariance we can simply set $\mathbf{x} = 0$, obtaining

$$\begin{aligned}\langle\rho_1\rho_2\rangle &= \frac{i^4}{4}\int\frac{d^3\mathbf{q}_i}{(2\pi)^{\frac{3}{2}}}\frac{\omega(q_1)\omega(q_2)\mathbf{q}_3\cdot\mathbf{q}_4}{2\omega(q_i)}\langle 0|(-a_{\mathbf{q}_1}+a_{\mathbf{q}_1}^\dagger)(-a_{\mathbf{q}_2}+a_{\mathbf{q}_2}^\dagger)(a_{\mathbf{q}_3}-a_{\mathbf{q}_3}^\dagger)(a_{\mathbf{q}_4}-a_{\mathbf{q}_4}^\dagger)|0\rangle \\ &= \frac{1}{4}\int\frac{d^3\mathbf{q}_1}{(2\pi)^3 2\omega(q_1)}\frac{d^3\mathbf{q}_2}{(2\pi)^3 2\omega(q_2)}\left[2\omega(q_1)\omega(q_2)\mathbf{q}_1\cdot\mathbf{q}_2 + \omega(q_1)^2\mathbf{q}_2^2\right].\end{aligned}\quad (4.9)$$

On the other hand, due to rotation invariance the integral over $\mathbf{q}_1\cdot\mathbf{q}_2$ vanishes and we obtain

$$\langle\rho_1\rho_2\rangle = \frac{1}{4}\int\frac{d^3\mathbf{q}_1}{(2\pi)^3}\frac{d^3\mathbf{q}_2}{(2\pi)^3}\frac{\omega(q_1)}{2}\frac{\mathbf{q}_2^2}{2\omega(q_2)} = \langle\rho_1\rangle\langle\rho_2\rangle. \quad (4.10)$$

Similarly, we obtain

$$\langle\rho_2\rho_3\rangle = \langle\rho_2\rangle\langle\rho_3\rangle. \quad (4.11)$$

Since $\dot{\phi}$ and ϕ do not commute, the calculation for $\langle\rho_1\rho_3\rangle$ is a bit different, yielding

$$\langle\rho_1\rho_3\rangle = \langle\rho_1\rangle\langle\rho_3\rangle - \frac{m^2}{2}\left(\int\frac{d^3\mathbf{q}_1}{(2\pi)^3}\right)^2. \quad (4.12)$$

Regularizing the last term above which is UV divergent by absorbing into counter terms, we obtain

$$\langle\rho_1\rho_3\rangle = \langle\rho_1\rangle\langle\rho_3\rangle = \langle\rho_3\rho_1\rangle. \quad (4.13)$$

Combining all we obtain

$$\begin{aligned}\delta\rho^2 &= 2\left(\langle\rho_1\rangle^2 + \langle\rho_2\rangle^2 + \langle\rho_3\rangle^2\right) - \frac{4}{3}\langle\rho_2\rangle^2 \\ &= \langle\rho\rangle^2 - 4\langle\rho_2\rangle\langle\rho_3\rangle - \frac{4}{3}\langle\rho_2\rangle^2,\end{aligned}\tag{4.14}$$

where to obtain the final result the relation (4.5) has been used.

Using the specific value of $\langle\rho_2\rangle$ and $\langle\rho_3\rangle$ from Eqs. (4.3) and (4.4) and taking both signs after the square root we obtain

$$\frac{\delta\rho}{\langle\rho\rangle} = \pm\left(1 - 4\frac{\langle\rho_2\rangle\langle\rho_3\rangle}{\langle\rho\rangle^2} - \frac{4\langle\rho_2\rangle^2}{3\langle\rho\rangle^2}\right)^{\frac{1}{2}} = \pm\sqrt{10}.\tag{4.15}$$

This is an interesting result. As argued before, the energy density of the background constructed from the zero point energy is very inhomogeneous and the system may collapse to black holes. This is in agreement with the intuition that a large number of incoherent patches of zero point energy can not cover the entire observable universe. Also note that both signs are allowed so we can have either over-density or under-density. Specifically, the quantity $\langle\rho\rangle + \delta\rho$ can take either signs so some regions of spacetime can be in dS-type and some other parts AdS-type. The parts of the spacetime which are in AdS-type may collapse due to instabilities.

The above conclusion that $\langle\rho\rangle + \delta\rho$ can be negative is based on the hidden assumption that the statistical distributions of ρ is mostly symmetric around its average value $\langle\rho\rangle$ so a large variance can yield to a negative local value of $\langle\rho\rangle + \delta\rho$. This is certainly the case for a Gaussian distribution while for other distribution like Poisson distribution it may not be the case. Therefore, the nature of the distribution of ρ is an important question. In our setup with a large uncorrelated dS patches of heavy fields created quantum mechanically from the vacuum it may be reasonable to assume that ρ follows a Normal (i. e. Gaussian) distribution. But it would be interesting to calculate the distribution of ρ from the first principle. In the following analysis we assume a Gaussian-type distribution such that $\langle\rho\rangle + \delta\rho$ can be negative for a large variance.

In the above analysis we have neglected the conventional matter and radiation energy density of the FLRW universe, ρ_F . Now, let us look at the total energy density from the combination of the FLRW matter-radiation ρ_F and the vacuum energy density, defined via $\rho_T = \rho_F + \rho_v$ in which ρ_v is the vacuum energy density given in Eq. (2.9). Note that ρ_F is independent of the zero point energy so its quantum expectation is trivial, i.e. $\langle\rho_F\rangle = \rho_F$. Correspondingly, $\langle\rho_T\rangle = \rho_F + \langle\rho_v\rangle$ and $\delta\rho = \delta\rho_v$ in which $\delta\rho_v$ is as given in Eq. (4.14). Therefore, the total density contrast is given by

$$\frac{\delta\rho_T}{\langle\rho_T\rangle} = \frac{\delta\rho_v}{\rho_F + \langle\rho_v\rangle} = \pm\sqrt{10}\frac{\langle\rho_v\rangle}{\rho_F + \langle\rho_v\rangle}.\tag{4.16}$$

Demanding that the amplitude of the total density contrast to be small, $|\frac{\delta\rho_T}{\langle\rho_T\rangle}| < 1$, we obtain

$$\langle\rho_v\rangle < \left(\sqrt{10} - 1\right)^{-1} \rho_F \sim \frac{\rho_F}{2}.\tag{4.17}$$

This means that in order for the background consisting of matter-radiation plus zero point energy to be stable for cosmological expansion one requires that the vacuum energy should be less than the FLRW matter-radiation energy density.

Motivated by the above analysis, it would be instructive to calculate the fractional contrast in T_μ^μ as well. This is a relevant question as T_μ^μ is a four-dimensional scalar which is related to the curvature of spacetime. More specifically, from the Einstein field equation we obtain $M_P^2 R = -T_\mu^\mu$ in which R is the Ricci scalar. Therefore, a variance in T_μ^μ is related to a variance in the curvature of spacetime. Using the perfect fluid form of the stress energy tensor, we have $T_\mu^\mu = \rho + 3p$. On the other hand, one can check that

$$p_v = \rho_1 - \frac{\rho_2}{3} - \rho_3, \quad (4.18)$$

so for the vacuum, we have

$$T_\mu^\mu = 4\rho_1 - 2\rho_3. \quad (4.19)$$

Now, repeating the same steps as in the case of ρ , one can show that $\langle T_\mu^\mu \rangle = -2\langle \rho_v \rangle$ and $\langle (T_\mu^\mu)^2 \rangle = 44\langle \rho_v \rangle^2$. Combining these two results we obtain

$$\delta T^2 \equiv \langle (T_\mu^\mu)^2 \rangle - \langle T_\mu^\mu \rangle^2 = 40\langle \rho_v \rangle^2, \quad (4.20)$$

and

$$\frac{\delta T}{\langle T_\mu^\mu \rangle} = \pm \sqrt{10}. \quad (4.21)$$

We see that the local curvature of the spacetime fluctuates rapidly in which some regions are expected to become AdS type and collapse to form black holes.

Having calculated the variance and the density contrast, it is also instructive to calculate the correlation length of the energy density. The correlation length is related to the connected part of the correlation function as follows. The two point correlation function is given by $\langle \rho(\mathbf{x})\rho(\mathbf{y}) \rangle$. Due to translation invariance the correlation function depends only on $\mathbf{x} - \mathbf{y}$ so we simply set $\mathbf{y} = \mathbf{0}$. The connected part of the two point correlation function is given by

$$\langle \rho(\mathbf{x})\rho(\mathbf{0}) \rangle_c \equiv \langle \rho(\mathbf{x})\rho(\mathbf{0}) \rangle - \langle \rho^2 \rangle. \quad (4.22)$$

Performing the analysis similar to the case of variance one can calculate $\langle \rho(\mathbf{x})\rho(\mathbf{0}) \rangle_c$. The analysis is somewhat involved. However, what we are interested in is the correlation length ξ which controls the exponential fall off of the connected correlation function on large distance,

$$\langle \rho(\mathbf{x})\rho(\mathbf{0}) \rangle_c \rightarrow e^{-\frac{r}{\xi}}, \quad (r \rightarrow \infty). \quad (4.23)$$

To estimate ξ let us look at $\langle \rho_3(\mathbf{x})\rho_3(\mathbf{0}) \rangle_c$ which is easier. Upon performing the various contractions we obtain

$$\langle \rho_3(\mathbf{x})\rho_3(\mathbf{0}) \rangle_c = \frac{m^4}{2} \left(\int \frac{d^3 \mathbf{q}}{(2\pi)^3 2\omega(q)} e^{-i\mathbf{q} \cdot \mathbf{x}} \right)^2. \quad (4.24)$$

The above integral is well-known in QFT which yields [22]

$$\langle \rho_3(\mathbf{x}) \rho_3(\mathbf{0}) \rangle_c = \frac{m^8}{32\pi^4} \left(\frac{K_1(mr)}{mr} \right)^2, \quad (4.25)$$

in which $K_1(x)$ is the modified Bessel function. Using the asymptotic behaviour $K_1(x) \sim e^{-x}$ we obtain $\langle \rho_3(\mathbf{x}) \rho_3(\mathbf{0}) \rangle_c \sim e^{-2mr}$ yielding the correlation length $\xi = 1/(2m)$. This is somewhat expected as m is the only mass scale relevant for this free QFT. Calculating the two point correlation functions for other components of the energy density yield the same exponential behaviour for large r .

Now having the correlation length at hand, we can compare the Hubble radius of zero point energy $H_{(m)}^{-1}$ and ξ , obtaining

$$\frac{\xi}{H_{(m)}^{-1}} \sim \frac{H_{(m)}}{m} \sim \frac{m}{M_P} \ll 1. \quad (4.26)$$

This is inline with our intuitive argument in previous section. The correlation length of the zero point energy density is much smaller than the Hubble radius of each dS patch so the dS patches are practically uncorrelated. As such, one can not cover the entire FLRW horizon by a vast number of uncorrelated dS patches (like the right panel of Fig. 1) which are created quantum mechanically and yet expect them to behave as a uniform cosmological constant.

4.2 Dirac Fermion Field

Here we present the analysis of variance and density contrast for a Dirac fermion field Ψ . In what follows we follow the notations of Weinberg [22].

Expressing the mode function in Fourier space, we have

$$\Psi = \int \frac{d^3\mathbf{q}}{(2\pi)^{\frac{3}{2}}} \sum_{\sigma} \left[u(\mathbf{q}, \sigma) e^{iq \cdot x} a(\mathbf{q}, \sigma) + v(\mathbf{q}, \sigma) e^{-iq \cdot x} b^{\dagger}(\mathbf{q}, \sigma) \right], \quad (4.27)$$

in which $a(\mathbf{q}, \sigma)$ and $b(\mathbf{q}, \sigma)$ are the annihilation operators for the “particle” and “antiparticle”, $\sigma = \pm \frac{1}{2}$ are two spin polarizations while $u(\mathbf{q}, \sigma)$ and $v(\mathbf{q}, \sigma)$ are the corresponding mode functions. As usual, $a(\mathbf{q}, \sigma)$ and $b(\mathbf{q}, \sigma)$ are anti commuting operators in which $\{a(\mathbf{q}, \sigma), a^{\dagger}(\mathbf{q}', \sigma')\} = \delta_{\sigma\sigma'} \delta(\mathbf{q} - \mathbf{q}')$ and $\{b(\mathbf{q}, \sigma), b^{\dagger}(\mathbf{q}', \sigma')\} = \delta_{\sigma\sigma'} \delta(\mathbf{q} - \mathbf{q}')$ while the rest of anti commutators are zero.

The action of the field with the mass m is given by

$$S = - \int d^4x \bar{\Psi} (\gamma^{\mu} \partial_{\mu} + m) \Psi, \quad (4.28)$$

in which γ^{μ} are the Dirac matrices satisfying the anti commutation relation

$$\{\gamma^{\mu}, \gamma^{\nu}\} = 2\eta^{\mu\nu}, \quad (4.29)$$

while $\bar{\Psi} \equiv \Psi^{\dagger} \beta$ in which the matrix β is given by $\beta \equiv i\gamma^0$.

The mode functions $u(\mathbf{q}, \sigma)$ and $v(\mathbf{q}, \sigma)$ satisfy the following relations

$$\sum_{\sigma} u_{\mu}(\mathbf{q}, \sigma) \bar{u}_{\nu}(\mathbf{q}, \sigma) = \left(\frac{-i\gamma^{\mu} q_{\mu} + m}{2q^0} \right)_{\mu\nu}, \quad (4.30)$$

and

$$\sum_{\sigma} v_{\mu}(\mathbf{q}, \sigma) \bar{v}_{\nu}(\mathbf{q}, \sigma) = \left(\frac{-i\gamma^{\mu} q_{\mu} - m}{2q^0} \right)_{\mu\nu}, \quad (4.31)$$

in which $q^0 = \sqrt{\mathbf{q}^2 + m^2}$ is the energy of the particle. Furthermore, the following relation is useful as well:

$$\sum_{\sigma} \bar{v}(\mathbf{q}, \sigma) \gamma^0 v(\mathbf{q}, \sigma) = -2i. \quad (4.32)$$

Finally, the energy density $\rho = T_{00}$ is given by

$$\rho = -\frac{1}{2} \bar{\Psi} \gamma^0 \partial_0 \Psi + \frac{1}{2} \partial_0 \bar{\Psi} \gamma^0 \Psi. \quad (4.33)$$

It is convenient to define the above two terms via $\rho_1 \equiv -\frac{1}{2} \bar{\Psi} \gamma^0 \partial_0 \Psi$ and $\rho_2 \equiv \frac{1}{2} \partial_0 \bar{\Psi} \gamma^0 \Psi$ (these values of ρ_i should not be confused with ρ_i in the case of a real scalar field studied in previous subsection). One can check that $\rho_2 = \rho_1^{\dagger}$ so $\rho = \rho_1 + \rho_1^{\dagger}$. This shows, as expected, that ρ is a Hermitian operator.

As a warm up, it is helpful to calculate $\langle \rho \rangle$ from Eq. (4.33). We obtain $\langle \rho \rangle = \langle \rho_1 \rangle + \langle \rho_1 \rangle^*$, so we only need to calculate $\langle \rho_1 \rangle$. Using the decomposition of the field in terms of the mode functions in Eq. (4.27), we have

$$\begin{aligned} \langle \rho_1 \rangle &= -\frac{1}{2} \langle 0 | \bar{\Psi} \gamma^0 \partial_0 \Psi | 0 \rangle \\ &= -\frac{i}{2} \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \sum_{\sigma} \bar{v}(\mathbf{q}, \sigma) \gamma^0 v(\mathbf{q}, \sigma) q^0 \\ &= -\int \frac{d^3 \mathbf{q}}{(2\pi)^3} q^0. \end{aligned} \quad (4.34)$$

As a result,

$$\langle \rho \rangle = 2\langle \rho_1 \rangle = -4 \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{q^0}{2} = -4 \left[\frac{m^4}{64\pi^2} \ln \left(\frac{m^2}{\mu^2} \right) \right]. \quad (4.35)$$

The above result confirms the factor 4 since we have four independent degrees of freedom in the Dirac field. In addition, compared to the case of a real scalar field given in Eq. (2.5) we have an overall minus sign which is the hallmark of the fermionic field.

To calculate $\langle \rho^2 \rangle$ note that $\langle \rho_2 \rangle = \langle \rho_1 \rangle^*$, $\langle \rho_2^2 \rangle = \langle \rho_1^2 \rangle^*$ and $\langle \rho_1 \rho_2 \rangle = \langle \rho_1 \rho_2 \rangle^* = \langle \rho_2 \rho_1 \rangle$ which will be handy in the following analysis. Furthermore, in the analysis below we show that $\langle \rho_1^2 \rangle$ and $\langle \rho_1 \rho_2 \rangle$ are real, so using the above mentioned relations, we obtain

$$\langle \rho^2 \rangle = 2\langle \rho_1^2 \rangle + 2\langle \rho_1 \rho_2 \rangle. \quad (4.36)$$

We calculate each term in turn. Due to translation invariance we set $x^\mu = 0$ without loss of generality.

Starting with $\langle \rho_1^2 \rangle$ we have

$$\langle \rho_1^2 \rangle = \frac{1}{4} \langle 0 | (\bar{\Psi} \gamma^0 \partial_0 \Psi) (\bar{\Psi} \gamma^0 \partial_0 \Psi) | 0 \rangle. \quad (4.37)$$

Using the decomposition in terms of mode functions from Eq. (4.27) and performing all the possible contractions involving the creation and annihilation operators, we end up with two types of contributions in $\langle \rho_1^2 \rangle$. Denoting these contributions as term A and term B , we have $\langle \rho_1^2 \rangle \equiv A + B$ in which

$$A \equiv \frac{1}{4} \sum_{\sigma_1 \sigma_2} \int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \frac{d^3 \mathbf{q}_2}{(2\pi)^3} q_1^0 q_2^0 \bar{v}(\mathbf{q}_1, \sigma_1) \gamma^0 u(\mathbf{q}_2, \sigma_2) \bar{u}(\mathbf{q}_2, \sigma_2) \gamma^0 v(\mathbf{q}_1, \sigma_1), \quad (4.38)$$

and

$$B \equiv -\frac{1}{4} \sum_{\sigma_1 \sigma_2} \int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \frac{d^3 \mathbf{q}_2}{(2\pi)^3} q_1^0 q_2^0 \bar{v}(\mathbf{q}_1, \sigma_1) \gamma^0 v(\mathbf{q}_1, \sigma_1) \bar{v}(\mathbf{q}_2, \sigma_2) \gamma^0 v(\mathbf{q}_2, \sigma_2). \quad (4.39)$$

Using the relations (4.30) and (4.31) we obtain

$$\begin{aligned} A &= \frac{1}{4} \int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \frac{d^3 \mathbf{q}_2}{(2\pi)^3} (m^2 - q_1^0 q_2^0) \\ &= \frac{m^2}{4} \left(\int \frac{d^3 \mathbf{q}}{(2\pi)^3} \right)^2 - \frac{1}{4} \langle \rho_1 \rangle^2, \end{aligned} \quad (4.40)$$

where to obtain the last result, Eq. (4.35) for $\langle \rho_1 \rangle$ has been used.

Similarly, using the relation (4.32), for the B term we obtain

$$B = \left(\int \frac{d^3 \mathbf{q}}{(2\pi)^3} q^0 \right)^2 = \langle \rho_1 \rangle^2. \quad (4.41)$$

Combining the above two contributions, we obtain

$$\langle \rho_1^2 \rangle = A + B = \frac{m^2}{4} \left(\int \frac{d^3 \mathbf{q}}{(2\pi)^3} \right)^2 + \frac{3}{4} \langle \rho_1 \rangle^2. \quad (4.42)$$

As promised, $\langle \rho_1^2 \rangle$ is real so we do not need to calculate $\langle \rho_2^2 \rangle$ since $\langle \rho_2^2 \rangle = \langle \rho_1^2 \rangle^* = \langle \rho_1^2 \rangle$. Furthermore, as in the case of scalar field, the first term above is UV divergent and has to be absorbed into counter terms via regularization so after regularization we obtain $\langle \rho_1^2 \rangle \cong \frac{3}{4} \langle \rho_1 \rangle^2$.

Now we calculate $\langle \rho_1 \rho_2 \rangle$ which is

$$\langle \rho_1 \rho_2 \rangle = -\frac{1}{4} \langle 0 | (\bar{\Psi} \gamma^0 \partial_0 \Psi) (\partial_0 \bar{\Psi} \gamma^0 \Psi) | 0 \rangle. \quad (4.43)$$

Like in the previous case, after performing all contractions involving the creation and the annihilation operators we have two different contributions in $\langle \rho_1 \rho_2 \rangle$. Incidentally, one of this

contribution is the B term above while the other contribution is new, denoted by the C term so $\langle \rho_1 \rho_2 \rangle = B + C$ in which

$$C \equiv -\frac{1}{4} \sum_{\sigma_1 \sigma_2} \int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \frac{d^3 \mathbf{q}_2}{(2\pi)^3} (q_2^0)^2 \bar{v}(\mathbf{q}_1, \sigma_1) \gamma^0 u(\mathbf{q}_2, \sigma_2) \bar{u}(\mathbf{q}_2, \sigma_2) \gamma^0 v(\mathbf{q}_1, \sigma_1). \quad (4.44)$$

Note that while the C term looks very similar to the A term, but they are different as the latter has the product $q_1^0 q_2^0$ while the former has $(q_2^0)^2$ inside the double integrals.

Using the relations (4.30) and (4.31) we obtain

$$C = -\frac{m^2}{4} \left(\int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \frac{1}{q_1^0} \right) \left(\int \frac{d^3 \mathbf{q}_2}{(2\pi)^3} q_2^0 \right) + \frac{1}{4} \int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \int \frac{d^3 \mathbf{q}_2}{(2\pi)^3} (q_2^0)^2. \quad (4.45)$$

Combining the two contributions B and C , we obtain

$$\langle \rho_1 \rho_2 \rangle = \langle \rho_1 \rangle^2 + \frac{m^2}{4} \left(\int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{q^0} \right) \langle \rho_1 \rangle + \frac{1}{4} \int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \int \frac{d^3 \mathbf{q}_2}{(2\pi)^3} (q_2^0)^2. \quad (4.46)$$

The last term above is again UV divergent and has to be absorbed into counter terms so we keep the first two terms above in the regularized $\langle \rho_1 \rho_2 \rangle$.

Having calculated $\langle \rho_1^2 \rangle$ and $\langle \rho_1 \rho_2 \rangle$ and discarding the two UV divergent terms as discussed above, from Eq. (4.36) $\langle \rho^2 \rangle$ is calculated to be

$$\langle \rho^2 \rangle = \frac{7}{2} \langle \rho_1 \rangle^2 + \frac{m^2}{2} \left(\int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{q^0} \right) \langle \rho_1 \rangle. \quad (4.47)$$

On the other hand, the integral in the second term above is already calculated in the case of the real scalar field in Eq. (4.4) which, combined with Eq. (4.35), can be expressed in terms of $\langle \rho_1 \rangle$ as

$$\frac{m^2}{2} \left(\int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{q^0} \right) = -2 \langle \rho_1 \rangle. \quad (4.48)$$

Now combining all the terms, we obtain

$$\langle \rho^2 \rangle = \frac{3}{2} \langle \rho_1 \rangle^2 = \frac{3}{8} \langle \rho \rangle^2. \quad (4.49)$$

Finally, the variance $\delta \rho^2 = \langle \rho^2 \rangle - \langle \rho \rangle^2$ is obtained to be

$$\delta \rho^2 = -\frac{10}{16} \langle \rho \rangle^2. \quad (4.50)$$

Surprisingly, the variance is negative! This may look absurd. However, we already encountered seemingly negative value for the expectation values of positive definite operators, such as the expectation value of $\langle \rho_2 \rangle$ in Eq. (4.3) for the case of real scalar field. This phenomena may be attributed to regularization which absorbs positive infinity terms leaving the finite possible negative terms.

The amplitude of the zero point density perturbation is obtained to be

$$\left| \frac{\delta\rho_v}{\rho_v} \right| = \frac{\sqrt{10}}{4}, \quad (4.51)$$

in which by $\left| \frac{\delta\rho_v}{\rho_v} \right|$ we mean the amplitude with the minus sign absorbed. Curiously, the factor $\sqrt{10}$ is common as in the case of a real scalar field, see Eq. (4.15). The factor $\frac{1}{4}$ may be attributed to the four degrees of freedom encoded in Dirac fermion field. Furthermore, calculating the variance of T_μ^μ we obtain $\left| \frac{\delta T_\mu^\mu}{\langle T_\mu^\mu \rangle} \right| = \frac{\sqrt{10}}{4}$.

As in the case of scalar field we see that the amplitude of density contrast associated to the zero point energy of the fermionic field is at the order unity. As before, we conclude that the vacuum zero point energy by itself can not be the dominant source of the cosmic energy density. One requires a dominant (classical) source of energy density like the matter-radiation energy density ρ_F with $\rho_F > \rho_v$ at each stage in cosmic expansion history to obtain a stable cosmological background.

Finally, calculating the correlation length ξ of the zero point energy, one can check that, as in the case of real scalar field, $\xi \sim m^{-1}$ as expected.

5 Zero Point Energy in dS Background

The vacuum zero point energy density Eq. (2.9) is calculated assuming a flat background. As discussed before, one expects this result to hold true for a curved background as well. This is because general relativity is a local theory and the combination of the Lorentz invariance and the equivalence principle guarantees the validity of Eq. (2.9) even in a curved field space. Having said this, it is a non-trivial exercise to see how this works in practice. Specifically, the accumulated zero point energy yielding to Eq. (2.9) generate a dS background. At the same time, one has to solve the mode function $\phi(x^\mu)$ in a dS spacetime and then construct $\langle T_{00} \rangle$. Technically speaking, it seems non-trivial how Eq. (2.9) emerges as the final result. This is more intriguing for a dS spacetime noting that a dS spacetime is characterized by the size of its horizon, $H_{(m)}^{-1}$ in our notation, while Eq. (2.9) is indifferent to this length scale.

In this Section we solve the mode function for a real scalar field in a dS background and calculate the vacuum zero point energy. To our knowledge the current analysis is new, however see also Refs. [15, 23, 24] and [25] (Chapter 6) who studied the analysis of zero point energy in a curved background.

To treat the regularization properly, we employ dimensional regularization for a free scalar field in a curved field space with the spacetime dimension d , with the action

$$S = \int d^d x \sqrt{-g} \left[-\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{m^2}{2} \phi^2 \right]. \quad (5.1)$$

The background metric is given by a FLRW type metric in d dimension,

$$ds^2 = -dt^2 + a(t)^2 d\mathbf{x}^2, \quad (5.2)$$

in which $a(t)$ is the scale factor.

Using the conformal time $d\tau = dt/a(t)$ and defining the canonically normalized field σ via $\sigma \equiv a^{\frac{d-2}{2}} \phi$, the action takes the following form

$$S = \frac{1}{2} \int d^d x \left[\sigma'^2 + \frac{(d-4)(d-2)}{4} \left(\frac{a'}{a} \right)^2 + \frac{d-2}{2} \frac{a''}{a} - m^2 a^2 \right], \quad (5.3)$$

in which a prime denotes the derivative with respect to the conformal time.

Going to the Fourier space and expanding the field σ in terms of the creation and annihilation operators we have

$$\sigma(x^\mu) = \int \frac{d^{d-1} \mathbf{k}}{(2\pi)^{(d-1)/2}} \left(\sigma_k(\tau) e^{i\mathbf{k} \cdot \mathbf{x}} a_{\mathbf{k}} + \sigma_k(\tau)^* e^{-i\mathbf{k} \cdot \mathbf{x}} a_{\mathbf{k}}^\dagger \right). \quad (5.4)$$

First we have to solve the mode function $\sigma_k(\tau)$ in a dS spacetime. Considering a dS spacetime with the Hubble expansion rate H , we have $aH\tau = -1$ where we have taken $-\infty < \tau < 0$. Note that since there is no rolling scalar field (i.e. there is no inflaton field), then the dS symmetry is exact and in particular $\dot{H} = 0$, yielding to $a' = Ha^2$ and $a'' = 2H^2 a^3$. Then the equation of motion for the mode function $\sigma_k(\tau)$ is given by

$$\sigma_k'' + \left[k^2 + \frac{m^2}{H^2 \tau^2} - \frac{(d-2)^2}{2\tau^2} \right] \sigma_k = 0. \quad (5.5)$$

As a check note that if we set $d = 4$, then the last term in the bracket above yields the well known contribution $-2/\tau^2$ which is the hallmark of gravitational particle production in a dS background for the light scalar field perturbations.

If we expect Eq. (2.9) to hold true, then $H \sim m^2/M_P$ and therefore $H/m \sim m/M_P \ll 1$. Therefore, we can safely ignore the last term in Eq. (5.5) compared to the mass term. Imposing the Bunch-Davies (Minkowski) vacuum for the modes deep inside the dS horizon $k\tau \rightarrow -\infty$,

$$\sigma_k = \frac{1}{\sqrt{2k}} e^{-ik\tau}, \quad (5.6)$$

the solution of the mode function is given in terms of the Hankel function of the first type as follows

$$\phi_k(t) = a^{\frac{2-d}{2}} \sigma_k = (-H\tau)^{\frac{d-1}{2}} \left(\frac{\pi}{4H} \right)^{\frac{1}{2}} e^{-\frac{\pi}{2}\nu} H_{i\nu}^{(1)}(-k\tau), \quad (5.7)$$

where

$$\nu \equiv \sqrt{\frac{m^2}{H^2} - 1} \simeq \frac{m}{H}. \quad (5.8)$$

Having calculated the mode function $\phi_k(\tau)$ we can calculate the vacuum energy density as follows

$$\langle \rho \rangle = \frac{\mu^{4-d}}{2a^2} \int \frac{d^{d-1} \mathbf{k}}{(2\pi)^{d-1}} \left[|\phi_k'(\tau)|^2 + (k^2 + m^2 a^2) |\phi_k(\tau)|^2 \right], \quad (5.9)$$

in which μ is a mass scale to properly take care of the energy dimension.

Defining the dimensionless variable $x \equiv -k\tau$, and separating the angular and the radial contributions of the integral, we obtain

$$\langle \rho \rangle = \frac{\pi \mu^{4-d}}{8(2\pi)^{d-1}} e^{-\pi\nu} H^d \left(\int d^{d-2}\Omega \int_0^\infty x^d \left[\left| \frac{d}{dx} H_{i\nu}^{(1)}(x) \right|^2 + \left(1 + \frac{\nu^2}{x^2}\right) \left| H_{i\nu}^{(1)}(x) \right|^2 \right] \right). \quad (5.10)$$

In deriving the above expression, we have neglected the time derivative of the prefactor $(-H\tau)^{(d-1)/2}$ in the mode function Eq. (5.7). One can check that this is a very good approximation in the limit of interest here where $\nu \gg 1$. The fractional errors induced in the final result is smaller by a factor $1/\nu^2 \ll 1$.

The integral over the azimuthal directions in Eq. (5.10) is given by [15]

$$\int d^{d-2}\Omega = \frac{2\pi^{(d-1)/2}}{\Gamma(\frac{d}{2} - \frac{1}{2})}, \quad (5.11)$$

where $\Gamma(x)$ is the gamma function.

The final step is to perform the following integral

$$I \equiv \int_0^\infty x^d \left[\left| \frac{d}{dx} H_{i\nu}^{(1)}(x) \right|^2 + \left(1 + \frac{\nu^2}{x^2}\right) \left| H_{i\nu}^{(1)}(x) \right|^2 \right]. \quad (5.12)$$

Fortunately the above integral can be taken exactly yielding ¹

$$I = \frac{(1-d+2i\nu)}{4\pi^{5/2} \sinh^2(\nu\pi)} \Gamma(-i\nu - \frac{1}{2} + \frac{d}{2}) \Gamma(i\nu + \frac{1}{2} + \frac{d}{2}) \Gamma(\frac{d}{2} - \frac{1}{2}) \Gamma(-\frac{d}{2}) \times \mathcal{C}, \quad (5.13)$$

where \mathcal{C} is given by

$$\begin{aligned} \mathcal{C} &\equiv 2 \cosh(\nu\pi) \cos\left(\frac{\pi(2i\nu-d)}{2}\right) \cos\left(\frac{\pi(2i\nu+d)}{2}\right) \\ &\quad - \cos\left(\frac{\pi d}{2}\right) \left[\cos\left(\frac{\pi(2i\nu+d)}{2}\right) + \cos\left(\frac{\pi(2i\nu-d)}{2}\right) \right]. \end{aligned} \quad (5.14)$$

The singular limit of the integral I governing the UV divergence of the zero point energy is controlled by the term $\Gamma(-\frac{d}{2})$ in Eq. (5.13).

To perform the dimensional regularization, we expand $d = 4 - \epsilon$ in $\langle \rho \rangle$, obtaining

$$\langle \rho \rangle = \frac{H^4}{2048\pi^2} (4\nu^2 + 1)(4\nu^2 + 9) \left[-\frac{4}{\epsilon} + 2 \ln\left(\frac{H^2}{4\pi\mu^2}\right) + \Delta + \mathcal{O}(\epsilon) \right], \quad (5.15)$$

where

$$\Delta \equiv 2\Psi\left(\frac{5}{2} + i\nu\right) + \Psi\left(\frac{3}{2} - i\nu\right) + \frac{(8\gamma - 12)\nu^2 + 8i\nu + 18\gamma - 15}{(4\nu^2 + 9)}, \quad (5.16)$$

¹We use Maple computational package to perform the integral.

in which $\Psi(x)$ is the Polygamma function and γ is the Euler-Mascheroni constant. One can check that for large ν limit (which is the case here) $\Delta \rightarrow 4 \ln(\nu)$. Finally, note that the $\frac{1}{\epsilon}$ term in Eq. (5.15) originates from $\Gamma(-\frac{d}{2})$ which is divergent near $d = 4$.

Now expanding $\langle \rho \rangle$ to leading order in ϵ and taking the limit $\nu \simeq m/H \gg 1$, we obtain

$$\langle \rho \rangle = -\frac{\nu^4 H^4}{64\pi^2} \left[\frac{2}{\epsilon} - \ln \left(\frac{H^2 \nu^2}{4\pi \mu^2} \right) + \dots \right], \quad (5.17)$$

where the terms in \dots are at the order ν^{-2} and smaller which are subleading compared to the log term.

As expected, the divergent $1/\epsilon$ term in Eq. (5.17) represents the logarithmic divergence of $\langle \rho \rangle$ with the UV scales. After regularizing this divergent term and absorbing the factor 4π into our renormalization scale and noting that $\nu \simeq m/H$, we recover the desired formula Eq. (2.9). It is intriguing that the factor H drops out non-trivially both in the prefactor and inside the log term in Eq. (5.17). This analysis reassures the validity of Eq. (2.9) in the dS spacetime.

Similar to the case of the flat background one can also calculate the correlation length associated with the zero point energy density fluctuations in the dS background. The analysis are somewhat complicated due to technicalities associated with the Hankel functions. Using the approximate relations of the Hankel functions we have verified that the correlation length of the zero point fluctuation in dS backgrounds is indeed $\xi \sim 1/m$ as in flat background. This is consistent with our notion that the intrinsic properties of the zero point energy density fluctuation such as its amplitude and the correlation length are local phenomena and are insensitive to the large scale structure of the spacetime.

Now equipped with the exact mode function in a dS background, we can calculate the accumulated energy density on super Compton scale $\delta\rho_C$ and compare the result with its flat counterpart Eq. (3.3). Setting simply $d = 4$ in Eq. (5.10) and $\nu \simeq m/H$, we obtain

$$\begin{aligned} \delta\rho_C &\simeq \frac{H^4}{16\pi} \int_0^\nu x^4 \left[\left| \frac{d}{dx} H_{i\nu}^{(1)}(x) \right|^2 + \left(1 + \frac{\nu^2}{x^2} \right) \left| H_{i\nu}^{(1)}(x) \right|^2 \right], \\ &\simeq \frac{H^4}{16\pi} \frac{\nu^4}{\pi} \simeq \frac{m^4}{16\pi^2}. \end{aligned} \quad (5.18)$$

Interestingly, we see that the above value of $\delta\rho_C$ is in good agreement with its flat counterpart given in Eq. (3.3).

As we have mentioned above the correlation length of zero point energy density is $\xi \sim 1/m$ so on length scale far beyond this correlation length the distribution of the zero point energy is expected to be uncorrelated. On the other hand, from Eq. (5.18) we see that the accumulated energy density on super Compton scales (i.e. for modes with wavelength larger than ξ) is comparable to the total zero point energy, $\delta\rho_C/\langle \rho_v \rangle \sim 1$. This again confirms that the zero point energy of the heavy fields can not be used to cover the entire spacetime as a background energy density. In other words, one needs to impose $\langle \rho_v \rangle \ll \rho_T \simeq \rho_F$ in order to have a sensible cosmological background.

6 Cosmological Implications

The results from the previous three Sections excludes the heavy fields with $\langle \rho_v \rangle \gg \rho_F$ to contribute to the observed dark energy today. Originally we came to this conclusion based on the fact that one requires an enormous number $N_{\text{patches}} \gg 1$ of incoherent small patches of size $H_{(m)}^{-1}$ to cover the FLRW horizon today. This conclusion was specifically confirmed in Section 4 by looking at the density contrast $|\frac{\delta \rho_v}{\langle \rho_v \rangle}| \sim 1$. A natural question now is if the heavy fields do not contribute to the current dark energy (cosmological constant) then what role in cosmology their vacuum energy density play?

To answer the above question we again employ the picture based on the scale of dS horizons as presented in Fig. 1. Consider the early stage in cosmic expansion history when $\rho_F \gg \rho_v$ so $H_F^{-1} \ll H_{(m)}^{-1}$ as in left panel of Fig. 1. In this situation the observable universe is inside the horizon of zero point energy but at the same time since $\rho_v \ll \rho_F$, then the contribution of ρ_v in expansion rate is too small to be important. As time proceeds and ρ_F decreases further we approach the epoch when $\rho_F \sim \rho_v$ and the zero point energy becomes relevant and we notice the effects of the vacuum energy. As time goes by further, ρ_F falls off rapidly so we end up with the situation $\rho_v \gg \rho_F$ as in right panel of Fig. 1. In this case the regions filled with the zero point energy develop strong inhomogeneities while falling into the FLRW Hubble horizon. The time scale for the dS patches to create inhomogeneities is about $2/H_{(m)}$ as it takes time of about $1/H_{(m)}$ for each dS horizon to enter the FLRW horizon. Once the FLRW horizon grows large enough (or equivalently when $\rho_F \ll \rho_v$) more and more of dS patches enter the FLRW Hubble radius. The variance condition (4.16) indicates that the mass inside the dS patches inside the FLRW Hubble radius collapse to form black holes. The time scale for the duration of collapse is expected to be $t_{\text{col}} \sim 1/m$. On the other hand we have $t_{\text{col}}/H_{(m)}^{-1} \sim m/M_P \ll 1$. Therefore, for our purpose we can assume that the masses inside the patches of the zero point energy shrink to black holes instantaneously. The crucial point is that once the masses inside these patches collapse into black holes, their contribution in background cosmological dynamics is expected to be in the form of matter. This indicates that the resulting mass behaves like dark matter or the seeds of dark matter.

Despite the above qualitative discussions, at this stage we can not be more specific about the mechanism of collapse for the regions filled with the zero point energy of heavy fields after the time when $H_F^{-1} > H_{(m)}^{-1}$. Depending on the mechanism of collapse, different scenarios are possible which we study below using a phenomenological fluid description.

6.1 Phenomenological Fluid Description

As mentioned above we do not know the detail mechanism which governs the collapse of the masses enclosed inside patches of zero point energy into the FLRW horizon. Here, we consider a phenomenological approach and treat the effects of heavy fields like a fluid with an undetermined equation of state.

To be more specific, let us denote the energy density of massive fields after the time when ρ_F is diluted enough and $\rho_F < \rho_m$ by $\tilde{\rho}_m$. We also assume that each components of effective

energy density to evolve separately by their own energy conservation: $\dot{\rho}_F + 3H(\rho_F + p_F) = 0$ and $\dot{\tilde{\rho}}_m + 3H(\tilde{\rho}_m + \tilde{p}_m) = 0$ in which \tilde{p}_m represents the effective pressure of the new matter-like energy component. On the physical ground we may expect $\tilde{p}_m \sim 0$ but for sake of generality we kept it undetermined. Defining the equation of states via $p_F \equiv w_F \rho_F$ and $\tilde{p}_m \equiv w_m \tilde{\rho}_m$, the total energy density now is $\rho_T = \rho_F + \tilde{\rho}_m$, yielding

$$\rho_T(t) = \left[\rho_m \left(\frac{a(t)}{a(t_m)} \right)^{-3(1+w_m)} + \rho_F(t_m) \left(\frac{a(t)}{a(t_m)} \right)^{-3(1+w_F)} \right], \quad (6.1)$$

in which t_m is the time when $\rho_F(t_m) \sim \rho_m$, i.e. when the two Hubble radius H_F^{-1} and $H_{(m)}^{-1}$ nearly coincide. Considering the numerical uncertainties, we set $\tilde{\rho}_m(t_m) \equiv \kappa' \rho_F(t_m)$ in which κ' is a numerical constant which may depend on m . Then, the total energy density is given by

$$\rho_T(t) = \rho_F(t_m) \left(\frac{a(t)}{a(t_m)} \right)^{-3(1+w_F)} \left[1 + \kappa' \left(\frac{a(t)}{a(t_m)} \right)^{3(w_F - w_m)} \right]. \quad (6.2)$$

The above is a phenomenological description of the contribution of the heavy fields after the time when $\rho_F < \rho_m$. As mentioned we have introduced the effective equation of state ω_m to capture the uncertain behaviour of the zero point energy associated with the massive fields after the time of collapse. If the mechanism of collapse is efficient then physically we expect $\omega_m = 0$ and $\tilde{\rho}_m$ to be dark matter. Of course this is a good news in which the nature of dark matter and dark energy is unified in this proposal as both being generated dynamically from the zero point energy. However, the difficulty with $\omega_m \simeq 0$ is that at the early time in cosmic history, the produced dark matter from the zero point energy of electron and heavier fields rapidly dominate over radiation energy density, long before the time of matter radiation equality at the temperature $T_{eq} \sim 3\text{eV}$, altering the hot big bang cosmology in various unwanted ways. One way out may be to consider a situation in which the phenomenological parameter κ' in Eq. (6.2) is (exponentially) small so it will take a long time for the seed of dark matter to take over the radiation energy density. On the other hand, if $\omega_m \simeq \frac{1}{3}$ then $\tilde{\rho}_m$ may behave like dark radiation. Considering the uncertainties involved in the process, ω_m may take different values for various fields so $\tilde{\rho}_m$ may behave differently during cosmic history so we may have dark radiation, dark matter or stiff fluid with $\omega_m = 1$.

Another possibility is that the mechanism of collapse for the mass enclosed inside the dS horizon of zero point energy is such that $\omega_m = \omega_F$ so $\tilde{\rho}_m(t) \propto \rho_F(t)$, i.e.

$$\tilde{\rho}_{(m)}(t) = 3\kappa M_P^2 H_F^2 = \kappa \rho_F, \quad (6.3)$$

in which κ is a numerical constant. This corresponds to a tracking scenario in which the energy density of the heavy fields at each stage in cosmic history tracks the background FLRW energy density. During the radiation dominated era their energy densities follow that of a radiation energy density while during the matter dominated era they behave like matter. This is an interesting case so below we concentrate on this scenario more closely.

The tracking scenario has interesting implications for the current energy density of dark matter. To see this, let us assume that $\rho_F = \rho_\gamma + \rho_b$ in which ρ_γ and ρ_b represent the energy

density of radiation and baryon. Specifically, we assume that there is no conventional (say WIMPs-like) dark matter in our setup. Let us define the time of matter-radiation equality at the redshift z_{eq} as when $\rho_b(z_{eq}) = \rho_\gamma(z_{eq})$. Now observe that what we mean by dark matter in Λ CDM model is the zero point energy density given by the heavy field in Eq. (6.3). Specifically, the current would-be dark matter energy density is actually $\rho_{DM}^{(0)} = \tilde{\rho}_m(t_0) = \kappa\rho_F^{(0)} = \kappa\rho_b^{(0)} + \kappa\rho_\gamma^{(0)}$. Taking into account that the energy density of radiation and baryon falls off like a^{-4} and a^{-3} respectively we obtain

$$1 + z_{eq} = \frac{1}{1 + \kappa} \left(\frac{\Omega_M^{(0)}}{\Omega_\gamma^{(0)}} - \kappa \right), \quad (6.4)$$

in which $\Omega_M^{(0)} \sim 0.31$ ($\Omega_\gamma^{(0)} \sim 10^{-4}$) is the fractional total matter (radiation) energy density today as inferred from Λ CDM model. Now defining $z_{eq}^{\Lambda CDM}$ as the time of matter-radiation equality in Λ CDM model and assuming $\kappa \sim \mathcal{O}(1)$ we obtain

$$z_{eq} \simeq \frac{z_{eq}^{\Lambda CDM}}{1 + \kappa}. \quad (6.5)$$

This means that in the current setup the time of matter-radiation equality is shifted towards later epoch in cosmic history. This is because we have assumed that the time of matter-radiation equality happens when $\rho_b(z_{eq}) = \rho_\gamma(z_{eq})$. Taking $\kappa \lesssim 1$ we may have $z_{eq} \sim \text{few} \times 1000$. However, note that shifting z_{eq} close to the time of CMB last scattering can be dangerous for the CMB observations. But we also note that shifting the time of matter-radiation equality towards later epoch can play important roles to solve the H_0 tension problem.

The above conclusion about the roles of the heavy fields may seem in conflict with our starting point that the vacuum energy should be locally Lorentz invariant and $\langle p \rangle = -\langle \rho \rangle$ which is trivially violated for both radiation and matter. The answer is that the relation $\langle p \rangle = -\langle \rho \rangle$ is enforced locally, say deep inside each dS patch. However, what we have in Eqs. (6.3) or (6.2) is the collective energy density of a large number of patches of zero point energy inside the FLRW horizon which may not be in conflict with the local requirement $\langle p \rangle = -\langle \rho \rangle$ deep inside each dS patch.

6.2 Selection Rules

Based on the results from the previous Sections, we end up with an interesting “selection rule” which works as follows. At each stage in cosmic epoch, only field with $H_{(m)} \sim H_F$ and energy density $\langle \rho_v \rangle \sim \rho_F$ can be relevant as the source of the dark energy. Fields which are much lighter ($H_{(m)} \ll H_F$) are irrelevant in cosmic expansion. This is simply because their contributions in $\langle \rho_v \rangle$ as given in Eq. (2.9) is exceedingly smaller than ρ_F to be noticeable. Finally, heavy fields with $H_{(m)} \gg H_F$ can not be the source of dark energy and depending on the mechanism of collapse they may form the seeds of dark matter from the start or to track the background FLRW energy density ρ_F and be the source of dark matter after the time of matter-radiation equality.

The above selection rule can easily address both the old and the new cosmological constant problems. Recall that the old cosmological constant problem is why ρ_v is not large i.e. why it is so small as $(10^{-3}\text{eV})^4$. The new cosmological constant problem is why the vacuum energy density becomes comparable to the matter energy density at the current stage of the cosmic history. The resolution is that there is a field in the SM field content, the lightest neutrino², with the vacuum zero point energy $\rho_{(\nu)} \sim m_\nu^4$ which happens to have a mass at the same order as $\rho_{F0}^{1/4} \simeq \rho_c^{1/4}$ in which $\rho_c \sim (10^{-3}\text{eV})^4$ is the critical energy density. The entire FLRW universe is currently within a single patch of the lightest neutrino with the horizon radius $H_{(\nu)}^{-1}$. As the universe expands, there will be a time in future when the dS horizon of the neutrino becomes smaller than the FLRW horizon and the effective mass (energy) enclosed inside the patches of the neutrino field will collapse into the FLRW horizon. The current dark energy survives in future for another period of roughly $1/H_{(\nu)} \sim 1/H_0 \sim 10^{10}$ years before the universe enters a matter dominated era (note that by that time the energy density of radiation is much smaller than that of matter). Of course, this conclusion about the future dynamics of the universe is based on the assumption that there is no light field in beyond SM sector (say like light axion) with mass lighter than 10^{-3}eV to contribute to the future dark energy. We comment that the relations between the zero point energy of neutrino and the observed dark energy (cosmological constant) have been studied in different contexts in the past in [29–33]

An immediate corollary of our selection rule is that at early stage in cosmic history (i.e. for all time prior to the time of matter-radiation equality) only fields with the masses at the order of the FLRW photon temperature could contribute to dark energy at that epoch. This is because the energy density of the FLRW background during the radiation dominated era is related to the photon temperature via

$$\rho_F = \frac{\pi^2}{30} g_* T^4, \quad (6.6)$$

in which g_* is the effective relativistic degrees of freedom at the temperature T . Now comparing the above equation with Eq. (2.9) for the vacuum energy density, and noting that for the field of interest $H_{(m)} \sim H_F$, we conclude that

$$m \sim T. \quad (6.7)$$

If a field is much lighter than T , then its vacuum energy density $\propto m^4$ is much smaller than T^4 and can not be important in cosmic energy density. On the other hand, if a field is much heavier than T , then based on the arguments mentioned in previous Sections the dS horizon radius associated with its vacuum energy density is much smaller than the Hubble radius H_F^{-1} and they collapse into FLRW horizon behaving like dark radiation or dark matter. Note that while formula (6.7) works well for the time prior to matter-radiation equality, but it also

²While the differences in neutrino mass squared are known, but the absolute masses and the mass of lightest neutrino are not exactly known, see [26]. Here we adopt the simple assumption that the lightest neutrino has the mass at the order 10^{-2}eV [27, 28].

gives a reasonable estimation for the relation between the mass of the lightest neutrino and the energy scale of the current dark energy. To see this note that $T_{\gamma_0} \sim 10^{-4}\text{eV}$ which is only about two orders of magnitude below the expected mass of lightest neutrino. Of course, this is easily understandable since after the time of matter-radiation equality the energy density of radiation falls off by an additional factor of $1/a$ compared to the matter energy density. This is why we had to use the critical mass density ρ_c instead of $T_{\gamma_0}^4$ to correctly estimate the magnitude of the current dark energy density.

Since only fields with $m \neq 0$ contribute to the vacuum energy we conclude that before the electroweak symmetry breaking at the temperature $T_{EW} \sim 160\text{ GeV}$, all SM fields were massless and they did not contribute to dark energy. Immediately after the electroweak phase transition the relevant fields are the top quark $m_t \simeq 170\text{ GeV}$, the Higgs field $m_H \sim 125\text{ GeV}$ and the three gauge bosons $m_Z \simeq 91\text{ GeV}$, $m_{W^\pm} \simeq 80\text{ GeV}$. The top quark and the Higgs field are somewhat heavy so they are at the threshold of being able to contribute into dark energy at that time, while the three gauge bosons have good chances to play important roles as the source of dark energy. Of course, as the temperature of the universe falls below the mass of vector bosons, their contribution in dark energy expires and they may contribute to dark radiation or dark matter depending on the mechanism of collapse of their zero point energy.

This story is repeated when the background temperature approaches the mass of other fundamental particles such as $m_\tau \simeq 1.7\text{ GeV}$, $m_\mu \simeq 105\text{ MeV}$ and $m_e \simeq 0.5\text{ MeV}$. Specially important is the time when the temperature is around MeV when the big bang nucleosynthesis (BBN) is at work. Changing the Hubble expansion rate of the universe at this epoch, either by contribution from dark energy of the electron field or its contributions in the form of dark radiation, can affect the dynamics of BBN. This issue requires careful investigation which can provide a non-trivial check for the consistency of the whole picture presented here. Another important time to look for is when the heavier neutrino fields (i.e. not the lightest neutrino) with the mass $m \sim 0.1\text{ eV}$ become relevant to contribute into dark energy. This is somewhat around $T \sim 10^{-2}\text{ eV}$ and redshifts of $z \sim 10^2$.

Based from the above discussions, we have transient periods of dark energy for a few e-folds anytime in cosmic history when the condition $m \sim T$ is met during which the Hubble expansion rate stays nearly constant and then it falls off as in conventional big bang cosmology. These can happen in cosmic history both after the surface of last scattering (for heavier neutrinos) at redshift $z \sim 10^2$ and before the surface of last scattering (for e , μ and τ fields) at much higher redshifts. The curious conclusion is that since the energy density is nearly constant for a few e-folds then the value of the Hubble expansion rate today (and at the time of last scattering) would be larger than what the standard ΛCDM model predicts. Multiple transient periods of dark energy both at early and intermediate cosmic expansion history is a non-trivial prediction of the model that has good potentials to solve the H_0 tension problem [34–36]. In this view, our proposal can incorporate the early dark energy (EDE) mechanism [37, 38] which is proposed to solve the H_0 tension problem.

We have presented the main picture of our proposal by now. In the next three Sections we consider a toy model studying the dynamics of bubbles of zero point energy expanding in

cosmological backgrounds which may shed some light on our proposal concerning the roles played by the dS horizon of zero point energy. The reader who is not interested in these technicalities may skip them and jump to Section 10.

7 Expansion of Bubble of Zero Point Energy

In the previous discussions we did not specify how patches of zero point energy with the size $H_{(m)}^{-1}$ are created from the vacuum. Obviously this is a non-trivial quantum process which is far beyond the scope of this work. However, here we simplify this picture by assuming that bubbles of zero point energy with the initial size $R_i \ll H_{(m)}^{-1}$ are continuously created from the vacuum fluctuations and then expand outwards. As before, the vacuum energy for each field of mass m is $\rho_v(m) \sim m^4$ yielding to a local Hubble radius $H_{(m)} \sim m^2/M_P$. Now the question is how these bubbles evolve in a given cosmological background? Can each bubble of the size $H_{(m)}^{-1}$ evolve to become as big as the observable universe before encountering another bubble of zero point energy? If not, then we end up with the picture where the bubbles of zero point energy merge with each other quickly and generate complicated inhomogeneities.

The above question is very similar to the question of bubble expansion in false vacuum decay which is vastly studied in the literature [39–49]. Below we follow the analysis of [49] which studied the evolution of bubble of true vacuum in an inflationary false vacuum background.

The initial size R_i of the bubble of zero point energy is undetermined. However, based on quantum consideration one may expect that the bubbles are pupped up from the vacuum with the radius of the Compton length, $R_i = m^{-1}$. We would like to study how the bubble universe expands in this background and in particular what is the time scale for it to encounter another bubble of zero point energy.

We consider two FLRW-types cosmologies for the bubble of zero point energy and our observable FLRW universe, separated at the surface which is expanding outwards. The metrics of the interior and the exterior geometries are given by

$$ds_{\pm}^2 = dt_{\pm} + a_{\pm}(t_{\pm})^2(dr_{\pm}^2 + r_{\pm}^2 d\Omega^2), \quad (7.1)$$

in which the $-(+)$ stands for the interior (exterior) cosmology equipped with their corresponding cosmic time t_{\mp} and the scale factor a_{\mp} . We have allowed for the cosmic times and the spatial radial coordinate r_{\pm} to be different. The azimuthal directions are covered by the coordinates (θ, ϕ) with the usual line element $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$. Because of the spherical symmetry the azimuthal geometries are the same for both backgrounds.

The two geometries are glued at the surface $R = R(\tau)$ in which τ is the time coordinate defined on the three-surface $R = \text{constant}$. The induced line element of the expanding three-surface is given by

$$ds^2 = -d\tau^2 + R(\tau)^2 d\Omega^2. \quad (7.2)$$

Our goal is to find the dynamics of $R(\tau)$ in terms of the properties of the bubble of zero energy and H_F . We solve the dynamics of the system by expressing various coordinates such as t_{\pm} and r_{\pm} on the surface of wall as functions of τ , i.e. $r_{\pm} = r_{\pm}(\tau)$ and so on.

The normal four-vector to the surface of wall is directed outward, given by

$$n_\mu^\pm = (-a_\pm r'_\pm, a_\pm t'_\pm, 0, 0). \quad (7.3)$$

In order for the two solutions to consistently match to each other one needs a distribution of local energy density on the surface of the expanding wall. Assuming local Lorentz invariance, a natural choice of the energy momentum tensor of the surface is in the form

$$S^{\mu\nu} = \sigma U^\mu U^\nu + \zeta (h^{\mu\nu} + U^\mu U^\nu), \quad (7.4)$$

in which $U^\mu = (1, 0, 0, 0)$ is the comoving four-velocity of the expanding wall. In addition, $h^{\mu\nu} = g^{\mu\nu} - n^\mu n^\nu$ is the induced metric on the surface of wall. In this view, σ is the surface energy density of the wall and ζ is its surface tension. In general, σ and ζ are function of time, $\sigma = \sigma(\tau)$, $\zeta = \zeta(\tau)$, satisfying the local energy conservation on the wall, $\sigma' + 2(R'/R)(\sigma + \zeta) = 0$ in which a prime here and below indicates the derivative with respect to τ . A simple possibility is that $\zeta = -\sigma$ yielding to a constant value of σ which will be assumed in the following analysis.

The continuity of the metric on the surface $r = R(\tau)$ requires that

$$R(\tau) = R_+(\tau) = R_-(\tau), \quad (7.5)$$

in which $R_\pm \equiv a_\pm(t_\pm)r_\pm$. Furthermore,

$$d\tau^2 = dt_+^2 - a_+(t_+)^2 dr_+^2 = dt_-^2 - a_-(t_-)^2 dr_-^2. \quad (7.6)$$

This equation can be simplified further to

$$t_\pm'^2 - a_\pm(t_\pm)^2 r_\pm'^2 = 1, \quad (7.7)$$

where a prime denotes the derivative with respect to τ .

From the above relations one can solve for t'_\pm , obtaining

$$t'_\pm = \frac{-RR'H_\pm + \sqrt{1 + R'^2 - R^2 H_\pm^2}}{1 - R^2 H_\pm^2}, \quad (7.8)$$

in which H_\pm represents the Hubble expansion rate on each side.

The other junction condition is related to the discontinuity of the extrinsic curvature across the wall. With the extrinsic curvature $K_{\mu\nu}$ given by $K_{\mu\nu} = n_{\mu;\nu}$ the other junction condition is given by [50]

$$[K_{ab}] = -8\pi G(S_{ab} - \frac{h_{ab}}{2}S), \quad (7.9)$$

in which the notation $[]$ indicates the discontinuity across Σ and the indices a, b runs across (r, θ, ϕ) on each side. From the $\theta - \theta$ component we have

$$K_{\theta\theta}^\pm = R(t'_\pm + RH_\pm a_\pm r'_\pm), \quad (7.10)$$

which after plugging into Eq. (7.9) yields

$$[t'_\pm + RH_\pm a_\pm r'_\pm] = -4\pi G\sigma. \quad (7.11)$$

After some manipulations this yields to our desired relation for the dynamics of $R(\tau)$ [49]

$$1 + R'^2 = A^2 R^2, \quad (7.12)$$

in which

$$\begin{aligned} A^2 &\equiv \frac{\sigma^2}{16M_P^4} + \frac{\rho_- + \rho_+}{6M_P^2} + \left(\frac{\rho_+ - \rho_-}{3\sigma}\right)^2 \\ &= \frac{\sigma^2}{16M_P^4} + \frac{H_-^2 + H_+^2}{2} + \frac{M_P^4}{\sigma^2} (H_+^2 - H_-^2)^2. \end{aligned} \quad (7.13)$$

Note that either ρ_+ or ρ_- represents the value of energy density in our FLRW universe so in general it may not be easy to solve Eq. (7.12) analytically as H_F is a function of time. Below, we consider the cases $\rho_- = \rho_v(m)$ and $\rho_+ = \rho_v(m)$ separately.

8 Case Where $H_{(m)}^{-1} < H_F^{-1}$

First consider the case where $H_{(m)}^{-1} < H_F^{-1}$ and the bubble of zero point energy furnishes the interior region, $\rho_- = \rho_v(m) \equiv 3M_P^2 H_{(m)}^2$, so the exterior region represents the FLRW universe with $\rho_+ = \rho_F \equiv 3M_P^2 H_F^2$. The bubble at the moment of formation has the size R_i and then it grows. We may assume $R_i \sim 1/m$, i.e. the bubble is created quantum mechanically from the vacuum with the Compton length and at the timescale $1/m$. Because of the causality, we always assume $R_i H_{(m)} \ll 1$ which means that the bubble formation happens deep inside the Hubble radius $H_{(m)}^{-1}$ associated with the zero point energy. Also, as times goes by and the radius of the bubble increases, the energy density of the exterior region falls off rapidly (like $1/a_+^3$ or $1/a_+^4$ for MD and RD era respectively) and we eventually end up with the situation where $H_{(m)}^{-1} \ll H_F^{-1}$ and the picture in the right panel in Fig. 1 becomes the natural outcome.

In order to solve Eq. (7.12) we need to estimate the surface energy density σ . Since the bubble is created from zero pointed energy and σ is the quantum vacuum energy density of the wall then one naturally expects that $\sigma = cm^3$ where c is a coefficient, like the prefactor in Eq. (2.9). Therefore, in the following analysis we simply estimate $\sigma \sim m^3$.

Before presenting the general analysis, first we consider the extreme case where $\rho_v \gg \rho_F$ corresponding to $H_{(m)} \gg H_F$. This is the case if the zero point energy is created from very massive fields. In this case, one can check that the third term in Eq. (7.13) dominates and $A \simeq m$. In this limit one can easily solve Eq. (7.12), obtaining

$$R(\tau) = \frac{1}{A} \cosh(A\tau). \quad (8.1)$$

We have neglected a phase term by assuming that at the time of formation $R(0) = 1/A \sim m^{-1}$. This is consistent with the intuition that the bubble of zero point energy are puffed out from the vacuum with the Compton length m^{-1} .

Once the bubble is formed from the vacuum it will expand. The important question is what is the timescale when a bubble of zero point energy meets another bubble created in different regions. If this timescale is much less than H_F^{-1} , then the corresponding bubble can not be a seed of our observed universe. In this case we end up with many small bubbles which have percolated throughout the observable universe as in the picture in the right panel of Fig. 1. They subsequently merge with each other generating complicated small scales inhomogeneities or collapse to form mini black holes.

Let us define $\tau \equiv \tau_H$ as the time when a given bubble approaches another nearby bubble, i.e. the timescale of bubble collision. A conservative estimation is the time when the radius of the bubble becomes comparable to the Hubble radius of its dS spacetime. This is because the Hubble radius is the boundary of another bubbles seated nearby. From Eq. (7.13) we obtain

$$\frac{1}{A} \cosh(A\tau_H) = H_{(m)}^{-1} \rightarrow \tau_H \sim \frac{1}{A} \ln\left(\frac{A}{H_{(m)}}\right), \quad (8.2)$$

which for $A \sim m^{-1}$, yields

$$\tau_H \simeq \frac{1}{m} \ln\left(\frac{M_P}{m}\right). \quad (8.3)$$

This indicates that the timescale of collision, as measured by a comoving observer on the surface of bubble, is simply m^{-1} which is very short for the massive fields. Therefore, one typically expects that bubble of zero point energy from massive fields collide more rapidly with each other and can not be relevant for the cosmological constant problem.

However, the above timescale is measured by the observer confined to the surface of the bubble. For a proper cosmological interpretation we have to measure the timescale of bubble collision from the point of view of the cosmological observer, i.e. the outside observer. Using Eq. (7.8) we obtain

$$\frac{dt_+}{d\tau} = \frac{\sqrt{A^2 - H_F^2} R - R R' H_F}{1 - R^2 H_F^2}. \quad (8.4)$$

We expect that $R H_F \ll 1$ since due to causality the created bubble can not be large compared to the FLRW Hubble radius. With $A \sim m^{-1}$, one approximately obtains

$$\frac{dt_+}{d\tau} \simeq R A \longrightarrow t_+ \sim \frac{1}{A} \sinh(A\tau_H). \quad (8.5)$$

Using the value of τ_H in Eqs. (8.2) or (8.3) we obtain

$$t_+(\tau_H) \sim \frac{1}{H_{(m)}}. \quad (8.6)$$

This is an interesting result. It means that from the clock of an outside observer (i.e. the cosmological observer) the timescale of bubble collision is at the order $1/H_{(m)} \ll 1/H_F$. Consequently, once the bubble of vacuum energy is popped out of vacuum, it expands and

very quickly collides with nearby bubbles. Therefore, as mentioned before, we end up with a configuration of many bubble collisions with complicated small structure inhomogeneities.

Also, using the solution for t_+ in Eq. (8.5) and the evolution of the bubble in Eq. (8.1) we obtain $R(\tau) \sim 1/t_+ \propto 1/H_F$. So as the bubble expands it keeps track of the Hubble radius of the FLRW region. However, this does not happen in practice as the bubble has collided with nearby bubbles before approaching the asymptotic solution $R \sim 1/H_F$.

After presenting the above special case, now let us consider the general case with no hierarchy assumed between ρ_v and ρ_F . The goal is to solve $R(\tau)$. The dynamics of evolution of $R(\tau)$ is given by Eq. (7.12) while the rate of change of the external observer's clock with respect to τ is given by Eq. (8.4). Combining Eqs. (7.12) and (8.4) one can eliminate $d\tau$ and calculate the rate $\frac{dR}{dt_+}$ as measured by the cosmological observer,

$$\frac{dR}{dt_+} = \frac{\sqrt{A^2 R^2 - 1}}{R} (1 - R^2 H_F^2) \left(\sqrt{A^2 - H_F^2} - H_F \sqrt{A^2 R^2 - 1} \right)^{-1}. \quad (8.7)$$

The above differential equation can not be solved analytically, except in some limiting cases of interests. The full numerical solution for the evolution of R is presented in Fig. 2 where the dimensionless variables are defined via

$$z \equiv H_{(m)} t_+, \quad y(z) \equiv R H_{(m)}. \quad (8.8)$$

Initially $z_i \sim \frac{H_{(m)}}{H_F} > 1$ while as the bubble expands and z increases towards larger values the energy density of the exterior region is diluted quickly. In this limit, one can check that $AR \gg 1$ and the differential equation (8.7) simplifies to

$$\frac{dy}{dz} = 1 + \lambda \frac{y(z)}{z}, \quad (8.9)$$

where the parameter $\lambda \equiv 2/(3(1+w_F))$ is defined via $H_F \equiv \frac{\lambda}{t_+}$. In particular, for a radiation dominated universe $\lambda = 1/2$ while for a matter dominated universe $\lambda = 2/3$. For the physically motivated matter fields we expect $0 < \lambda < 1$.

The above differential equation with the initial condition $y(z_i) = z_i$ can be solved as

$$y(z) \simeq \frac{z}{(1-\lambda)} + \left(\frac{z}{z_i}\right)^\lambda \left(y_i - \frac{z_i}{(1-\lambda)}\right). \quad (8.10)$$

We have checked that the above analytic solution is in excellent agreement with the full numerical solution as can be seen in Fig. 2.

As time goes by z increases and the first term in Eq. (8.10) is the attractor solution for $z \rightarrow \infty$ yielding $y(z) \simeq \frac{z}{1-\lambda}$. In particular we obtain $R \simeq \frac{\lambda}{(1-\lambda)H_F}$ which is consistent with the limiting case $\rho_v \gg \rho_F$ studied at the start of this Section. However, the second term in Eq. (8.10) is important for early stage when $z \gtrsim z_i$. Also note that for the physically relevant configurations we expect $y_i/z_i \sim R(z_i)H_F(z_i) \ll 1$. This is because we expect the bubble to form initially deep inside the dS horizon, i.e $R(z_i) \ll H_{(m)}^{-1} \ll H_F(z_i)^{-1}$.

As before, the important question is what is the timescale when the bubble grows to the size $1/H_{(m)}$, corresponding to the time $z = z_c$ when $y(z_c) = 1$. Since $y(z)$ grows quickly, one




Figure 2: The evolution of $y(z) \equiv RH_{(m)}$ vs. $z \equiv H_{(m)}t_+$ for the setup where $\rho_- = \rho_v$ with $z_i > 1$. Initially, the bubble is formed deep inside the Hubble radius $H_{(m)}^{-1}$ with $y_i \ll 1$ but as time goes by and $H_F \rightarrow 0$ the bubble grows like $R \simeq \lambda/((1-\lambda)H_F)$. The solid red curve represents the full numerical solution while the blue dots represent the analytic solution Eq. (8.10).

expects that z_c is not far from z_i . Using the approximation $z_c = z_i(1 + \delta)$ with $\delta \ll 1$ and plugging this into Eq. (8.10) we obtain $\delta \simeq 1/x_i$, yielding

$$z_c \simeq z_i + 1. \quad (8.11)$$

From the above expression we easily see that from the point of view of the external FLRW observer it takes timescale of order $\Delta t_+ \sim 1/H_{(m)}$ for the bubble to grow from the initial size R_i to the dS horizon size $1/H_{(m)}$. This is consistent with the result obtained in Eq. (8.6) where $\rho_v \gg \rho_F^i$.

Similarly, after another time of $\Delta t_+ \sim 1/H_{(m)}$, the size of the bubble increases from $1/H_{(m)}$ to $2/H_{(m)}$ and so on. In other words, after time $n/H_{(m)}$ has elapsed, n dS patches of size $1/H_{(m)}$ enter the bubble. This is well supported from the linear growth of the first term in Eq. (8.10) as also seen in Fig. 2. Therefore, one quickly ends up with a picture like the right panel of Fig 1.

9 Case Where $H_{(m)}^{-1} \gg H_F^{-1}$

Now we consider the interesting case where the FLRW universe is encompassed by the dS horizon of zero point energy as in the left panel of Fig. 1. This happens when initially $H_{(m)}^{-1} \gg H_F^{-1}$ which is naturally the case where the field is very light. Our goal here is to

estimate the timescale of starting from the left panel of Fig. 1 and ending to a configuration like the right panel of Fig. 1.

In this case, we also have the additional requirement that

$$\rho_- = \rho_v(m) + \rho_F, \quad (9.1)$$

where $\rho_v(m) = \rho_+ \equiv 3M_P^2 H_{(m)}^2$ and ρ_F represents the energy density of the FLRW universe in the absence of dark energy (i.e. radiation and matter contributions). Defining $H_F^2 \equiv \rho_F/3M_P^2$, from Eq. (9.1) we have $H_-^2 = H_F^2 + H_{(m)}^2$ so $H_- \neq H_F$.

Note that the condition (9.1) is the requirement that the vacuum energy fills not only the exterior region but also the interior region. This is not in conflict with causality as we imagine that the FLRW bubble is inside the horizon of the dS spacetime generated from the zero point energy, i.e. $H_{(m)}^{-1} \gg H_F^{-1}$. The bubble is an FLRW universe filled with matter, radiation and dark energy which is pushing towards the exterior region filled with the zero point energy. The equations are the same as before except that now we would like to measure the rate of the expansion of $R(\tau)$ as viewed from the internal observer with the clock t_- which is the cosmological observer.

The evolution of $R(\tau)$ is given by Eq. (7.12) while the rate of change of clock of the internal observer with respect to τ is given by

$$\frac{dt_-}{d\tau} = \frac{\sqrt{A^2 - H_-^2} R - R R' H_-}{1 - R^2 H_-^2}. \quad (9.2)$$

Combining Eqs. (7.12) and (9.2) one can eliminate $d\tau$ and calculate the rate $\frac{dR}{dt_-}$ as measured by the cosmological observer,

$$\frac{dR}{dt_-} = \frac{\sqrt{A^2 R^2 - 1}}{R} (1 - R^2 H_-^2) \left(\sqrt{A^2 - H_-^2} - H_- \sqrt{A^2 R^2 - 1} \right)^{-1}. \quad (9.3)$$

Needless to say the above equation can not be solved analytically. However, in the limit of interest where $AR \gg 1$ and $A \gg H_-$, the above equation simplifies significantly into

$$\frac{dR}{dt_-} \simeq 1 + R H_-. \quad (9.4)$$

On the other hand, since $\rho_- = \rho_+ + \rho_F$ and ρ_F is evolving, the relation between t_- and H_- is not a simple power law form. However, we can find an analytic relation between H_- and t_- as follows.

Using the energy conservation $\dot{\rho}_- + 3H_-(\rho_- + p_-) = 0$, we obtain

$$dt_- = -\frac{1}{3H_-} \frac{d\rho_-}{(1 + w_F)\rho_F} = -\frac{1}{3H_-} \frac{d\rho_-}{(1 + w_F)(\rho_- - \rho_+)}. \quad (9.5)$$

Note that in obtaining the above expression the relations $\rho_- = \rho_F + \rho_+$, $p_- = p_F + p_+$ and $\rho_+ + p_+ = 0$ have been used. Furthermore, $w_F \equiv p_F/\rho_F$. Finally, using the Friedmann equation $\rho_- = 3M_P^2 H_-^2$ we can trade $d\rho_-$ in terms of dH_- , obtaining

$$dt_- = -\lambda \frac{dH_-}{H_-^2 - H_{(m)}^2}, \quad (9.6)$$

where as before, $\lambda \equiv 2/(3(1 + w_F))$. The above equation can be solved, yielding

$$t_- = \frac{\lambda}{2H_{(m)}} \ln \left(\frac{H_- + H_{(m)}}{H_- - H_{(m)}} \right). \quad (9.7)$$

Alternatively, the above equation can be inverted to solve H_- in term of t_-

$$H_- = H_{(m)} \coth \left(\frac{H_{(m)} t_-}{\lambda} \right). \quad (9.8)$$

The above solution is easy to understand. As time proceeds, the FLRW energy density inside the bubble is diluted rapidly so asymptotically the interior of the bubble approaches a dS background.

Now we go back to look for the solution of Eq. (9.4). At the onset of formation the bubble is deep inside the dS horizon, $R_i H_{(m)} \ll 1$, so at the early stage when $H_{(m)} t_i \ll 1$ the bubble grows linearly with time $R \simeq R_i + t_-$. However, as time proceeds and $H_{(m)} t_- > 1$ the bubble grows exponentially and

$$R \simeq \frac{e^{H_{(m)} t_-}}{H_{(m)}}, \quad (H_{(m)} t_- > 1). \quad (9.9)$$

Alternatively, from the above equation one can express t_- in term of R as follows

$$t_- \simeq \frac{1}{H_{(m)}} \ln (RH_{(m)}). \quad (9.10)$$

As the bubble grows, more and more of dS patches of size $1/H_{(m)}$ enters into the bubble. The above formula indicates that the timescale for the first few successive entrance of the patches is approximately $1/H_{(m)}$. In other words, it takes time of roughly $1/H_{(m)}$ for the bubble size to grow from $1/H_{(m)}$ to $2/H_{(m)}$. However, the above ideal picture of indefinite expansion of the bubble is not trusted after time of roughly $t_- \sim 2/H_{(m)}$. This is because after this timescale the FLRW energy density has fallen significantly below $\rho_v(m)$ and based on the variance analysis in Section 4 the density contrast of zero point energy develops nonlinearities. After that, the dS patches are expected to collapse into black holes and the above idealized picture can not be trusted.

Having said this, it would be insightful to calculate H_F and ρ_F as the bubble expands. As we mentioned, originally $\rho_- \sim \rho_F$. Beyond the time $t_- \sim 1/H_{(m)}$ we have $H_F \ll H_{(m)}$. Using $\rho_- = \rho_v(m) + \rho_F$ and $H_-^2 = H_{(m)}^2 + H_F^2$ we obtain $H_F^2 \simeq 2H_{(m)}(H_- - H_{(m)})$. Using Eq. (9.8) to express H_- in terms of $H_{(m)}$, we obtain

$$H_F \simeq 2H_{(m)} e^{-\frac{1}{\lambda} H_{(m)} t_-} \quad (t_- > H_{(m)}^{-1}). \quad (9.11)$$

Now, combining the above equation with the solution of $R(t_-)$ in Eq. (9.9) we obtain

$$R \simeq \frac{1}{H_{(m)}} \left(\frac{H_F}{2H_{(m)}} \right)^{-\frac{2}{3(1+w_F)}}. \quad (9.12)$$

Interestingly we see that the radius of the bubble evolves like the scale factor of the FLRW universe in the absence of the cosmological constant, $R \propto a_F \propto H_F^{-\frac{2}{3(1+w_F)}}$.

The conclusion is that starting as in the left panel of Fig. 1, the energy density inside the bubble keeps track of the vacuum energy for time scale of roughly $1/H_{(m)}$ to $2/H_{(m)}$ before the bubble is filled with numerous patches of size $1/H_{(m)}$ ending to a configuration like in the right panel of Fig. 1.

10 Summary and Discussions

In this work we have revisited the quantum cosmological constant problem. As already known in the literature, the conventional approach of imposing a hard UV momentum cutoff violates the underlying Lorentz invariance of the vacuum. Employing a regularization scheme which respects the underlying Lorentz invariance of the theory, such as the dimensional regularization method, one obtains that the energy density of each field scales like $\rho_v \sim m^4$. Furthermore, the condition of Lorentz invariance of the vacuum, $\langle p \rangle = -\langle \rho \rangle$ is manifest. Some noticeable conclusions are that the massless fields such as gravitons, photons and gluons do not contribute to the vacuum energy density. Furthermore, $\langle \rho \rangle$ runs logarithmically with the renormalization scale μ and depending on the ratio m/μ the value of cosmological constant can be either positive (dS) or negative (AdS).

We have highlighted that the dS horizon associated with the zero point energy plays important roles. For a field with mass m the associated dS space has a horizon radius $H_{(m)}^{-1} \sim M_P/m^2$ which was not taken into account in previous studies of the cosmological constant problems. In the conventional approach, it is naively assumed that the entire observable FLRW patch is covered by the vacuum energy density $\langle \rho \rangle \sim m^4$ irrespective of the vast hierarchy between the two Hubble radii H_F^{-1} and $H_{(m)}^{-1}$. For the heavy fields with the large hierarchy $H_{(m)} \gg H_F$ this requires as many as $\left(\frac{H_{(m)}}{H_F}\right)^3 \gg 1$ independent patches of zero point energy to fill the FLRW patch. However, since the patches of zero point energy are created quantum mechanically they are uncorrelated. Therefore, one expects that the space filled with as many as $\left(\frac{H_{(m)}}{H_F}\right)^3$ uncorrelated patches of zero point energy to be highly chaotic and prone to collapse into small scale inhomogeneities and possibly to black holes. To support this picture, we have calculated the variance of the zero point energy for both the real scalar field and the Dirac fermion field. In both cases we have found that $|\delta\rho_v/\langle\rho_v\rangle| \sim 1$ supporting the above intuitive picture. Furthermore, the correlation length of the fluctuations of the zero point energy is calculated to be at the order $\xi \sim m^{-1}$. This results in the vast hierarchy $\xi \ll H_{(m)}^{-1} \ll H_F^{-1}$. It is shown that the accumulated energy density on super Compton scales, i.e. scales which are far beyond the correlation length, is comparable to the would-be background vacuum energy density. This conclusion, in conjunction with the above hierarchies amongst the three scales ξ , $H_{(m)}^{-1}$ and H_F^{-1} , is another sign that one can not use the vacuum energy density $\langle\rho_v\rangle$ as the dominant background energy density. To have a viable cosmological background, we have to impose $\langle\rho_v\rangle \lesssim \rho_F$ at each stage in cosmic expansion history in which ρ_F is the non-vacuum energy density (i.e. classical energy density) of the background FLRW

cosmology.

We have established a “selection rule” stating that at each stage in cosmic expansion history, only fields with mass m satisfying the condition $H_{(m)} \sim H_F$ can contribute to dark energy at that epoch. Specifically, only fields with the mass at the order of the background photon temperature, $m \sim T$, are relevant as the source for dark energy. Lighter fields carry so little energy ($m^4 \ll T^4$) to be important in cosmic energy budget. On the other hand, heavy fields with $H_{(m)} \gg H_F$ (or $m \gg T$) can not be the source of dark energy while they can be the seeds of dark matter. Our proposal provides a unifying platform for the nature of both dark energy and dark matter in which the zero point energy of light fields are the source of dark energy while dark matter is sourced by the zero point energy of heavy fields. As both dark energy and dark matter emerge dynamically from the competition of the horizon radius of zero point energy and the FLRW Hubble radius, we may call this scenario as the “emergent Λ CDM” or e Λ CDM for short.

The above e Λ CDM proposal has far reaching cosmological implications. In this proposal both the old and new cosmological constant problems can be addressed readily. Specifically, these puzzled are solved by noting that there is a field in SM spectrum, the lightest neutrino, which happens to have a mass at the order of $\rho_c^{1/4} \sim 10^{-2}\text{eV}$. Another important implication is that at various stages in cosmic history the energy density experiences loitering stages of dark energy in which the Hubble expansion rate stays nearly constant for about one or two e-folds in expansion rate and then the energy density falls off as in Λ CDM model. As a consequence, the Hubble expansion rate at the time of CMB last scattering is expected to be higher than what is inferred from Λ CDM model. Currently we are at the last transient stage associated with the lightest neutrino. However, prior to this stage there were two transient phases of dark energy happening at around the mass scale of the heavier neutrinos at $T \sim 10^{-2}\text{eV}$ corresponding to the redshift $z \sim 10^2$. This period will be sometime after the time of CMB decoupling. Multiple transient phases of dark energy, both before and after the surface of last scattering, yield to a higher values of the current Hubble expansion rate compared to what is inferred from the Λ CDM model. In this view our proposal can incorporate the EDE proposal and has the potential to solve the H_0 tension problem. Another interesting prediction of the model (as mentioned above) is that there may be no dark matter. All matter are in the form of known baryonic matter while the roles of dark matter are played by the heavy fields which may track the background FLRW energy density. As such, the time of matter-radiation equality is shifted towards later time in cosmic history.

There are a number of open questions in this study. The immediate one is how the collapse for the patches of zero point energy associated to heavy fields happen. We provided only a rough phenomenological picture of this mechanism. This is an important question which controls whether or not we have the tracking energy or a fluid with the equation of state $\omega_m \sim 0$. Another open question is what is the natural value of the renormalization scale μ ? Note that the vacuum energy runs logarithmically with μ . Furthermore, depending on the statistics of the field (i.e. being a boson or a fermion) and on the ratio m/μ , the vacuum energy can be either positive (dS) or negative (AdS). To solve the cosmological constant problems we have implicitly assumed that the resulting vacuum energy is positive so taking

the neutrinos as the relevant fields for this purpose, we require $\mu > \text{eV}$. If we take μ to be the scale of electroweak symmetry breaking, say $\mu \gtrsim 10^2 \text{GeV}$, then all fermionic (bosonic) fields add positive (negative) contribution to the vacuum energy. If the energy density is AdS-type, then there maybe a fall off period (instead of loitering phase) in Hubble expansion rate at the corresponding stage in cosmic history. This will modify the inferred value of the Hubble expansion rate at the time of CMB last scattering. Therefore, the dS and AdS contributions for all SM degrees of freedom from the mass scale 10^2GeV after the electroweak symmetry breaking down to the neutrino scale should be carefully included to see how dark energy and dark matter behave at each stage in cosmic history.

In Sections 7, 8 and 9 we have presented a dynamical toy model involving bubbles of zero point energy which may mimic the setup studied here. It is assumed that the bubbles of zero point energy with the typical radius of Compton length m^{-1} are created quantum mechanically from the vacuum and then expand in the surrounding cosmological background. We have seen that this simple toy model can capture the main properties of our proposal concerning the roles of the dS horizon associated with the zero point energy. In particular, we have verified that it takes time of order $1/H_{(m)}$ for a dS patch to enter FLRW patch and roughly another time of $1/H_{(m)}$ for two dS patches to enter the FLRW patch.

Finally, it is important to examine the implications of the Λ CDM scenario for other aspects of early universe cosmology. Specifically, the interplay between this proposal and cosmic inflation is an open and interesting question. However, we comment that our proposal is not in conflict with an early stage of inflation. More specifically, during the slow-roll inflation driven by an inflaton field of mass m and the potential V , the dominant energy is given by the inflaton classical potential V_c while the vacuum zero point energy is at the order m^4 . Since for a rolling inflaton $V_c \simeq 3M_P^2 H^2 \gg m^4$ the induced energy density from the zero point fluctuations during inflation is negligible. Therefore, the quantum cosmological constant is not an issue during inflation and a large enough classical vacuum energy during inflation is stable against quantum zero point corrections. As a result, a period of slow-roll inflation can take place at early universe followed by Λ CDM setup after electroweak symmetry breaking.

Note added: After the completion of this work we have become aware³ of a series of papers in [51–54] who questioned the assumption of the homogeneity of the spacetime in the presence of the vacuum zero point energy. It was concluded, among other things, that a uniform cosmological constant can not cover the large scale spacetime and the local spacetime is very inhomogeneous as in Wheeler’s spacetime foam. Using the semiclassical GR, they have studied the cosmological implications of their proposal such as in resolving the old cosmological constant problem. Our analysis in Section 4 for the fluctuations in the zero point energy density and large density contrast is conceptually in line with their investigation.

³We are grateful to Jerome Martin for bringing to our attention the papers by Unruh and his collaborators [51–54].

Acknowledgments

We would like to thank Xingang Chen, Jerome Martin, Shinji Mukohyama, Mohammad Hossein Namjoo and Misao Sasaki for insightful comments about the draft. We also thank Hossein Moshafi for the preliminary investigations of the predictions of the proposal for the H_0 tension problem and Omid Sameie for useful discussions.

References

- [1] S. Weinberg, “Cosmology,” Oxford University Press, 2008.
- [2] N. Aghanim *et al.* [Planck], Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)].
- [3] V. Sahni and A. A. Starobinsky, Int. J. Mod. Phys. D **9**, 373-444 (2000).
- [4] P. J. E. Peebles and B. Ratra, Rev. Mod. Phys. **75**, 559-606 (2003).
- [5] E. J. Copeland, M. Sami and S. Tsujikawa, Int. J. Mod. Phys. D **15**, 1753-1936 (2006).
- [6] S. Weinberg, “The Cosmological Constant Problem,” Rev. Mod. Phys. **61**, 1-23 (1989).
- [7] C. Armendariz-Picon, V. F. Mukhanov and P. J. Steinhardt, Phys. Rev. Lett. **85**, 4438-4441 (2000).
- [8] I. Zlatev, L. M. Wang and P. J. Steinhardt, Phys. Rev. Lett. **82**, 896-899 (1999).
- [9] N. Arkani-Hamed, S. Dimopoulos, N. Kaloper and R. Sundrum, Phys. Lett. B **480**, 193-199 (2000).
- [10] S. Kachru, M. B. Schulz and E. Silverstein, Phys. Rev. D **62**, 045021 (2000).
- [11] C. Csaki, J. Erlich and C. Grojean, Nucl. Phys. B **604**, 312-342 (2001).
- [12] J. M. Cline and H. Firouzjahi, Phys. Lett. B **514**, 205-212 (2001).
- [13] J. M. Cline and H. Firouzjahi, Phys. Rev. D **65**, 043501 (2002).
- [14] S. Weinberg, Phys. Rev. Lett. **59**, 2607 (1987).
- [15] J. Martin, Comptes Rendus Physique **13**, 566-665 (2012), [arXiv:1205.3365 [astro-ph.CO]].
- [16] S. Perlmutter *et al.* [Supernova Cosmology Project], Astrophys. J. **517**, 565-586 (1999).
- [17] A. G. Riess *et al.* [Supernova Search Team], Astron. J. **116**, 1009-1038 (1998).

- [18] E. K. Akhmedov, [arXiv:hep-th/0204048 [hep-th]].
- [19] J. F. Koksma and T. Prokopec, [arXiv:1105.6296 [gr-qc]].
- [20] G. Ossola and A. Sirlin, Eur. Phys. J. C **31**, 165-175 (2003).
- [21] M. Visser, Particles **1**, no.1, 138-154 (2018).
- [22] S. Weinberg, “The Quantum theory of fields. Vol. 1: Foundations.”
- [23] C. Moreno-Pulido and J. Sola, Eur. Phys. J. C **80**, no.8, 692 (2020).
- [24] C. Moreno-Pulido and J. S. Peracaula, [arXiv:2110.08070 [gr-qc]].
- [25] N. D. Birrell and P. C. W. Davies, “Quantum Fields in Curved Space,”
- [26] P.A. Zyla *et al.* [Particle Data Group], PTEP **2020**, no.8, 083C01 (2020).
- [27] A. Loureiro, *et al.* Phys. Rev. Lett. **123**, no.8, 081301 (2019).
- [28] P. Stöcker *et al.*, Phys. Rev. D **103**, no.12, 123508 (2021).
- [29] I. L. Shapiro and J. Sola, JHEP **02**, 006 (2002).
- [30] I. L. Shapiro and J. Sola, Phys. Lett. B **475**, 236-246 (2000).
- [31] D. Glavan and T. Prokopec, [arXiv:1504.00842 [gr-qc]].
- [32] L. E. Ibanez, V. Martin-Lozano and I. Valenzuela, JHEP **11**, 066 (2017).
- [33] M. Blasone, A. Capolupo, S. Capozziello, S. Carloni and G. Vitiello, Braz. J. Phys. **35**, 455-461 (2005).
- [34] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri and D. Scolnic, Astrophys. J. **876**, no.1, 85 (2019).
- [35] A. G. Riess, *et al.*, [arXiv:2112.04510 [astro-ph.CO]].
- [36] E. Di Valentino et al, Class. Quant. Grav. **38**, no.15, 153001 (2021).
- [37] V. Poulin, T. L. Smith, T. Karwal and M. Kamionkowski, Phys. Rev. Lett. **122**, no.22, 221301 (2019).
- [38] T. L. Smith, V. Poulin and M. A. Amin, Phys. Rev. D **101**, no.6, 063523 (2020).
- [39] S. K. Blau, E. I. Guendelman and A. H. Guth, Phys. Rev. D **35**, 1747 (1987).
- [40] V. A. Berezin, V. A. Kuzmin and I. I. Tkachev, Phys. Rev. D **36**, 2919 (1987).
- [41] V. A. Berezin, V. A. Kuzmin and I. I. Tkachev, Phys. Lett. **120B**, 91 (1983).

- [42] A. Aguirre and M. C. Johnson, Phys. Rev. D **72**, 103525 (2005).
- [43] K. Lake, Phys. Rev. D **29**, 1861 (1984).
- [44] J. Garriga, A. Vilenkin and J. Zhang, JCAP **1602**, no. 02, 064 (2016).
- [45] H. Deng, J. Garriga and A. Vilenkin, JCAP **1704**, no. 04, 050 (2017).
- [46] J. Garriga, A. H. Guth and A. Vilenkin, Phys. Rev. D **76**, 123512 (2007).
- [47] S. Ansoldi and T. Tanaka, J. Exp. Theor. Phys. **120**, no. 3, 460 (2015).
- [48] P. Chen, Y. C. Hu and D. h. Yeom, Phys. Rev. D **94**, 024044 (2016).
- [49] H. Firouzjahi, S. Jazayeri, A. Karami and T. Rostami, JCAP **12**, 029 (2017).
- [50] W. Israel, Nuovo Cim. B **44S10**, 1 (1966) [Nuovo Cim. B **44**, 1 (1966)] Erratum: [Nuovo Cim. B **48**, 463 (1967)].
- [51] Q. Wang, Z. Zhu and W. G. Unruh, Phys. Rev. D **95**, no.10, 103504 (2017).
- [52] S. S. Cree, T. M. Davis, T. C. Ralph, Q. Wang, Z. Zhu and W. G. Unruh, Phys. Rev. D **98**, no.6, 063506 (2018).
- [53] Q. Wang and W. G. Unruh, Phys. Rev. D **102**, no.2, 023537 (2020).
- [54] Q. Wang, Phys. Rev. Lett. **125**, no.5, 051301 (2020).