

# Anisotropic Special Relativity

Hassan Ganjitabar\*

School of Engineering, Newcastle University,  
Newcastle upon Tyne, NE1 7RU, UK

(Dated: January 10, 2022)

Anisotropic Special Relativity (ASR) is the relativistic theory of nature with a preferred direction in space-time. By relaxing the *full-isotropy* constraint on space-time to the *preference of one direction*, we get a perturbative modification of the Minkowski metric as  $g_{\mu\nu} = \eta_{\mu\nu} + b\epsilon_{\mu\nu}$  leading to an extension to the geometrical objects such as line element. The symmetry group in ASR is obtained to be the space-time translations together with the group of 2-dimensional complex special linear transformations,  $SL(2, C)$ , generated by a complexification of  $A = K_x + K_y$  and  $B = J_y - J_x$  and  $C = K_z$ , in the case of  $z$ -direction preference. A procedure to make the anisotropic quantum field theory is provided where the Lorentz invariant Lagrangians need to be replaced with their ASR version in which the inner product of any pair of covariant/contravariant indices is mediated by the anisotropic metric  $g_{\mu\nu}$ .

**PACS numbers:** 11.10.-z, 03.30.+p, 11.30.Cp

## INTRODUCTION

One of the widely accepted fundamental theories of physics is Special Relativity (SR) in which Quantum Field Theory (QFT) and Standard Model (SM) of the universe are grounded. In SR, some quantities, like the line element  $ds^2 = (dt^2 - dx^2 - dy^2 - dz^2)$ , are invariant under the Lorentz group transformations. Despite the very strict restrictions on departure from Lorentz symmetry [1–3], the Lorentz Invariance Violation (LIV) idea [4, 5] is still popular. One of the scenarios for LIV is Standard Model Extension (SME) [6] in which the Lorentz violation is allowed by adding non-relativistic terms to the Lagrangian. On the other side, there are theories wherein the LIV does not require the complete breakdown of relativistic symmetry. The examples of latter are perturbation approach of Coleman and Glashow [7] and Very Special Relativity (VSR) proposed by Cohen and Glashow [8].

There are two more theories allowing for LIV, Finslerian structure of the space-time [9] and Non-Commutative Quantum Field Theory [10], in context of which VSR can be realised [11, 12]. In fact, a deformation of VSR symmetry can be seen in the case of a Finsler space-time in which the fundamental metric  $g$  can be obtained from a function  $F = (\eta_{\mu\nu} dx^\mu dx^\nu)^{(1-b)/2} (n_\alpha dx^\alpha)^b$  using  $g_{\mu\nu} = \frac{1}{2} \partial_\mu \partial_\nu F^2$ , where  $\eta_{\mu\nu}$  is the Minkowski metric of SR,  $n^\alpha = (1, 0, 0, 1)$  is a null vector and  $b$  is a constant parameter. VSR suggests a subgroup of the Lorentz group,  $SIM(2)$ , in addition to the space-time translations to be the maximal symmetry of nature wherein one of the three spatial directions is preferred. The anisotropy of space is one of the possible scenarios for LIV, whose upper bound has already been investigated by analysing the cosmological data [13] as well as performing terres-

trial experiments [14, 15].

On the track of building an anisotropic version of SR, Bogoslovsky suggested the Finslerian line element  $ds = (\eta_{\mu\nu} dx^\mu dx^\nu)^{(1-b)/2} (n_\alpha dx^\alpha)^b$  which is not SR invariant but VSR invariant [16, 17]. In this letter, we ask if there may be any other possible form of line element that is invariant under a reduced Lorentz symmetry but not under the transformations of the full Lorentz group?

## RESULTS AND DISCUSSION

We obtain such a line element by easing the isotropy constraint on the Minkowski metric, but considering a preferred direction,  $n^\mu$ , in space-time instead. In an anisotropic theory we need non-zero off-diagonal elements in the metric. But, the only part of the Minkowski metric we are promoting here is its  $x - y$  block because we want  $n^\mu$  to be yet invariant. The promotion of the  $x - y$  block must be done in a symmetric way as we are not looking for a full isotropy breakdown; the isotropy in  $x$ - $y$  plane is still reserved. A modification of Minkowski metric as

$$\eta_{\mu\nu} \rightarrow g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1+b & -b & 0 \\ 0 & -b & -1+b & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad (1)$$

is showing such a desirable anisotropy in space-time with the  $t$ - and  $z$ -elements being left as same as before, leading to a preferred direction determined by the null vector  $n^\mu = (1, 0, 0, 1)$ . By Anisotropic Special Relativity (ASR) we mean a relativistic theory in which the inner product of a pair of covariant/contravariant indices is mediated by the anisotropic metric  $g$ .

Applying metric (1) gives our version of Lorentz violating line element as

$$ds^2 = (dt^2 - dx^2 - dy^2 - dz^2) + b(dx - dy)^2 \quad (2)$$

which can be considered as a perturbation of the Lorentz invariant line element with  $b$  factor to be a constant parameter controlling the perturbation. Obviously, setting  $b = 0$  turns the situation back to the full Lorentz symmetry. The metric  $g$  itself can be also written as a perturbation of the Minkowski metric

$$g_{\mu\nu} = \eta_{\mu\nu} + b\epsilon_{\mu\nu} \quad (3)$$

where

$$\epsilon_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (4)$$

or can be obtained through the Finsler approach using a function of the form

$$F = \sqrt{(\eta_{\mu\nu} + b\epsilon_{\mu\nu})dx^\mu dx^\nu} \quad (5)$$

Moreover, it can be checked that the metric introduced above is compatible with the  $SL(2, C)$  symmetry group. Starting from the line element invariance principle  $g'_{\mu\nu}dx'^\mu dx'^\nu = g_{\mu\nu}dx^\mu dx^\nu$  and considering  $\Lambda$  as the most general transformation of the theory (excluding the space-time translations), one can write  $\Lambda^T(\Lambda g \Lambda^T)\Lambda = g$  in the matrix form. Defining  $\Lambda = e^\omega$ , and in terms of infinitesimal transformation,  $\Lambda \simeq \mathbb{1} + \omega$ , we have  $(\omega^T + \omega)g(\omega^T + \omega) = g$  suggesting  $\omega$  as

$$\omega = \begin{pmatrix} 0 & \alpha & \alpha & \xi \\ \alpha & 0 & \omega_{12} & \omega_{13} \\ \alpha & -\omega_{12} & 0 & \omega_{23} \\ \xi & -\omega_{13} & -\omega_{23} & 0 \end{pmatrix} \quad (6)$$

where  $\alpha$  and  $\xi$  are two constant parameter. The metric  $g$  given by (1) has been used in the last step. Eq. (6) can be written as  $\omega = -i\alpha(K_x + K_y) - i\xi K_z + i\omega_{23}J_x - i\omega_{13}J_y + i\omega_{12}J_z$  where  $K_i$  and  $J_i$  are respectively the boost and rotation generators of the Lorentz group. The appearance of  $K_x + K_y$  and its commutation with  $K_z$  constraints the parameters  $\omega_{13}$  and  $\omega_{23}$  to be equal, say  $\omega_{13} = \omega_{23} = \beta$ , and sets  $\omega_{12} = 0$  in order to keep the algebra closed. Therefore, we can rewrite (6) as  $\omega = -i\alpha A - i\beta B - i\xi C$  with the generators  $A = K_x + K_y$ ,  $B = J_y - J_x$  and  $C = K_z$  satisfying the commutator relations  $[A, B] = i2C$ ,  $[A, C] = -iB$  and  $[B, C] = iA$ . A complexification of these generators in the form of

$$G^+ = \frac{A + iB}{\sqrt{2}}, \quad G^- = \frac{A - iB}{\sqrt{2}}, \quad X = 2C \quad (7)$$

gives the below algebra

$$[G^+, G^-] = X, \quad [X, G^\pm] = \pm 2G^\pm \quad (8)$$

which is the algebra of the group of complex special linear transformation in two dimension,  $SL(2, C)$ .

Since  $SL(2, C)$  has no abelian minimal subgroup, we can expect non-trivial irreducible representations in ASR. As a matter of fact,  $SL(2, C)$  is isomorphic to the entire Lorentz group,  $SO(1,3)$ , which in turn means that the representations of ASR not only do exist but also are the same as those of the full Lorentz group.

As another consequence of our metric  $g$  given by (1), we can infer to introducing a Lorentz violating term into the fermionic Lagrangian to explain the neutrino mass in a natural way without violating the lepton number or adding sterile right-handed neutrinos. This has been discussed for the first time in [18, 19]; however, we ask if we can add a different Lorentz violating term as an implication of our metric  $g$  which preserve the features. We consider the initially massless Majorana Lagrangian of neutrino and modify it just by applying metric  $g_{\mu\nu}$  to lower the index of one of the contravariant vectors e.i.  $\mathcal{L}^{ASR} = i\bar{\nu}_M \gamma^\alpha g_{\alpha\beta} \partial^\beta \nu_M$  leading to

$$\mathcal{L}^{ASR} = \mathcal{L}^{SR} + ib\bar{\nu}_M(\gamma^1 - \gamma^2)(\partial^1 - \partial^2)\nu_M \quad (9)$$

where  $\mathcal{L}^{SR}$  is the massless Majorana Lagrangian of neutrino in SR given by  $i\bar{\nu}_M(\gamma^0\partial^0 - \gamma^1\partial^1 - \gamma^2\partial^2 - \gamma^3\partial^3)\nu_M$  with  $\nu_M$  to be the four-component Majorana neutrino and  $\gamma^\alpha$ ,  $\alpha = 0 - 3$ , to be the Dirac matrices. Eq. (9) gives the dispersion relation as  $(m^2)^{ASR} = (p_\alpha p^\alpha)^{SR} + b(p_x - p_y)^2$  for which  $(p_\alpha p^\alpha)^{SR} = (m^2)^{SR}$  is assumed to be zero; even in such a case, it is still possible for  $(m^2)^{ASR}$  to be not zero due to the appearance of the perturbative term  $b(p_x - p_y)^2$ . However, for the neutrinos moving parallel to the  $z$  direction or, more generally, those carrying the same momentum in  $x$ - and  $y$ -direction,  $p_x = p_y$ , the corrected mass yet remains zero. In total, the neutrino mass in anisotropic space-time will depend on the direction of its movement which in turn means that the neutrino flavour oscillation allowed by anisotropic theories like ASR and VSR [20] could be direction-dependent as well.

It can also be noticed that the field theory (FT) obtained in ASR remains *local* in contrast to the non-local FT emerging from the correction term introduced by Cohen and Glashow in VSR [18]. The term they suggested to correct the equations of motion with is  $\frac{m^2 \eta_{\alpha\beta} \gamma^\alpha n^\beta}{2n \cdot \partial} \nu_M$  where the appearance of  $\partial$  operator in the denominator makes the theory non-local. However, ASR-FT remains local as only the lower orders of the derivative operator appear in the Lagrangian.

In general, the mass of any field in ASR can be modified just similar to the line element in Eq. (2) i.e.

$$(m^2)^{SR} \rightarrow (m^2)^{ASR} = (m^2)^{SR} + b(p_x - p_y)^2 \quad (10)$$

where  $m^{SR}$  is the mass of the field in SR given by  $(E^2 - p_x^2 - p_y^2 - p_z^2)$  with  $E$  and  $p^i$ ,  $i = 1 - 3$ , to be respectively the energy and momentum of the field.

In ASR, we just need to lower the indices of any contravariant vector or tensor with the anisotropic metric  $g$ . This gives the inner product of two arbitrary vectors  $v^\mu$  and  $u^\mu$  as

$$(v_\mu u^\mu)^{ASR} = (v_\mu u^\mu)^{SR} + b(v^1 - v^2)(u^1 - u^2) \quad (11)$$

where  $(v_\mu u^\mu)^{SR}$  is the inner product of the vectors in SR. In the case that at least one of the two vectors is parallel to the preferred direction ( $v^1 = v^2 = 0$  or  $u^1 = u^2 = 0$ ) or its  $x$ - and  $y$ -component are equal ( $v^1 = v^2$  or  $u^1 = u^2$ ), the inner product reduces down to its SR version.

In order to make an anisotropic quantum field theory, one can promote any Lagrangian by replacing the isotropic inner product of any pair of covariant/contravariant indices with its anisotropic version. For example, the Dirac Lagrangian for the fermions can be modified to

$$\mathcal{L}_D^{ASR} = \mathcal{L}_D^{SR} + ib\bar{\psi}(\gamma^1 - \gamma^2)(\partial^1 - \partial^2)\psi \quad (12)$$

a consequence of which is the extension of the dispersion relation as (10). Similarly, in the electrodynamics Lagrangian  $\mathcal{L}_E^{SR} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$  we just need to lower the indices using  $g$  i.e.  $\mathcal{L}_E^{ASR} = -\frac{1}{4}g_{\mu\alpha}g_{\nu\beta}F^{\alpha\beta}F^{\mu\nu}$  resulting in equations of motion as

$$(\partial_\mu F^{\mu\nu})^{SR} + b\epsilon_{\mu\alpha}\partial^\alpha F^{\mu\nu} = 0 \quad (13)$$

from which the anisotropically modified Maxwell equations can be derived.

The shape of gauge symmetry in ASR remains the same as in SR

$$\begin{cases} \psi \rightarrow e^{iq\vartheta}\psi \\ \partial_\mu \rightarrow D_\mu = \partial_\mu - iqA_\mu \\ A_\mu \rightarrow A_\mu + \partial_\mu\vartheta \end{cases} \quad (14)$$

with the difference that, here, all the covariant indices are made from the contravariant ones by applying the metric  $g$ . For instance, the ASR version of  $\partial_\mu$  and  $A_\mu$  are

$$\begin{cases} \partial_\mu^{ASR} = \partial_\mu^{SR} + b(0, \partial^1 - \partial^2, \partial^2 - \partial^1, 0) \\ A_\mu^{ASR} = A_\mu^{SR} + b(0, A^1 - A^2, A^2 - A^1, 0) \end{cases} \quad (15)$$

## CONCLUSION

In this letter, we proposed an anisotropic relativistic theory, ASR, as an alternative to the special theory of relativity. ASR can be formalised by replacing the Minkowski metric with a perturbative anisotropic metric in which the space-time isotropy constraint has been replaced with the preference of a direction determined by the null vector  $n^\mu$ . On the one hand, the partial breakdown of Lorentz symmetry is allowed while on the other hand, the symmetry group  $SL(2, C)$  guarantees the irreducible representations of the theory to be just the same

as those of SR. Despite a direction-dependent modification to the dispersion relation, the QFT built up in ASR remains local.

As one of the interesting applications of ASR, we can infer to redesigning of neutrino oscillation experiments. ASR suggests the neutrino mass, hence the neutrino oscillation, to be direction dependent; so, designing a neutrino oscillation experiment in which the direction of propagation of neutrinos can sweep different spatial directions could be of special interest. In addition, as the transformation of ASR, presented by  $SL(2, C)$  group, are direction dependent, we expect most of the relativistic phenomena like length contraction, time dilation, Doppler effect and aberration of light, etc. to be direction dependent too. A possible origin for the space-time anisotropy could be the anisotropic mass/energy distribution in the universe, which can be reflected in cosmic microwave background. This means that different locations in the universe could be expected to experience different level or direction of anisotropy. In such a case, a universal version of ASR can be achieved by promoting the perturbative parameter  $b$  from a constant to a field.

---

\* Hassan.ganjitabar@newcastle.ac.uk

- [1] V. A. Kostelecky and N. Russell, "Data tables for lorentz and cpt violation," *Reviews of Modern Physics*, vol. 83, no. 1, pp. 11–32, 2011.
- [2] D. Mattingly, "Modern tests of lorentz invariance," *Living Reviews in Relativity*, vol. 8, no. 1, p. 5, 2005.
- [3] S. Liberati, "Tests of lorentz invariance: a 2013 update," *Classical and Quantum Gravity*, vol. 30, no. 13, p. 133001, 2013.
- [4] D. A. Kirzhnits and V. A. Chechin, "The ultra-high-energy cosmic rays and a possible generalization of the relativistic theory," *Yadern Fiz*, vol. 15, no. 5, pp. 1051–1059, 1972.
- [5] T. G. Pavlopoulos, "Breakdown of lorentz invariance," *Physical Review*, vol. 159, no. 5, pp. 1106–1110, 1967.
- [6] D. Colladay and V. A. Kostelecký, "Lorentz-violating extension of the standard model," *Physical Review D*, vol. 58, no. 11, p. 116002, 1998.
- [7] S. Coleman and S. L. Glashow, "High-energy tests of lorentz invariance," *Physical Review D*, vol. 59, no. 11, p. 116008, 1999.
- [8] A. G. Cohen and S. L. Glashow, "Very special relativity," *Physical Review Letters*, vol. 97, no. 2, p. 021601, 2006.
- [9] P. Antonelli, R. Ingarden, and M. Matsumoto, *The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology*. 1993.
- [10] M. R. Douglas and N. A. Nekrasov, "Noncommutative field theory," *Rev. Mod. Phys.*, vol. 73, pp. 977–1029, Nov 2001.
- [11] L. Zhang and X. Xue, "The finsler type of space-time realization of deformed very special relativity," 2012.

- [12] M. M. Sheikh-Jabbari and A. Tureanu, “Realization of cohen-glashow very special relativity on noncommutative space-time,” *Physical Review Letters*, vol. 101, no. 26, p. 261601, 2008.
- [13] E. Komatsu, J. Dunkley, M. R. Nolta, C. L. Bennett, B. Gold, G. Hinshaw, N. Jarosik, D. Larson, M. Limon, L. Page, D. N. Spergel, M. Halpern, R. S. Hill, A. Kogut, S. S. Meyer, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright, “Five-year wilkinson microwave anisotropy probe observations: Cosmological interpretation,” *The Astrophysical Journal Supplement Series*, vol. 180, no. 2, pp. 330–376, 2009.
- [14] C. Eisele, A. Y. Nevsky, and S. Schiller, “Laboratory test of the isotropy of light propagation at the  $10^{-17}$  level,” *Physical Review Letters*, vol. 103, no. 9, p. 090401, 2009.
- [15] B. Altschul, “Bounding isotropic lorentz violation using synchrotron losses at lep,” *Physical Review D*, vol. 80, no. 9, p. 091901, 2009.
- [16] G. Y. Bogoslovsky, “A special-relativistic theory of the locally anisotropic space-time,” *Il Nuovo Cimento B (1971-1996)*, vol. 40, no. 1, pp. 99–115, 1977.
- [17] G. Y. Bogoslovsky, “A special-relativistic theory of the locally anisotropic space-time,” *Il Nuovo Cimento B (1971-1996)*, vol. 40, no. 1, pp. 116–134, 1977.
- [18] A. Cohen and S. Glashow, “A lorentz-violating origin of neutrino mass?,” 2006.
- [19] A. E. Bernardini, “Dirac neutrino mass from the beta-decay end point modified by the dynamics of a lorentz-violating equation of motion,” *Physical Review D*, vol. 75, no. 9, p. 097901, 2007.
- [20] J. Alfaro, P. González, and R. Ávila, “Electroweak standard model with very special relativity,” *Physical Review D*, vol. 91, no. 10, p. 105007, 2015.