Merger rate of charged black holes from the two-body dynamical capture

Lang Liu,^a Sang Pyo Kim^a

^aDepartment of Physics, Kunsan National University, Kunsan 54150, Korea

E-mail: liulang915871@gmail.com, sangkim@kunsan.ac.kr

Abstract. We consider the two-body dynamical capture of black holes carrying U(1) charge which can not only correspond to electric or magnetic charge but also have other physical interpretations such as dark or hidden charge. In the low-velocity and weak-field regime, we study gravitational and electromagnetic radiations from point masses with U(1) charges in a hyperbolic orbit, and we develop a formalism to derive the merger rate of charged black holes from the two-body dynamical capture. We apply the formalism to find the effects of the charge-to-mass ratio on the merger rate for possible different cases and discover that the effects depend on the models.

Contents

1	Introduction	1
2	Gravitational and electromagnetic radiations from point masses in a hyperbolic orbit	- 2
3	Merger rate of charged black holes from the two-body dynamical capture	5
4	 Effects of charge-to-mass ratio on the merger rate 4.1 Astrophysical black hole 4.2 Primordial black hole 	7 7 8
5	Conclusion	10
A	Appendix A.1 Electromagnetic dipole and quadrupole radiations	10 10

1 Introduction

Nearly one hundred years after the theoretical prediction of gravitational waves (GWs) in the general relativity (GR) by Einstein [1, 2], the first direct measurement of GWs [3] by the Advanced Laser Interferometer Gravitational-Wave Observatory (LIGO) detectors established gravitational wave astronomy. The LIGO-Virgo detection opened a new window to observe the universe and marked the dawn of multi-messenger astronomy [4–6]. So far, the LIGO-Virgo detection results have already provided extremely accurate confirmation that all gravitational merger events can be described by GR [7–9].

The no-hair theorem of black holes (BHs) in GR states that four-dimensional stationary BHs in the Einstein-Maxwell theory can be completely described by a Kerr-Newman metric [10-12], and the theorem allows one to characterize all BH solutions in asymptotically flat spacetime by three physical quantities: the mass, spin and charge. When Hawking radiation is not considered, in comparison with Schwarzschild BHs, charged BHs emit not only gravitational radiation but also electromagnetic radiation, and have attracted much attention [13-32]. Recently, assuming that the influence of BH spins can be neglected, Bozzola et al found that the charge-to-mass ratios of up to 0.3 are compatible with Binary BH merger event GW150914 [21]. Later, they also reported general relativistic simulations of the inspiral and merger of non-spinning charged binary BHs [27].

In the universe, the two-body dynamical capture is a fairly common and effective way to form the binary BH systems including astrophysical black holes (ABHs) and primordial black holes (PBHs) [33, 34]. Therefore, it is important and meaningful to work out the merger rate of charged BHs from the two-body dynamical capture. To do so, we analytically find gravitational and electromagnetic radiations from a binary of point masses with charges in a hyperbolic orbit, and calculate the merger rate of BHs from the two-body dynamical capture with charges and a general mass function by taking into account gravitational and electromagnetic radiation. The U(1) charge considered in this paper can correspond to the following physical interpretations: (1) electric charges, (2) magnetic charges [35–37], (3) hidden or dark charges interacting with dark electromagnetism [38-41], (4) modified theories of gravity with additional scalar or vector fields [42-44], and (5) the fifth force [45-47].

The paper is organized as follows. In Section 2, in the low-velocity and weak-field regime, we calculate gravitational and electromagnetic radiations from point masses with U(1) charges in a hyperbolic orbit. In Section 3, we develop a formalism to derive the merger rate of charged BHs from the two-body dynamical capture via gravitational and electromagnetic radiations. In Section 4, we show the effects of the charge-to-mass ratio on the merger rate of ABH and PBH binaries from the two-body dynamical capture. Finally, we summarize our results and conclude with physical implications in the last section.

In this paper, we set $G = c = 4\pi\varepsilon_0 = \frac{\mu_0}{4\pi} = 1$ unless otherwise specified. Although the U(1) charges are intended to have different physical interpretations, we will refer to their quantities by using "electromagnetic" for the sake of simplification through this paper.

2 Gravitational and electromagnetic radiations from point masses in a hyperbolic orbit

We study an encounter of two unbounded massive charges and the gravitational and electromagnetic radiations from the encounter. The unbound system under gravitational and electric forces follows a hyperbolic orbit and dominantly emits gravitational and electromagnetic radiations near the closest approach and loses energy and angular momentum. To describe the dynamical evolution, we assume that the orbit lies in x-y plane, and that the coordinates of the point masses m_1 with charge Q_1 and m_2 with charge Q_2 are $(d_1 \cos \psi, d_1 \sin \psi)$ and $(-d_2 \cos \psi, -d_2 \sin \psi)$, respectively. Choosing the origin at the center of mass, we have

$$d_1 = \left(\frac{m_2}{m_1 + m_2}\right) d, \quad d_2 = \left(\frac{m_1}{m_1 + m_2}\right) d.$$
 (2.1)

As shown in Fig. 1, the equation for a hyperbolic orbit is

$$d = \frac{a(e^2 - 1)}{1 - e\cos\psi},$$
(2.2)

where a and e, which can be interpreted as the semi-major axis and eccentricity, and are defined as

$$a \equiv \frac{m_1 m_2 (1 - \lambda)}{2E}, \quad e \equiv \left(1 + \frac{2EL^2}{(m_1 + m_2) m_1 m_2 (1 - \lambda)^2}\right)^{1/2}.$$
 (2.3)

Here, $\lambda \equiv \frac{Q_1 Q_2}{m_1 m_2}$ represents the ratio of the Coulomb force to the gravitational force, E is the total energy of charged binary system including the gravitational, electrostatic and kinetic energy, and L is the angular momentum of binary system. And the angular velocity along the orbit is given by

$$\dot{\psi} = \frac{\left[(m_1 + m_2) a \left(e^2 - 1\right) (1 - \lambda)\right]^{1/2}}{d^2}.$$
(2.4)

Firstly, we calculate the gravitational radiation from the point masses m_1 with charge Q_1 and m_2 with charge Q_2 in the hyperbolic orbit. The non-vanishing second mass moment is a 2×2 matrix that is given by

$$M_{ab} = \mu d^2 \begin{pmatrix} \cos^2 \psi & \sin \psi \cos \psi \\ \sin \psi \cos \psi & \sin^2 \psi \end{pmatrix}_{ab},$$
(2.5)



Figure 1. A schematic picture of a hyperbolic orbit.

where subscripts (a, b = 1, 2) refer to indices in the x-y plane and $\mu = m_1 m_2/(m_1 + m_2)$ is the reduced mass. Note that M_{ij} is reducible while the traceless quadrupole moment $Q_{ij} \equiv M_{ij} - \frac{1}{3}\delta_{ij}M_{kk}$ is irreducible. Following [48], the rate of energy loss from a system by gravitational quadrupole radiation is given by

$$\frac{dE_{GW}^{quad}}{dt} \equiv -\frac{1}{5} \left(\ddot{Q}_{ij} \ddot{Q}_{ij} \right) = -\frac{2}{15} \left[\left(\ddot{M}_{11} + \ddot{M}_{22} \right)^2 - 3 \left(\ddot{M}_{11} \ddot{M}_{22} - \ddot{M}_{12}^2 \right) \right].$$
(2.6)

The derivatives of the components of the second mass moment tensor can be calculated by the aid of Eqs. (2.2) and (2.4); putting those expressions into Eq. (2.6), we obtain

$$\frac{dE_{GW}^{quad}}{dt} = -\frac{4(1-\lambda)^3 m_1^2 m_2^2 (m_1 + m_2) (e\cos(\psi) - 1)^4 \left(11e^2 \cos(2\psi) + 13e^2 - 48e\cos(\psi) + 24\right)}{15a^5 (e^2 - 1)^5}$$
(2.7)

The post-Newtonian formalism gives the rate of angular momentum emission due to gravitational quadrupole radiation as [49]

$$\frac{dL_{GW}^{i,quad}}{dt} \equiv -\frac{2}{5} \epsilon^{ikl} \left(\ddot{Q}_{ka} \ddot{Q}_{la} \right) = -\frac{2}{5} \epsilon^{ikl} \left(\ddot{M}_{ka} \ddot{M}_{la} \right), \qquad (2.8)$$

where ϵ^{ikl} is the Levi-Civita symbol ($\epsilon^{123} = 1$). As the orbit is assumed to lie in the *x-y* plane and the gravitational and Coulomb forces keep the same orbital plane, we have $L^z = L$ and $L^x = L^y = 0$. Substituting Eqs. (2.2) and (2.4) into the components of the second mass moment tensor for the unbound system in the hyperbolic orbit, the non-vanishing rate of angular momentum emission due to gravitational quadrupole radiation is

$$\frac{dL_{GW}^{quad}}{dt} = \frac{dL_{GW}^{z,quad}}{dt} = -\frac{8(1-\lambda)^{5/2}m_1^2m_2^2\sqrt{m_1+m_2}}{5a^{7/2}(e^2-1)^{7/2}}\sin^2(\psi)(e\cos(\psi)-1)^2 \times (e(\cos(2\psi)+3)-4\cos(\psi))(e(3\cos(2\psi)+4)-8\cos(\psi)).$$
(2.9)

Secondly, we compute the energy and angular momentum loss rates due to electromagnetic dipole radiation. In our reference frame, the electric dipole is given by

$$\boldsymbol{p} \equiv Q_1 \boldsymbol{d_1} + Q_2 \boldsymbol{d_2} = \frac{m_2 Q_1 - m_1 Q_2}{m_1 + m_2} (d\cos\psi \hat{\boldsymbol{x}} + d\sin\psi \hat{\boldsymbol{y}}).$$
(2.10)

The overall factor can be written as $\mu(\frac{Q_1}{m_1} - \frac{Q_2}{m_2})$. Then, according to [50, 51], the energy and angular momentum loss rates due to electromagnetic dipole radiation are

$$\frac{dE_{EM}^{dip}}{dt} = -\frac{2\ddot{p}^2}{3} = -\frac{2(1-\lambda)^2 \left(m_2 Q_1 - m_1 Q_2\right)^2 \left(e\cos(\psi) - 1\right)^4}{3a^4 \left(e^2 - 1\right)^4},$$
(2.11)

$$\frac{dL_{EM}^{dip}}{dt} = -\frac{2}{3}(\dot{p}_2\ddot{p}_1 - \dot{p}_1\ddot{p}_2) = -\frac{2(1-\lambda)^{3/2}(m_2Q_1 - m_1Q_2)^2(1 - e\cos(\psi))^3}{3a^{5/2}(e^2 - 1)^{5/2}\sqrt{m_1 + m_2}}.$$
 (2.12)

In Appendix A.1, we calculate the quadrupole term of electromagnetic radiation and find that its contribution is always smaller than the quadrupole term of gravitational radiation. The total energy and angular momentum radiated due to gravitational quadrupole radiation and electromagnetic dipole radiation are then the integral of Eqs. (2.7), (2.9), (2.11) and (2.12) over the orbit. Notice the asymptotes of the hyperbolic orbit are $y = \pm \tan(\psi_0)x$, where ψ_0 is determined by $\cos(\psi_0) = 1/e$, i.e, $\psi \in (\psi_0, 2\pi - \psi_0)$; thus we find

$$\Delta E_{GW}^{quad} = \int_{-\infty}^{\infty} \frac{dE_{GW}^{quad}}{dt} dt = \int_{\psi_0}^{2\pi - \psi_0} \frac{dE_{GW}^{quad}}{dt} (\frac{d\psi}{dt})^{-1} d\psi = -\frac{2(1-\lambda)^{5/2} m_1^2 m_2^2 \sqrt{m_1 + m_2}}{45a^{7/2} (e^2 - 1)^{7/2}} \\ \times \left(3 \left(37e^4 + 292e^2 + 96\right) \left(\pi - \cos^{-1}\left(\frac{1}{e}\right)\right) \pi + \sqrt{1 - \frac{1}{e^2}}e \left(673e^2 + 602\right)\right),$$

$$(2.13)$$

$$\Delta L_{GW}^{quad} = \int_{-\infty}^{\infty} \frac{dL_{GW}^{quad}}{dt} dt = \int_{\psi_0}^{2\pi - \psi_0} \frac{dL_{GW}^{quad}}{dt} (\frac{d\psi}{dt})^{-1} d\psi = \frac{8(1 - \lambda)^2 m_1^2 m_2^2}{5a^2 (e^2 - 1)^2} \\ \times \left(\left(7e^2 + 8\right) \left(\pi - \cos^{-1}\left(\frac{1}{e}\right)\right) + \frac{\sqrt{1 - \frac{1}{e^2}} \left(21e^4 - 10e^2 + 4\right)}{e^3} \right),$$

$$\Delta E_{EM}^{dip} = \int_{-\infty}^{\infty} \frac{dE_{EM}^{dip}}{dt} dt = \int_{\psi_0}^{2\pi - \psi_0} \frac{dE_{EM}^{dip}}{dt} (\frac{d\psi}{dt})^{-1} d\psi = -\frac{(1 - \lambda)^{3/2} (m_2 Q_1 - m_1 Q_2)^2}{3a^{5/2} (e^2 - 1)^{5/2} \sqrt{m_1 + m_2}} \\ \times \left(\pi \left(e^2 + 2\right) + 6e\sqrt{1 - \frac{1}{e^2}} + 2\left(e^2 + 2\right) \csc^{-1}(e)\right),$$

$$(2.14)$$

$$(-2) + 6e\sqrt{1 - \frac{1}{e^2} + 2(e^2 + 2)\csc^{-1}(e)}$$

(2.15)

$$\Delta L_{EM}^{dip} = \int_{-\infty}^{\infty} \frac{dL_{EM}^{dip}}{dt} dt = \int_{\psi_0}^{2\pi - \psi_0} \frac{dL_{EM}^{dip}}{dt} (\frac{d\psi}{dt})^{-1} d\psi = -\frac{4(1 - \lambda) (m_2 Q_1 - m_1 Q_2)^2}{3a (e^2 - 1) (m_1 + m_2)} \times \left(e\sqrt{1 - \frac{1}{e^2}} - \cos^{-1}\left(\frac{1}{e}\right) + \pi \right).$$
(2.16)

In the case of either charge neutral masses $(Q_1 = Q_2 = \lambda = 0)$ or the same charge to mass ratio $(\frac{Q_1}{m_1} = \frac{Q_2}{m_2})$, we find that the electric dipole radiation vanishes, $\Delta E_{EM}^{dip} = \Delta L_{EM}^{dip} = 0$, and then the total energy and angular momentum are radiated through the gravitational quadrupole radiation and agree with the results given by Hansen [52]. In the parabolic limit, i.e., $e \to 1$, Eqs. (2.13), (2.14), (2.15) and (2.16) can be simplified as

$$\Delta E_{EM}^{dip} = -\frac{\pi (1-\lambda)^{3/2} \left(m_2 Q_1 - m_1 Q_2\right)^2}{2\sqrt{2}\sqrt{m_1 + m_2} r_p^{5/2}},$$
(2.17)

$$\Delta E_{GW}^{quad} = -\frac{85\pi (1-\lambda)^{5/2} m_1^2 m_2^2 \sqrt{m_1 + m_2}}{12\sqrt{2}r_p^{7/2}},$$
(2.18)

$$\Delta L_{EM}^{dip} = -\frac{2\pi (1-\lambda) \left(m_2 Q_1 - m_1 Q_2\right)^2}{3 \left(m_1 + m_2\right) r_p},\tag{2.19}$$

$$\Delta L_{GW}^{quad} = -\frac{6\pi (1-\lambda)^2 m_1^2 m_2^2}{r_p^2},$$
(2.20)

where $r_p \equiv a(e-1)$ is the distance of periastron. Another way to estimate the total energy loss is to approximate the trajectory of a close encounter by taking the e = 1 limit of the elliptical orbit since the radiation is most dominant when the orbit is closest to the focus; thereby the total energy loss is given by $\Delta E = T \langle \frac{dE}{dt} \rangle$ where T is orbital period and $\langle \frac{dE}{dt} \rangle$ is the time-averaged energy loss rate of the binary in the Keplerian orbit which is given by [51]. The gravitational and electromagnetic radiations from binary BHs with electric and magnetic charges were found in [53–55].

3 Merger rate of charged black holes from the two-body dynamical capture

If two charged BHs get closer and closer, the energy loss due to gravitational and electromagnetic radiations can exceed the orbital kinetic energy, and thus the unbound system cannot escape to infinity anymore and form a binary with a negative orbital energy. This binary immediately merges through consequent gravitational and electromagnetic radiations[51]. From such a process, we can estimate the cross section and calculate the rate coefficient.

Let us consider the interaction of two charged BHs with masses m_1 , m_2 and charges Q_1 , Q_2 and assume that they have an initial relative velocity v, the distance of periastron r_p and the impact parameter b. We can approximate the trajectory of a close encounter by the hyperbolic with $e \to 1$ since when the two charged BHs pass by closely, the true trajectory is physically indistinguishable from a parabolic one near the periastron, in which gravitational and electromagnetic radiations dominantly occur. According to Sec. 2 and Appendix A.1, the total energy loss due to the electromagnetic dipole and quadrupole radiations and the gravitational quadrupole radiation by the close-encounter can be evaluated by using $e \to 1$ and the definition of $r_p \equiv a(e-1)$,

$$\Delta E = \Delta E_{EM}^{dip} + (1 + \Lambda) \,\Delta E_{GW}^{quad},\tag{3.1}$$

where we denote $\Lambda = \frac{\mu^2 (Q_1/m_1^2 + Q_2/m_2^2)^2}{4}$ for short and ΔE_{EM}^{dip} , ΔE_{GW}^{quad} are given by Eqs. (2.17) and (2.18). Similarly, the total angular momentum loss due to the electromagnetic dipole and quadrupole radiations and the gravitational quadrupole radiation is given by

$$\Delta L = \Delta L_{EM}^{dip} + \left(1 + \frac{\Lambda}{2}\right) \Delta L_{GW}^{quad}, \qquad (3.2)$$

where ΔL_{EM}^{dip} , ΔL_{GW}^{quad} are given by Eqs. (2.19) and (2.20). The relation between r_p and b is $b^2 = r_p^2 + \frac{2(1-\lambda)(m_i+m_j)r_p}{v^2}$. In the limit of a strong gravitational and electromagnetic focusing (i.e. $r_p \ll b$), then the distance of closest approach r_p is given by

$$r_p = \frac{b^2 v^2}{2(1-\lambda)(m_i + m_j)}.$$
(3.3)

The condition for the two charged BHs to form a binary is that the total energy loss due to radiations is larger than the kinetic energy $\mu v^2/2$, i.e.,

$$\Delta E + \frac{\mu v^2}{2} < 0. \tag{3.4}$$

Equations (3.1), (3.3) and (3.4) yield the maximum impact parameter b_{max} for the charged BHs to form a bound system, and then we obtain the merging cross section as $\sigma = \pi b_{max}^2$, where b_{max} is determined by

$$\frac{2\pi(1-\lambda)^4 \left(m_1+m_2\right)^2 \left(m_2 Q_1-m_1 Q_2\right)^2}{b_{max}^5 v^5} + \frac{170\pi(1-\lambda)^6 \Lambda m_1^2 m_2^2 \left(m_1+m_2\right)^4}{3b_{max}^7 v^7} = \frac{\mu v^2}{2}.$$
 (3.5)

If the dipole radiation is dominant (i.e. $\Delta E_{EM}^{dip} > (1 + \Lambda) \Delta E_{GW}^{quad}$), the merging cross section is

$$\sigma_{dip} \approx \frac{2^{4/5} \pi^{7/5} (1-\lambda)^{8/5} (m_1 + m_2)^{6/5} (m_1 Q_2 - m_2 Q_1)^{4/5}}{m_1^{2/5} m_2^{2/5} v^{14/5}},$$
(3.6)

whereas if the total quadrupole radiation is dominant (i.e. $\Delta E_{EM}^{dip} < (1 + \Lambda) \Delta E_{GW}^{quad}$), the merging cross section is

$$\sigma_{quad} \approx \frac{\left(\frac{85}{3}\right)^{2/7} 2^{4/7} \pi^{9/7} (1-\lambda)^{12/7} m_1^{2/7} m_2^{2/7} (m_1+m_2)^{10/7}}{v^{18/7}} \left(1+\Lambda\right)^{2/7}, \qquad (3.7)$$

which is consistent with [33]. Notice that the Schwarzschild radius $\sigma_{Sch} \simeq \pi (2Gm)^2 \simeq (v)^{18/7} \sigma_{GW}$, $v \ll 1$ and $\sigma \geq \sigma_{GW}$; thereby the Newtonian approximation is sufficiently accurate no matter whether electromagnetic radiation is dominant or gravitational radiation is dominant. The differential merger rate of charged BHs per comoving volume reads

$$dR = n(m_1, Q_1)n(m_2, Q_2) \langle \sigma v \rangle dm_1 dm_2 dQ_1 dQ_2$$
(3.8)

where $n(m_1, Q_1)$ and $n(m_2, Q_2)$ are the comoving average number density of charged BHs with masses m_1, m_2 and charges Q_1, Q_2 , and $\langle \sigma v \rangle$ denotes the average over relative velocity distribution with $\sigma = \pi b_{max}^2$ given in Eq. (3.5). In this section, we have computed the merging cross section of charged BHs without assuming the origin of those charged BHs.

4 Effects of charge-to-mass ratio on the merger rate

4.1 Astrophysical black hole

In astrophysics, there are two main astrophysical mechanisms for ABH binary formation¹. One channel is isolated (or 'field') binary evolution and the other is dynamical capture in dense stellar environments. Dynamical capture can become absolutely effective in deeply dense stellar environments. Especially, it is shown in Ref. [57] that the vast majority ($\sim 80\%$) of the merger events detected by LIGO-Virgo comes from dynamical capture².

According to the initial mass function that describes the number distribution of stars [59], we assume the mass function of ABHs takes a truncated power-law form as [60]

$$p(m) \propto m^{\zeta} \theta \left(m - m_{\min}\right) \theta \left(m_{\max} - m\right) \tag{4.1}$$

with $\zeta = -1.3$, and the normalization of mass function $\int p(m)dm = 1$ will be taken. Here, we adopt $m_{\min} = 3.0M_{\odot}$ which corresponds to the lower mass bound of ABHs [61], and $m_{\max} = 55M_{\odot}$ which corresponds to the beginning of pair-instability supernova (PISN) gap [62]. We consider the following two simple but different models

- model (a): all ABHs have the same charge-to-mass ratio ι in magnitude and sign,
- model (b): a half of ABHs have the same charge-to-mass ratio ι and the other half ABHs have a charge-to-mass ratio equal in magnitude but oppositely charged.

In the following calculation, the Maxwell-Boltzmann distribution $P(v) \propto v^2 \exp\left(-v^2/v_0^2\right)$ will be used for the velocity distribution of BHs with the most probable velocity $v_0 = 100$ km/s.

To show the effects of charge-to-mass ratio ι on the merger rate of ABH binaries from the two-body dynamical capture, we define a function of ι as

$$\eta(\iota) \equiv \frac{R_{\rm ABH}(\iota)}{R_{\rm ABH}(0)},\tag{4.2}$$

where $R_{ABH}(\iota)$ is the total merger rate of charged ABHs with charge-to-mass ratio ι and $R_{ABH}(0)$ is the total merger rate of ABHs in charge neutral case.

In the model a, we have the number density of charged ABHs

$$n(m,Q) \propto \delta(\frac{Q}{m}-\iota)\frac{p(m)}{m},$$
(4.3)

where δ is the Dirac delta function. In such a case, we have $Q_1/m_1 = Q_2/m_2 = \iota$. Thus, the dipole radiation vanish and the energy loss due to total quadrupole radiation is given by $\Delta E = (1 + \iota^2/4) \Delta E_{GW}^{quad}$. Therefore, according to Eqs. (3.7) and (3.8) and from the definition of η , we obtain

$$\eta(\iota) = (1 - \iota^2)^{12/7} \left(1 + \iota^2/4 \right)^{2/7}.$$
(4.4)

In the model b, we have the number density of charged ABHs

$$n(m,Q) \propto \frac{1}{2} \left(\delta(\frac{Q}{m} - \iota) + \delta(\frac{Q}{m} + \iota) \right) \frac{p(m)}{m}.$$
(4.5)

¹There are others channels like mergers in triple systems assisted by the Kozai-Lidov mechanism (See a recent review [56] for details).

²There are currently some disagreements about the contributions of different channels. It is argued that in Ref. [58] that $\sim 90\%$ is provided by isolated binary evolution channel.

Therefore, the merger events can be divided into two parts. One half of merger events are those binary BHs with the same sign charges while the other half of merger events are those binary BHs with the opposite charges. When the binary BHs have the opposite charges, we have $Q_1/m_1 = -Q_2/m_2 = \iota$ and $\Lambda = \left(1 + \frac{\iota^2(m_1-m_2)^2}{4(m_1+m_2)^2}\right)$. The merging cross section in this case is $\sigma \simeq \pi b_{max}^2$, where b_{max} is a root of

$$\frac{8\pi G^4 \iota^2 \left(\iota^2+1\right)^4 m_1^2 m_2^2 \left(m_1+m_2\right)^2}{b_{max}^5 v^5} + \frac{170\pi G^7 (1+\iota^2)^6 \Lambda m_1^2 m_2^2 \left(m_1+m_2\right)^4}{3 b_{max}^7 v^7} = \frac{\mu v^2}{2}.$$
 (4.6)

Hence, in the model b, from the Eqs. (3.8) and (4.2), we have

$$\eta(\iota) = \frac{1}{2} (1 - \iota^2)^{12/7} \left(1 + \iota^2/4 \right)^{2/7} + \frac{1}{2} \frac{\int \int \int P(v) \sigma v \frac{p(m_1)}{m_1} \frac{p(m_2)}{m_2} dv dm_1 dm_2}{\int \int \int \int P(v) \sigma_0 v \frac{p(m_1)}{m_1} \frac{p(m_2)}{m_2} dv dm_1 dm_2}$$
(4.7)

where

$$\sigma_0 = \frac{2^{4/7} \left(\frac{85}{3}\right)^{2/7} \pi^{9/7} m_1^{2/7} m_2^{2/7} \left(m_1 + m_2\right)^{10/7}}{v^{18/7}} \tag{4.8}$$

represents the merging cross section for charge neutral BHs. From the definition above, we have $\eta(\iota) = \eta(-\iota)$ and $\eta(0) = 1$ for either the model a or b. Therefore, we only need to consider $\iota \in [0, 1]$. In Fig. 2, we plot $\eta(\iota)$ as the function of ι for two models. In the model a, we find that $\eta(\iota)$ decreases as ι increases and $\eta(1) = 0$. For the model b, we exactly show that $\eta(\iota)$ increases as ι increases and reaches the maximum value of $\eta(1) \approx 8.0$.



Figure 2. Left: The plots of $\eta(\iota)$ as the function of ι in the model a (red) and model b (purple). Right: The normalized merger rate of charged PBH $R_{\text{PBH}}(\iota)/R_{\text{PBH}}(0)$ as the function of ι in the model a (red) and model b (purple).

4.2 Primordial black hole

In contrast with ABHs, PBHs are those BHs that are formed in the radiation-dominated era of the early universe due to the collapse of large energy density fluctuations [63–65]. The issue of how to distinguish ABHs and PBHs has been studied in [66–69]. In recent years, PBHs have attracted a lot of attention and have been extensively studied, not only because they can be candidates of dark matter [70], but also because they can be the sources for LIGO-Virgo detections [34, 71] and the seeds for galaxy formation [72–74]. There are two primary mechanisms for PBH binary formation, which can be classified by the cosmological epoch when they work. One mechanism is that the PBH binary would have formed in the late universe [34, 75, 76] while the other is that PBH binary would have formed in the early universe [71, 77–89]. Recently, it is shown in Ref [29] that PBHs can be near-extremally charged. In this subsection, we will investigate the effects of charge-to-mass ratio ι on the merger rate of PBH binaries which would have formed in the late universe. Similarly to ABHs, we also consider two special models by assuming that all PBHs have the same mass $M_{\rm PBH}$:

- model (a): all PBHs have the same charge-to-mass ratio ι in magnitude and sign,
- model (b): a half of PBHs have the same charge-to-mass ratio ι and the other half of PBHs have the same magnitude but with the opposite charges.

In the model a, the number density of charged PBHs is given by

$$n(m,Q) = \frac{f\rho_{\rm DM}}{M_{\rm PBH}} \delta(\frac{Q}{m} - \iota)\delta(m - M_{\rm PBH}), \qquad (4.9)$$

where f is the fraction of PBHs in the dark matter and $\rho_{\rm DM}$ is the dark matter energy density in the present time. According to Sec. 3, the merger rate of the charged PBHs that would have formed in the late universe is

$$R_{\rm PBH}(\iota) \approx 1.5 \times 10^{-8} f^2 (1 - \iota^2)^{12/7} \left(1 + \iota^2/4\right)^{2/7} \rm Gpc^{-3} yr^{-1}, \qquad (4.10)$$

and is independent of the mass of the charged PBHs. For the model b, the number density of charged PBHs is

$$n(m,Q) = \frac{1}{2} \frac{f\rho_{\rm DM}}{M_{\rm PBH}} \left(\delta(\frac{Q}{m} - \iota) + \delta(\frac{Q}{m} + \iota) \right) \delta(m - M_{\rm PBH}).$$
(4.11)

Following Sec. 3 again, the merger rate of the charged PBHs that would have formed in the late universe can be simplified as

$$R_{\rm PBH}(\iota) \approx 0.75 \times 10^{-8} f^2 (1-\iota^2)^{12/7} \left(1+\iota^2/4\right)^{2/7} \rm Gpc^{-3} yr^{-1} + \frac{f^2 \rho_{\rm DM}^2}{2M_{\rm PBH}^2} \int P(v) \sigma_1 v dv, \quad (4.12)$$

where $\sigma_1 = \pi b_{max}^2$ denotes the merging cross section for charged PBHs and b_{max} is determined by

$$-\frac{2720\pi \left(\iota^2+1\right)^6 M_{\rm PBH}^8}{3b_{max}^7 v^7} - \frac{32\pi \iota^2 \left(\iota^2+1\right)^4 M_{\rm PBH}^6}{b_{max}^5 v^5} + \frac{M_{\rm PBH} v^2}{4} = 0.$$
(4.13)

Notice that from Eq. (4.13) it follows $b_{max} \propto M_{\rm PBH}$ and $\sigma_1 \propto M_{\rm PBH}^2$. Therefore, we find that $R_{\rm PBH}(\iota)$ in (4.12) is independent of $M_{\rm PBH}$ and scales as f^2 . So, $R_{\rm PBH}(\iota)/R_{\rm PBH}(0)$ is only a function of ι where $R_{\rm PBH}(0) \approx 1.5 \times 10^{-8} f^2 {\rm Gpc}^{-3} {\rm yr}^{-1}$. In Fig. 2, we plot the normalized merger rate $R_{\rm PBH}(\iota)/R_{\rm PBH}(0)$ of charged PBHs as the function of ι for different models. Similarly, we notice that $\eta(\iota)$ decreases as ι increases and $\eta(1) = 0$ in the model a while $\eta(\iota)$ increases as ι increases and reaches $\eta(1) \approx 8.4$ in the model b.

In this section, we have shown the effects of the charge-to-mass ratio on the merger rate of ABH and PBH binaries from the two-body dynamical capture for possible different models. Here, we want to discuss the understandings of the different models. As mentioned in the introduction, the U(1) charge considered in this paper can correspond to the different physical models and have interesting interpretations. When the U(1) charge corresponds to modified theories of gravity with additional scalar or vector fields, we can choose model a as a representative model. On the other hand, when the U(1) charge corresponds to electric, magnetic or dark charge, we can choose model b as a representative model, which suggests that the part of the universe under study is charge neutral but charges are locally separated by some mechanisms which were found in Refs. [17, 90].

5 Conclusion

In this work, we have calculated gravitational and electromagnetic radiations from binaries of point masses with U(1) charges in a hyperbolic orbit in the low-velocity and weak-field regime, and applied the result to derive the merger rate of charged BHs from the two-body dynamical capture via gravitational and electromagnetic radiations. We have then shown the effects of the charge-to-mass ratio on the merger rate of ABH and PBH binaries from the two-body dynamical capture. We have also found that the effects of the charge-to-mass ratio on the merger rate depend on the models.

There are various possible extensions and applications of BHs with U(1) charges. Firstly, the U(1) charges can be interpreted as electric charges. As shown in [91, 92], the mergers of electrically charged BHs may potentially offer an explanation for the still-mysterious fast radio bursts and gamma-ray bursts. In Refs. [93, 94], the coalescence of earth-mass primordial BHs with the charge-to-mass ratio $\sim 10^{-4}$ is proposed to be a candidate for sources of the observed fast radio bursts. Here, we have confirmed that the effect of the charge-tomass ratio of $\sim 10^{-4}$ is negligible and the merger scenarios of charged BHs are similar to those without charges. Secondly, the U(1) charges can be interpreted as magnetic charges. Magnetic charges, if they would have existed in the early universe, will offer a new window to explore fundamental physics. Recently, Maldacena extensively discussed the spectacular electroweak symmetry restoration of magnetically charged BHs [35]. Since no evidence of magnetic charges or monopoles has been found yet from terrestrial experiments [95, 96], the merger events of binary BHs detected by LIGO-Virgo may provide us with a new way to investigate whether these BHs indeed have magnetic charges. Finally, the U(1) charges can be interpreted as dark or hidden charges with an extremely weak coupling constant. It is shown in Refs [20, 21] that some merger events detected by LIGO-Virgo are compatible with charged BHs while some are compatible with uncharged BHs. An interesting suggestion to this controversy is that those merger events would have originated from different scenarios. Therefore, a key question is how those BHs would carry dark or hidden charges. We leave all those interesting issues for future works.

Acknowledgments

We would like to thank Daniel Marin for the useful conversations. This work was supported in part by the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2019R1I1A3A01063183).

A Appendix

A.1 Electromagnetic dipole and quadrupole radiations

In the case of the same charge to mass ratio $(Q_1/m_1 = Q_2/m_2)$, the electromagnetic dipole radiation vanishes and the quadrupole radiation becomes the leading term. This is true for any system of charges [50]. Here, we will calculate the energy and angular momentum loss rates due to the electromagnetic dipole and quadrupole radiations. From the definition of traceless charge quadrupole, we find

$$D^{ab} = \sum_{i=1,2} Q_i d_i^a d_i^b - \frac{1}{3} \sum_{i=1,2} Q_i d_i^c d_i^c \delta_{ab} = \mu \left(\frac{Q_1}{m_1^2} + \frac{Q_1}{m_1^2} \right) Q^{ab}.$$
 (A.1)

We extend the magnetic vector potential \boldsymbol{A} as the quadrupole term,

$$A_{i} = \frac{P_{ij}(n)}{r} (\dot{p}^{j} + \frac{1}{2} \ddot{D}^{jk} n_{k}), \qquad (A.2)$$

where $P_{ij} = \delta_{ij} - n_i n_j$ is the projection operator that realizes the transversal gauge in the *n*-direction. According to [50], the total energy emitted due to the electromagnetic dipole and quadrupole radiations is

$$\frac{dE_{EM}}{dt} = -\frac{r^2}{4\pi} \int \boldsymbol{H}^2 d\Omega \tag{A.3}$$

where $H^i = \epsilon^i_{jk} \dot{A}^j n^k$. Adding Eq. (A.2) and using

$$\int n^i d\Omega = \int n^i n^j n^k d\Omega = 0, \tag{A.4}$$

$$\int n^{i}n^{j}n^{k}n^{l}d\Omega = \frac{4\pi}{15} \left(\delta^{ij}\delta^{kl} + \delta^{ik}\delta^{jl} + \delta^{il}\delta^{jk} \right), \quad \int n^{d}n^{k}d\Omega = \frac{4}{3}\pi\delta^{dk}, \tag{A.5}$$

we find

$$\frac{dE_{EM}}{dt} = \frac{dE_{EM}^{dip}}{dt} + \frac{dE_{EM}^{quad}}{dt},\tag{A.6}$$

where

$$\frac{dE_{EM}^{dip}}{dt} = -\frac{2\ddot{p}^2}{3},$$
 (A.7)

$$\frac{dE_{EM}^{quad}}{dt} = -\frac{\overrightarrow{D}_{ij}\overrightarrow{D}_{ij}}{20}.$$
(A.8)

Similarly, the loss rate of angular momentum due to electromagnetic radiation is given by

$$\frac{dL_{EM}^i}{dt} = -\frac{r^2}{4\pi} \int \left(-\epsilon^{ikl} \left(\partial_0 A_j\right) x^k \partial^l A_j + \epsilon^{ikl} A_k \partial_0 A_l\right) d\Omega.$$
(A.9)

After very careful calculation, we obtain

$$\frac{dL_{EM}^i}{dt} = \frac{dL_{EM}^{i,dip}}{dt} + \frac{dL_{EM}^{i,quad}}{dt},\tag{A.10}$$

where

$$\frac{dL_{EM}^{i,dip}}{dt} = -\frac{2\epsilon^{ikl}}{3}\dot{p}_k\ddot{p}_l \tag{A.11}$$

$$\frac{dL_{EM}^{i,quad}}{dt} = -\frac{1}{20} \epsilon^{ikl} \left(\ddot{D}_{ka} \ddot{D}_{la} \right) \tag{A.12}$$

Noting

$$\frac{dE_{GW}^{quad}}{dt} \equiv -\frac{1}{5} \left(\ddot{Q}_{ij} \ddot{Q}_{ij} \right), \quad \frac{dL_{GW}^{i,quad}}{dt} \equiv -\frac{2}{5} \epsilon^{ikl} \left(\ddot{Q}_{ka} \ddot{Q}_{la} \right)$$
(A.13)

and using the definition of traceless charge quadrupole (A.1), we find the relations between quadrupole radiations:

$$\frac{dE_{EM}^{quad}}{dt} \equiv \frac{\mu^2 \left(Q_1/m_1^2 + Q_2/m_2^2\right)^2}{4} \frac{dE_{GW}^{quad}}{dt},\tag{A.14}$$

$$\frac{dL_{EM}^{i,quad}}{dt} \equiv \frac{\mu^2 \left(Q_1/m_1^2 + Q_2/m_2^2\right)^2}{8} \frac{dL_{GW}^{i,quad}}{dt}.$$
(A.15)

Under the condition of gravitational attraction dominance over the electric repulsion, i.e $|Q_1| \leq m_1$ and $|Q_2| \leq m_2$, we always have $\frac{dE_{EM}^{quad}}{dt} \leq \frac{1}{4} \frac{dE_{GW}^{quad}}{dt}$ and $\frac{dL_{EM}^{i,quad}}{dt} \leq \frac{1}{8} \frac{dL_{GW}^{i,quad}}{dt}$.

References

- A. Einstein, Approximative Integration of the Field Equations of Gravitation, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) 1916 (1916) 688.
- [2] A. Einstein, Über Gravitationswellen, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) 1918 (1918) 154.
- [3] LIGO SCIENTIFIC, VIRGO collaboration, Observation of Gravitational Waves from a Binary Black Hole Merger, Phys. Rev. Lett. 116 (2016) 061102 [1602.03837].
- [4] LIGO SCIENTIFIC, VIRGO collaboration, GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs, Phys. Rev. X9 (2019) 031040 [1811.12907].
- [5] LIGO SCIENTIFIC, VIRGO collaboration, GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run, Phys. Rev. X 11 (2021) 021053 [2010.14527].
- [6] LIGO SCIENTIFIC, VIRGO, KAGRA collaboration, Gwtc-3: Compact binary coalescences observed by ligo and virgo during the second part of the third observing run, 2111.03606.
- [7] LIGO SCIENTIFIC, VIRGO collaboration, Tests of General Relativity with the Binary Black Hole Signals from the LIGO-Virgo Catalog GWTC-1, Phys. Rev. D 100 (2019) 104036
 [1903.04467].
- [8] LIGO SCIENTIFIC, VIRGO collaboration, Tests of general relativity with binary black holes from the second LIGO-Virgo gravitational-wave transient catalog, Phys. Rev. D 103 (2021) 122002 [2010.14529].
- [9] LIGO SCIENTIFIC, VIRGO, KAGRA collaboration, Tests of General Relativity with GWTC-3, 2112.06861.
- [10] S. W. Hawking, Black holes in general relativity, Commun. Math. Phys. 25 (1972) 152.
- [11] D. C. Robinson, Uniqueness of the Kerr black hole, Phys. Rev. Lett. 34 (1975) 905.
- [12] V. Cardoso and L. Gualtieri, Testing the black hole 'no-hair' hypothesis, Class. Quant. Grav. 33 (2016) 174001 [1607.03133].
- [13] M. Zilhao, V. Cardoso, C. Herdeiro, L. Lehner and U. Sperhake, Collisions of charged black holes, Phys. Rev. D 85 (2012) 124062 [1205.1063].
- [14] M. Zilhão, V. Cardoso, C. Herdeiro, L. Lehner and U. Sperhake, Collisions of oppositely charged black holes, Phys. Rev. D 89 (2014) 044008 [1311.6483].
- [15] S. L. Liebling and C. Palenzuela, Electromagnetic Luminosity of the Coalescence of Charged Black Hole Binaries, Phys. Rev. D94 (2016) 064046 [1607.02140].

- [16] B. Toshmatov, Z. Stuchlík, J. Schee and B. Ahmedov, Electromagnetic perturbations of black holes in general relativity coupled to nonlinear electrodynamics, Phys. Rev. D 97 (2018) 084058 [1805.00240].
- [17] Y. Bai and N. Orlofsky, Primordial Extremal Black Holes as Dark Matter, Phys. Rev. D 101 (2020) 055006 [1906.04858].
- [18] A. Allahyari, M. Khodadi, S. Vagnozzi and D. F. Mota, Magnetically charged black holes from non-linear electrodynamics and the Event Horizon Telescope, JCAP 02 (2020) 003 [1912.08231].
- [19] O. Christiansen, J. Beltrán Jiménez and D. F. Mota, Charged Black Hole Mergers: Orbit Circularisation and Chirp Mass Bias, Class. Quant. Grav. 38 (2021) 075017 [2003.11452].
- [20] H.-T. Wang, P.-C. Li, J.-L. Jiang, Y.-M. Hu and Y.-Z. Fan, Post-Newtonian waveform for charged binary black hole inspirals and analysis with GWTC-1 events, 2004.12421.
- [21] G. Bozzola and V. Paschalidis, General Relativistic Simulations of the Quasicircular Inspiral and Merger of Charged Black Holes: GW150914 and Fundamental Physics Implications, Phys. Rev. Lett. 126 (2021) 041103 [2006.15764].
- [22] Y. Kim and A. Kobakhidze, Topologically induced black hole charge and its astrophysical manifestations, 2008.04506.
- [23] V. Cardoso, W.-D. Guo, C. F. B. Macedo and P. Pani, The tune of the Universe: the role of plasma in tests of strong-field gravity, Mon. Not. Roy. Astron. Soc. 503 (2021) 563 [2009.07287].
- [24] B. McInnes, About Magnetic AdS Black Holes, JHEP 03 (2021) 068 [2011.07700].
- [25] Y. Bai and M. Korwar, Hairy Magnetic and Dyonic Black Holes in the Standard Model, JHEP 04 (2021) 119 [2012.15430].
- [26] M. D. Diamond and D. E. Kaplan, Constraints on Relic Magnetic Black Holes, 2103.01850.
- [27] G. Bozzola and V. Paschalidis, Numerical-relativity simulations of the quasicircular inspiral and merger of nonspinning, charged black holes: Methods and comparison with approximate approaches, Phys. Rev. D 104 (2021) 044004 [2104.06978].
- [28] B. McInnes, The weak gravity conjecture requires the existence of exotic AdS black holes, Nucl. Phys. B 971 (2021) 115525 [2104.07373].
- [29] K. Kritos and J. Silk, Mergers of Maximally Charged Primordial Black Holes, 2109.09769.
- [30] S. Hou, S. Tian, S. Cao and Z.-H. Zhu, Dark photon bursts from compact binary systems and constraints, 2110.05084.
- [31] C. A. Benavides-Gallego and W.-B. Han, *Phenomenological model for the electromagnetic response of a black hole binary immersed in magnetic field*, 2111.04323.
- [32] M. D. Diamond, D. E. Kaplan and S. Rajendran, Binary Collisions of Dark Matter Blobs, 2112.09147.
- [33] H. Mouri and Y. Taniguchi, Runaway merging of black holes: analytical constraint on the timescale, Astrophys. J. Lett. 566 (2002) L17 [astro-ph/0201102].
- [34] S. Bird, I. Cholis, J. B. Muñoz, Y. Ali-Haïmoud, M. Kamionkowski, E. D. Kovetz et al., *Did LIGO detect dark matter?*, *Phys. Rev. Lett.* **116** (2016) 201301 [1603.00464].
- [35] J. Maldacena, Comments on magnetic black holes, JHEP 04 (2021) 079 [2004.06084].
- [36] Y. Bai, J. Berger, M. Korwar and N. Orlofsky, Phenomenology of magnetic black holes with electroweak-symmetric coronas, JHEP 10 (2020) 210 [2007.03703].
- [37] D. Ghosh, A. Thalapillil and F. Ullah, Astrophysical hints for magnetic black holes, Phys. Rev. D 103 (2021) 023006 [2009.03363].

- [38] L. Ackerman, M. R. Buckley, S. M. Carroll and M. Kamionkowski, Dark Matter and Dark Radiation, Phys. Rev. D 79 (2009) 023519 [0810.5126].
- [39] J. L. Feng, M. Kaplinghat, H. Tu and H.-B. Yu, *Hidden Charged Dark Matter*, JCAP 07 (2009) 004 [0905.3039].
- [40] R. Foot and S. Vagnozzi, Dissipative hidden sector dark matter, Phys. Rev. D 91 (2015) 023512 [1409.7174].
- [41] R. Foot and S. Vagnozzi, Diurnal modulation signal from dissipative hidden sector dark matter, Phys. Lett. B 748 (2015) 61 [1412.0762].
- [42] J. W. Moffat, Scalar-tensor-vector gravity theory, JCAP 03 (2006) 004 [gr-qc/0506021].
- [43] V. Cardoso, C. F. B. Macedo, P. Pani and V. Ferrari, Black holes and gravitational waves in models of minicharged dark matter, JCAP 1605 (2016) 054 [1604.07845].
- [44] V. Cardoso, C. F. B. Macedo and R. Vicente, Eccentricity evolution of compact binaries and applications to gravitational-wave physics, Phys. Rev. D 103 (2021) 023015 [2010.15151].
- [45] E. Fischbach and C. Talmadge, Six years of the fifth force, Nature 356 (1992) 207.
- [46] G. R. Dvali and M. Zaldarriaga, Changing alpha with time: Implications for fifth force type experiments and quintessence, Phys. Rev. Lett. 88 (2002) 091303 [hep-ph/0108217].
- [47] S. S. Gubser and J. Khoury, Scalar self-interactions loosen constraints from fifth force searches, Phys. Rev. D 70 (2004) 104001 [hep-ph/0405231].
- [48] P. C. Peters and J. Mathews, Gravitational radiation from point masses in a Keplerian orbit, Phys. Rev. 131 (1963) 435.
- [49] P. C. Peters, Gravitational Radiation and the Motion of Two Point Masses, Phys. Rev. 136 (1964) B1224.
- [50] L. D. Landau and E. M. Lifschits, The Classical Theory of Fields, vol. Volume 2 of Course of Theoretical Physics. Pergamon Press, Oxford, 1975.
- [51] L. Liu, Z.-K. Guo, R.-G. Cai and S. P. Kim, Merger rate distribution of primordial black hole binaries with electric charges, Phys. Rev. D 102 (2020) 043508 [2001.02984].
- [52] R. O. Hansen, Postnewtonian gravitational radiation from point masses in a hyperbolic kepler orbit, Phys. Rev. D 5 (1972) 1021.
- [53] L. Liu, O. Christiansen, Z.-K. Guo, R.-G. Cai and S. P. Kim, Gravitational and electromagnetic radiation from binary black holes with electric and magnetic charges: Circular orbits on a cone, Phys. Rev. D 102 (2020) 103520 [2008.02326].
- [54] L. Liu, O. Christiansen, W.-H. Ruan, Z.-K. Guo, R.-G. Cai and S. P. Kim, Gravitational and electromagnetic radiation from binary black holes with electric and magnetic charges: elliptical orbits on a cone, Eur. Phys. J. C 81 (2021) 1048 [2011.13586].
- [55] L. Liu and S. P. Kim, Gravitational and electromagnetic radiations from binary black holes with electric and magnetic charges, in 17th Italian-Korean Symposium on Relativistic Astrophysics, 1, 2022, 2201.01138.
- [56] L. Barack et al., Black holes, gravitational waves and fundamental physics: a roadmap, Class. Quant. Grav. 36 (2019) 143001 [1806.05195].
- [57] K. W. K. Wong, K. Breivik, K. Kremer and T. Callister, Joint constraints on the field-cluster mixing fraction, common envelope efficiency, and globular cluster radii from a population of binary hole mergers via deep learning, Phys. Rev. D 103 (2021) 083021 [2011.03564].
- [58] M. Zevin, S. S. Bavera, C. P. L. Berry, V. Kalogera, T. Fragos, P. Marchant et al., One Channel to Rule Them All? Constraining the Origins of Binary Black Holes Using Multiple Formation Pathways, Astrophys. J. 910 (2021) 152 [2011.10057].

- [59] P. Kroupa, On the variation of the initial mass function, Mon. Not. Roy. Astron. Soc. 322 (2001) 231 [astro-ph/0009005].
- [60] LIGO SCIENTIFIC, VIRGO collaboration, Population Properties of Compact Objects from the Second LIGO-Virgo Gravitational-Wave Transient Catalog, Astrophys. J. Lett. 913 (2021) L7 [2010.14533].
- [61] C. E. Rhoades, Jr. and R. Ruffini, Maximum mass of a neutron star, Phys. Rev. Lett. 32 (1974) 324.
- [62] S. E. Woosley, A. Heger and T. A. Weaver, The evolution and explosion of massive stars, Rev. Mod. Phys. 74 (2002) 1015.
- [63] Y. B. N. Zel'dovich, I. D., The Hypothesis of Cores Retarded during Expansion and the Hot Cosmological Model, Soviet Astron. AJ (Engl. Transl.), 10 (1967) 602.
- [64] S. Hawking, Gravitationally collapsed objects of very low mass, Mon. Not. Roy. Astron. Soc. 152 (1971) 75.
- [65] B. J. Carr and S. W. Hawking, Black holes in the early Universe, Mon. Not. Roy. Astron. Soc. 168 (1974) 399.
- [66] Z.-C. Chen and Q.-G. Huang, Distinguishing Primordial Black Holes from Astrophysical Black Holes by Einstein Telescope and Cosmic Explorer, JCAP 08 (2020) 039 [1904.02396].
- [67] S. Mukherjee and J. Silk, Can we distinguish astrophysical from primordial black holes via the stochastic gravitational wave background?, Mon. Not. Roy. Astron. Soc. 506 (2021) 3977 [2105.11139].
- [68] L. Liu, X.-Y. Yang, Z.-K. Guo and R.-G. Cai, Testing primordial black hole and measuring the Hubble constant with multiband gravitational-wave observations, 2112.05473.
- [69] G. Franciolini, R. Cotesta, N. Loutrel, E. Berti, P. Pani and A. Riotto, *How to assess the primordial origin of single gravitational-wave events with mass, spin, eccentricity, and deformability measurements*, 2112.10660.
- [70] B. Carr, F. Kuhnel and M. Sandstad, Primordial Black Holes as Dark Matter, Phys. Rev. D94 (2016) 083504 [1607.06077].
- [71] M. Sasaki, T. Suyama, T. Tanaka and S. Yokoyama, Primordial Black Hole Scenario for the Gravitational-Wave Event GW150914, Phys. Rev. Lett. 117 (2016) 061101 [1603.08338].
- [72] R. Bean and J. Magueijo, Could supermassive black holes be quintessential primordial black holes?, Phys. Rev. D 66 (2002) 063505 [astro-ph/0204486].
- [73] M. Kawasaki, A. Kusenko and T. T. Yanagida, Primordial seeds of supermassive black holes, Phys. Lett. B 711 (2012) 1 [1202.3848].
- [74] B. Carr and J. Silk, Primordial Black Holes as Generators of Cosmic Structures, Mon. Not. Roy. Astron. Soc. 478 (2018) 3756 [1801.00672].
- [75] S. Clesse and J. García-Bellido, The clustering of massive Primordial Black Holes as Dark Matter: measuring their mass distribution with Advanced LIGO, Phys. Dark Univ. 15 (2017) 142 [1603.05234].
- [76] H. Nishikawa, E. D. Kovetz, M. Kamionkowski and J. Silk, Primordial-black-hole mergers in dark-matter spikes, Phys. Rev. D99 (2019) 043533 [1708.08449].
- [77] T. Nakamura, M. Sasaki, T. Tanaka and K. S. Thorne, Gravitational waves from coalescing black hole MACHO binaries, Astrophys. J. 487 (1997) L139 [astro-ph/9708060].
- [78] K. Ioka, T. Chiba, T. Tanaka and T. Nakamura, Black hole binary formation in the expanding universe: Three body problem approximation, Phys. Rev. D58 (1998) 063003 [astro-ph/9807018].

- [79] M. Raidal, V. Vaskonen and H. Veermäe, Gravitational Waves from Primordial Black Hole Mergers, JCAP 1709 (2017) 037 [1707.01480].
- [80] B. Kocsis, T. Suyama, T. Tanaka and S. Yokoyama, Hidden universality in the merger rate distribution in the primordial black hole scenario, Astrophys. J. 854 (2018) 41 [1709.09007].
- [81] Y. Ali-Haïmoud, E. D. Kovetz and M. Kamionkowski, Merger rate of primordial black-hole binaries, Phys. Rev. D96 (2017) 123523 [1709.06576].
- [82] Z.-C. Chen and Q.-G. Huang, Merger rate distribution of primordial-black-hole binaries, Astrophys. J. 864 (2018) 61 [1801.10327].
- [83] G. Ballesteros, P. D. Serpico and M. Taoso, On the merger rate of primordial black holes: effects of nearest neighbours distribution and clustering, JCAP 1810 (2018) 043 [1807.02084].
- [84] M. Raidal, C. Spethmann, V. Vaskonen and H. Veermäe, Formation and Evolution of Primordial Black Hole Binaries in the Early Universe, JCAP 02 (2019) 018 [1812.01930].
- [85] L. Liu, Z.-K. Guo and R.-G. Cai, Effects of the surrounding primordial black holes on the merger rate of primordial black hole binaries, Phys. Rev. D 99 (2019) 063523 [1812.05376].
- [86] L. Liu, Z.-K. Guo and R.-G. Cai, Effects of the merger history on the merger rate density of primordial black hole binaries, Eur. Phys. J. C 79 (2019) 717 [1901.07672].
- [87] S. Young and C. T. Byrnes, *Initial clustering and the primordial black hole merger rate*, JCAP 03 (2020) 004 [1910.06077].
- [88] V. Vaskonen and H. Veermäe, Lower bound on the primordial black hole merger rate, Phys. Rev. D 101 (2020) 043015 [1908.09752].
- [89] J. Garriga and N. Triantafyllou, Enhanced cosmological perturbations and the merger rate of PBH binaries, JCAP 1909 (2019) 043 [1907.01455].
- [90] R. M. Wald, Black hole in a uniform magnetic field, Phys. Rev. D 10 (1974) 1680.
- [91] B. Zhang, Mergers of Charged Black Holes: Gravitational Wave Events, Short Gamma-Ray Bursts, and Fast Radio Bursts, Astrophys. J. 827 (2016) L31 [1602.04542].
- [92] T. Liu, G. E. Romero, M.-L. Liu and A. Li, Fast Radio Bursts and Their Gamma-ray or Radio Afterglows as Kerr-newman Black Hole Binaries, Astrophys. J. 826 (2016) 82 [1602.06907].
- [93] C.-M. Deng, Y. Cai, X.-F. Wu and E.-W. Liang, Fast Radio Bursts From Primordial Black Hole Binaries Coalescence, Phys. Rev. D 98 (2018) 123016 [1812.00113].
- [94] C.-M. Deng, Earth-mass primordial black hole mergers as sources for nonrepeating fast radio bursts, Phys. Rev. D 103 (2021) 123030 [2110.08981].
- [95] MOEDAL collaboration, Recent Results and Future Plans of the MoEDAL Experiment, in Meeting of the Division of Particles and Fields of the American Physical Society, 10, 2019, 1910.05772.
- [96] N. E. Mavromatos and V. A. Mitsou, Magnetic monopoles revisited: Models and searches at colliders and in the Cosmos, Int. J. Mod. Phys. A 35 (2020) 2030012 [2005.05100].