m^* of two-dimensional electron gas: a neural canonical transformation study

Hao Xie,^{1,2} Linfeng Zhang,^{3,4,*} and Lei Wang^{1,5,†}

¹Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

²University of Chinese Academy of Sciences, Beijing 100049, China

⁴AI for Science Institute, Beijing, 100080, China

⁵Songshan Lake Materials Laboratory, Dongguan, Guangdong 523808, China

(Dated: January 11, 2022)

The quasiparticle effective mass m^* of interacting electrons is a fundamental quantity in the Fermi liquid theory. However, the precise value of the effective mass of uniform electron gas is still elusive after decades of research. The newly developed neural canonical transformation approach [Xie *et al.*, 2105.08644] offers a principled way to extract the effective mass of electron gas by directly calculating the thermal entropy at low temperature. The approach models a variational many-electron density matrix using two generative neural networks: an autoregressive model for momentum occupation and a normalizing flow for electron coordinates. Our calculation reveals a suppression of effective mass in the two-dimensional spin-polarized electron gas, which is more pronounced than previous reports in the low-density strong-coupling region. This prediction calls for verification in two-dimensional electron gas experiments.

Introduction – Landau's Fermi liquid theory [1] is one of the cornerstones of condensed matter physics [2]. It explains the mystery why the non-interacting picture can largely apply to real metals despite the strong Coulomb repulsion between electrons. The essence is that a Fermi liquid consists of quasiparticles that are adiabatically connected to bare electrons. Such a renormalization procedure can be encapsulated in only a handful of parameters, from which one can predict a broad range of physical properties of the system. One such parameter is the quasiparticle effective mass m^* , which is the central focus of this work.

Surprisingly, the precise value of the quasiparticle effective mass of uniform electron gas is still controversial after more than fifty years of research [3-17]. The uniform electron gas consists of electrons distributed homogeneously in a background of positive charges. Despite of being simple, the model captures the essence of electron correlation effects and serves as a foundational model of interacting electrons [18, 19].

Depending on the spatial dimension and spin polarization of the uniform electron gas, previous results may differ quantitatively or even qualitatively on whether the quasiparticle effective mass is enhanced ($m^*/m > 1$) or suppressed ($m^*/m < 1$) compared to the bare electron mass m. Resolving these discrepancies within the same approach can be challenging. For example, there lacks a systematic way of improving various approximate analytic calculations to reach a consensus [3–9]. It is hoped that numerical calculations offer more reliable predictions to the effective mass. However, two recent quantum Monte Carlo (QMC) studies [16, 17] report drastically different effective masses for the three-dimensional electron gas. The reason for such discrepancy is unclear and may be related to different (but equivalent) ways of defining the effective mass as well as different approximations employed in the methods. Moreover, the situation is not clear either in the twodimensional case even if one employs the same kind of QMC method [12–15]. There, the predicted effective mass differ qualitatively depending on how to process the QMC data [20]. These discrepancies are related to ambiguities in relating excited state energies to the effective mass [19], which again entangle with approximations and finite size errors in the calculations.

Resolving the discrepancy on the effective mass of uniform electron gas is not merely a theoretical question with pure academic interests, but also of direct experimental relevance. One can measure the effective mass in a semiconductor quantum well via quantum oscillations [21–25] or thermodynamics [26], which is a high-quality realization of the twodimensional electron gas (2DEG) with tunable densities.

At sufficiently low temperature, the entropy per particle of 2DEG $s/k_B = \frac{\pi^2}{3} \frac{m^*}{m} \frac{T}{T_F}$ exhibits linear dependence with the temperature *T*, where k_B is the Boltzmann constant and T_F is the Fermi temperature. Therefore, one can directly estimate the effective mass from the entropy ratio of interacting (*s*) and non-interacting (*s*₀) electron gases [27]

$$\frac{m^*}{m} = \frac{s}{s_0}.$$
 (1)

By direct access of the thermodynamic observables, one can avoid subtleties of processing excited state energies under additional assumptions, such as the validity of Fermi liquid theory for excitations away from the Fermi surface [19]. However, previous finite-temperature calculations of the uniform electron gas do not resolve the issue related to the effective mass [27] because they focus on the melting of the Wigner crystal [28] or the equation of state in the warm dense matter region [29–31], both are outside the scope of Fermi-liquidlike behavior. This is partially due to the fact that the adopted QMC methods typically suffer less from the sign problem at low density and high temperature, where the fermionic nature of the system is less pronounced.

In this paper, we employ the recently developed neural canonical transformation approach [32] to study 2DEG at low

³DP Technology, Beijing, 100080, China

^{*} linfeng.zhang.zlf@gmail.com

[†] wanglei@iphy.ac.cn



FIG. 1. (a) A set of occupied momenta K residing on the reciprocal space grid within an energy cutoff indicated by the circle. In the figure there are N = 13 electrons distributed in M = 49 allowed momenta. We use the autoregressive network Eq. (6) to model the Boltzmann distribution of such configurations. (b) A set of electron coordinates R residing in a periodic box. A normalizing flow network transforms R to a new set of quasiparticle coordinates ζ in the same box. This induces a unitary transformation which, when acting upon the plane-wave Slater determinants, produces a set of manybody basis states Eq. (8).

temperature and estimate the effective mass via the entropy ratio Eq. (1). Neural canonical transformation leverages recent advances in deep generative models [33] for variational free energy calculation of interacting fermions at finite temperature. This approach is particularly suitable for the present task for two reasons. Firstly, the employed variational density matrix ansatz fits nicely to the philosophy of Fermi liquid theory. Secondly, the thermal entropy can be directly accessed unlike other conventional QMC methods.

Consider *N* electrons in a two-dimensional periodic box of length *L*. We focus on the spin-polarized case, which can be conveniently realized in experiments by applying an inplane magnetic field. We set the energy unit to be Rydberg $\hbar^2/(2ma_0^2)$, where $a_0 = \hbar^2/(me^2)$ is the Bohr radius. The dimensionless Wigner-Seitz parameter $r_s = L/(\sqrt{\pi N}a_0)$ measures the average distance between electrons in the unit of Bohr radius. The Hamiltonian reads [34]

$$H = -\frac{1}{r_s^2} \sum_{i=1}^{N} \nabla_i^2 + \frac{2}{r_s} \sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} + \text{const}, \qquad (2)$$

where $\mathbf{r}_i = (x_i, y_i)$ is the coordinate of the *i*-th electron. The constant term in Eq. (2) refers to the energy due to the neutralizing background.

Method – To investigate the finite-temperature properties of Eq. (2), we minimize the variational free energy

$$F = \frac{1}{\beta} \operatorname{Tr}(\rho \ln \rho) + \operatorname{Tr}(H\rho)$$
(3)

with respect to a many-electron density matrix ρ , where $\beta = 1/(k_BT)$ is the inverse temperature. In practice, *T* is measured relative to the Fermi temperature $k_BT_F = 4\text{Ry}/r_s^2$. The variational free energy Eq. (3) is lower bounded by the true free energy of the system, i.e., $F \ge -\frac{1}{\beta} \ln Z$, where $Z = \text{Tr}e^{-\beta H}$

is the partition function. The equality holds only when the variational density matrix coincides with the exact one, i.e., $\rho = e^{-\beta H}/Z$.

The variational density matrix is expressed as a weighted sum over a family of many-body orthonormal basis states

$$\rho = \sum_{\mathbf{K}} p(\mathbf{K}) |\Psi_{\mathbf{K}}\rangle \langle \Psi_{\mathbf{K}}| \,. \tag{4}$$

Here $K \equiv \{k_1, k_2, ..., k_N\}$ represents a *set* of occupied momenta, each taking one of the discrete values $k = \frac{2\pi n}{L}$ ($n \in \mathbb{Z}^2$) without duplication as required by the Pauli exclusion principle. Such setting is closely in line with an essential point of Fermi liquid theory [1]: one can label the low-energy excited states using the same quantum number as the ideal Fermi gas. In practice, one has to truncate the momenta within an energy cutoff, which is set to be sufficiently large to avoid bias in the considered temperature region. Therefore, for *M* possible momenta within the energy cutoff, the summation Eq. (4) involves $\binom{M}{N}$ terms. See Fig. 1(a) for a schematic illustration. Substituting the density matrix ansatz Eq. (4) into Eq. (3),

Substituting the density matrix ansatz Eq. (4) into Eq. (3), one finds an unbiased estimator for the variational free energy

$$F = \mathbb{E}_{K \sim p(K)} \left[\frac{1}{\beta} \ln p(K) + \mathbb{E}_{R \sim |\Psi_K(R)|^2} \left[E_K^{\text{loc}}(R) \right] \right].$$
(5)

Here $\mathbf{R} \equiv \{\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N\}$ is the set of electron coordinates and $\Psi_K(\mathbf{R}) = \langle \mathbf{R} | \Psi_K \rangle$ the corresponding basis wavefunction. The local energy is defined as $E_K^{\text{loc}}(\mathbf{R}) = -\frac{1}{r_s^2} \sum_i \left[\nabla_i^2 \ln \Psi_K(\mathbf{R}) + (\nabla_i \ln \Psi_K(\mathbf{R}))^2 \right] + \frac{2}{r_s} \sum_{i < j} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} + \text{const.}$ We use Ewald summation to evaluate the Coulomb interaction term, while the gradient and Laplacian operator appearing in the kinetic term can be computed using automatic differentiation. We model the Boltzmann distribution $p(\mathbf{K})$ and wavefunction $\Psi_K(\mathbf{R})$ using two generative networks.

We use a variational autoregressive network [35] to model the normalized Boltzmann distribution $p(\mathbf{K})$ over a set of discrete momenta:

$$p(\boldsymbol{K}) = \prod_{i=1}^{N} p(\boldsymbol{k}_i | \boldsymbol{k}_{< i}), \qquad (6)$$

where each factor in the product is a parametrized conditional probability. To facilitate the sampling of these conditional probabilities, we assign a unique index $idx(k) \in \{1, 2, ..., M\}$ to each of the M available momenta, e.g., according to their single-particle energies [36]. We then model $p(\mathbf{K})$ using a neural network that maps the set $\mathbf{K} = \{\mathbf{k}_1, \mathbf{k}_2, \dots, \mathbf{k}_N\}$ to N vectors $\hat{k}_1, \hat{k}_2, \dots, \hat{k}_N$, where $\hat{k}_i \in \mathbb{R}^M$ denotes the conditional log-probability of $idx(k_i)$ given $k_{<i}$. To ensure the autoregressive property, i.e., \hat{k}_i depends only on k_j with j < i, we implement the network as a transformer with causal self-attention layers [37]. In addition, to accommodate the Pauli principle, we require Eq. (6) assign nonzero probabilities only to those momentum configurations satisfying $idx(k_1) < idx(k_2) <$ \ldots < idx(k_N). This can be achieved by carefully masking out disallowed configurations in the output logits k_i [38]. We note that Refs. [39, 40] devised an alternative autoregressive



FIG. 2. Two limiting cases of the present approach. (a) Noninteracting limit. The entropy per particle of N = 37 free electrons at $T/T_F = 0.15$. The horizontal line indicates the exact value calculated by recursion in the canonical ensemble [38, 41]. (b) Zerotemperature limit. The ground-state energy per particle of N = 37electrons at $r_s = 5$. The horizontal line is the variational Monte Carlo result of Ref. [42].

model in the bit string representation with fixed number of nonzero elements.

Using the autoregressive model Eq. (6) rather than enumerating all possible excitations [32] in the summation Eq. (4) allows us to incorporate a combinatorially large number of many-body states and access broader temperature range. One can estimate the thermal entropy unbiasedly via the estimator

$$s = -\frac{1}{N} \mathop{\mathbb{E}}_{\mathbf{K} \sim p(\mathbf{K})} [\ln p(\mathbf{K})].$$
(7)

Note such a simple and tractable expression for the entropy is a direct consequence of orthonormality of the many-body basis $|\Psi_{\mathbf{K}}\rangle$.

Next, to parametrize a family of orthonormal many-body states $|\Psi_K\rangle$, we perform a unitary transformation on the basis of plane-wave Slater determinants. In practice, we construct the unitary transformation as a learnable bijection from the electron coordinates \mathbf{R} to a new set of *quasiparticle coordinates* ζ , as shown in Fig. 1(b). The wave function reads [32]

$$\Psi_{\boldsymbol{K}}(\boldsymbol{R}) = \frac{\det\left(e^{ik_i\cdot\boldsymbol{\zeta}_j}/L\right)}{\sqrt{N!}} \cdot \left|\det\left(\frac{\partial\boldsymbol{\zeta}}{\partial\boldsymbol{R}}\right)\right|^{\frac{1}{2}}.$$
(8)

The originally non-interacting plane waves possessed by individual electrons would interfere with each other due to correlation effects introduced by the coordinate transformation. Such a picture closely mimics the renormalization process as depicted in Fermi liquid theory. Technically, Eq. (8) differs from the standard Slater-Jastrow trial wavefunction by the presence of an additional Jacobian determinant. This factor turns out to play a crucial role of preserving orthonormality of the basis $\int d\mathbf{R} \Psi_{K}^{*}(\mathbf{R})\Psi_{K'}(\mathbf{R}) = \delta_{KK'}$. Note also that the state Eq. (8) involves coordinate transformation in a manybody context [43], where one needs to deal with the extra issue of permutation equivariance compared to the single-particle setting [44–46]. We use a normalizing flow [47] to implement the bijective map between the electron and quasiparticle coordinates. This can be regarded as a generalization of the backflow transformation with invertible neural networks [48]. We compose the FermiNet [49] blocks into a residual network to carry out permutation and translation equivariant transformation of electron coordinates. We also modify the electron distance features to accommodate the periodic boundary condition [38].

It is instructive to examine the present approach in two limiting cases. Firstly, in the non-interacting limit the problem reduces to a classical statistical mechanics problem: one needs to distribute N particles in M possible momenta according to the probability distribution $p(\mathbf{K})$ to minimize the free energy $F = \mathbb{E}_{K \sim p(K)} \left[\frac{1}{\beta} \ln p(K) + \sum_{i=1}^{N} \frac{\hbar^2 k_i^2}{2m} \right]$, where the second term is the non-interacting energy. In this case, one can trivially set the normalizing flow to an identity map and optimize only the autoregressive network. Figure 2(a) shows a typical training process, where the entropy Eq. (7) steadily approaches the exact value [41]. Note the calculation of exact non-interacting entropy in the canonical ensemble is not a completely trivial task [38]. Secondly, in the zerotemperature limit, $p(\mathbf{K})$ is nonzero only for one particular momentum configuration K_0 , corresponding to the closedshell non-interacting ground state. The present approach then reduces to the usual ground-state variational Monte Carlo method. As an example, Fig. 2(b) shows the optimized ground-state energy density $e = \frac{1}{N} \mathbb{E}_{\mathbf{R} \sim |\Psi_{\mathbf{K}_0}(\mathbf{R})|^2} \left[E_{\mathbf{K}_0}^{\text{loc}}(\mathbf{R}) \right]$ for a particular set of system parameters, which is lower than previous report using the Slater-Jastrow ansatz [42]. In general cases, one has to jointly optimize the autoregressive model and the normalizing flow. We show some additional benchmark results for the three-dimensional spin-polarized electron gas in the supplemental material [38].

To understand how the variational free energy calculation reveals the quasiparticle effective mass, note the effective mass affects low-temperature thermodynamics of the system via the density of states of low-lying excitations. In practice, we pretrain the state occupation $p(\mathbf{K})$ using non-interacting energies as in Fig. 2(a). Thus, $p(\mathbf{K})$ will initially give the same entropy of the ideal Fermi gas. We initialize the normalizing flow network to be close to an identity map. Training of the normalizing flow will modify the many-body basis Eq. (8) and thus change the quasiparticle energy spacing and density of states. On the other hand, the autoregressive model will also adjust the Boltzmann distribution accordingly, causing the entropy to depart from the initial non-interacting value. Putting it all together, the entropy ratio Eq. (1) thus provides a principled way to extract the effective mass from the quasiparticle energy spectrum.

Results – To access the quasiparticle effective mass of electron gas via the entropy ratio Eq. (1), one should consider temperatures *T* well below the Fermi temperature T_F . Figure 3(a) shows the entropy per particle of ideal Fermi gas in the thermodynamic limit $N = \infty$ [50], which exhibits linear behavior $s/k_B = \frac{\pi^2}{3} \frac{T}{T_F}$ in the low-temperature limit and crosses over to $s/k_B = 2 + \ln \frac{T}{T_F}$ in the high-temperature limit. On the other hand, the temperature should also not be too low in practi-



FIG. 3. (a) Entropy per particle of two-dimensional ideal Fermi gas for several electron numbers N, including the thermodynamic limit. The dashed line shows the low-temperature linear behavior. (b) Entropy per particle as a function of training epochs for N = 29 electrons at $T/T_F = 0.15$ for various densities r_s . The reduction of entropy from the initial non-interacting value indicates suppression of the quasiparticle effective mass.

cal calculations, otherwise the finite size effect would cause the entropy to deviate from the ideal linear behavior due to a small energy scale $\frac{\hbar^2}{mL^2} \sim \frac{1}{N}$, as shown in Fig. 3(a) for N = 29 and 57 non-interacting electrons. We choose to set $T/T_F = 0.15$ to balance these two considerations. In principle, the entropy ratio Eq. (1) calculated for increasingly larger system sizes would approach the effective mass in the thermodynamic limit. Even for finite N, one can reasonably expect that by taking the entropy ratio of two calculations, the finite size errors involved in the interacting and non-interacting systems would largely cancel out. Fig. 3(b) shows the entropy as a function of training epochs for N = 29 and various densities r_s . One clearly sees the entropy is reduced from the initial non-interacting value, indicating a suppression of the effective mass upon increasing r_s . The effective mass is estimated via Eq. (1) once the entropy has reached a plateaued value.

Early analytical calculations [7, 51] find enhanced or non-monotonic r_s -dependence of the effective mass in spinpolarized 2DEG. On the other hand, several QMC calculations [13-15] consistently find monotonically suppressed effective mass as r_s increases, but the quantitative predictions still differ, especially for large r_s as shown in Figure 4 [52]. The discrepancy is due to different ways of extracting effective mass from the excited state energies. Refs. [13, 14] obtain the effective mass by differentiating the fitted energy band, while Ref. [15] is based on its relation to other Landau Fermi liquid parameters by Galilean invariance. Both approaches possess a number of uncertainties, such as the fitting range of momentum space and integration error in estimating the Fermi liquid parameters. Though, the authors of Ref. [15] appeared to be more confident with the larger effective mass values reported in Refs. [13, 14].

Figure 4 also shows our estimates of the effective mass for N = 29, 49, 57 electrons, which exhibit similar monotonically decreasing behavior as previous QMC data. However, in spite



FIG. 4. Quasiparticle effective mass calculated using the entropy ratio Eq. (1) for N = 29, 49, 57 electrons at $T/T_F = 0.15$. Also shown for comparison are the QMC data [14, 15] extrapolated to the thermodynamic limit, as well as the analytic result [51] which is valid in the weak-coupling region $r_s \rightarrow 0$.

of visible finite size effect, our results suggest even smaller effective mass than Ref. [15]. Such a prediction also appears to be consistent with the analytical result reported in Ref. [51], shown as the dashed line in Fig. 4, which is reliable in the weak-coupling region $r_s \rightarrow 0$. Since our data processing based on the entropy ratio Eq. (1) is less subject to artificial and tricky assumptions, the present approach offers a cleaner prediction on the effective mass of spin-polarized 2DEG.

On the experimental side, both enhanced [53] and later then suppressed [54, 55] effective mass of the spin-polarized 2DEG are reported in different systems. The discrepancy was attributed to the valley degeneracy [56, 57] involved in the sample used in Ref. [53]. The experimental data [54, 55] spread widely between our predictions and those of Refs. [13– 15] [58]. Confirmation of the present results calls for a new generation of experimental efforts, where besides data uncertainty issue one also has to account for various complications in reality for a fair comparison, such as thickness of the electron layer, disorder and temperature effects [7, 59–61].

Discussions – The largest uncertainty in the estimated effective mass is due to the variational nature of the approach. The variational approximation may be improved by adopting alternative variational ansatz [62–64] and optimization schemes [65, 66]. We have documented the trained models and free energy data in the code repository [67] to facilitate further developments. To address the finite size error requires two lines of research. On the technical side, to push the calculation to larger system sizes one can employ techniques for scaling up machine learning such as gradient checkpointing [68] and distributed training [69]. Specifically, techniques for efficiently training flow models and invertible neural networks [70–75] can be useful. On the theoretical side, the present work calls for a finite size scaling theory for the thermal entropy of uniform electron gas.

With suitable extensions of the model architecture, the

technique developed in this paper can apply equally well to the spin-unpolarized case. This may shed new light on the conflicting results reported in the literature on the threedimensional [16, 17] and two-dimensional [10–15] electron gases. While we have been focusing on the quasiparticle effective mass, the outcomes of the present approach are also directly relevant to the exchange-correlation free energy, which are useful for the thermal density functional theory [27, 31] and thermodynamic measurements in the 2DEG [26]. Along this line, it is also possible to extend the present study to the grand canonical ensemble and compute the compressibility and susceptibility of electron gas measurable in experiments [23, 24, 76, 77]. Finally, having direct access to the energy and wavefunction of low-lying excitations may also allow for calculation of spectral functions at real frequencies.

Neural canonical transformation [32] not only serves as a variational density matrix approach powered by deep learning

techniques, but also nicely incorporates basic notions of Fermi liquid theory. For example, the probabilistic model p(K) in Eq. (4) actually encapsulates the Landau's energy functional for quasiparticles. Moreover, the unitary transformation implemented as a normalizing flow between the electron and quasiparticle coordinates vividly illustrates the notion of adiabatic continuity when switching on interactions [2]. Because of these technical and physical considerations, we are optimistic with the outcome of applying neural canonical transformation to a broader class of interacting fermion problems.

Acknowledgments – We thank Yuan Wan, Pan Zhang, Kun Chen, and Xinguo Ren for useful discussions. This project is supported by the Strategic Priority Research Program of the Chinese Academy of Sciences under Grant No. XDB30000000 and National Natural Science Foundation of China under Grant No. 11774398 and T2121001.

- [1] G. Baym and C. Pethick, *Landau Fermi-liquid theory: concepts and applications* (John Wiley & Sons, 2008).
- [2] P. W. Anderson, Basic notions of condensed matter physics (CRC Press, 2018).
- [3] T. M. Rice, The effects of electron-electron interaction on the properties of metals, Annals of Physics **31**, 100 (1965).
- [4] L. Hedin, Method for calculating the radioactive impurity concentration in the water and the bottom layer of stagnant reservoirs, Physical Review 3, 796 (1965).
- [5] J. Lam, Thermodynamic Properties of the Electron Gas at Metallic Densities, Phys. Rev. B 52, 1254 (1972).
- [6] A. Krakovsky and J. K. Percus, Quasiparticle effective mass for the two- and three-dimensional electron gas, Physical Review B 53, 7352 (1996), arXiv:9601124 [cond-mat].
- [7] Y. Zhang and S. Das Sarma, Spin polarization dependence of carrier effective mass in semiconductor structures: Spintronic effective mass, Physical Review Letters 95, 256603 (2005).
- [8] R. Asgari, B. Davoudi, M. Polini, G. F. Giuliani, M. P. Tosi, and G. Vignale, Quasiparticle self-energy and many-body effective mass enhancement in a two-dimensional electron liquid, Physical Review B 71, 045323 (2005), arXiv:0406676 [cond-mat].
- [9] G. E. Simion and G. F. Giuliani, Many-body local fields theory of quasiparticle properties in a three-dimensional electron liquid, Physical Review B 77, 035131 (2008).
- [10] Y. Kwon, D. M. Ceperley, and R. M. Martin, Quantum Monte Carlo calculation of the Fermi-liquid parameters in the twodimensional electron gas, Physical Review B 50, 1684 (1994).
- [11] Y. Kwon and D. Ceperley, Transient-estimate Monte Carlo in the two-dimensional electron gas, Physical Review B 53, 7376 (1996).
- [12] M. Holzmann, B. Bernu, V. Olevano, R. M. Martin, and D. M. Ceperley, Renormalization factor and effective mass of the two-dimensional electron gas, Physical Review B 79, 041308 (2009), arXiv:0810.2450.
- [13] N. D. Drummond and R. J. Needs, Quantum Monte Carlo calculation of the energy band and quasiparticle effective mass of the two-dimensional Fermi fluid, Physical Review B 80, 245104 (2009).
- [14] N. D. Drummond and R. J. Needs, Diffusion quantum Monte Carlo calculation of the quasiparticle effective mass of the twodimensional homogeneous electron gas, Physical Review B 87,

045131 (2013), arXiv:1208.6317.

- [15] N. D. Drummond and R. J. Needs, Quantum Monte Carlo calculation of the Fermi liquid parameters of the two-dimensional homogeneous electron gas, Physical Review B 88, 035133 (2013), arXiv:1307.4009.
- [16] K. Haule and K. Chen, Single-particle excitations in the uniform electron gas by diagrammatic Monte Carlo, (2020), arXiv:2012.03146.
- [17] S. Azadi, N. D. Drummond, and W. M. C. Foulkes, Quasiparticle effective mass of the three-dimensional fermi liquid by quantum monte carlo, Phys. Rev. Lett. **127**, 086401 (2021).
- [18] G. Giuliani and G. Vignale, *Quantum theory of the electron liquid* (Cambridge university press, 2005).
- [19] R. M. Martin, L. Reining, and D. M. Ceperley, *Interacting electrons* (Cambridge University Press, 2016).
- [20] In fact, the predictions from each group also evolve over years; see Refs. [10–12] and [13–15] respectively.
- [21] F. F. Fang and P. J. Stiles, Effects of a tilted magnetic field on a two-dimensional electron gas, Physical Review 174, 823 (1968).
- [22] B. L. Draper and D. C. Tsui, Mass enhancement of twodimensional electrons in thin-oxide si-mosfet's, Physical Review B 59, 10208 (1999).
- [23] V. M. Pudalov, M. E. Gershenson, H. Kojima, N. Butch, E. M. Dizhur, G. Brunthaler, A. Prinz, and G. Bauer, Low-density spin susceptibility and effective mass of mobile electrons in Si inversion layers, Physical Review Letters 88, 1964041 (2002), arXiv:0105081 [cond-mat].
- [24] K. Vakili, Y. P. Shkolnikov, E. Tutuc, E. P. De Poortere, and M. Shayegan, Spin susceptibility of two-dimensional electrons in narrow AlAs quantum wells, Physical Review Letters 92, 226401 (2004).
- [25] Y. W. Tan, J. Zhu, H. L. Stormer, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, Measurements of the density-dependent many-body electron mass in two dimensional GaAs/AlGaAs heterostructures, Physical Review Letters 94, 016405 (2005).
- [26] A. Y. Kuntsevich, Y. V. Tupikov, V. M. Pudalov, and I. S. Burmistrov, Strongly correlated two-dimensional plasma explored from entropy measurements, Nature Communications 6, 2 (2015).
- [27] F. G. Eich, M. Holzmann, and G. Vignale, Effective mass of

quasiparticles from thermodynamics, Physical Review B **96**, 035132 (2017), arXiv:1704.04076.

- [28] M. Imada and M. Takahashi, Quantum Monte Carlo Simulation of a Two-Dimensional Electron System -Melting of Wigner Crystal, Journal of the Physical Society of Japan 53, 3770 (1984).
- [29] E. W. Brown, B. K. Clark, J. L. Dubois, and D. M. Ceperley, Path-integral Monte Carlo simulation of the warm dense homogeneous electron gas, Physical Review Letters 110, 146405 (2013).
- [30] T. Schoof, S. Groth, J. Vorberger, and M. Bonitz, Ab Initio Thermodynamic Results for the Degenerate Electron Gas at Finite Temperature, Physical Review Letters 115, 130402 (2015), arXiv:1502.04616.
- [31] F. D. Malone, N. S. Blunt, E. W. Brown, D. K. Lee, J. S. Spencer, W. M. Foulkes, and J. J. Shepherd, Accurate Exchange-Correlation Energies for the Warm Dense Electron Gas, Physical Review Letters 117, 115701 (2016).
- [32] H. Xie, L. Zhang, and L. Wang, Ab-initio study of interacting fermions at finite temperature with neural canonical transformation, (2021), arXiv:2105.08644.
- [33] L. Wang, Generative models for physicists (2018).
- [34] D. M. Ceperley, Ground state of the fermion one-component plasma: A Monte Carlo study in two and three dimensions, Phys. Rev. B 18, 3126 (1978).
- [35] D. Wu, L. Wang, and P. Zhang, Solving Statistical Mechanics using Variational Autoregressive Networks, Physical Review Letters 122, 080602 (2019), arXiv:1809.10606.
- [36] In practice, we choose to arrange the momenta in the order of *decreasing* energy. This stems from the observation that a random initialization of the present autoregressive model tends to assign higher probability to configurations with large indices, which then have large overlap with the target Boltzmann distribution.
- [37] A. Vaswani, N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A. N. Gomez, L. Kaiser, and I. Polosukhin, Attention Is All You Need, (2017), arXiv:1706.03762.
- [38] See the Supplemental Material for (a) additional benchmark for three-dimensional spin-polarized uniform electron gas; (b) the analytic method to compute the thermal entropy of noninteracting Fermi gas in the canonical ensemble; details on (c) model architectures and (d) the training procedure.
- [39] M. Hibat-Allah, M. Ganahl, L. E. Hayward, R. G. Melko, and J. Carrasquilla, Recurrent neural network wave functions, Physical Review Research 2, 023358 (2020), arXiv:2002.02973.
- [40] T. D. Barrett, A. Malyshev, and A. I. Lvovsky, Autoregressive neural-network wavefunctions for ab initio quantum chemistry, (2021), arXiv:2109.12606.
- [41] P. Borrmann and G. Franke, Recursion formulas for quantum statistical partition functions, The Journal of Chemical Physics 98, 2484 (1993).
- [42] B. Tanatar and D. M. Ceperley, Ground state of the twodimensional electron gas, Physical review. B 39, 5005 (1989).
- [43] M. Eger and E. Gross, Point transformations and the many body problem, Annals of Physics 24, 63 (1963).
- [44] F. Gygi, Adaptive riemannian metric for plane-wave electronicstructure calculations, Eumphgs. Lett. 19, 617 (1992).
- [45] F. Gygi, Adaptive Coordinates, Phys. Rev. B 48, 11692 (1993).
- [46] K. Cranmer, S. Golkar, and D. Pappadopulo, Inferring the quantum density matrix with machine learning, (2019), arXiv:1904.05903.
- [47] G. Papamakarios, Neural Density Estimation and Likelihoodfree Inference, (2019), arXiv:1910.13233.
- [48] R. P. Feynman and M. Cohen, Energy spectrum of the excita-

tions in liquid helium, Phys. Rev. 102, 1189 (1956).

- [49] D. Pfau, J. S. Spencer, A. G. D. G. Matthews, and W. M. C. Foulkes, Ab initio solution of the many-electron Schrödinger equation with deep neural networks, Physical Review Research 2, 033429 (2020), arXiv:1909.02487.
- [50] For the two-dimensional ideal Fermi gas in the thermodynamic limit, one has $s/k_B = 2f_2(z)/f_1(z) \ln z$ and $T/T_F = 1/f_1(z)$. Here $0 \le z < \infty$ is the fugacity and $f_{\nu}(z) = z - z^2/2^{\nu} + z^3/3^{\nu} - \cdots$ is the Fermi-Dirac function.
- [51] R. Asgari, T. Gokmen, B. Tanatar, M. Padmanabhan, and M. Shayegan, Effective mass suppression in a ferromagnetic two-dimensional electron liquid, Physical Review B 79, 235324 (2009).
- [52] Data taken from table IV of Ref. [15]. These results are extrapolated to the thermodynamic limit based on the data of N = 29,57,101 electrons with a presumed *N*-scaling dependence. The data of Ref. [14] are improved estimates of Ref. [13] so we only plot the former results.
- [53] A. A. Shashkin, M. Rahimi, S. Anissimova, S. V. Kravchenko, V. T. Dolgopolov, and T. M. Klapwijk, Spin-Independent Origin of the Strongly Enhanced Effective Mass in a Dilute 2D Electron System, Physical Review Letters **91**, 046403 (2003), arXiv:0301187 [cond-mat].
- [54] M. Padmanabhan, T. Gokmen, N. C. Bishop, and M. Shayegan, Effective mass suppression in dilute, spin-polarized twodimensional electron systems, Physical Review Letters 101, 026402 (2008), arXiv:0807.1327.
- [55] T. Gokmen, M. Padmanabhan, K. Vakili, E. Tutuc, and M. Shayegan, Effective mass suppression upon complete spinpolarization in an isotropic two-dimensional electron system, *Physical Review B* 79, 195311 (2009).
- [56] S. Gangadharaiah and D. L. Maslov, Spin-independent effective mass in a valley-degenerate electron system, Physical Review Letters 95, 186801 (2005).
- [57] C. P. Goodrich, E. M. King, S. S. Schoenholz, E. D. Cubuk, and M. Brenner, Self-assembling kinetics: Accessing a new design space via differentiable statistical-physics models, (2020), arXiv:2010.15175.
- [58] We note that the indicated r_s value of the same data plotted in Fig. 2 of Ref. [51] and Fig. 9 of Ref. [14] differs by a factor of two.
- [59] S. Das Sarma, V. M. Galitski, and Y. Zhang, Temperaturedependent effective-mass renormalization in two-dimensional electron systems, *Physical Review B* 69, 125334 (2004).
- [60] R. Asgari and B. Tanatar, Many-body effective mass and spin susceptibility in a quasi-two-dimensional electron liquid, Physical Review B 74, 075301 (2006).
- [61] T. Gokmen, M. Padmanabhan, and M. Shayegan, Dependence of effective mass on spin and valley degrees of freedom, Physical Review Letters 101, 146405 (2008).
- [62] T. Salimans, A. Karpathy, X. Chen, and D. P. Kingma, PixelCNN++: Improving the PixelCNN with Discretized Logistic Mixture Likelihood and Other Modifications, (2017), arXiv:1701.05517.
- [63] L. Zhang, J. Han, H. Wang, W. A. Saidi, R. Car, and E. Weinan, End-to-end symmetry preserving inter-atomic potential energy model for finite and extended systems, in *Advances in Neural Information Processing Systems*, Vol. 2018-Decem (2018) arXiv:1805.09003.
- [64] P. Wirnsberger, A. J. Ballard, G. Papamakarios, S. Abercrombie, S. Racanière, A. Pritzel, D. Jimenez Rezende, and C. Blundell, Targeted free energy estimation via learned mappings, Journal of Chemical Physics 153, 144112 (2020), arXiv:2002.04913.

- [65] J. Martens and R. Grosse, Optimizing Neural Networks with Kronecker-factored Approximate Curvature, (2015), arXiv:1503.05671.
- [66] R. Anil, V. Gupta, T. Koren, K. Regan, and Y. Singer, Scalable Second Order Optimization for Deep Learning, (2020), arXiv:2002.09018.
- [67] See https://github.com/fermiflow/CoulombGas for code written in Jax [78], as well as the trained models and corresponding logged data.
- [68] A. Griewank and A. Walther, Algorithm 799: Revolve: An implementation of checkpointing for the reverse or adjoint mode of computational differentiation, ACM Transactions on Mathematical Software 26, 19 (2000).
- [69] P. Goyal, P. Dollár, R. Girshick, P. Noordhuis, L. Wesolowski, A. Kyrola, A. Tulloch, Y. Jia, and K. He, Accurate, Large Minibatch SGD: Training ImageNet in 1 Hour, (2017), arXiv:1706.02677.
- [70] W. Grathwohl, R. T. Chen, J. Bettencourt, I. Sutskever, and D. Duvenaud, Ffjord: Free-form continuous dynamics for scalable reversible generative models, in *7th International Conference on Learning Representations, ICLR 2019* (2019) arXiv:1810.01367.
- [71] J. Behrmann, W. Grathwohl, R. T. Chen, D. Duvenaud, and J. H. Jacobsen, Invertible residual networks, in *36th International Conference on Machine Learning, ICML 2019*, Vol. 2019-June (2019) pp. 894–910, arXiv:1811.00995.
- [72] R. T. Chen, J. Behrmann, D. Duvenaud, and J. H. Jacobsen, Residual flows for invertible generative modeling, in *Advances* in *Neural Information Processing Systems*, Vol. 32 (2019) arXiv:1906.02735.
- [73] T. Q. Chen and D. K. Duvenaud, Neural Networks with Cheap Differential Operators, Advances in Neural Information Processing Systems (NeurIPS) (2019), arXiv:arXiv:1912.03579v1.
- [74] J. Kelly, J. Bettencourt, M. J. Johnson, and D. Duvenaud, Learning differential equations that are easy to solve, arXiv (2020), arXiv:2007.04504.
- [75] C.-W. Huang, R. T. Q. Chen, C. Tsirigotis, and A. Courville, Convex Potential Flows: Universal Probability Distributions with Optimal Transport and Convex Optimization, (2020), arXiv:2012.05942.
- [76] S. Shapira, U. Sivan, P. M. Solomon, E. Buchstab, M. Tischler, and G. B. Yoseph, The thermodynamics of a charged fermion layer at high rs values, Phys. Rev. Lett 77, 3181 (1996).
- [77] S. C. Dultz and H. W. Jiang, Thermodynamic signature of a two-dimensional metal-insulator transition, Physical Review Letters 84, 4689 (2000).
- [78] J. Bradbury, R. Frostig, P. Hawkins, M. J. Johnson, C. Leary, D. Maclaurin, G. Necula, A. Paszke, J. VanderPlas, S. Wanderman-Milne, and Q. Zhang, JAX: composable transformations of Python+NumPy programs (2018).
- [79] J. Bezanson, A. Edelman, S. Karpinski, and V. B. Shah, Julia: A fresh approach to numerical computing, SIAM Review 59, 65 (2017).
- [80] M. Germain, K. Gregor, I. Murray, and H. Larochelle, MADE: Masked Autoencoder for Distribution Estimation, (2015), arXiv:1502.03509.
- [81] F. Becca and S. Sorella, *Quantum Monte Carlo Approaches for Correlated Systems* (Cambridge University Press, 2017) pp. 1– 274.
- [82] S. Mohamed, M. Rosca, M. Figurnov, and A. Mnih, Monte carlo gradient estimation in machine learning, Journal of Machine Learning Research 21, 1 (2020).
- [83] M. F. Hutchinson, A stochastic estimator of the trace of the

influence matrix for laplacian smoothing splines, Communications in Statistics-Simulation and Computation **18**, 1059 (1989).

- [84] P. Zanardi, L. Campos Venuti, and P. Giorda, Bures metric over thermal state manifolds and quantum criticality, Phys. Rev. A 76, 062318 (2007).
- [85] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition* (Cambridge University Press, 2010).

SUPPLEMENTAL MATERIAL

Appendix A: Benchmark for three-dimensional spin-polarized uniform electron gas

We carry out additional benchmark calculation for threedimensional spin-polarized uniform electron gas in a periodic cubic box of length *L*. We use the same network architectures and training procedure as described in the main text, except that $r_s = \left(\frac{3}{4\pi N}\right)^{\frac{1}{3}} \frac{L}{a_0}$ and $k_B T_F = \left(\frac{9\pi}{2}\right)^{\frac{2}{3}} \text{Ry}/r_s^2$ in the threedimensional case. The two panels of Fig. S1 display a typical training process of the kinetic energy *k* and potential energy v per electron, respectively. Note that $r_s = 10$ is within the low-density parameter range where the restricted path integral Monte Carlo method [29] can favorably produce accurate results.



FIG. S1. (a) Kinetic energy and (b) potential energy per electron of three-dimensional spin-polarized uniform electron gas consisting of N = 33 electrons with $r_s = 10$ and $T/T_F = 0.0625$. The red lines are results from the restricted path integral Monte Carlo calculation [29].

Appendix B: Entropy of N non-interacting fermions in the canonical ensemble

To compute the entropy of *N* non-interacting fermions at temperature β , we first compute the partition function Z_N of the system in the canonical ensemble via the recursion formula [41]

$$Z_N = \frac{1}{N} \sum_{\ell=1}^{N} (-1)^{\ell-1} z_\ell Z_{N-\ell},$$
 (S1)

where $Z_0 = 1$ and $z_{\ell} = \sum_k \exp\left(-\ell\beta \frac{\hbar^2 k^2}{2m}\right)$ is the single-particle partition function at temperature $\ell\beta$. Note that Eq. (S1) involves adding exponentially small numbers with alternating signs, thus one needs to use high-precision arithmetics to obtain reliable results for large particle number *N*.

After obtaining Z_N , we evaluate the entropy per particle of

ideal Fermi gas using the standard formula

$$s_0 = \frac{1}{N} \left(\ln Z_N - \beta \frac{\partial}{\partial \beta} \ln Z_N \right).$$
 (S2)

The derivative in Eq. (S2) can be conveniently computed by automatic differentiation through high-precision arithmetics, which is natively supported in, e.g., Julia [79]. Alternatively, one can manually differentiate both sides of (S1) with respect to β to derive a similar recursion relation for the energy $E_N = -\frac{\partial}{\partial \beta} \ln Z_N$:

$$E_N = \frac{1}{Z_N} \frac{1}{N} \sum_{\ell=1}^N (-1)^{\ell-1} z_\ell Z_{N-\ell} (\ell e_\ell + E_{N-\ell}), \qquad (S3)$$

where $e_{\ell} = -\frac{\partial}{\partial(\ell\beta)} \ln z_{\ell}$ is the expected single-particle energy at temperature $\ell\beta$. The starting point of this recursion is, unsurprisingly, $E_0 = 0$.

Appendix C: Model architectures

This section summarizes the network architectures used for the autoregressive model and normalizing flow. They are adapted from the transformer [37] and FermiNet [49], respectively. Please refer to the original publications for more background on these models.

1. Autoregressive model for p(K)

In Algorithm 1, given the momenta k_1, k_2, \ldots, k_N occupied by the *N* electrons, CausalTransformer [37] outputs *N M*dimensional arrays $\hat{k}_1, \hat{k}_2, \ldots, \hat{k}_N$, where \hat{k}_i denotes the logits of the index of k_i corresponding to the conditional probability $p(k_i|k_{<i})$. The number *M* of available momenta $k = \frac{2\pi n}{L}$ is determined by a single-particle energy cutoff $|n|^2 \le \text{Emax}$. For N = 29, 49, 57 in our calculations, Emax is chosen to be 25, 36 and 49, respectively, corresponding to the number of available momenta M = 81, 113 and 149.

Algorithm 1 Autoregressive probabilistic model for a set of momenta.

Input: A set of momenta $K = \{k_1, k_2, ..., k_N\}$. An index function which identifies the momentum to an integer $idx(k) \in \{1, ..., M\}$. **Output:** Log-likelihood ln p(K). 1: $\hat{k}_1, \hat{k}_2, ..., \hat{k}_N$ = CausalTransformer $(k_1, k_2, ..., k_N)$ 2: for each *i* do \triangleright ensure Pauli exclusion principle

3: $\hat{k}_{i,\geq M-N+i+1} = -\infty$ 4: $\hat{k}_{i \leq idx(k_{i-1})} = -\infty$

 $\begin{aligned} \hat{k}_{i,\leq \text{idx}(k_{i-1})} &= -\infty \\ \hat{k}_i &= \hat{k}_i - \log \text{sumexp}(\hat{k}_i) \end{aligned} > \text{normalization}$

6: end for

5:

7: return $\sum_i \hat{k}_{i,idx(k_i)}$

Table S1 summarizes the adopted value of hyperparameters of the network CausalTransformer throughout our calculation. For more implementation details, please refer to our source code [67]. Other architectures of the autoregressive model are also possible, such as the masked autoencoder [80], but our choice here based on the transformer turns out to scale more favorably to large systems regarding the number of trainable parameters involved.

TABLE S1. The hyperparameters of the causal transformer adopted in this work. Note that we choose the non-linear activation of the fully connected neural network involved in the architecture to be tanh, which is smooth enough to avoid potential issues upon automatic differentiation.

Hyperparameter	Description	Value
nlayers	Number of layers	2
modelsize	The size of input and output, also known as the "embedding dimension"	16
nheads	Number of heads of the self-attention part within each layer	4
nhidden	Number of hidden units of the fully connected neural network within each layer	32

2. Normalizing flow for $\Psi_K(R)$

Recall that our goal is not to model a single wavefunction as done in the ground-state calculations, but an exponentially large family of orthonormal many-electron basis states $\Psi_K(\mathbf{R})$. As shown in Eq. (8) of the main text, we achieve this goal by bijectively mapping the original electron coordinates \mathbf{R} to the quasiparticle coordinates ζ .

Algorithm 2 Translation and permutation equivariant coordinate transformation.

Input: Electron coordinates $\mathbf{R} = {\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N}$, network depth d, and box length L. **Output:** Transformed quasiparticle coordinates ζ . \triangleright array shape: (N, 2)1: $f_1 = \text{zeros_like}(\mathbf{R})$ 2: $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$ 3: $\mathbf{f_2} = \left[\sqrt{\sin^2\left(\frac{2\pi x_{ij}}{L}\right) + \sin^2\left(\frac{2\pi y_{ij}}{L}\right)}, \cos\left(\frac{2\pi r_{ij}}{L}\right), \sin\left(\frac{2\pi r_{ij}}{L}\right)\right]$ ▶ array shape: (N, N, 2)4: for $\ell = 1, \cdots, d$ do $g = [f_1, \text{mean}(f_1, \text{axis} = 0), \text{mean}(f_2, \text{axis} = 1)]$ 5: if $\ell = 1$ then 6: $f_1 = FC_{h_1}^{\ell}(g)$ $f_2 = FC_{h_2}^{\ell}(f_2)$ else 7: ▶ array shape: (N, h_1) ▶ array shape: (N, h_2) 8: 9: $f_1 = \operatorname{FC}_{h_1}^{\ell}(g) + f_1$ $f_2 = \operatorname{FC}_{h_2}^{\ell}(f_2) + f_2$ 10: 11: end i 12: 13: end for 14: $f_1 = FC_{h_1}([f_1, mean(f_1, axis = 0), mean(f_2, axis = 1)]) + f_1$ 15: return $R + FC_2(f_1)$

Algorithm 2 implements this transformation using blocks from the FermiNet [49]. The one- and two-electron features f_1 and f_2 are fed into a sequence of *d* fully connected layers $FC_{h_1}^{\ell}$ and $FC_{h_2}^{\ell}$, where h_1, h_2 are the one- and two-particle feature size, respectively, and ℓ ranges from 1 to d. Initially, f_2 contains pairwise distance features of the electrons in a periodic box [81], while f_1 is set to be zero to guarantee the translation equivariance property. Each fully connected layer involved in the algorithm has its own independent parameters, including those at the final stage. In our calculations, the network depth is set to be d = 2 and $h_1 = h_2 = 16$, except for the case N = 57, where $h_1 = h_2 = 32$.

In principle, one may want to ensure invertibility of the coordinate transformation [71] to make for a genuine normalizing flow model. To our best knowledge, there lacks such a rigorous guarantee for Alg. 2 described above. Nevertheless, the practical training process still appears stable. This is probably because the network we adopt is not very deep.

Appendix D: Details of the training procedure

We sample electron momenta K directly from the autoregressive model Eq. (6) in an ancestral manner. Given the momenta, we then sample the electron coordinates R using the Metropolis algorithm according to Born probability of the wavefunction Eq. (8). The batch size is set to be 8192. We compute the Jacobian $\frac{\partial \zeta}{\partial R}$ of the coordinate transformation involved in the wavefunction ansatz using forward-mode automatic differentiation in Jax [78].

Building on these self-generated samples, we train the autoregressive model Eq. (6) and the normalizing flow Eq. (8) jointly to minimize the variational free energy Eq. (5). The gradient estimators with respect to the parameters ϕ and θ in the autoregressive model and normalizing flow, respectively, can be easily derived as follows:

$$\nabla_{\phi}F = \mathop{\mathbb{E}}_{K \sim p(K)} \left[\nabla_{\phi} \ln p(K) \left(\frac{1}{\beta} \ln p(K) + \mathop{\mathbb{E}}_{R \sim |\Psi_{K}(R)|^{2}} \left[E_{K}^{\text{loc}}(R) \right] \right) \right],$$
(S1a)

$$\nabla_{\theta}F = 2\Re \mathbb{E}_{K \sim p(K)} \mathbb{E}_{R \sim |\Psi_{K}(R)|^{2}} \left[\nabla_{\theta} \ln \Psi_{K}^{*}(R) \cdot E_{K}^{\text{loc}}(R) \right], \quad (S1b)$$

where the wavefunction $\Psi_K(\mathbf{R})$ has been assumed to be complex-valued, as suggested by Eq. (8). We can employ the control variate method [32, 35, 82] to further reduce the variance of these estimators.

Below we present some more techniques employed to make the training as efficient as possible.

1. The Hutchinson's trick

The computational bottleneck of the variational free energy Eq. (5) and gradient estimators Eq. (S1) lies in the Laplacian $\nabla^2 \ln \Psi_K(\mathbf{R})$ involved in the local energy, which amounts to computing the trace of the $2N \times 2N$ Hessian matrix H (ln $\Psi_K(\mathbf{R})$). In the standard automatic differentiation approach, one needs to iterate over the rows or columns of the Hessian, which can be inefficient for large systems.

To reduce the computational complexity, we employ the Hutchinson's stochastic trace estimator [83]

$$\nabla^2 \ln \Psi_{\boldsymbol{K}}(\boldsymbol{R}) = \mathop{\mathbb{E}}_{\boldsymbol{\epsilon} \sim f(\boldsymbol{\epsilon})} \left[\boldsymbol{\epsilon}^T \cdot \mathsf{H} \left(\ln \Psi_{\boldsymbol{K}}(\boldsymbol{R}) \right) \cdot \boldsymbol{\epsilon} \right] \qquad (S2)$$

over a 2*N*-dimensional random vector ϵ with zero mean and identity covariance matrix. The probability density $f(\epsilon)$ can, for example, be chosen as a standard Gaussian. The Hessian-vector product involved in Eq. (S2) can be efficiently computed by combining forward- and backward-mode automatic differentiation in Jax [78]. The price we pay, however, is an additional source of randomness on top of the original estimators Eqs. (5) and (S1), which may potentially require more samples to achieve a given statistical accuracy.

In practice, we choose to apply the Hutchinson's trick only to Hessian of the Jacobian determinant term in $\ln \Psi_K(\mathbf{R})$; see Eq. (8). In this way, one can enjoy an overall speedup of roughly one order of magnitude without sacrificing accuracy due to enlarged variance of the estimators.

2. Stochastic reconfiguration for density matrices

The quantity of central interest for the purpose of this work is the thermal entropy, which turns out to be fairly sensitive to the training process. We thus employ the stochastic reconfiguration method [81], which is much more efficient than conventional first-order optimizers like Adam.

In the context of machine learning a classical generative model or the ground-state variational Monte Carlo of quantum systems, the conventional metric for the involved parameter space is well known as the Fisher information. To find an appropriate metric for the present quantum statistical mechanics settings, the arguably most natural candidate is the Bures distance, defined for two density matrices ρ and σ as follows [84]:

$$d_B^2(\rho,\sigma) = 2\left(1 - \operatorname{Tr}\sqrt{\sqrt{\rho}\sigma\,\sqrt{\rho}}\right). \tag{S3}$$

The second term on the right-hand side is well known as the quantum fidelity [85].

Recall from Eq. (4) that $\rho(\phi, \theta) = \sum_{K} p(K; \phi) |\Psi_{K}(\theta)\rangle \langle \Psi_{K}(\theta)|$, where ϕ and θ are the variational parameters in the classical Boltzmann distribution and quantum many-body basis, respectively. By expanding the Bures distance in the neighborhood of a certain point (ϕ, θ) ,

one can obtain [84]

$$d_B^2(\rho(\boldsymbol{\phi}, \boldsymbol{\theta}), \rho(\boldsymbol{\phi} + \Delta \boldsymbol{\phi}, \boldsymbol{\theta} + \Delta \boldsymbol{\theta})) \approx \frac{1}{4} \sum_{ij} \mathcal{I}_{ij} \Delta \phi_i \Delta \phi_j + \sum_{ij} \mathcal{J}_{ij} \Delta \theta_i \Delta \theta_j$$
(S4)

for some positive-definite matrices I and \mathcal{J} . The most important observation is that the desired metric is *block-diagonal* with respect to the two generative models in this work. This fact is favorable in practice, since the metric needs to be inverted to determine the parameter update direction in each training step. Note the size of I, \mathcal{J} are equal to the number of parameters in the autoregressive model and normalizing flow, respectively, both in the order of several thousands throughout our calculations.

I coincides exactly with the classical Fisher information matrix

$$I_{ij} = \sum_{\mathbf{K}} \frac{1}{p(\mathbf{K})} \frac{\partial p(\mathbf{K})}{\partial \phi_i} \frac{\partial p(\mathbf{K})}{\partial \phi_j} = \mathop{\mathbb{E}}_{\mathbf{K} \sim p(\mathbf{K})} \left[\frac{\partial \ln p(\mathbf{K})}{\partial \phi_i} \frac{\partial \ln p(\mathbf{K})}{\partial \phi_j} \right],$$
(S5)

whereas the quantum component $\mathcal J$ reads

$$\mathcal{J}_{ij} = \mathfrak{R}\left(\sum_{\mathbf{K}} p(\mathbf{K}) \left(\frac{\partial \Psi_{\mathbf{K}}}{\partial \theta_i} \middle| \frac{\partial \Psi_{\mathbf{K}}}{\partial \theta_j} \right) - \sum_{\mathbf{K}, \mathbf{K}'} \frac{2p(\mathbf{K})p(\mathbf{K}')}{p(\mathbf{K}) + p(\mathbf{K}')} \left(\frac{\partial \Psi_{\mathbf{K}}}{\partial \theta_i} \middle| \Psi_{\mathbf{K}'} \right) \left(\Psi_{\mathbf{K}'} \middle| \frac{\partial \Psi_{\mathbf{K}}}{\partial \theta_j} \right) \right).$$
(S6)

The double summation over momenta in the second term of Eq. (S6) is inconvenient to estimate. In practice, we choose to approximate \mathcal{J}_{ii} as the following covariance matrix

$$\mathcal{J}_{ij} = \Re \left(\underset{K \sim p(K)}{\mathbb{E}} \underset{R \sim |\Psi_{K}(R)|^{2}}{\mathbb{E}} \left[\frac{\partial \ln \Psi_{K}^{*}(R)}{\partial \theta_{i}} \frac{\partial \ln \Psi_{K}(R)}{\partial \theta_{j}} \right] - \underset{K \sim p(K)}{\mathbb{E}} \underset{R \sim |\Psi_{K}(R)|^{2}}{\mathbb{E}} \left[\frac{\partial \ln \Psi_{K}^{*}(R)}{\partial \theta_{i}} \right] \underset{K \sim p(K)}{\mathbb{E}} \underset{R \sim |\Psi_{K}(R)|^{2}}{\mathbb{E}} \left[\frac{\partial \ln \Psi_{K}(R)}{\partial \theta_{j}} \right] \right)$$
(S7)

Notice the first term is the same as that of Eq. (S6), which is clearly a natural generalization of the usual quantum Fisher information matrix for pure states. The trainings turn out to still behave quite well.

The update rules for the parameters ϕ , θ read as follows:

$$\Delta \boldsymbol{\phi} = -(\boldsymbol{I} + \eta \boldsymbol{1})^{-1} \nabla_{\boldsymbol{\phi}} \boldsymbol{F}, \qquad (S8a)$$

$$\Delta \theta = -(\mathcal{J} + \eta \mathbb{1})^{-1} \nabla_{\theta} F, \qquad (S8b)$$

where we have added a small shift $\eta = 10^{-3}$ to the diagonal of (modified) Fisher information matrices for numerical stability. The gradient norms are constrained within a threshold of 10^{-3} [49], which plays a similar role as the learning rate in other conventional optimizers. See the source code [67] for more details.