

# RIGOROUS COMPUTATION OF LINEAR RESPONSE FOR INTERMITTENT MAPS

ISAIA NISOLI AND TOBY TAYLOR-CRUSH

ABSTRACT. We present a rigorous numerical scheme for the approximation of the linear response of the invariant density of a map with an indifferent fixed point, with explicit and computed estimates for the error and all the involved constants.

## 1. INTRODUCTION

In [38] Ruelle proved that for certain perturbations of uniformly hyperbolic deterministic dynamical systems the underlying SRB measure changes smoothly. He also obtained a formula for the derivative of the SRB measure, called the *linear response formula* [38]<sup>1</sup>. Since then, the topic of linear response has been a very active direction of research in smooth ergodic theory. Indeed, the work of Ruelle was refined in the uniformly hyperbolic setting [12, 25], extended to the partially hyperbolic setting [15], and has been a topic of deep investigation for unimodal maps, see [8], the survey article [7], the recent works [3, 9, 14, 39] and references therein. More recently, the topic of linear response was also studied in the context of random or extended systems [6, 16, 18, 23, 31, 40, 44]. Optimisation of statistical properties through linear response was developed in [1, 2, 22, 30].

Numerical algorithms for the approximation of linear response for uniformly expanding maps, via finite rank transfer operators was obtained in [4] and via dynamical determinants and periodic orbits in [37]<sup>2</sup>, and for uniformly hyperbolic systems [26, 35, 36].

Our work extends the methods in [4] to intermittent maps far from the boundary, allowing us to compute the linear response for LSV maps, a Version of the Manneville-Pomeau family, [34] as the exponent at the indifferent fixed point changes.

Linear response for indifferent fixed point maps has been investigated in [5, 10, 32], but three important questions have to be addressed to obtain a rigorous numerical approximation scheme:

- (1) how to approximate efficiently the involved discretized operators;
- (2) how to bound the approximation errors involved in the discretization;
- (3) how to bound explicitly and efficiently the constants used in the proofs of [5, 10, 32].

---

*Date:* January 12, 2022.

*2020 Mathematics Subject Classification.* Primary 37A05, 37E05.

*Key words and phrases.* Linear Response, Intermittent Maps, Transfer Operators, Rigorous Approximations.

<sup>1</sup>See also earlier related work [29]. See also [24] for a comprehensive historical account including literature from physics.

<sup>2</sup>See also [28] for related work on dynamical determinants.

In our paper we provide answers to the three questions above for general intermittent maps and present an explicit computation for LSV type maps. Our scheme and techniques are very flexible and can be easily adapted to other one dimensional nonuniformly expanding maps whose associated transfer operators do not admit a spectral gap (or a uniform spectral gap) as long as the linear response formula can be obtained via inducing with the first return map.

In the text are presented some numerical remarks, that allow the reader to get an overview of some of the delicate points of the implementation.

The paper is divided as follows: in Section 2 we state the hypothesis on the dynamical system and state our results, in Section 3 we present the theory behind the approximation of the density for the induced map, in Section 4 we discuss the approximation of the linear response for the induced map, in Section 5 we discuss pulling back the measure to the original map and normalizing the density, in Section 6 we give a proof of the fact that the error may be made as small as wanted, in Section 7 we compute an approximation with an explicit error of the linear response for an LSV map; section 8 is devoted to computing effective bounds for the constants in [5, 32] and section 9 explains the technique we use to compute some of the functions involved in our approximation.

#### ACKNOWLEDGEMENTS

The authors would like to thank Prof. Bahsoun and Prof. Galatolo for their guidance and assistance, their patience and attention. Isaia Nisoli was partially supported by CNPq, UFRJ, CAPES (through the programs PROEX and the CAPES-STINT project "Contemporary topics in non uniformly hyperbolic dynamics").

## 2. HYPOTHESIS ON THE MAP AND STATEMENT OF THE RESULTS

We are interested in approximating the invariant density and linear response for one dimensional interval maps with an indeterminate fixed point by inducing. In particular we wish to gain explicitly calculable error bounds in the  $L^1$  norm. We use an induced map on  $[0.5, 1]$ , to gain a map with good statistical properties to approximate an invariant density and linear response, and then using a formula used in [5] to pull back our approximation to the invariant density and linear response of the full map. We apply this method to a family of Pomeau-Manneville maps to gain an approximation of the statistics with explicit error.

**2.1. Interval maps with an inducing scheme.** We introduce now a class (family) of interval maps which are non-uniformly expanding with two branches, for which one can construct an inducing scheme which allow it to inherit the linear response formula from the one for the induced system.

- Let  $V$  be a neighbourhood of 0. For any  $\epsilon \in V$ ,  $T_\epsilon: [0, 1] \rightarrow [0, 1]$  is a non-singular map, with respect to Lebesgue measure,  $m$ , with two onto branches  $T_{0,\epsilon}: [0, 0.5] \rightarrow [0, 1]$  and  $T_{1,\epsilon}: [0.5, 1] \rightarrow [0, 1]$ . The inverse branches of  $T_{0,\epsilon}$ ,  $T_{1,\epsilon}$  are respectively denoted by  $g_{0,\epsilon}$  and  $g_{1,\epsilon}$ . We call  $T_0 := T$  the unperturbed map, and  $T_\epsilon$ , for  $\epsilon \neq 0$ , the perturbed map.
- We assume that for each  $i = 0, 1$  and  $j = 0, 1, 2$  the following partial derivatives exist and satisfy the commutation relation

$$(2.1) \quad \partial_\epsilon g_{i,\epsilon}^{(j)} = (\partial_\epsilon g_{i,\epsilon})^{(j)}.$$

- We assume that  $T_\epsilon$  has a unique absolutely continuous invariant measure<sup>3</sup> (up to multiplication) whose Radon-Nikodym derivative will be denoted by  $h_\epsilon$ , and we denote for simplicity  $h = h_0$ .
- Let  $\hat{T}_\epsilon$ , be the first return map of  $T_\epsilon$  to  $\Delta$ , where  $\Delta := [0.5, 1]$ ; i.e., for  $x \in \Delta$

$$\hat{T}_\epsilon(x) = T_\epsilon^{R_\epsilon(x)}(x),$$

where

$$R_\epsilon(x) = \inf\{n \geq 1 : T_\epsilon^n(x) \in \Delta\}.$$

We assume that  $\hat{T}_\epsilon$  has a unique absolutely continuous invariant measure (up to multiplication) with a continuous density denoted  $\hat{h}_\epsilon \in C^0$ .

- Let  $\Omega$  be the set of finite sequences of the form  $\omega = 10^n$ , for  $n \in \mathbb{N} \cup \{0\}$ . We set  $g_{\omega,\epsilon} = g_{1,\epsilon} \circ g_{0,\epsilon}^n$ . Then for  $x \in [0, 1]$  we have  $T_\epsilon^{n+1} \circ g_{\omega,\epsilon}(x) = x$ . The cylinder sets  $[\omega]_\epsilon = g_{\omega,\epsilon}(\Delta)$ , form a partition of  $\Delta \pmod{0}$ . For  $x \in [0, 1]$ , we assume

$$(2.2) \quad \sup_{\epsilon \in V} \sup_{x \in [0,1]} |g'_{\omega,\epsilon}(x)| < \infty;$$

$$(2.3) \quad \sup_{\epsilon \in V} \sup_{x \in [0,1]} |\partial_\epsilon g_{\omega,\epsilon}(x)| < \infty;$$

$$(2.4) \quad \sum_{\omega} \sup_{\epsilon \in V} \|g'_{\omega,\epsilon}\|_{\mathcal{B}} < \infty;$$

and

$$(2.5) \quad \sum_{\omega} \sup_{\epsilon \in V} \|\partial_\epsilon g'_{\omega,\epsilon}\|_{\mathcal{B}} < \infty,$$

where  $\mathcal{B}$  denotes the set of continuous functions on  $(0, 1]$  with the norm

$$\|f\|_{\mathcal{B}} = \sup_{x \in (0,1]} |x^\gamma f(x)|,$$

for a fixed<sup>4</sup>  $\gamma > 0$ . When equipped with the norm  $\|\cdot\|_{\mathcal{B}}$ ,  $\mathcal{B}$  is a Banach space.

For  $\Phi \in L^1$ , let

$$(2.6) \quad F_\epsilon(\Phi) := 1_\Delta \Phi + (1 - 1_\Delta) \sum_{\omega \in \Omega} \Phi \circ g_{\omega,\epsilon} g'_{\omega,\epsilon}.$$

Note that  $F_\epsilon$  is a linear operator. In fact, for  $x \in [0, 1] \setminus \Delta$ , the formula of  $F_\epsilon$  can be re-written using the Perron-Frobenius operator of  $T_\epsilon$ :

$$(2.7) \quad F_\epsilon(\Phi) := 1_\Delta \Phi + (1 - 1_\Delta) \sum_{k \geq 1} L_\epsilon^k(\Phi \cdot 1_{\{R_\epsilon > k\}}),$$

where  $L_\epsilon$  is the Perron-Frobenius operator associated with  $T_\epsilon$ ; i.e., for  $\varphi \in L^\infty$  and  $\psi \in L^1$

$$\int \varphi \circ T_\epsilon \cdot \psi dm = \int \varphi \cdot L_\epsilon \psi dm.$$

It is given in [5] that the densities of the original system and the induced one are related (modulo normalization in the finite measure case) by

$$(2.8) \quad h_\epsilon = F_\epsilon(\hat{h}_\epsilon).$$

<sup>3</sup>The  $T_\epsilon$  absolutely continuous invariant measure is not assumed to be probabilistic; we allow for  $T_\epsilon$  to admit a  $\sigma$ -finite absolutely continuous invariant measure.

<sup>4</sup>In (2.4) and (2.5) we need the assumptions to hold only for a single  $\gamma$ .

We also define the following operator, which represents  $\partial_\epsilon F_\epsilon \Phi|_{\epsilon=0}$

$$(2.9) \quad Q\Phi = (1 - 1_\Delta) \sum_\omega \Phi' \circ g_\omega \cdot a_\omega g'_\omega + \Phi \circ g_\omega \cdot b_\omega,$$

where  $a_\omega = \partial_\epsilon g_{\omega,\epsilon}|_{\epsilon=0}$  and  $b_\omega = \partial_\epsilon g'_{\omega,\epsilon}|_{\epsilon=0}$ .

**2.2. Interval maps with countable number of branches.** We introduce here a class of interval maps which are uniformly expanding, with a finite or countable number of branches, for which we will be able to prove a linear response formula. The induced map in Subsection 2.1 is a particular case of such uniformly expanding maps.

Let  $\Delta$  be an interval and  $V$  be a neighborhood of 0. Let  $\Omega$  be a finite or countable set. We assume that the maps  $\hat{T}_\epsilon: \Delta \rightarrow \Delta$  satisfy

- For each  $\epsilon \in V$ , there exists a partition (mod 0) of  $\Delta$  into open intervals  $\Delta_{\omega,\epsilon}$ ,  $\omega \in \Omega$  such that the restriction of  $\hat{T}_\epsilon$  to  $\Delta_{\omega,\epsilon}$  is piecewise  $C^3$ , onto and uniformly expanding in the sense that  $\inf_\omega \inf_{\Delta_{\omega,\epsilon}} |\hat{T}'_{\omega,\epsilon}| > 1$ . We denote by  $g_{\omega,\epsilon}$  the inverse branches of  $\hat{T}_\epsilon$  on  $\Delta_{\omega,\epsilon}$ .
- We assume that for each  $\omega \in \Omega$  and  $j = 0, 1, 2$  the following partial derivatives exist and satisfy the commutation relation<sup>5</sup>

$$(2.10) \quad \partial_\epsilon g_{\omega,\epsilon}^{(j)} = (\partial_\epsilon g_{\omega,\epsilon})^{(j)}.$$

- We assume

$$(2.11) \quad \sup_\omega \sup_{\epsilon \in V} \sup_{x \in \Delta} \left| \frac{g''_{\omega,\epsilon}(x)}{g'_{\omega,\epsilon}(x)} \right| < \infty;$$

and for  $i = 2, 3$

$$(2.12) \quad \sum_\omega \sup_{\epsilon \in V} \sup_{x \in \Delta} |g_{\omega,\epsilon}^{(i)}(x)| < \infty;$$

and for  $i = 1, 2$

$$(2.13) \quad \sum_\omega \sup_{\epsilon \in V} \sup_{x \in \Delta} |\partial_\epsilon g_{\omega,\epsilon}^i(x)| < \infty.$$

Let  $\hat{L}_\epsilon$  denote the transfer operator of the map  $\hat{T}_\epsilon$ ; i.e., for  $\Phi \in L^1(\Delta)$

$$\hat{L}_\epsilon \Phi(x) := \sum_{\omega \in \Omega} \Phi \circ g_{\omega,\epsilon}(x) g'_{\omega,\epsilon}(x)$$

for a.e.  $x \in \Delta$ . Under these conditions it is well known that  $\hat{T}_\epsilon$  admits a unique (up to multiplication) finite absolutely continuous invariant measure. We denote its density by  $\hat{h}_\epsilon$ . Hence  $\hat{L}_\epsilon \hat{h}_\epsilon = \hat{h}_\epsilon$ . Moreover,  $\hat{L}_\epsilon$  has a spectral gap when acting on  $C^k$  and  $W^{k,1}$ ,  $k = 1, 2$ . We denote the Perron-Frobenius operator of the unperturbed map  $\hat{T}$  by  $\hat{L}$ ; i.e.,  $\hat{L} := \hat{L}_0$  and let  $\hat{h} := \hat{h}_0$ .

<sup>5</sup>Note that (2.10) is satisfied when  $\hat{T}_\epsilon$  is an induced map as in Subsection 2.1. In particular, for each  $i = 0, 1$  and  $j = 0, 1, 2$  the following partial derivatives exist and satisfy the commutation relation  $\partial_\epsilon g_{i,\epsilon}^{(j)} = (\partial_\epsilon g_{i,\epsilon})^{(j)}$ .

**2.3. Linear response formula.** In [5] it is shown that the invariant density  $\hat{h}_\epsilon$  of the induced map  $\hat{T}_\epsilon$  is differentiable as a  $C^0$  element and its linear response formula is given by

$$(2.14) \quad \hat{h}^* := (I - \hat{L})^{-1} \hat{L}[A_0 \hat{h}' + B_0 \hat{h}],$$

where  $\hat{h}'$  is the spatial derivative of  $\hat{h}$  and

$$A_0 = - \left( \frac{\partial_\epsilon \hat{T}_\epsilon}{\hat{T}'_\epsilon} \right) \Big|_{\epsilon=0}, \quad B_0 = \left( \frac{\partial_\epsilon \hat{T}_\epsilon \cdot \hat{T}_\epsilon''}{\hat{T}'_\epsilon{}^2} - \frac{\partial_\epsilon \hat{T}'_\epsilon}{\hat{T}'_\epsilon} \right) \Big|_{\epsilon=0}.$$

Moreover, for the original map,  $\epsilon \mapsto h_\epsilon$  is differentiable as an element of  $\mathcal{B}$ ; in particular, if the conditions hold for some  $\gamma < 1$

$$\lim_{\epsilon \rightarrow 0} \left\| \frac{h_\epsilon - h}{\epsilon} - h^* \right\|_1 = 0,$$

and  $h^*$  is given by <sup>6</sup>

$$(2.15) \quad h^* = F_0 \hat{h}^* + Q \hat{h}.$$

**2.4. Main result and explicit strategy.** We focus on the case  $\gamma < 1$ . The goal of this work is to provide a numerical scheme that can rigorously approximate  $h^*$ , up to a pre-specified error  $\tau > 0$ , in the  $L^1$ -norm. To obtain such a result we follow the following steps:

- (1) first provide a sequence of finite rank operators  $\hat{L}_\eta$  that can be used to approximate the linear response for the induced map  $\hat{h}^*$  in  $L^1(\Delta)$ . Since the formula of  $\hat{h}^*$  involves  $\hat{h}$  and  $\hat{h}'$ , we will design  $\hat{L}_\eta$  so that its invariant density,  $\hat{h}_\eta$ , well approximates  $h_\eta$  in the  $C^1$ -norm,
- (2) we pull-back to the original map by defining  $F_0^{\text{app}}$  and  $Q_0^{\text{app}}$  by truncating (2.6) and (2.9); i.e., for  $\Phi \in L^1$ ,

$$F_0^{\text{app}}(\Phi) := 1_\Delta \Phi + (1 - 1_\Delta) \sum_{\omega=1}^{N^*} \Phi \circ g_{\omega,0} g'_{\omega,0}$$

and

$$Q^{\text{app}}\Phi = (1 - 1_\Delta) \sum_{\omega=1}^{N^*} \Phi' \circ g_\omega \cdot a_\omega g'_\omega + \Phi \circ g_\omega \cdot b_\omega$$

- (3) finally, find  $N^*$  large enough and set

$$(2.16) \quad h_\eta^* := F_0^{\text{app}} \hat{h}_\eta^* + Q^{\text{app}} \hat{h}_\eta$$

so that

$$\|h_\eta^* - h^*\|_1 \leq \tau.$$

This strategy allows us to prove the following theorem.

<sup>6</sup>Note that in the finite measure case,  $h^*$  is the derivative of the non-normalized density  $h_\epsilon$ . The advantage in working with  $h_\epsilon$  is reflected in keeping the operator  $F_\epsilon$  linear and to accommodate the infinite measure preserving case. In the finite measure case, once the derivative of  $h_\epsilon$  is obtained, the derivative of the normalized density can be easily computed. Indeed,  $h_\epsilon = h + \epsilon h^* + o(\epsilon)$ . Consequently,  $\int h_\epsilon = \int h + \epsilon \int h^* + o(\epsilon)$ . Hence,  $\partial_\epsilon \left( \frac{h_\epsilon}{\int h_\epsilon} \right) \Big|_{\epsilon=0} = h^* - h \int h^*$ .

**Theorem 2.1.** *For any  $\tau > 0$ , there exists a sequence of finite rank operators  $\hat{L}_\eta : L^1(\Delta) \rightarrow L^1(\Delta)$  such that for  $\eta > 0$  small enough and  $N^* > 0$  large enough*

$$\|h_\eta^* - h^*\|_1 \leq \tau.$$

**2.5. The validated numerics toolbox.** While the strategy for the approximation of the linear response may seem quite simple, to make it rigorous, i.e., with a certified control on the error terms so that the results have the strength of proofs, many different quantities have to be estimated explicitly by means of a priori and a posteriori estimates.

The main toolbox we use for these validated estimates consists in

- (1) Interval Arithmetics and rigorous contractors as the Interval Newton Method and the Shooting Method [42]
- (2) discretization of the transfer operator, using the Ulam and Chebyshev basis [20, 19, 43]
- (3) a priori estimate on the tail of a series and rigorous bounds for a finite number of terms.

We will introduce these methods and some of their implementation details during the proof of our result, showing the difference with the cited references when needed.

### 3. APPROXIMATING THE INVARIANT DENSITY OF THE INDUCED MAP

To approximate the invariant density for the induced map two approximation steps are needed. First we need to approximate the induced map, which has countable branches with a map with a finite number of branches. Then, we will discretize the transfer operator of this map by using a Chebyshev approximation scheme.

**3.1. From countable branches to finite branches.** Let  $\delta_k > 0$  with  $\delta_k = |\cup_{n=k}^\infty [\omega]|$ . To simplify notation we assume without loss of generality that  $\frac{1}{2} \in \cup_{n=k}^\infty [\omega]$ . Let

$$\hat{T}_{\delta_k}(x) = \begin{cases} \hat{T}(x) & , \text{ if } x \in [\delta_k, 1], \\ \frac{1}{2}\delta_k^{-1}(x - \frac{1}{2}) + \frac{1}{2} & , \text{ if } x \in [0.5, \delta_k]. \end{cases}$$

Then the transfer operator  $\hat{L}_{\delta_k}$ , associated with  $T_{\delta_k}$  is acting on  $\Phi \in L^1(\Delta)$  as:

$$\hat{L}_{\delta_k} \Phi(x) := \sum_{\substack{\omega \in \Omega \\ n < k}} \Phi \circ g_{\omega, \epsilon}(x) g'_{\omega, \epsilon}(x) + \Phi \left( \delta_k(2x - 1) + \frac{1}{2} \right) 2\delta_k$$

for a.e.  $x \in \Delta$ .

**Lemma 3.1.** *Let  $\Phi \in C^1$ , then*

$$\|(\hat{L} - \hat{L}_{\delta_k})\Phi\|_{C^1} \leq (D + D_0 D + 2)\|\Phi\|_{C^1} \delta_k,$$

where  $D_0 = \left\| \frac{g''_\omega}{g'_\omega} \right\|_\infty$  and  $D \geq 2 \sup_\omega \frac{|g'_\omega(x)|}{|g'_\omega(y)|}$  for all<sup>7</sup>  $x, y$  in  $[0.5, 1]$ .

<sup>7</sup>The existence of a uniform constant  $D > 0$  is implied by condition (2.11).

*Proof.* First notice that

$$(3.1) \quad \begin{aligned} \left| (\hat{L} - \hat{L}_{\delta_k})\Phi \right| &= \left| \sum_{\substack{\omega \in \Omega \\ n \geq k}} \Phi \circ g_\omega(x) g'_\omega(x) - \Phi(\delta_k(2x-1) + \frac{1}{2}) 2\delta_k \right| \\ &\leq \sum_{\substack{\omega \in \Omega \\ n \geq k}} |\Phi \circ g_\omega(x)| \cdot |g'_\omega(x)| + |\Phi(\delta_k(2x-1) + \frac{1}{2})| 2\delta_k \end{aligned}$$

and

$$(3.2) \quad \begin{aligned} &\left| \left( (\hat{L} - \hat{L}_{\delta_k})\Phi \right)' \right| \\ &= \left| \sum_{\substack{\omega \in \Omega \\ n \geq k}} (\Phi' \circ g_\omega(x) (g'_\omega(x))^2 + \Phi \circ g_\omega(x) g''_\omega(x)) - |\Phi'(\delta_k(2x-1) + \frac{1}{2})| 4\delta_k^2 \right| \\ &\leq \sum_{\substack{\omega \in \Omega \\ n \geq k}} |\Phi' \circ g_\omega(x)| \cdot (g'_\omega(x))^2 + \sup_\omega \left\| \frac{g''_\omega}{g'_\omega} \right\|_\infty \sum_{\substack{\omega \in \Omega \\ n \geq k}} |\Phi \circ g_\omega(x)| \cdot |g'_\omega(x)| \\ &\quad + |\Phi'(\delta_k(2x-1) + \frac{1}{2})| 4\delta_k^2. \end{aligned}$$

Now notice that by the Mean Value Theorem,  $\exists \xi_\omega \in (\frac{1}{2}, 1)$  such that

$$|g_\omega(1) - g_\omega(\frac{1}{2})| = |g'_\omega(\xi_\omega)|/2.$$

Therefore,

$$(3.3) \quad |g'_\omega(x)| \leq 2|g_\omega(1) - g_\omega(\frac{1}{2})| \cdot \sup_\omega \frac{|g'_\omega(x)|}{|g'_\omega(\xi_\omega)|} := D \cdot |g_\omega(1) - g_\omega(\frac{1}{2})|.$$

Thus, using (6.4) in (6.2) and (6.3), we obtain

$$(3.4) \quad \begin{aligned} &\left| (\hat{L} - \hat{L}_{\delta_k})\Phi \right| + \left| \left( (\hat{L} - \hat{L}_{\delta_k})\Phi \right)' \right| \leq \|\Phi\|_{C^0} D \sum_{\substack{\omega \in \Omega \\ n \geq k}} |g_\omega(1) - g_\omega(\frac{1}{2})| + \|\Phi\|_{C^0} 2\delta_k \\ &\quad + \left( \|\Phi'\|_{C^0} D + \|\Phi\|_{C^0} D \sup_\omega \left\| \frac{g''_\omega}{g'_\omega} \right\|_\infty \right) \sum_{\substack{\omega \in \Omega \\ n \geq k}} |g_\omega(1) - g_\omega(\frac{1}{2})| + \|\Phi'\|_{C^0} 4\delta_k^2 \\ &= (D + D_0 D + 2) \|\Phi\|_{C^0} \delta_k + \|\Phi'\|_{C^0} 4\delta_k^2 \leq (D + D_0 D + 2) \|\Phi\|_{C^1} \delta_k. \end{aligned}$$

□

The next lemma shows that using the above information, the densities  $\hat{h}$  and  $\hat{h}_{\delta_k}$  can be made arbitrarily close in  $C^1$ .

**Lemma 3.2.** *For two operators,  $L_1$  and  $L_2$ , with fixed points  $h_1$  and  $h_2$  normalised with respect to  $\|\cdot\|_1$ , and a shared Lasota-Yorke inequality*

$$\|L_i^n f\|_s \leq A\lambda^n \|f\|_s + B\|f\|_w$$

for  $i \in \{1, 2\}$ , there is a  $C^*$  such that  $\|L_i^N f\|_s \leq C^* \|f\|_s$ , and for any  $N \geq 1$  we have

$$\|h_1 - h_2\|_s \leq \|L_1^N(h_1 - h_2)\|_s + NC^* \|(L_1 - L_2)h_2\|_s.$$

Furthermore if  $\|L_1^N|_{U^0}\|_s \leq C_N < 1$  then we can have

$$\|h_1 - h_2\|_s \leq \frac{NC^* \|(L_1 - L_2)h_2\|_s}{1 - C_N}.$$

*Proof.* The value of  $C^*$  is given by  $A\lambda + B$ , and the distance between the two fixed points is shown as follows,

$$\begin{aligned} \|h_1 - h_2\|_s &\leq \|L_1^N h_1 - L_2^N h_2\|_s \\ &\leq \|L_1^N(h_1 - h_2)\|_s + \|(L_1^N - L_2^N)h_2\|_s. \end{aligned}$$

Note that

$$\begin{aligned} (L_1^N - L_2^N)h_2 &= \sum_{k=1}^N L_1^{N-k}(L_1 - L_2)L_2^{k-1}h_2 \\ &= \sum_{k=1}^N L_1^{N-k}(L_1 - L_2)h_2. \end{aligned}$$

Consequently,

$$\begin{aligned} \|(L_1^N - L_2^N)h_2\|_s &\leq \sum_{k=1}^N C^* \|(L_1 - L_2)h_2\|_s \\ &\leq NC^* \|(L_1 - L_2)h_2\|_s. \end{aligned}$$

Given that  $\|h_1\|_1 = \|h_2\|_1$  we have that  $h_1 - h_2 \in U^0$  and therefore we can bound  $\|L_1^N(h_1 - h_2)\|_s$  by  $C_N \|h_1 - h_2\|_s$ , rearranging gives us the last result.  $\square$

**Remark 3.3.** The operators  $\hat{L}$  and  $\hat{L}_{\delta_k}$  admit a uniform Lasota-Yorke inequality,

$$\|\hat{L}^n f\|_{C^1} \leq A\lambda^n \|f\|_{C^1} + B \|f\|_{C^0}$$

as shown in section 8.2.1, where a value for  $C^*$  is found. Bounds on  $\|L_{\delta_k}|_{U^0}\|_{C^1}$  can be found by techniques described in section 3.2.4. The  $C^1$  norms of  $\hat{h}_{\delta_k}$  and  $\hat{h}$  can be estimated using the Lasota-Yorke inequalities. We can then use lemma 3.1 to make the error in  $\|\hat{h}_{\delta_k} - \hat{h}\|_{C^1}$  as small as we like.

Next we define a finite rank operator to obtain  $\hat{h}_n$  so that  $\|\hat{h}_n - \hat{h}_{\delta_k}\|_{C^1}$  can be made as small as required.

**3.2. Approximating the invariant density for  $\hat{T}_{\delta_k}$ .** To approximate the invariant density, we will discretize the operator  $\hat{L}_{\delta_k}$  using the basis of the Chebyshev polynomials of the first kind. The Chebyshev basis is a basis for the space of polynomials with a main advantage: given a continuous function  $f$  on  $[-1, 1]$  the interpolating polynomial on the Chebyshev points are "near-best" approximants with respect to  $\|\cdot\|_\infty$  [41, Theorem 16.1]; moreover if the function  $f$  is regular enough the coefficients of the interpolant decay "fast" and are easily computed by means of the Fast Fourier Transform.

Before going forward, some observations are in order, since Chebyshev polynomials do not solve all the problems involved with approximation: to apply this approximation scheme we need to prove a priori that our stationary density is

regular enough and keep track of all the errors involved in the computation of the coefficients. Moreover, evaluating a Chebyshev polynomial of high degree rigorously is a delicate matter [33].

3.2.1. *Chebyshev interpolation and projection.* The material for this section comes from [41], [27] and [45].

Given an  $f \in W^{k,p}$  from  $[-1, 1] \rightarrow \mathbb{C}$ , we can define a function  $\mathcal{F}(\theta)$  on  $[0, 2\pi]$  by

$$\mathcal{F}(\theta) = f(\cos(\theta)).$$

The Fast Fourier Transform (FFT) on a grid of size  $2N$  allows us to compute the coefficients  $a_k$  of the trigonometric polynomial interpolating  $\mathcal{F}$  on an equispaced grid  $\theta_i = (2\pi i)/(2N)$ , for  $i$  in  $0, \dots, 2N - 1$ .

Let  $x_i = \cos(\theta_i)$  for  $i$  in  $1, \dots, N$ ; observe that  $\mathcal{F}(\theta_i) = \mathcal{F}(\theta_{2N-i}) = f(x_i)$ . We define the Chebyshev polynomials by the relation

$$T_n(\cos(\theta)) = \cos(n\theta),$$

then, if we denote by  $b_0 = a_0/2$ ,  $b_{N-1} = a_{N-1}/2$  and  $b_i = a_i$  for all  $i = 1, \dots, N-2$ :

$$p(x) = \sum_{k=0}^{N-1} b_k T_k(x),$$

where the  $a_k$  are the ones computed by the FFT is the interpolating polynomial of  $f$  on the grid given by the  $x_i$ .

**Definition 3.4.** Let  $f \in W^{k,1}$ ,  $k > 1$ , we define the **Chebyshev (interpolating) projection**

$$\pi_n f = \sum_{k=0}^n a_k T_k(x).$$

**Remark 3.5.** If, instead of the FFT we had taken the Fourier transform of  $\mathcal{F}$ , the Fourier coefficients  $\hat{a}_k$  would define coefficients  $\hat{b}_k$ , the Chebyshev orthogonal expansion

$$f(x) = \sum_{k=0}^{+\infty} \hat{b}_k T_k(x),$$

and the **Chebyshev projection**

$$\hat{\pi}_n f = \sum_{k=0}^n \hat{b}_k T_k(x).$$

The coefficients  $a_k$  and  $\hat{a}_k$  are related by the aliasing relation:

$$a_k = \sum_{p \in \mathbb{Z}} \hat{a}_{k+p2n},$$

a-priori knowledge of the regularity of  $f$  allows to estimate the aliasing error above.

This foundational Theorem from [41] estimates the decay rate of the Chebyshev coefficients.

**Theorem 3.6.** For an integer  $\nu \geq 0$ , let  $f$  and its derivatives through  $f^{(\nu-1)}$  be absolutely continuous on  $[-1, 1]$  and suppose the  $\nu^{\text{th}}$  derivative  $f^{(\nu)}$  is of bounded variation  $V$ . Then for  $k \geq \nu + 1$ , the Chebyshev coefficients of  $f$  satisfy

$$|\hat{b}_k| \leq \frac{2V}{\pi k(k-1)\dots(k-\nu)} \leq \frac{2V}{\pi(k-\nu)^{\nu+1}}.$$

The decay rate of Chebyshev coefficients allows us to estimate the projection error in  $C^0$  and  $C^1$  norm.

**Theorem 3.7.** If  $f$  satisfies the conditions of Theorem 3.6, with  $V$  again the total variation of  $f^{(\nu)}$  for some  $\nu \geq 1$ , then for any  $n \geq \nu$ , its Chebyshev projection satisfies

$$\|f - \pi_n f\|_\infty \leq \frac{2V}{\pi \nu n(n-1)\dots(n+1-\nu)}$$

The following theorem is a consequence of combining theorem 3.6, with the proof of theorem 2.3 from [45],

**Theorem 3.8.** If  $f, f', \dots, f^{(\nu-1)}$  are absolutely continuous on  $[-1, 1]$  and if  $\|f^{(\nu)}\|_1 = V < \infty$  for some  $\nu \geq 0$ , then for each  $n \geq \nu + 1$ , we have that for  $\nu > 2$

$$\|f' - (\pi_n f)'\|_\infty \leq \frac{4(n+1)V}{n(\nu-2)\pi(n-2)(n-3)\dots(n+1-\nu)}.$$

*Proof.* From the proof of theorem 2.3 from [45] we have

$$\|f' - (\pi_n f)'\|_\infty \leq 2 \sum_{k=n+1}^{\infty} |a_k| k^2$$

and theorem 3.6 then gives

$$\begin{aligned} \|f' - (\pi_n f)'\|_\infty &\leq \sum_{k=n+1}^{\infty} \frac{4Vk^2}{\pi k(k-1)\dots(k-\nu)} \\ &\leq \frac{4(n+1)V}{n(k-2)\pi(n-2)(n-3)\dots(n+1-k)}. \end{aligned}$$

□

We can use these theorems to bound the error of Chebyshev projections in the  $C_1$  norm.

**3.2.2. Numerical remarks: FFT and Chebyshev.** It is important to have an explicit estimate of the error on the coefficients introduced by the FFT. The main issue here is that, when we computing Chebyshev points and evaluating the function  $f$  are not exact operation. To compute rigorous inclusions of the true mathematical value, we use Interval Arithmetics [42].

This means that we need to compute the FFT of a vector of intervals, not of floating point numbers. The following is the consequence of a classical result from [27] that allows us to find a vector of intervals that encloses the Fast Fourier Transform of any element of the vector of the values. This allows us to use optimized implementations of the FFT algorithm as FFTW [17].

**Lemma 3.9.** *Let  $\tilde{f}$  be a vector of intervals of dimension  $N$ ,  $f_m$  the vector of their midpoints,  $f_r$  the vector of their radiuses. Let  $\hat{a}$  be the computed FFT of  $f_m$ . Then*

$$\|\hat{a} - \bar{a}\|_2 \leq \frac{\log_2(N)}{\sqrt{N}} \left( \frac{\eta}{1-\eta} \|f_m\|_2 + \|f_r\|_2 \right).$$

where  $\bar{a}$  is the exact FFT for any  $f \in \tilde{f}$ ,  $\eta = \mu + \gamma_4(\sqrt{2} + \mu)$  with  $\mu$  the absolute error in the computation of the twiddle factors and  $\gamma_4 = 4u/(1-4u)$  where  $u$  is the unit roundoff.

3.2.3. *Bounding the error on the invariant density.* Let  $\pi_n$  be the Chebyshev projection and let

$$\hat{L}_\eta = \pi_n \hat{L}_{\delta_k} \pi_n$$

be our finite rank approximation of  $\hat{L}_{\delta_k}$ .

**Lemma 3.10.** *If  $1/\hat{T}_{\delta_k}$  is of class  $C^\nu$  then  $\hat{L}_{\delta_k}$  admits Lasota-Yorke like inequalities of the form*

$$\left\| (\hat{L}_{\delta_k}^n f)^{(k)} \right\|_1 \leq (\lambda^k)^n \left\| f^{(k)} \right\|_1 + \tilde{A}_k \|f\|_{W^{k-1,1}},$$

for some  $\nu \in \mathbb{N}$  for  $k = 1, \dots, \nu$ . This implies that if  $h_{\delta_k}$  is a fixed point of  $\hat{L}_{\delta_k}$

$$\|h_{\delta_k}\|_{W^{k,1}} \leq \frac{A_k}{1-\lambda^k} \|h_{\delta_k}\|_{W^{k-1,1}}$$

**Remark 3.11.** *The Lasota-Yorke inequalities give us an upper bound on the  $W^{k,1}$  norm of the fixed point. This, together with Theorems 3.7 and 3.8 permit us to control the discretization error. To estimate our error, we need to compute the constants of this Lasota-Yorke inequality explicitly, we refer to Subsection 7.2.3 for the technique we used.*

**Remark 3.12.** *We can use theorems 3.7 and 3.8, together with lemma 3.10 to get a bound on  $\|\hat{L}_{\delta_k} - \hat{L}_n\|_{C^1}$  and the techniques from section 3.2.4 to with lemma 3.2 using operators  $\hat{L}_{\delta_k}$  and  $\hat{L}_n$  in order to bound  $\|\hat{h}_{\delta_k} - \hat{h}_n\|_{C^1}$  explicitly.*

*This approach is now quite established, a full treatment can be found in [20, 19].*

**Remark 3.13.** *The discretized operator obtained by the Chebyshev discretization does not preserve the value of the integral. To solve this issue, as in [19] we correct the behaviour of the discretized operator by defining a new operator*

$$\hat{Q}_n f = \hat{L}_n f + 1 \cdot \left( \int f dx - \int \hat{L}_n f dx \right)$$

which is guaranteed to preserve the space of average 0 measure and has eigenvalue 1, since the row vector that contains the integrals of the basis elements is preserved by multiplication on the left.

3.2.4. *Numerical remarks: convergence rates.* The problem of bounding the error in  $C^1$  is now reduced to estimate  $c_N$  such that  $\|\hat{L}_n^N|_{U^0}\|_{C^1} \leq c_N$ . Since the operator  $L_n$  is of finite rank, we can use numerical methods to compute these quantities in a rigorous way.

Given a basis  $\{e_i\}_i$  of  $U^0$  normalized with respect to the  $C^1$  norm, a generic function  $v$  in  $U^0$  is written as  $v = \sum_{i=1}^N b_i e_i$ . We want to find a constants  $C_k$  such that

$$\|\hat{L}_n^k v\|_{C^1} \leq C_k \|v\|_{C^1}.$$

If a  $c_k$  exists such that for all basis elements  $e_i$  we have

$$\|\hat{L}_n^k e_i\|_{C^1} \leq c_k.$$

then

$$\|\hat{L}_n^k v\|_{C^1} \leq c_k \sum_{i=1}^N |b_i| \leq c_k \|b\|_{\ell^1}$$

where the  $\ell_1$  norm is the linear algebra norm on the coefficients; we will exhibit a constant  $D$  such that  $\|\cdot\|_{\ell^1} \leq D \|\cdot\|_{C^1}$ ; then  $C_k \leq D c_k$ .

A basis of  $U^0$  in the Chebyshev basis is given by

$$e_i = \frac{g_i}{\|g_i\|_{C^1}}, \quad \text{where } g_i(x) = T_i(x) - \int_{-1}^1 T_i(x) dx.$$

We can link the Chebyshev coefficients  $a_i$  and  $b_i$  by

$$a_0 = - \sum_{i=1}^N b_i \int_{-1}^1 T_i(x) dx$$

and  $a_i = b_i$  for  $i > 0$ . We can use theorem 3.6 to say

$$|a_k| \leq \frac{\|v\|_{C^1}}{k}$$

and therefore

$$\|b\|_{\ell^1} = \sum_{k=1}^N |b_k| \leq \sum_{k=1}^N \frac{\|v\|_{C^1}}{k} \leq \log(N+1) \|v\|_{C^1}$$

so we have  $D = \log(N+1)$  and

$$\|\hat{L}_n^k v\|_{C^1} \leq c_k \log(N+1) \|v\|_{C^1}.$$

Computationally if we take  $N$  functions

$$\hat{e}_i = \frac{g_i}{\|g_i\|_{C^1}^-},$$

where  $\|g_i\|_{C^1}^-$  is a lower bound on the  $C^1$  norm of  $g_i$  and if we calculate each

$$\|\hat{L}_n^k e_i\|_{C^1}$$

then take the maximum value and call it  $\hat{c}_k$ , then  $\hat{c}_k \log(N+1)$  is an upper bound on the  $C^1$  contraction of  $\|\hat{L}_n^k|_{U^0}\|_{C^1}$ .

It is important to explain how we compute an upper bound for the  $C^1$  norm: we use a classical optimization algorithm in IntervalArithmetics [42] that allows us to give a certified upper bound, implemented in the Julia package IntervalOptimisation.jl. The main issue here is that the Clenshaw algorithm is prone to overestimation when evaluated on intervals [33]; to solve this we extended the algorithms in [33] to get tighter bound for the maximum of a Chebyshev polynomial and its derivative.

## 4. APPROXIMATING THE LINEAR RESPONSE FOR THE INDUCED SYSTEM

We now provide an approximation of  $\hat{h}^*$  in the  $L^1$ -norm, through the use of the Ulam approximation; we refer to [19, 20] for an in deep treatment of the Ulam discretization.

**Definition 4.1.** *The Ulam projection, is a projection  $\Pi_u : L^1([0.5, 1]) \rightarrow L^1([0.5, 1])$  over a partition of  $[0.5, 1]$ ,  $\mathcal{P}$ ,*

$$\Pi_u f(x) = \frac{1}{|A|} \int_A f(x) dx$$

where  $x \in A \in \mathcal{P}$ . The Ulam discretisation of the transfer operator  $L_{\delta_k}$  is

$$\hat{L}_u := \Pi_u \hat{L}_{\delta_k} \Pi_u.$$

**4.1. Error in approximating the linear response.** Set

$$(4.1) \quad \hat{h}_\eta^* := \sum_{n=0}^{l^*} \hat{L}_u^n \hat{L}_{\delta_k} [A_0 \hat{h}'_n + B_0 \hat{h}_n].$$

where  $\hat{L}_u$  is the Ulam approximation of  $\hat{L}_{\delta_k}$ .

**Lemma 4.2.** *Recall from (2.14) that*

$$\hat{h}^* = (I - \hat{L})^{-1} \hat{L} [A_0 \hat{h}' + B_0 \hat{h}]$$

We have

$$(4.2) \quad \begin{aligned} \|\hat{h}_\eta^* - \hat{h}^*\|_1 &\leq \sum_{n=0}^{l^*} \sum_{i=0}^n \left\| (\hat{L}_u - \hat{L}_{\delta_k}) \hat{L}_u^{n-i} W_u \right\|_1 + \sum_{n=0}^{l^*} \sum_{i=0}^n \left\| (\hat{L}_{\delta_k} - \hat{L}) \hat{L}_u^{n-i} W_u \right\|_1 \\ &\quad + l^* \|(W_u - W)\|_1 + \sum_{n=l^*+1}^{\infty} \left\| \hat{L}^n W \right\|_1 \end{aligned}$$

where  $W = \hat{L} [A_0 \hat{h}' + B_0 \hat{h}]$  and

$$\begin{aligned} W_u(x) &= \sum_{\omega \leq k} \frac{1}{\eta} \int_{g_\omega(I_i)} A_0(z) \cdot \hat{h}'_n(\zeta) + B_0(z) \hat{h}_n(\zeta) dz \\ &\quad + \frac{1}{\eta} \int_{\delta_k(2I_i-1)+1/2} [A_0 \hat{h}'_\eta + B_0 \hat{h}_\eta](z) dz \end{aligned}$$

for  $x \in I_i$  and  $\zeta \in g_\omega(I_i)$ .

*Proof.* First we must recall from (4.1) that

$$\hat{h}_\eta^* = \sum_{n=0}^{l^*} \hat{L}_u^n \Pi_u L_{\delta_k} [A_0 \hat{h}'_n + B_0 \hat{h}_n].$$

Take  $x \in I_i$ , then we have

$$\begin{aligned} & \Pi_u L_{\delta_k} [A_0 \hat{h}'_n + B_0 \hat{h}_n](x) \\ &= \frac{1}{\eta} \int_{I_i} \sum_{\omega \leq k} [A_0 \hat{h}'_n + B_0 \hat{h}_n] \circ g_\omega(y) g'_\omega(y) + [A_0 \hat{h}'_n + B_0 \hat{h}_n] \circ (\delta_k(2y-1) + 1/2) \delta_k dy \\ &= \frac{1}{\eta} \sum_{\omega \leq k} \int_{g_\omega(I_i)} [A_0 \hat{h}'_n + B_0 \hat{h}_n](z) dz + \frac{1}{\eta} \int_{\delta_k(2I_i-1)+1/2} [A_0 \hat{h}'_n + B_0 \hat{h}_n](z) dz. \end{aligned}$$

The estimate follows by direct calculation. Indeed, by (2.14) and the definition of  $\hat{h}_\eta^*$  we have

$$\begin{aligned} \left\| \hat{h}^* - \hat{h}_\eta^* \right\|_1 &= \left\| \sum_{n=0}^{\infty} \hat{L}^n W - \sum_{n=0}^{l^*} \hat{L}_u^n W_u \right\|_1 \\ &\leq \left\| \sum_{n=0}^{l^*} \hat{L}^n W - \sum_{n=0}^{l^*} \hat{L}_u^n W_u \right\|_1 + \sum_{n=l^*+1}^{\infty} \left\| \hat{L}^n W \right\|_1 \end{aligned}$$

which, by using the triangle inequality and the second resolvent identity, is bounded above by

$$\begin{aligned} & \left\| \sum_{n=0}^{l^*} \hat{L}^n W_u - \sum_{n=0}^{l^*} \hat{L}_u^n W_u \right\|_1 + \left\| \sum_{n=0}^{l^*} (\hat{L}^n W - \hat{L}_u^n W_u) \right\|_1 + \sum_{n=l^*+1}^{\infty} \left\| \hat{L}^n W \right\|_1 \\ &\leq \sum_{n=0}^{l^*} \sum_{i=0}^n \left\| \hat{L}^i (\hat{L}_u - \hat{L}) \hat{L}_u^{n-i} W_u \right\|_1 + \sum_{n=0}^{l^*} \left\| \hat{L}^n (W_u - W) \right\|_1 + \sum_{n=l^*+1}^{\infty} \left\| \hat{L}^n W \right\|_1 \\ &\leq \sum_{n=0}^{l^*} \sum_{i=0}^n \left\| (\hat{L}_u - \hat{L}_{\delta_k}) \hat{L}_u^{n-i} W_u \right\|_1 + \sum_{n=0}^{l^*} \sum_{i=0}^n \left\| (\hat{L}_{\delta_k} - \hat{L}) \hat{L}_u^{n-i} W_u \right\|_1 \\ &+ l^* \|(W_u - W)\|_1 + \sum_{n=l^*+1}^{\infty} \left\| \hat{L}^n W \right\|_{C^1} \end{aligned}$$

□

**Remark 4.3.** *The estimates in Lemma 4.2 can all be made as small as desired. Indeed, notice that  $\hat{L}[A_0 \hat{h}' + B_0 \hat{h}]$  is a zero average  $C^1$  function; therefore*

- *the last summand  $\sum_{n=l^*+1}^{\infty} \|\hat{L}^{n+1} W\|_{C^1}$  can be made, for sufficiently large  $l^*$ , small since  $\hat{L}$  admits a spectral gap when acting on  $C^1$ . Once this term is estimated,  $l^*$  is fixed once and for all;*
- *the summand  $\sum_{n=0}^{l^*} \sum_{i=1}^n \|(\hat{L} - \hat{L}_{\delta_k}) \hat{L}_u^{n-i} W_u\|_1$  can be made small by choosing  $\delta_k$  small enough;*
- *the summand  $\sum_{n=0}^{l^*} \sum_{i=1}^n \|(\hat{L}_{\delta_k} - \hat{L}_u) \hat{L}_u^{n-i} W_u\|_1$  can be made small by choosing  $\eta$ , the size of the Ulam discretization, small enough;*
- *the term  $l^* \|(W_u - W)\|_1$  can be made small by reducing  $\delta_k$ .*

## 5. NORMALISING THE DENSITY AND THE LINEAR RESPONSE.

Ultimately the goal is to approximate the dynamics of the system, so we would like the invariant measure to be a probability measure. This is not always possible

for maps with indeterminate fixed points, however it was shown in [10] that a fixed point of the transfer operator of an LSV map is bounded above by  $Cx^{-\alpha}$ , for some constant  $C$ , giving a maximum integral of  $\frac{C}{1-\alpha}$ , so we can make our calculated density a probability density by normalising with respect to its integral, which will give us a new error of

$$\begin{aligned} \left\| \frac{h_n}{\int h_n dm} - \frac{h}{\int h dm} \right\|_1 &= \left\| \frac{h_n}{\int h_n dm} - \frac{h}{\int h_\eta dm} + h \frac{\int h dm - \int h_n dm}{\int h dm \int h_n dm} \right\|_1 \\ &\leq \frac{\|h - h_n\|_1}{\int h_n dm} + \frac{\|h\|_1}{\int h dm} \frac{\|h - h_n\|_1}{\int h_n dm} \leq 2 \frac{\|h - h_n\|_1}{\int h_n dm} \end{aligned}$$

and if we ensure that the integral is preserved throughout the approximation then the error is  $\frac{\|h^* - h_\eta^*\|_1}{\int h_\eta^* dm}$ .

Since the Chebyshev approximation does not preserve the integral we use the first estimate to bound

$$\left\| \frac{h_n}{\int h_n dm} - \frac{h}{\int h dm} \right\|_1$$

where we calculate the integral of  $h_n$  by

$$\int_0^1 F^{\alpha pp} \hat{h}_n dx = \int_{0.5}^1 \hat{h}_n dx + \sum_{|\omega|=1}^{N^*} H_n \circ g_\omega(1) - H_n \circ g_\omega(0.5)$$

where  $H_n(x) = \int_{0.5}^x h_n dx$ .

The linear response for the normalised invariant density is then  $\rho^*$  such that

$$\lim_{\epsilon \rightarrow 0} \left\| \frac{\frac{h}{\int h dm} - \frac{h_\epsilon}{\int h_\epsilon dm}}{\epsilon} - \rho^* \right\|_1 = 0$$

we get from this

$$\begin{aligned} \left\| \frac{\frac{h}{\int h dm} - \frac{h_\epsilon}{\int h_\epsilon dm}}{\epsilon} - \rho^* \right\|_1 &= \left\| \frac{\frac{h}{\int h dm} - \frac{h_\epsilon}{\int h + \epsilon h^* + o(\epsilon^2) dm}}{\epsilon} - \rho^* \right\|_1 \\ &= \left\| \frac{\frac{h}{\int h dm} - \frac{h_\epsilon}{\int h dm}}{\epsilon} - \frac{h_\epsilon \int [h^* + o(\epsilon)] dm}{\int h dm (\int [h + \epsilon h^* + o(\epsilon^2)] dm)} - \rho^* \right\|_1 \end{aligned}$$

which tells us that

$$\rho^* = \frac{h^*}{\int h dm} - \frac{h \int h^* dm}{(\int h dm)^2}$$

Letting

$$A = \frac{h^*}{\int h dm} - \frac{h_\eta^*}{\int h_\eta dm} \quad B = -\frac{h \int h^* dm}{(\int h dm)^2} + \frac{h_n \int h_\eta^* dm}{(\int h_n dm)^2}$$

the error on the normalised linear response is calculated as follows,

$$\begin{aligned}
& \left\| \frac{h^*}{\int h dm} - \frac{h \int h^* dm}{(\int h dm)^2} - \frac{h_n^*}{\int h_n dm} + \frac{h_n \int h_n^* dm}{(\int h_n dm)^2} \right\|_1 \\
&= \left\| \frac{h^*}{\int h_n dm} - \frac{h_n^*}{\int h_n dm} + \frac{h^* \int (h_n - h) dm}{\int h dm \int h_n dm} + B \right\|_1 \\
&= \left\| \frac{h^* - h_n^*}{\int h_n dm} + \frac{h^* \int (h_n - h) dm}{\int h dm \int h_n dm} + B \right\|_1 \\
&= \left\| A + \frac{h_n \int h_n^* dm}{(\int h_n dm)^2} - \frac{h \int h^* dm}{(\int h dm)^2} \right\|_1 \\
&\leq \left\| A + \frac{h_n \int h_n^* dm}{(\int h_n dm)^2} - \frac{(h_n + (h - h_n))(\int h_n^* dm - \|h^* - h_n^*\|_1)}{(\int h_n dm + \|h - h_n\|_1)^2} \right\|_1 \\
&= \left\| A - \frac{(h - h_n)(\int h_n^* dm - \|h^* - h_n^*\|_1) - h_n \|h^* - h_n^*\|_1}{(\int h_n dm + \|h - h_n\|_1)^2} \right. \\
&\quad \left. + \frac{h_n \int h_n^* dm (2 \int h_n dm \|h - h_n\|_1 + \|h - h_n\|_1^2)}{(\int h_n dm + \|h - h_n\|_1)^2 (\int h_n)^2} \right\|_1.
\end{aligned}$$

This allows us to bound the  $L^1$  error of the normalised linear response by

$$\begin{aligned}
(5.1) \quad & \frac{\|h^* - h_n^*\|_1}{\|h_n\|_1} + \frac{(\|h_n^*\|_1 + \|h^* - h_n^*\|_1) \|h - h_n\|_1}{(\|h_n\|_1 - \|h - h_n\|_1) \|h_n\|_1} \\
& + \frac{\|h - h_n\|_1 (\|h_n^*\|_1 - \|h^* - h_n^*\|_1)}{(\|h_n\|_1 + \|h - h_n\|_1)^2} + \frac{\|h_n\|_1 \|h^* - h_n^*\|_1}{(\|h_n\|_1 + \|h - h_n\|_1)^2} \\
& + \frac{2 \|h_n^*\|_1 \|h - h_n\|_1}{(\|h_n\|_1 + \|h - h_n\|_1)^2} + \frac{\|h_n^*\|_1 \|h - h_n\|_1^2}{(\|h_n\|_1 + \|h - h_n\|_1)^2 \|h_n\|_1}
\end{aligned}$$

## 6. PROOF OF THEOREM 2.1

In this section we give a proof of the main result in the paper, i.e., that we can approximate as well as we want the linear response.

*Proof of Theorem 2.1.* Using (2.16) we have

$$\begin{aligned}
(6.1) \quad & \|h^* - h_n^*\|_1 \leq \|F_0 \hat{h}^* - F_0^{\text{app}} \hat{h}_n^*\|_1 + \|Q \hat{h} - Q^{\text{app}} \hat{h}_n\|_1 \\
& \leq \|F_0 \hat{h}^* - F_0 \hat{h}_n^*\|_1 + \|F_0 \hat{h}_n^* - F_0^{\text{app}} \hat{h}_n^*\|_1 \\
& \quad + \|Q \hat{h} - Q \hat{h}_n\|_1 + \|Q \hat{h}_n - Q^{\text{app}} \hat{h}_n\|_1 \\
& := (I) + (II) + (III) + (IV).
\end{aligned}$$

By (2.6), we get

$$\begin{aligned}
(6.2) \quad & (I) \leq \|\hat{h}^* - \hat{h}_n^*\|_1 + \sum_{\omega \in \Omega} \int_{\Delta^c} \left| (\hat{h}^* \circ g_\omega - \hat{h}_n^* \circ g_\omega) g'_\omega \right| dx \\
& \leq \|\hat{h}^* - \hat{h}_n^*\|_1 + \|\hat{h}^* - \hat{h}_n^*\|_{C^0} \cdot \sum_{\omega} \|g'_\omega\|_{\mathcal{B}} \int_{\Delta^c} x^{-\gamma} dx.
\end{aligned}$$

Using (2.6) again, we have

$$(6.3) \quad \begin{aligned} (II) &\leq \sum_{|\omega| > N^*} \int_{\Delta^c} \left| \hat{h}_\eta^* \circ g_\omega g'_\omega \right| dx \leq \|\hat{h}_\eta^*\|_{C^0} \cdot \sum_{|\omega| > N^*} \|g'_\omega\|_{\mathcal{B}} \int_{\Delta^c} x^{-\gamma} dx \\ &\leq \frac{1}{2^{1-\gamma}(1-\gamma)} \cdot \|\hat{h}_\eta^*\|_{C^0} \cdot \sum_{|\omega| > N^*} \|g'_\omega\|_{\mathcal{B}}. \end{aligned}$$

Note that by (2.4), one can choose  $N^*$  large enough so that (II) is sufficiently small. Using (2.9), we have

$$(III) \leq \sum_{\omega \in \Omega} \int_{\Delta^c} \left| (\hat{h}' \circ g_\omega - \hat{h}'_n \circ g_\omega) \cdot a_\omega g'_\omega \right| dx + \sum_{\omega \in \Omega} \int_{\Delta^c} \left| (\hat{h} \circ g_\omega - \hat{h}_n \circ g_\omega) \cdot b_\omega \right| dx.$$

Now using (2.3), (2.5), and the change of variables  $y_\omega = g_\omega(x)$  we get

$$(6.4) \quad \begin{aligned} (III) &\leq \sup_{\omega} |a_\omega| \sum_{\omega \in \Omega} \int_{[\omega]_0} \left| \hat{h}'(y_\omega) - \hat{h}'_n(y_\omega) \right| dy_\omega + \|\hat{h} - \hat{h}_n\|_{C^0} \cdot \sum_{\omega \in \Omega} \|b_\omega\|_{\mathcal{B}} \int_{\Delta^c} x^{-\gamma} dx \\ &= \sup_{\omega} |a_\omega| \cdot \|\hat{h}' - \hat{h}'_n\|_1 + \frac{1}{2^{1-\gamma}(1-\gamma)} \cdot \sum_{\omega \in \Omega} \|b_\omega\|_{\mathcal{B}} \cdot \|\hat{h} - \hat{h}_n\|_{C^0} \\ &\leq \max \left\{ \sup_{\omega} |a_\omega|, \frac{1}{2^{1-\gamma}(1-\gamma)} \cdot \sum_{\omega \in \Omega} \|b_\omega\|_{\mathcal{B}} \right\} \cdot \|\hat{h} - \hat{h}_n\|_{C^1}. \end{aligned}$$

Finally, using (2.9) again, we have

$$(IV) \leq \sum_{|\omega| > N^*} \int_{\Delta^c} \left| \hat{h}'_n \circ g_\omega \cdot a_\omega g'_\omega \right| dx + \sum_{|\omega| > N^*} \int_{\Delta^c} \left| \hat{h}_n \circ g_\omega \cdot b_\omega \right| dx.$$


Using (2.3) and (2.4) in the first integral, and using (2.5) in the second integral, we choose  $N^*$  large enough and get

$$(6.5) \quad (IV) \leq \frac{1}{2^{1-\gamma}(1-\gamma)} \left[ \sup_{\omega} |a_\omega| \cdot \|\hat{h}'_n\|_{C^0} \cdot \sum_{|\omega| > N^*} \|g'_\omega\|_{\mathcal{B}} + \|\hat{h}_n\|_{C^0} \cdot \sum_{|\omega| > N^*} \|b_\omega\|_{\mathcal{B}} \right].$$

Choosing  $l^*$  in 4.1 to make  $\sum_{n=l^*+1} \|\hat{L}^n W\|_{C^1}$  small enough, followed by  $k$  and  $\eta$  to make (6.2) and (6.4) small enough, then choosing  $N^*$  in (6.3) and (6.5) so  $(I) + (II) + (III) + (IV) \leq \tau$  completing the proof.  $\square$

## 7. APPLICATION TO AN EXAMPLE

In this section we will apply our algorithm to a classical example of maps with an indifferent fixed point, strictly related to Pomeau-Manneville maps, the Liverani-Saussol-Vaienti map. The behaviour of this map is determined by the exponent  $\alpha$ ; if  $\alpha \in (0, 1)$  it is a non-uniformly expanding map with an absolutely continuous invariant probability measure; if  $\alpha \geq 1$  there is an absolutely continuous invariant infinite measure.

FIGURE 1. The inducing scheme for  $T$  in (7.1).

**7.1. Definition of the map and the induced map.** The equation of the map is

$$(7.1) \quad T(x) = \begin{cases} x(1 + 2^\alpha x^\alpha) & \text{if } x \in [0, \frac{1}{2}] \\ 2x - 1 & \text{if } x \in (\frac{1}{2}, 1] \end{cases} .$$

**Numerical assumption 7.1.** We fix  $\alpha = 0.125$  in our example. This is the value corresponding to  $\epsilon = 0$  in the previous section.

We construct the inducing scheme as in Subsection 2.1. Let  $x'_0 = 1$ ,  $x'_1 = \frac{3}{4}$ , and

$$x'_n = g_\omega\left(\frac{1}{2}\right) \text{ for } n \geq 2.$$

Letting  $\omega = 10^n$  and  $g_\omega = g_1 \circ g_0^n$ , then cylinder set  $[\omega]$  is given by  $g_\omega([0.5, 1]) = (x'_n, x'_{n-1}]$ .

Then  $\hat{T} : \Delta \rightarrow \Delta$  is a piecewise smooth and onto map with countable number of branches and it satisfies all the assumptions of subsection 2.1. See Figure 1 for a pictorial representation of the above inducing scheme.

**7.1.1. Numerical remark: the Shooting Method.** To approximate rigorously the operators in this paper we need a rigorous way to approximate long orbits given a coding, i.e., we need to be able to compute

$$x = g_\omega(y) = g_1 \circ g_0^{n-1}(y)$$


i.e. we need to be able to compute  $x \in [0.5, 1]$  such that

$$T^{n+1}(x) = y, \quad T^i(x) \in [0, 0.5]$$

for  $i \in \{1, \dots, n\}$ .

To solve this problem efficiently and obtain tight bounds on  $x$  is tricky taking preimages sequentially leads to propagation of errors and the computed interval ends up being not usable.

The main idea is to substitute the equation above with the following system of equations (this technique is called the Shooting Method, and we were introduced to


 FIGURE 2. The approximation map with  $k = 200$ 

it by W. Tucker)

$$\begin{cases} T(x_1) - x_2 = 0 & x_1 \in [0.5, 1] \\ T(x_2) - x_3 = 0 & x_2 \in [0, 0.5] \\ T(x_3) - x_4 = 0 & x_3 \in [0, 0.5] \\ \vdots \\ T(x_n) - y = 0 & x_n \in [0, 0.5]. \end{cases}$$

We will use the rigorous Newton method [42], to simultaneously enclose  $x_1, \dots, x_n$ . This way we are solving a unique system of equations instead of propagating backwards the error through solving equations with a “fat” variable. Given a function  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  and a vector of intervals  $\hat{x} = (\hat{x}_1, \dots, \hat{x}_n)$  the rigorous Newton step is given by

$$N(\hat{x}) = \hat{x} \cap \text{mid}(\hat{x}) - D\phi(\hat{x})^{-1}\phi(\text{mid}(\hat{x})),$$

where the intersection between interval vectors is meant componentwise and  $\text{mid}$  is a function that sends a vector of intervals to the vector of their midpoints [42].

In our specific case, the shooting method is numerically well behaved: denoting by  $\phi(x_1, \dots, x_n) = (T(x_1) - x_2, \dots, T(x_n) - y)^T$ , the Jacobian  $D\phi$  is given by a bidiagonal matrix, whose  $i$ -th diagonal entry is  $T'(x_i)$  and the superdiagonal entries are constant and equal to  $-1$ . In particular, this guarantees us that the Jacobian is invertible, since its eigenvalues correspond to the diagonal elements and these are bounded away from 0. Moreover a bidiagonal system is solved in time  $O(n)$  by backsubstitution, with small numerical error, and these assumptions guarantee that the interval Newton method converges.

This allows us to compute tight enclosure of  $g_\omega(y)$ ,  $g'_\omega(y)$ , which allows us to compute discretizations of the transfer operator.

**7.2. Computing the error when taking a finite number of branches.** Since we cannot calculate values for maps with infinitely many branches on the computer we use an approximating map as described in subsection 3.1, this is depicted in figure 2 for  $\alpha = 0.125$ . To calculate bounds on the  $C^1$  distance between the systems these maps define we use lemma 3.1 and find  $D$  and  $D_0$  for LSV maps. Estimating these bounds efficiently is delicate since it involves estimating the sum (and the tail) of converging series whose general term is going to zero slowly. The estimates in literature [5, 32] give rise to values that are impractical for our computations;

as an example, the value of the constant  $C_8$  in [32] computed according to their proof is of the order of  $10^{269}$ , which makes its use in our computations unfeasible, therefore some work is needed to give sharper bounds for the constants. Since these estimates are quite technical and need the introduction of specific notations, we separate them in the appendix not to hinder the flow of the sections.

In section 8.2 we bound  $D_0 \leq 0.2513$  and  $D \leq 1.286$ , so

$$\left\| (\hat{L} - \hat{L}_{\delta_k}) \right\|_{C^1} \leq (D + D_0 D + 2) \delta_k \leq 3.609 \delta_k$$

can be made as small as needed by increasing  $k$ .

Choosing  $k = 200$  gives  $\left\| \hat{L} - \hat{L}_{\delta_k} \right\|_{C^1} \leq 7.743 \cdot 10^{-12}$ .

7.2.1. *Bounding  $\left\| \hat{h}_{\delta_k} - \hat{h} \right\|_{C^1}$ .* In section 8.2.1 we prove the following Lasota-Yorke inequality

$$\left\| \hat{L}_{\delta_k}^n f \right\|_{C^1} \leq 2.491 \cdot (0.5)^n \|f\|_{C^1} + 6.206 \|f\|_{\infty}$$

The Lasota-Yorke inequality implies that

$$\left\| \hat{L}_{\delta_k}^n f \right\|_{C^1} \leq 7.452,$$

which together with Lemma 3.3, and the fact proved in section 7.3.5 by using the methods from [21] that

$$\left\| \hat{L}_{\delta_k}^4 |U^0| \right\|_{C^1} \leq 0.1557$$

allowing us to bound

$$(7.2) \quad \left\| \hat{h} - \hat{h}_{\delta_k} \right\|_{C^1} \leq \frac{4 \cdot 7.452}{1 - 0.1557} \left\| (\hat{L}_{\delta_k} - \hat{L}) \hat{h} \right\|_{C^1} \leq 2.734 \cdot 10^{-10} \left\| \hat{h} \right\|_{C^1}$$

Observing that  $\left\| \hat{h} \right\|_{C^1} \leq \left\| \hat{h}_{\delta_k} \right\|_{C^1} + \left\| \hat{h} - \hat{h}_{\delta_k} \right\|_{C^1}$  and a bound on  $\left\| \hat{h}_{\delta_k} \right\|_{C^1}$  in section 8.2.1 gives us a final error of  $2.113 \cdot 10^{-9}$ .

7.2.2. *Computing the discretization error.* The truncated operator  $\hat{L}_{\delta_k}$  satisfies the following Lasota-Yorke like inequalities<sup>8</sup>

$$\begin{aligned} \left\| (\hat{L}_{\delta_k}^n f)' \right\|_1 &\leq \lambda^n \|f'\|_1 + 1.785 \|f\|_1 \\ \left\| (\hat{L}_{\delta_k}^n f)'' \right\|_1 &\leq \lambda^n \|f''\|_1 + 0.3076 \|f'\|_1 + 6.57 \|f\|_1 \\ \left\| (\hat{L}_{\delta_k}^n f)''' \right\|_1 &\leq \lambda^{3n} \|f'''\|_1 + 0.145 \|f''\|_1 + 1.98 \|f'\|_1 + 36.96 \|f\|_1 \\ \left\| (\hat{L}_{\delta_k}^n f)^{(4)} \right\|_1 &\leq \lambda^{4n} \|f^{(4)}\|_1 + 0.057 \|f^{(3)}\|_1 + 1.49 \|f''\|_1 + 16.97 \|f'\|_1 + 559.4 \|f\|_1 \\ \left\| (\hat{L}_{\delta_k}^n f)^{(5)} \right\|_1 &\leq \lambda^{5n} \|f^{(5)}\|_1 + 0.0199 \|f^{(4)}\|_1 + 0.85 \|f^{(3)}\|_1 \\ &\quad + 17.57 \|f''\|_1 + 794.59 \|f'\|_1 + 10086 \|f\|_1 \\ \left\| (\hat{L}_{\delta_k}^n f)^{(6)} \right\|_1 &\leq \lambda^{6n} \|f^{(6)}\|_1 + 0.0066 \|f^{(5)}\|_1 + 0.41 \|f^{(4)}\|_1 \\ &\quad + 13.33 \|f^{(3)}\|_1 + 895 \|f''\|_1 + 24840.2 \|f'\|_1 + 684431 \|f\|_1. \end{aligned}$$

Since we know  $\|h_{\delta_k}\|_1 = 1$  for  $f$  a probability density, we can use these to get a bound on  $\left\| h_{\delta_k}^{(6)} \right\|_1$  which is calculated to be  $7.953 \cdot 10^5$ . Denoting by  $\hat{L}_n = \pi_n \hat{L}_{\delta_k} \pi_n$

<sup>8</sup>it is straightforward to see that these inequalities imply Lasota-Yorke inequalities on  $W^{k,1}$  with weak norm  $W^{k-1,1}$ .




FIGURE 3. The invariant density of the induced map as calculated according to section 3.1.

the discretized operator on the base of Chebyshev polynomials of the first kind of degree up to  $n$ , the same Lasota-Yorke inequalities allow us to compute

$$\left\| \hat{L}_{\delta_k} - \hat{L}_n \right\|_{C^1} \leq 3.297 \cdot 10^{-11} .$$

This, together with the computed bounds on the  $C^1$  mixing rate in table 1 gives

TABLE 1. Calculated contraction rates of our discretised operators.

$\mathbf{k}$	$\left\  \hat{L}_n^k _{U^0} \right\ _{C^1}$	$\left\  \hat{L}_u^k _{U^0} \right\ _1$
1	3.674	1
2	1.254	1
3	0.4237	1
4	0.1427	1
5	0.04799	1
6	0.01613	1
7	0.005421	1
8	0.001821	1
9	0.0006119	1
10	0.0002056	0.09782
11	0.0007551	0.02349

us an error of

$$\left\| \hat{h} - \hat{h}_n \right\|_{C^1} \leq 3.833 \cdot 10^{-9} ;$$

in figure 3 a plot of the approximated density is presented.

7.2.3. *Numerical remark: automated Lasota-Yorke inequalities.* We detail a way to automatically calculate Lasota-Yorke type inequalities for transfer operators in  $W^{k,1}$ . Following [11] let

$$L_k f = \sum_{y \in T^{-1}(x)} \frac{f(y)}{|T'(y)|^k}.$$

From this follows

$$(7.3) \quad (L_k f)' = L_{k+1} f' + k L_k (fD),$$




FIGURE 4. The linear response of the induced map as calculated according to Section 4.

where  $D = (1/T'(x))'$  is the distortion. We use the formula above to compute symbolical expressions for the derivatives  $(L_1 f)^{(l)}$ .

We use Interval Arithmetic and higher order Automatic Differentiation [42] (as implemented in TaylorSeries.jl) to compute bounds for

$$\left\| (1/T')^{(l)} \right\|_{\infty}.$$

This allows us to bound the coefficients of the Lasota-Yorke inequalities.

**7.3. Approximating the linear response for the induced map.** We approximate the linear response of our induced map using (4.1) which uses the Ulam approximation of  $\hat{L}_u = \Pi_u \hat{L}_{\delta_k} \Pi_u$ , where  $\Pi_u$  is the Ulam discretisation with partition size  $\eta = 9.537 \cdot 10^{-7}$ . We get our error from lemma 4.2, which gives us four terms that need to be bound, each of which is done in the appendix, 7.3.2, 7.3.3, 7.3.4 and 7.3.7 for  $l^* = 988$  and  $k = 200$ :

- (1)  $\sum_{n=0}^{l^*} \sum_{i=0}^n \left\| (\hat{L}_u - \hat{L}_{\delta_k}) \hat{L}_u^{n-i} W_u \right\|_1 \leq 0.0007662$
- (2)  $\sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\| (\hat{L}_{\delta_k} - \hat{L}) \hat{L}_u^{n-i} W_u \right\|_1 \leq 9.616 \cdot 10^{-8}$
- (3)  $l^* \|W_u - W\|_1 \leq 3.055 \cdot 10^{-7}$
- (4)  $\sum_{n=0}^{l^*} \left\| \hat{L}^n W \right\|_1 \leq 2 \cdot 10^{-289}$

this gives us a total error  $\left\| \hat{h}^* - \hat{h}_{\eta}^* \right\|_1 \leq 0.0007666$

**7.3.1. The contraction rates of  $\hat{L}_{\delta_k}$  in the BV norm.** In order to bound  $\left\| \hat{L}_{\delta_k}^n |_{U^0} \right\|_{BV}$  we use lemma 7.13 from [4] to bound

$$\left\| (\hat{L}_{\delta_k}^n - \hat{L}_u^n) f \right\|_1 \leq A_n \|f\|_{BV} + B_n \|f\|_1$$

from which we can use  $C_{u,n}$  to bound  $\left\| \hat{L}_{\delta_k}^n |_{U^0} \right\|_1 \leq A \|f\|_{BV} + (B + C_{u,n}) \|f\|_1$  and the Lasota-Yorke inequality (1) from section 8.2.1, and use the small matrix method from [20]. We have

$$\begin{pmatrix} \left\| \hat{L}_{\delta_k}^n f \right\|_{BV} \\ \left\| \hat{L}_{\delta_k}^n f \right\|_1 \end{pmatrix} \leq \begin{pmatrix} \lambda^n & M \\ A_n & B_n + C_{u,n} \end{pmatrix} \begin{pmatrix} \|f\|_{BV} \\ \|f\|_1 \end{pmatrix}.$$

We take  $C_{u,11} \leq 0.02349$  from table 1, together with the calculation  $\left\| (\hat{L}_{\delta_k}^{11} - \hat{L}_u^{11})f \right\|_1 \leq 0.002927 \|f\|_{BV} + 0.0182 \|f\|_1$ , which gives the largest eigenvalue of the small matrix  $\rho = 0.09785$ .

7.3.2. *Bounding Item (1)*. We can bound  $\sum_{n=0}^{l^*} \sum_{i=0}^n \left\| (\hat{L}_u - \hat{L}_{\delta_k}) \hat{L}_u^{n-i} W_u \right\|_1$  by theorem 7.13 of [4], from (1) we have

$$\left\| \hat{L}_{\delta_k}^n f \right\|_{BV} \leq (0.5)^n \|f\|_{BV} + 2(0.2513 + 1) \|f\|_1$$

We refer to [20] for the proof of

- $\|(\Pi_u - I)f\|_1 \leq \eta \text{Var}(f) \leq \eta \|f\|_{BV}$ ,
- $\|\Pi_u\|_1 \leq 1$ ,
- $\left\| \hat{L}_{\delta_k} \right\|_1 \leq 1$ .

In the theorem the value of  $C_0 = 1$ , so we have all of the values we need for the theorem's  $N = 1$  case. The theorem then gives

$$\sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\| (\hat{L}_u - \hat{L}_{\delta_k}) \hat{L}_u^{n-i} W_u \right\|_1 \leq \eta \frac{3}{2} \sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\| \hat{L}_u^{n-i} W_u \right\|_{BV} + \eta \frac{5}{2} M \sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\| \hat{L}_u^{n-i} W_u \right\|_1.$$

We can calculate  $\left\| \hat{L}_u^{n-i} W_u \right\|_1$  and  $\left\| \hat{L}_u^{n-i} W_u \right\|_{BV}$  explicitly by using validated numerical methods, since  $W_u$  explicitly represented on the computer, so we can compute an enclosure of  $\hat{L}_u W_u$  by rigorous matrix multiplication; for a function  $f$  in the Ulam basis with coefficients  $v_i$  we have explicit functions that allow us to compute the  $L_1$ ,  $BV$  and  $L^\infty$  norms, i.e.:

$$\|f\|_{L^1} = \eta \sum |v_i|, \quad \text{Var}(f) = \sum |v_{i+1} - v_i|, \quad \|f\|_\infty = \max_i |v_i|.$$

We compute a bound

$$\sum_{n=0}^{l^*} \sum_{i=0}^n \left\| (\hat{L}_u - \hat{L}_{\delta_k}) \hat{L}_u^{n-i} W_u \right\|_1 \leq 0.0007662.$$

7.3.3. *Bounding Item (2)*. To bound  $\sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\| (\hat{L}_{\delta_k} - \hat{L}) \hat{L}_u^{n-i} W_u \right\|_1$  we observe that as in lemma 3.1

$$\begin{aligned} \|(L - L_{\delta_k})f\|_1 &= \left\| \sum_{|\omega| > N^*} f \circ g_\omega |g'_\omega| + f(\delta_k(2x-1) + \frac{1}{2}) 2\delta_k \right\|_1 \\ &\leq \sum_{|\omega| > N^*} \|f \circ g_\omega |g'_\omega|\|_1 + \left\| f(\delta_k(2x-1) + \frac{1}{2}) 2\delta_k \right\|_1 \\ &\leq \|f\|_{BV} \left( \sum_{|\omega| > N^*} |g'_\omega| + 2\delta_k \right), \end{aligned}$$

therefore

$$\sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\| (\hat{L}_{\delta_k} - \hat{L}) \hat{L}_u^{n-i} W_u \right\|_1 \leq \left\| (\hat{L}_{\delta_k} - \hat{L}) \right\|_{BV \rightarrow L^1} \sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\| \hat{L}_u^{n-i} W_u \right\|_{BV}$$

As in the estimate for item (1) we can compute  $\left\|\hat{L}_u^{n-i}W_u\right\|_1$  and  $\left\|\hat{L}_u^{n-i}W_u\right\|_{BV}$  explicitly, which gives us

$$\sum_{n=1}^{l^*} \sum_{i=0}^{n-1} \left\|(\hat{L}_{\delta_k} - \hat{L})\hat{L}_u^{n-i}W_u\right\|_1 \leq 9.616 \cdot 10^{-8} \quad .$$

7.3.4. *Bounding Item (3).* We bound  $\|W_u - W\|_1$  by the following

$$\|(W_u - W)\|_1 \leq \|(W_u - W_{u_\infty})\|_1 + \|(W_{u_\infty} - W)\|_1$$

where

$$W_{u_\infty} = \sum_{\omega} \frac{1}{\eta} \int_{g_\omega(I_i)} A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta) dx$$

for  $x \in I_i$  and  $\zeta \in g_\omega(I_i)$ <sup>9</sup> and the sum is over all  $\omega$ .

$$\begin{aligned} & \|W - W_{u_\infty}\|_1 \\ &= \left\| \hat{L}[A_0 \hat{h}' + B_0 \hat{h}](x) - \sum_{\omega} \frac{1}{|I_i|} \int_{g_\omega(I_i)} A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta) dx \right\|_{L^1} \\ &\leq \int_{I_i} \sum_{\omega} \left| [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}] \circ g_\omega(y) g'_\omega(y) - \frac{1}{|I_i|} \int_{g_\omega(I_i)} A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta) dx \right| dy \\ &= \sum_{\omega} \int_{g_\omega(I_i)} \left| [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}](z) - \frac{1}{g'_\omega(\hat{T}(z))} \frac{1}{|I_i|} \int_{g_\omega(I_i)} A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta) dx \right| dz \end{aligned}$$

where we used the change of variables  $z = g_\omega(y)$ . The expression above is then equal to

$$\begin{aligned} & \sum_{\omega} \int_{g_\omega(I_i)} \left| \int_{g_\omega(I_i)} \frac{1}{|g_\omega(I_i)|} [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}](z) - \hat{T}'(z) \frac{A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta)}{|I_i|} dx \right| dz \\ &\leq \sum_{\omega} \int_{g_\omega(I_i)} \int_{g_\omega(I_i)} \left| \frac{1}{|g_\omega(I_i)|} [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}](z) - \hat{T}'(z) \frac{A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta)}{|I_i|} \right| dx dz \\ &= \sum_{\omega} \sum_{* \in \{+, -\}} \int_{g_\omega(I_i)} \left| \int_{g_\omega(I_i)} 1_* \frac{1}{|g_\omega(I_i)|} [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}](z) - \hat{T}'(z) \frac{A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta)}{|I_i|} dz \right| dx \end{aligned}$$

Here  $1_+$  is the indicator function on the set where

$$\frac{1}{|g_\omega(I_i)|} [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}](z) - \hat{T}'(z) \frac{A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta)}{|I_i|}$$

is positive, and  $1_-$  is the set on which it is negative.

In the following use  $1_{+,+}$  to be the indicator function for the set where the above function is positive and

$$\int_{g_\omega(I_i)} 1_* \frac{1}{|g_\omega(I_i)|} [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}](z) dz - A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta)$$

<sup>9</sup>It should be noted that finding the integral of  $A_0 \hat{h}'_n$  and  $B_0 \hat{h}_n$  is not easy so in our calculations instead of a true Ulam approximation where  $\zeta$  corresponds to the value that gives the integral we simply use the midpoint; this error is taken into account explicitly and depends on the regularity estimates we have on  $h_\eta$ .

is positive,  $1_{+,-}$  where they are positive and negative,  $1_{-,+}$  where they are negative and positive and  $1_{-,-}$  to be the indicator of the set on which they are both negative. Then continuing our inequality we have that the expression above is equal to

$$\begin{aligned}
& \sum_{\omega} \sum_{*,* \in \{+,-\}^2} \left| \int_{g_{\omega}(I_i)} \int_{g_{\omega}(I_i)} 1_{*,*} \frac{1}{|g_{\omega}(I_i)|} [A_0 \cdot \hat{h}' + B_0 \cdot \hat{h}](z) dz - A_0(x) \cdot \hat{h}'_n(\zeta) + B_0(x) \hat{h}_n(\zeta) dx \right| \\
&= \sum_{\omega} \sum_{*,* \in \{+,-\}^2} \left| \int_{g_{\omega}(I_i)} 1_{*,*} A_0(z) \cdot (\hat{h}'(z) - \hat{h}'_n(\zeta)) dz + \int_{g_{\omega}(I_i)} 1_{*,*} B_0(z) \cdot (\hat{h}(z) - \hat{h}_n(\zeta)) dz \right| \\
&\leq \sum_{\omega} \sum_{*,* \in \{+,-\}^2} \left( |g_{\omega}(I_i)| \|\hat{h}''\|_{\infty} + \|\hat{h} - \hat{h}_{\eta}\|_{C^1} \right) \left| \int_{g_{\omega}(I_i)} 1_{*,*} A_0(z) dz \right| \\
&\quad + (|g_{\omega}(I_i)| \|\hat{h}'\|_{\infty} + \|\hat{h} - \hat{h}_{\eta}\|_{C^1}) \left| \int_{g_{\omega}(I_i)} 1_{*,*} B_0(z) dz \right| \\
&\leq |I_i| \left( \frac{\|\hat{h}''\|_{\infty} \|A_0\|_1}{2} + \frac{\|\hat{h}'\|_{\infty} \|B_0\|_1}{2} \right) + \|\hat{h} - \hat{h}_{\eta}\|_{C^1} (\|A_0\|_1 + \|B_0\|_1)
\end{aligned}$$

We can calculate this using a computer to for  $\eta = 9.537 \cdot 10^{-7}$  to get

$$\|(W - W_{u_{\infty}})\|_1 \leq 3.046 \cdot 10^{-7}$$

We need now to estimate this for  $b$ , the linear branch of  $\hat{T}_{\delta_k}$ :

$$\begin{aligned}
& \sum_{I_i} \left\| \frac{1_{I_i}(x)}{|I_i|} \int_{b^{-1}(I_i)} A_0(y) \hat{h}'_n(\zeta) dy + B_0(y) \hat{h}_n(\zeta) dy - \frac{1_{I_i}(x)}{|I_i|} \sum_{\omega > k} \int_{g_{\omega}(I_i)} A_0(y) \hat{h}'_n(\zeta) + B_0(y) \hat{h}_n(\zeta) dy \right\|_1 \\
&\leq \sum_{I_i} \int_{b^{-1}(I_i)} |A_0(y)| \|\hat{h}'_n\|_{\infty} + B_0(y) \|\hat{h}_n\|_{\infty} |dy + \sum_{\omega > k} \int_{g_{\omega}(I_i)} |A_0(y)| \|\hat{h}'_n\|_{\infty} + B_0(y) \|\hat{h}_n\|_{\infty} |dy \\
&= \int_{0.5}^{g_k(1)} |A_0(y)| \|\hat{h}'_n\|_{\infty} + B_0(y) \|\hat{h}_n\|_{\infty} |dy + \int_{0.5}^{g_k(1)} |A_0(y)| \|\hat{h}'_n\|_{\infty} + B_0(y) \|\hat{h}_n\|_{\infty} |dy
\end{aligned}$$

where we have used that both  $\{g_{\omega}(I_i)\}_{1 \leq i \leq N, \omega > k}$  and  $\{b^{-1}(I_i)\}_{1 \leq i \leq N}$  form a disjoint cover of  $(0.5, g_k(1))$ ; then we have that the expression above is equal to

$$\begin{aligned}
& 2 \int_{0.5}^{g_k(1)} |A_0(y)| \|\hat{h}'_n\|_{\infty} + B_0(y) \|\hat{h}_n\|_{\infty} |dy \\
&\leq 2 \|\hat{h}'_n\|_{\infty} (g_k(1) - 0.5) A_0(g_k(1)) + 2 \|\hat{h}_n\|_{\infty} \text{Var}(1_{(0.5, g_k(1))} \cdot A_0).
\end{aligned}$$

We can make this arbitrarily small by increasing  $k$  as much as is needed. We calculate  $\text{Var}(1_{(0.5, g_k(1))} \cdot A_0) = \|1_{(0.5, g_k(1))} \cdot B_0\|_1$  and according to section 7.3.8 and  $A_0(g_k(1))$  according to section 9.

Taking  $k = 200$  gives us  $\|W_u - W_{u_{\infty}}\|_1 \leq 9.007 \cdot 10^{-10}$ . This together with the first bound gives  $l^* \|W_u - W\|_1 \leq 3.055 \cdot 10^{-7}$ .

7.3.5. *The contraction rates of  $\hat{L}_{\delta_k}$  in the  $C^1$  norm.* In order to bound  $\|\hat{L}_{\delta_k}^n|_{U^0}\|_{C^1}$  we use lemma 7.13 from [4] to bound

$$\|(\hat{L}_{\delta_k}^m - \hat{L}_n^m)f\|_{C^1} \leq A_m \|f\|_{C^2} + B_m \|f\|_{C^1}$$

from which we can use  $C_{c,m}$  to bound  $\left\| \hat{L}_{\delta_k}^m |_{U^0} \right\|_{C^1} \leq A \|f\|_{C^2} + (B + C_{c,m}) \|f\|_{C^1}$  and the Lasota-Yorke inequality (3) from section 8.2.1, and use the small matrix method from [20]. We have

$$\begin{pmatrix} \left\| \hat{L}_{\delta_k}^m f \right\|_{C^2} \\ \left\| \hat{L}_{\delta_k}^m f \right\|_{C^1} \end{pmatrix} \leq \begin{pmatrix} M\lambda^{2m} & D \\ A_n & B_n + C_{c,m} \end{pmatrix} \begin{pmatrix} \|f\|_{C^2} \\ \|f\|_{C^1} \end{pmatrix}.$$

Choosing the value of  $n$  that minimises equation (7.2), we take  $C_{c,4} \leq 0.1427$  from section 7.2.2, together with the calculation  $\left\| (\hat{L}_{\delta_k}^4 - \hat{L}_n^4) f \right\|_1 \leq 8.5 \cdot 10^{-7} \|f\|_{C^2} + 3.203 \cdot 10^{-6} \|f\|_{C^1}$ , which gives the largest eigenvalue of the small matrix  $\rho = 0.1557$ .

7.3.6. *The contraction rates of  $\hat{L}$  in the  $C^1$  norm.* We can use  $\left\| \hat{L}_{\delta_k}^n |_{U^0} \right\|_{C^1} \leq \rho_{n,C^1}$  and lemma 3.1 to get

$$\left\| \hat{L}^n |_{U^0} \right\|_{C^1} \leq \left\| \hat{L}_{\delta_k}^n |_{U^0} \right\|_{C^1} + \left\| \hat{L}_{\delta_k}^n - \hat{L}^n \right\|_{C^1} \leq nC^*(2 + D_0D + D)\delta_k + \rho_{n,C^1}.$$

These give us  $\left\| \hat{L}^n |_{U^0} \right\|_{C^1} \leq 2.991 \cdot 10^{-7}$  for  $n = 22$ .

7.3.7. *Bounding Item (4).* From subsection 7.3.6 we have bounds on  $\left\| \hat{L}^n |_{U^0} \right\|_1 \leq C_n$  and we can then write

$$\sum_{n=l^*}^{\infty} \left\| \hat{L}^n W \right\|_1 \leq l^* \frac{C_{l^*} \|W\|_1}{1 - C_{l^*}}.$$

Calculating  $\|W\|_1$  gives us

$$\begin{aligned} & \left\| \hat{L}[A_0 \hat{h}' + B_0 \hat{h}] \right\|_1 \\ & \leq \left\| [A_0 \hat{h}' + B_0 \hat{h}] \right\|_1 \\ & \leq \|A_0\|_1 \left\| \hat{h}' \right\|_{\infty} + \|B_0\|_1 \left\| \hat{h} \right\|_{\infty} \\ & \leq \max\{\|A_0\|_1, \|B_0\|_1\} \left\| \hat{h} \right\|_{C^1} \end{aligned}$$

which we calculate  $\|A_0\|_1$  and  $\|B_0\|_1$  as described in section 7.3.8, which gives a bound  $\|W\|_1 \leq 968.7$ . We have from subsection 7.3.6 that  $\left\| \hat{L}^n |_{U^0} \right\|_{C^1} \leq 2.991 \cdot 10^{-7}$ , for  $n = 22$  so we choose  $l^*$  to be a multiple of 22 which gives for  $l^* = 988$

$$\sum_{n=l^*}^{\infty} \left\| \hat{L}^n W \right\|_1 \leq 2 \cdot 10^{-289}$$

7.3.8. *Calculating  $\|A_0\|_1$  and  $\|B_0\|_1$ .* For calculating  $\left\| \hat{h}^* - \hat{h}_{\eta}^* \right\|_1$  we need bounds on  $\|A_0\|_1$  and  $\|B_0\|_1$ , as used in subsections 7.3.4 and 7.3.7. We have a method to calculate the values of  $A_0$  and  $B_0$  from section 9, since  $B_0 = A_0'$  we can calculate the integral of  $\int_{[a,b]} B_0(x) dx = A_0(b) - A_0(a)$ . In order to calculate the integral of  $A_0$  we approximate it by taking  $k=10$  evenly spaced values in each partition element  $I_i$ , we then take  $\frac{|I_j|}{k} \sum_{j=1}^k A_0(x_j)$  as the value of the integral on  $I_i$ . This




FIGURE 5. The invariant density of the LSV map for  $\alpha = 0.125$  calculated according to section 2.1 with  $L^1$  error of  $9.047 \cdot 10^{-9}$ .

has an  $L^1$  error of  $\frac{|I_i|}{k} \text{Var}(A_0)$ . Taking our approximation of the integral of  $A_0$  and adding  $\frac{|I_i|}{k} \text{Var}(A_0)$  gives an upper bound of  $\|A_0\|_1$ .

**7.4. Pulling back to the original map.** To get the invariant density and linear response for the full map we must pull them back to the unit interval with  $F$  and  $Q$  from subsection 2.1. The invariant density is fairly straight forward to calculate and find the error. We want a bound on  $\|h - F_0^{app} \hat{h}_n\|_1$  for which we can use a bound from (I) in the proof of theorem 2.1.

$$\|F_0 \hat{h} - F_0^{app} \hat{h}_n\|_1 \leq 2 \|\hat{h}_n - \hat{h}\|_1 + \frac{1}{2^{1-\gamma}(1-\gamma)} \|\hat{h}_n\|_{C^0} \sum_{\omega > N^*} \|g'_\omega\|_B$$

We use the bounds for  $\gamma = 0.5$   $\frac{1}{2^{1-\gamma}(1-\gamma)} \leq 1.414$ ,  $\sum_{\omega > N^*} \|g'_\omega\|_B \leq 6.51 \cdot 10^{-10}$  as calculated in section 8.1 and  $\|\hat{h}_n\|_{C^0} \leq 1.5$ , which gives the second term to be bounded by  $1.381 \cdot 10^{-9}$ . The first term we can bound by  $2 \|\hat{h}_n - \hat{h}\|_{C^1} \leq 7.666 \cdot 10^{-9}$  as calculated in section 7.2.2, giving us  $\|h - F_0^{app} \hat{h}_n\|_1 \leq 9.047 \cdot 10^{-9}$ .

As seen in theorem 2.1 pulling back the linear response requires the following bounds

$$\begin{aligned} \|h^* - h_\eta^*\|_1 &\leq \|F_0 \hat{h}^* - F_0 \hat{h}_\eta^*\|_1 + \|F_0 \hat{h}_\eta^* - F_0^{app} \hat{h}_\eta^*\|_1 \\ &\quad + \|Q \hat{h} - Q \hat{h}_n\|_1 + \|Q \hat{h}_n - Q^{app} \hat{h}_n\|_1. \end{aligned}$$

We bound these in section 7.4.1 giving

- (1)  $\|F_0 \hat{h}^* - F_0 \hat{h}_\eta^*\|_1 \leq 0.001533$
- (2)  $\|F_0 \hat{h}_\eta^* - F_0^{app} \hat{h}_\eta^*\|_1 \leq 4.603 \cdot 10^{-10}$
- (3)  $\|Q \hat{h} - Q \hat{h}_n\|_1 \leq 6.818 \cdot 10^{-6}$
- (4)  $\|Q \hat{h}_n - Q^{app} \hat{h}_n\|_1 \leq 0.006225$

This gives us  $\|h^* - h_\eta^*\|_1 \leq 0.007765$ .

7.4.1. *Bounding Items (1) and (2).* It is given in theorem 2.1 that

$$\left\| F_0 \hat{h}^* - F_0 \hat{h}_\eta^* \right\|_1 \leq 2 \left\| \hat{h}^* - \hat{h}_\eta^* \right\|_1,$$

which we have from subsection 7.3 is bounded by  $2 \cdot 0.0007666 = 0.001533$ .

We also have from theorem 2.1 that  $\left\| F_0 \hat{h}_\eta^* - F_0^{app} \hat{h}_\eta^* \right\|_1$  is bounded by

$$\left\| \hat{h}_\eta^* \right\|_{C^0} \frac{1}{2^{1-\gamma}(1-\gamma)} \sum_{\omega > N^*} \|g'_\omega\|_B.$$

We can compute explicitly  $\left\| \hat{h}_\eta^* \right\|_{C^0}$  and it is bounded by  $0.5$ ,  $\frac{1}{2^{1-\gamma}(1-\gamma)} \leq 1.414$  and for  $N^* = 1000$ ,  $\sum_{\omega > N^*} \|g'_\omega\|_B \leq 6.51 \cdot 10^{-10}$  as is shown in section 8.1. These give us the bound  $\left\| F_0 \hat{h}_\eta^* - F_0^{app} \hat{h}_\eta^* \right\|_1 \leq 4.603 \cdot 10^{-10}$ .

7.4.2. *Bounding Items (3) and (4).* We can bound  $\left\| Q\hat{h} - Q\hat{h}_n \right\|_1$  by

$$\frac{1}{2^{1-\gamma}(1-\gamma)} \sum_{\omega} \|b_\omega\|_B \left\| \hat{h} - \hat{h}_n \right\|_{C^1}$$

for which we will need a bound on  $\sum_{\omega} \|b_\omega\|_B$ . In section 8.3 we show is less than 1258; the bounds from earlier  $\frac{1}{2^{1-\gamma}(1-\gamma)} \leq 1.414$  and  $\left\| \hat{h} - \hat{h}_n \right\|_{C^1} \leq 3.833 \cdot 10^{-9}$  allow us to prove that

$$\left\| Q\hat{h} - Q\hat{h}_n \right\|_1 \leq 6.818 \cdot 10^{-6}.$$

We now bound

$$\left\| Q\hat{h}_n - Q^{app} \hat{h}_n \right\|_1 \leq \frac{1}{2^{1-\gamma}(1-\gamma)} \left[ \sup_{\omega} |a_\omega| \cdot \left\| \hat{h}'_n \right\|_{C^0} \cdot \sum_{|\omega| > N^*} \|g'_\omega\|_B + \left\| \hat{h}_n \right\|_{C^0} \sum_{\omega > N^*} \|b_\omega\|_B \right].$$

We need a bound on  $\sum_{\omega} |a_\omega|$  which we show is bounded by 7107 in section 8.4, and the bounds from computer approximations of  $\hat{h}_n$  which gives us  $\left\| \hat{h}_n \right\|_{C^0} \leq 1.5$  and  $\left\| \hat{h}'_n \right\|_{C^0} \leq 1$ .

We can use the same method from section 8.3 to calculate  $\sum_{\omega > N^*} \|b_\omega\|_B \leq 0.002931$ . All this together gives us

$$\left\| Q\hat{h}_n - Q^{app} \hat{h}_n \right\|_1 \leq 0.006225.$$

7.5. **Normalizing the density and the linear response.** In this subsection we follow the estimates in Section 5.

First of all, we compute

$$\left\| \frac{h_n}{\int h_n dm} - \frac{h}{\int h dm} \right\|_1 \leq 4.11 \cdot 10^{-7}.$$

Following through the calculations we bound (5.1), the  $L^1$  error on the normalized linear response by 0.01501.




FIGURE 6. The linear response and invariant density of the LSV map for  $\alpha = 0.125$ , calculated according to section 2.1 with  $L^1$  error for the linear response of  $0.007765$ .




FIGURE 7. This shows the normalised invariant density of the LSV map at  $\alpha = 0.125$  with the linear response of the normalised density.

## 8. APPENDIX: EFFECTIVE BOUNDS FOR [32, 5]

In this section we will use often the following notation following [32]. Let  $T_0$  be the left branch of the map  $T$ , let  $z \in [0, 1]$  and

$$z_r := T_0^{-r}(z).$$

By  $(\cdot)'$  we denote the derivative with respect to  $z$ . To simplify the lookup of constants, they are presented in table 2.

8.1. **Estimating the tail**  $\sum_{\omega > N^*} \|g'_\omega\|_B$ . For this we look at [5] lemma 5.2 which gives

$$\|g'_\omega\|_B \leq C_8 \sup_{z \in (0, 0.5]} z^\gamma (1 + nz^\alpha \alpha 2^\alpha)^{-1/\alpha-1}.$$

using calculations from section 8.3

$$\begin{aligned} & \sup_{z \in (0, 0.5]} z^\gamma (1 + nz^\alpha \alpha 2^\alpha)^{-1/\alpha-1} \\ & \leq \sup_{z \in (0, 0.5]} \frac{z^{\gamma-1-\alpha} (\alpha 2^\alpha)^{-1/\alpha-1}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \\ & \leq C_{sum} n^{-\gamma/\alpha} \end{aligned}$$

TABLE 2. Table of constants.

Label	Description	Value
$\alpha$	Parameter for the LSV map.	0.125
$\gamma$	Parameter for $\ \cdot\ _B$ .	0.5
$\delta_k$	The size of the interval different from the true induced map.	$2.146 \cdot 10^{-12}$
$\eta$	Partition size for Ulam discretisation.	$9.537 \cdot 10^{-7}$
$N^*$	The number of branches used to approximate operators $F$ and $Q$ .	1000
$l^*$	The number of iterations used to calculate $\hat{h}_\eta^*$ .	988
$\nu$	The number of Lasota-Yorke inequalities used to bound error in the Chebyshev projection.	5
$n$	The number of Chebyshev polynomials used for the Chebyshev discretisation.	1024
$C_1$		0.88
$C_2$		16.77
$C_3$		9.022
$C_4$	$\frac{1}{\alpha} + \log(C_1^{-1/\alpha})$	9.475
$C_5$	$\frac{1}{\alpha} - \frac{\log(C_1^{1/\alpha})}{\log(2)}$	1.091
$C_6$	$2^\alpha$	1.227
$C_7$	$(\alpha + 1)2^\alpha$	0.1534
$C_8$	$\alpha(\alpha + 1)2^\alpha$	2.766
$C_{10}$	A computed value from section 8.1.	50.58
$C_{11}$	$C_2 \cdot C_6 \cdot C_8$	1739
$C_{12}$	$C_2 \cdot C_6 \cdot C_8 \frac{-\log(1/C_2)+1}{\log(1/C_2)+1}$	1232
$C_{sum}$	$2^\alpha C_2^2 \cdot C_4 \cdot C_7 \cdot C_8 \cdot \frac{\alpha}{(\frac{1+\alpha-\gamma}{\gamma})^{-\gamma/\alpha+1/\alpha+1}(\alpha 2^\alpha)^{-\gamma/\alpha}}$	5.981
$D_0$	A bound on the distortion of the branches of the induced map.	0.2513
$D$	A bound on the distortion of the inverse of the branches of the induced map.	1.286

so  $\sum_{n>N^*} \|g'_\omega\|_B \leq C_8 \cdot C_{sum}[\zeta(\gamma/\alpha) - \sum_{j=1}^{N^*} j^{-\gamma/\alpha}]$ .

The constant  $C_8$  comes from [32] where it is shown to be finite, but, when we calculate  $C_8$  according to their proof we get  $C_8 = \exp(1 + (\alpha + 1)^2 2^{2\alpha} C_2^2 \frac{\pi^2}{6})$ , which is of order  $10^{269}$ .

Therefore we need a sharper bound for  $C_8$ . We start similarly

$$\begin{aligned} z'_n &= \prod_{j=1}^n \frac{1}{1 + (\alpha + 1)2^\alpha z_j^\alpha} = \exp\left(\sum_{j=1}^n -\log(1 + (\alpha + 1)2^\alpha z_j^\alpha)\right) \\ &= \exp\left(\sum_{j=1}^n -(\alpha + 1)2^\alpha z_j^\alpha + \sum_{j=1}^n [-\log(1 + (\alpha + 1)2^\alpha z_j^\alpha) + (\alpha + 1)2^\alpha z_j^\alpha]\right) \\ &\leq e \cdot (1 + nz_0^\alpha \alpha 2^\alpha)^{-(\alpha+1)/\alpha} \cdot \exp\sum_{j=1}^n [-\log(1 + (\alpha + 1)2^\alpha z_j^\alpha) + (\alpha + 1)2^\alpha z_j^\alpha] \end{aligned}$$

where in the last line we use the calculation in [32] following equation (5.7) which gives,

$$-(\alpha + 1)2^\alpha \sum_{j=1}^n z_j^\alpha \leq -\frac{\alpha + 1}{\alpha} (\log(1 + nz_0^\alpha \alpha 2^\alpha) + C)$$

where  $C$  comes from

$$\sum_{j=1}^r \frac{1}{z_0^{-\alpha} + j\alpha 2^\alpha} \geq \int_1^r \frac{z_0^\alpha}{1 + tz_0^\alpha \alpha 2^\alpha} dt - C.$$

Since the function in the integral is monotonically decreasing and  $\frac{z_0^\alpha}{1 + tz_0^\alpha \alpha 2^\alpha} \leq 1$  we can bound  $C$  by 1, which gives us the factor of  $e$ .

In the next paragraph we will use the Taylor expansion of  $-\log(1 + x)$ , however this is only convergent for  $x \in (-1, 1)$ , so first we choose a  $j^*$  large enough that  $\frac{-(\alpha+1)2^\alpha C_2}{j^*} \in (-1, 1)$ .

$$\begin{aligned} z'_n &\leq e \cdot (1 + nz_0^\alpha \alpha 2^\alpha)^{-(\alpha+1)/\alpha} \cdot \exp\sum_{j=1}^n [-\log(1 + (\alpha + 1)2^\alpha z_j^\alpha) + (\alpha + 1)2^\alpha z_j^\alpha] \\ &= e \cdot (1 + nz_0^\alpha \alpha 2^\alpha)^{-(\alpha+1)/\alpha} \cdot \exp\sum_{j=1}^{j^*-1} [-\log(1 + (\alpha + 1)2^\alpha z_j^\alpha) + (\alpha + 1)2^\alpha z_j^\alpha] \\ &\quad \cdot \exp\sum_{j=j^*}^n [-\log(1 + (\alpha + 1)2^\alpha z_j^\alpha) + (\alpha + 1)2^\alpha z_j^\alpha] \end{aligned}$$

For  $\exp(\sum_{j=j^*}^n [-\log(1 + (\alpha + 1)2^\alpha z_j^\alpha) + (\alpha + 1)2^\alpha z_j^\alpha])$  we use

$$\begin{aligned} &\exp\left(\sum_{j=j^*}^n [-\log(1 + (\alpha + 1)2^\alpha z_j^\alpha) + (\alpha + 1)2^\alpha z_j^\alpha]\right) \\ &\leq \exp\left(\sum_{j=j^*}^n [-\log(1 + (\alpha + 1)2^\alpha C_2 j^{-1}) + (\alpha + 1)2^\alpha C_2 j^{-1}]\right). \end{aligned}$$

Substituting in the Taylor expansion of  $\log(1+x)$  where  $x = (\alpha+1)2^\alpha C_2 j^{-1}$  gives

$$\begin{aligned}
& \exp\left(\sum_{j=j^*}^n \sum_{m=2}^{\infty} \frac{-(\alpha+1)2^\alpha C_2^m}{m} j^{-m}\right) \\
&= \exp\left(\sum_{m=2}^{\infty} \sum_{j=j^*}^n \frac{-(\alpha+1)2^\alpha C_2^m}{m} j^{-m}\right) \\
&= \exp\left(\sum_{m=2}^{\infty} \frac{-(\alpha+1)2^\alpha C_2^m}{m} \sum_{j=j^*}^n j^{-m}\right) \\
&\leq \exp\left(\sum_{m=2}^{\infty} \frac{-(\alpha+1)2^\alpha C_2^m}{m} \left[\zeta(m) - \sum_{j=1}^{j^*} \frac{1}{j^m}\right]\right) \\
&\leq \exp\left(\sum_{m=2}^{\infty} \frac{-(\alpha+1)2^\alpha C_2^m}{m} \left[\zeta(2) - \sum_{j=1}^{j^*} \frac{1}{j^2}\right]\right) \\
&\leq \exp\left(-\left[\zeta(2) - \sum_{j=1}^{j^*} \frac{1}{j^2}\right] \cdot \left[\log(1 + (\alpha+1)2^\alpha C_2) - (\alpha+1)2^\alpha C_2\right]\right) \\
&= \exp\left((\alpha+1)2^\alpha C_2 \left[\zeta(2) - \sum_{j=1}^{j^*} \frac{1}{j^2}\right] \cdot (1 + (\alpha+1)2^\alpha C_2)^{-\left[\zeta(2) - \sum_{j=1}^{j^*} \frac{1}{j^2}\right]}\right)
\end{aligned}$$

To get our final estimate we need to bound

$$\exp \sum_{j=1}^{j^*-1} [-\log(1 + (\alpha+1)2^\alpha z_j^\alpha) + (\alpha+1)2^\alpha z_j^\alpha],$$

Since there are a finite number of terms we can bound it from above through the use of rigorous numerical methods.

Choosing  $j^* = 1000$  gives us that  $\frac{-(\alpha+1)2^\alpha C_2}{j^*} = -0.02057 \in (-1, 1)$  and  $C_8 \geq 2.766$  is an upper bound.

This gives us that for  $N^* = 1000$ ,  $\sum_{\omega > N^*} \|g'_\omega\|_B \leq 6.51 \cdot 10^{-10}$ .

**8.2. Bounds for lemma 3.1.** We want a bound on  $D_0 = \left\| \frac{g''_\omega}{g'_\omega} \right\|_\infty$ . In [32] they have bounds for  $z_r$  where  $g'_{r+1} = 0.5z'_r$  and  $g''_{r+1} = 0.5z''_r$  so we may use the bound from [32, Lemma 5.4] which gives

$$\frac{z''_r}{z'_r} = \frac{\alpha(\alpha+1)2^\alpha z_{r+1}^{\alpha-1} z'_{r+1}}{1 + (\alpha+1)2^\alpha z_{r+1}^\alpha} + \frac{z''_{r+1}}{z'_{r+1}}$$

which we can use to get a bound on  $\frac{z''_r}{z'_r} - \frac{z''_{r+1}}{z'_{r+1}}$ . We use lemma 5.2 and 5.3 from [32] to get

$$z'_r \leq C_8 (\alpha 2^\alpha)^{-(\alpha+1)/\alpha} r^{-(\alpha+1)/\alpha} z_0^{-\alpha-1}$$

where we calculate  $C_8$  in section 8.1, and

$$z_r^{\alpha-1} = (z_r^\alpha)^{(\alpha-1)/\alpha} \leq \left( \frac{2^{1-\alpha}}{\alpha(1-\alpha)} \right)^{(\alpha-1)/\alpha} r^{-(\alpha-1)/\alpha}$$

giving us

$$z_r^{\alpha-1} z'_r \leq C_8 \frac{\alpha^{-2} 2^{1-1/\alpha-2\alpha}}{(1-\alpha)^{(\alpha-1)/\alpha}} r^{-2} z_0^{-\alpha-1}.$$

Then

$$\begin{aligned} \frac{z''_r}{z'_r} - \frac{z''_{r+1}}{z'_{r+1}} &\leq \alpha(\alpha+1) 2^\alpha C_8 \frac{\alpha^{-2} 2^{1-1/\alpha-2\alpha}}{(1-\alpha)^{(\alpha-1)/\alpha}} r^{-2} z_0^{-\alpha-1} \\ &\leq (\alpha+1) C_8 \frac{\alpha^{-1} 2^{1-1/\alpha-\alpha}}{(1-\alpha)^{(\alpha-1)/\alpha}} r^{-2} 0.5^{-\alpha-1} \end{aligned}$$

from which follows that

$$\sup_r \left\| \frac{z''_r}{z'_r} \right\|_\infty \leq \pi^2 \frac{0.5^{-\alpha-1}}{6} (\alpha+1) C_8 \frac{\alpha^{-1} 2^{1-1/\alpha-\alpha}}{(1-\alpha)^{(\alpha-1)/\alpha}}$$

which for  $\alpha = 0.125$  gives  $D_0 = 0.2513$ .

Since

$$\begin{aligned} \log\left(\frac{g'_\omega(x)}{g'_\omega(y)}\right) &= [\log(g'_\omega(\zeta))]'(x-y) \\ &= \frac{g''_\omega(\zeta)}{g'_\omega(\zeta)}(x-y) \leq \frac{g''_\omega(\zeta)}{g'_\omega(\zeta)}, \end{aligned}$$

we have that  $D \leq \exp(D_0) \leq 1.286$ .

8.2.1. *Lasota-Yorke inequalitys for  $\hat{L}_{\delta_k}$  and  $\hat{L}$ .* We use some estimates from [4];

- (1) From proposition 7.2 we have  $\text{var}(\hat{L}f) \leq \lambda \text{var}(f) + B \|f\|_1$  with  $B := \left\| \hat{T}'' / (\hat{T}')^2 \right\|_\infty$  and  $\lambda := 1 / \inf_x |D_x \hat{T}|$ ;
- (2) From proposition 7.4  $\left\| \hat{L}^n f \right\|_{C^1} \leq M \lambda^n \|f\|_{C^1} + M^2 \|f\|_\infty$  with  $M := 1 + \frac{B}{1-\lambda}$ ;
- (3) From proposition 7.6  $\left\| \hat{L}^n f \right\|_{C^2} \leq M(\lambda^2)^n \|f\|_{C^2} + D \|f\|_{C^1}$  where  $D := \max\{3 \frac{\lambda B M}{1-\lambda}, 3M \left(\frac{B}{1-\lambda}\right)^2 + MZ\} + M\lambda + M^2$ .  $Z$  being

$$\frac{1}{1-\lambda^2} \left( \left\| \hat{T}''' / (\hat{T}')^3 \right\|_\infty + \frac{3\lambda}{1-\lambda} \left\| \hat{T}'' / (\hat{T}')^2 \right\|_\infty \right).$$

By the construction of  $\hat{T}_{\delta_k}$  we know  $1 / \inf_x |D_x \hat{T}_{\delta_k}| = 1 / \inf_x |D_x \hat{T}|$ ,  $\left\| \hat{T}_{\delta_k}'' / (\hat{T}_{\delta_k}')^2 \right\|_\infty \leq \left\| \hat{T}'' / (\hat{T}')^2 \right\|_\infty$  and  $\left\| \hat{T}_{\delta_k}''' / (\hat{T}_{\delta_k}')^3 \right\|_\infty \leq \left\| \hat{T}''' / (\hat{T}')^3 \right\|_\infty$  so bounding these values for  $\hat{T}$  gives us inequalitys that are true for both.

We note that  $\left\| \hat{T}'' / (\hat{T}')^2 \right\|_\infty = \sup_\omega \|g''_\omega / (g'_\omega)\|_\infty = D_0$  which is calculated in section 8.2. We can calculate  $\left\| \hat{T}''' / (\hat{T}')^3 \right\|_\infty$  a similar way as follows,

$$\left\| \frac{T'''}{(T')^3} \right\|_\infty = \left\| \left( \frac{g'''}{(g')^4} + 3 \frac{(g'')^2}{(g')^5} (g')^3 \right) \right\|_\infty \leq \left\| \frac{g'''}{g'} \right\|_\infty + 3 \left\| \left( \frac{g''}{g'} \right)^2 \right\|_\infty$$

so we need to bound  $\left\| \frac{g'''}{g'} \right\|_\infty$  and  $\left\| \left( \frac{g''}{g'} \right)^2 \right\|_\infty$ , the second of which is  $D_0^2$  from section

8.2. In [32] it is proven that  $\left\| \frac{g'''}{g'} \right\|_\infty$  is bounded and their method gives that it is

less than

$$\sum_{r=0}^{\infty} (\alpha - 1)\alpha(\alpha + 1)2^\alpha z_{r+1}^{\alpha-2} (z'_{r+1})^2 + 3\alpha(\alpha + 1)2^\alpha z_{r+1}^{\alpha-1} z'_{r+1} \frac{z''_{r+1}}{z'_{r+1}}$$

where  $g_\omega = z_r \circ g_1$ , so  $z'_r = \frac{g'_\omega}{2}$  and  $z''_r = \frac{g''_\omega}{2}$ . We have bounds on  $\frac{z''_{r+1}}{z'_{r+1}}$  and  $z_{r+1}^{\alpha-1} z'_{r+1}$  from section 8.2, and we bound  $z_{r+1}^{\alpha-2}$  using lemma 5.2 of [32] to get  $z_{r+1}^{\alpha-2} \leq (\frac{C_2}{r+1})^{(\alpha-2)/\alpha}$ . We bound  $(z'_{r+1})^2$  using lemma 5.3 of [32] to get  $(z'_{r+1})^2 \leq C_8^2 (2^\alpha \alpha)^{-2(\alpha+1)/\alpha} z_0^{-2(\alpha+1)} (r+1)^{-2(\alpha+1)/\alpha}$  so we can bound  $\left\| \frac{g''_\omega}{g'_\omega} \right\|_\infty$  by

$$\left( z_0^{-\alpha-4} \sum_{r=0}^{\infty} (r+1)^{-3} \right) [(\alpha-1)\alpha(\alpha+1)2^\alpha C_2^{(\alpha-2)/\alpha} C_8^2 (2^\alpha \alpha)^{-2(\alpha+1)/\alpha} + 3\alpha(\alpha+1)2^\alpha C]$$

where  $C$  is the product of the values from section 8.2. Substituting in the maximizing value of  $z_0 = 0.5$  and note  $\sum_{r=0}^{\infty} (r+1)^{-3} = \zeta(3)$  to get a bound of 0.08016

This gives us a bound of  $\left\| \frac{T'''}{(T')^3} \right\|_\infty \leq 0.2696$  and we have

- $\lambda = 0.5$ ,
- $B \leq 0.2513$ ,
- $M \leq 1.503$ ,
- $Z \leq 1.365$ ,
- $D \leq 12.19$ .

These values give us the explicit bounds

- (1)  $\text{Var}(\hat{L}_{\delta_k} f) \leq 0.5 \text{Var}(f) + 0.2513 \|f\|_1$ ;
- (2)  $\left\| \hat{L}_{\delta_k}^n f \right\|_{C^1} \leq 1.503 \cdot 0.5^n \|f\|_{C^1} + 2.258 \|f\|_\infty$ ;
- (3)  $\left\| \hat{L}_{\delta_k}^n f \right\|_{C^2} \leq 1.503 \cdot 0.25^n \|f\|_{C^2} + 12.19 \|f\|_{C^1}$ .

These Lasota-Yorke inequalities give us the bounds  $C^* = M\lambda + M^2 = 3.009$  for lemma 3.3.

For a bound on  $\left\| \hat{h}_{\delta_k} \right\|_{C^1}$  and  $\left\| \hat{h}_{\delta_k} \right\|_{C^1}$  we observe that  $\left\| \hat{h}_{\delta_k} \right\|_1 = 1$ , and  $\hat{L}_{\delta_k} \hat{h}_{\delta_k} = \hat{h}_{\delta_k}$ , the inequalities above give us

- (1)  $\text{var}(\hat{h}_{\delta_k}) \leq 0.2513 \implies \left\| \hat{h}_{\delta_k} \right\|_{BV} \leq 1 + 0.2513 = 1.251$
- (2)  $\left\| \hat{h}_{\delta_k} \right\|_{C^1} \leq 2.258 \left\| \hat{h}_{\delta_k} \right\|_\infty \leq 2.258 \left\| \hat{h}_{\delta_k} \right\|_{BV} = 2.825$
- (3)  $\left\| \hat{h}_{\delta_k} \right\|_{C^2} \leq 12.19 \left\| \hat{h}_{\delta_k} \right\|_{C^1} \leq 34.45$

8.3. **Bounding**  $\sum_\omega \|\partial_\epsilon g'_\omega\|_B$ . For this we use from Lemma 5.2. [5]

$$\begin{aligned} \sum_\omega \|\partial_\epsilon g'_\omega\|_B &\leq C_5 \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} |z^\gamma \cdot z'_n| \sum_{j=1}^n z_j^\alpha \\ &\quad + C_6 \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} |z^\gamma \cdot z'_n| \sum_{j=1}^n z_j^\alpha |\log z_j| \\ &\quad + C_7 \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} |z^\gamma \cdot z'_n| \sum_{j=1}^n z_j^{\alpha-1} |\partial_\alpha z_j| \end{aligned}$$

where  $C_5 = 2^\alpha$ ,  $C_6 = (\alpha + 1)2^\alpha$  and  $C_7 = \alpha(\alpha + 1)2^\alpha$ . We now use our bound from 8.1 get  $C_8$  for  $z'_n \leq C_8(1 + rz_0^\alpha 2^\alpha)^{-\frac{\alpha+1}{\alpha}}$ . We use this to bound  $|z^\gamma \cdot z'_n|$  by  $C_8 z^\gamma (1 + nz_0^\alpha 2^\alpha)^{-\frac{\alpha+1}{\alpha}}$ .

We then use  $z_n^\alpha < \frac{C_2}{n}$  to get  $z_j^\alpha \leq C_2 j^{-1}$ ,  $\log(z_j) \leq \frac{-\log(1/C_2)+1}{\alpha} \log j$ . Using the fact that  $z_n^{\alpha-1} |\partial_\alpha z_n| = z_n^\alpha \frac{|\partial_\alpha z_n|}{z_n}$  and inequality (5.9) from [32]

$$z_n^{\alpha-1} |\partial_\alpha z_n| \leq z_n^\alpha \sum_{j=1}^n 2^\alpha z_j^\alpha (-\log(2z_j)) \leq 2^\alpha \frac{C_2^2}{n} \sum_{j=1}^n j^{-1} (-\log(2z_j))$$

We now use the following

$$-\log(2z_j) \leq C_4 \log(j)$$

To get a value on  $C_4$  we do the following,

$$\begin{aligned} \frac{-\log(2z_j)}{\log(j)} &\leq \frac{-\log(2(\frac{C_1}{j})^{1/\alpha} z_0)}{\log(j)} \leq \frac{-\log(2C_1^{1/\alpha} z_0) + \frac{1}{\alpha} \log(j)}{\log(j)} \\ &\leq \frac{1}{\alpha} - \frac{\log(2C_1^{1/\alpha} z_0)}{\log(j)} \leq \frac{1}{\alpha} - \frac{\log(2C_1^{1/\alpha} z_0)}{\log(2)} \\ &= C_4 \leq 9.475 \end{aligned}$$

where we note that  $\partial_\alpha z_j = 0$  for  $|\omega| = 1$  and so  $j = 1$  this is still a valid bound for  $z_j^{\alpha-1} |\partial_\alpha z_j|$ .

which gives us  $\sum_\omega \|b_\omega\|_B \leq (I) + (II) + (III)$  where

$$\begin{aligned} (I) &\leq C_{10} \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} z^\gamma (1 + nz^\alpha \alpha 2^\alpha)^{-1/\alpha-1} \sum_{j=1}^n (j^{-1}) \\ (II) &\leq C_{11} \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} z^\gamma (1 + nz^\alpha \alpha 2^\alpha)^{-1/\alpha-1} \sum_{j=1}^n (j^{-1} \log j) \\ (III) &\leq C_{12} \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} z^\gamma (1 + nz^\alpha \alpha 2^\alpha)^{-1/\alpha-1} \sum_{j=1}^n (j^{-1} \sum_{k=1}^j k^{-1} \log k) \end{aligned}$$

where  $C_{10} = C_5 \cdot C_8 \cdot C_2$ ,  $C_{11} = C_6 \cdot C_8 \cdot C_2 \cdot \frac{-\log(1/C_2)+1}{\alpha}$  and  $C_{12} = 2^\alpha C_4 \cdot C_7 \cdot C_8 \cdot C_2^2$ . To bound (I) we use that  $\sum_{j=1}^n j^{-1} \leq 1 + \log(n)$  to get

$$\begin{aligned} z^\gamma (1 + nz^\alpha \alpha 2^\alpha)^{-1/\alpha-1} \sum_{j=1}^n (j^{-1}) &\leq \frac{z^\gamma (1 + \log(n))}{(1 + nz^\alpha \alpha 2^\alpha)^{1/\alpha+1}} \\ &\leq \frac{z^{\gamma-\alpha(1/\alpha+1)} (\alpha 2^\alpha)^{-1/\alpha-1} (1 + \log(n))}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \leq \frac{z^{\gamma-1-\alpha} (\alpha 2^\alpha)^{-1/\alpha-1} (1 + \log(n))}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}}. \end{aligned}$$

We can use this to bound

$$\begin{aligned}
& \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} z^{\gamma} (1 + nz^{\alpha} \alpha 2^{\alpha})^{-1/\alpha-1} (1 + \log(n)) \\
& \leq \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} \frac{z^{\gamma-1-\alpha} (\alpha 2^{\alpha})^{-1/\alpha-1} (1 + \log(n))}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \\
& = \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} \frac{z^{\gamma-1-\alpha} (\alpha 2^{\alpha})^{-1/\alpha-1}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} + \frac{z^{\gamma-1-\alpha} (\alpha 2^{\alpha})^{-1/\alpha-1} \log(n)}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}}
\end{aligned}$$

In order to calculate bounds we must find the  $z \in (0, 0.5]$  that gives us the maximum value, which we do by finding the zero of the derivative of the part that depends on  $z$ ,

$$\begin{aligned}
& \partial_z \frac{z^{\gamma-1-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \\
& = \frac{\partial_z z^{\gamma-1-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} + z^{\gamma-1-\alpha} \partial_z \frac{1}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \\
& = \frac{(\gamma-1-\alpha)z^{\gamma-2-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} + z^{\gamma-1-\alpha} \frac{-(1/\alpha+1) \cdot -\alpha \cdot z^{-\alpha-1} \cdot \alpha^{-1} 2^{-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+2}} \\
& = \frac{(\gamma-1-\alpha)z^{\gamma-2-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} + \frac{(1+\alpha) \cdot z^{\gamma-2\alpha-2} \cdot \alpha^{-1} 2^{-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+2}} \\
& = \frac{z^{\gamma-\alpha-2}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \left( (\gamma-1-\alpha) + \frac{(1+\alpha)z^{-\alpha} \alpha^{-1} 2^{-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)} \right)
\end{aligned}$$

which is zero when  $-(\gamma-1-\alpha) = \frac{(1+\alpha)z^{-\alpha} \alpha^{-1} 2^{-\alpha}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)}$ , we let  $y = z^{-\alpha} \alpha^{-1} 2^{-\alpha}$  which gives

$$\begin{aligned}
& -(\gamma-1-\alpha) = (1+\alpha) \frac{y}{y+n} \\
& \implies -(\gamma-1-\alpha)n + (1+\alpha)y - \gamma y = (1+\alpha)y \\
& \implies -(\gamma-1-\alpha)n - \gamma y = 0
\end{aligned}$$

Therefore  $y = \frac{(1+\alpha-\gamma)}{\gamma} n$  and

$$z = (\alpha 2^{\alpha} \frac{(1+\alpha-\gamma)}{\gamma} n)^{-1/\alpha}.$$

We substitute this into the first sum

$$\begin{aligned}
& \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0, 0.5]} \frac{z^{\gamma-1-\alpha} (\alpha 2^{\alpha})^{-1/\alpha-1}}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \\
& = \sum_{n=1}^{\infty} \sup_{\alpha \in U} \frac{(\alpha 2^{\alpha} \frac{(1+\alpha-\gamma)}{\gamma} n)^{-\gamma/\alpha+1/\alpha+1} (\alpha 2^{\alpha})^{-1/\alpha-1}}{(\alpha 2^{\alpha} \frac{(1+\alpha-\gamma)}{\gamma} n \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}},
\end{aligned}$$

therefore

$$\begin{aligned}
& \sum_{n=1}^{\infty} \sup_{\alpha \in U} \frac{\left(\frac{(1+\alpha-\gamma)}{\gamma}\right)^{-\gamma/\alpha+1/\alpha+1} n^{-\gamma/\alpha} (\alpha 2^\alpha)^{-\gamma/\alpha}}{\left(\frac{1+\alpha-\gamma}{\gamma} + 1\right)^{1/\alpha+1}} \\
&= \sup_{\alpha \in U} \frac{\left(\frac{(1+\alpha-\gamma)}{\gamma}\right)^{-\gamma/\alpha+1/\alpha+1} (\alpha 2^\alpha)^{-\gamma/\alpha}}{\left(\frac{1+\alpha-\gamma}{\gamma} + 1\right)^{1/\alpha+1}} \sum_{n=1}^{\infty} n^{-\gamma/\alpha} \\
&= \sup_{\alpha \in U} \frac{\left(\frac{(1+\alpha-\gamma)}{\gamma}\right)^{-\gamma/\alpha+1/\alpha+1} (\alpha 2^\alpha)^{-\gamma/\alpha}}{\left(\frac{1+\alpha-\gamma}{\gamma} + 1\right)^{1/\alpha+1}} \zeta(\gamma/\alpha).
\end{aligned}$$

By the same calculation we have the second sum is bounded by

$$\begin{aligned}
& \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0,0.5]} \frac{z^{\gamma-1-\alpha} (\alpha 2^\alpha)^{-1/\alpha-1} \log(n)}{(z^{-\alpha} \alpha^{-1} 2^{-\alpha} + n)^{1/\alpha+1}} \\
&\leq \sup_{\alpha \in U} \frac{\left(\frac{(1+\alpha-\gamma)}{\gamma}\right)^{-\gamma/\alpha+1/\alpha+1} (\alpha 2^\alpha)^{-\gamma/\alpha}}{\left(\frac{1+\alpha-\gamma}{\gamma} + 1\right)^{1/\alpha+1}} \sum_{n=1}^{\infty} n^{-\gamma/\alpha} \log(n) \\
&= \sup_{\alpha \in U} \frac{\left(\frac{(1+\alpha-\gamma)}{\gamma}\right)^{-\gamma/\alpha+1/\alpha+1} (\alpha 2^\alpha)^{-\gamma/\alpha}}{\left(\frac{1+\alpha-\gamma}{\gamma} + 1\right)^{1/\alpha+1}} |\zeta'(\gamma/\alpha)|
\end{aligned}$$

which gives us

$$(I) \leq C_{10} \cdot C_{sum} (\zeta(\gamma/\alpha) + |\zeta'(\gamma/\alpha)|).$$

where  $C_{sum} = \sup_{\alpha \in U} \frac{\left(\frac{(1+\alpha-\gamma)}{\gamma}\right)^{-\gamma/\alpha+1/\alpha+1} (\alpha 2^\alpha)^{-\gamma/\alpha}}{\left(\frac{1+\alpha-\gamma}{\gamma} + 1\right)^{1/\alpha+1}} \leq 5.981$ .

To bound (II) we do the same, but using  $\sum_{j=1}^n \log(j) j^{-1} \leq \log^2(n)$

$$\begin{aligned}
& \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0,0.5]} z^\gamma (1 + n z^\alpha \alpha 2^\alpha)^{-1/\alpha-1} \sum_{j=1}^n (j^{-1} \log j) \\
&\leq C_{sum} \sum_{n=1}^{\infty} n^{-\gamma/\alpha} \log^2(n) \\
&\leq C_{sum} \zeta''(\gamma/\alpha)
\end{aligned}$$

giving

$$(II) \leq C_{11} \cdot C_{sum} \zeta''(\gamma/\alpha).$$

For (III) we use  $\sum_{j=1}^n (j^{-1} \sum_{k=1}^j k^{-1} \log k) \leq \sum_{j=1}^n (j^{-1} \log^2(j)) \leq \log^3(n)$  to get

$$\begin{aligned}
& \sum_{n=1}^{\infty} \sup_{\alpha \in U} \sup_{z \in (0,0.5]} z^\gamma (1 + n z^\alpha \alpha 2^\alpha)^{-1/\alpha-1} \sum_{j=1}^n (j^{-1} \sum_{k=1}^j k^{-1} \log k) \\
&\leq C_{sum} \sum_{n=1}^{\infty} n^{-\gamma/\alpha} \log^3(n) \\
&\leq C_{sum} |\zeta'''(\gamma/\alpha)|
\end{aligned}$$

This implies directly that

$$(III) \leq C_{12} \cdot C_{sum} \zeta'''(\gamma/\alpha).$$

In order to get the bound closer, we can use the technique of calculating the first  $N$  terms of  $\sum_{n=1}^{\infty} \|b_{\omega}\|_B$  using the computer calculations from 9 and the range estimation method from [42].

$$\begin{aligned} \sum_{\omega} \|b_{\omega}\|_B &\leq C_{10} \cdot C_{sum}(\zeta(\gamma/\alpha) + |\zeta'(\gamma/\alpha)| - \sum_{j=1}^N [j^{-\gamma/\alpha} + j^{-\gamma/\alpha} \log(j)]) \\ &\quad + C_{11} \cdot C_{sum}(\zeta''(\gamma/\alpha) - \sum_{j=1}^N [j^{-\gamma/\alpha} \log^2(j)]) \\ &\quad + C_{12} \cdot C_{sum}(|\zeta'''(\gamma/\alpha)| - \sum_{j=1}^N [j^{-\gamma/\alpha} \log^3(j)]) \\ &\quad + \sum_{1 \leq |\omega| \leq N} \|b_{\omega}\|_B. \end{aligned}$$

We calculate upper bounds on the derivatives of  $\zeta(x)$  using methods from [13].

Choosing  $N = 1000$  and  $j^* = 1000$  gives us  $\sum_{\omega} \|b_{\omega}\|_B \leq 1258$ . The tail of the sum starting at  $n = 1000$  gives  $\sum_{|\omega| \geq n} \|b_{\omega}\|_B \leq 0.002931$ .

**8.4. Bounding  $\sup_{\omega} |a_{\omega}|$ .** For  $\sup_{\omega} |a_{\omega}|$  we use lemma 5.2 from [32]. The proof of this lemma gives us

$$\frac{z_0^{\alpha}}{n} \cdot \frac{1}{1 + \alpha 2^{\alpha}} \leq z_n^{\alpha} \leq \frac{1}{z_0^{-\alpha} + n\alpha(1 - \alpha)2^{\alpha-1}}$$

from which we get  $C_1 = \frac{1}{1 + \alpha 2^{\alpha}}$  and  $C_2 = \frac{1}{\alpha(1 - \alpha)2^{\alpha-1}}$

$$(8.1) \quad \frac{z_0^{\alpha}}{n} \cdot C_1 \leq z_n^{\alpha} \leq \frac{1}{n} \cdot C_2.$$

Then  $z_0^{\alpha} \frac{C_1}{n} \leq z_n^{\alpha}$  gives us  $-\log z_0 \leq \frac{-1}{\alpha} \log \frac{C_1}{n} - \log z_0$ . To get a  $C_3$  such that  $\frac{-1}{\alpha} \log \frac{C_1}{n} \leq C_3 \log(n)$  we take  $C_3 = \log(C_1^{-1/\alpha}) + \frac{1}{\alpha}$ . Since  $C_3 > 1$

$$-\log z_0 \leq C_3(\log(n) - \log z_0).$$

Then from the proof of lemma 5.2 from [5] we have

$$(8.2) \quad \partial_{\alpha} z_{n+1} \leq 2^{\alpha} \sum_{j=1}^{n+1} z_j^{\alpha+1} (-\log 2z_j)$$

where  $\sup_{\omega} |a_{\omega}| \leq \sup_{z_0 \in [0, 0.5]} \partial_{\alpha} z_{n+1}$ . We use the fact that  $x^{\alpha+1}(-\log 2x)$  is monotonically increasing below  $x = 0.5 \exp(\frac{-1}{\alpha+1})$  to say that if  $C_2^{1/\alpha} j^{*-1/\alpha} \leq 0.5 \exp(\frac{-1}{\alpha+1})$  then

$$\begin{aligned} \partial_{\alpha} z_{n+1} &\leq 2^{\alpha} \sum_{j=1}^{n+1} z_j^{\alpha+1} (-\log 2z_j) \\ &= 2^{\alpha} \sum_{j=j^*}^{n+1} z_j^{\alpha+1} (-\log 2z_j) + 2^{\alpha} \sum_{j=1}^{j^*-1} z_j^{\alpha+1} (-\log 2z_j). \end{aligned}$$

We may use a computer to calculate the sum up to  $j^* - 1$  and we bound the rest as follows,

$$\begin{aligned}
& 2^\alpha \sum_{j=j^*}^{n+1} z_j^{\alpha+1} (-\log 2z_j) \\
& \leq 2^\alpha \sum_{j=j^*}^{n+1} [C_2^{1/\alpha} j^{-1/\alpha}]^{\alpha+1} (-\log (2C_2^{1/\alpha} j^{-1/\alpha})) \\
& = 2^\alpha \sum_{j=j^*}^{n+1} [C_2^{1/\alpha} j^{-1/\alpha}]^{\alpha+1} (-\log (j^{-1/\alpha}) - \log (2C_2^{1/\alpha})) \\
& \leq 2^\alpha \sum_{j=j^*}^{n+1} [C_2^{1/\alpha} j^{-1/\alpha}]^{\alpha+1} (-\log (j^{-1/\alpha})) \\
& \leq \frac{2^\alpha C_2^{(\alpha+1)/\alpha}}{\alpha} \sum_{j=j^*}^{n+1} j^{-1-1/\alpha} \log j.
\end{aligned}$$

Noticing that  $\sum_{j=1}^{\infty} j^{-1-1/\alpha} \log j = -\zeta'(1+1/\alpha)$  which can be calculated by methods from [13] gives us

(8.3)

$$\sup_{\omega} |a_{\omega}| \leq \frac{[-\zeta'(1+1/\alpha) - \sum_{j=1}^{j^*-1} j^{-1-1/\alpha} \log j] 2^\alpha}{\alpha(\alpha(1-\alpha)2^{\alpha-1})^{(\alpha+1)/\alpha}} + 2^\alpha \sum_{j=1}^{j^*-1} z_j^{\alpha+1} (-\log 2z_j)$$

which for  $\alpha = 0.125$  and taking  $j^* = 1000$  gives  $\sup_{\omega} |a_{\omega}| \leq 7107$

## 9. APPENDIX: COMPUTING DERIVATIVES

In order to calculate  $A_0$ ,  $B_0$ ,  $a_{\omega}$  and  $b_{\omega}$  we use an iterative formula. We start with

$$g_{\omega} \circ T_{\omega}(x) = x$$

from which we get

$$\begin{aligned}
(g_{\omega} \circ T_{\omega})'(x) &= g'_{\omega} \circ T_{\omega}(x) \cdot T'_{\omega}(x) = 1 \\
\implies g'_{\omega}(x) &= \frac{1}{T'_{\omega} \circ g_{\omega}(x)}
\end{aligned}$$

and

$$\begin{aligned}
\partial_{\alpha}(g_{\omega} \circ T_{\omega})(x) &= \partial_{\alpha} g_{\omega} \circ T_{\omega}(x) + g'_{\omega} \circ T_{\omega}(x) \cdot \partial_{\alpha} T_{\omega}(x) = 0 \\
\implies \partial_{\alpha} g_{\omega}(x) &= -\frac{\partial_{\alpha} T_{\omega} \circ g_{\omega}(x)}{T'_{\omega} \circ g_{\omega}(x)}.
\end{aligned}$$

We then use these to get

$$\begin{aligned}
(g'_{\omega} \circ T_{\omega})'(x) &= g''_{\omega} \circ T_{\omega}(x) \cdot T'_{\omega}(x) = -\frac{T''_{\omega}(x)}{(T'_{\omega}(x))^2} \\
\implies g''_{\omega}(x) &= -\frac{T''_{\omega} \circ g_{\omega}(x)}{(T'_{\omega} \circ g_{\omega}(x))^3}
\end{aligned}$$

and

$$\begin{aligned}\partial_\alpha(g'_\omega \circ T_\omega)(x) &= \partial_\alpha g'_\omega \circ T_\omega(x) + g''_\omega \circ T_\omega(x) \cdot \partial_\alpha T_\omega(x) = \partial_\alpha \frac{1}{T'_\omega(x)} \\ \implies \partial_\alpha g'_\omega(x) &= \frac{T''_\omega \circ g_\omega(x) \cdot \partial_\alpha T_\omega \circ g_\omega(x)}{(T'_\omega \circ g_\omega)^3} - \frac{\partial_\alpha T'_\omega \circ g_\omega(x)}{(T'_\omega \circ g_\omega(x))^2}.\end{aligned}$$

We already can calculate  $g_\omega$  so we need to calculate  $\partial_\alpha T_\omega$ ,  $T'_\omega$ ,  $\partial_\alpha T'_\omega$  and  $T''_\omega$ . Note that  $T_\omega = T_0^n \circ T_1$  where  $|\omega| = n$ , so

$$\begin{aligned}(T_0^n \circ T_1)' &= (T_0^n)' \circ T_1 \cdot T_1' \\ \partial_\alpha(T_0^n \circ T_1) &= \partial_\alpha(T_0^n) \circ T_1 + (T_0^n)' \circ T_1 \cdot \partial_\alpha T_1 \\ (T_0^n \circ T_1)'' &= (T_0^n)'' \circ T_1 \cdot (T_1')^2 + (T_0^n)' \circ T_1 \cdot T_1'' \\ \partial_\alpha(T_0^n \circ T_1)' &= \partial_\alpha(T_0^n)' \circ T_1 \cdot T_1' + (T_0^n)'' \circ T_1 \cdot T_1' \cdot \partial_\alpha T_1 + (T_0^n)' \circ T_1 \cdot \partial_\alpha T_1'\end{aligned}$$

which we may write as a matrix

$$\begin{pmatrix} T'_\omega \\ \partial_\alpha T_\omega \\ T''_\omega \\ \partial_\alpha T'_\omega \end{pmatrix} = \begin{pmatrix} T'_1 & 0 & 0 & 0 \\ \partial_\alpha T_1 & 1 & 0 & 0 \\ T''_1 & 0 & (T'_1)^2 & 0 \\ \partial_\alpha T'_1 & 0 & T'_1 \partial_\alpha T_1 & T'_1 \end{pmatrix} \cdot \begin{pmatrix} (T_0^n)' \circ T_1 \\ \partial_\alpha(T_0^n) \circ T_1 \\ (T_0^n)'' \circ T_1 \\ \partial_\alpha(T_0^n)' \circ T_1 \end{pmatrix}.$$

By the same logic we may write

$$\begin{aligned}\begin{pmatrix} T'_\omega \\ \partial_\alpha T_\omega \\ T''_\omega \\ \partial_\alpha T'_\omega \end{pmatrix} &= \begin{pmatrix} T'_1 & 0 & 0 & 0 \\ \partial_\alpha T_1 & 1 & 0 & 0 \\ T''_1 & 0 & (T'_1)^2 & 0 \\ \partial_\alpha T'_1 & 0 & T'_1 \partial_\alpha T_1 & T'_1 \end{pmatrix} \\ \cdot \begin{pmatrix} T'_0 \circ T_1 & 0 & 0 & 0 \\ \partial_\alpha T_0 \circ T_1 & 1 & 0 & 0 \\ T''_0 \circ T_1 & 0 & (T'_0 \circ T_1)^2 & 0 \\ \partial_\alpha T'_0 \circ T_1 & 0 & T'_0 \circ T_1 \partial_\alpha T_0 \circ T_1 & T'_0 \circ T_1 \end{pmatrix} &\begin{pmatrix} (T_0^{n-1})' \circ T_0 \circ T_1 \\ \partial_\alpha(T_0^{n-1}) \circ T_0 \circ T_1 \\ (T_0^{n-1})'' \circ T_0 \circ T_1 \\ \partial_\alpha(T_0^{n-1})' \circ T_0 \circ T_1 \end{pmatrix}.\end{aligned}$$

and use induction to give a series of matrices such that

$$\begin{aligned}\begin{pmatrix} T'_\omega \\ \partial_\alpha T_\omega \\ T''_\omega \\ \partial_\alpha T'_\omega \end{pmatrix} &= \begin{pmatrix} T'_1 & 0 & 0 & 0 \\ \partial_\alpha T_1 & 1 & 0 & 0 \\ T''_1 & 0 & (T'_1)^2 & 0 \\ \partial_\alpha T'_1 & 0 & T'_1 \partial_\alpha T_1 & T'_1 \end{pmatrix} \\ \cdot \begin{pmatrix} T'_0 \circ T_1 & 0 & 0 & 0 \\ \partial_\alpha T_0 \circ T_1 & 1 & 0 & 0 \\ T''_0 \circ T_1 & 0 & (T'_0 \circ T_1)^2 & 0 \\ \partial_\alpha T'_0 \circ T_1 & 0 & T'_0 \circ T_1 \partial_\alpha T_0 \circ T_1 & T'_0 \circ T_1 \end{pmatrix} \cdots \begin{pmatrix} T'_0 \circ T_0^{n-1} \circ T_1 \\ \partial_\alpha T_0 \circ T_0^{n-1} \circ T_1 \\ T''_0 \circ T_0^{n-1} \circ T_1 \\ \partial_\alpha T'_0 \circ T_0^{n-1} \circ T_1 \end{pmatrix}.\end{aligned}$$

Using

$$\begin{aligned}
T_0 &= x(1 + (2x)^\alpha) \\
T_1 &= 2x - 1 \\
T'_0 &= 1 + (1 + \alpha)(2x)^\alpha \\
T'_1 &= 2 \\
\partial_\alpha T_0 &= (\log(x) + \log(2))2^\alpha x^{\alpha+1} \\
\partial_\alpha T_1 &= 0 \\
T''_0 &= \alpha(1 + \alpha)2^\alpha x^{\alpha-1} \\
T''_1 &= 0 \\
\partial_\alpha T'_0 &= (2x)^\alpha((\alpha + 1)(\log(x) + \log(2)) + 1) \\
\partial_\alpha T'_1 &= 0
\end{aligned}$$

we are able to calculate explicitly the values  $A_0$ ,  $B_0$ ,  $a_\omega$  and  $b_\omega$ . To calculate  $a_\omega$  and  $b_\omega$  we use  $g_\omega = g_1 \circ g_0^{n-1}$  and we use  $T_0^m \circ T_1 \circ g_1 \circ g_0^{n-1} = g_0^{n-1-m}$  to calculate

$$\begin{aligned}
\begin{pmatrix} T'_\omega \circ g_\omega \\ \partial_\alpha T'_\omega \circ g_\omega \\ T''_\omega \circ g_\omega \\ \partial_\alpha T''_\omega \circ g_\omega \end{pmatrix} &= \begin{pmatrix} T'_1 \circ g_\omega & 0 & 0 & 0 \\ \partial_\alpha T'_1 \circ g_\omega & 1 & 0 & 0 \\ T''_1 \circ g_\omega & 0 & (T'_1)^2 \circ g_\omega & 0 \\ \partial_\alpha T''_1 \circ g_\omega & 0 & T'_1 \circ g_\omega \partial_\alpha T'_1 \circ g_\omega & T'_1 \circ g_\omega \end{pmatrix} \\
\cdot \begin{pmatrix} T'_0 \circ g_0^n & 0 & 0 & 0 \\ \partial_\alpha T'_0 \circ g_0^n & 1 & 0 & 0 \\ T''_0 \circ g_0^n & 0 & (T'_0 \circ g_0^n)^2 & 0 \\ \partial_\alpha T''_0 \circ g_0^n & 0 & T'_0 \circ g_0^n \partial_\alpha T'_0 \circ g_0^n & T'_0 \circ g_0^n \end{pmatrix} &\cdots \begin{pmatrix} T'_0 \circ g_0 \\ \partial_\alpha T'_0 \circ g_0 \\ T''_0 \circ g_0 \\ \partial_\alpha T''_0 \circ g_0 \end{pmatrix}
\end{aligned}$$

where we calculate  $g_0^m$  using the shooting method from section 7.1.1.

## REFERENCES

- [1] Antown, F., Dragičević, D., Froyland, G. Optimal linear responses for Markov chains and stochastically perturbed dynamical systems. *J. Stat. Phys.* 170 (2018), no. 6, 1051Ð1087.
- [2] Antown, F., Froyland, G., Junge, O. Linear response for the dynamic Laplacian and finite-time coherent sets. (2019) arXiv.
- [3] Aspenberg, M., Baladi, V., Leppänen, J., Persson, T. On the fractional susceptibility function of piecewise expanding maps. (2019) arXiv.
- [4] Bahsoun, W., Galatolo, S., Nisoli, I. Niu, X. A rigorous computational approach to linear response. *Nonlinearity* 31 (2018), no. 3, 1073–1109.
- [5] Bahsoun, W., Saussol, B., Linear response in the intermittent family: differentiation in a weighted  $C^0$ -norm. *Discrete Contin. Dynam. Syst.* 36 (12) (2016) 6657–6668.
- [6] Bahsoun, W., Ruziboev, M., Saussol, B. Linear response for random dynamical systems. *Adv. Math.* 364 (2020), 107011, 44 pp.
- [7] Baladi, V. (2014) Linear response, or else. Proceedings of the International Congress of Mathematicians, Seoul 2014. Vol. III, 525–545, Kyung Moon Sa, Seoul.
- [8] Baladi, V. On the susceptibility function of piecewise expanding interval maps, *Comm. Math. Phys.*, (2007) 839-859.
- [9] Baladi, V., Smiana, D. Fractional susceptibility functions for the quadratic family: Misiurewicz-Thurston parameters. (2020) arXiv .
- [10] Baladi, V., Todd, M., Linear response for intermittent maps. *Comm. Math. Phys.* 347 (2016), no. 3, 857–874. Dimension. (*Birkhäuser Boston*).
- [11] Butterly, O., Kiamari, N., Liverani, C., Locating Ruelle-Pollicott resonances. arXiv:2012.13145

- [12] Butterley, O., Liverani, C. Smooth Anosov flows: correlation spectra and stability. *J. Mod. Dyn.* (2007), 301–322.
- [13] Choudhury, B. The Riemann zeta-function and its derivatives. *Proceedings: Mathematical and Physical Sciences*, 450(1995), 477–499.
- [14] de Lima, A., Smania, D. Central limit theorem for the modulus of continuity of averages of observables on transversal families of piecewise expanding unimodal maps. *J. Inst. Math. Jussieu* 17 (2018), no. 3, 673–733.
- [15] Dolgopyat, D. On differentiability of SRB states for partially hyperbolic systems. *Invent. Math.* (2004), 389–449.
- [16] Dragičević, D., Sedro, J. Statistical stability and linear response for random hyperbolic dynamics. (2020) arXiv.
- [17] Frigo M., Johnson S. G., The Design and Implementation of FFTW3. *Proceedings of the IEEE* 93 (2), 216–231 (2005).
- [18] Galatolo, S.; Giulietti, P. A linear response for dynamical systems with additive noise. *Nonlinearity* 32 (2019), no. 6, 2269–2301.
- [19] S. Galatolo, M. Monge, I. Nisoli, F. Poloni. A general framework for the rigorous computation of invariant densities and the course-fine strategy. (preprint)
- [20] Galatolo, S., Nisoli, I., An elementary approach to rigorous approximation of invariant measures. *SIAM J. Appl. Dyn. Syst.* 13 (2014), no. 2, 958–985.
- [21] Galatolo, S., Nisoli, I. and Saussol, S., An elementary way to rigorously estimate convergence to equilibrium and escape rates. *J. Comput. Dyn.* 2 (2015), no. 1, 51–64.
- [22] Galatolo S., Pollicott M, Controlling the statistical properties of expanding maps. *Nonlinearity* 30 (2017), no. 7, 2737–2751.
- [23] S. Galatolo, J. Sedro (2019) Quadratic response of random and deterministic dynamical systems. arXiv:1908.00025
- [24] Gottwald, G. Introduction to focus issue: linear response theory: potentials and limits. *Chaos* 30(2), 020401 (2020). <https://doi.org/10.1063/5.0003135>.
- [25] Gouëzel, S., Liverani, C., Banach spaces adapted to Anosov systems, *Ergodic Theory Dynam. Systems* 26 (2006), 189–217.
- [26] Gutiérrez M. S., Lucarini V. Response and sensitivity using Markov chains. *J. Stat. Phys.* 179 (2020), no. 5-6, 1572–1593.
- [27] Higham N. J., Accuracy and Stability of Numerical Algorithms. Second Edition, SIAM, (2002)
- [28] Jézéquel, M. Parameter regularity of dynamical determinants of expanding maps of the circle and an application to linear response. *Discrete Contin. Dyn. Syst.*, 39 (2) (2019), 927–958.
- [29] Katok, A., Knieper, G., Pollicott, M., Weiss, H., Differentiability and analyticity of topological entropy for Anosov and geodesic flows. *Invent. Math.* 98 (1989), no. 3, 581–597.
- [30] KloECKner, B. The linear request problem. *Proc. Amer. Math. Soc.* 146 (2018), no. 7, 2953–2962.
- [31] Koltai, P, Lie, H-C., Plonka, M. Fréchet differentiable drift dependence of Perron-Frobenius and Koopman operators for non-deterministic dynamics. *Nonlinearity* 32 (2019), no. 11, 4232–4257.
- [32] Korepanov, A. Linear response for intermittent maps with summable and nonsummable decay of correlations. *Nonlinearity* 29, (2016), no. 6, 1735–1754.
- [33] Ledoux V., Moroz G. Evaluation of Chebyshev polynomials on intervals and application to root finding. Mathematical Aspects of Computer and Information Sciences 2019, Nov 2019, Gebze, Turkey. hal-02405752
- [34] Liverani, C., Saussol, B., Vaienti, S. A probabilistic approach to intermittency, *Ergodic theory Dynam. Systems*, 19, (1999), 671–685.
- [35] Lucarini, V. Response operators for Markov processes in a finite state space: radius of convergence and link to the response theory for Axiom A systems. *J. Stat. Phys.* 162 (2016), no. 2, 312–333.
- [36] Ni, A. Linear response algorithm for differentiating stationary measures of chaos.(2020) arXiv.
- [37] Pollicott, M., Vytanova P., *Linear response and periodic points. Nonlinearity* 29 (2016), no. 10, 3047–3066.
- [38] Ruelle, D., Differentiation of SRB states. *Comm. Math. Phys.* 187 (1997) 227–241.
- [39] Sedro, J. Pre-threshold fractional susceptibility functions at Misiurewicz parameters. (2020) arXiv.

- [40] Sélley, F., Tanzi, M. Linear response for a family of self-consistent transfer operators. (2020) arXiv.
- [41] Lloyd N. Trefethen, Approximation Theory and Approximation Practice. (2018) SIAM
- [42] Tucker W. *Validated Numerics: A Short Introduction to Rigorous Computations* PRINCETON; OXFORD: Princeton University (2011).
- [43] Wormell, C.L., Spectral Galerkin methods for transfer operators in uniformly expanding dynamics. *Numerische Mathematik* 14 (2019) 421–463
- [44] Wormell, C., Gottwald, G. Linear response for macroscopic observables in high-dimensional systems. *Chaos* 29 (2019), no. 11, 113127, 18 pp.
- [45] S. Xiang, X. Chen, H. Wang, Error bounds for approximation in Chebyshev points, *Numer. Math.* (2010) 116:463–491 DOI 10.1007/s00211-010-0309-4

INSTITUTO DE MATEMATICA - UFRJ Av. ATHOS DA SILVEIRA RAMOS 149, CENTRO DE TECNOLOGIA - BLOCO C CIDADE UNIVERSITARIA - ILHA DO FUNDÃO. CAIXA POSTAL 68530 21941-909 RIO DE JANEIRO - RJ - BRASIL

*Email address:* `nisoli@im.ufrj.br`

DEPARTMENT OF MATHEMATICAL SCIENCES, LOUGHBOROUGH UNIVERSITY, LOUGHBOROUGH, LEICESTERSHIRE, LE11 3TU, UK

*Email address:* `T.Taylor-Crush@lboro.ac.uk`