

A Note on Numerical Fluxes Conserving a Member of Harten's One-Parameter Family of Entropies for the Compressible Euler Equations

Hendrik Ranocha^{*1}

¹Applied Mathematics, University of Hamburg, Germany

March 8, 2022

Entropy-conserving numerical fluxes are a cornerstone of modern high-order entropy-dissipative discretizations of conservation laws. In addition to entropy conservation, other structural properties mimicking the continuous level such as pressure equilibrium and kinetic energy preservation are important. This note proves that there are no numerical fluxes conserving (one of) Harten's entropies for the compressible Euler equations that also preserve pressure equilibria and have a density flux independent of the pressure. This is in contrast to fluxes based on the physical entropy, where even kinetic energy preservation can be achieved in addition.

Key words. entropy stability, numerical fluxes, flux differencing, kinetic energy preservation, pressure equilibrium preservation, local linear stability

1 Introduction

Ever since the seminal work of Tadmor [22], researchers have been interested in entropy-dissipative numerical methods for conservation laws and related models. Usually, these methods have improved robustness properties, even for underresolved simulations. Nowadays, several means have been explored to ensure entropy stability. One of the most popular and successful approaches is based on entropy-conservative (EC) numerical fluxes. These can be used to construct high-order central-type methods using flux differencing [6] to which appropriate dissipation can be added. A recent alternative is the general algebraic approach of [1, 2].

There are several numerical fluxes for the compressible Euler equations conserving the physical (logarithmic) entropy [4, 10, 12]. Harten [8] studied another family of entropies for the compressible Euler equations. Although these entropies do not symmetrize the heat flux terms in the compressible Navier-Stokes equations [9], they are still of interest, for example to construct entropy splitting methods or related EC fluxes [21].

Entropy conservation alone is often insufficient to construct good numerical methods. Of course, dissipation and related issues such as shock capturing and positivity of the density and internal energy are also important but are not the focus of this contribution. Instead, preservation of the kinetic energy [11, 13, 14] and pressure equilibria [16, 20] is considered. Moreover, the numerical density flux should not depend on the pressure, in accordance with physical expectations. This is

*ORCID: 0000-0002-3456-2277

discussed further in [5, 12], where positivity failure could be identified for certain setups with large pressure jumps and constant densities and velocities, even in the presence of strong (numerical) dissipation.

The main contribution of this note is to prove that there are no numerical fluxes conserving Harten's entropies for the compressible Euler equations that also preserve pressure equilibria and have a density flux independent of the pressure (Section 2). Further discussion of this result is presented in Section 3.

2 Main result

It suffices to concentrate on the 1D compressible Euler equations

$$\partial_t \mathbf{u} + \partial_x \mathbf{f}(\mathbf{u}) = \mathbf{0}, \quad \mathbf{u} = (\varrho, \varrho v, \varrho e)^T, \quad \mathbf{f}(\mathbf{u}) = (\varrho v, \varrho v^2 + p, (\varrho e + p)v)^T, \quad (2.1)$$

where ϱ is the density, v the velocity, ϱe the total energy, and the pressure $p = (\gamma - 1)(\varrho e - \varrho v^2/2)$ is given by the ideal gas law with ratio of specific heats $\gamma > 1$. Harten [8] considered entropies of the form $U(\mathbf{u}) = -\varrho h(s)$, where $s = \log(p/\varrho^\gamma)$ and h is a smooth function satisfying $h''(s)/h'(s) < 1/\gamma$. A special one-parameter family of these entropies is given by

$$U(\mathbf{u}) = -\frac{\gamma + \alpha}{\gamma - 1} \varrho (p/\varrho^\gamma)^{1/(\alpha + \gamma)}, \quad \alpha > 0 \text{ or } \alpha < -\gamma, \quad (2.2)$$

where the restriction of the parameter α ensures convexity of U [21]. The members of this one-parameter family (2.2) of entropy functions for the compressible Euler equations are often referred to as Harten's entropies in the literature, e.g., in [21]. The associated entropy variables $\mathbf{w} = U'(\mathbf{u})$ and the flux potential ψ [22] are given by

$$\mathbf{w} = \frac{\varrho}{p} (p/\varrho^\gamma)^{1/(\alpha + \gamma)} \left(-\frac{\alpha}{\gamma - 1} \frac{p}{\varrho} - \frac{1}{2} v^2, v, -1 \right), \quad \psi = \varrho (p/\varrho^\gamma)^{1/(\alpha + \gamma)} v. \quad (2.3)$$

This note focuses on two-point numerical fluxes for the compressible Euler equations. Such a two-point numerical flux \mathbf{f}^{num} is characterized as follows.

Definition 2.1 (Entropy conservation [22]). The numerical flux \mathbf{f}^{num} is EC if $[[\mathbf{w}]] \cdot \mathbf{f}^{\text{num}} - [[\psi]] = 0$, where $[[a]] = a_+ - a_-$ is the common jump operator. \triangleleft

Definition 2.2 (Pressure equilibrium preservation [16]). A numerical flux $\mathbf{f}^{\text{num}} = (\mathbf{f}_\varrho^{\text{num}}, \mathbf{f}_{\varrho v}^{\text{num}}, \mathbf{f}_{\varrho e}^{\text{num}})$ is pressure equilibrium preserving (PEP) if $\mathbf{f}_{\varrho v}^{\text{num}} = v \mathbf{f}_\varrho^{\text{num}} + \text{const}_1(p, v)$ and $\mathbf{f}_{\varrho e}^{\text{num}} = \frac{1}{2} v^2 \mathbf{f}_\varrho^{\text{num}} + \text{const}_2(p, v)$ whenever the velocity v and the pressure p are constant. Here, $\text{const}_i(p, v)$, $i \in \{1, 2\}$, denote some generic constants depending only on the constants v and p . \triangleleft

As discussed in [7, 16, 20], pressure equilibria are important setups where the velocity and the pressure are constant. In this case, the compressible Euler equations are reduced to linear advection equations. Pressure equilibrium preserving schemes keep this property at the discrete level [16].

The main result of this note is

Theorem 2.3. *There is no two-point numerical flux for the compressible Euler equations (2.1) that is entropy-conserving in the sense of Tadmor (Def. 2.1) for a member of Harten's one-parameter family of entropies (2.2), pressure equilibrium preserving (Def. 2.2), and has a density flux that does not depend on the pressure.*

The proof of Theorem 2.3 is divided into the following steps.

Lemma 2.4. *For $p \equiv \text{const}$ and $v \equiv \text{const}$, a numerical flux that is PEP and EC for a member of Harten's one-parameter family of entropies (2.2) has a density flux of the form*

$$\mathbf{f}_\varrho^{\text{num}} = -\frac{\gamma}{\alpha} \frac{[[\varrho^{\alpha/(\alpha + \gamma)}]]}{[[\varrho^{-\gamma/(\alpha + \gamma)}]]} v. \quad (2.4)$$

Proof. For constant pressure and velocity,

$$\begin{aligned} \llbracket \mathbf{w} \rrbracket \cdot \mathbf{f}^{\text{num}} - \llbracket \psi \rrbracket &= \left(-\frac{\alpha}{\gamma-1} p^{1/(\alpha+\gamma)} \llbracket \varrho^{-\gamma/(\alpha+\gamma)} \rrbracket - \frac{1}{2} p^{(1-\alpha-\gamma)/(\alpha+\gamma)} v^2 \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \right) \mathbf{f}_\varrho^{\text{num}} \\ &+ p^{(1-\alpha-\gamma)/(\alpha+\gamma)} v \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \mathbf{f}_{\varrho v}^{\text{num}} - p^{(1-\alpha-\gamma)/(\alpha+\gamma)} \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \mathbf{f}_{\varrho e}^{\text{num}} \\ &- p^{1/(\alpha+\gamma)} v \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket. \end{aligned} \quad (2.5)$$

A PEP flux is of the form $\mathbf{f}_{\varrho v}^{\text{num}} = v \mathbf{f}_\varrho^{\text{num}} + p$, $\mathbf{f}_{\varrho e}^{\text{num}} = \frac{1}{2} v^2 \mathbf{f}_\varrho^{\text{num}} + p v \gamma / (\gamma - 1)$. Thus,

$$\begin{aligned} \llbracket \mathbf{w} \rrbracket \cdot \mathbf{f}^{\text{num}} - \llbracket \psi \rrbracket &= \left(-\frac{\alpha}{\gamma-1} p^{1/(\alpha+\gamma)} \llbracket \varrho^{-\gamma/(\alpha+\gamma)} \rrbracket - \frac{1}{2} p^{(1-\alpha-\gamma)/(\alpha+\gamma)} v^2 \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \right) \mathbf{f}_\varrho^{\text{num}} \\ &+ p^{(1-\alpha-\gamma)/(\alpha+\gamma)} v^2 \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \mathbf{f}_\varrho^{\text{num}} + p^{1/(\alpha+\gamma)} v \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \\ &- \frac{1}{2} p^{(1-\alpha-\gamma)/(\alpha+\gamma)} v^2 \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \mathbf{f}_\varrho^{\text{num}} - \frac{\gamma}{\gamma-1} p^{1/(\alpha+\gamma)} v \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \\ &- p^{1/(\alpha+\gamma)} v \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket \\ &= -\frac{\alpha}{\gamma-1} p^{1/(\alpha+\gamma)} \llbracket \varrho^{-\gamma/(\alpha+\gamma)} \rrbracket \mathbf{f}_\varrho^{\text{num}} - \frac{\gamma}{\gamma-1} p^{1/(\alpha+\gamma)} v \llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket. \end{aligned} \quad (2.6)$$

Hence, the EC condition is equivalent to (2.4). \square

Lemma 2.5. For $v \equiv \text{const}$ and $p_- = p_+ (\varrho_- / \varrho_+)^{\alpha/(\alpha+\gamma-1)}$, a numerical flux that is EC for a member of Harten's one-parameter family of entropies (2.2) has a density flux of the form

$$\mathbf{f}_\varrho^{\text{num}} = -\frac{\gamma-1}{\alpha} \frac{\llbracket \varrho^{\alpha/(\alpha+\gamma-1)} \rrbracket}{\llbracket \varrho^{(1-\gamma)/(\alpha+\gamma-1)} \rrbracket} v. \quad (2.7)$$

Proof. For this special choice of the pressure,

$$\begin{aligned} \frac{\varrho_-}{p_-} (p_- / \varrho_-^\gamma)^{1/(\alpha+\gamma)} &= \varrho_-^{\alpha/(\alpha+\gamma)} p_-^{(1-\alpha-\gamma)/(\alpha+\gamma)} = \varrho_-^{\alpha/(\alpha+\gamma)} (\varrho_- / \varrho_+)^{-\alpha/(\alpha+\gamma)} p_+^{(1-\alpha-\gamma)/(\alpha+\gamma)} \\ &= \varrho_+^{\alpha/(\alpha+\gamma)} p_+^{(1-\alpha-\gamma)/(\alpha+\gamma)} = \frac{\varrho_+}{p_+} (p_+ / \varrho_+^\gamma)^{1/(\alpha+\gamma)}, \end{aligned} \quad (2.8)$$

i.e., $\llbracket (\varrho/p)(p/\varrho^\gamma)^{1/(\alpha+\gamma)} \rrbracket = 0$. Thus, for $v \equiv \text{const}$, (2.3) yields

$$\llbracket \mathbf{w} \rrbracket \cdot \mathbf{f}^{\text{num}} - \llbracket \psi \rrbracket = -\frac{\alpha}{\gamma-1} \llbracket (p/\varrho^\gamma)^{1/(\alpha+\gamma)} \rrbracket \mathbf{f}_\varrho^{\text{num}} - \llbracket (\varrho/p)(p/\varrho^\gamma)^{1/(\alpha+\gamma)} \rrbracket v. \quad (2.9)$$

Consequently, an EC flux must be of the form

$$\mathbf{f}_\varrho^{\text{num}} = -\frac{\gamma-1}{\alpha} \frac{\llbracket (\varrho/p)(p/\varrho^\gamma)^{1/(\alpha+\gamma)} \rrbracket}{\llbracket (p/\varrho^\gamma)^{1/(\alpha+\gamma)} \rrbracket} v. \quad (2.10)$$

Inserting the pressure ratio, the fraction of the jump terms can be written as

$$\begin{aligned} \frac{\varrho_+^{\alpha/(\alpha+\gamma)} p_+^{1/(\alpha+\gamma)} - \varrho_-^{\alpha/(\alpha+\gamma)} p_-^{1/(\alpha+\gamma)}}{\varrho_+^{-\gamma/(\alpha+\gamma)} p_+^{1/(\alpha+\gamma)} - \varrho_-^{-\gamma/(\alpha+\gamma)} p_-^{1/(\alpha+\gamma)}} &= \frac{\varrho_+^{\alpha/(\alpha+\gamma)} - \varrho_-^{\alpha/(\alpha+\gamma)} (\varrho_- / \varrho_+)^{\alpha/((\alpha+\gamma-1)(\alpha+\gamma))}}{\varrho_+^{-\gamma/(\alpha+\gamma)} - \varrho_-^{-\gamma/(\alpha+\gamma)} (\varrho_- / \varrho_+)^{\alpha/((\alpha+\gamma-1)(\alpha+\gamma))}} \\ &= \frac{\varrho_+^{\alpha/(\alpha+\gamma-1)} - \varrho_-^{\alpha/(\alpha+\gamma-1)}}{\varrho_+^{-\gamma/(\alpha+\gamma-1)} - \varrho_-^{-\gamma/(\alpha+\gamma-1)}}, \end{aligned} \quad (2.11)$$

proving (2.7). \square

Proof of Theorem 2.3. For given α, γ , and ϱ_{\pm} , such a flux needs to satisfy both (2.4) and (2.7), i.e.,

$$\gamma \frac{\llbracket \varrho^{\alpha/(\alpha+\gamma)} \rrbracket}{\llbracket \varrho^{-\gamma/(\alpha+\gamma)} \rrbracket} = (\gamma - 1) \frac{\llbracket \varrho^{\alpha/(\alpha+\gamma-1)} \rrbracket}{\llbracket \varrho^{(1-\gamma)/(\alpha+\gamma-1)} \rrbracket}. \quad (2.12)$$

For fixed α and γ , it is easy to find $\varrho_- \neq \varrho_+$ such that this equation is not satisfied. Hence, a numerical flux with all properties listed in Theorem 2.3 cannot exist. \square

3 Discussion

There are no numerical fluxes conserving a member of Harten’s one-parameter family of entropies (2.2) for the compressible Euler equations that also preserve pressure equilibria and have a density flux independent of the pressure (Theorem 2.3). This result is in contrast to fluxes conserving the physical (logarithmic) entropy, where kinetic energy preservation can be achieved in addition to all of these properties [13, 14], resulting in an essentially unique numerical flux [16].

Following the uniqueness proof of such a numerical flux based on the logarithmic entropy presented in [16], it is tempting to choose a density flux based on a simplified setting, e.g., for constant velocity and pressure as in Lemma 2.4. Then, kinetic energy preservation determines the momentum flux accordingly [13, 14] and one might expect to be able to use the EC criterion to find an energy flux. However, this does not work in general since the energy part of the numerical flux can be orthogonal to the jump of the entropy variables as in Lemma 2.5. In this case, such an approach will often lead to a blow-up of the derived component for the total energy flux. Hence, one cannot impose a form of the density flux in general.




Figure 1: Spectra of discontinuous Galerkin semidiscretizations of the 2D compressible Euler equations conserving Harten’s entropy (2.2) with parameter α using some values of $\beta = (\alpha + \gamma)/(1 - \gamma)$ recommended for entropy splitting discretizations [21] and $\alpha = 1$. The EC discretizations use the flux of [21] and tensor product Lobatto Legendre bases with polynomials of degree five on 4×4 elements. Using the central flux instead results in a spectrum with negligible imaginary part [7, 16].

Moreover, it is interesting to note that EC fluxes based on Harten’s one-parameter family of entropies do not solve the local linear/energy stability issues discussed in [7, 16]. This is demonstrated by the spectra of discontinuous Galerkin semidiscretizations of the 2D compressible Euler equations shown in Figure 1. The setup uses the density wave example described in [16]. If the central flux $\mathbf{f}^{\text{num}} = \{\{\mathbf{f}\}\}$ is used instead of an EC flux, the spectrum of the resulting semidiscretization is essentially purely imaginary (ignoring floating point errors etc.), indicating (marginal) local linear/energy stability of the method [7, 16]. This property is desired and reasonable for this physical setup, since the density wave reduces the compressible Euler equations to linear advection equations, see also the extended discussion in [7, 16].

The semidiscretizations used to compute the spectra above are implemented using Trixi.jl [17, 19]. The Jacobian is computed using automatic differentiation [18] in Julia [3]. All source code required to reproduce the examples as well as additional material verifying the implementation and some calculations presented in this article are available online [15].

Acknowledgments

Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2044-390685587, Mathematics Münster: Dynamics-Geometry-Structure. Special thanks to Gregor Gassner and Ayaboe Edoh for discussions related to this manuscript.

References

- [1] R. Abgrall. "A general framework to construct schemes satisfying additional conservation relations. Application to entropy conservative and entropy dissipative schemes." In: *Journal of Computational Physics* 372 (2018), pp. 640–666. doi: 10.1016/j.jcp.2018.06.031. arXiv: 1711.10358 [math.NA].
- [2] R. Abgrall, P. Öffner, and H. Ranocha. "Reinterpretation and Extension of Entropy Correction Terms for Residual Distribution and Discontinuous Galerkin Schemes: Application to Structure Preserving Discretization." In: *Journal of Computational Physics* 453 (Mar. 2022), p. 110955. doi: 10.1016/j.jcp.2022.110955. arXiv: 1908.04556 [math.NA].
- [3] J. Bežanson, A. Edelman, S. Karpinski, and V. B. Shah. "Julia: A Fresh Approach to Numerical Computing." In: *SIAM Review* 59.1 (2017), pp. 65–98. doi: 10.1137/141000671. arXiv: 1411.1607 [cs.MS].
- [4] P. Chandrashekar. "Kinetic Energy Preserving and Entropy Stable Finite Volume Schemes for Compressible Euler and Navier-Stokes Equations." In: *Communications in Computational Physics* 14.5 (2013), pp. 1252–1286. doi: 10.4208/cicp.170712.010313a.
- [5] D. Derigs, A. R. Winters, G. J. Gassner, and S. Walch. "A novel averaging technique for discrete entropy-stable dissipation operators for ideal MHD." In: *Journal of Computational Physics* 330 (2017), pp. 624–632. doi: 10.1016/j.jcp.2016.10.055.
- [6] T. C. Fisher and M. H. Carpenter. "High-order entropy stable finite difference schemes for nonlinear conservation laws: Finite domains." In: *Journal of Computational Physics* 252 (2013), pp. 518–557. doi: 10.1016/j.jcp.2013.06.014.
- [7] G. J. Gassner, M. Svärd, and F. J. Hindenlang. "Stability Issues of Entropy-Stable and/or Split-form High-order Schemes." In: *Journal of Scientific Computing* 90.3 (2022), pp. 1–36. doi: 10.1007/s10915-021-01720-8. arXiv: 2007.09026 [math.NA].
- [8] A. Harten. "On the symmetric form of systems of conservation laws with entropy." In: *Journal of Computational Physics* 49.1 (1983), pp. 151–164. doi: 10.1016/0021-9991(83)90118-3.
- [9] T. J. R. Hughes, L. P. Franca, and M. Mallet. "A new finite element formulation for computational fluid dynamics: I. Symmetric forms of the compressible Euler and Navier-Stokes equations and the second law of thermodynamics." In: *Computer Methods in Applied Mechanics and Engineering* 54.2 (1986), pp. 223–234. doi: 10.1016/0045-7825(86)90127-1.
- [10] F. Ismail and P. L. Roe. "Affordable, entropy-consistent Euler flux functions II: Entropy production at shocks." In: *Journal of Computational Physics* 228.15 (2009), pp. 5410–5436. doi: 10.1016/j.jcp.2009.04.021.
- [11] A. Jameson. "Formulation of Kinetic Energy Preserving Conservative Schemes for Gas Dynamics and Direct Numerical Simulation of One-Dimensional Viscous Compressible Flow in a Shock Tube Using Entropy and Kinetic Energy Preserving Schemes." In: *Journal of Scientific Computing* 34.2 (2008), pp. 188–208. doi: 10.1007/s10915-007-9172-6.
- [12] H. Ranocha. "Comparison of Some Entropy Conservative Numerical Fluxes for the Euler Equations." In: *Journal of Scientific Computing* 76.1 (July 2018), pp. 216–242. doi: 10.1007/s10915-017-0618-1. arXiv: 1701.02264 [math.NA].

- [13] H. Ranocha. “Entropy Conserving and Kinetic Energy Preserving Numerical Methods for the Euler Equations Using Summation-by-Parts Operators.” In: *Spectral and High Order Methods for Partial Differential Equations ICOSAHOM 2018*. Ed. by S. J. Sherwin, D. Moxey, J. Peiró, P. E. Vincent, and C. Schwab. Vol. 134. Lecture Notes in Computational Science and Engineering. Cham: Springer, Aug. 2020, pp. 525–535. DOI: 10.1007/978-3-030-39647-3_42.
- [14] H. Ranocha. “Generalised Summation-by-Parts Operators and Entropy Stability of Numerical Methods for Hyperbolic Balance Laws.” PhD thesis. TU Braunschweig, Feb. 2018.
- [15] H. Ranocha. *Reproducibility repository for “A Note on Numerical Fluxes Conserving Harten’s Entropies for the Compressible Euler Equations”*. https://github.com/ranocha/paper-2022-Euler_Harten_EC. 2022. DOI: 10.5281/zenodo.5837898.
- [16] H. Ranocha and G. J. Gassner. “Preventing pressure oscillations does not fix local linear stability issues of entropy-based split-form high-order schemes.” In: *Communications on Applied Mathematics and Computation* (Aug. 2021). DOI: 10.1007/s42967-021-00148-z. arXiv: 2009.13139 [math.NA].
- [17] H. Ranocha, M. Schlottke-Lakemper, A. R. Winters, E. Faulhaber, J. Chan, and G. Gassner. “Adaptive numerical simulations with Trixi.jl: A case study of Julia for scientific computing.” In: *Proceedings of the JuliaCon Conferences 1.1* (Jan. 2022), p. 77. DOI: 10.21105/jcon.00077. arXiv: 2108.06476 [cs.MS].
- [18] J. Revels, M. Lubin, and T. Papamarkou. *Forward-Mode Automatic Differentiation in Julia*. July 2016. arXiv: 1607.07892 [cs.MS].
- [19] M. Schlottke-Lakemper, A. R. Winters, H. Ranocha, and G. J. Gassner. “A purely hyperbolic discontinuous Galerkin approach for self-gravitating gas dynamics.” In: *Journal of Computational Physics* 442 (June 2021), p. 110467. DOI: 10.1016/j.jcp.2021.110467. arXiv: 2008.10593 [math.NA].
- [20] N. Shima, Y. Kuya, Y. Tamaki, and S. Kawai. “Preventing spurious pressure oscillations in split convective form discretization for compressible flows.” In: *Journal of Computational Physics* 427 (2021), p. 110060. DOI: 10.1016/j.jcp.2020.110060.
- [21] B. Sjögren and H. Yee. “Entropy Stable Method for the Euler Equations Revisited: Central Differencing via Entropy Splitting and SBP.” In: *Journal of Scientific Computing* 81.3 (2019), pp. 1359–1385. DOI: 10.1007/s10915-019-01013-1.
- [22] E. Tadmor. “The numerical viscosity of entropy stable schemes for systems of conservation laws. I.” In: *Mathematics of Computation* 49.179 (1987), pp. 91–103. DOI: 10.1090/S0025-5718-1987-0890255-3.