# Numerical evaluation of ODE solutions by Monte Carlo enumeration of Butcher series

Guillaume Penent<sup>∗</sup> Nicolas Privault†

Division of Mathematical Sciences School of Physical and Mathematical Sciences Nanyang Technological University 21 Nanyang Link, Singapore 637371

January 19, 2022

#### Abstract

We present an algorithm for the numerical solution of ordinary differential equations by random enumeration of the Butcher trees used in the implementation of the Runge-Kutta method. Our Monte Carlo scheme allows for the direct numerical evaluation of an ODE solution at any given time within a certain interval, without iteration through multiple time steps. In particular, this approach does not involve a discretization step size, and it does not require the truncation of Taylor series.

Keywords: Ordinary differential equations, Runge-Kutta method, Butcher series, random trees, Monte Carlo method.

Mathematics Subject Classification (2020): 65L06, 34A25, 34-04, 05C05, 65C05.

### 1 Introduction

Taylor expansions appearing in the Runge–Kutta methods for the numerical solution of ordinary differential equations (ODEs) have been expressed by rooted trees enumeration in terms of Butcher series [\[But63\]](#page-21-0), [\[But16\]](#page-21-1), see Chapters 4-6 of [\[DB02\]](#page-21-2), and [\[MMMKV17\]](#page-21-3) for a recent review starting from the early work of  $[\text{Cay57}]$ . It is known that the solution  $y(t)$ of the autonomous ODE

$$
\begin{cases}\ny'(t) = f(y(t)) \\
y(0) = y_0 \in \mathbb{R}, & t \in \mathbb{R}_+, \n\end{cases}
$$

<sup>∗</sup>[pene0001@e.ntu.edu.sg](mailto:PENE0001@e.ntu.edu.sg)

<sup>†</sup>[nprivault@ntu.edu.sg](mailto:nprivault@ntu.edu.sg)

in  $\mathbb{R}^d$ , where  $f(y) = (f^1(y), \ldots, f^d(y))$  is a smooth  $\mathbb{R}^d$ -valued function of y in a domain of  $\mathbb{R}^d$ , can be expressed as

<span id="page-1-0"></span>
$$
y(t) = y_0 + tf(y_0) + \frac{t^2}{2}f'f(y_0) + \frac{t^3}{6}f'f'f(y_0) + \frac{t^3}{6}f''[f,f](y_0) + \cdots
$$
 (1.1)

,

where we use the notation

$$
f'f(y_0) = \left(\sum_{k=1}^d \frac{\partial f^i}{\partial x_k}(y_0) f^k(y_0)\right)_{i=1,\dots,d},
$$
  

$$
f'f'(y_0) = \left(\sum_{k,l=1}^d \frac{\partial f^i}{\partial x_k}(y_0) \frac{\partial f^i}{\partial x_l}(y_0) f^l(y_0)\right)_{i=1,\dots,d},
$$
  

$$
f''[f, f](y_0) = \left(\sum_{k,l=1}^d \frac{\partial^2 f^i}{\partial x_k \partial x_l}(y_0) f^k(y_0) f^l(y_0)\right)_{i=1,\dots,d}
$$

etc. In addition, the expansion [\(1.1\)](#page-1-0) can be coded and enumerated using the following sequence of Butcher trees.



The numerical evaluation of Butcher series involves tree enumeration up to a certain order that determines the level of precision of the algorithm.

In this paper, we propose an alternative method to the numerical evaluation of ODE solutions, based on a random enumeration of Butcher trees by Monte Carlo simulation. Probabilistic methods based on the Feynman-Kac formula provide alternatives to finite difference schemes, and have been successfully applied to the solution of partial differential equations. In particular, stochastic branching mechanisms have been used to represent the solutions of partial differential equations in [\[Sko64\]](#page-21-5), [\[INW69\]](#page-21-6), [\[NS69\]](#page-21-7), [\[McK75\]](#page-21-8), [\[LM96\]](#page-21-9), [\[CLM08\]](#page-21-10). This branching argument has been recently extended in  $[HLOT+19]$  to the treatment polynomial non-linearities in gradient terms, see also [\[PP21\]](#page-21-12) for nonlocal and fractional PDEs.

In Theorem [3.2,](#page-6-0) under suitable integrability conditions we express ODE solutions as the expected value of a functional of random Butcher trees which encode nonlinearities. Then in Proposition [3.3](#page-8-0) we provide sufficient conditions ensuring that the representation formula of Theorem [3.2](#page-6-0) holds at any time within certain interval. Numerical values of ODE solutions can be computed beyond that initial interval by iterating the method and by piecing together the solutions obtained on adjacent intervals.

This approach complements the use of the Feynman-Kac formula for the numerical estimation of the solutions of partial differential equations, see also [\[SH21\]](#page-21-13) for a different approach to the Feynman-Kac representation of ODE solutions. Other links between Butcher trees and probability theory have been pointed out in [\[Maz04\]](#page-21-14), see also [\[Ski92\]](#page-21-15) for the numerical solution of ODEs as an inference problem by Bayesian techniques.

In Section [2](#page-2-0) we introduce the construction of coding trees that will be used for the numerical solution of ODEs. Section [3](#page-6-1) presents the probabilistic representation formula of ODE solutions obtained by the random generation of coding trees. In Section [4](#page-10-0) we consider examples and in Section [5](#page-13-0) we describe the correspondance between Butcher trees and coding trees, namely we show how any Butcher tree can be recovered by performing a depth first search on the corresponding coding tree. Section [6](#page-16-0) considers numerical applications.

### <span id="page-2-0"></span>2 Coding trees

This section introduces the random coding trees used for the probabilistic representation of ODE solutions. Let  $y(t)$  be the solution of an ODE of the form

<span id="page-2-2"></span><span id="page-2-1"></span>
$$
y(t) = y_0 + \int_0^t f(s, y(s))ds, \qquad t \in \mathbb{R}_+,
$$
\n(2.1)

where f is a smooth function defined on a domain of  $\mathbb{R}_+ \times \mathbb{R}$ . In order to solve [\(2.1\)](#page-2-1), we expand  $f(s, y(s))$  as

$$
f(s, y(s)) = f(0, y_0) + \int_0^s (\partial_0 f(u, y(u)) + f(u, y(u))\partial_1 f(u, y(u)))du
$$
 (2.2)

by differentiating  $v(s) := f(s, y(s))$ , where

$$
\partial_0 f(t, y) := \frac{\partial f}{\partial t}(t, y)
$$
 and  $\partial_1 f(t, y) := \frac{\partial f}{\partial y}(t, y).$ 

In the sequel, given g a function on  $\mathbb{R}_+ \times \mathbb{R}$  we let  $g^*$  denote the mapping

<span id="page-3-0"></span>
$$
g^*: \mathbb{R}^{\mathbb{R}_+} \longrightarrow \mathbb{R}^{\mathbb{R}_+}
$$
  

$$
(t \mapsto y(t)) \longmapsto g^*(y) := (t \mapsto g(t, y(t))),
$$
 (2.3)

where  $\mathbb{R}^{\mathbb{R}_+}$  represents the set of functions from  $\mathbb{R}_+$  to  $\mathbb{R}$ . In order to formalize and extend the iteration initiated in [\(2.2\)](#page-2-2), we introduce the following definition.

**Definition 2.1** We let C denote the set of functions from  $\mathbb{R}^{\mathbb{R}_+}$  to  $\mathbb{R}^{\mathbb{R}_+}$  called codes, defined as

$$
\mathcal{C} := \left\{ \mathrm{Id}, \, \left( \partial_0^k \partial_1^l f \right)^*, \, k, l \ge 0 \right\},
$$

where Id denotes the identity on  $\mathbb{R}^{\mathbb{R}_+}$ .

By [\(2.3\)](#page-3-0), the elements of C are operators mapping a function  $h \in \mathbb{R}^{\mathbb{R}_+}$  to another function  $c(h) \in \mathbb{R}^{\mathbb{R}_+}$ . We also consider a mapping M, called the *mechanism*, defined on C by matching a code  $c \in \mathcal{C}$  to a set  $\mathcal{M}(c)$  of code tuples.

**Definition 2.2** The mechanism M is defined by  $\mathcal{M}(\text{Id}) = \{f^*\}\$ and

<span id="page-3-2"></span><span id="page-3-1"></span>
$$
\mathcal{M}(g^*) = \{ (\partial_0 g)^*, (f^*, (\partial_1 g)^*) \},
$$
\n(2.4)

for g a smooth function on  $\mathbb{R}_+ \times \mathbb{R}$ .

In the next lemma we show that  $c(y)$  satisfies a system of equations indexed by  $c \in \mathcal{C}$ .

**Lemma 2.3** For any code  $c \in \mathcal{C}$  we have

$$
c(y)(t) = c(y)(0) + \sum_{Z \in \mathcal{M}(c)} \int_0^t \prod_{z \in Z} z(y)(s)ds, \qquad t \in \mathbb{R}_+.
$$
 (2.5)

*Proof.* When  $c = Id$  we have

$$
c(y)(t) = y(t) = y_0 + \int_0^t f(s, y(s))ds = y_0 + \int_0^t f^*(y)(s)ds,
$$

hence [\(2.5\)](#page-3-1) holds since  $\mathcal{M}(\text{Id}) = \{f^*\}.$  When  $c = g^* \in \mathcal{C}$  with  $c \neq \text{Id}$ , the equation

$$
g(t, y(t)) = g(0, y_0) + \int_0^t \partial_0 g(s, y(s)) ds + \int_0^t f(s, y(s)) \partial_1 g(s, y(s))
$$

satisfied by  $g^*(y)(t)$  reads

$$
g^*(y)(t) = g^*(y)(0) + \int_0^t \partial_0 g^*(y)(s)ds + \int_0^t f^*(y)(s)\partial_1 g^*(y)(s),
$$

and  $(2.5)$  follows by the definition  $(2.4)$  of M.

<span id="page-3-3"></span>

We note that for any  $g^* \in \mathcal{C}$  it is always possible to compute  $g^*(y)(0)$  by applying the code g<sup>\*</sup> to the solution y of the ODE and then evaluating it at time 0 as  $g^*(y)(0) = g(0, y_0)$ . In particular, the full knowledge of the function y is not necessary to compute  $g^*(y)(0)$ .

#### Random trees

For each code  $c \in \mathcal{C}$  we denote by  $I_c$  a uniformly distributed random variable on  $\mathcal{M}(c)$ . For example, when  $c = f^*$  we have

$$
\mathbb{P}(I_{f^*} = (f^*, (\partial_1 f)^*)) = \mathbb{P}(I_{f^*} = (\partial_0 f)^*) = \frac{1}{2}.
$$

In the sequel we will use the notation

$$
q_c(b) := \mathbb{P}(I_c = b), \qquad b \in \mathcal{M}(c), \quad c \in \mathcal{C}.
$$

In addition, we consider a probability density function  $\rho : \mathbb{R}_+ \to (0, \infty)$  and

- an i.i.d. family  $(\tau^{i,j})_{i,j\geq 1}$  of random variables with distribution  $\rho(t)dt$  on  $\mathbb{R}_+$ ,
- for each  $c \in \mathcal{C}$ , an i.i.d. family  $(I_c^{i,j})_{i,j\geq 1}$  of discrete random variables, with

$$
\mathbb{P}(I_c^{i,j} = b) = q_c(b) > 0, \qquad b \in \mathcal{M}(c),
$$

where the sequences  $(\tau^{i,j})_{i,j\geq 1}$  and  $(I_c^{i,j})_{c\in\mathcal{C},i,j\geq 1}$  are assumed to be mutually independent. For each  $n \in \mathbb{N}$  we also consider an injection  $\pi_n : \mathbb{N}^n \to \mathbb{N}$ .

#### Coding trees

Starting from time 0, a particle labelled  $\overline{1} := (1)$  bearing the code Id lives up to a random time  $\tau^{1,1}$  distributed according to  $\rho$ . If  $\tau^{1,1} > t$ , the branching process stops. Otherwise, if  $\tau^{1,1} \leq t$ , a new particle with label  $(1, 1)$  is created, and bears the code  $f^*$  since  $\mathcal{M}(\text{Id}) = \{f^*\}.$  This new branch lives during the time  $\tau^{2,\pi_2(1,1)}$ . If  $\tau^{(1,1)} + \tau^{2,\pi_2(1,1)} > t$  the tree stops branching, otherwise, the particle branches into two new offsprings, one bearing the code f and the other one bearing the code  $f'$ .

Otherwise, if  $\tau^{1,1} \leq t$ , a new particle with label  $(1, 1)$  is created, bearing the code  $f^*$ , and independently follows the same pattern as the first one, lives for a time  $\tau^{2,\pi_2(1,1)}$  and so on until the random tree formed by these particles reaches the horizon time t. If  $\tau^{1,1} + \tau^{2,\pi_2(1,1)} > t$ , the tree stops branching. Otherwise, if  $\tau^{1,1} + \tau^{2,\pi_2(1,1)} \leq t$ , the particle branches in two different ways. Namely, with probability 1/2, either

- into two new offsprings, one bearing the code  $f^*$  and the other one bearing the code  $(\partial_1 f)^*$ , or
- into a single offspring bearing the code  $(\partial_0 f)^*$ ,

since  $\mathcal{M}(f^*) = \left\{ (f^*,(\partial_1 f)^*),(\partial_0 f)^* \right\}.$ 

More generally, a particle with code  $c \in \mathcal{C}$  at the generation  $n \geq 1$  is assigned a label of the form  $\bar{k} = (1, k_2, \ldots, k_n) \in \mathbb{N}^n$ , and its parent is labeled  $\bar{k} - := (1, k_2, \ldots, k_{n-1})$ . The birth time of particle  $\bar{k}$  is denoted by  $T_{\bar{k}-1}$ , and its lifetime  $\tau^{n,\pi_n(\bar{k})}$  is the element of index  $\pi_n(\bar{k})$  in the i.i.d. sequence  $(\tau^{n,j})_{j\geq 1}$ . If  $T_{\bar{k}} := T_{\bar{k}-} + \tau^{n,\pi_n(\bar{k})} < t$ , we draw a sample  $I_c^{n,\pi_n(\bar{k})} = (c_1,\ldots,c_l)$  uniformly in  $\mathcal{M}(c)$ , and the particle  $\bar{k}$  branches into  $\left|I_c^{n,\pi_n(\bar{k})}\right|$  offsprings at generation  $(n+1)$ , which are labeled by  $\bar{k} = (1, \ldots, k_n, i)$ ,  $i = 1, \ldots, |I_c^{n, \pi_n(\bar{k})}|$ . The particle with label ending with an integer i will carry the code  $c_i$ . Finally, the code of particle  $\overline{k}$ will be denoted by  $c_{\bar{k}} \in \mathcal{C}$ . Note that the labels are only used to distinguish the particles in the branching process, and we sometimes abusively use the word branch to speak about its label and vice-versa. The set of particles dying before time t is denoted by  $\mathcal{K}^{\circ}$ , whereas those dying after t form a set denoted by  $\mathcal{K}^{\partial}$ . The death time of the particle  $\bar{k}$  is denoted by  $T_{\overline{k}}$ , and its birth time is  $T_{\overline{k}-}$ .

**Definition 2.4** When started at time  $t \in [0, T]$  and a code  $c \in \mathcal{C}$  on its first branch, the above construction yields a marked branching process called a random coding tree, and denoted by  $\mathcal{T}_{t,c}.$ 

The tree  $\mathcal{T}_{t,Id}$  will be used for the stochastic representation of the solution  $y(t)$  of the ODE [\(2.1\)](#page-2-1), while the trees  $\mathcal{T}_{t,c}$  will be used for the stochastic representation of  $c(y)(t)$ . The next table summarizes the notation introduced so far.

Object	Notation
Initial time	
Tree rooted at $t$ with initial code $c$	$\mathcal{T}_{t.c}$
Particle (or label) of generation $n \geq 1$	$\overline{k}=(1,k_2,\ldots,k_n)$
First branching time	$T_{\overline{1}}$
Lifespan of a particle	$T_{\overline{k}}-T_{\overline{k}-}$
Birth time of a particle $k$	$T_{\overline{k}-}$
Death time of a particle $k$	$T_{\overline{k}}$
Code of a particle $k$	$c_{\overline{k}}$

The following is a visual representation of a sample of the random tree  $\mathcal{T}_{t,Id}$  we just created.



We denote by  $K^{\circ}$  the particles that do not reach t and  $K^{\partial}$  the ones reaching t (in this specific case, there is only one particle in  $\mathcal{K}^{\partial}$ . The above construction also allows us to build a coding tree  $\mathcal{T}_{t,c}$  started from any code  $c \in \mathcal{C}$  at any time  $t \geq 0$ .

### <span id="page-6-1"></span>3 Probabilistic representation of ODE solutions

Next, we introduce the random multiplicative functional which will be used to represent ODE solutions.

**Definition 3.1** Given  $\mathcal{T}_{t,c}$  a random coding tree started at time  $t \geq 0$  with code  $c \in \mathcal{C}$ , we define the universal multiplicative functional  $H$  by

<span id="page-6-0"></span>
$$
\mathcal{H}(\mathcal{T}_{t,c}) := \prod_{\overline{k} \in \mathcal{K}^{\circ}} \frac{1}{q_{c_{\overline{k}}}(I_{c_{\overline{k}}}) \rho(T_{\overline{k}} - T_{\overline{k}-})} \prod_{\overline{k} \in \mathcal{K}^{\partial}} \frac{c_{\overline{k}}(y)(0)}{\overline{F}(t - T_{\overline{k}-})}.
$$

The next result gives the probabilistic representation of ODE solutions as an expectation over random coding trees.

**Theorem 3.2** Let  $T > 0$  for which there exists  $K_0 > 0$  such that

$$
\mathbb{E}\big[\big|\mathcal{H}(\mathcal{T}_{t,c})\big|\big]\leq K_0,\qquad c\in\mathcal{C},\quad t\in[0,T].
$$

Then, for any  $c \in \mathcal{C}$  we have the probabilistic representation

<span id="page-6-2"></span>
$$
c(y)(t) = \mathbb{E}\big[\mathcal{H}(\mathcal{T}_{t,c})\big], \qquad t \in [0,T], \tag{3.1}
$$

where y is the solution of the ODE

$$
y(t) = y_0 + \int_0^t f(s, y(s))ds,
$$
  $t \in [0, T].$ 

In particular, taking  $c = Id$ , we have

<span id="page-7-1"></span>
$$
y(t) = \mathbb{E}\big[\mathcal{H}(\mathcal{T}_{t,\text{Id}})\big], \qquad t \in [0,T].
$$

*Proof.* For  $c \in \mathcal{C}$  we let

$$
y_c(t) := \mathbb{E}\big[\mathcal{H}(\mathcal{T}_{t,c})\big], \qquad t \in [0,T].
$$

By conditioning on the first branching time  $T_{\overline{1}}$ , the first particle bearing the code Id branches at time  $T_{\overline{1}}$  into a new particle bearing the code  $f^*$  as  $\mathcal{M}(\mathrm{Id}) = \{f^*\},\$ hence

$$
y_{\text{Id}}(t) = \mathbb{E}\left[\mathcal{H}(\mathcal{T}_{t,\text{Id}})\mathbb{1}_{\{T_{\overline{1}}>t\}} + \mathcal{H}(\mathcal{T}_{t,\text{Id}})\mathbb{1}_{\{T_{\overline{1}}\leq t\}}\right]
$$
  
\n
$$
= \mathbb{E}\left[\frac{y_0}{\overline{F}(t)}\mathbb{1}_{\{T_{\overline{1}}>t\}}\right] + \mathbb{E}\left[\frac{y_{f^*}(t-T_{\overline{1}})}{\rho(T_{\overline{1}})}\mathbb{1}_{\{T_{\overline{1}}\leq t\}}\right]
$$
  
\n
$$
= y_0 \frac{\mathbb{P}(T_{\overline{1}}>t)}{\overline{F}(t)} + \int_0^t \frac{y_{f^*}(t-s)}{\rho(s)}\rho(s)ds
$$
  
\n
$$
= y_0 + \int_0^t y_{f^*}(s)ds, \qquad t \in [0,T].
$$

Similarly, starting from any code  $g^* \in \mathcal{C}$  different from Id, the particle branches at time  $T_1$ into either one particle with code  $(\partial_0 g)^*$  or into two particles with codes  $(f^*, (\partial_1 g)^*)$ , hence we have

$$
y_{g^*}(t) = \mathbb{E}\left[\mathcal{H}(\mathcal{T}_{t,g^*})\mathbb{1}_{\{T_{\overline{1}} > t\}} + \mathcal{H}(\mathcal{T}_{t,g^*})\mathbb{1}_{\{T_{\overline{1}} \leq t\}}\right]
$$
  
\n
$$
= \mathbb{E}\left[\frac{g^*(y_0)}{\overline{F}(t)}\mathbb{1}_{\{T_{\overline{1}} > t\}} + \mathbb{1}_{\{T_{\overline{1}} \leq t\}}\left(\mathbb{1}_{\{I_{g^*} = (\partial_0 g)^* \}}\frac{y_{(\partial_0 g)^*}(t - T_{\overline{1}})}{q_{g^*}(I_{g^*})\rho(T_{\overline{1}})} + \mathbb{1}_{\{I_{g^*} = (g^*, (\partial_1 g)^*)\}}\frac{y_{f^*}(t - T_{\overline{1}})y_{(\partial_1 g)^*}(t - T_{\overline{1}})}{q_{g^*}(I_{g^*})\rho(T_{\overline{1}})}\right)\right]
$$
  
\n
$$
= \mathbb{E}\left[\frac{g^*(y_0)}{\overline{F}(t)}\mathbb{1}_{\{T_{\overline{1}} > t\}} + \mathbb{1}_{\{T_{\overline{1}} \leq t\}}\left(q_{g^*}((\partial_0 g)^*)\frac{y_{(\partial_0 g)^*}(t - T_{\overline{1}})}{q_{g^*}((\partial_0 g)^*)\rho(T_{\overline{1}})} + q_{g^*}((g^*, (\partial_1 g)^*))\frac{y_{f^*}(t - T_{\overline{1}})y_{(\partial_1 g)^*}(t - T_{\overline{1}})}{q_{g^*}((g^*, (\partial_1 g)^*))\rho(T_{\overline{1}})}\right)\right]
$$
  
\n
$$
= \mathbb{E}\left[\frac{g^*(y_0)}{\overline{F}(t)}\mathbb{1}_{\{T_{\overline{1}} > t\}}\right] + \sum_{Z \in \mathcal{M}(g^*)} \mathbb{E}\left[\mathbb{1}_{\{T_{\overline{1}} \leq t\}}\frac{1}{\rho(T_{\overline{1}})}\prod_{z \in Z}y_z(t - T_{\overline{1}})\right]
$$
  
\n
$$
= g^*(y)(0) + \sum_{Z \in \math
$$

which yields a system of equations

<span id="page-7-0"></span>
$$
y_c(t) = c(y)(0) + \sum_{Z \in \mathcal{M}(c)} \int_0^t \prod_{z \in Z} y_z(s) ds, \qquad t \in [0, T], \quad c \in \mathcal{C}.
$$
 (3.3)

By the Cauchy-Lipschitz theorem on the Banach space of sequences  $\ell^{\infty}$ , this system admits a unique maximal solution. We conclude by noting that from Lemma [2.3,](#page-3-3) the family of functions  $(c(y))_{c\in\mathcal{C}}$  is the solution of the system  $(3.3)$ , hence  $(c(y))_{c\in\mathcal{C}} = (y_c)_{c\in\mathcal{C}}$ , and

$$
\mathbb{E}[\mathcal{H}(\mathcal{T}_{t,c})] = y_c(t) = c(y)(t), \qquad t \in [0,T].
$$

#### Integrability condition

The following proposition provides sufficient conditions ensuring that the representation formula [\(3.1\)](#page-6-2) of Theorem [3.2](#page-6-0) holds at any time within certain interval. In order to represent an ODE solution beyond that interval we may reuse a numerical value obtained close the boundary as a new initial value in order to represent the solution on an extended time interval. The error on numerical values can be controlled by the Central Limit theorem.

**Proposition 3.3** Assume that there exists  $K > 0$  such that  $c(y)(0) \leq K$  for any  $c \in \mathcal{C}$ . Then, for any  $T > 0$  such that

<span id="page-8-2"></span><span id="page-8-1"></span><span id="page-8-0"></span>
$$
T < \log\left(1 + \frac{1}{K}\right),\tag{3.4}
$$

or such that the density function  $\rho$  is nonincreasing and

<span id="page-8-3"></span>
$$
\rho(T) \ge 2, \quad K \le \overline{F}(T),\tag{3.5}
$$

there exists  $K(T) > 0$  such that

$$
\mathbb{E}\big[\big|\mathcal{H}(\mathcal{T}_{s,c})\big|\big] \leq K(T), \quad c \in \mathcal{C}, \quad t \in [0,T].\tag{3.6}
$$

Proof. We have

$$
\mathbb{E}\big[\big|\mathcal{H}(\mathcal{T}_{t,c})\big|\big]\leq S_c(t):=\mathbb{E}_{t,c}\Bigg[\prod_{\overline{k}\in\mathcal{K}^{\circ}}\frac{1}{q_{c_{\overline{k}}}(I_{c_{\overline{k}}})\rho(\tau_{\overline{k}})}\prod_{\overline{k}\in\mathcal{K}^{\partial}}\frac{K}{\overline{F}(t-T_{\overline{k}-})}\Bigg].
$$

We note that we have  $S_c(t) = S_{f^*}(t)$  for all  $c \in \mathcal{C}$  of the form  $c = (\partial_0^k \partial_1^l f)^*, k, l \ge 0$ , since the corresponding trees have the same distribution. Therefore, the system of equations satisfied by  $S_c(t)$  may be written as

$$
\begin{cases}\nS_{\text{Id}}(t) = K + \int_0^t S_{f^*}(s)ds \\
S_{f^*}(t) = K + \int_0^t (S_{f^*}(s))^2 ds + \int_0^t S_{f^*}(s)ds,\n\end{cases}
$$

which can be solved as

$$
S_{f^*}(t) = \frac{Ke^t}{1 + K(1 - e^t)},
$$
  $S_{\text{Id}}(t) = K - \log(1 + K(1 - e^t)).$ 

Therefore, integrability holds when

$$
0 \le t \le T < \log\left(1 + \frac{1}{K}\right).
$$

Under Condition [\(3.5\)](#page-8-1), since  $q_{\min} := \min_{c \in \mathcal{C}} q_c(I_c) = 1/2$ , we have

$$
\prod_{\overline{k}\in \mathcal{K}^\circ}\frac{1}{q_{\min}\rho(T_{\overline{k}}-T_{\overline{k}-})}\prod_{\overline{k}\in \mathcal{K}^\partial}\frac{K}{\overline{F}(t-T_{\overline{k}-})}\leq 1,
$$

hence

$$
\mathbb{E}\big[\big|\mathcal{H}(\mathcal{T}_{t,c})\big|\big] \leq \mathbb{E}_{t,c}\left[\prod_{\overline{k}\in\mathcal{K}^{\circ}}\frac{1}{q_{\min}\rho(T_{\overline{k}}-T_{\overline{k}-})}\prod_{\overline{k}\in\mathcal{K}^{\partial}}\frac{K}{\overline{F}(t-T_{\overline{k}-})}\right] \leq 1, \qquad t\in[0,T].
$$

 $\Box$ 

In the case of a higher dimensional autonomous system of the form

$$
\begin{cases}\ny^1(t) = y_0^1 + \int_0^t f_1(y^1(s), \dots, y^d(s)) ds \\
y^2(t) = y_0^2 + \int_0^t f_2(y^1(s), \dots, y^d(s)) ds \\
\vdots \\
y^d(t) = y_0^d + \int_0^t f_n(y^1(s), \dots, y^d(s)) ds\n\end{cases}
$$

it is possible to create n different coding trees, where each tree codes for a different  $y^i$  for  $i \in \{1, \ldots, n\}$ . In this case, the set of codes  $C$  is defined as

$$
\mathcal{C} := \left\{ \mathrm{Id}_i, \ \partial_1^{i_1} \cdots \partial_d^{i_d} f_i \ : \ i_1, \ldots, i_d \geq 0, \ i = 1, \ldots, d \right\}
$$

and the mechanism is obtained using the chain rule and is defined as

$$
\mathcal{M}(\mathrm{Id}_i)=\{f_i\},\quad \mathcal{M}(g)=\big\{(f_1,\partial_1g),(f_2,\partial_2g),\ldots,(f_d,\partial_dg)\big\}.
$$

and by definition we let  $\mathrm{Id}_i(y_0^1,\ldots,y_0^d) := y_0^i$ . By an argument similar to  $(3.2)$ , for  $c \in \mathcal{C}$  we have

$$
y_c(t) := \mathbb{E}\big[\mathcal{H}(\mathcal{T}_{t,c})\big] = c(y)(0) + \sum_{Z \in \mathcal{M}(c)} \int_0^t \prod_{z \in Z} y_z(s)ds
$$

and  $y_{\text{Id}_i}(t) = y^i(t)$ ,  $i = 1, \ldots, d$ . In addition, any non autonomous system can be transformed into an autonomous system by addition of a dimension. In particular, any higher order ordinary differential equation of the form

$$
y^{(d)}(t) = f(t, y(t), y'(t), \dots, y^{(d-1)}(t))
$$

can be written as a system of the above form by taking  $f_i(x_0, x_1, \ldots, x_d) := x_i, \ i =$  $0, 1, \ldots, d-1$  and  $f_d(x_0, x_1, \ldots, x_d) := f(x_0, x_1, \ldots, x_d)$ , with  $y^0(t) = t$  and  $y^i(t) = y^{(i-1)}(t)$ ,  $i=1,\ldots,d.$ 

### <span id="page-10-0"></span>4 Examples

#### Exponential series

We first consider the equation

<span id="page-10-1"></span>
$$
\begin{cases} y'(t) = y(t) \\ y(0) = y_0 \end{cases}
$$

rewritten in integral form as

$$
y(t) = y_0 + \int_0^t y(s)ds, \quad t \in \mathbb{R}_+,
$$

whose solution admits the power series expansion

$$
y(t) = y_0 e^t = y_0 \sum_{n=0}^{\infty} \frac{t^n}{n!}, \qquad t \in \mathbb{R}_+.
$$

Here we have  $C = \{Id\}$ , and the mechanism M satisfies  $\mathcal{M}(Id) = \{Id\}$ . The particle of generation *n* bears the label  $\overline{k} = (1, \ldots, 1) \in \mathbb{N}^n$ , and its parent is the particle  $\overline{k-} =$  $(1,\ldots,1) \in \mathbb{N}^{n-1}$ . When the random times  $(\tau^n)_{n\geq 1}$  are independent and exponentially distributed, i.e.  $\rho(s) = e^{-s}\mathbb{1}_{[0,\infty)}(s)$  with  $\overline{F}(t) = e^{-t}$ , the total number of branches in the random tree  $\mathcal{T}_t$  is given by  $N_t + 1$  where  $(N_t)_{t \geq 0}$  is a standard Poisson process with unit intensity.

$$
\begin{array}{c|c}\n\overline{1} & (1,1) \\
\hline\n\text{Id} & \overline{1} \\
\end{array}\n\qquad\n\begin{array}{c|c}\n(1,1) & (1,1,1) \\
\hline\n\text{Id} & \text{Id}\n\end{array}\n\qquad\n\begin{array}{c|c}\n(1,1,1,1) & (1,1,1,1) \\
\hline\n\text{Id} & \text{Id}\n\end{array}
$$

In this case, the multiplicative functional

$$
\mathcal{H}(\mathcal{T}_t) := \prod_{k \in \mathcal{K}^{\circ}} \frac{1}{\rho(T_{\overline{k}} - T_{\overline{k}-})} \prod_{k \in \mathcal{K}^{\partial}} \frac{y_0}{\overline{F}(t - T_{\overline{k}-})}
$$

simplifies to the deterministic expression

$$
\mathcal{H}(\mathcal{T}_t) = y_0 e^{t - \sum_{n=1}^{N_t} \tau^n} \prod_{n=1}^{N_t} e^{\tau^n} = y_0 e^t, \qquad t \in \mathbb{R}_+,
$$

in which we take  $\sum_{n=1}^{0} 1 := 0$  and  $\prod_{n=1}^{0} 1 := 1$ .

#### Autonomous ODEs

Consider the autonomous equation

$$
y(t) = y_0 + \int_0^t f(y(s))ds, \qquad t \in \mathbb{R}_+,
$$
\n(4.1)

where  $f \in C^{\infty}(\mathbb{R}; \mathbb{R})$  is bounded together with its derivatives  $f^{(k)}$  of order  $k \geq 1$ , with

<span id="page-11-0"></span>
$$
\left|f^{(k)}(y_0)\right|_{\infty} \le K, \qquad k \ge 0,\tag{4.2}
$$

for some  $K > 0$ . For any  $k \geq 0$  we have the integral equation

$$
f^{(k)}(y(t)) = f^{(k)}(y_0) + \int_0^t f(y(s))f^{(k+1)}(y(s))ds, \qquad t \in \mathbb{R}_+,
$$

and the set of codes is given by

$$
\mathcal{C}:=\big\{\mathrm{Id},\ f^{(k)},\ k\geq 1\big\},
$$

where  $f^{(k)}$ ,  $k \geq 0$ , denotes the operator acting as

$$
f^{(k)}(y)(s) := (f^{(k)}(y))(s), \qquad s \in \mathbb{R}.
$$

In this case, the mechanism  $\mathcal M$  is defined by

$$
\mathcal{M}(\mathrm{Id}) := \{f\}, \quad \mathcal{M}(f) := \{(f, f')\}, \quad \mathcal{M}(f^{(k)}) := \{(f, f^{(k+1)})\}, \qquad k \ge 1.
$$

Below is a visual representation of a sample of the random tree  $\mathcal{T}_{t,Id}$ .



Here, the multiplicative functional on the random tree  $\mathcal{T}_{t,Id}$  is given by

<span id="page-12-0"></span>
$$
\mathcal{H}(\mathcal{T}_{t,Id}) := \prod_{\overline{k} \in \mathcal{K}^{\circ}} \frac{1}{\rho(T_{\overline{k}} - T_{\overline{k}-})} \prod_{\overline{k} \in \mathcal{K}^{\partial}} \frac{c_{\overline{k}}(y)(0)}{\overline{F}(t - T_{\overline{k}-})}
$$

.

We note that the above product is well defined, and in particular we are able to compute  $c_{\overline{k}}(y)(0) = c_{\overline{k}}(y_0)$  by interpreting the operator  $c_{\overline{k}}$  as a function from  $\mathbb R$  to  $\mathbb R$ . With  $y_c(t) :=$  $\mathbb{E}[\mathcal{H}(\mathcal{T}_{t,c})], c \in \mathcal{C}$ , the system  $(3.2)$  reads

$$
\begin{cases}\ny_{\text{Id}}(t) = y_0 + \int_0^t y_f(s)ds \\
y_{f^{(k)}}(t) = f^{(k)}(y_0) + \int_0^t y_f(s)y_{f^{(k+1)}}(s)ds, \quad k \ge 0.\n\end{cases}
$$

By [\(4.2\)](#page-11-0), for any code  $c \in \mathcal{C}$  we have

$$
y_c(t) \leq S_c(t) := \mathbb{E}_c \left[ \prod_{\overline{k} \in \mathcal{K}^\circ} \frac{1}{\rho(T_{\overline{k}} - T_{\overline{k}-})} \prod_{\overline{k} \in \mathcal{K}^\partial} \frac{K}{\overline{F}(t - T_{\overline{k}-})} \right],
$$

where the right-hand side of the above inequality is solution of

$$
\begin{cases}\nS_{\text{Id}}(t) = K + \int_0^t S_f(s)ds \\
S_{f^{(k)}}(t) = K + \int_0^t S_f(s)S_{f^{(k+1)}}(s)ds, \qquad k \ge 0,\n\end{cases}
$$
\n(4.3)

and by construction we have  $S_f = S_{f^{(k)}}, k \ge 0$ , since the corresponding trees starting with the codes f and  $f^{(k)}$ ,  $k \geq 1$ , have the same distribution. Thus, [\(4.3\)](#page-12-0) reduces to

$$
\begin{cases}\nS_{\text{Id}}(t) = K + \int_0^t S_f(s)ds \\
S_f(t) = K + \int_0^t S_f^2(s)ds, \quad k \ge 0,\n\end{cases}
$$

which yields

$$
S_{\text{Id}}(t) = K - \log(1 - Kt)
$$
 and  $S_f(t) = \frac{K}{1 - Kt}$ .

Hence the probabilistic representation  $(3.1)$  holds if

<span id="page-12-1"></span>
$$
T < \frac{1}{K},\tag{4.4}
$$

which is weaker than Condition  $(3.4)$  in Proposition [3.3.](#page-8-0) Another way to ensure the integrability of  $\mathcal{H}(\mathcal{T}_{t,Id})$  is to choose a decreasing probability density function  $\rho(t)$  satisfying the condition

$$
\rho(T) \ge 1
$$
 and  $K \le \overline{F}(T)$ ,

which is also weaker than  $(3.5)$ .

### <span id="page-13-0"></span>5 Coding trees vs Butcher trees

In this section we describe the connection between coding trees, Butcher series and Butcher trees, by showing bow any Butcher tree can be recovered by performing a depth first search on the corresponding coding tree.

#### Autonomous case

In the case of the autonomous ODE

$$
y(t) = y_0 + \int_0^t f(y(s))ds, \qquad t \in \mathbb{R}_+,
$$

we write the Taylor expansion of  $y(t)$  as the Butcher series

$$
y(t) = y_0 + tf(y_0) + \frac{1}{2}t^2 f(y_0) f'(y_0)
$$
  
+ 
$$
\frac{1}{3!}t^3 (f''(y_0) f^2(y_0) + f'^2(y_0) f(y_0))
$$
  
+ 
$$
\frac{1}{4!}t^4 (f'''(y_0) f^3(y_0) + f''(y_0) f'(y_0) f^2(y_0) + f'(y_0) f'(y_0) f^2(y_0) + f'^3(y_0) f(y_0)) + \cdots
$$
  
= 
$$
y_0 + \sum_{\mathcal{B}} \frac{t^{|\mathcal{B}|}}{\nu(\mathcal{B})} c(\mathcal{B})
$$

where the summation is over Butcher trees, the order  $|\mathcal{B}|$  denotes the number of vertices of the tree B,  $c(\mathcal{B}, f, y_0)$  is a term depending on the derivatives of f at  $y_0$ , and  $\nu(\mathcal{B})$  is a coefficient which is defined recursively, see (4) in [\[But10\]](#page-21-16). In the following tables we show how Butcher trees  $\beta$  in the above sum can be identified to a coding trees  $\mathcal{T}$ .

















Namely, every Butcher tree can be recovered by performing a depth first search on the corresponding coding tree by matching leaves on Butcher trees to branches of the same color in coding trees.

If a node in the initial coding tree branches into two new coding trees  $\mathcal{T}_1$  (above) and  $\mathcal{T}_2$ (below), the Butcher tree for the initial coding tree is obtained by sticking the Butcher tree obtained from  $\mathcal{T}_1$  to the root of the Butcher tree obtained from  $\mathcal{T}_2$ . If a tree is a leaf then the Butcher tree obtained is simply a node with the code of the leaf.

#### Non-autonomous case

In the case of a non autonomous ODE of the form

$$
y(t) = y_0 + \int_0^t f(s, y(s))ds
$$
,  $t \in \mathbb{R}_+$ ,

multiple coding trees are needed to represent a given Butcher tree. For example, the following Butcher tree



is represented using the following two coding trees:



On the other hand, the following Butcher tree



will be represented using the following four coding trees:



# <span id="page-16-0"></span>6 Numerical application

i) Taking  $f(y) := y^2$ , We start with the quadratic ODE

<span id="page-16-1"></span>
$$
y'(t) = y2(t), \t y(0) = y0 > 0, \t (6.1)
$$

with solution

$$
y(t) = \frac{y_0}{1 - y_0 t}, \qquad t \in [0, 1/y_0).
$$

In the framework of  $(4.1)$  we have

$$
\mathcal{C} = \{ \text{Id}, \ f^{(k)}, \ k \ge 0 \} = \{ \text{Id}, \ x \mapsto 0, \ x \mapsto 2, \ x \mapsto 2x, \ x \mapsto x^2 \},
$$

<span id="page-17-0"></span>hence we have  $c(y)(0) \leq K := \max(1, 2y_0, y_0^2)$  for all  $c \in \mathcal{C}$ , hence by Proposition [3.3](#page-8-0) and [\(4.4\)](#page-12-1) the representation formula [\(3.1\)](#page-6-2) of Theorem [3.2](#page-6-0) holds for all  $t \in [0, 1/2)$ , see Figure [1.](#page-17-0) The following figures are plotted with one million Monte Carlo samples.



Figure 1: Numerical solution of  $(6.1)$  with  $y_0 = 1$ .

In this example and the next one we take  $\rho$  to be the exponential probability density function  $\rho(t) = e^{-t}, t \geq 0$ .

ii) Next, we take  $f(y) := \cos(y)$  and consider the equation

<span id="page-17-1"></span>
$$
y'(t) = \cos(y(t)), \quad y(0) = y_0,\tag{6.2}
$$

with solution

$$
y(t) = 2 \tan^{-1} \left( \tanh \left( \frac{t + 2 \tanh^{-1}(\tan(y_0/2))}{2} \right) \right), \quad t \in \mathbb{R}_+.
$$

in the framework of [\(4.1\)](#page-10-1). When  $y_0 = 1$  we have  $\sup_{k \in \mathbb{N}} f^{(k)}(1) = 1$  hence by Proposi-tion [3.3](#page-8-0) and [\(4.4\)](#page-12-1) the representation formula [\(3.1\)](#page-6-2) of Theorem [3.2](#page-6-0) holds for all  $t \in [0, 1)$ , see Figure [2.](#page-18-0)

<span id="page-18-0"></span>

Figure 2: Numerical solution of [\(6.2\)](#page-17-1).

<span id="page-18-3"></span>iii) Taking  $f(t, y) := (y + t)/(y - t)$  we find Equation (201a) in [\[But16\]](#page-21-1), i.e.

$$
y'(t) = \frac{y(t) + t}{y(t) - t}, \qquad y(0) = 1,
$$
\n(6.3)

with solution

<span id="page-18-2"></span>
$$
y(t) = t + \sqrt{1 + 2t^2}.
$$

<span id="page-18-1"></span>In this case, the time interval of validity may not be determined explicitly because  $\sup_{k,l\geq 0} |\partial_0^k \partial_1^l f(0,1/2)| = \infty$  and the finiteness of this supremum is only a sufficient condition for [\(3.6\)](#page-8-3) to hold in Proposition [3.3.](#page-8-0) Figure [3](#page-18-1) shows the convergence of the Monte Carlo algorithm until  $t = 0.25$ .



Figure 3: Numerical solution of [\(6.3\)](#page-18-2).

iv) Taking  $f(t, y) := (y - t)/(y + t)$  yields Equation (316e) in [\[But16\]](#page-21-1), i.e.

<span id="page-18-4"></span>
$$
y'(t) = \frac{y(t) - t}{y(t) + t}, \qquad y(0) = 1,\tag{6.4}
$$

<span id="page-19-0"></span>whose solution is given in parametric form as  $(t(u), y(u)) = (u \sin \log(u), u \cos \log(u)).$ As in Example [iii\)](#page-18-3) above, the time interval of validity may not be determined explicitly, see Figure [4.](#page-19-0)



Figure 4: Numerical solution of [\(6.4\)](#page-18-4).

In this example and the next one we take  $\rho$  to be the gamma probability density function  $\rho(t) = t^{-1/2} e^{-t} / \Gamma(1/2), t \ge 0.$ 

v) Taking  $f(t, y) := yt + y^2$  we find Equation (223a) in [\[But16\]](#page-21-1), i.e.

$$
y'(t) = ty(t) + y2(t), \t y(0) = 1/2,
$$
\t(6.5)

with solution

<span id="page-19-1"></span>
$$
y(t) = \frac{e^{t^2/2}}{2 - \int_0^t e^{s^2/2} ds}.
$$

In this case we have  $\sup_{k,l\geq 0} |\partial_0^k \partial_1^l f(0,1/2)| \leq 2$ , hence by Proposition [3.3](#page-8-0) and [\(4.4\)](#page-12-1) the representation formula  $(3.1)$  of Theorem [3.2](#page-6-0) is valid on the time interval  $[0, 0.5)$ , see Figure [5.](#page-20-0)

<span id="page-20-0"></span>

Figure 5: Numerical solution of [\(6.5\)](#page-19-1).

In addition, after running the algorithm on the time interval  $[0, 0.5]$  we may reuse the numerical evaluation at time  $t = 0.5$  as a new initial condition and iterate the algorithm on the time interval [0.5, 1] with a better stability of estimates, as shown in Figure [5.](#page-20-0)

## A Maple and Mathematica codes

The following codes implement the algorithm of Theorem [3.2](#page-6-0) using an exponential distribution  $\rho(t) = e^{-t}, t \geq 0$ .

```
codetofunction := proc(f, c, t0, y0) if nops(c) = 0 then return y0; end if; if c = [0, 0] then return
    f(t0, y0); else return eval(eval(diff(f(t, y), t \f(c[1], y \f(c[2]), t = t0), y = y0; end if; end
    proc;
mcsample := proc(f, t, t0, y0, c, h) local A, tau; tau := random[exponential[1]](1); if t - t0 < tau thenreturn h*codetofunction(f, c, t0, y0)/exp(-t + t0); else if nops(c) = 0 then return mcsample(f, t -
    tau, t0, y0, [0, 0], h/exp(-tau)); else if random[uniform](1) < 0.5 then return mcsample(f, t - tau,
    t0, y0, [c[1] + 1, c[2]], 2*h/exp(-tau)); else A := mcsample(f, t - tau, t0, y0, [0, 0], 1); return
    mcsample(f, t - tau, t0, y0, [c[1], c[2] + 1], 2*A*h/exp(-tau)); end if; end if; end if; end proc;
solution := proc(f, t, t0, y0, n) local i, temp; temp := 0; for i to n do temp := temp + mcsample(f, t,
    t0, y0, [], 1); end do; return evalf(temp/n); end proc;
f := (t, y) -> y^2; solution(f, 0.5, 0, 1, 10000);
```
Maple code.

```
\text{codetofunction}[f_-, c_-, t0_-, y0_+] := (\text{If } [c == {} \{\}, \text{ Return } [y0],Return[D[D[f[t, y], {t, c[[1]]}], {y, c[[2]]}] /. {t -> t0} /. {y -> y0}]])
MCSample[f_-, t_-, t_0, y_0, c_-, h_+] := (Module[fA, tau], tau = RandomVariable[Exponential Distribution[1]];If [tau > t - t0, Return [h*codetofunction[f, c, t0, y0]/E^(-(t - t0))],
   If [c == \{\}, Return[MCsample[f, t - tau, t0, y0, {0, 0}, h/E^(-tau)]],
     If[RandomVariate[UniformDistribution[1]][[1]] <= 0.5,
      Return[MCsample[f, t - tau, t0, y0, {c[[1]] + 1, c[[2]]}, 2*h/E^(-tau)]],A = MCsample[f, t - tau, t0, y0, \{0, 0\}, 1];Return[MCsample[f, t - tau, t0, y0, {c[[1]], c[[2]] + 1}, 2*A*h/E^(-tau)]]]]])
Solution[f_-, t_-, t_-, y_-, n_+] := (temp = 0; For[i = 1, i \le n, i^+, \text{temp} += \text{MCsample}[f, t, t_0, y_0, \{\},1]]; Return[temp/n])
f[t_-, y_+] := y^2;Solution[f, 0.5, 0, 1, 100000]
```
Mathematica code.

# References

- <span id="page-21-0"></span>[But63] J.C. Butcher. Coefficients for the study of Runge-Kutta integration processes. J. Austral. Math. Soc., 3:185–201, 1963.
- <span id="page-21-16"></span>[But10] J.C. Butcher. Trees and numerical methods for ordinary differential equations. Numerical Algorithms, 53:153–170, 2010.
- <span id="page-21-1"></span>[But16] J.C. Butcher. Numerical methods for ordinary differential equations. John Wiley & Sons, Ltd., Chichester, third edition, 2016.
- <span id="page-21-4"></span>[Cay57] A. Cayley. On the theory of the analytical forms called trees. Philosophical Magazine, 13(85):172–176, 1857.
- <span id="page-21-10"></span>[CLM08] S. Chakraborty and J.A. L´opez-Mimbela. Nonexplosion of a class of semilinear equations via branching particle representations. Advances in Appl. Probability, 40:250–272, 2008.
- <span id="page-21-2"></span>[DB02] P. Deuflhard and F. Bornemann. Scientific Computing with Ordinary Differential Equations, volume 42 of Texts in Applied Mathematics. Springer-Verlag, New York, 2002.
- <span id="page-21-11"></span>[HLOT+19] P. Henry-Labord`ere, N. Oudjane, X. Tan, N. Touzi, and X. Warin. Branching diffusion representation of semilinear PDEs and Monte Carlo approximation. Ann. Inst. H. Poincaré Probab. Statist., 55(1):184–210, 2019.
- <span id="page-21-6"></span>[INW69] N. Ikeda, M. Nagasawa, and S. Watanabe. Branching Markov processes I, II, III. J. Math. Kyoto Univ., 8-9:233–278, 365–410, 95–160, 1968-1969.
- <span id="page-21-9"></span>[LM96] J.A. López-Mimbela. A probabilistic approach to existence of global solutions of a system of nonlinear differential equations. In Fourth Symposium on Probability Theory and Stochastic Processes (Spanish) (Guanajuato, 1996), volume 12 of Aportaciones Mat. Notas Investigación, pages 147–155. Soc. Mat. Mexicana, México, 1996.
- <span id="page-21-14"></span>[Maz04] C. Mazza. Simply generated trees, B-series and Wigner processes. Random Structures Algorithms, 25(3):293–310, 2004.
- <span id="page-21-8"></span>[McK75] H.P. McKean. Application of Brownian motion to the equation of Kolmogorov-Petrovskii-Piskunov. Comm. Pure Appl. Math., 28(3):323–331, 1975.
- <span id="page-21-3"></span>[MMMKV17] R.I. McLachlan, K. Modin, H. Munthe-Kaas, and O. Verdier. Butcher series: a story of rooted trees and numerical methods for evolution equations. Asia Pac. Math. Newsl., 7(1):1–11, 2017.
- <span id="page-21-7"></span>[NS69] M. Nagasawa and T. Sirao. Probabilistic treatment of the blowing up of solutions for a nonlinear integral equation. Trans. Amer. Math. Soc., 139:301–310, 1969.
- <span id="page-21-12"></span>[PP21] G. Penent and N. Privault. Existence and probabilistic representation of the solutions of semilinear parabolic PDEs with fractional Laplacians. To appear in Stochastics and Partial Differential Equations: Analysis and Computations, 30 pages, 2021.
- <span id="page-21-13"></span>[SH21] Z. Selk and H. Harsha. A Feynman-Kac type theorem for ODEs: Solutions of second order ODEs as modes of diffusions. Preprint arXiv:2106.08525, 16 pages, 2021.
- <span id="page-21-15"></span>[Ski92] J. Skilling. Bayesian solution of ordinary differential equations. In C. R. Smith, G. J. Erickson, and P. O. Neudorfer, editors, Proceedings of the Eleventh International Workshop on Maximum Entropy and Bayesian Methods of Statistical Analysis, volume 50 of Fundamental Theories of Physics, pages 23–38. Kluwer Academic Publishers Group, Dordrecht, 1992.
- <span id="page-21-5"></span>[Sko64] A.V. Skorokhod. Branching diffusion processes. Teor. Verojatnost. i. Primenen., 9:492–497, 1964.