# The storage effect is not about bet-hedging or population stage-structure

Evan C. Johnson<sup>1,2,\*</sup> and Alan Hastings<sup>1</sup>

<sup>1</sup>, Department of Environmental Science and Policy; University of California Davis; Davis, California 95616 USA <sup>2</sup>, Center for Population Biology; University of California Davis; Davis, California 95616 USA \* Corresponding author: Evan Johnson, evcjohnson@ucdavis.edu

January 19, 2022

# Abstract

The storage effect is a well-known explanation for the coexistence of competing species in temporally varying environments. Like many complex ecological theories, the storage effect is often used as an explanation for observed coexistence on the basis of heuristic understanding, rather than careful application of a detailed model. But, a careful examination of some widely employed heuristic representations of the storage effect shows that they can be misleading. One interpretation of the storage effect states that species coexist by specializing on a small slice of all environmental states, and therefore must have a robust life-stage (e.g., long-lived adults, a seedbank) in order to "wait it out" for a good year. Another more general interpretation states that *buffering* helps species "wait it out", where *buffering* means populations are protected from large losses when the environment is poor and competition is high. Here, we show that both of these conventional interpretations are imperfect. Multiple models show that stage-structure, long lifespans, and overlapping generations are not required for the storage effect. Further, a species that experiences buffering necessarily grows poorly when the environment is favorable and competition is high. We review the empirical literature and conclude that species are likely to find themselves in this good-environment/high-competition scenario, and therefore, that buffering tends to hurt individual species. The buffering interpretation of the storage effect can be thought of as conflating a community-level criterion for coexistence with a population-level criterion for persistence; it is like claiming that species can persist by limiting their own growth rates, since the Lotka-Volterra model tells us that intraspecific competition must be greater than interspecific competition. To build a better understanding of the storage effect, we examine how it manifests in particular models and we demonstrate the connections between the mathematical and verbal/textual descriptions of the storage effect. While it is possible to improve one's understanding of the storage effect, the storage effect is a very general phenomenon, and so a simple ecological interpretation (in terms of a small set of life-history characteristics like stage-structure or dormancy) will not be forthcoming.

Keywords: storage effect, coexistence mechanisms, modern coexistence theory, bet-hedging, stagestructure, dormancy

## Contents

1	Introduction	3
2	<ul> <li>A crash course on the storage effect</li> <li>2.1 A simple interpretation of the storage effect</li></ul>	<b>3</b> 6 9 10 10 12
3	A critique of the conventional interpretations of the storage effect3.1The origin of the conventional interpretation3.2The perpetuation of the conventional interpretation	<b>13</b> 16 16
4	Discussion	17
5	Acknowledgements	18
6	Appendixes         6.1       The storage effect in the lottery model         6.2       An alternative definition of the storage effect	<b>18</b> 18 20
7	References	20

# 1 Introduction

The temporal storage effect (often simply called the storage effect) is a general explanation for how species can coexist by specializing on different states of a temporally fluctuating environment. The storage effect has an impressive resume. First, it formalized the concept of environmental niche partitioning, which has long been thought (e.g., Grinell, 1917) to promote coexistence. Second, by showing that many species can coexist on just a single resource (Chesson, 1994, Eq. 91; Miller and Klausmeier, 2017), the storage effect provided a potential resolution to Hutchinson's (1961) paradox of the plankton. Third, the storage effect revealed the beneficent side of environmental variation, which historically was thought to undermine the ability of species to persist (Lewontin and Cohen, 1969; May, 1974). A number of recent papers have synthesized the simultaneously stabilizing and destabilizing effects of environmental variation (Adler and Drake, 2008; Schreiber et al., 2019; Pande et al., 2020, Dean and Shnerb, 2020).

Unfortunately, the storage effect is not easy to understand. This is partially because the storage effect is inherently complicated, with a full explanation involving abstractions like statistical interaction effects and cross-species comparisons. One limitation to understanding is the fact that there are three ways to describe the storage effect — a mathematical definition, a list of conditions, and the verbal/textual interpretation — and it is not clear how these descriptions relate to each other. In Section 2, we provide an explanation of the storage effect that explicitly relates the three ways of describing the storage effect.

It is worth belaboring the distinction between the *definition* and *interpretation* of the storage effect. The *definition* is codified in a mathematical formula, which shows how the storage effect can be quantified. The *interpretation* is a verbal or textual story about how species coexist via the storage effect, and serves several important purposes. 1) It imbues the math with biological meaning. 2) It is useful in data-poor situations; it allows one to hypothesize about whether or not the storage effect is operating in a specific system, without having to analyze a model or conduct a pilot experiment. 3) It is not always clear how the pieces of the mathematical definition of the storage effect should be defined (specifically  $E_j$ ,  $C_j$ ,  $E_j^*$ , and  $C_j^*$ ; read on to Section 2.2). The interpretation of the storage effect can provide guidance for making these choices, which in turn can inform experimental design.

The storage effect is often given an interpretation — which we will call the conventional interpretation v1 — paraphrased as follows: species coexist by specializing on different states of a temporally fluctuating environment, so species must have a robust life stage in order to "wait it out" for a favorable time period; "Storage" refers to a robust life stage can "wait it out". The conventional interpretation v1 is incorrect, or at the very least, imprecise. One can construct models where the storage effect is positive, despite there being no stage structure (e.g., Loreau, 1989; Loreau, 1992; Klausmeier, 2010; Li and Chesson, 2016; Letten et al., 2018; Schreiber, 2021). Despite these counterexamples, the imprecision of the conventional interpretation v1 is not widely recognized. In Section 3, we explain why the storage effect has nothing to do with storing population vitality in a robust life stage.

Some authors present a more general interpretation, which we will call the conventional interpretation v2: "Storage" can be more generally understood as *buffering*, which is a negative interaction effect (analogous to a negative coefficient for an interaction term in multiple regression) of environment and competition on per capita growth rates; this buffering helps out rare species because it prevents extreme losses when the environment is unfavorable and competition is high. In Section 3, we explain why the conventional interpretations v2 is also imprecise. We also speculate on the reasons for the genesis and perpetuation of these interpretations (Sections 3.1 & 3.2), which serves as a geneological debunking argument (Srinivasan, 2015; Kahane, 2011), i.e., undermining a proposition by showing that common belief in said proposition can be explained by historical or psychological factors.

# 2 A crash course on the storage effect

Here, we provide a comprehensive description of the storage effect. This section builds the groundwork for our critique of the conventional interpretations of the storage effect, but experts may skip to Section 3).

The storage effect is well-demonstrated with a toy model of coral reef fish dynamics. The *lottery hypothesis* (Sale, 1977) states that the local diversity of coral reef fishes is generated by the random allocation of space: when an adult fish dies, the various fish species enter a lottery for the open territory with a number of tickets equal to the number of larvae that each fish species produces. The lottery hypothesis was motivated by the fact that coral reef fishes do not appear to finely partition food types, but do appear to be limited by space. Space limitation is evidenced by the observed territoriality of adults (Warner and Hoffman, 1980), the production of larvae in massive numbers, and the weak correlation between adult population size and the subsequent number of recruits (Cushing, 1971, Szuwalski et al., 2015).

While Sale's lottery hypothesis does a fine job at explaining local biodiversity, it cannot explain the maintenance of biodiversity — coexistence. Chesson and Warner (1981) were able to attain coexistence with the addition of a single feature: temporal variability in per capita larval production. The resulting model is now known as the *lottery model*, and the more general process permitting coexistence is known as *the storage effect*. The exposition here follows the lottery model of Chesson (1994), as opposed to original lottery model (Chesson and Warner, 1981), which is more complex due to stochasticity in both per capita adult mortality and larval production.

Imagine a guild of fish species inhabiting discrete territories on a coral reef. Several events occur in each time-step of the lottery model, here presented in chronological order

- 1. The fish spawn. Per capita larval-production, (i.e., per capita fecundity) fluctuates from timestep to time-step, putatively due to dependency on environmental factors that also fluctuate. Like the larvae of many marine fish, our hypothetical larvae disperse offshore (ostensibly to avoid predation) and return some time later, though still within the time-step.
- 2. Adult fish die with some density-independent probability, leaving behind an empty territory. The death probability may vary across species, but unlike fecundity, does not vary across time.
- 3. The larvae return to the reef and inherit the empty territories with a recruitment probability for any given larva being equal to the number of empty sites, divided by the total number of larvae. This uniform probability of per larva recruitment is the *lottery* in the lottery model. The unrecruited larvae die before the next time-step begins.

The above dynamics are expressed in the difference equations,

$$n_{j}(t+1) = n_{j}(t) \begin{bmatrix} \text{survival prob.} & \text{per capita fecundity} \\ \widehat{s_{j}} & + & \widehat{\eta_{j}(t)} \end{bmatrix} \begin{bmatrix} \overbrace{\sum_{j\neq i}^{S}(1-s_{j})n_{j}(t)}^{\text{open territories}} \\ \overbrace{\sum_{j\neq i}^{S}\eta_{j}(t)n_{j}(t)}^{S} \\ \overbrace{\sum_{j\neq i}^{S}\eta_{j}(t)n_{j}(t)}^{S} \end{bmatrix} \end{bmatrix}, \quad j = (1, 2, \dots, S), \quad (1)$$

where  $n_j(t)$  is the density of species j at time t,  $s_j$  is the adult survival probability, and  $\eta_j(t)$  is the time-varying per capita larval production.

In the lottery model, empty space is the only limiting resource, so the competitive exclusion principle (which states that no more than N species can coexist on N regulating factors; (Volterra, 1926, Lotka, 1932, Gause, 1934; Levin, 1970) is transcended if even two fish species are able to coexist. Intuitively, if a species becomes rare for whatever reason (e.g., competition, catastrophes), then it must have a positive per capita growth rate if it is to recover from rarity. This is a slight simplification (see Barabás et al., 2018, p. 293), but for our current purposes, we say that coexistence is related to a *rare-species advantage*, operationalized by the *invasion growth rate*: the long-term average per capita growth rate of a species that has been perturbed to low density. The practice of determining coexistence based on invasion growth rates is called an *invasion analysis* (Turelli, 1978; Grainger et al., 2019).



a) Environment favors  $\rightarrow$  fecundity

b) Environment favors  $\bigodot$  fecundity



Figure 1: Diagrammatic representation of the lottery model

Consider a two-species lottery model with a red species and a blue species (Fig. 2). The red species produces many larvae during hot years and few larvae during cold years. The blue species responds to temperature oppositely: it produces few larvae during hot years and many larvae during cold years. If the blue fish species becomes rare, will it be able to recover?

We first ask the question, what happens when the blue fish species experiences a good (i.e., cold) year (Fig. 2a)? There are many blue larvae produced per capita, but there are few larvae produced in total because blue fish are rare, and although the red fish are common, their per capita fecundity is low because the environment is less favorable to them. Each blue fish larva thus experiences relatively little competition, which we may measure as larvae per empty site. In this scenario, we see that a rare species is able to capitalize on a good environment.

To uncover a potential rare-species advantage, we must now examine an analogous scenario from the perspective of the common species: what happens when the red fish species experiences a good (i.e., warm) year (Fig. 2b)? Many red larvae are produced in total, since there are many red fish and the environment favors the red fish. However, there is now an excess of red larvae, which significantly decreases the probability of any one larva winning a territory. The consequence of this high competition is a small or zero-valued per capita growth rate; the last panel of Fig. 2b shows no net change). Unlike the rare blue fish, the common red fish is unable to capitalize on a good environment.

We will now frame this concrete scenario (i.e., the red fish experiencing a hot year) in slightly more general terms: For a common species, a good response to the environment (e.g., high per capita larval fecundity) causes high competition (e.g., many larvae per open site), which ultimately undermines the good response to the environment. For a rare species, a good response to the environment does not lead to as much competition. It is this asymmetry between rare and common species that drives the storage effect.

## 2.1 A simple interpretation of the storage effect

Good environments lead to high competition for common species, but less so for rare species. Since high competition undermines the positive effects of a good environment (via a negative interaction effect of environment and competition on per capita growth rates), rare species are better able to take advantage of a good environment than common species.

This interpretation is correct but leaves out some details. The simple interpretation corresponds to a positively-valued storage effect, but what does a negatively-valued storage effect look like? How can the asymmetry between rare species and common species be represented mathematically? For that matter, how can the negative interaction effect be represented mathematically? What, in the simple interpretation, is necessary and/or sufficient for the storage effect? Is the storage effect a species-level or community-level characteristic? These questions are best answered with the mathematical definition of the storage effect and its textual analogue: an ingredient list of conditions for the storage effect.

## 2.2 The mathematical definition

The mathematical definition of the storage effect is embedded within *Modern Coexistence Theory* (Chesson, 1994; Chesson, 2000a; Barabás et al., 2018), a framework for partitioning invasion growth rates into additive terms; these terms correspond to different explanations for coexistence, and are therefore called *coexistence mechanisms*. The storage effect is one of several coexistence mechanisms. In order to arrive at the mathematical definition of the storage effect, we provide a step-wise summary of the derivation of coexistence mechanisms:

1. Choose one species to be the rare species. This species is called the *invader* and the remaining species are called *residents*. Set the invader's density to zero, and let residents attain their limiting dynamics, i.e., let them equilibrate to their typical densities.

The invader is denoted by the subscript i, the residents are denoted by the subscript s, and a generic species is denoted by the subscript j.

2. Write the per capita growth rates in terms of the environmental and competition. Let the per capita growth rate of species j be some function  $g_j$  of the environmental parameter  $E_j(t)$  and competition  $C_j(t)$ , i.e.,  $dn_j(t)/(n_j(t)dt) = r_j(t) = g_j(E_j(t), C_j(t))$ . In discrete-time models like the lottery model, the effective per capita growth rate is the logged *finite rate of increase*, i.e.,  $r_j(t) = \log(\lambda_j(t))$ , where  $\lambda_j(t) = n_j(t+1)/n_j(t)$ . Extensions to structured populations can be found in Ellner et al. (2019). For notational simplicity, we drop the explicit dependence on time t.

The parameter  $E_j$  is called the environmental parameter, the environmentally-dependent parameter, or simply the environment. It is typically a demographic parameter, belonging to species j, that depends on the abiotic environment (e.g., germination probability depends on temperature). More generally,  $E_j$  may represent the effects of density-independent factors. The parameter  $C_j$  is called the competition parameter, but it may represent the effects of density-dependent factors. Of particular interest is the case where  $C_j$  is a function of shared predators, potentially leading to the storage effect due to predation (Kuang and Chesson, 2010; Chesson and Kuang, 2010; Stump and Chesson, 2017).

The invasion growth rate is the long-term average per capita growth rate of the invader. More generally, the invasion growth rate is the dominant Lyapunov exponent of the dynamical system representing population dynamics (Metz et al., 1992; **dennis2003**).

3. Expand growth rates with respect to  $E_j$  and  $C_j$ . First, select equilibrium values of the environment and competition,  $E_j^*$  and  $C_j^*$ , such that  $g_j(E_j^*, C_j^*) = 0$ . Next, perform a second-order Taylor series expansion of  $g_j(E_j, C_j)$  about  $E_j^*$  and  $C_j^*$ .

The result is

$$r_{j}(E_{j},C_{j})\Big|_{\substack{C_{j}=C_{j}^{*}\\E_{j}=E_{j}^{*}}} \approx \alpha_{j}^{(1)}(E_{j}-E_{j}^{*}) + \beta_{j}^{(1)}(C_{j}-C_{j}^{*}) + \frac{1}{2}\alpha_{j}^{(2)}(E_{j}-E_{j}^{*})^{2} + \frac{1}{2}\beta_{j}^{(2)}(C_{j}-C_{j}^{*})^{2} + \zeta_{j}(E_{j}-E_{j}^{*})(C_{j}-C_{j}^{*}),$$

$$(2)$$

where the coefficients of the Taylor series,

$$\alpha_{j}^{(1)} = \frac{\partial g_{j}(E_{j}^{*},C_{j}^{*})}{\partial E_{j}}, \quad \beta_{j}^{(1)} = \frac{\partial g_{j}(E_{j}^{*},C_{j}^{*})}{\partial C_{j}}, \quad \alpha_{j}^{(2)} = \frac{\partial^{2} g_{j}(E_{j}^{*},C_{j}^{*})}{\partial E_{j}^{2}}, \quad \beta_{j}^{(2)} = \frac{\partial^{2} g_{j}(E_{j}^{*},C_{j}^{*})}{\partial C_{j},^{2}} \quad \zeta_{j} = \frac{\partial^{2} g(E_{j}^{*},C_{j}^{*})}{\partial E_{j}\partial C_{j}} \quad (3)$$

are all evaluated at  $E_j = E_j^*$  and  $C_j = C_j^*$ .

4. **Time-averaging**. Invasibility is determined by what happens in the long-run, so our next step is to take the temporal average of Eq.2. Temporal averages are denoted with "bars"; e.g., the average per capita growth rate of species j is  $\overline{r}_j$ .

$$\overline{r}_{j}\Big|_{\substack{C_{j}=C_{j}^{*}\\E_{j}=E_{j}^{*}}} \approx \alpha_{j}^{(1)}(\overline{E_{j}}-E_{j}^{*}) + \beta_{j}^{(1)}(\overline{C_{j}}-C_{j}^{*}) + \frac{1}{2}\alpha_{j}^{(2)}\operatorname{Var}(E_{j}) + \frac{1}{2}\beta_{j}^{(2)}\operatorname{Var}(C_{j}) + \zeta_{j}\operatorname{Cov}(E_{j},C_{j}).$$

$$(4)$$

The above expression above rests on several assumptions about the magnitude of environmental fluctuations and the relationship between environment, population density, and competition (details can be found in Chesson (1994) and Chesson (2000a)). Most crucially, we assume that environmental fluctuations,  $|E_j - E_j^*|$ , are very small, and that average environmental fluctuations  $|\overline{E_j} - E_j^*|$ , are even smaller. These *small-noise assumptions* ensure that the expression Eq.4 is a good approximation of the true invasion growth rate, thus justifying the truncation of the Taylor series at second order. The small-noise assumptions also justify the replacement of second-order polynomial terms with central moments, e.g.,  $(E_j - E_j^*)^2$  is replaced by  $\operatorname{Var}(E_j)$ : the equilibrium value  $E_j^*$  is assumed to be very close the temporal average  $\overline{E_j}$ , such that little growth rate is lost by replacing the former with the latter.

#### 5. Invader–resident comparisons

The long-term average growth rate of each resident must be zero (otherwise residents would go extinct or explode to infinity), so the value of the invasion growth rate is unaltered if we subtract a linear combination of the residents' long-term average growth rates.

$$\overline{r}_i = \overline{r}_i - \sum_{s \neq i}^S q_{is} \overline{r}_s \tag{5}$$

The  $q_{is}$  are called *scaling factors* (Barabás et al., 2018) or comparison quotients (Chesson, 2019). The original definition, provided by Chesson (1994), is  $q_{is} = \frac{\beta_i^{(1)}}{\beta_s^{(1)}} \frac{\partial C_i}{\partial C_s}$  (evaluated at the equilibrium values of environment and competition). Ellner et al. (2016, 2019) has suggested scaling resident growth rates to create a simple average over resident species: in Eq.5, replace the  $q_{is}$  with 1/(S-1). Elsewhere, we have argued for scaling resident growth rates by a ratio of species' generation times, so as to convert the residents' population-dynamical speeds to that of the invader (Johnson and Hastings, 2022). Incidentally, this approach is equivalent to the  $q_{is}$  scaling factors for the simple models that we examine in this paper. Additionally, our arguments against the conventional interpretations of the storage effect do not hinge on how resident growth rates are scaled. For these reasons and for the sake of convention, we use Chesson's original scaling factors.

Though the average growth rate of each resident is zero, the components of the average growth rate (i.e., the additive terms in Eq.4) are not necessarily zero. Therefore, we can draw meaningful comparisons between the invader and the residents by substituting the right-hand side of the Taylor series expansion (Eq.4) into the invader–resident comparison (Eq.5) and grouping like-terms:

$$\overline{r}_{i} \approx \alpha_{i}^{(1)}(\overline{E_{i}} - E_{i}^{*}) + \frac{1}{2}\alpha_{i}^{(2)}\operatorname{Var}(E_{i}) + \beta_{i}^{(1)}C_{i}^{*} - \sum_{s\neq i}^{S} q_{is}\left((\overline{E_{s}} - E_{s}^{*}) + \frac{1}{2}\alpha_{s}^{(2)}\operatorname{Var}(E_{s}) + \beta_{s}^{(1)}C_{s}^{*}\right)$$

 $r'_i$ , Density-independent effects

$$+ \underbrace{\beta_{i}^{(1)}\overline{C_{i}} - \sum_{s \neq i}^{S} q_{is}\beta_{s}^{(1)}\overline{C_{s}}}_{\Delta \rho_{i},\text{Linear density-dependent effects}} \\ + \underbrace{\frac{1}{2} \left[ \beta_{i}^{(2)}\text{Var}(C_{i}) - \sum_{s \neq i}^{S} q_{is}\beta_{s}^{(2)}\text{Var}(C_{s}) \right]}_{\Delta N_{i},\text{Relative nonlinearity}} \\ + \underbrace{\zeta_{i}\text{Cov}(E_{i},C_{i}) - \sum_{s \neq i}^{S} q_{is}\zeta_{s}\text{Cov}(E_{s},C_{s})}_{\Delta I_{i},\text{The storage effect}}.$$

(6)

The new symbols  $(r'_i, \Delta \rho_i, \Delta N_i, \text{ and } \Delta I_i)$  denote coexistence mechanisms.

Because the coexistence mechanisms constitute the invasion growth rate, they measure a *rare-species advantage*. Because the coexistence mechanisms make cross-species comparisons, they capture (respectively) different types of *specialization*. Because explanations for coexistence have historically featured types of specialization that result in a rare-species advantage, it is fair to say that coexistence mechanisms measure the importance of different classes of explanations for coexistence.

The mathematical definition of the storage effect, denoted  $\Delta I_i$ , is

$$\Delta I_i = \zeta_i \operatorname{Cov}(E_i, C_i) - \sum_{s \neq i}^{S} q_{is} \zeta_j \operatorname{Cov}(E_s, C_s).$$
(7)

When ecologists talk colloquially about a storage effect, they are typically talking about a positive  $\Delta I_i$  that is mediated through competition. However, the storage effect can also be negative, and/or mediated through apparent competition. In the case of a negative storage effect, there is a tendency for rarity to cause lower per capita growth rates. Therefore, negative storage effects can mediate a stochastic priority effect (Chesson, 1988; Schreiber, 2021).

Coexistence mechanisms are often divided by the invader's sensitivity to competition, which we may operationalize here as  $|\beta_i^{(1)}|$ . The rationale here is that  $|\beta_i^{(1)}|$  can be interpreted as the speed of population dynamics (at least in the lottery model and annual plant model; sensu Chesson, 1994), so dividing by it enables a better comparison of species with slow and fast life-cycles (Chesson, 2018). Note that this type of scaling is distinct from the aforementioned  $q_{is}$  scaling factors (though they are related; Johnson and Hastings, 2022).

Scaled coexistence mechanisms are sometimes averaged over species (see Chesson, 2003, Barabás et al., 2018), either to make comparisons between communities or to quantify how a mechanism affects species *in general*. The *community-average storage effect* is defined as

$$\overline{\left(\frac{\Delta I}{\left|\beta^{(1)}\right|}\right)} = \frac{1}{S} \sum_{i=1}^{S} \frac{\Delta I_i}{\left|\beta_i^{(1)}\right|}.$$
(8)

### 2.3 The ingredient-list definition

The storage effect depends on three ingredients:

- 1. species-specific responses to the environment,
- 2. a non-zero interaction effect with respect to fluctuations in the environment and competition (also known as *nonadditivity*), and
- 3. covariance between environment and competition (EC covariance for short).

This ingredient-list definition (Chesson and Huntly, 1989; Chesson, 1994) describes what is generally required for the storage effect to be positive (or negative) for many species. To see that the ingredients are associated with a systematically positive (or negative) storage effect, consider a model with symmetric species: each species responds the environment in accordance with a symmetric covariance matrix with diagonal elements  $\sigma^2$  and off-diagonal elements  $\rho\sigma^2$ , where  $\rho$  is the between-species correlation in  $E_j$ ; species otherwise have identical demographic parameters. Just below (in Section 2.3.1), we show that the storage effect is then

$$\Delta I = -\zeta (1 - \rho)\theta \tag{9}$$

for every species. The three ingredients are captured in the above formula: species-specific responses to the environment is  $(1 - \rho)$ ; the interaction effect is  $\zeta$ ; and covariance is  $\sigma^2 \theta$ , where  $\theta$  is a constant that converts the environmental responses of residents into competition (see Section 2.3.1 below). Mathematical expressions for the storage effect are generally more complicated in the non-symmetric case (see Eq. 29 in Chesson, 1994).

Contrary to the claims of previous research, the three ingredients are neither necessary nor sufficient for a single species' storage effect to be non-zero. To see why the ingredients are not necessary, consider the following non-symmetric scenario in which all species respond to the environment identically, but only species k has a non-zero interaction effect (i.e.,  $\zeta_k \neq 0$ ). In this scenario, the condition # 1 (species-specific responses to the environment) is not satisfied, yet species k may nonetheless have a non-zero storage effect. In Appendix 6.1, we show that the ingredients are not sufficient for the storage effect: when all ingredients are present in the lottery model, the storage effect can be zero if different species have different adult survival probabilities.

While it may seem pedantic to emphasize that the ingredients are neither necessary nor sufficient, we will see later (Section 4) that this has implications for the interpretation of the storage effect. We do not want to give the impression that the ingredients are unimportant. Very few high-level features of the world have necessary and sufficient properties; in everyday life and in ecology, concepts are fuzzy, being held together by an open-ended set of correlated properties (Wittgenstein famously called these family resemblance concepts; Wittgenstein, 1968). Because the storage effect is generally related to the three ingredients, we can better understand the the storage effect by studying the three ingredients in-depth.

#### 2.3.1 Ingredient #1: Species-specific responses to the environment

The function of ingredient #1 is rather obvious: to establish the presence of niche differences, which are necessary for coexistence via any mechanism (Gause, 1934; Chesson, 1991). In the absence of *species-specific responses to the environment*, there would be no rare-species advantage. In terms of the lottery model, a good (bad) year for the blue species would automatically be a good (bad) year for the common red species, such that both species would always experience the same level of competition. In our story about coral reef fishes, species had negatively correlated responses to the environment. However, *species-specific responses to the environment* only entails that species' responses not be perfectly correlated. As we will see (in Section 3), real-world species tend to have positively correlated responses to the environment.

To see how species-specific responses to the environment are implied by the mathematical definition, we must re-express the competition parameter in terms of the environment. For illustrative purposes, suppose that there are only two species, and that both share a competition parameter C that can be written as a smooth function h of the resident's density and environmental response:  $C = h(E_s, n_s)$ . The the competition parameter can then be approximated as a first-order Taylor series expansion: C = $C^* + \frac{\partial h(E_s^*, n_s^*)}{\partial E_s}(E_s - E_s^*) + \frac{\partial h(E_s^*, n_s^*)}{\partial n_s}(n_s - n_s^*)$ . With the additional assumption that environmental states are uncorrelated across time (which precludes covariance between environment and population density), this approximation of C can be substituted into the EC covariance. For the invader,  $\text{Cov}(E_i, C) \approx$  $\frac{\partial h(E_s^*, n_s^*)}{\partial E_s} \text{Cov}(E_i, E_s) = \frac{\partial h(E_s^*, n_s^*)}{\partial E_s} \rho_{\sigma_i \sigma_s}$ , where  $\sigma_i$  is the scale of fluctuations in  $E_i$ ,  $\sigma_s$  is the scale of fluctuations in  $E_s$ , and  $\rho$  is the cross-species correlation between  $E_i$  and  $E_s$ . By contrast, the residents' covariance term is  $\text{Cov}(E_s, C) \approx \frac{\partial h(E_s^*, n_s^*)}{\partial E_s} \text{Var}(E_s) = \frac{\partial h(E_s^*, n_s^*)}{\partial E_s} \sigma_2^2$ . Next, we plug these covariance approximations into the definition of the storage effect, define

Next, we plug these covariance approximations into the definition of the storage effect, define  $\theta = \frac{\partial h(E_2^*, n_2^*)}{\partial E_2}$ , and assume that species have identical  $\zeta_j$ ,  $\sigma_j$ ,  $\beta_j^{(1)}$  (which implies that  $q_{is} = 1$ ). Then, we arrive at Eq.9 above:  $\Delta I_i = -\zeta(1-\rho)\theta$ . Generalizing to the multi-species case involves redefining the h so that competition is a function of all residents' environmental responses  $\vec{E}$ , and population densities  $\vec{n}$ ; redefining  $\theta = \frac{1}{S-1}\sum_{s\neq i}^{S} \frac{\partial h(\vec{E^*}, \vec{n^*})}{\partial E_s}$ . If  $\zeta$  is negative (which indicates *buffering*), then the storage effect is proportional to  $(1-\rho)$ , demonstrating that (all else being equal) the storage effect increases as environmental niche differences increase. asdf

## 2.3.2 Ingredient #2: An interaction effect between environment and competition

When a red fish experiences a hot year, it is not merely the case that the negative effects of high competition offset the positive effects of a good environment. Rather, the environment and competition act synergistically to reduce per capita growth rates further. This synergy is the *interaction effect*, akin to an interaction effect in multiple regression. In fact, the coefficient  $\zeta_j$  in the mathematical definition of the storage effect (Eq.7) is the interaction effect of a multiple regression where  $E_j - E_j^*$ and  $C_j - C_j^*$  are predictors, in the limit of small environmental noise. The causal interpretation of an interaction effect in multiple regression is that the level of one predictor modulates another predictor's effect on the response variable (Gelman and Hill, 2007), which is why the simple interpretation of the storage effect (Section 2.1) states that "high competition undermines the positive effects of a good environment". Put yet another way, high competition means that the population is less sensitive to changes in the environment. In our exposition thus far, we have described a negative interaction effect. However, both negative or positive interaction effect can lead to either a positive or negative storage effect. In the jargon of Modern Coexistence Theory, a negative interaction effect (i.e.,  $\zeta < 0$ ) is called *subadditivity* or *buffering* (Chesson, 1994). The term *subadditive* comes from the fact that the joint effects of environment and competition are less than than the sum of their parts. The term *buffering* comes from the fact that the doubly deleterious effect of a poor environment and high competition is somewhat abated: the term  $\zeta_j(E_j - E_j^*)(C_j - C_j^*)$  is positive when  $\zeta_j < 0$ ,  $(E_j - E_j^*) < 0$ , and  $(C_j - C_j^*) > 0$ . A positive interaction effect (i.e.,  $\zeta_j > 0$ ) is synonymous with *superadditivity* or *amplifying* (Chesson and Ellner, 1989). More generally, both positive and negative interaction effects are referred to as *nonadditivity*. The storage effect is generally positive in systems with subadditivity and positive *EC* covariances, *or* systems with superadditivity and negative *EC* covariances. Conversely, the storage effect is generally negative in systems with subadditivity and negative *EC* covariances, *or* systems with superadditivity and positive *EC* covariances.

At a high level of abstraction, the interaction effect can be thought of combining the environment and competition into a large number of density-dependent factors. Coexistence requires negative feedback loops where species demonstrate some degree of specialization on density-dependent factors (Meszéna et al., 2006), but some communities do not have enough density-dependent factors to be specialized upon (Hutchinson, 1959; Hutchinson, 1961; but see Levin, 1970; Haigh and Smith, 1972; Abrams, 1988). On the other hand, species may readily specialize on different environmental states, but environmental variation alone cannot promote coexistence (Chesson and Huntly, 1997). The interaction effect combines the competition parameter with the environmental parameter to get the best of both worlds: the density-dependent factors (implicit in the competition parameter) provide the negative feedback while the species-specific environmental parameter provides the specialization.

But what is an interaction effect in more concrete terms? What ecological concepts lead to an interaction effect in particular models? In coexistence theory, a negative interaction effect has primarily been associated with differential sensitivities of different life-stages. Chesson and Huntly (1988) write, "... iteroparous plant and sessile marine organisms, can buffer by participating in reproduction over a number of years... Semelparous species can experience these buffering effects if the offspring of an individual mature over a range of years..." More generally, a negative interaction effect may arise from other forms of population structure: dormancy (Cáceres, 1997; Ellner, 1987), phenotypic variation (Chesson, 2000b), or spatial variation (Chesson, 2000a). In studies of the population genetic storage effect (which promotes allelic diversity), negative interaction effects can be produced by heterozygotes (Dempster, 1955; Haldane and Jayakar, 1963), sex-linked alleles (Reinhold, 2000), epistasis (Gulisija et al., 2016), and maternal effects (Yamamichi and Hoso, 2017).

Though there are plenty of models in which population structure produces an interaction effect, population structure is neither necessary nor sufficient for an interaction effect. To see why it is not sufficient, consider a modified version of the lottery model in which adult mortality occurs before the spawning event. In this model, the logged finite rate of increase (i.e., the discrete-time analogue of per capita growth rate) is  $r_j = \log(s_j(1 - \exp\{E_j - C\}))$  (compare with the original lottery model, Eq.14), and therefore,  $\zeta_j = 0$ . Here, a storage effect is impossible, even though there is stage structure, and even though the larval stage is more sensitive than the adult stage to both environment and competition (as in the original lottery model).

To see why population structure is not necessary for an interaction effect, consider a phytoplankton species that Monod (1949) dynamics,

$$\frac{dn_j(t)}{dt} = n_j(t) \left[ b_j(t) \frac{R(t)}{K_j + R(t)} - d_j \right],$$
(10)

where  $n_j(t)$  is the density of phytoplankton species j,  $b_j(t)$  is the temporally-fluctuating maximum rate of increase, R(t) is nitrogen concentration,  $K_j$  is the half-saturation constant, and  $d_j$  is the death rate. Defining  $E_j = \log(b_j)$  and  $C_j = \log\left(\frac{K_j+R}{R}\right)$ , we find that  $\zeta_j = d_j$ . Here, we see an interaction effect (buffering increases as the death rate increases), despite the lack of stage structure. This phytoplankton model also demonstrates that "buffering" (i.e., a negative interaction effect) is not necessarily a result of life-history traits (e.g., dormancy, iteroparity) that have a clear adaptive purpose, but rather a by-product of banal population dynamics: the maximum uptake rate varies with the environment (which follows from the fact that the rate of a chemical reaction tends to increase with temperature; Eyring, 1935; Kingsolver, 2009), growth rates are a saturating function of resource concentration (which follows from ion channels having a nutrient handling time; Aksnes and Egge, 1991). The consequence of these non-adaptive and putatively inescapable features is a negative interaction effect.

We conclude that at the current moment, there is no ecological interpretation of the interaction effect that is satisfactorily general. That is, an interaction effect cannot be intuited purely from knowledge about species' life-history characteristics, such as dormancy, iteroparity/non-overlapping generations, stage-structure, or more generally, a verbal description of a new model. Instead, the ecological interpretation of an interaction effect (or lack thereof) must be determined on a model-by-model basis, either with mathematical analysis or analogy with previously-studied models.

## 2.3.3 Ingredient #3: Covariance between environment and competition

The final ingredient, covariance between environment and competition, is immediately evident in the mathematical definition of the storage effect (Eq.7). In more biological terms, the covariance captures the causal relationship between environment and competition. The most obvious way in which a good environment causes high competition (and vice-versa) is through intergenerational population growth: a good environment produces a larger population, and a larger population usually corresponds to higher competition. However, temporal autocorrelation in the environment is required for ECcovariance via intergenerational population growth (Li and Chesson, 2016; Letten et al., 2018; Ellner et al., 2019; Schreiber, 2021); the past environment determines the present competition, but the ECcovariation involves the current environment and current competition, so the current environment must resemble the past environment. The storage effect also arises when species have phenology differences in periodic environments (Loreau, 1989; Loreau, 1992; Klausmeier, 2010), since a periodic environment is just a special case of a temporally autocorrelated environment. In a stage-structured model, a good environment can lead to high competition within a single time-step (Chesson and Huntly, 1988). Consider the lottery model: a good environment (i.e., high per capita fecundity) at the spawning stage leads to high competition (i.e., many larvae per territory) at the recruitment stage. Note here that there is still temporal autocorrelation in the sense that the larvae carry the effects of the environment through time.

Although the archetypical storage effect is mediated through resource competition, the storage effect may also be mediated through apparent competition; the parameter  $C_j$  may be generally understood as the effects of all density-dependent factors. When the storage effect is mediated through resource competition, the *EC* covariance is generically positive, though it may be negative when the environment is negatively autocorrelated (Schreiber, 2021).

When the storage effect is mediated through apparent competition, a negative, positive or zerovalued covariance is possible. For the sake of the current discussion, assume that the competition parameter is the density of the shared predator P, times the predator's functional response  $f_j(N_j)$ , all divided by prey density i.e.,  $C_j = P * (f_j(N_j)/N_j)$ . Stump and Chesson (2017) analyzed a variant of the annual plant model and found that a type 2 functional response leads to a negative EC covariance: good environments lead to a large number of seeds, which satiate predators, thus lowering the per-seed predation pressure. Kuang and Chesson (2010, 2010) found that the a type 3 functional response (i.e., frequency-dependent predation) leads to a positive EC covariance: good environments lead to a large number of seeds, which are then preferentially consumed. In the previous two examples, the predator demonstrates a fast behavioral response to changes in prey density. If the predator demonstrates only a numerical response to prey density (corresponding to a type 1 functional response) and the environment is temporally autocorrelated, then the covariance will be positive. If instead the environment is uncorrelated through time, the storage effect due to the predation is not possible (Kuang and Chesson, 2009): the environment changes before it appreciably affects predation pressure, and therefore does not produce the necessary covariance.

# 3 A critique of the conventional interpretations of the storage effect

The conventional interpretation v1 of the storage effect is built around the idea that species have special adaptations or life-history traits that help them persist through temporarily hostile environments. This theme is evidenced by various quotes:

- "...adults must be able to survive over periods of poor recruitment, such that the population declines only slowly during these periods. Under these conditions, a species tends to recover from low densities, and competitive exclusion is opposed. .... We refer to this phenomenon as the storage effect because strong recruitments are essentially stored in the adult population, and are capable of contributing to reproduction when favorable conditions return (Chesson, 1985).
- "... the storage-effect coexistence mechanism relies on such buffering effects of persistent stages, because these prevent catastrophic population decline when poor recruitment occurs." (Chesson, 2003)
- "Persistence of adults limits the damage from unfavourable conditions, but does not prevent strong growth at other times. ... Similarly, the dormant seeds of annual plants are relatively insensitive to environmental factors and competition in comparison with the actively growing plants." (Chesson et al., 2004)
- "Seed banks or long-lived adults "store" the effects of favorable years, which buffer the effects of bad years when population sizes may decline." (Sears and Chesson, 2007)
- "First, organisms must have some mechanism for persisting during unfavourable periods, such as a seedbank, quiescence or diapause. This condition, which gives the storage effect its name, buffers negative population growth; without it, populations would go extinct after a brief unfavourable period and environmental variation could never promote coexistence." (Adler, 2014)

To be clear, we are not making any judgements about what the quoted authors do or do not know about the storage effect; it is often pedagogically useful to present definitions that are evocative but not 100% precise. Our purpose in quoting the above authors is to provide evidence that there exists an interpretation of the storage effect, paraphrased as ...

The conventional interpretation of the storage effect v1: Species coexist by specializing on different parts of a fluctuating environment, so species must have a robust life stage in order to "wait it out" for a favorable time period. Thus, "storage" refers to a robust life stage can "wait it out".

By showing that age or stage structure is not necessary for the storage effect, a number of papers have shown that the conventional interpretation v1 is imprecise. The temporal storage effect can be generated in models without any age or stage structure if the environment is temporally autocorrelated (Loreau, 1989; Loreau, 1992; Klausmeier, 2010; Li and Chesson, 2016; Letten et al., 2018; Ellner et al., 2019; Schreiber, 2021). All of these unstructured population models can be interpreted as having non-overlapping generations; thus, the notion of longevity or an individual "waiting it out" for a good year is not necessary for the storage effect.

A simple argument shows why the conventional interpretation v1 is fundamentally flawed: An invader population has functionally has no individuals in which population vitality or years of good recruitment can be stored. The fact that we can approximate the dynamics of a low density population with a zero density population means that the presence of conspecific individuals has a negligible effect on per capita growth rates. To be clear, in structured population models, invasion growth rates depend on the quasi-stable age/stage distribution (Caswell, 2001), even if the invader is at infinitesimal density. However, in the lottery model, all larvae die within a single time-step, which means the stable stage distribution in the lottery model is entirely adult fish, regardless of the adult survival parameter. Again, the conventional interpretation v1 cannot be correct because the whole premise of invasion analysis is that individuals of a rare species have negligible effects on per capita growth rates.

It is true that in the lottery model and the annual plant model, coexistence is not possible when generations are non-overlapping (i.e., adult or seed survival equals zero). However, this is because the particular structure of these models do not allow for an interaction effect when the survival probability equals zero, not because anything is stored in a particular life stage. To see this more clearly, we will consider the classic annual plant model (also called the seedbank model; Chesson, 1994),

$$X_j(t+1) = X_j(t) \left[ s_j(1 - G_j(t)) + \frac{G_j(t)Y_j}{\sum_{k=1}^S c_k G_k(t)X_k(t)} \right],$$
(11)

where  $X_j$  is the density of seeds of species j,  $s_j$  is the probability that a seed survives the growing season if it does not germinate,  $G_j(t)$  is the time-varying germination probability,  $c_k$  are competition coefficients, and  $Y_j$  is proportional to yield: the number of seeds per germinant. Defining  $C_j = C =$  $\log\left(\sum_{k=1}^{S} c_k G_k(t) X_k(t)\right)$  and  $E_j = \log(G_j(t))$ , we find that  $\zeta_j = -s_j$ : As previously stated, the interaction effect is zero when seed survival is zero. In general, the interaction effect will vanish for any growth rate function that takes the form  $r = \log(u(E) * v(C)) = \log(u(E)) + \log(v(C))$ , where uand v are arbitrary but smooth functions.

Now, consider the slightly-modified annual plant model, where competition is a piece-wise function:

$$X_j(t+1) = X_j(t) \left[ s_j(1 - G_j(t)) + G_j(t) Y_j - C_j(t) \right]$$
(12)

$$C_{j}(t) = \begin{cases} \sum_{k=1}^{S} c_{k}G_{k}(t)X_{k}(t) & \text{if } \epsilon + \sum_{k=1}^{S} c_{k}G_{k}(t)X_{k}(t) \leq s_{j}(1 - G_{j}(t)) \\ \epsilon & \text{if } \epsilon + \sum_{k=1}^{S} c_{k}G_{k}(t)X_{k}(t) > s_{j}(1 - G_{j}(t)). \end{cases}$$
(13)

Like the original annual plant model, the per capita yield is  $G_j Y_j$  when  $C_j = 0$ . Unlike the original annual plant model, the finite rate of increase,  $\lambda_j$ , decreases linearly with competition. As competition increases further, the finite rate of increase levels-out at some small number  $\epsilon$ , which prevents  $\log(\lambda_j) = -\infty$  (a mathematical nuisance), but may have no influence on actual dynamics if competition never increases to that point. In this modified annual plant model, the interaction effect is  $\zeta_j = Y_j$  when when there is no seedbank (i.e.,  $s_j = 0$ ). This demonstrates that stage structure is by no means necessary for the storage effect, it is just one of many ways in which an interaction effect may arise. Although the meaning of an interaction effect can be stated in abstract terms, it is difficult to explain in terms of ecological or population dynamical concepts (see Section 2.3.2).

Some authors employ a more general interpretation of the storage effect based on the abstract notion of *buffering*. Chesson et al. (2004) explains: "Buffered population growth means that less is lost during unfavorable conditions (high competition and an unfavorable environment) than is gained during favorable conditions (low competition and a favorable environment)...". Buffering can be measured as minus the interaction effect between environment and competition, which is, in mathematical terms, the cross partial derivative of per capita growth rates with respect to environment and competition, evaluated around equilibrium values, i.e., buffering =  $-\zeta = -\partial^2 r(E^*, C^*)/(\partial E\partial C)$  (Chesson, 1985; Chesson, 2000b; Snyder, 2012). Again, quotes provide evidence that this more general interpretation exists in the ecology milieu:

- "...there is some way to "store" the effects of good times, to get organisms through bad ones." (Barabás et al., 2018)
- "More generally, mechanisms leading a positive  $\Delta I$  value involve storage of the benefits of favorable periods in the population, whether this storage can be traced to a seed bank of something else. The term storage is a metaphor for the potential for periods of strong positive growth rate that cannot be canceled by negative growth at other times." (Chesson, 1994)
- "Storage effects happen when the invader experiences low competition in favorable environments and has the ability to store that double benefit." (ete:Snyder)
- "However, these gains by the rare come to nothing if they are wiped out in bad years. The storage effect can therefore maintain coexistence only if species are buffered against sudden rapid

declines. One natural way for this to occur is if generations overlap and established individuals are immune to the causes of temporal variation (e.g., viability selection on offspring, no selection on adults)." (Messer et al., 2016)

The conventional interpretation storage effect v2: "Storage" can be more generally understood as 'buffering', which is a negative interaction effect of environment and competition on per capita growth rates. This buffering helps out rare species because it prevents extreme losses when the environment is unfavorable and competition is high.

The convention interpretation v2 is imprecise because the storage effect tends to decrease as invader "storage" (operationalized as buffering) increases. To see this, we derive a mathematical expression for the storage effect in the two-species lottery model (Appendix 6.1). When species-specific responses to the environment are partially correlated (i.e.,  $0 < \rho < 1$ ), the storage effect for the species 1 is proportional to  $[s_2 - \rho s_1]$ , where  $s_j$  is the adult survival probability of species *j*. As the invader's adult survival probability  $s_1$  increases, the storage effect decreases. Since the invader's adult survival probability is measure of the invader's "storage" or "buffering", we see that storage can decrease the storage effect.

If species have positively correlated environmental responses, then species *i*'s storage effect will tend to increase as species *i* experiences more buffering (i.e., as  $\zeta_i$  becomes more negative). This is precisely what we expect to see in nature. After all, it is well-known that plants tend to have strong and positive responses to increases in temperature and precipitation (Rosenzweig, 1968; Lieth and Whittaker, 1973; Sala et al., 1988. The probability of germination — which is often identified as the environmental response in models of annual plants — can display a complex interdependency on temperature and precipitation (Facelli et al., 2005), but nevertheless tends to increases as either abiotic variable increases (Baskin and Baskin, 1998).

Indeed, there is good empirical evidence that species do have positively correlated environmental responses. We reviewed empirical studies that explicitly attempted to quantify or provide evidence for/against the storage effect (Cáceres, 1997; Venable et al., 1993; Pake and Venable, 1995; Pake and Venable, 1996; Adler et al., 2006; Sears and Chesson, 2007; Descamps-Julien and Gonzalez, 2005; Angert et al., 2009; Usinowicz et al., 2012; Facelli et al., 2005; Chesson et al., 2012; Kelly and Bowler, 2002; Kelly and Bowler, 2005; Usinowicz et al., 2017; Ignace et al., 2018; Hallett et al., 2019; Armitage and Jones, 2019; Armitage and Jones, 2020; Zepeda and Martorell, 2019; Zepeda and Martorell, 2019; Towers et al., 2020; Jiang and Morin, 2007; Holt and Chesson, 2014; Ellner et al., 2016). In the 24 studies we were able to find, there were 15 distinct communities. Of these 15 communities, 7/15 (47%) showed evidence of positive correlations in species' environmental responses, 3/15 (20%) showed zero correlation on average, 2/15 (13%) showed negative correlations, and 3/15 (20%) did not provide sufficient information to make a determination about the average sign of pairwise correlations. Two of the communities were only studied in the context of microcosm experiments. When we only consider natural communities for which sufficient information is available, 7/10 (70%) communities showed positive correlations and 2/10 (20%) show uncorrelated responses. Only 1 community showed evidence of negative correlations, which is what is needed for buffering to help rare species. For more details on our analysis of the literature, see *empirical* E correlations.pdf at https://github.com/ ejohnson6767/storage effect critique.

To be clear, it *is* true that in the lottery model (and the annual plant model), no species can have a positive storage effect if all species have zero adult (or seed) survival probability. However, *this* is a community-level condition for coexistence, reflected properly in the ingredient list definition of the storage effect (Section 2.3). The conventional interpretations v2 may be thought of as conflating the community-level condition for coexistence (i.e., some species must have some "storage" for some species to coexist via the storage effect) with a species-level condition for persistence (i.e., one species must have "storage" in order for said species' storage effect to be positive). The conflation is analogous to falsely claiming that a species can persist by strongly competing with itself, since the competitive Lotka-Volterra model shows us that coexistence occurs when intraspecific competition is greater than interspecific competition.

## 3.1 The origin of the conventional interpretation

The storage effect was discovered by Chesson and Warner (1981) and coined by Chesson (1983), though the phenomenon had been discovered much earlier in the context of population genetics (Dempster, 1955;Haldane and Jayakar, 1963; Lloyd and Dybas, 1966; Harper and White, 1971; Gillespie, 1977). In the 1980's, a sequence of papers (Chesson and Warner, 1981; Chesson, 1982; Chesson, 1983; Chesson, 1984; Warner and Chesson, 1985) analyzed the two-species lottery model and repeatedly highlighted an interesting result: coexistence is not possible if both species have non-overlapping generations (i.e., the adult survival probability equals zero). Additionally, the species with the lower average per capita fecundity cannot persist if it has non-overlapping generations. Analogous results were found for a model of annual plants (Ellner, 1985a; Ellner, 1985b; Ellner, 1987; Chesson, 1994): coexistence is not possible if neither species has a seed bank (i.e., the seed survival probability is zero). At first glance, these results seems to support the conventional interpretation v1 and v2.

This superficial paradox can be resolved by first recognizing that coexistence mechanisms are not independent (Song et al., 2020; Kuang and Chesson, 2010; Yuan and Chesson, 2015). For example, every coexistence mechanism in the lottery model is modulated by the survival parameter. When invader survival increases, the invasion growth rate unsurprisingly increases, despite the fact that the storage effect may decrease. Now the question becomes, why did early expositions of the storage effect conflate the contemporary storage effect with the invasion growth rate?

When the storage effect was first discovered, Modern Coexistence Theory (the framework for partitioning invasion growth rates into coexistence mechanisms) had not yet been invented. The storage effect could not be separated out from other coexistence mechanisms. Consequentially, early formulae defined the storage effect as the sum of multiple coexistence mechanisms. For example, Eq.10 of Warner and Chesson (1985) suggests measuring the storage effect as the combined effects of relative nonlinearity (a different coexistence mechanism; Armstrong and McGehee, 1980; Chesson, 1994) and the contemporary storage effect, which would indeed imply that increasing invader survival increases the storage effect (Appendix 6.1).

## **3.2** The perpetuation of the conventional interpretation

Why is the conventional interpretation perpetuated? We have two speculative answers: 1) the conventional interpretation is "mentally sticky": the ability to persist is intuitively tied to the ability to weather bad years. 2) Coexistence is often studied in models where species behave symmetrically. In such models, the formula for the species-level storage effect (e.g., Eq.9) increases with a shared (across species) survival parameter, thus giving the misleading impression that in general, the storage effect increases with invader survival.

It is entirely reasonable to think that if species specialize on a fluctuating environment, they must have some way to slow the exponential loss of individuals over a sequence of bad years, and that this must be particularly important for rare species, which are inherently extinction-prone (MacArthur, 1967; Lande, 1998). However important this phenomenon might be, it is not what the storage effect is measuring. Most models used to demonstrate the storage effect feature infinite populations (an assumption made for mathematical/computational convenience), which obviates the possibility of stochastic extinction. Unless  $\lambda = 0$  or  $r = -\infty$  (which in most models is only possible as  $n \to \infty$ , a biological impossibility), extinction for infinite populations will take an infinite amount of time even without any "storage".

The notion of "waiting it out" implies that the finite rate of increase (i.e.,  $\lambda$ , the discrete-time analogue of the per capita growth rate) is drawn from a distribution with right skew. Intuition dictates that right skew in  $\lambda$  — many slightly bad years and few extremely good years — should increase the risk of stochastic extinction. However, using a Taylor series of  $\lambda$  about its mean, one can show that right skew increases the invasion growth rate when fluctuations in  $\lambda$  are small. This apparent contradiction suggests that a theory of large environmental fluctuations in finite populations is needed to understand the phenomenon which the conventional interpretation v1 points to.

Because coexistence mechanisms are defined as invader–resident comparisons (5), coexistence mechanisms capture both a rare-species advantage *and* intrinsic differences between species (Johnson and Hastings, 2022). To isolate the rare-species advantage, and to make the math easier, theoretical ecologists often study the storage effect in models where all species are identical except for their responses to the environment. Such *symmetrical species* models produce formulas for the storage effect where the storage effect increases as a shared survival parameter increases (see Eq.9). This may give the misleading impression that in general, an individual species' storage effect increases with said species' survival probability.

# 4 Discussion

The storage effect is an important concept in community ecology, but it remains widely misunderstood. In Section 2.1, we gave a simple interpretation of the storage effect: Good environments lead to high competition for common species, but less so for rare species; since high competition undermines the positive effects of a good environment, rare species are better able to take advantage of a good environment. This interpretation is a good starting point, but is lacking crucial details. It can be supplemented with the mathematical and ingredient-list definitions of the storage effect, but the math can be confusingly abstract, the ingredient list recapitulates the math, and toy models (though more concrete) can lead to confusion if one overgeneralizes from the results of particular models. In this paper, we have taken a pluralistic approach to explanation, providing many details with the hope that some will be cause the storage effect to "click" for any given reader. At the very least, we hope to give the impression that the storage effect is an intrinsically tricky concept, and therefore, that one ought to have a baseline of skepticism towards heuristic uses of the storage effect.

The conventional interpretation of the storage effect v1 first operationalizes "storage" as a robust or long-lived life stage, and then claims that such "storage" helps rare species. A number of models act as counterexamples to this claim. Further, because we measure the strength of coexistence by perturbing species to zero density, there are effectively no individuals in which years of good recruitment can be stored. While the invasion growth rate often increases with an invader's survival probability, this merely reflects lower per capita death rates, not any sort of bet-hedging strategy as implied by the conventional interpretation v1.

The conventional interpretation v2 first operationalizes "storage" as *buffering* — protection against the double-whammy of a bad environment and high competition (i.e.,  $\zeta_j < 0$ ; Section 2.3.2) — and then claims that such storage helps rare species. If a species experiences buffering, then by definition, it grows comparatively poorly in a good environment with high competition. According to our review of papers that have attempted to quantify the storage effect, species are more likely to find themselves in the good environment/high competition scenario than the good environment/low competition scenario, and thus, buffering tends to hurt rare species.

An imprecise definition or interpretation can often serve a pedagogical purpose. One example comes from Godfrey-Smith (2009): Many evolution textbooks state that there are three conditions for evolution by natural selection: phenotypic variation, heritability, and fitness differences. However, these conditions are not sufficient for evolution, since all conditions can be met, and yet allele frequencies remain stagnant in the face of stabilizing selection. Does the conventional interpretation v2 of the storage effect similarly fall into the category of imperfect-but-useful? By questioning the generality of the conventional interpretation v2, are we undermining its (implicitly pedagogical) purpose? Perhaps; buffering promotes persistence when species' responses to the environment are negatively correlated, and perhaps it is easier to think of niche differentiation (a necessary ingredient for coexistence via any mechanism) as a negative correlation between environmental responses, rather than positively but not perfectly correlated responses.

Our critique of the conventional interpretation v2 (but not v1) implicitly assumes that the contemporary mathematical definition of the storage effect is the right way to define the storage effect. Elsewhere (Johnson and Hastings, 2022), we have argued that the definition is essentially correct (though the scaling factors can be problematic in certain contexts), since the invader-resident comparison captures the notion of *specialization* / *ecological differentiation*, a key element in historical explanations for coexistence. It is possible to devise a definition such that a species' storage effect will always increase with buffering, but such a definition would not automatically validate the conventional interpretation v2 (Appendix 6.2). There is a tendency for the storage effect to increase as the buffering of all species increases. This can be seen clearly in formulas for the community-average storage effect (see Eq.20, Appendix 6.1), and the storage effect in the symmetrical case (see Eq.9). The specious similarity between this tendency and the conventional interpretation v2 suggests that 1) the conventional interpretation v2 can be thought of as taking a community-level criterion for coexistence and mistaking it for a species-level explanation for persistence, and 2) that the storage effect is best understood in a community context. The ingredient-list definition of the storage effect (Section 2.3) gives the community-level "conditions", in the sense that the ingredients *tend* to lead to a positive (or negative) storage effect for many species.

It is possible that much of the confusion around the storage effect can be explained by a psychological tendency towards explaining coexistence in terms of how a single species is able to recover from rarity. We speculate that this tendency exists for the same reason that we use an invasion analysis: it breaks up an abstract property — coexistence — into a set of simpler, concrete scenarios (i.e., can a species invade in the context determined by the limiting dynamics of resident species?). When formulating heuristic explanations for coexistence, one naturally considers a scenario in which a single species is rare, and then asks what allows this species to recover from rarity. In the case of coexistence via classical mechanisms, the heuristic explanation is, "If a species were ever to become rare, the resource that it specializes on will become more abundant, thus increasing per capita growth rates". Here, the density-dependent feedback loop contains intuitive state variables (i.e., species densities and resource concentrations, specifically their mean levels) which interact in an obvious way (i.e., more resources per capita  $\rightarrow$  higher per capita growth rates). The analogous heuristic explanation for coexistence via the storage effect is "If a species were ever to become rare, then the environmental states it specializes on will not lead to as much competition (as they did previously), the covariance between environment and competition will decrease, thus increasing per capita growth rates". Here, the density-dependent feedback loop contains an abstract state variable (i.e., the covariance between environment and competition) that interacts with a species' density in a non-obvious way (i.e., a negative interaction effect, deduced via a Taylor series). This explanation is unintuitive and requires an understanding of the math behind the storage effect. It is simply easier to say that rare species "store good years of recruitment" or "are buffered against unfavorable environmental conditions".

The generality of the storage effect is both a strength and weakness. It is a strength because it gives us a way to analyze, talk about, and quantify a phenomenon that occurs in disparate systems. It is a weakness because a fully-general description of the storage effect requires abstractions like "interaction effects" and "EC covariance", which makes it difficult to formulate a general ecological interpretation of the storage effect — to connect the storage effect to an ecological concept like bet-hedging, or a small set of life-history traits like age-structure or phenotypic variation. Sure, the storage effect can be associated with stage structure, or non-overlapping generations, or dormancy. But it can also be associated by none of those things, or temporal autocorrelation, or maternal effects, or whatever mechanism theorists will dream up next.

# 5 Acknowledgements

We would like to thank Karen Abbott for helpful suggestions.

# 6 Appendixes

## 6.1 The storage effect in the lottery model

The lottery model with pure temporal variation (sensu Chesson, 1994) is written as

$$n_j(t+1) = n_j(t) \left[ s_j + \eta_j(t) \left( \frac{\sum\limits_{k=1}^{S} (1-s_k)n_k(t)}{\sum\limits_{k=1}^{S} \eta_k(t)n_k(t)} \right) \right], \quad j = (1, 2, \dots, S).$$
(14)

Note that we do not need extra equations to track larvae because they die if they are not recruited. In order to fit the lottery model into the mold of Modern Coexistence Theory, we must define the environ-

mental parameter and the competition parameter. Selecting  $E_j = \log(\eta_j)$  and  $C = \log\left(\frac{\sum\limits_{k=1}^{S} \eta_k n_k}{\sum\limits_{j=1}^{S} (1-s_k)n_k}\right)$ ,

the effective per capita growth rate takes the form

$$r_j(E_k, C_j) = \log(s_j + \exp\{E_j - C\})$$
(15)

With the log-scale specifications of  $E_j$  and C, all Taylor series coefficients can be expressed purely as a function of  $s_i$ , which leads to tidy expressions of the coexistence mechanisms. If one uses the non-log-scale specification, the results are qualitatively identical.

Next, we choose the equilibrium parameters  $E_i^*$  and  $C_i^*$ . The shared equilibrium level of competition is defined as the temporal average of competition experienced by the invader, averaged over all species acting as the invader:  $C^* = \frac{1}{S} \sum_{i=1}^{S} \overline{C_i}$ . With this choice made, the equilibrium environmental parameters are fixed at  $E_j^* = \log(1 - \overline{s_j}) + C^*$ .

With the equilibrium parameters defined, the partial derivatives of  $g_j$  may be computed:

$$\alpha_{j}^{(1)} = \frac{\partial g_{j}(E_{j}^{*}, C_{j}^{*})}{\partial E_{j}} = 1 - s_{j}$$

$$\beta_{j}^{(1)} = \frac{\partial g_{j}(E_{j}^{*}, C_{j}^{*})}{\partial C_{j}} = -(1 - s_{j})$$

$$\alpha_{j}^{(2)} = \frac{\partial^{2} g_{j}(E_{j}^{*}, C_{j}^{*})}{\partial E_{j}^{2}} = s_{j}(1 - s_{j})$$

$$\beta_{j}^{(2)} = \frac{\partial^{2} g_{j}(E_{j}^{*}, C_{j}^{*})}{\partial C_{j}^{2}} = s_{j}(1 - s_{j})$$

$$\zeta_{j} = \frac{\partial^{2} g(E_{j}^{*}, C_{j}^{*})}{\partial E_{j} \partial C_{j}} = -s_{j}(1 - s_{j}).$$
(16)

The general definition of the scaling factors is  $q_{is} = \frac{\beta_i^{(1)}}{\beta_s^{(1)}} \frac{\partial C_i}{\partial C_s}$  (see Section 2.2). In the case where species share a single competition parameter, this reduces to  $q_{is} = \frac{\beta_i^{(1)}}{\beta_s^{(1)}}$ . Under the standard small-noise assumptions of Modern Coexistence Theory, the competition pa-

rameter can be approximated by a function of residents' environmental responses:  $(C - C^*) \approx$  $\sum_{s\neq i}^{S} \frac{\partial h(\vec{E^*}, \vec{n^*})}{\partial E_s} (E_s - E_s^*).$  In the lottery model with only one invader and one resident, there is a one-to-one conversion between environment and competition:  $\frac{\partial h(E_s^*, n_s^*)}{\partial E_s} = 1$ . In a two-species system with species 1 as the invader and species 2 as the resident, the covariance

between species 1's environment response and competition is  $\operatorname{Cov}(E_1, C) = \operatorname{Cov}(E_1, E_2) = \rho \sigma^2$ , where  $\rho$  is the correlation between the two species' responses to the environment, and  $\sigma^2$  is the variance of environmental responses (shared by both species for maximum simplicity).

Utilizing the general mathematical expression of the storage effect (Eq.7), we find that the storage effect of species 1 is

$$\Delta I_1 \approx \sigma^2 \left( s_2 (1 - s_1) - \rho \, s_1 (1 - s_1) \right). \tag{17}$$

Setting  $\Delta I_1$  equal to zero and solving for  $s_2$ , we find that  $s_2 = \rho s_1$ , which has a solution when  $\rho$  is positive. This shows that the storage effect can be zero even when there are species-specific responses to the environment (i.e.,  $\rho < 1$ ), subadditivity (i.e.,  $s_i > 0$ ), and covariance between environment and competition (i.e.,  $\sigma^2 \neq 0$ ), as claimed in Section 2.3.

Coexistence mechanisms are often divided by species' sensitivity to competition, here operationalized as  $|\beta_i^{(1)}|$ . This scaled version of species 1's storage effect is

$$\frac{\Delta I_1}{(1-s_1)} \approx \sigma^2 \left(s_2 - \rho \ s_1\right). \tag{18}$$

Following the definitions of coexistence mechanisms (Eq.6), the sum of the scaled storage effect and the scaled relative nonlinearity for species 1 is

$$\frac{\Delta I_1}{(1-s_1)} + \frac{\Delta N_1}{(1-s_1)} \approx \sigma^2 s_1 (1-\rho), \tag{19}$$

which clearly increases with invader survival, as claimed in the main text (Section 3.1).

In the two-species lottery model, the scaled community-average storage effect (Eq.8) is

$$\overline{\left(\frac{\Delta I}{1-s}\right)} = \frac{\sigma^2}{2}(s_1+s_2)(1-\rho), \tag{20}$$

which demonstrates that at the community-level, survival increases the (community-average) storage effect, as claimed in Section 4.

## 6.2 An alternative definition of the storage effect

An alternative definition of the storage effect could use an *invader-invader comparison* (Johnson and Hastings, 2022). Instead of comparing an invader to residents (as in Eq.5), we could compare a single focal species to itself at high versus low density. Using this alternative definition of the storage effect in the two species lottery model, we would find that the storage effect increases with adult survival: when going from the high-density state to the low-density state, the *EC* covariance decreases (if there are species-specific responses to the environment) but the survival parameter is unchanged.

There are problems with the invader–invader comparison that make it a poor method for defining coexistence mechanisms generally: it is not defined when the invasion growth rate is negative (a community with the focal species at high density cannot be prepared for the purpose of the comparison), and it does not necessarily capture the notion of specialization/differentiation. However, even if these problems did not exist, the invader–invader comparison would not validate the conventional interpretation v2. The invader–invader comparison does not say that buffering helps rare species. Rather, the invader–invader comparison shows that buffering helps a rare species relative to said species at high density: it is as much about self-limitation when abundant as it is about recovery when rare.

# 7 References

# References

- Abrams, P. A. (1988). How should resources be counted? *Theoretical Population Biology*, 33(2), 226–242.
- Adler, P. (2014). Testing the storage effect with long term observational data. Temporal dynamics and ecological process, 82–101.
- Adler, P. B., & Drake, J. M. (2008). Environmental variation, stochastic extinction, and competitive coexistence. The American Naturalist, 172(5), E186–E195.
- Adler, P. B., HilleRisLambers, J., Kyriakidis, P. C., Guan, Q., & Levine, J. M. (2006). Climate variability has a stabilizing effect on the coexistence of prairie grasses. *Proceedings of the National Academy of Sciences*, 103(34), 12793–12798.
- Aksnes, D., & Egge, J. (1991). A theoretical model for nutrient uptake in phytoplankton. Marine ecology progress series. Oldendorf, 70(1), 65–72.
- Angert, A. L., Huxman, T. E., Chesson, P., & Venable, D. L. (2009). Functional tradeoffs determine species coexistence via the storage effect. *Proceedings of the National Academy of Sciences*, 106(28), 11641–11645.
- Armitage, D. W., & Jones, S. E. (2019). Negative frequency-dependent growth underlies the stable coexistence of two cosmopolitan aquatic plants. *Ecology*, 100(5), e02657.
- Armitage, D. W., & Jones, S. E. (2020). Coexistence barriers confine the poleward range of a globally distributed plant. *Ecology Letters*, 23(12), 1838–1848.

- Armstrong, R. A., & McGehee, R. (1980). Competitive exclusion. The American Naturalist, 115(2), 151–170.
- Barabás, G., D'Andrea, R., & Stump, S. M. (2018). Chesson's coexistence theory. *Ecological Mono-graphs*, 88(3), 277–303.
- Baskin, C. C., & Baskin, J. M. (1998). Seeds: Ecology, biogeography, and, evolution of dormancy and germination. Elsevier.
- Cáceres, C. E. (1997). Temporal variation, dormancy, and coexistence: A field test of the storage effect. Proceedings of the National Academy of Sciences, 94(17), 9171–9175.
- Caswell, H. (2001). *Matrix population models: Construction, analysis, and interpretation* (2nd ed.). Sinauer Associates.
- Chesson, P. L., & Ellner, S. (1989). Invasibility and stochastic boundedness in monotonic competition models. Journal of Mathematical Biology, 27(2), 117–138.
- Chesson, P. (1991). A need for niches? Trends in ecology & evolution, 6(1), 26–28.
- Chesson, P. (1994). Multispecies Competition in Variable Environments. Theoretical Population Biology, 45(3), 227–276.
- Chesson, P. (2000a). General theory of competitive coexistence in spatially-varying environments. Theoretical Population Biology, 58(3), 211–237.
- Chesson, P. (2000b). Mechanisms of maintenance of species diversity. Annual review of Ecology and Systematics, 31(1), 343–366.
- Chesson, P. (2003). Quantifying and testing coexistence mechanisms arising from recruitment fluctuations. *Theoretical Population Biology*, 64(3), 345–357.
- Chesson, P. (2018). Updates on mechanisms of maintenance of species diversity. *Journal of Ecology*, 106(5), 1773–1794.
- Chesson, P. (2019). Chesson's coexistence theory: Comment. Ecology,  $\theta(0)$ , 1–7.
- Chesson, P., Gebauer, R. L., Schwinning, S., Huntly, N., Wiegand, K., Ernest, M. S., Sher, A., Novoplansky, A., & Weltzin, J. F. (2004). Resource pulses, species interactions, and diversity maintenance in arid and semi-arid environments. *Oecologia*, 141(2), 236–253.
- Chesson, P., & Huntly, N. (1989). Short-term instabilities and long-term community dynamics. Trends in Ecology & Evolution, 4(10), 293–298.
- Chesson, P., & Huntly, N. (1997). The roles of harsh and fluctuating conditions in the dynamics of ecological communities. The American Naturalist, 150(5), 519–553.
- Chesson, P., Huntly, N. J., Roxburgh, S. H., Pantastico-Caldas, M., & Facelli, J. M. (2012). The storage effect: Definition and tests in two plant communities. *Temporal dynamics and ecological process* (pp. 11–40). Cambridge University Press.
- Chesson, P., & Kuang, J. J. (2010). The storage effect due to frequency-dependent predation in multispecies plant communities. *Theoretical Population Biology*, 78(2), 148–164.
- Chesson, P. L. (1982). The stabilizing effect of a random environment. *Journal of Mathematical Biology*, 15(1), 1–36.
- Chesson, P. L. (1983). Coexistence of competitors in a stochastic environment: The storage effect. *Population biology* (pp. 188–198). Springer.
- Chesson, P. L. (1984). The storage effect in stochastic population models. *Mathematical ecology* (pp. 76–89). Springer.
- Chesson, P. L. (1985). Coexistence of competitors in spatially and temporally varying environments: A look at the combined effects of different sorts of variability. *Theoretical Population Biology*, 28(3), 263–287.
- Chesson, P. L. (1988). Interactions between environment and competition: How fluctuations mediate coexistence and competitive exclusion. *Community ecology* (pp. 51–71). Springer.
- Chesson, P. L., & Huntly, N. (1988). Community consequences of life-history traits in a variable environment. Annales Zoologici Fennici, 5–16.
- Chesson, P. L., & Warner, R. R. (1981). Environmental variability promotes coexistence in lottery competitive systems. The American Naturalist, 117(6), 923–943.
- Cushing, D. (1971). The dependence of recruitment on parent stock in different groups of fishes. *ICES Journal of Marine Science*, 33(3), 340–362.

- Dean, A. M., & Shnerb, N. M. (2020). Stochasticity-induced stabilization in ecology and evolution: a new synthesis. *Ecology*, 101(9), 1–14.
- Dempster, E. R. (1955). Maintenance of genetic heterogeneity. Cold Spring Harbor symposia on quantitative biology, 20.
- Descamps-Julien, B., & Gonzalez, A. (2005). Stable coexistence in a fluctuating environment: An experimental demonstration. *Ecology*, 86(10), 2815–2824.
- Ellner, S. (1985a). Ess germination strategies in randomly varying environments. i. logistic-type models. *Theoretical Population Biology*, 28(1), 50–79.
- Ellner, S. (1985b). Ess germination strategies in randomly varying environments. ii. reciprocal yield-law models. *Theoretical Population Biology*, 28(1), 80–116.
- Ellner, S. (1987). Alternate plant life history strategies and coexistence in randomly varying environments. Vegetatio, 69(1), 199–208.
- Ellner, S. P., Snyder, R. E., & Adler, P. B. (2016). How to quantify the temporal storage effect using simulations instead of math. *Ecology letters*, 19(11), 1333–1342.
- Ellner, S. P., Snyder, R. E., Adler, P. B., & Hooker, G. (2019). An expanded modern coexistence theory for empirical applications. *Ecology Letters*, 22(1), 3–18.
- Eyring, H. (1935). The activated complex in chemical reactions. The Journal of Chemical Physics, 3(2), 107–115.
- Facelli, J. M., Chesson, P., & Barnes, N. (2005). Differences in seed biology of annual plants in arid lands: A key ingredient of the storage effect. *Ecology*, 86(11), 2998–3006.
- Gause, G. F. (1934). The struggle for existence. Williams & Wilkins.
- Gelman, A., & Hill, J. (2007). Data analysis using regression and multilevelhierarchical models (Vol. 1). Cambridge University Press New York, NY, USA.
- Gillespie, J. H. (1977). Natural Selection for Variances in Offspring Numbers: A New Evolutionary Principle. *The American Naturalist*, 111(981), 1010–1014.
- Godfrey-Smith, P. (2009). Darwinian populations and natural selection. Oxford University Press.
- Grainger, T. N., Levine, J. M., & Gilbert, B. (2019). The Invasion Criterion: A Common Currency for Ecological Research. Trends in Ecology and Evolution, 34 (10), 925–935.
- Grinell, J. (1917). The niche relationship of california thrsher. Auk, 1, 64–82.
- Gulisija, D., Kim, Y., & Plotkin, J. B. (2016). Phenotypic plasticity promotes balanced polymorphism in periodic environments by a genomic storage effect. *Genetics*, 202(4), 1437–1448.
- Haigh, J., & Smith, J. M. (1972). Can there be more predators than prey? Theoretical Population Biology, 3(3), 290–299.
- Haldane, J. B., & Jayakar, S. D. (1963). Polymorphism due to selection of varying direction. Journal of Genetics, 58(2), 237–242.
- Hallett, L. M., Shoemaker, L. G., White, C. T., & Suding, K. N. (2019). Rainfall variability maintains grass-forb species coexistence. *Ecology Letters*, 22(10), 1658–1667.
- Harper, J. L., & White, J. (1971). The dynamics of plant populations. Proc. Adv. Study Inst. Dynamics Numbers Popul. (Oosterbeek), (September 1970), 41–63.
- Holt, G., & Chesson, P. (2014). Variation in moisture duration as a driver of coexistence by the storage effect in desert annual plants. *Theoretical Population Biology*, 92, 36–50.
- Hutchinson, G. E. (1959). Homage to Santa Rosalia or Why are there so many kinds of animals? 93(870).
- Hutchinson, G. E. (1961). The paradox of the plankton. The American Naturalist, 95 (882), 137-145.
- Ignace, D. D., Huntly, N., & Chesson, P. (2018). The role of climate in the dynamics of annual plants in a chihuahuan desert ecosystem. *Evolutionary Ecology Research*, 19(3), 279–297.
- Jiang, L., & Morin, P. J. (2007). Temperature fluctuation facilitates coexistence of competing species in experimental microbial communities. *Journal of animal ecology*, 76(4), 660–668.
- Johnson, E. C., & Hastings, A. (2022). Methods for calculating coexistence mechanisms: Beyond scaling factors. Unpublished Manuscript.
- Kahane, G. (2011). Evolutionary debunking arguments. Noûs, 45(1), 103–125.
- Kelly, C. K., & Bowler, M. G. (2002). Coexistence and relative abundance in forest trees. Nature, 417(6887), 437–440.

- Kelly, C. K., & Bowler, M. G. (2005). A new application of storage dynamics: Differential sensitivity, diffuse competition, and temporal niches. *Ecology*, 86(4), 1012–1022.
- Kingsolver, J. G. (2009). The well-temperatured biologist: (american society of naturalists presidential address). The American Naturalist, 174(6), 755–768.
- Klausmeier, C. A. (2010). Successional state dynamics: A novel approach to modeling nonequilibrium foodweb dynamics. *Journal of Theoretical Biology*, 262(4), 584–595.
- Kuang, J. J., & Chesson, P. (2009). Coexistence of annual plants: Generalist seed predation weakens the storage effect. *Ecology*, 90(1), 170–182.
- Kuang, J. J., & Chesson, P. (2010). Interacting coexistence mechanisms in annual plant communities: Frequency-dependent predation and the storage effect. *Theoretical Population Biology*, 77(1), 56–70.
- Lande, R. (1998). Demographic stochasticity and allee effect on a scale with isotropic noise. *Oikos*, 353–358.
- Letten, A. D., Dhami, M. K., Ke, P.-J., & Fukami, T. (2018). Species coexistence through simultaneous fluctuation-dependent mechanisms. *Proceedings of the National Academy of Sciences*, 115(26), 6745–6750.
- Levin, S. A. (1970). Community equilibria and stability, and an extension of the competitive exclusion principle. The American Naturalist, 104(939), 413–423.
- Lewontin, R. C., & Cohen, D. (1969). On population growth in a randomly varying environment. Proceedings of the National Academy of Sciences of the United States of America, 62(4), 1056–1060.
- Li, L., & Chesson, P. (2016). The effects of dynamical rates on species coexistence in a variable environment: The paradox of the plankton revisited. *The American Naturalist*, 188(2), E46– E58.
- Lieth, H., & Whittaker, R. H. (1973). Primary productivity of the biosphere. Springer.
- Lloyd, M., & Dybas, H. S. (1966). the Periodical Cicada Problem. I. Population Ecology. Evolution, 20(2), 133–149.
- Loreau, M. (1989). Coexistence of temporally segregated competitors in a cyclic environment. Theoretical Population Biology, 36(2), 181–201.
- Loreau, M. (1992). Time scale of resource dynamics and coexistence through time partitioning. Theoretical Population Biology, 41(3), 401–412.
- Lotka, A. J. (1932). The growth of mixed populations: Two species competing for a common food supply. Journal of the Washington Academy of Sciences, 22(16-17), 461–469.
- MacArthur, R. H. (1967). The theory of island biogeography. Princeton University Press.
- May, R. M. (1974). On the theory of niche overlap. *Theoretical Population Biology*, 5(3), 297–332.
- Messer, P. W., Ellner, S. P., & Hairston Jr, N. G. (2016). Can population genetics adapt to rapid evolution? *Trends in Genetics*, 32(7), 408–418.
- Meszéna, G., Gyllenberg, M., Pásztor, L., & Metz, J. A. (2006). Competitive exclusion and limiting similarity: A unified theory. *Theoretical Population Biology*, 69(1), 68–87.
- Metz, J., Nisbet, R., & Geritz, S. (1992). How should we define 'fitness' for general ecological scenarios? Trends in Ecology & Evolution, 7(6), 198–202.
- Miller, E. T., & Klausmeier, C. A. (2017). Evolutionary stability of coexistence due to the storage effect in a two-season model. *Theoretical Ecology*, 10(1), 91–103.
- Monod, J. (1949). The growth of bacterial cultures. Annual review of microbiology, 3(1), 371–394.
- Pake, C. E., & Venable, D. L. (1995). Is coexistence of sonoran desert annuals mediated by temporal variability reproductive success. *Ecology*, 76(1), 246–261.
- Pake, C. E., & Venable, D. L. (1996). Seed banks in desert annuals: Implications for persistence and coexistence in variable environments. *Ecology*, 77(5), 1427–1435.
- Pande, J., Fung, T., Chisholm, R., & Shnerb, N. M. (2020). Mean growth rate when rare is not a reliable metric for persistence of species. *Ecology letters*, 23(2), 274–282.
- Reinhold, K. (2000). Maintenance of a genetic polymorphism by fluctuating selection on sex-limited traits. Journal of Evolutionary Biology, 13(6), 1009–1014.
- Rosenzweig, M. L. (1968). Net primary productivity of terrestrial communities: Prediction from climatological data. The American Naturalist, 102(923), 67–74.

- Sala, O. E., Parton, W. J., Joyce, L., & Lauenroth, W. (1988). Primary production of the central grassland region of the united states. *Ecology*, 69(1), 40–45.
- Sale, P. F. (1977). Maintenance of High Diversity in Coral Reef Fish Communities. The American Naturalist, 111 (978), 337–359.
- Schreiber, S. J. (2021). Positively and negatively autocorrelated environmental fluctuations have opposing effects on species coexistence. The American Naturalist, 197(4), 000–000.
- Schreiber, S. J., Yamamichi, M., & Strauss, S. Y. (2019). When rarity has costs: Coexistence under positive frequency-dependence and environmental stochasticity. *Ecology*, 100(7), e02664.
- Sears, A. L., & Chesson, P. (2007). New methods for quantifying the spatial storage effect: An illustration with desert annuals. *Ecology*, 88(9), 2240–2247.
- Snyder, R. (2012). Storage effect. Encyclopedia of theoretical ecology (1st ed., pp. 722–726). University of California Press.
- Song, C., Von Ahn, S., Rohr, R. P., & Saavedra, S. (2020). Towards a Probabilistic Understanding About the Context-Dependency of Species Interactions. *Trends in Ecology and Evolution*, 35(5), 384–396.
- Srinivasan, A. (2015). The archimedean urge. *Philosophical Perspectives*, 29, 325–362.
- Stump, S. M., & Chesson, P. (2017). How optimally foraging predators promote prey coexistence in a variable environment. *Theoretical Population Biology*, 114, 40–58.
- Szuwalski, C. S., Vert-Pre, K. A., Punt, A. E., Branch, T. A., & Hilborn, R. (2015). Examining common assumptions about recruitment: A meta-analysis of recruitment dynamics for worldwide marine fisheries. *Fish and Fisheries*, 16(4), 633–648.
- Towers, I. R., Bowler, C. H., Mayfield, M. M., & Dwyer, J. M. (2020). Requirements for the spatial storage effect are weakly evident for common species in natural annual plant assemblages. *Ecology*, 101(12), e03185.
- Turelli, M. (1978). A reexamination of stability in randomly varying versus deterministic environments with comments on the stochastic theory of limiting similarity. *Theoretical Population Biology*, 13(2), 244–267.
- Usinowicz, J., Chang-Yang, C.-H., Chen, Y.-Y., Clark, J. S., Fletcher, C., Garwood, N. C., Hao, Z., Johnstone, J., Lin, Y., Metz, M. R., Masaki, T., Nakashizuka, T., Sun, I.-F., Valencia, R., Wang, Y., Zimmerman, J. K., Ives, A. R., & Wright, S. J. (2017). Temporal coexistence mechanisms contribute to the latitudinal gradient in forest diversity. *Nature*, 550(7674), 105– 108.
- Usinowicz, J., Wright, S. J., & Ives, A. R. (2012). Coexistence in tropical forests through asynchronous variation in annual seed production. *Ecology*, 93(9), 2073–2084.
- Venable, D. L., Pake, C. E., & Caprio, A. C. (1993). Diversity and coexistence of sonoran desert winter annuals. *Plant Species Biology*, 8(2-3), 207–216.
- Volterra, V. (1926). Variations and fluctuations of the number of individuals in animal species living together. Animal Ecology, 409–448.
- Warner, R. R., & Chesson, P. L. (1985). Coexistence mediated by recruitment fluctuations: A field guide to the storage effect. *The American Naturalist*, 125(6), 769–787.
- Warner, R. R., & Hoffman, S. G. (1980). Population density and the economics of territorial defense in a coral reef fish. *Ecology*, 61(4), 772–780.
- Wittgenstein, L. (1968). Philosophical investigations. Basil Blackwell.
- Yamamichi, M., & Hoso, M. (2017). Roles of maternal effects in maintaining genetic variation: Maternal storage effect. *Evolution*, 71(2), 449–457.
- Yuan, C., & Chesson, P. (2015). The relative importance of relative nonlinearity and the storage effect in the lottery model. *Theoretical Population Biology*, 105, 39–52.
- Zepeda, V., & Martorell, C. (2019). Fluctuation-independent niche differentiation and relative nonlinearity drive coexistence in a species-rich grassland. *Ecology*, 100(8), e02726.