

# Black Holes in a Cavity: Heat engine and Joule-Thomson Expansion

Yihe Cao,<sup>\*</sup> Hanwen Feng,<sup>†</sup> Jun Tao,<sup>‡</sup> and Yadong Xue<sup>§</sup>

*Center for Theoretical Physics, College of Physics,  
Sichuan University, Chengdu, 610065, China*

## Abstract

We consider the charged  $d$ -dimensional black holes in a cavity in extended phase space and investigate the heat engine and the Joule-Thomson (JT) expansion. Since the phase structure of black holes in a cavity is similar to anti-de-sitter (AdS) cases, we take black holes in a cavity as the working substance in the heat engine and calculate their efficiency in Carnot cycle and rectangular cycle. Also, we discuss whether the JT expansion of charged black holes in a cavity are consistent with AdS cases and conclude the effect of different boundary conditions on black hole thermodynamics.

---

<sup>\*</sup>Electronic address: yihe@stu.scu.edu.cn

<sup>†</sup>Electronic address: fenghanwen@stu.scu.edu.cn

<sup>‡</sup>Electronic address: taojun@scu.edu.cn

<sup>§</sup>Electronic address: xueyadong@stu.scu.edu.cn

## I. INTRODUCTION

The thermal quantum radiance of black hole reveals the connection among gravity, quantum theory and black hole thermodynamics [1–3]. The idea to place black holes in a cavity is proposed to solve the problem of investigating thermodynamically unstable black hole in asymptotically flat space, and it attracts interests in the study of black hole thermodynamics and quantum gravity [4]. The temperature of the surface of a cavity is fixed with a heat bath surrounded. According to the semi-classical asymptotic relationship between the on-shell Euclidean action and the partition function, we can obtain the expression of free energy. Furthermore, we define the volume of cavity as the one of the system. In this framework, the phase transition, the thermodynamic geometry and the critical behaviour of vary BHs in a cavity are discussed in [5–13]. Another boundary condition to make the black hole thermodynamically stable is to put it in the anti-de sitter(AdS) space, and related work on black hole thermodynamics are studied in [14–21]. These results shows the thermodynamic behaviours of AdS-BHs are similar to those of BHs in cavities [6].

Due to this fact, we can investigate its properties as a heat engine like AdS-BH, which is to define a cycle in the extended thermodynamical space and the AdS black holes are considered as the working substance, and the related work can be found in [22–37]. Moreover, a cavity has explicit boundaries, which makes the correspondence between the classic heat engine and the black hole one more reasonable. We considered charged black holes in a cavity as the heat engine which works in the Carnot cycle and rectangular cycle. The Carnot cycle consists of two isothermal paths and two adiabatic paths, it is always theoretically highest according to the second law of thermodynamics, and thus it provides an upper bound for us to check the calculation. The rectangular cycle is made up of two isochoric paths and two isobaric paths, which can be taken as the smallest unit of a more complicated cycle [37].

As an important property of van der Waals system, the Joule-Thomson (JT) expansion is introduced to AdS-BHs and firstly investigated in [38], which shows both cooling and heating states of the RN-AdS black hole exist during the isenthalpic process. Furthermore, it's found that the JT expansion for different black holes were found have similarities in isenthalpic curves and inversion curves [39–69], which means the isenthalpic curves divides the heating and the cooling regions and the extreme points are coincident with the inversion curves. Moreover, [70] shows different types of isenthalpic curves when it comes to study the

JT expansion of the FRW universe. In this paper, we prove the JT expansion of black holes in a cavity does exist, but differ from AdS-BHs. We find the JT coefficient always keeps positive, which means the temperature decrease during this kind of adiabatic expansion. In other words, the black hole can only become colder after the isenthalpic process. And it is different from the situations ever investigated before.

In section II, we derive the thermodynamic quantities of the d-dimensional charged BH in the cavities. We calculate the heat engine efficiency of 4-dimensional Schwarzschild black holes and RN black holes in a cavity in section III. In section IV, the JT expansion of the RN black holes and the higher dimensional cases in a cavity are investigated. We summarize our results in section V.

## II. BLACK HOLES IN A CAVITY

The  $d$ -dimensional Einstein-Maxwell action according to the ref [71, 72] is given by

$$I = I_{bulk} + I_{surf} = -\frac{1}{16\pi} \int_{\mathcal{M}} d^d x \sqrt{-g} R + \frac{1}{8\pi} \int_{\partial\mathcal{M}} d^{d-1} x \sqrt{-\gamma} (K - K_0) + \frac{1}{16\pi} \int_{\mathcal{M}} d^d x \sqrt{-g} F^2 + \frac{1}{16\pi} \int_{\partial\mathcal{M}} d^{d-1} x \sqrt{-\gamma} n_\nu F^{\mu\nu} A_\mu, \quad (1)$$

where  $n_\nu$  is the unit normal vector of the boundary,  $\gamma$  is the metric on the boundary,  $K$  is the trace of the extrinsic curvature, and  $K_0$  is a subtraction term to ensure the Gibbons-Hawking-York term vanish in flat space-time. The metric of the spherical d-dimensional charged BH reads [73]

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2 d\omega_{d-2}^2, \quad A_t dt = -\sqrt{\frac{d-2}{2d-6}} \frac{Q}{r^{d-3}} dt. \quad (2)$$

By varying the action Eq.(1), we have the metric function as

$$f(r) = 1 + \frac{Q^2}{r^{2(d-3)}} - \frac{Q^2}{r^{d-3} r_+^{d-3}} - \frac{r_+^{d-3}}{r^{d-3}}, \quad (3)$$

where the radius of the event horizon is defined by  $f(r_+) = 0$  and the parameter  $Q$  is the charge of the black hole. Corresponding to the Hawking temperature  $T_H$  by  $T = \frac{T_H}{\sqrt{f(r_B)}}$  in [5], the temperature of a cavity is defined as

$$T = \frac{(d-3)(1 - Q^2 r_+^{6-2d})}{4\pi r_+ \sqrt{f(r_B)}}, \quad (4)$$

where  $r_B$  is the radius of a cavity. Furthermore, the Euclidean action corresponds to the action Eq.(1) by  $I_E = iI$ . The Euclidean time  $\tau$  is related to  $t$  by  $\tau = it$  and the metric becomes positive infinite [4, 8]. In this way, we derive the Euclidean action by substituting Euclidean metric into the action, which gives

$$I_E = \frac{d-2}{8\pi} \frac{\omega_{d-2} r_B^{d-3}}{T} (1 - \sqrt{f(r_B)}) - S, \quad (5)$$

where  $S = \frac{\omega_{d-2}}{4} r_+^{d-2}$  is the entropy of the black hole.  $\omega_d = \frac{2\pi^{\frac{d+1}{2}}}{\Gamma(\frac{d+1}{2})}$  is the volume of the unit  $d$ -sphere where  $\Gamma$  represents the gamma function. The free energy is related to the Euclidean action in an approximately semi-classical way as  $F = -T \ln Z = T I_E$  [7], and thus the thermal energy of the black hole in a cavity is  $E = -T^2 \frac{\partial F/T}{\partial T}$ , which yields as

$$E = \frac{(d-2) r_B^{d-3} \omega_{d-2} \left(1 - \sqrt{f(r_B)}\right)}{8\pi}. \quad (6)$$

The thermodynamic volume  $V$  of the system is defined as the volume of the cavity

$$V \equiv \frac{r_B^{d-1} \omega_{d-2}}{d-1}, \quad (7)$$

and the conjugate thermodynamic pressure naturally arises according to  $P = -\frac{\partial E}{\partial V}$ . In this new extended phase space, the enthalpy  $H$  can be derived by  $H = E + PV$ .

### III. BLACK HOLES AS HEAT ENGINES IN A CAVITY

For the purpose of comparing the heat engine properties of black hole in a cavity with AdS-BH, we consider a cavity as the container and take black holes as the working substance in the pressure-volume space.

#### A. Schwarzschild black holes in a cavity

In the following sections, the black hole is set in thermodynamical cycles. For simplicity, we focus on the case of  $d = 4$ . Four-dimensional space-time metric of RN black hole yields

$$ds^2 = -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (8)$$

where  $f(r)$  is expressed as

$$f(r) = \left(1 - \frac{r_+}{r}\right) \left(1 - \frac{Q_b^2}{r_+ r}\right), \quad (9)$$

and the effective potential satisfies  $A = A_t dt = -\frac{Q_b}{r} dt$ . The event horizon radius is represented by  $r_+$ , meanwhile the charge of the black hole is represented by  $Q_b$ . Subsequently the temperature of this system with radius  $r = r_B > r_+$  had a definition in [5], and for  $d = 4$ ,

$$T(r_B, x, Q) = \frac{T_H}{\sqrt{f(r_B)}} = \frac{1 - \frac{Q^2}{r_B^2 x^2}}{4\pi r_B x \sqrt{f(r_B)}}, \quad (10)$$

where  $Q_b = Q$ ,  $r_B$  is the radius of a cavity and  $T_H = \frac{1}{4\pi} f'(r_+)$  is the Hawking temperature of the black hole. It is worth noting that the physical range of the event horizon is constrained as  $\frac{r_e}{r_B} \leq x \equiv \frac{r_+}{r_B} \leq 1$  in which  $r_e = Q$  is the horizon radius of the extremal RN-BH.

First, we start from the Schwarzschild black holes, which is the form of RN black holes reduce to when  $Q = 0$ . Under these conditions, Eq. (10) takes the form as

$$T(r_B, x) = \frac{T_H}{\sqrt{f(r_B)}} = \frac{1}{4\pi r_B x \sqrt{1-x}}. \quad (11)$$

and the conjugate pressure is

$$P(r_B, x) = \frac{2r_B^2 x - r_B^2 x^2}{8\pi r_B^4 x \sqrt{1-x}} - \frac{1}{4\pi r_B^2}. \quad (12)$$

We can derive the expression for  $x$  in terms of  $r_B$  and  $P$  with Eq. (12),

$$x = 4 \left( (4\pi P r_B^3 + r_B) \sqrt{2\pi P (2\pi P r_B^2 + 1)} - 4\pi P r_B^2 (2\pi P r_B^2 + 1) \right). \quad (13)$$

With this result, the temperature can be rewritten as

$$T(r_B, P) = (16\pi r_B \Gamma \sqrt{1+4\Gamma})^{-1}, \quad (14)$$

$$\Gamma = \sqrt{2\pi} \sqrt{P (2\pi P r_B^2 + 1) (4\pi P r_B^3 + r_B)^2 - 4\pi P r_B^2 (2\pi P r_B^2 + 1)}. \quad (15)$$

We denote the heat absorbed as  $Q_H$ , and the heat delivered as  $Q_C$ , so that the mechanical work is  $W = Q_H - Q_C$ . The efficiency is the ratio of mechanical work  $W$  to heat absorbed  $Q_H$

$$\eta = \frac{W}{Q_H} = 1 - \frac{Q_C}{Q_H}. \quad (16)$$

The Carnot cycle consists of two isothermal paths and two adiabatic paths which means that the Carnot heat engine is between two different temperatures. We define the high temperature as  $T_H(r_{B2}, P_2)$  and the cold one as  $T_L(r_{B1}, P_1)$ , which are connected through the isochoric paths. The efficiency yields as

$$\eta = 1 - \frac{T_L(r_{B1}, P_1)}{T_H(r_{B4}, P_4)}, \quad (17)$$




FIG. 1: The Carnot cycle efficiency of Schwarzschild black hole heat engine with varying radius  $R$ , where we set  $r_{B1} = 1$ ,  $P_1 = 1$ ,  $P_4 = 4$ .

by which we plot how it varies with respect to the radius of a cavity in FIG. 1.

Moreover, we define a rectangular cycle with two isochoric and two isobaric paths and compare its efficiency with that of Carnot cycle, and the efficiency for the rectangle cycle can be calculated by a formula deduced in [74, 75]

$$\eta = \frac{W}{Q} = \frac{(P_1 - P_4)(V_4 - V_1)}{Q}, \quad (18)$$

where

$$Q = \Delta E + P\Delta V. \quad (19)$$

Thus the Eq. (18) yield as

$$\eta = \frac{4\pi(P_1 - P_4)(r_{B1}^3 - r_{B4}^3)}{r_{B4}(-4\pi P_1 r_{B4}^2 + 3\Delta_4 - 3) + 4\pi P_1 r_{B1}^3 - 3r_{B1}(\Delta_1 - 1)}, \quad (20)$$

where we define

$$\Delta_\alpha = \sqrt{16\pi P_\alpha r_{B\alpha}^2 (2\pi P_\alpha r_{B\alpha}^2 + 1) - 4\sqrt{2\pi} \sqrt{P_\alpha (2\pi P_\alpha r_{B\alpha}^2 + 1) (4\pi P_\alpha r_{B\alpha}^3 + r_{B\alpha})^2 + 1}}.$$

and  $\alpha = 1, 4$ .

In this way, we can show how the efficiency changes as well.

## B. RN black holes in a cavity

Then we discuss a more complicated situation. Different from the case of Schwarzschild, we can't analytically derive the expression of  $x$  in terms of  $r_B$  and  $P$ . Alternatively, we calculate the efficiency numerically.




FIG. 2: The rectangular cycle efficiency in the Schwarzschild case with varying radius  $R$ , where we set  $r_{B1} = 1$ ,  $P_1 = 1$ ,  $P_4 = 4$

In the 4-dimension, the conjugate thermodynamic pressure is

$$P(r_B, x, Q) = -\frac{\partial E}{\partial V} = \frac{2r_B^2 x - Q^2 - r_B^2 x^2}{8\pi r_B^4 x \sqrt{f(r_B)}} - \frac{1}{4\pi r_B^2}. \quad (21)$$

For every pressure we have fixed, we can solve the corresponding  $x$  by Eq. (21) with conditions  $0 < x \leq 1$  and  $x > \frac{Q^2}{R^2}$  which guarantee  $P$  and  $T$  are real, and then we can obtain the temperature with  $x$  to derive the efficiency of Carnot cycle. Similarly, we can plot the efficiency curve in the rectangular cycle.




FIG. 3: Left Plane: The Carnot cycle efficiency of RN black hole heat engine; Right Plane: The rectangular cycle efficiency of RN black hole heat engine. Here we set  $r_{B1} = 1$ ,

$$P_1 = 0.01, P_4 = 0.02$$

#### IV. THE JT EXPANSION OF RN BLACK HOLE IN A CAVITY

The JT expansion remains a characteristic of the Van der Waals system and the system undergoes an isenthalpic process. As the Joule-Thomson expansion takes place in an isenthalpic process, the enthalpy has an important role in the issues.

First of all, four dimensional temperature graph in FIG.4 according to Eq.(4) shows how the temperature  $T$  varies with respect to  $x$ . It is similar to the temperature graph of RN-AdS black holes [19] only with one difference, that is when  $x$  approaches 1, the temperature diverges. That will be unexpected to have a infinitely high temperature, so we would like to investigate with lower value of  $x$ .




FIG. 4: We set  $Q = 0.035, 0.1, 0.2$  and  $0.35$  from top to the bottom. And  $r_B = 1$  while  $x$  have a range of  $[0, 1]$  in  $T - x$  plane.

For further study, we derive the relating thermodynamic quantities. The thermal energy in this system have been discussed in [76] and from Eq.(9), we obtain the equations of the thermal energy of the system, which yields

$$E(r_B, x, Q) = r_B(1 - \sqrt{f(r_B)}). \quad (22)$$

Moreover, the volume of the system is defined to be related to the radius of a cavity, which is

$$V = \frac{4}{3}\pi r_B^3, \quad (23)$$

whose conjugate thermodynamic pressure naturally arises as

$$P(r_B, x, Q) = -\frac{\partial E}{\partial V} = \frac{2r_B^2x - Q^2 - r_B^2x^2}{8\pi r_B^4x\sqrt{f(r_B)}} - \frac{1}{4\pi r_B^2}. \quad (24)$$



With the three equations above (22), (23) and (24), the enthalpy of the system is obtained by employing the relation  $H = E + PV$ ,

$$H(r_B, x, Q) = \frac{-r_B^2 x \left( 6f(r_B) - 4\sqrt{f(r_B)} + x - 2 \right) - Q^2}{6\sqrt{f(r_B)}r_B x}. \quad (25)$$

We have obtained all the quantities required and are able to study the JT expansion of RN black hole in a cavity. The isenthalpic curves can be obtained from Eqs.(10), (24) and (25) numerically in FIG. 5.




FIG. 5: Isenthalpic curves in  $T - P$  plane.  $H = 1, 2, 3$  and  $5$ .  $Q = 0.035$ ,  $d = 4$ , and  $r_B$  varies from  $0.2$  to  $10$ .

As we can see, there are no extreme points in the curves and the temperature decreases with lower pressure. Furthermore, when the pressure and temperature are high, the lines from bottom to the top show the isenthalpic curves of increasing value of the enthalpy,  $H = 1, 2, 3$  and  $5$ , which is obvious in FIG.5(a). Moreover, in FIG.5(b), we show the situation of lower temperature and pressure. The part of the curves that decreases as the enthalpy rise corresponds to smaller  $x$ , that is, when the ratio of event horizon  $r_+$  and cavity radius  $r_B$  is small, and it is in line with the range we want when we study the temperature curves. And the orange line coincide with the orange one in FIG.7 which represent the four dimensional cases when the enthalpy equals 1.

We can investigate the project further. For  $d$ -dimensional cases, the explicit form of  $P(r_+, r_B, q, d)$  and  $H(r_+, r_B, q, d)$  takes the form as

$$P = -\frac{\partial E}{\partial V} = \frac{(d-2)(d-3)}{8\pi}\Xi, \quad (26)$$

$$\Xi = -\frac{q^2}{2\sqrt{f(r_B)}r_B^{3d-6}r_+^{d-3}} - \frac{1}{2\sqrt{f(r_B)}r_B^{d-1}r_+} + \frac{1}{\sqrt{f(r_B)}r_B^2} - \frac{1}{r_B^2}.$$

The enthalpy  $H$  is derived by  $H = E + PV$ , which yields

$$H = \frac{(d-2)r_B^{d-3}\omega_{d-2}}{8\pi}(1 - \sqrt{f(r_B)}) + \frac{(d-3)r_B^2}{d-1}\Xi. \quad (27)$$

We obtain the isenthalpic curves in numerically way with Eqs.(4), (26) and (27), and then we plot how the enthalpy varies from 1 to 5 when  $d = 5, Q = 0.035$  in FIG.6.




FIG. 6: The isenthalpic curves of charged black hole in cavity.  $H = 1, 2, 3$  and  $5$ .  $d = 5$ ,  $Q = 0.035$ , and  $r_B$  varies from  $0.2$  to  $3$ .

The temperature decreases with pressure as shown in FIG.6, which shows that the process only cools the system. The graph can be divided with the curves and the left part represents the low pressure region, while the right part indicates the high pressure region. Additionally, from bottom to the top, the temperature in low pressure region increases with decreasing enthalpy and increases with the increasing enthalpy in the high pressure region. Similar to the four-dimensional case, in the low pressure region, the ratio of  $r_+$  and  $r_B$  is small.

After studying the four and five dimensional cases, we would like to consider the impact of dimensions on the isenthalpic curves.




FIG. 7: The isenthalpic curves of charged black holes in a cavity for high dimensional cases. We set the dimension  $d$  equals 4, 5, 6 and 7 respectively, the enthalpy  $H$  equals 1, and the charge  $Q$  equals 0.035.

The picture FIG.7 presents different dimensional cases where the red and the orange curve coincides with the orange one in FIG.6 and the orange one in FIG.5 respectively. We can see from the picture that, in the low pressure area, higher dimensions lead to higher temperatures, which is the opposite in the high pressure region. The influence of the dimension parameter  $d$  and the enthalpy  $H$  on the temperature during the throttling process is opposite.

As we know, the JT coefficient of classical ideal gas always equals 0, which means the isenthalpic process doesn't change the temperature of ideal gas, while the process can cool or heat the classical van der Waals gas only depending on the initial temperature and pressure. With Eqs.(10) and (24), we derive JT coefficient for 4 dimensional case by

$$\mu = \left(\frac{\partial T}{\partial P}\right)_H, \quad (28)$$

and show how it changes with  $r_B$  in FIG.8 with the red line. The JT coefficient  $\mu$  obtained numerically with Eqs.(4) and (26) in different dimensions are shown in FIG.8 too. When  $\mu = \left(\frac{\partial T}{\partial P}\right)_H = 0$ , the transition between cooling and heating region occurs. But for  $d$ -dimensional charged black holes in a cavity, the Joule-Thomson coefficient is positive infinite.

We conclude Joule-Thomson expansion is characterized by the invariance of enthalpy and Joule-Thomson coefficient  $\mu$  determines the final change of temperature during the process. One can use the sign of  $\mu$  to divide the heating and cooling zone. We can see from FIG.8 that

$\mu$  is always positive, which means no transition occurs. Comparing the isenthalpic curves above with the JT coefficient graph, the property of positive definite and the constant phase of cooling are obvious.




FIG. 8: The Joule-Thomson coefficient of the charged RN black hole in a cavity for  $d = 4, 5, 6$  and 7. The enthalpy  $H = 1$ ,  $r_B$  varies from 0.5 to 2.5, and  $Q = 0.035$ .

## V. CONCLUSIONS

Based on the thermodynamics of black holes in a cavity, we derive the temperature, pressure and other thermodynamic quantities in  $d$ -dimension to discuss the heat engine and the Joule-Thomson expansion of charged black holes. As a cavity defines an explicit boundary for the working substance, i.e. black holes, it's more rational to consider a heat engine in cavities rather than in AdS space, whose scale corresponds to the pressure in the extended space. Therefore, we derive the efficiency of Schwarzschild and RN black hole heat engines in the Carnot cycle and rectangular cycle and show how it changes with respect to the radius of a cavity. Moreover, we find the Joule-Thomson expansion of charged black holes exist in a cavity but differ from the AdS cases. We conclude the black hole always cools down during the isenthalpic process with the decreasing pressure, while the JT coefficient  $\mu$  keeps positive infinite.

## Acknowledgments

We are grateful to Yucheng Huang and Peng Wang for useful discussions and valuable comments. This work is supported by NSFC (Grant No.11947408 and 12047573).

- 
- [1] James W. York, Jr. Dynamical Origin of Black Hole Radiance. *Phys. Rev. D*, 28:2929, 1983.
  - [2] S. W. Hawking. Particle Creation by Black Holes. *Commun. Math. Phys.*, 43:199–220, 1975.
  - [3] J. D. Bekenstein. Statistical Black Hole Thermodynamics. *Phys. Rev. D*, 12:3077–3085, 1975.
  - [4] James W. York, Jr. Black hole thermodynamics and the Euclidean Einstein action. *Phys. Rev. D*, 33:2092–2099, 1986.
  - [5] Harry W. Braden, J. David Brown, Bernard F. Whiting, and James W. York, Jr. Charged black hole in a grand canonical ensemble. *Phys. Rev. D*, 42:3376–3385, 1990.
  - [6] Peng Wang, Houwen Wu, Haitang Yang, and Feiyu Yao. Extended Phase Space Thermodynamics for Black Holes in a Cavity. *JHEP*, 09:154, 2020.
  - [7] Peng Wang, Haitang Yang, and Shuxuan Ying. Thermodynamics and phase transition of a Gauss-Bonnet black hole in a cavity. *Phys. Rev. D*, 101(6):064045, 2020.
  - [8] Peng Wang, Houwen Wu, and Haitang Yang. Thermodynamics and Phase Transition of a Nonlinear Electrodynamics Black Hole in a Cavity. *JHEP*, 07:002, 2019.
  - [9] Peng Wang, Houwen Wu, and Haitang Yang. Thermodynamic Geometry of AdS Black Holes and Black Holes in a Cavity. *Eur. Phys. J. C*, 80(3):216, 2020.
  - [10] Peng Wang, Houwen Wu, and Shuxuan Ying. Validity of Thermodynamic Laws and Weak Cosmic Censorship for AdS Black Holes and Black Holes in a Cavity. *Chin. Phys. C*, 45(5):055105, 2021.
  - [11] Wen-Bo Zhao, Guo-Rong Liu, and Nan Li. Hawking–Page phase transitions of the black holes in a cavity. *Eur. Phys. J. Plus*, 136:981, 2021.
  - [12] Fil Simovic and Robert B. Mann. Critical Phenomena of Born-Infeld-de Sitter Black Holes in Cavities. *JHEP*, 05:136, 2019.
  - [13] Oscar J. C. Dias and Ramon Masachs. Charged black hole bombs in a Minkowski cavity. *Class. Quant. Grav.*, 35(18):184001, 2018.
  - [14] Marco M. Caldarelli, Guido Cognola, and Dietmar Klemm. Thermodynamics of Kerr-

- Newman-AdS black holes and conformal field theories. *Class. Quant. Grav.*, 17:399–420, 2000.
- [15] T. Padmanabhan. Classical and quantum thermodynamics of horizons in spherically symmetric space-times. *Class. Quant. Grav.*, 19:5387–5408, 2002.
- [16] David Kastor, Sourya Ray, and Jennie Traschen. Enthalpy and the Mechanics of AdS Black Holes. *Class. Quant. Grav.*, 26:195011, 2009.
- [17] S. W. Hawking and Don N. Page. Thermodynamics of Black Holes in anti-De Sitter Space. *Commun. Math. Phys.*, 87:577, 1983.
- [18] Andrew Chamblin, Roberto Emparan, Clifford V. Johnson, and Robert C. Myers. Charged AdS black holes and catastrophic holography. *Phys. Rev. D*, 60:064018, 1999.
- [19] Andrew Chamblin, Roberto Emparan, Clifford V. Johnson, and Robert C. Myers. Holography, thermodynamics and fluctuations of charged AdS black holes. *Phys. Rev. D*, 60:104026, 1999.
- [20] David Kubiznak and Robert B. Mann. P-V criticality of charged AdS black holes. *JHEP*, 07:033, 2012.
- [21] Song He, Li-Fang Li, and Xiao-Xiong Zeng. Holographic Van der Waals-like phase transition in the Gauss–Bonnet gravity. *Nucl. Phys. B*, 915:243–261, 2017.
- [22] Jie-Xiong Mo and Shan-Quan Lan. Phase transition and heat engine efficiency of phantom AdS black holes. *Eur. Phys. J. C*, 78(8):666, 2018.
- [23] K. V. Rajani, C. L. Ahmed Rizwan, A. Naveena Kumara, Deepak Vaid, and K. M. Ajith. Regular Bardeen AdS black hole as a heat engine. *Nucl. Phys. B*, 960:115166, 2020.
- [24] Ujjal Debnath. The General Class of Accelerating, Rotating and Charged Plebanski-Demianski Black Holes as Heat Engine. *arXiv.2006.02920*.
- [25] Clifford V. Johnson. de Sitter Black Holes, Schottky Peaks, and Continuous Heat Engines. *arXiv.1907.05883*.
- [26] Leonardo Balart and Sharmanthie Fernando. Non-linear black holes in 2+1 dimensions as heat engines. *Phys. Lett. B*, 795:638–643, 2019.
- [27] Pavan Kumar Yerra and Bhamidipati Chandrasekhar. Heat engines at criticality for nonlinearly charged black holes. *Mod. Phys. Lett. A*, 34(27):1950216, 2019.
- [28] H. Ghaffarnejad, E. Yaraie, M. Farsam, and K. Bamba. Hairy black holes and holographic heat engine. *Nucl. Phys. B*, 952:114941, 2020.
- [29] Jialin Zhang, Yanjun Li, and Hongwei Yu. Accelerating AdS black holes as the holographic

- heat engines in a benchmarking scheme. *Eur. Phys. J. C*, 78(8):645, 2018.
- [30] Jialin Zhang, Yanjun Li, and Hongwei Yu. Thermodynamics of charged accelerating AdS black holes and holographic heat engines. *JHEP*, 02:144, 2019.
- [31] Pavan Kumar Yerra and Chandrasekhar Bhamidipati. Critical heat engines in massive gravity. *Class. Quant. Grav.*, 37(20):205020, 2020.
- [32] Sharmanthie Fernando. Massive gravity with Lorentz symmetry breaking: black holes as heat engines. *Mod. Phys. Lett. A*, 33(31):1850177, 2018.
- [33] S. H. Hendi, B. Eslam Panah, S. Panahiyan, H. Liu, and X. H. Meng. Black holes in massive gravity as heat engines. *Phys. Lett. B*, 781:40–47, 2018.
- [34] Avik Chakraborty and Clifford V. Johnson. Benchmarking Black Hole Heat Engines, II. *Int. J. Mod. Phys. D*, 27(16):1950006, 2018.
- [35] Bhamidipati Chandrasekhar and Pavan Kumar Yerra. Heat engines for dilatonic Born–Infeld black holes. *Eur. Phys. J. C*, 77(8):534, 2017.
- [36] Bhamidipati Chandrasekhar and Pavan Kumar Yerra. A note on Gauss–Bonnet black holes at criticality. *Phys. Lett. B*, 772:800–807, 2017.
- [37] Avik Chakraborty and Clifford V. Johnson. Benchmarking black hole heat engines, I. *Int. J. Mod. Phys. D*, 27(16):1950012, 2018.
- [38] Özgür Ökcü and Ekrem Aydiner. Joule–Thomson expansion of the charged AdS black holes. *Eur. Phys. J. C*, 77(1):24, 2017.
- [39] Yihe Cao, Hanwen Feng, Wei Hong, and Jun Tao. Joule–Thomson expansion of RN-AdS black hole immersed in perfect fluid dark matter. *Commun. Theor. Phys.*, 73(9):095403, 2021.
- [40] Shihao Bi, Minghao Du, Jun Tao, and Feiyu Yao. Joule-Thomson expansion of Born-Infeld AdS black holes. *Chin. Phys. C*, 45(2):025109, 2021.
- [41] Sen Guo, Yan Han, and Guo-Ping Li. Joule–Thomson expansion of a specific black hole in  $f(R)$  gravity coupled with Yang–Mills field. *Class. Quant. Grav.*, 37(8):085016, 2020.
- [42] Kartheek Hegde, A. Naveena Kumara, C. L. Ahmed Rizwan, Ajith K. M., and Md Sabir Ali. Thermodynamics, Phase Transition and Joule Thomson Expansion of novel 4-D Gauss Bonnet AdS Black Hole. *arXiv.2003.08778*.
- [43] K. V. Rajani, C. L. Ahmed Rizwan, A. Naveena Kumara, Md. Sabir Ali, and Deepak Vaid. Joule–Thomson expansion of regular Bardeen AdS black hole surrounded by static anisotropic matter field. *Phys. Dark Univ.*, 32:100825, 2021.

- [44] M. Rostami, J. Sadeghi, S. Miraboutalebi, A. A. Masoudi, and B. Pourhassan. Charged accelerating AdS black hole of  $f(R)$  gravity and the Joule–Thomson expansion. *Int. J. Geom. Meth. Mod. Phys.*, 17(09):2050136, 2020.
- [45] Cao Hoang Nam. Heat engine efficiency and Joule–Thomson expansion of nonlinear charged AdS black hole in massive gravity. *Gen. Rel. Grav.*, 53(3):30, 2021.
- [46] Davood Mahdavian Yekta, Arezoo Hadikhani, and Özgür Ökcü. Joule-Thomson expansion of charged AdS black holes in Rainbow gravity. *Phys. Lett. B*, 795:521–527, 2019.
- [47] Shan-Quan Lan. Joule-Thomson expansion of charged Gauss-Bonnet black holes in AdS space. *Phys. Rev. D*, 98(8):084014, 2018.
- [48] H. Ghaffarnejad, E. Yaraie, and M. Farsam. Quintessence Reissner Nordström Anti de Sitter Black Holes and Joule Thomson effect. *Int. J. Theor. Phys.*, 57(6):1671–1682, 2018.
- [49] Jin Pu, Sen Guo, Qing-Quan Jiang, and Xiao-Tao Zu. Joule-Thomson expansion of the regular(Bardeen)-AdS black hole. *Chin. Phys. C*, 44(3):035102, 2020.
- [50] Jie-Xiong Mo and Gu-Qiang Li. Effects of Lovelock gravity on the Joule–Thomson expansion. *Class. Quant. Grav.*, 37(4):045009, 2020.
- [51] Adolfo Cisterna, Shi-Qian Hu, and Xiao-Mei Kuang. Joule-Thomson expansion in AdS black holes with momentum relaxation. *Phys. Lett. B*, 797:134883, 2019.
- [52] Cong Li, Pengzhang He, Ping Li, and Jian-Bo Deng. Joule-Thomson expansion of the Bardeen-AdS black holes. *Gen. Rel. Grav.*, 52(5):50, 2020.
- [53] Ahmed Rizwan C. L., Naveena Kumara A., Deepak Vaid, and K. M. Ajith. Joule-Thomson expansion in AdS black hole with a global monopole. *Int. J. Mod. Phys. A*, 33(35):1850210, 2019.
- [54] M. Chabab, H. El Moumni, S. Iraoui, K. Masmarr, and S. Zhizeh. Joule-Thomson Expansion of RN-AdS Black Holes in  $f(R)$  gravity. *LHEP*, 02:05, 2018.
- [55] Jie-Xiong Mo, Gu-Qiang Li, Shan-Quan Lan, and Xiao-Bao Xu. Joule-Thomson expansion of  $d$ -dimensional charged AdS black holes. *Phys. Rev. D*, 98(12):124032, 2018.
- [56] Özgür Ökcü and Ekrem Aydiner. Joule–Thomson expansion of Kerr–AdS black holes. *Eur. Phys. J. C*, 78(2):123, 2018.
- [57] Amritendu Haldar and Ritabrata Biswas. Joule-Thomson expansion of five-dimensional Einstein-Maxwell-Gauss-Bonnet-AdS black holes. *EPL*, 123(4):40005, 2018.
- [58] Zhong-Wen Feng, Xia Zhou, Guansheng He, Shi-Qi Zhou, and Shu-Zheng Yang.



- Joule–Thomson expansion of higher dimensional nonlinearly AdS black hole with power Maxwell invariant source. *Commun. Theor. Phys.*, 73(6):065401, 2021.
- [59] Cao H. Nam. Effect of massive gravity on Joule–Thomson expansion of the charged AdS black hole. *Eur. Phys. J. Plus*, 135(2):259, 2020.
- [60] Yuan Meng, Jin Pu, and Qing-Quan Jiang. P-V criticality and Joule-Thomson expansion of charged AdS black holes in the Rastall gravity. *Chin. Phys. C*, 44(6):065105, 2020.
- [61] Ming Zhang, Chao-Ming Zhang, De-Cheng Zou, and Rui-Hong Yue. P-V criticality and Joule-Thomson expansion of Hayward-AdS black holes in 4D Einstein-Gauss-Bonnet gravity. *Nucl. Phys. B*, 973:115608, 2021.
- [62] Jing Liang, Wei Lin, and Benrong Mu. Joule–Thomson expansion of the torus-like black hole. *Eur. Phys. J. Plus*, 136(11):1169, 2021.
- [63] Jing Liang, Benrong Mu, and Peng Wang. Joule-Thomson expansion of lower-dimensional black holes. *Phys. Rev. D*, 104(12):124003, 2021.
- [64] J. P. Morais Graça, Eduardo Folco Capossoli, and Henrique Boschi-Filho. Joule-Thomson expansion for quantum corrected AdS-Reissner-Nordstrom black holes in Kiselev spacetime. *arXiv.2105.04689*.
- [65] Behrouz Mirza, Fatemeh Naeimipour, and Masoumeh Tavakoli. Joule-Thomson Expansion of the Quasitopological Black Holes. *Front. in Phys.*, 9:33, 2021.
- [66] Rui Yin, Jing Liang, and Benrong Mu. Joule–Thomson expansion of Reissner–Nordström–Anti-de Sitter black holes with cloud of strings and quintessence. *Phys. Dark Univ.*, 34:100884, 2021.
- [67] Chao-Ming Zhang, Ming Zhang, and De-Cheng Zou. Joule-Thomson Expansion of Born-Infeld AdS Black Holes in 4D Einstein-Gauss-Bonnet gravity. *arXiv.2106.00183*.
- [68] Anindya Biswas. Joule-Thomson expansion of AdS black holes in Einstein Power-Yang-mills gravity. *Phys. Scripta*, 96(12):125310, 2021.
- [69] J. P. Morais Graça, Eduardo Folco Capossoli, and Henrique Boschi-Filho. Joule-Thomson expansion for noncommutative uncharged black holes. *EPL*, 135(4):41002, 2021.
- [70] Haximjan Abdusattar, Shi-Bei Kong, Wen-Long You, Hongsheng Zhang, and Ya-Peng Hu. Joule-Thomson Expansion and Heat Engine of the FRW Universe. *arXiv.2108.09407*.
- [71] James York. Boundary terms in the action principles of general relativity. *Found. Phys.*, 16:249–257, 1986.

- [72] Andrew P. Lundgren. Charged black hole in a canonical ensemble. *Phys. Rev. D*, 77:044014, 2008.
- [73] M. Chabab, H. El Moumni, S. Iraoui, and K. Masmar. Behavior of quasinormal modes and high dimension RN–AdS black hole phase transition. *Eur. Phys. J. C*, 76(12):676, 2016.
- [74] Clifford V. Johnson. An Exact Efficiency Formula for Holographic Heat Engines. *Entropy*, 18:120, 2016.
- [75] Felipe Rosso. Holographic heat engines and static black holes: a general efficiency formula. *Int. J. Mod. Phys. D*, 28(02):02, 2018.
- [76] Steven Carlip and S. Vaidya. Phase transitions and critical behavior for charged black holes. *Class. Quant. Grav.*, 20:3827–3838, 2003.
- [77] James W. York, Jr. Black Hole in Thermal Equilibrium With a Scalar Field: The Back Reaction. *Phys. Rev. D*, 31:775, 1985.
- [78] Claudia S. Peca and Jose P. S. Lemos. Thermodynamics of Reissner-Nordstrom anti-de Sitter black holes in the grand canonical ensemble. *Phys. Rev. D*, 59:124007, 1999.