# A Riemann-Roch theorem on a weighted infinite graph

Atsushi Atsuji<sup>†</sup> Hiroshi Kaneko<sup>‡</sup>

<sup>†</sup>Department of Mathematics, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama, Kanagawa 223-8522, Japan

<sup>‡</sup>Department of Mathematics, Tokyo University of Science, 1-3 Kagurazaka, Shinjuku, Tokyo, 162-8601, Japan

## Abstract

A Riemann-Roch theorem on graph was initiated by M. Baker and S. Norine. In their article [2], a Riemann-Roch theorem on a finite graph with uniform vertex-weight and uniform edge-weight was established and it was suggested a Riemann-Roch theorem on an infinite graph was feasible. In this article, we take an edge-weighted infinite graph and focus on the importance of the spectral gaps of the Laplace operators defined on its finite subgraphs naturally given by Q-valued positive weights on the edges. We build a potential theoretic scheme for proof of a Riemann-Roch theorem on the edge-weighted infinite graph.

## 1 Introduction

A Riemann-Roch theorem on a connected finite graph was initiated by M. Baker and S. Norine in [2], where connected graph with finite vertices was investigated and unit weight was given on each edge of the graph. Since a counterpart of the lowest exponents of the complex variable in the Laurent series was directly highlighted in the Riemann-Roch theorem on graph, research on its relationships with tropical geometry was undertaken before other non-tropical complex analytical developments were implemented on the graphs. In fact, A. Gathmann and M. Kerber showed Riemann-Roch theorem on metric graphs (i.e. graphs where edge lengths are not required to be rational numbers) and then proceeded to the graphs on tropical curves (i.e. graphs with possibly unbounded edges) in [8].

In [3], M. Baker and F. Shokrieh revealed a close relationships between chip-firing games and potential theory on graphs, by characterizing reduced divisors on graphs as the solution to an energy minimization problem. In their article, an algorithm to find random spanning trees was proposed where its running times, as well as other algorithms, were analyzed by using potential theoretic ideas. A new proof of Kirchhoff's matrix-tree theorem was also covered in their article. R. James and R. Miranda showed an approach to Riemann-Roch theorem on an edge-weighted graph by proposing an alternative notion of dimension for divisors where unit weight is assigned for each vertex. Recently, S. Backman introduced in [1], the notions of edge pivot, cut reversal and cycle reversal and found a procedure to produce an acyclic partial orientation as an alternative proof of the Riemann-Roch theorem for a finite graph, where validity of two stronger hypotheses than (RR1) and (RR2) in [2] was given and their importance was highlighted. On the other hand, Riemann-Roch theorem on an infinite graph has been suggested by M. Baker and S. Norine in [2].

The objective of this article is to demonstrate a Riemann-Roch theorem on an edge-weighted infinite graph where weights on vertices are determined by the sum of the Q-valued positive weights given on its adjacent edges. Since the edge-weights give the Laplace operator which is also regarded as a generator of a Markov process, the proof is performed by a potential theoretic method including the Dirichlet space theory. We show a sufficient condition for the spectral gap in terms of the transition probability of the Markov process and prove a Riemann-Roch theorem under certain probabilisitic assumptions.

In the second section, we reinvestigate the method in [2] for Riemann-Roch theorem to accommodate procedures in [2] with  $\mathbb{Q}$ -valued edge-weighted finite graph by rethinking the canonical divisor in the existing Riemann-Roch theorem in [2]. For the sake of self-contained article presentation, we attentively look at several assertions updated for a weighted finite graph and pay attention to geometric aspects and characteristics of the weighted graph. As for various important relationships between networks with weighted edges and observations in other research subjects, see [10] and [5]. In the third section, since we can find a potential theoretic aspect in the proof of the Riemann-Roch theorem on weighted finite graph, we take validity of Poincaré inequality and Riemann-Roch theorem on any edge-weighted finite graph into account and use potential theoretical techniques to approach Poincaré inequality on edge-weighted infinite graph. We focus also on probabilistic significance in a decay of edge-weights responding to increase in its distance from a fixed reference point in graph metric basis and establish a Poincaré inequality on the edge-weighted infinite graph by taking an exhaustive sequence of finite subgraphs. The aim of the section is to validate a spectral gap on the basis of the method as in [11] for widely accepted discussions on  $L^2$ -boundedness of 0-order resolvent of a probabilistically natural Laplacian associated with the edge-weights. In the final section, we finalize the proof of our Riemann-Roch theorem on the infinite graph with Q-valued positive weights on edges. In the proof, we note that the spectral gap yields the uniform norm boundedness of a family of functions on the infinite graph (each function in the family represents an accumulation of chip-firings on a subgraph) to obtain a characteristic value of a given divisor. Along with a counterpart of the Euler characteristic for the weighted infinite graph, the characteristic values of divisors are involved directly in our Riemann-Roch theorem.

## 2 Riemann-Roch theorem on a weighted finite graph

Let  $G = (V_G, E_G)$  be a connected graph consisting of a finite set  $V_G$  of vertices and a finite set  $E_G$  of edges without loops. To be more precise,  $E_G$  is given as a subset of  $(V_G \times V_G \setminus \{\{x,x\} \mid x \in V_G\}) / \sim$ , where  $\{x,y\} \sim \{y,x\} \in E_G$  for any pair  $x, y \in V_G$ . Accordingly,  $\{x, y\}$  and  $\{y, x\}$  are regarded as an identical element in  $E_G$ . We assume that a Q-valued positive weight  $C_{x,y}$  is given at every edge  $\{x, y\} \in E_G$ with the symmetricity  $C_{x,y} = C_{y,x}$  for any  $\{x, y\} \in E_G$ , according to the identification  $\{x,y\} = \{y,x\}$ . At each point x in  $V_G$ , the set N(x) consisting of its neighbor is defined by  $N(x) = \{y \in V_G \mid \{x, y\} \in E_G\}$  of x. For any  $\mathbb{Z}$ -valued function f on  $V_G$ , the Laplacian  $\Delta f$  of the function is defined by  $\Delta f(x) = \sum_{y \in N(x)} C_{x,y}(f(x) - f(y))$ as a function on  $V_G$ . When G is regarded as the electrical circuit given by installing a resistor with the resistance  $1/C_{x,y}$  at every edge  $\{x, y\} \in E_G$ , the value f(x) is viewed as the voltage level at  $x \in V_G$ . We can focus on another function i on  $V_G$ defined by  $i(x) = \min\{|\Delta f(x)| \mid f: V_G \to \mathbb{Z} \text{ satisfying } f(x) = 0 \text{ and } \Delta f(x) \neq 0\},\$ which is viewed as the minimum positive absolute value of feasible current flows at the grounded vertex x given by integer valued voltage f(y) with  $y \in N(x)$ . In what follows, similarly to M. Baker and S. Norine's article [2], f persistently stands for  $\mathbb{Z}$ -valued function on  $V_G$ .

We use notation  $\ell$  for another Z-valued function to assign an integer multiple of i(x) at each  $x \in V_G$ . A divisor on the graph G is given by  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$ and its degree deg(D) is defined by deg $(D) = \sum_{x \in V_G} \ell(x)i(x)$ . A divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  is said to be effective if  $\ell(x) \ge 0$  for all  $x \in V_G$ . Since f is Z- valued,  $\Delta f(x)$  is given as an integer multiple of i(x) at each  $x \in V_G$  and the Laplacian  $\Delta f$  will be identified with the divisor  $\sum_{x \in V_G} \Delta f(x) \mathbb{1}_{\{x\}}$ . This identification makes it possible to add the Laplacian  $\Delta f$  to any divisor.

Along with this addition, we need two materials characterizing the graph G for a Riemann-Roch theorem on the weighted graph: one is the positive real value  $\min\{|\sum_{x \in V_G} \ell(x)i(x)| \in (0, \infty) | \ell : V_G \to \mathbb{Z}\},$  which is denoted by  $i_{(G,C)}$ ; and the other is the canonical divisor  $K_G$  on the weighted graph G, which is given by  $K_G =$  $\sum_{x \in V_G} \{\sum_{y \in N(x)} C_{x,y} - 2i(x)\} \mathbf{1}_{\{x\}}.$  Similar to [2], for our goal, we take advantage of total orders on  $V_G$  and denote the family of total orders on  $V_G$  by  $\mathcal{O}$ . For each  $\mathcal{O} \in \mathcal{O}$ , we introduce the divisor  $\nu_O$  given by

$$\sum_{x \in V_G} \nu_O(x) \mathbb{1}_{\{x\}},$$

with  $\nu_O(x) = \sum_{y \in N(x), y <_O x} C_{x,y} - i(x)$  for  $x \in V_G$ , and its reversed total order  $\overline{O}$  is defined by  $x <_{\overline{O}} y$  for any  $x, y \in V_G$  satisfying  $y <_O x$ .

The importance of such divisors is found in its close relationship with the canonical divisor  $K_G$  as shown in the first assertion of the following lemma, similar to [2].

**Lemma 2.1.** For any divisor  $\nu_{\overline{O}}$  determined by  $\overline{O} \in \mathcal{O}$  and any divisor  $\Delta f$  determined by  $\mathbb{Z}$ -valued function f, one sees the followings:

- (i)  $\nu_{\overline{O}} = K_G \nu_O$ ,
- (ii)  $\deg(\Delta f) = 0.$

*Proof.* (i) The assertion is clear from the identity  $\nu_{\overline{O}} + \nu_O = K_G$ . (ii) The identity follows from the equivalence between  $y \in N(x)$  and  $x \in N(y)$  and the trivial identity  $C_{x,y}(f(x) - f(y)) = -C_{y,x}(f(y) - f(x))$ .

If a pair of divisors D and D' satisfies  $D' = D + \Delta f$  for some  $\mathbb{Z}$ -valued function f, D and D' are called equivalent and the relationship is denoted by  $D \sim D'$ . As in [2], we introduce the linear system  $|D| = \{D' \mid D' \text{ is effective and equivalent with } D\}$  for any divisor D and the following condition on the graph  $G = (V_G, E_G)$ :

(**RR**) For each divisor D, there exists an  $O \in \mathcal{O}$  such that exactly one of two linear systems |D| and  $|\nu_O - D|$  is empty.

The family of total orders satisfying this condition with respect to D will be denoted by  $\mathcal{O}_D$ .

We introduce  $\mathfrak{e}_{(G,C)} = \sum_{x \in V_G} i(x) - \sum_{\{x,y\} \in E_G} C_{x,y}$  as a counterpart of 1 - g with the genus g of G in M. Baker and S. Norine's article [2]. In fact, we see that

$$\deg(K_G) = \sum_{x \in V_G} \{ \sum_{y \in N(x)} C_{x,y} - 2i(x) \} = -2\mathfrak{e}_{(G,C)}.$$

**Lemma 2.2.** For each  $O \in \mathcal{O}$ ,  $\nu_O \in \mathcal{N}$ , where  $\mathcal{N}$  stands for the family of divisors of degree  $-\mathfrak{e}_{(G,C)}$  admitting only non-effective equivalent divisors.

*Proof.* It is easy to see that  $\deg(\nu_O) = -\mathfrak{e}_{(G,C)}$  and  $\nu_O$  is not effective as in [2]. For any divisor D given by  $D = \nu_O - \Delta f$  with some non-constant  $\mathbb{Z}$ -valued function f, we take  $A_f = \{x \in V_G \mid f(x) = \max_{y \in V_G} f(y)\}$  and the minimal element z in  $A_f$  with respect to the total order O. Since  $y <_O z$  implies  $f(y) \leq f(z) - 1$  and  $f(y) \leq f(z)$  for all  $y \in V_G$ , it turns out that

$$D(z) = \left(\sum_{y \in N(z), y <_O z} C_{z,y} - i(z)\right) - \left(\sum_{y \in N(z)} C_{z,y}(f(z) - f(y))\right)$$
  
=  $-i(z) + \sum_{y \in N(z), y >_O z} C_{z,y}(f(y) - f(z)) + \sum_{y \in N(z), y <_O z} C_{z,y}(f(y) - f(z) + 1)$   
 $\leq -i(z).$ 

This implies that  $\nu_O$  can not be equivalent to any effective divisor.

For any divisor D and non-negative integer k, we introduce  $E_k(D) = \{$  effective divisors  $E \mid \text{deg } (E) = k i_{(G,C)}$  satisfying  $|D - E| \neq \emptyset \}$ . We note that  $E_0(D)$  is either the empty set or the set consisting only of the zero divisor vanishing on  $V_G$ . We also define a  $\{-i_{(G,C)}, 0, i_{(G,C)}, 2i_{(G,C)}, \dots\}$ -valued function r on the set consisting of all divisors by

$$r(D) = \begin{cases} -i_{(G,C)}, & \text{if } E_0(D) = \emptyset, \\ \max\{k \, i_{(G,C)} & \mid E_k(D) \text{ consists of all effective divisors of degree } k \, i_{(G,C)}\}, \\ & \text{otherwise,} \end{cases}$$

similar to [2].

Our objective is to establish the following identity as a natural counterpart of the Riemann-Roch theorem on a weighted graph:

$$r(D) - r(K_G - D) = \deg(D) + \mathfrak{e}_{(G,C)}.$$

Let us briefly look back at the procedures for the proof of Riemann-Roch theorem presented in the article based on the notions in our settings involving edge-weight.

For a divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$ , the subset  $\{x \in V_G \mid \ell(x) \neq 0\}$  is called a support of D and denoted by  $\operatorname{supp}[D]$ . It is straightforward that any divisor D is uniquely decomposed as D = E - E' with effective divisors E, E' without intersection of their supports. In what follows, the degree of E in the decomposition will be denoted by  $\operatorname{deg}^+(D)$  and the degree of E' in the decomposition will be denoted by  $\operatorname{deg}^-(D)$ .

**Proposition 2.3.** Condition (**RR**) implies that

$$r(D) = \left(\min_{D' \sim D, O \in \mathcal{O}} \deg^+(D' - \nu_O)\right) - i_{(G,C)}$$

for any divisor D.

*Proof.* If  $|D-E| = \emptyset$  for some effective divisor E, (**RR**) implies that there exists a total order  $O \in \mathcal{O}_{D-E}$  such that  $\nu_O - (D-E) \sim E'$  equivalent to  $\nu_O - D \sim E' - E$ with some effective divisor E'. This shows that  $D' - \nu_O \sim E - E'$  with the effective divisors E, E' independent of the choice of  $D' \sim D$ .

Conversely, if  $D' - \nu_O \sim E - E'$  for some effective divisors  $E, E', O \in \mathcal{O}$  and D' satisfying  $D' \sim D$ , then  $D - E \sim \nu_O - E'$ . However, Lemma 2.2 implies that  $\nu_O - E'$  is not equivalent to any effective divisor. This shows that  $|D - E| = \emptyset$ .

Originally, by taking  $I_{(G,C)} = \{k i_{(G,C)} \mid k \in \{0, 1, 2, ...\}\}$ , the value r(D) is characterized by r(D) < s if and only if there exists some effective divisor E with  $\deg(E) = s$  such that  $|D - E| = \emptyset$ , where  $s \in I_{(G,C)}$ . Another characterization can now be admitted r(D) < s if and only if  $s \ge \deg^+(D' - \nu_O)$  for some  $D' \sim D$  and  $O \in \mathcal{O}$ , by using non-negative integer multiple s of  $i_{(G,C)}$ . The assertion has been proved.

This proposition shows that, for a divisor D on  $V_G$ , its equivalent divisor D'and a total order O on  $V_G$  are taken so that the minimum in the proposition is attained. We introduce the involution  $\overline{D}'$  of D' by  $\overline{D}' = K_G - D'$  and then see that  $\overline{D}' - \nu_{\overline{O}} = K_G - D' - (K_G - \nu_O) = \nu_O - D'$ , which gives an alternative expression  $\overline{D}' - \nu_{\overline{O}} \sim E' - E$ , the decomposition into the sum of effective divisors as in the proof of the last proposition. Cororally 2.4. Condition (RR) implies

(i) for any divisor D,

$$r(\overline{D}) = \left(\min_{D' \sim D, O \in \mathcal{O}} \deg^+(\overline{D}' - \nu_{\overline{O}})\right) - i_{G,C},$$

where  $\overline{D} = K_G - D$ , the involution of D,

(ii) min{deg(E) | D' - ν<sub>O</sub> ~ E - E' with some effective divisors E, E'} and min{ deg(E') | D' - ν<sub>O</sub> ~ E' - E with some effective divisors E, E'} are both attained by the effective divisors E = E<sub>0</sub>, E' = E'<sub>0</sub>, which are characterized by a unique decomposition D' - ν<sub>O</sub> ~ E<sub>0</sub> - E'<sub>0</sub> into the sum of effective divisors of minimal degrees, equivalently given as D' - ν<sub>O</sub> ~ E'<sub>0</sub> - E<sub>0</sub>,

(iii) for any divisor D with  $\deg(D) = -\mathfrak{e}_{(G,C)}, D \in \mathcal{N}$  if and only if  $K_G - D \in \mathcal{N}$ .

*Proof.* (i) The identity follows from the equivalence between  $D' \sim D$  and  $\overline{D}' \sim \overline{D}$ and from the equivalence between the choice of O and the choice of  $\overline{O}$ . (ii) Since  $\deg(E) = \mathfrak{e}_{(G,C)} + \deg(D) + \deg(E')$ , the assertion is straightforward and justified. (iii) The assertion follows from  $\deg(K_G - D) = -\mathfrak{e}_{(G,C)}$  and the equivalence between  $D \notin \mathcal{N}$  and  $K_G - D = \overline{D} \notin \mathcal{N}$ , i.e., the equivalence between  $r(D') = r(D) \ge 0$  and  $r(\overline{D}') = r(\overline{D}) \ge 0$ .

**Remark 2.5.** Condition (RR2) in [2] is affirmatively asserted in (iii) in the corollary.

We introduce the graph metric d which is defined by

$$d(v,w) = \begin{cases} \min\{j \mid \{w_0, w_1\}, \dots, \{w_{j-1}, w_j\} \in E_G, v = w_0 \text{ and } w = w_j\} & \text{if } v \neq w, \\ 0 & \text{if } v = w. \end{cases}$$

We take a fixed base (reference) vertex  $v_0 \in V_G$ , then the distance between v and  $v_0$  is given by  $d(v_0, v)$ , which will be denoted by  $d_G(v)$ .

We introduce  $d_G = \max\{d(v) \mid v \in V_G\}$ , and for any given divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  on  $V_G$ , we introduce a  $d_G - 1$ -dimensional vector

$$\mathbf{V}_{1}(D) = \Big(\sum_{z \in S_{d_{G}}, \ell(z) < 0} \ell(z)i(z), \sum_{z \in S_{d_{G}-1}, \ell(z) < 0} \ell(z)i(z), \dots, \sum_{z \in S_{1}, \ell(z) < 0} \ell(z)i(z)\Big)$$

and a  $d_G$ -dimensional vector

$$\mathbf{V}_2(D) = \Big(\sum_{z \in S_0} \ell(z)i(z), \sum_{z \in S_1} \ell(z)i(z), \dots, \sum_{z \in S_{d_G}} \ell(z)i(z)\Big),$$

where  $S_j = \{v \in V_G \mid d_G(v) = j\}$  for any non-negative integer j.

If a divisor  $D = \sum_{x \in V_G} \ell(x) i(x) \mathbb{1}_{\{x\}}$  satisfies the following two conditions

- (P1)  $\ell(z)i(z) \ge 0$  for all  $z \in V_G \setminus \{v_0\},\$
- (P2) for every non-empty set  $A \subset V_G \setminus \{v_0\}$ , there exists a vertex  $x \in A$  such that  $\ell(x)i(x) < \operatorname{outdeg}_A(x) = \sum_{y \in N(x) \cap A^c} C_{x,y}$ ,

with respect to the base vertex  $v_0$ , then D is said to be  $v_0$ -reduced.

**Proposition 2.6.** For any divisor D, there exists a unique  $v_0$ -reduced divisor  $D_0$  such that  $D_0 \sim D$ .

*Proof.* By starting with the originally given divisor D, we first take the subfamily  $\mathcal{D}'$  of its equivalent divisors given by

 $\mathcal{D}' = \{ D' \mid D' \text{ attains the maximum } \max_{D' \sim D} \mathbf{V}_1(D') \text{ in the sense of the lexicographical order} \},$ 

and in the next, we take the subfamily  $\mathcal{D}''$  of  $\mathcal{D}'$  given by

$$\mathcal{D}'' = \{ D'' \in \mathcal{D}' \mid D'' \text{ attains the maximum } \max_{D'' \sim D} \mathbf{V}_2(D'') \text{ in the sense of the lexicographical order} \}.$$

We see that any divisor in  $\mathcal{D}''$  attains both maxima and satisfies the conditions (P1) and (P2) with respect to the base (reference) vertex  $v_0$  as in the proof of Proposition 3.1 in [2].

For the uniqueness of the  $v_0$ -reduced divisor, we assume that there exist two distinct  $v_0$ -reduced divisors  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}, D' = \sum_{x \in V_G} \ell'(x)i(x)1_{\{x\}}$  such that  $D' = D - \Delta f$  for some non-constant  $\mathbb{Z}$ -valued function f. Without losing these general settings, we may assume that  $\max_{y \in V_G} f(y) > f(v_0)$  as in the proof of uniqueness in the proposition in [2], from which we can derive that  $v_0 \notin A_f = \{x \in V_G \mid f(x) = \max_{y \in V_G} f(y)\}$  and

$$0 \le \ell'(x)i(x) = \ell(x)i(x) - \sum_{\{x,y\} \in E_G} C_{x,y}(f(x) - f(y)) \le \ell(x)i(x) - \text{outdeg}_{A_f}(x)$$

for any  $x \in A_f$ . This contradicts the condition (P2) originally imposed on D.

The following theorem shows that (RR1) in [2] still holds valid even for our weighted graph.

### **Theorem 2.7.** For any divisor D, $\mathcal{O}_D \neq \emptyset$ , namely Condition (**RR**) is satisfied.

*Proof.* A similar procedure to the construction of a total order in the proof of Theorem 3.3 in [2] works to deduce  $\mathcal{O}_D \neq \emptyset$ .

**Remark 2.8.** For any divisor D with  $\deg(D) = -\mathfrak{e}_{(G,C)}$ ,  $D \in \mathcal{N}$  if and only if  $D \sim \nu_O$  for some  $O \in \mathcal{O}$ . In fact, thanks to Lemma 2.2, it suffices to show that  $D \in \mathcal{N}$  implies the existence of  $O \in \mathcal{O}$  satisfying  $D \sim \nu_O$ . This implication is verified by combining  $\mathcal{O}_D \neq \emptyset$  with the fact that any  $D \in \mathcal{N}$  does not admit any equivalent effective divisors.

**Remark 2.9.** Our weight function i(x) defined on the set  $V_G$  of vertices may differ from the uniform vertex-weight assigned in [2]. On the other hand, we see that  $C_{x,y}$ admits representation as  $C_{x,y} = \frac{n_{x,y}}{m_G}$  with positive integers  $n_{x,y}$  and  $m_G$ , where mcan be taken independently of edges  $\{x, y\} \in E_G$ . Accordingly, we can install  $n_{x,y}$ -ple edges with a weight between x and y in the network discussed in [2] so that essentially the same network is obtained up to the difference in minimum edge weights, namely, such a difference between  $i_{(G,C)}$  and  $1/m_G$ . This shows us that the procedure in [2] for finding the  $v_0$ -reduced divisor is still valid. This is because the image  $\{\Delta f \mid f :$  $V_G \to \mathbb{Z}\}$  of our Laplacian is linearly isomorphic to the one in [2].

**Theorem 2.10.** (Riemann-Roch theorem on a weighted finite graph). For any divisor D,

$$r(D) - r(K_G - D) = \deg(D) + \mathfrak{e}_{(G,C)}.$$

*Proof.* Theorem 2.7 shows that  $\mathcal{O}_D \neq \emptyset$  for any divisor D. Therefore, one can take a divisor D' satisfying  $D' \sim D$  and a total order  $O \in \mathcal{O}_{D-E_0}$  for any effective divisor  $E_0$  satisfying  $|D - E_0| = \emptyset$  and the minimal degree condition  $\deg(E_0) = \min\{\deg(E) \mid |D-E| = \emptyset, E \text{ is effective}\}$ . By the discussion in the proof of Corollary 2.4, we have the decomposition  $D' - \nu_O = E_0 - E'_0$  with effective divisors  $E_0, E'_0$  of the minimal degree. Accordingly, we see that

$$deg^{+}(D' - \nu_{O}) - deg^{+}(\overline{D}' - \nu_{\overline{O}}) = deg^{+}(D' - \nu_{O}) - deg^{+}((K_{G} - D') - (K_{G} - \nu_{O}))$$
  
=  $deg^{+}(D' - \nu_{O}) - deg^{+}(\nu_{O} - D')$   
=  $deg(E_{0}) - deg(E'_{0})$   
=  $deg(D' - \nu_{O})$   
=  $deg(D) + \mathfrak{e}_{(G,C)}.$ 

Since the left-hand side is equal to  $r(D) - r(K_G - D)$  due to Proposition 2.3, the identity in the assertion is derived.

We close this section with the following fundamental properties of r(D) which are utilized later.

**Lemma 2.11.** (i) If D' is effective, then  $r(D) + \deg(D') \ge r(D + D')$ , (ii) If -D'' is effective, then  $r(D) + \deg(D'') \le r(D + D'')$ .

*Proof.* (i) Since there exists an effective divisor E with  $\deg(E) > r(D)$  satisfying  $|D - E| = \emptyset$ , it turns out that the effective divisor D' + E satisfies  $\deg(E + D') > r(D) + \deg(D')$  and  $|(D + D') - (E + D')| = \emptyset$ . Accordingly, r(D + D') does not exceed  $r(D) + \deg(D')$ .

(ii) It suffices to show that  $r(D) \leq r(D + D'') - \deg(D'')$ . This follows from (i) by regarding D + D'' and -D'' as D and D' respectively in the identity of (i).

# 3 Poincaré inequality for spectral gap on an infinite graph

We shift our attention from finite graphs to an infinite graph. Throughout this section, we consider a connected infinite graph  $G = (V_G, E_G)$  consisting of countably

infinite sets  $V_G$  and  $E_G$  of vertices and edges, respectively. We assume that the graph  $G = (V_G, E_G)$  does not admit any loops and is locally finite, namely  $N(x) = \{y \in V_G \mid \{x, y\} \in E_G\}$  is a finite set for every  $x \in V_G$ . Let  $C_{x,y}$  denote Q-valued positive weight given at every edge  $\{x, y\} \in E_G$  satisfying that  $C_{x,y} = C_{y,x}$  for every  $x, y \in V_G$ , according to the identification  $\{x, y\} = \{y, x\}$ . Define a measure m on  $V_G$  by

$$m(A) = \sum_{x \in A} \sum_{y \in N(x)} C_{x,y},$$

for  $A \subset V_G$ . We also assume finiteness of total volume on the graph G, namely,  $m(V_G) < \infty$ .

Here, we describe the basic idea of obtaining our Riemann-Roch theorem on an infinite graph from the facts obtained in the case of finite graphs. We take the graph metric d that is exactly the same in the previous section and fix a base (reference) vertex  $v_0$ , define  $V_n = \{x \in V_G \mid d(v_0, x) \leq n\}$  and  $E_n = \{\{x, y\} \mid x, y \in V_n\}$  to determine the subgraph  $G_n = (V_n, E_n)$  and introduce the Laplacian  $\Delta_n f(x) = \sum_{y \in N(x) \cap V_n} C_{x,y}(f(x) - f(y))$  on  $V_n$ .

As will be seen later, a divisor D can be given globally on G and it admits a sequence  $\{(D)_n\}$  of divisors, *n*-th divisor of which is a restriction of D to  $G_n$ . We already have the Riemann-Roch theorem on  $G_n$  in Section 2. Thus, by the equivalence  $D' \stackrel{n}{\sim} D''$  between divisors D' and D'' on  $V_n$  defined by  $D'' = D' + \Delta_n f$ for some  $\mathbb{Z}$ -valued function f on  $V_n$  and with the same notations in the last section, we have  $r_n((D)_n)$  on  $G_n$  satisfying

$$r_n((D)_n) = \left(\min_{D' \sim (D)_n, O_n \in \mathcal{O}_n} \deg^+ (D' - \nu_{O_n})\right) - i_{(G_n, C_n)},$$

where  $\mathcal{O}_n$  stands for the set consisting of all total orders on  $V_n$ . To be more precise, bearing the difference between  $\Delta_n f(x)$  and  $\Delta f(x) = \sum_{y \in N(x)} C_{x,y}(f(x) - f(y))$  for x with  $d(v_0, x) = n$  in our mind, we notice that  $\sum_{x \in V_n} \ell(x)i(x)\mathbf{1}_{\{x\}}$  can not always be regarded as any divisor on  $G_n$ , and in contrast  $\sum_{x \in V_{n-1}} \ell(x)i(x)\mathbf{1}_{\{x\}}$  is regarded as a divisor vanishing outside  $V_{n-1}$  as detailed in Remark 3.1 below. We take the latter vanishing outside  $V_{n-1}$  as the restriction  $(D)_n$  to  $G_n$ , when a divisor  $D = \sum_{x \in V_G} \ell(x)i(x)\mathbf{1}_{\{x\}}$  is globally given.

**Remark 3.1.** For any positive integer n and  $x \in V_n$ , not only  $i(x) = \min\{|\Delta f(x)| \mid f : V_G \to \mathbb{Z} \text{ with } \Delta f(x) \neq 0\}$  but  $i_n(x) = \min\{|\Delta_n f(x)| \mid f : V_n \to \mathbb{Z} \text{ with } \Delta_n f(x) \neq 0\}$  is defined. We note that i(x) does not always coincide with  $i_n(x)$  for x with  $d(v_0, x) = n$ , and that  $D' = \sum_{x \in V_n} \ell(x)i(x)1_{\{x\}}$  does not always admit the

expression  $\sum_{x \in V_n} \ell(x) i_n(x) \mathbb{1}_{\{x\}}$ . However, for any pair of positive integers j, n with  $j \geq n$ , any divisor  $D = \sum_{x \in V_{n-1}} \ell(x) i(x) \mathbb{1}_{\{x\}}$  vanishing outside  $V_{n-1}$  is regarded as a divisor on  $G_j$ .

An integer-valued function  $f_n$  on  $V_n$ , such that  $D' = (D)_n + \Delta_n f_n$  attains the minimum  $\min_{D' \sim (D)_n, O_n \in \mathcal{O}_n} \deg^+ (D' - \nu_{O_n})$ , is called a minimizer for  $r_n((D)_n)$ . Then, one of the problems to be considered here is the global existence of such a function, namely, the solution  $f_n$  to the equation  $\Delta_n f = D' - (D)_n$  and extension of the function to larger subgraph. In the case of finite graph, we always have the solution to  $\Delta_n f = g$  on  $G_n$ , since the solution can be given by the 0-order resolvent of  $\Delta_n$ . If we can expect a stability of the correspondence from  $D' - (D)_n$  to  $f_n$  as  $n \to \infty$ , using 0-order resolvent as a main tool still in infinite graph, we carry out the following procedure: First, we take a minimizer  $f_n$  by using resolvent of  $\Delta_n$  on  $G_n$ . Second, by deriving the convergence of  $\{f_n\}$  as  $n \to \infty$  from the stability, we discuss the existence of  $\lim_{n\to\infty} r_n((D)_n)$ , and finally we define r(D) as the limit and complete a proof of a Riemann-Roch theorem on infinite graphs.

However, these procedures are not actually straightforward and we need to carefully discuss the convergence by selecting subsequences  $\{(D)_n\}$  and  $\{f_n\}$ , linked with a choice of subsequence of  $\{O_n\}$  with  $O_n \in \mathcal{O}_n$ . Precise discussion on this problem will be given in Section 4.

The more basic problem here is that the 0-order resolvent does not always exist on infinite graphs and operator theoretical treatment of the resolvent  $\Delta_n^{-1}$  for our purpose is unclear within general operator theory. The existence of a spectral gap of the Laplacian are known as a powerful condition for the existence of the resolvent (see [11] for the general theory of spectral gaps and related functional inequalities) and we can also clear the convergence problem. On the other hand, the resolvent can be defined from a natural Markov process on the graphs associated with the weight  $C_{x,y}$ , which we will give in Section 3.1. To ensure the existence of the spectral gap, we propose a condition of infinite graphs using a quantity  $\rho_n$  on  $G_n$  in terms of the Markov process as defined in (3) below. In fact, we will show in Section 3.2 that the spectral gap exists if  $\limsup_{n\to\infty}\rho_n$  is sufficiently small. Roughly speaking, this condition implies a strong recurrence property of the process and then a graph satisfying the condition looks closely like a finite graph from the view of the Markov process. To show the existence of a spectral gap of our Laplacian under the condition, we introduce a Dirichlet form  $\mathcal{E}$  associated with  $\Delta$  in Section 3.1 and show a Poincaré type inequality of  $\mathcal{E}$  over some  $L^2$  space in Section 3.2. We remark that there are one to one correspondence among the Markov process, the Laplacian and the Dirichlet form.

## 3.1 Basic properties of weighted Laplacians

In this section, we use the following notions and notations. Let  $U = (V_U, E_U)$  be a connected subgraph of G, i.e. U is a graph consisting of set  $V_U$  of vertices with  $V_U \subset V_G$  and set  $E_U$  of edges with  $E_U = \{\{x, y\} \in E_G \mid x, y \in V_U\}$ , and  $m_U$  be the measure on  $V_U$  determined by  $m_U(A) = \sum_{x \in V_U \cap A} m_U(\{x\})$  with  $m_U(\{x\}) = \sum_{y \in N(x) \cap V_U} C_{x,y}$ . Here and throughout this subsection, we denote  $V_G$  by V.

Recall the definition of the Laplacians on graphs G and U;

$$\Delta \phi(x) = \sum_{y \in N(x)} C_{x,y}(\phi(x) - \phi(y)),$$
$$\Delta_U u(x) = \sum_{y \in N(x) \cap V_U} C_{x,y}(u(x) - u(y)),$$

for real-valued functions  $\phi$  and u on V and  $V_U$  respectively.

Define the operators by

$$L = \frac{1}{m}\Delta$$
 and  $L_U = \frac{1}{m_U}\Delta_U$ .

Here and in the sequel, m and  $m_U$  stand for the function taking  $m(\{x\})$  at every x and the one taking  $m_U(\{x\})$  at every  $x \in U$ , respectively. In what follows,  $m(\{x\})$  and  $m_U(\{x\})$  will be denoted by m(x) and  $m_U(x)$ , respectively.

Note that L and  $L_U$  are symmetric, respectively in the sense that

$$(L\phi,\psi)_{L^2(m)} = (\phi,L\psi)_{L^2(m)}$$
 and  $(L_Uu,v)_{L^2(m_U)} = (u,L_Uv)_{L^2(m_U)}$ 

for  $\phi, \psi \in L^2(m)$  with  $L\phi, L\psi \in L^2(m)$  and for  $u, v \in L^2(m_U)$  with  $Lu, Lv \in L^2(m_U)$ .

We see that those operators are associated with Dirichlet forms:

$$\mathcal{E}(f,g) = \sum_{\{x,y\}\in E_G} C_{x,y}(f(x) - f(y))(g(x) - g(y))$$

for functions f, g on V with  $f, g \in \mathcal{F} = \{h \in L^2(m) \mid \mathcal{E}(h, h) = \sum_{\{x,y\} \in E_G} C_{x,y}(h(x) - h(y))^2 < \infty\}$ , and

$$\mathcal{E}_{U}(u,v) = \sum_{\{x,y\}\in E_{U}, x,y\in V_{U}} C_{x,y}(u(x) - u(y))(v(x) - v(y))$$

for functions u, v on U with  $u, v \in \mathcal{F}_U = \{w \in L^2(m_U) \mid \mathcal{E}_U(w, w) = \sum_{\{x,y\} \in E_U} C_{x,y}$  $(w(x) - w(y))^2 < \infty\}$ , in the sense that

$$(Lf, g)_{L^2(m)} = \mathcal{E}(f, g) \text{ and } (L_U u, v)_{L^2(m_U)} = \mathcal{E}_U(u, v).$$

For a function  $\phi$  on V

$$L\phi(x) = \frac{1}{m(x)} \sum_{y \in N(x)} C_{x,y}(\phi(x) - \phi(y))$$
  
=  $\frac{1}{m(x)} \sum_{y \in N(x) \cap V_U} C_{x,y}(\phi(x) - \phi(y)) + \frac{1}{m(x)} \sum_{y \in N(x) \cap V_{U^c}} C_{x,y}(\phi(x) - \phi(y))$   
=  $\frac{m_U(x)}{m(x)} L_U \phi(x) + \frac{1}{m(x)} \sum_{y \in N(x) \cap V_{U^c}} C_{x,y}(\phi(x) - \phi(y)).$ 

By identifying function  $\phi$  on V with its restriction to  $V_U$ , we have the following relationship between L and  $L_U$  (which is used frequently below):

$$L\phi(x) = \frac{m_U(x)}{m(x)} L_U\phi(x) + \frac{\phi(x)}{m(x)} (m(x) - m_U(x)),$$
(1)

for any function  $\phi$  satisfying  $\phi = 0$  on  $V_U^c$ . The last term of the right hand side may be regarded as a boundary operator on  $\partial U = \{x \in V \mid N(x) \cap V_U \neq \emptyset \text{ and } N(x) \cap V_U^c \neq \emptyset\}.$ 

In what follows, we need the probability measure  $\mu$  proportional to m given by the normalization  $\mu(A) = \frac{m(A)}{m(V)}$  for  $A \subset V$ . We use the following estimate with the function  $\mu(x) = \frac{m(x)}{m(V)}$  on V later:

**Lemma 3.2.** Let U be a connected subgraph of G and  $f \in L^2(\mu)$ . For any  $\epsilon > 0$ 

$$\sum_{x \in V_U} f(x) \frac{1}{m(x)} \Delta_U f(x) \mu(x) \le \frac{\epsilon}{2} ||f||_{L^2(\mu)}^2 + \frac{1}{\epsilon m(V)} \mathcal{E}(f, f).$$

*Proof.* By Fubini's theorem and the Cauchy-Schwarz inequality, we have

$$\sum_{x \in V_U} f(x) \frac{1}{m(x)} \Delta_U f(x) \mu(x)$$
  
=  $m(V) \sum_{x \in V_U} \sum_{y \in V_U} 1_{\{x,y\} \in E_U\}} \frac{1}{m(x)m(y)} C_{x,y}(f(x) - f(y))f(x)\mu(x)\mu(y)$   
 $\leq m(V) \Big(\sum_{x \in V_U} \sum_{\{x,y\} \in E_U} f(x)^2 \frac{1}{m(x)m(y)} C_{x,y}\mu(x)\mu(y) \Big)^{1/2}$   
 $\times \Big(2 \sum_{\{x,y\} \in E_U} (f(x) - f(y))^2 \frac{1}{m(x)m(y)} C_{x,y}\mu(x)\mu(y) \Big)^{1/2}.$ 

Since  $\sum_{\{x,y\}\in E_U} \frac{1}{m(x)m(y)} C_{x,y}\mu(y) \leq \frac{1}{m(V)} \frac{1}{m(x)} \sum_{\{x,y\}\in E_U} C_{x,y} \leq \frac{1}{m(V)}$ , for any  $\epsilon > 0$ , we see that the right-hand side is dominated by  $\epsilon^{1/2} \left(\frac{2}{\epsilon m(V)}\right)^{1/2} ||f||_{L^2(\mu)} \mathcal{E}(f,f)^{1/2}$ , which does not exceed  $\frac{\epsilon}{2} ||f||_{L^2(\mu)}^2 + \frac{1}{\epsilon m(V)} \mathcal{E}(f,f)$ .

Here, we give a remark on the relationship between the weighted Laplacian L and some stochastic processes: a reversible Markov chain  $\{X_n\}_{n=0,1,2,\ldots}$  and a Hunt process  $\{Y_t\}_{t\in[0,\infty)}$ . See [12] for reversible Markov chains and see [6] for Hunt processes and the theory of Dirichlet forms.

**Remark 3.3.** The Markov chain  $\{X_n\}$  can be determined by the transition matrix  $P = (p(x, y))_{x,y \in V}$ , where

$$p(x,y) = C_{x,y}/m(x).$$
 (2)

Thus for  $A \subset V$ ,  $P_x(X_1 \in A) = \sum_{x \in A} C_{x,y}/m(x)$ . The Hunt process  $\{Y_t\}$  can be determined by transition semigroup  $\{P_t\}$  associated with L as its infinitesimal generator.  $\{P_t\}$  is a symmetric semigroup on  $L^2(\mu)$ ; heuristically  $\{P_t\}$  can be expressed by  $P_t = e^{-tL}$  for any t > 0.

## 3.2 Spectral gaps and Poincaré inequality

As in the second section, the graph metric between x and y is denoted by d(x, y)and  $i(x) = \min\{|\Delta f(x)| \mid f : V_G \to \mathbb{Z} \text{ with } \Delta f(x) \neq 0\}$  is defined for every  $x \in V_G$ , due to the local finiteness of G. We fix a base point  $v_0$  and then take an exhaustive sequence  $G_1 \subset G_2 \subset \cdots$  of subgraphs of  $G = (V_G, E_G)$  determined by  $V_n = \{x \in V_G \mid d(v_0, x) \le n\}, E_n = \{\{x, y\} \in E_G \mid x, y \in V_n\}$  and  $G_n = (V_n, E_n)$  for each positive integer n.

We note that any divisor  $D = \sum_{x \in V_{n-1}} \ell(x)i(x)1_{\{x\}}$  on the finite subgraph  $G_n$  is viewed as the one on  $G_j$  for any j with j > n. In other words,  $D = \sum_{x \in V_{n-1}} \ell(x)i(x)1_{\{x\}}$  can be consistently viewed as a divisor on  $V_j$ .

A formal linear combination  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  of the family  $\{1_{\{x\}} \mid x \in V_G\}$  of indicator functions with at most countably many non-zero real coefficients  $\{\ell(x)i(x) \mid x \in V_G\}$  is called a divisor on G. Then, the divisor  $(D)_n = \sum_{x \in V_{n-1}} \ell(x)i(x)1_{\{x\}}$  called the restriction of D to  $G_n$  is regarded as a divisor vanishing outside  $V_{n-1}$  on the finite graph  $G_j$  for any integer j with  $j \geq n$ .

Our objective is to extend the Riemann-Roch theorem on finite graphs in [2] to the one on an infinite graph for divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  on G satisfying  $\sum_{x \in V_G} |\ell(x)|i(x) < \infty$ . To achieve this, it is necessary to discuss the convergence of  $r_n(D)$  defined in the second section on each  $G_n$ , at least in the case that  $\operatorname{supp}[D] = \{x \in V_G \mid \ell(x)i(x) \neq 0\}$  is a finite set. One of the key properties of the graphs for our aim is so-called a spectral gap of L. We assume some conditions on the infinity of G for taking our approach on the basis of spectral gap theory.

We denote the finite measure  $m_{G_n}$  given in the previous subsection by  $m_n$  and introduce the probability measure  $\mu_n$  given by  $\mu_n(A) = \frac{m_n(A)}{m_n(V_n)}$  for  $A \subset V_n$ . For  $x \in V_n, m_n(\{x\})$  is denoted briefly by  $m_n(x)$ .

In terms of a reversible Markov chain  $\{X_n\}$  determined by the transition matrix defined in (2), for any vertex x with  $d(x, v_0) = n$ , we see that  $P_x(X_1 \in V_n) = m_n(x)/m(x)$ ,

$$P_x(X_1 \in V_{n-1}) = \frac{1}{m(x)} \sum_{y \in N(x) \cap V_{n-1}} C_{x,y}$$

and these two identities imply  $P_x(X_1 \in V_n) \ge P_x(X_1 \in V_{n-1})$ . From those relationships among the probabilities,

$$P_x(X_1 \in V_{n-1}^c) \ge \frac{m(x) - m_n(x)}{m(x)}$$

Define

$$\rho_n = \sup_{x \in S_n} P_x(X_1 \in V_{n-1}^c) \tag{3}$$

for any positive integer n where  $S_n = \{x \in V_G \mid d(v_0, x) = n\}$ . In the sequel, we assume that

$$\limsup_{n \to \infty} \rho_n < 1. \tag{4}$$

By the assumption (4), we can take  $\rho < 1$  such that  $\limsup_{n\to\infty} \rho_n < \rho$ . Then we have

$$m(x) - m_n(x) \le \rho_n m(x) \text{ and } m(x) \le \frac{1}{1 - \rho} m_n(x)$$
 (5)

for any  $x \in S_n$  with sufficiently large n. Since  $m(x) = m_n(x)$  for any  $x \in V_{n-1}$ , the first estimate and the finiteness of the measure m imply that  $m(V_n) - m_n(V_n) \to 0$  as  $n \to \infty$ .

In what follows, we denote  $V_G$  by V, when it is preferable for the notation to require a shorter description.

Lemma 3.4. (4) implies the inequality

$$\sum_{x \in V_n, y \in N(x) \cap V_n^c} f(x)^2 C_{x,y} \le \rho_n m(V) ||f||_{L^2(\mu)}^2.$$

on real-valued function f on  $V_n^c$  for sufficiently large n.

*Proof.* The left hand side equals  $\sum_{x \in S_n} f(x)^2(m(x) - m_n(x))$  and (5) shows that this value is dominated by  $\rho_n \sum_{x \in S_n} f(x)^2 m(x)$  for sufficiently large n.

The following assertion is crucial to establish Riemann-Roch theorem in our approach.

**Theorem 3.5.** There exists a positive constant A with A < 1 such that

$$\limsup_{n \to \infty} \rho_n < A$$

implies the existence of a positive constant C such that

$$||f||^2_{L^2(\mu)} \le C\mathcal{E}(f, f),$$
 (6)

for any  $f \in L^2(\mu)$  with  $(f, 1)_{L^2(\mu)} = 0$ .

*Proof.* We first divide the sum of the left-hand side of (6) into two terms as

$$||f||_{L^{2}(\mu)}^{2} = \sum_{x \in V_{n}} f(x)^{2} \mu(x) + \sum_{x \in V_{n}^{c}} f(x)^{2} \mu(x).$$
(7)

The first term can be written as

$$\sum_{x \in V_n} f(x)^2 \mu(x) = \frac{m_n(V_n)}{m(V)} ||f||_{L^2(\mu_n)}^2 + \frac{1}{m(V)} \sum_{x \in V_n} f(x)^2 (m(x) - m_n(x)).$$

Since  $G_n$  is a finite graph, there exists a positive constant  $\lambda_n$  such that

$$\sum_{x \in V_n} f(x)^2 \mu(x) \le \frac{m_n(V_n)}{m(V)\lambda_n} \mathcal{E}^{(n)}(f,f) + \frac{m_n(V_n)}{m(V)} (f,1)_{L^2(\mu_n)}^2 + \nu_n(f)$$
$$\le \frac{m_n(V_n)}{m(V)\lambda_n} \mathcal{E}(f,f) + \frac{m_n(V_n)}{m(V)} (f,1)_{L^2(\mu_n)}^2 + \nu_n(f),$$

where  $\mathcal{E}^{(n)}(f,f) = \mathcal{E}_{G_n}(f,f)$  and  $\nu_n(f) = \frac{1}{m(V)} \sum_{x \in V_n} f(x)^2 (m(x) - m_n(x))$ . Since  $m(x) = m_n(x)$  for any  $x \in V_{n-1}$ , by Lemma 3.4 we obtain

$$\nu_n(f) \le \rho_n ||f||_{L^2(\mu)}^2$$

and by  $(f, 1)_{L^2(\mu)} = 0$ ,

$$(f,1)_{L^{2}(\mu_{n})}^{2} = \left(\frac{m(V)}{m(V_{n})}\sum_{x\in V_{n}^{c}}|f(x)|\mu(x)\right)^{2} \le \left(\frac{m(V)}{m(V_{n})}\right)^{2}\mu(V_{n}^{c})||f||_{L^{2}(\mu)}^{2}$$

Combining these estimates, we have

$$\sum_{x \in V_n} f(x)^2 \mu(x) \le \frac{m_n(V_n)}{m(V)} \lambda_n^{-1} \mathcal{E}(f, f) + \alpha_n ||f||_{L^2(\mu)}^2, \tag{8}$$

where  $\alpha_n = \rho_n + (\frac{m(V)}{m(V_n)})^2 \mu(V_n^c)$ .

Next, we estimate the second term of the right hand side of (7) by taking the function  $e^{ar(x)}$  on V with  $r(x) = d(x, v_0)$  and an arbitrarily fixed a > 0. Since

$$\Delta e^{ar(x)} = \sum_{y \in N(x), r(y) = r(x)+1} C_{x,y}(e^{ar(x)} - e^{ar(y)}) + \sum_{y \in N(x), r(y) = r(x)-1} C_{x,y}(e^{ar(x)} - e^{ar(y)})$$
$$= -(e^a - 1)e^{ar(x)}m_+(x) + (1 - e^{-a})e^{ar(x)}m_-(x),$$

where

$$m_+(x) = \sum_{y \in N(x), r(y) = r(x)+1} C_{x,y}$$
 and  $m_-(x) = \sum_{y \in N(x), r(y) = r(x)-1} C_{x,y}$ ,

we have  $Le^{ar(x)} = -(e^a - 1)e^{ar(x)}\frac{m_+(x)}{m(x)} + (1 - e^{-a})e^{ar(x)}\frac{m_-(x)}{m(x)}$ . Noting that

$$\frac{m_{-}(x)}{m(x)} = P_x(X_1 \in V_{n-1})$$

for any  $x \in S_n$  with  $n \ge 1$ , we have another expression on  $\rho_n$ ,  $\rho_n = \sup\{1 - \frac{m_-(x)}{m(x)} \mid x \in S_n\}$ , and thus

$$\frac{m_+(x)}{m(x)} \le 1 - \frac{m_-(x)}{m(x)} \le \rho_n$$

for any  $x \in S_n$  with  $n \ge 1$ .

From these observations, we can derive that

$$e^{-ar(x)}Le^{ar(x)} = -(e^a - 1)\frac{m_+(x)}{m(x)} + (1 - e^{-a})\frac{m_-(x)}{m(x)}$$
  

$$\geq -(e^a - 1)(1 - \frac{m_-(x)}{m(x)}) - (1 - e^{-a})(1 - \frac{m_-(x)}{m(x)}) + 1 - e^{-a}$$
  

$$\geq -(e^a - e^{-a})\rho_n + 1 - e^{-a}.$$

For the sequence  $\{\eta_n\}$  defined by  $\eta_n = -(e^a - e^{-a})\rho_n + 1 - e^{-a}$  (n = 1, 2, ...), we may assume that  $\eta_n > 0$  for any  $n \ge n_0$  with some positive integer  $n_0$ , more specifically,

$$\rho_n < \frac{1 - e^{-a}}{e^a - e^{-a}} \quad (n \ge n_0).$$
(9)

Let us apply (1) to the subgraph  $U = G_n^c$  determined by taking  $V_n^c$  as  $V_U$  and the function  $\phi$  defined by

$$\phi(x) = \begin{cases} e^{ar(x)} & (x \in V_{n-1}^c), \\ 0 & (x \in V_{n-1}). \end{cases}$$

Then, we have  $e^{-ar(x)}L\phi(x) = \frac{m_{G_n^c}(x)}{m(x)}e^{-ar(x)}L_{G_n^c}\phi(x) + \frac{1}{m(x)}\sum_{y\in N(x)\cap V_{n-1}}C_{x,y}$ , which

implies

$$\sum_{x \in V_n^c} f(x)^2 \mu(x) = \frac{1}{\eta_n} \sum_{x \in V_{n-1}^c} f(x)^2 e^{-ar(x)} L\phi(x) \mu(x) - \sum_{x \in S_n} f(x)^2 \mu(x)$$
  
$$= \frac{1}{\eta_n m(V)} \sum_{x \in V_{n-1}^c} f(x)^2 e^{-ar(x)} L_{G_{n-1}^c} \phi(x) m_{G_{n-1}^c}(x)$$
  
$$+ \frac{1}{m(V)} \Big\{ \frac{1}{\eta_n} \sum_{x \in V_{n-1}^c} f(x)^2 \sum_{y \in N(x) \cap V_{n-1}} C_{x,y} - \sum_{x \in S_n} f(x)^2 m(x) \Big\}.$$
  
(10)

We can regard the second term in the right-hand side as the one involving a boundary operator. We estimate the first term in the right hand side of (10). By the symmetry of  $L_{G_{n-1}^c}$  with respect to the inner product of  $L^2(m_{G_{n-1}^c})$ ,

$$\sum_{x \in V_{n-1}^c} f(x)^2 e^{-ar(x)} L_{G_{n-1}^c} \phi(x) m_{G_{n-1}^c}(x) = \sum_{x \in V_{n-1}^c} e^{ar(x)} L_{G_{n-1}^c}(f(x)^2 e^{-ar(x)}) m_{G_{n-1}^c}(x)$$
$$= \sum_{x \in V_{n-1}^c} e^{ar(x)} \Delta_{G_{n-1}^c}(f(x)^2 e^{-ar(x)}).$$

By direct calculations, we have

$$\Delta_{G_{n-1}^c}(f(x)^2 e^{-ar(x)}) = e^{-ar(x)} \Delta_{G_{n-1}^c}(f(x)^2) + f(x)^2 \Delta_{G_{n-1}^c}(e^{-ar(x)}) - \Gamma(f(x)^2, e^{-ar(x)}),$$
  
where  $\Gamma(u, v)(x) = \sum_{y \in N(x) \cap V_{n-1}^c} C_{x,y}(u(x) - u(y))(v(x) - v(y)).$  It is easy to see that

$$\Delta_{G_{n-1}^c}(f(x)^2) = 2f(x)\Delta_{G_{n-1}^c}f(x) - \Gamma(f, f)$$
(11)

and

$$\Delta_{G_{n-1}^c}(e^{-ar(x)}) = e^{-ar(x)}(1-e^{-a})m_+^{G_{n-1}^c}(x) - (e^a-1)e^{-ar(x)}m_-^{G_{n-1}^c}(x)$$
  
$$\leq e^{-ar(x)}(1-e^{-a})\rho_n m(x), \qquad (12)$$

where  $m_{+}^{G_{n-1}^{c}}(x) = \sum_{r(y)=r(x)+1, y \in V_{n-1}^{c}} C_{x,y}$ . We also see that

$$|\Gamma(f(x)^2, e^{-ar(x)})| \le (e^a - 1)e^{-ar(x)} \sum_{y \in N(x) \cap V_{n-1}^c} C_{x,y} |f(x) + f(y)| |f(x) - f(y)|.$$

From these observations, we have

$$\begin{split} \frac{1}{\eta_n m(V)} &\sum_{x \in V_{n-1}^c} e^{ar(x)} \Delta_{G_{n-1}^c}(f(x)^2 e^{-ar(x)}) \\ &\leq \frac{1}{\eta_n m(V)} \Big\{ \sum_{x \in V_{n-1}^c} (2f(x) \Delta_{G_{n-1}^c} f(x) - \Gamma(f, f)) \\ &+ (1 - e^{-a}) \rho_n \sum_{x \in V_{n-1}^c} f(x)^2 m(x) \\ &+ (e^a - 1) \sum_{x \in V_{n-1}^c} \sum_{y \in N(x) \cap G_{n-1}^c} C_{x,y} |f(x) + f(y)| |f(x) - f(y)| \Big\} \\ &\leq \frac{1}{\eta_n} \Big\{ \sum_{x \in V_{n-1}^c} 2f(x) \frac{1}{m(x)} \Delta_{G_{n-1}^c} f(x) \mu(x) \\ &+ (1 - e^{-a}) \rho_n \sum_{x \in V_{n-1}^c} f(x)^2 \mu(x) \\ &+ (e^a - 1) \sum_{x \in V_{n-1}^c} \sum_{y \in N(x) \cap V_{n-1}^c} C_{x,y} |f(x) + f(y)| |f(x) - f(y)| \Big\}. \end{split}$$

Lemma 3.2 implies that the first term in the right hand side is dominated by

$$\frac{\epsilon}{\eta_n}||f||^2_{L^2(\mu)} + \frac{2}{\epsilon m(V)\eta_n}\mathcal{E}(f,f)$$

for any  $\epsilon > 0$ . The second term is dominated by

$$\frac{(1-e^{-a})\rho_n}{\eta_n}||f||_{L^2(\mu)}^2.$$

As for the third term, by the Cauchy-Schwarz inequality, we have

$$\sum_{x \in V_{n-1}^c} \sum_{y \in N(x) \cap V_{n-1}^c} C_{x,y} |f(x) + f(y)| |f(x) - f(y)|$$
  
= 
$$\sum_{x \in V_{n-1}^c} \sum_{y \in V_{n-1}^c} 1_{\{x,y\} \in E_{V_{n-1}^c}} C_{x,y} |f(x) + f(y)| |f(x) - f(y)|$$
  
$$\leq \Big(\sum_{x \in V_{n-1}^c} \sum_{y \in V_{n-1}^c} 1_{\{x,y\} \in E_{V_{n-1}^c}} C_{x,y} |f(x) + f(y)|^2 \Big)^{1/2} (2\mathcal{E}(f,f))^{1/2}.$$

Fubini's theorem implies

$$\sum_{x \in V_{n-1}^c} \sum_{y \in V_{n-1}^c} \mathbb{1}_{\{x,y\} \in E_{V_{n-1}^c}} C_{x,y} |f(x) + f(y)|^2 \le 4m(V) ||f||_{L^2(\mu)}^2$$

Thus, the third term does not exceed

$$\frac{(e^a-1)\epsilon}{\eta_n}||f||_{L^2(\mu)}^2 + \frac{4(e^a-1)m(V)}{\epsilon\eta_n}\mathcal{E}(f,f).$$

We next estimate the term with the boundary operator in (10) as

$$\frac{1}{m(V)} \left\{ \frac{1}{\eta_n} \sum_{x \in V_{n-1}^c} f(x)^2 \sum_{y \in N(x) \cap V_{n-1}} C_{x,y} - \sum_{x \in S_n} f(x)^2 m(x) \right\}$$
$$= \frac{1}{m(V)} \sum_{x \in S_n} f(x)^2 \left( \frac{1}{\eta_n} \sum_{y \in N(x) \cap V_{n-1}} C_{x,y} - m(x) \right)$$
$$\leq \frac{1}{m(V)} \sum_{x \in S_n} f(x)^2 \left( \frac{1}{\eta_n} - 1 \right) m_-(x)$$
$$\leq \left( \frac{1}{\eta_n} - 1 \right) ||f||_{L^2(\mu)}^2.$$

Combining these estimates, we have

$$\sum_{x \in V_n^c} f(x)^2 \mu(x) \le \beta(n,\epsilon) ||f||_{L^2(\mu)}^2 + \gamma(\epsilon,n) \mathcal{E}(f,f),$$
(13)

where  $\beta(\epsilon, n) = \frac{(e^a+1)\epsilon+2(1-e^{-a})\rho_n}{2\eta_n} + \frac{1}{\eta_n} - 1$  and  $\gamma(\epsilon, n) = \frac{2m(V)^{-1}+4(e^a-1)m(V)}{\epsilon\eta_n}$ . From (8) and (13), we can derive

$$||f||_{L^{2}(\mu)}^{2} \leq (1 + \lambda_{n}^{-1} + \gamma(\epsilon, n)) \mathcal{E}(f, f) + (\alpha_{n} + \beta(\epsilon, n)) ||f||_{L^{2}(\mu)}^{2}.$$

Here, we consider a condition on  $\rho_n$  ensuring that  $\alpha_n + \beta(\epsilon, n) < 1$ . Let B(a), B'(a) be the smaller and larger solutions of the quadratic equation

$$(e^{a} - e^{-a})t^{2} - 2(e^{a} - 2e^{-a} + 1)t + 1 - 2e^{-a} = 0,$$

respectively. Then  $\rho_n < B(a)$  implies

$$\rho_n + \frac{(1 - e^{-a})\rho_n}{\eta_n} + \frac{1}{\eta_n} - 1 < 1.$$

We can take the maximum of B(a) subject to  $a > \log 2$  as the positive constant A in the statement of Theorem 3.5. In fact, B(a) > 0 for  $a > \log 2$  and under the condition that  $\limsup_{n\to\infty} \rho_n < \max\{B(a) \mid a > \log 2\}$ , we can easily take a positive real a such that the condition (9) is satisfied for sufficiently large n. (This is because  $B(a) < \frac{1-e^{-a}}{e^a - e^{-a}} < B'(a)$  for any a). Consequently, there exist  $n_G$  and  $\epsilon_0 > 0$  such that  $\alpha_{n_G} + \beta(\epsilon_0, n_G) < 1$ . Therefore, we have

$$||f||_{L^{2}(\mu)}^{2} \leq \frac{1 + \lambda_{n_{G}}^{-1} + \gamma(\epsilon_{0}, n_{G})}{1 - \alpha_{n_{G}} - \beta(\epsilon_{0}, n_{G})} \mathcal{E}(f, f).$$
(14)

**Remark 3.6.** When we apply this theorem for a function f on subgraph  $G_n = (V_n, E_n)$ , i.e., a function with its support contained in  $V_n$  and  $(1, f)_{L^2(\mu_n)} = 0$ , we need to restrict our attention to such sufficiently large n that  $\rho_n < A$  holds. This point will be used when we apply Lemma 3.10 below in Section 4.

**Remark 3.7.** By a numerical calculation, one sees that  $A = \max\{B(a) \mid a > \log 2\} \approx 0.0569$ .

For a connected subgraph U of G, we introduce

$$\lambda_U = \inf\{\tilde{\mathcal{E}}_U(f,f) \mid f \in L^2(\mu_U), (f,1)_{L^2(\mu_U)} = 0 \text{ and } \|f\|_{L^2(\mu_U)} = 1\}$$

called a spectral gap of  $L_U$ , where  $\tilde{\mathcal{E}}_U(u, v) = m_U(V_U)^{-1}\mathcal{E}_U(u, v)$ . Similarly to the relationship between the Laplace operator  $L_U$  on  $L^2(m_U)$  and  $\mathcal{E}_U$ , we note that  $(L_U u, v)_{L^2(\mu_U)} = \tilde{\mathcal{E}}_U(u, v)$ . It is well known that if U is a finite graph, then  $\lambda_U > 0$ (cf. [7]). The spectral gap  $\lambda_n$  of  $L_{G_n}$  is taken in the first paragraph of the proof of Theorem 3.5, the assertion of which shows the positivity of the spectral gap of L.

**Cororally 3.8.** Under the assumption of Theorem 3.5, we have

$$\lambda_G \ge \frac{1}{Cm(V_G)}$$

**Lemma 3.9.** Let  $\lambda_n$  be the spectral gap of  $L_{G_n}$  and assume  $\lambda_G > 0$ . If  $\limsup_{n \to \infty} \frac{\rho_n}{1-\rho_n} < \lambda_G$ , then  $\liminf_{n \to \infty} \lambda_n > 0$ . Moreover, if  $\lim_{n \to \infty} \rho_n = 0$ , then

$$\liminf_{n \to \infty} \lambda_n \ge \lambda_G.$$

Proof. Since  $G_n$  is a finite graph,  $\lambda_n$  is the smallest non-zero eigenvalue of  $L_{G_n}$ and any eigenfunction  $\psi_n$  associated with  $\lambda_n$  satisfies  $(1, \psi_n)_{L^2(\mu_n)} = 0$ . We assume that the eigenfunction is normalized as  $||\psi_n||_{L^2(\mu_n)} = 1$  beforehand. We extend this function to the one defined on V taking identically zero on  $V_n^c$  and denote it again by  $\psi_n$ . Let  $\overline{\psi_n} = \psi_n - (1, \psi_n)_{L^2(\mu)}$  and we can first show that  $||\overline{\psi_n}||_{L^2(\mu)} \ge m(V_n)m(V)^{-1}$ . In fact, the identities

$$||\overline{\psi_n}||_{L^2(\mu)} = \sum \left(\psi_n(x) - (1, \psi_n)_{L^2(\mu)}\right)^2 \mu(x)$$
  
=  $\sum \psi_n(x)^2 \mu(x) - (1, \psi_n)_{L^2(\mu)}^2$   
=  $m(V)^{-1} \sum \psi_n(x)^2 (m(x) - m_n(x)) + m_n(V_n) m(V)^{-1}$   
 $- \left(m(V)^{-1} \sum \psi_n(x) (m(x) - m_n(x))\right)^2$ 

and the Cauchy-Schwarz inequality yield a lower bound

$$m(V)^{-1} \sum \psi_n(x)^2 (m(x) - m_n(x)) + m_n(V_n) m(V)^{-1} - m(V)^{-2} (m(V) - m_n(V_n)) \sum \psi_n(x)^2 (m(x) - m_n(x)) \geq m_n(V_n) m(V)^{-1}$$

of the right-hand side. Since  $m(x) - m_n(x) \le \frac{\rho_n}{1-\rho_n} m_n(x)$ ,

$$\mathcal{E}(\overline{\psi_n}, \overline{\psi_n}) = \mathcal{E}_{G_n}(\overline{\psi_n}, \overline{\psi_n}) + \sum_{x \in G_n, y \in V_n^c} C_{x,y} \psi_n(x)^2$$
$$= m_n(V_n)\lambda_n + \sum_{x \in V_n} \psi_n(x)^2 (m(x) - m_n(x))$$
$$\leq m_n(V_n)\lambda_n + \frac{\rho_n}{1 - \rho_n} m_n(V_n).$$

As a result, it turns out that

$$m_n(V_n)^{-1}m(V)\lambda_G \le \lambda_n + \frac{\rho_n}{1-\rho_n}.$$

**Lemma 3.10.** If  $\limsup_{n\to\infty} \frac{\rho_n}{1-\rho_n} < \lambda_G$  as assumed in Lemma 3.9, then there exists some positive constant K such that any sequence  $\{g_n\}$  of functions  $g_n$  on  $V_n$  satisfying  $(1, g_n)_{L^2(\mu_n)} = 0$  for sufficiently large n yields

$$||R^{(n)}g_n||_{L^2(\mu_n)} \le K||g_n||_{L^2(\mu_n)}$$

for sufficiently large n, where  $R^{(n)}$  stands for the 0-order resolvent of  $L_{G_n}$  on  $L^2(\mu_n)$ .

*Proof.* For the transition operator  $P_t^{(n)}$  associated with the generator  $L_{G_n}$  on  $L^2(\mu_n)$ , the positivity of spectral gaps implies (cf.[11])

$$||P_t^{(n)}g_n||_{L^2(\mu_n)} \le e^{-\lambda_n t} ||g_n||_{L^2(\mu_n)}.$$

Note that  $R^{(n)}g_n(x) = \int_0^\infty P_t^{(n)}g_n(x)dt$  and

$$R^{(n)}g_n(x) = \sum_{k=1}^{\infty} R_1^k g_n(x)$$

where  $R_1g_n(x) = \int_0^\infty e^{-t}P_t^{(n)}g_n(x)dt$  and  $R_1^k$  denotes the operator given as the ktimes iteration of  $R_1$  (cf. [6]). Lemma 3.9 implies that there exists  $\delta > 0$  such that  $\lambda_n \geq \delta$  for all n. By applying Jensen's inequality and Fubini's theorem, we obtain

$$||R_1g_n||_{L^2(\mu_n)}^2 \le \frac{1}{2\lambda_n + 1} ||g_n||_{L^2(\mu_n)}^2 \le \frac{1}{2\delta + 1} ||g_n||_{L^2(\mu_n)}^2$$

for sufficiently large n. The desired result follows from this estimate.

4 Proof of Riemann-Roch theorem on an infinite graph.

In this section, we establish a Riemann-Roch theorem on the connected infinite graph  $G = (V_G, E_G)$  with local finiteness and finite volume as in the last section, by applying  $L^2$ -boundedness of the 0-order resolvent derived from spectral gap theory to a sequence of functions in the images of the Laplace operator. For that purpose, we first take a divisor  $D = \sum_{x \in V_{n-1}} \ell(x)i(x)1_{\{x\}}$  on some subgraph  $G_n = (V_n, E_n)$  of  $G = (V_G, E_G)$  as given in the last section and denote  $\Delta_{G_n}$  and  $L_{G_n}$  by  $\Delta_n$  and by  $L_n$ , respectively. The equivalence between divisors  $D' = \sum_{x \in V_n} \ell'(x)i(x)1_{\{x\}}$  and  $D'' = \sum_{x \in V_n} \ell''(x)i(x)1_{\{x\}}$  with  $\mathbb{Z}$ -valued functions  $\ell'$  and  $\ell''$  is defined by  $D'' = D' + \Delta_n f$  on  $V_n$  for some  $\mathbb{Z}$ -valued function f with  $\operatorname{supp}[f] \subset V_n$ . This relationship will be denoted by  $D' \stackrel{n}{\sim} D''$  and will be called n-equivalence. The family of total orders on  $V_n$  is denoted by  $\mathcal{O}_n$ .

For a divisor D on the finite graph  $G_n = (V_n, E_n)$ ,  $r_n(D)$  is defined on the finite subgraph  $G_n$ , by replacing "~" in the second section with the *n*-equivalence "~", more specifically given by

$$r_n(D) = \left(\min_{D' \stackrel{n}{\sim} D, O_n \in \mathcal{O}_n} \deg^+(D' - \nu_{O_n})\right) - i_{(G_n, C_n)}$$

When a divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  satisfying  $\sum_{x \in V_G} |\ell(x)|i(x)| < \infty$  is given, we denote  $\sum_{x \in V_G, \ell(x) > 0} \ell(x)i(x)$  by deg<sup>+</sup>(D) and  $-\sum_{x \in V_G, \ell(x) < 0} \ell(x)i(x)$  by deg<sup>-</sup>(D). The family  $\{(D)_n\}$  of divisors is consistent in the sense that  $((D)_j)_n = (D)_n$  whenever j > n. Later, by taking control over the sequence  $\{O_j\}$  of total orders each of which is taken in the minimization of  $r_j((D)_n) = \min_{D' \stackrel{j}{\sim} (D)_n, O_j \in \mathcal{O}_j} \deg^+(D' - \nu_{O_j}) - i_{(G_j, C_j)}$ on every subgraph  $G_j$  satisfying  $V_j \supset V_{n-1} \supset \operatorname{supp}[(D)_n]$ , we facilitate successive procedures of taking limits as  $j \to \infty$  and  $n \to \infty$  in  $r_j((D)_n)$ . In fact, thanks to Proposition 2.11, we see that  $|r_j((D)_n) - r_j((D)_{n'})|$  is sufficiently small as long as n, n' are both large enough, independently of j with  $j > \max\{n, n'\}$ . This is because deg<sup>+</sup> $((D)_n - (D)_{n'}) + \operatorname{deg}^-((D)_n - (D)_{n'})$  is sufficiently small as long as n, n' are both large enough.

In the following two lemmas, we focus only on a divisor D satisfying  $\operatorname{supp}[D] \subset V_{j_0}$ for some  $j_0$  and the case that  $\rho_n < \min\{A, \frac{\lambda_G}{\lambda_G+1}\}$  is satisfied for sufficiently large  $n > j_0$ . We note that, in this case, the uniform boundedness of the family  $\{R^{(n)}\}$ of operators is derived from the proof of Lemma 3.10. We denote a minimizer for  $r_n(D)$  by  $f_n$ . To be more precise,  $f_n$  is an integer-valued function on  $V_n$  such that  $D' = D + \Delta_n f_n$  attains the minimum

$$r_n(D) = \left(\min_{D' \stackrel{n}{\sim} D, O_n \in \mathcal{O}_n} \deg^+(D' - \nu_{O_n})\right) - i_{(G_n, C_n)}$$

with some total order  $O_n \in \mathcal{O}_n$ .

For such a divisor D the divisor  $(D)_j$  is equal to D for any j with  $j \ge n$ . To show the convergence of a subsequence of  $\{r_j(D)\}$ , we will need an upper bound of  $|f_n|$  on  $V_n$ . In fact, the boundedness enables us to take the real-valued harmonic extension  $h_j^{(n)}$  of  $f_n$  on  $V_j \setminus V_n$  as seen after the following lemma, which yields a sequence of integer-valued functions used for deriving the convergence of a subsequence later.

**Lemma 4.1.** For sufficiently large integer n,  $r_n(D)$  admits a minimizer  $f_n$  satisfying  $||f_n||_{L^2(\mu_n)}^2 \leq 4m(V)K((\deg^+(D) + \deg^-(D) + m_n(V_n))^2 / \min\{m(x) \mid x \in V_n\})$ , where K stands for the constant given in Lemma 3.10. In particular,

$$\max_{x \in V_n} |f_n(x)| \le 2\sqrt{m(V)K} (\deg^+(D) + \deg^-(D) + 1) / \min\{m(x) \mid x \in V_n\}$$

for sufficiently large n.

*Proof.* Any  $\mathbb{Z}$ -valued function  $f_D^{(n)}$  on  $V_n$  minimizing deg<sup>+</sup> $(D + \Delta_n f_D^{(n)})$  satisfies deg<sup>+</sup> $(D + \Delta_n f_D^{(n)}) \leq$  deg<sup>+</sup>D. Accordingly, Lemma 2.1 (ii) shows that deg<sup>-</sup> $(D + \Delta_n f_D^{(n)}) \leq$ 

 $\Delta_n f_D^{(n)}) \leq \deg^- D, \text{ which implies } |\Delta_n f_D^{(n)}(x)| \text{ does not exceed } 2(\deg^+(D) + \deg^-(D))$ for any  $x \in V_n$  and  $f_D^{(n)}$  admits the estimate  $\|\frac{1}{m}\Delta_n f_D^{(n)}\|_{L^2(\mu_n)}^2 \leq 4m(V)(\deg^+(D) + \deg^-(D))^2 / \min\{m(x) \mid x \in V_n\}.$  Since  $|\nu_{O_n}(x)| \leq m(x)$  for every  $x \in V_n$  and  $O_n \in \mathcal{O}_n$ , any minimizer  $f_n$  for  $r_n(D) = \deg^+(D + \Delta_n f_n - \nu_{O_n})$ , which is viewed as  $f_{D+\nu_{O_n}}^{(n)}$ , satisfies

$$\left\|\frac{1}{m}\Delta_n f_n\right\|_{L^2(\mu_n)}^2 \le 4m(V)(\deg^+(D) + \deg^-(D) + m_n(V_n))^2 / \min\{m(x) \mid x \in V_n\}.$$

Thanks to Lemma 3.10, these inequalities imply  $||R^{(n)}L_n f_n||^2_{L^2(\mu_n)} \leq 4m(V)$  $K((\deg^+(D) + \deg^-(D) + m_n(V_n))^2 / \min\{m(x) \mid x \in V_n\})$ .

The uniqueness of the solution to the Poisson equation on connected finite graphs shows that  $y, z \in V_n$  implies  $R^{(n)}L_nf_n(y) - R^{(n)}L_nf_n(z) \in \mathbb{Z}$ . This and the uniqueness up to difference given by constants allow us to take such a real constant c with  $0 \leq c < 1$  that  $R^{(n)}L_nf_n+c$  is not only  $\mathbb{Z}$ -valued but a minimizer for  $r_n(D)$ . We focus only on this minimizer obtained by this procedure and denote it by  $f_n$ , then  $f_n$  admits the estimate in the assertion and  $\max_{x \in V_n} |f_n(x)| \leq 2\sqrt{m(V)K} ((\deg^+(D) + \deg^-(D) + 1)/\min\{m(x) \mid x \in V_n\})$  is satisfied for sufficiently large n. In fact, this follows from  $|f_n(x)|^2 m(x) \leq 4m(V)K ((\deg^+(D) + \deg^-(D) + 1)^2/\min\{m(x) \mid x \in V_n\})$  for any  $x \in V_n$  as long as n is sufficiently large.  $\square$ 

**Lemma 4.2.** For sufficiently large integer n, the minimizer  $f_n$  in the last lemma for  $r_n(D)$  admits a sequence  $\{f_i^{(n)}\}_{j>n}$  of functions such that

- (i)  $f_n = f_j^{(n)} \text{ on } V_n$ ,
- (ii)  $\max_{x \in V_j} |f_j^{(n)}(x)| \le 2\sqrt{m(V)K}(\deg^+(D) + \deg^-(D) + 1)/\min\{m(x) \mid x \in V_n\} + 1,$
- (iii)  $\lim_{n\to\infty} \sup_{j>n} \|m_j(\cdot)^{-1} (\Delta_j f_j^{(n)} D)\|_{L^1(V_j \setminus V_n;\mu)} = 0.$

*Proof.* We take the harmonic extension  $h_j^{(n)}$  of  $f_n$  on  $V_j \setminus V_n$  determined by  $\Delta_j h_j^{(n)} = 0$  on  $V_j \setminus V_n$  and  $h_j^{(n)} = f_n$  on  $V_n$ . The function  $g_j^{(n)}$  defined as the integer part of  $h_j^{(n)}$  admits the estimate

$$\frac{|\Delta_j g_j^{(n)}(x)|}{m(x)}\Big|_{V_j \setminus V_n} \le 21_{V_j \setminus V_n}$$

on  $\Delta_j g_j^{(n)}(x) = \sum_{y \in N(x) \cap V_j} C_{x,y}(g_j^{(n)}(x) - g_j^{(n)}(y)).$ 

In fact, since  $\max_{x \in V_j \setminus V_{n-1}} |h_j^{(n)}(x) - g_j^{(n)}(x)| \le 1$ , we see

$$C_{x,y}|g_j^{(n)}(x) - h_j^{(n)}(x) - g_j^{(n)}(y) - h_j^{(n)}(y)| \le 2C_{x,y}$$

for any  $x \in V_j \setminus V_n$  and  $y \in N(x)$ . By combining this with the harmonicity of  $h_j^{(n)}$ on  $V_j \setminus V_{n-1}$ , equivalently  $x \in V_j \setminus V_n \Rightarrow \sum_{y \in N(x) \cap V_j} C_{x,y}(h_j^{(n)}(x) - h_j^{(n)}(y)) = 0$ , we see that  $\frac{|\sum_{y \in N(x) \cap V_j} C_{x,y}(g_j^{(n)}(x) - g_j^{(n)}(y))|}{m(x)} \leq 2$  for any  $x \in V_j \setminus V_n$ .

By applying the maximal principle to the function  $h_j^{(n)}$ , we observe  $\max_{x \in V_j} |h_j^{(n)}(x)| \le 2\sqrt{m(V)K}(\deg^+(D) + \deg^-(D) + 1)/\min\{m(x) \mid x \in V_n\}$  which implies  $\max_{x \in V_j} |g_j^{(n)}(x)| \le 2\sqrt{m(V)K}(\deg^+(D) + \deg^-(D) + 1)/\min\{m(x) \mid x \in V_n\} + 1.$ 

Accordingly, we see that the function  $f_j^{(n)}$  taking  $f_n$  on  $V_n$  and  $g_j^{(n)}$  on  $V_n^c$  enjoys  $\|\frac{1}{m}\Delta_j f_j^{(n)} \mathbf{1}_{V_n^c}\|_{L^1(\nu)} \leq 2m(V_j \setminus V_n)$ . In other words,  $f_j^{(n)}$  meets the conditions (i)-(iii) in the assertion.

In what follows, we discuss divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  satisfying  $D = \sum_{x \in V_G} |\ell(x)|i(x) < \infty$ . In the next lemma, we start with a divisor  $(D)_{n_0(\varepsilon)}$  satisfying deg<sup>+</sup> $(D - (D)_{n_0(\varepsilon)}) + \deg^-(D - (D)_{n_0(\varepsilon)}) < \varepsilon$  for a given  $\varepsilon > 0$  and we only focus on the subgraphs  $G_n = (V_n, E_n)$  with  $n \ge n_0(\varepsilon)$ . Here and in the sequel, we assume that  $n_0(\varepsilon)$  is sufficiently large as mentioned at the beginning of this section. For any pair of positive integers j, n with  $j > n \ge n_0(\varepsilon)$  and total order  $O_j \in \mathcal{O}_j$ ,

the restriction of the total order of  $O_i$  to  $V_n$  is denoted by  $O_i|_{V_n}$ .

Here, we make an attempt on choice of subsequence  $\{n_k\}$  of positive integers so that the consistency in the sense  $O_{n_j}|_{V_k} = O_{n_k}$  is satisfied for any pair j, k of integers with j > k. Then, not only is the convergence of subsequence of  $\{r_j((D)_n)\}$  for an arbitrarily fixed n required, but a procedure of taking limit as  $j \to \infty$  and  $n \to \infty$  in the doubly indexed sequence  $\{r_j((D)_n)\}$  should be reasonably organized to determine a characteristic value of such a divisor D for our Riemann-Roch theorem. In the following lemma, we obtain such reasonable subsequence that we can eventually obtain the limit which makes sense for that purpose.

**Lemma 4.3.** For any  $\varepsilon > 0$ , there exists a sequence  $\{O_{N(\varepsilon/2^l)}\}$  of total orders satisfying  $O_{N(\varepsilon/2^j)} \in \mathcal{O}_{N(\varepsilon/2^j)}$  with  $m(V_{N(\varepsilon/2^j)}^c) < \varepsilon/2^j$  for any non-negative integer j and a sequence  $\{n_j\}$  satisfying  $n_1 < n_2 < \ldots$  and  $n_{j+1} \ge N(\varepsilon/2^j)$  for any non-negative integer j such that

$$r_{n_k}((D)_{n_l}) = \left(\min_{D' \stackrel{n_k}{\sim} (D)_{n_l}, O_{N(\varepsilon/2^l)} = O_{n_k}|_{V_{N(\varepsilon/2^l)}}, O_{n_k} \in \mathcal{O}_{n_k}} \deg^+ (D' - \nu_{O_{n_k}})\right) - i_{(G_{n_k}, C_{n_k})},$$
(15)

whenever k > l. In particular, k > l implies  $O_{N(\varepsilon/2^l)} = O_{n_k}|_{V_{N(\varepsilon/2^l)}}$  and

$$\deg^{+}(\nu_{O_{n_{l}}} - \nu_{O_{n_{k}}}) + \deg^{-}(\nu_{O_{n_{l}}} - \nu_{O_{n_{k}}}) < m(V_{N(\varepsilon/2^{\min\{k,l\}})}^{c})) < \varepsilon/2^{\min\{k,l\}}$$
(16)

for any positive integers k and l.

*Proof.* We take a divisor  $(D)_{n_0(\varepsilon)}$  satisfying  $\deg^+(D - (D)_{n_0(\varepsilon)}) + \deg^-(D - (D)_{n_0(\varepsilon)}) < \varepsilon$ . Since the finiteness  $m(V) < \infty$  implies  $\lim_{n\to\infty} m(V_n) = m(V)$ , there exists a positive integer  $N(\varepsilon)$  such that  $m(V_{N(\varepsilon)}^c) < \varepsilon$ . We may assume that  $n_0(\varepsilon) \ge N(\varepsilon)$ . Since the cardinality of  $\mathcal{O}_{N(\varepsilon)}$  is finite, the sequence  $O_{N(\varepsilon)+1} \in \mathcal{O}_{N(\varepsilon)+1}, O_{N(\varepsilon)+2} \in \mathcal{O}_{N(\varepsilon)+2}, \ldots$  admits a subsequence  $O_{n_1(\varepsilon)}, O_{n_2(\varepsilon)}, \ldots$  with  $N(\varepsilon) \le n_0(\varepsilon) \le n_1(\varepsilon) < n_2(\varepsilon) < \ldots$  such that  $O_{N(\varepsilon)} = O_{n_k(\varepsilon)}|_{V_{N(\varepsilon)}}$  and

$$r_{n_{k}(\varepsilon)}((D)_{n_{0}(\varepsilon)}) = \left(\min_{D'^{n_{k}(\varepsilon)}(D)_{n_{0}(\varepsilon)}, O_{N(\varepsilon)}=O_{n_{k}(\varepsilon)}|_{V_{N(\varepsilon)}}, O_{n_{k}(\varepsilon)}\in\mathcal{O}_{n_{k}(\varepsilon)}} \deg^{+}(D'-\nu_{O_{n_{k}(\varepsilon)}})\right)$$
$$-i_{(G_{n_{k}(\varepsilon)}, C_{n_{k}(\varepsilon)})}$$

for any  $k \geq 1$ . We may assume that  $\deg^+(D-(D)_{n_1(\varepsilon)}) + \deg^-(D-(D)_{n_1(\varepsilon)}) < \varepsilon/2$ 

By taking sufficiently large  $n_2(\varepsilon)$ , we may concentrate our attention to the case  $n_2(\varepsilon) \ge N(\varepsilon/2)$ . Since the cardinality of  $\mathcal{O}_{N(\varepsilon/2)}$  is finite, the sequence  $O_{n_2(\varepsilon)}, O_{n_3(\varepsilon)}, O_{n_4(\varepsilon)}, \ldots$  admits a subsequence  $O_{n_2(\varepsilon/2)}, O_{n_3(\varepsilon/2)}, \ldots$  with  $N(\varepsilon/2) \le n_2(\varepsilon/2) < n_3(\varepsilon/2) < \ldots$  such that  $O_{N(\varepsilon/2)} = O_{n_k(\varepsilon/2)}|_{V_{N(\varepsilon/2)}}$  and

$$r_{n_{k}(\varepsilon/2)}((D)_{n_{1}(\varepsilon)}) = \left(\min_{\substack{D'^{n_{k}(\varepsilon/2)}(D)_{n_{1}(\varepsilon)}, O_{N(\varepsilon/2)} = O_{n_{k}(\varepsilon/2)}|_{V_{N(\varepsilon/2)}}, O_{n_{k}(\varepsilon/2)} \in \mathcal{O}_{n_{k}(\varepsilon/2)}} \operatorname{deg}^{+}(D' - \nu_{O_{n_{k}(\varepsilon/2)}})\right)$$
$$- i_{(G_{n_{k}(\varepsilon/2)}, C_{n_{k}(\varepsilon/2)})}$$

for any  $k \geq 2$ . We may assume that  $\deg^+(D-(D)_{n_2(\varepsilon/2)}) + \deg^-(D-(D)_{n_2(\varepsilon/2)}) < \varepsilon/4$ 

By repeating this procedure, we obtain a subsequence  $n_{j+1}(\varepsilon/2^j), n_{j+2}(\varepsilon/2^j), \cdots$ of  $n_{j+1}(\varepsilon/2^{j-1}), n_{j+2}(\varepsilon/2^{j-1}), \cdots$  with  $N(\varepsilon/2^j) \leq n_{j+1}(\varepsilon/2^j) < n_{j+2}(\varepsilon/2^j) < \cdots$ such that  $O_{N(\varepsilon/2^j)} = O_{n_k(\varepsilon/2^j)}|_{V_{N(\varepsilon/2^j)}}$  and

$$\begin{split} r_{n_{k}(\varepsilon/2^{j})}((D)_{n_{j}(\varepsilon/2^{j-1})}) \\ &= \Big(\min_{D'^{n_{k}(\varepsilon/2^{j})}(D)_{n_{j}(\varepsilon/2^{j-1})}, O_{N(\varepsilon/2^{j})} = O_{n_{k}(\varepsilon/2^{j})}|_{V_{N(\varepsilon/2^{j})}, O_{n_{k}(\varepsilon/2^{j})} \in \mathcal{O}_{n_{k}(\varepsilon/2^{j})}} } \operatorname{deg}^{+}(D' - \nu_{O_{n_{k}(\varepsilon/2^{j})}})\Big) \\ &- i_{(G_{n_{k}(\varepsilon/2^{j})}, C_{n_{k}(\varepsilon/2^{j})})} \end{split}$$

for any k > j. We may assume that  $\deg^+(D-(D)_{n_{j+1}(\varepsilon/2^j)}) + \deg^-(D-(D)_{n_{j+1}(\varepsilon/2^j)}) < \varepsilon/2^j$ .

As a result, by taking  $n_1 = n_1(\varepsilon), n_2 = n_2(\varepsilon/2), \ldots$ , we obtain the sequence  $n_1, n_2, \ldots$  which meets all conditions in the assertion.

(16) follows from the straightforward estimate  $|\nu_{O_n}(x) - \nu_{O_{n'}}(x)| \le m(x)$  for any  $x \in V_n \cap V_{n'}$  with  $O_n|_{V_{N(\varepsilon)}} = O_{n'}|_{V_{N(\varepsilon)}}$ .

We are now in position to assert the convergence of the sequence  $\{r_{n_k}((D)_{n_l})\}_{k=l+1}^{\infty}$ as  $k \to \infty$ , for any fixed positive integer l.

### Proposition 4.4. If

$$\rho_n m(S_n) / \min_{x \in V_n} m(x) \to 0 \text{ as } n \to \infty,$$
(17)

then  $\{r_{n_k}((D)_{n_l})\}_{k=l+1}^{\infty}$  converges as  $k \to \infty$ , for any fixed non-negative integer l, where  $n_1, n_2, \ldots$  is the subsequence satisfying (15) associated with a sequence  $\{O_{N(\varepsilon/2^j)}\}$  of total orders in Lemma 4.3.

*Proof.* First we note that, the assumptions in Theorem 3.5 and Lemma 3.9 are satisfied since (17) implies  $\limsup_{n\to\infty} \rho_n < \min\{A, \frac{\lambda_G}{\lambda_G+1}\}$ .

On the other hand, we also note that it suffices to show the assertion in the case l = 1 because the following proof can be performed for any other positive integer l. In what follows,  $(D)_{n_1}$  will be denoted simply by D. From now on, we use notation  $\ell$  persistently for assigning an integer  $\ell$  satisfying  $\ell > k$  so that confusion with the use of notation l assigning integer less than k is avoided for comparing  $r_{n_\ell}(D)$  with  $r_{n_k}(D)$ . Here, we take a minimizer  $f_{n_\ell}$  for  $r_{n_\ell}(D)$  and a minimizer  $f_{n_k}$  for the other one.

By applying the identity just before (1) for the operators  $\Delta_{n_{\ell}}$  and  $\Delta_{n_k}$  instead of L and  $L_U$ , we have  $\Delta_{n_{\ell}}\phi(x) = \Delta_{n_k}\phi(x) + \left(\sum_{y \in N(x) \cap S_{n_k+1}} C_{x,y}(\phi(x) - \phi(y))\right) \mathbf{1}_{S_{n_k}} + C_{x,y}(\phi(x) - \phi(y))$ 

 $\Delta_{n_{\ell}}\phi(x)1_{V_{n_{\ell}}\setminus V_{n_{k}}}$  on  $V_{n_{k}}$  for any function  $\phi$  defined on  $V_{n_{\ell}}$ , where we recall  $S_{n_{k}} = \{x|d(v_{0},x) = n_{k}\}$ . Accordingly, for the function  $f_{n_{\ell}}^{(n_{k})}$  taken in Lemma 4.2 by starting with the minimizer  $f_{n_{k}}$ , we have

$$\begin{aligned} r_{n_{\ell}}(D) &= \deg^{+}(D'_{n_{\ell}} - \nu_{O_{n_{\ell}}}) - i(G_{n_{\ell}}, C_{n_{\ell}}) \\ &= \deg^{+}(D + \Delta_{n_{\ell}}f_{n_{\ell}} - \nu_{O_{n_{\ell}}}) - i(G_{n_{\ell}}, C_{n_{\ell}}) \\ &\leq \deg^{+}(D + \Delta_{n_{\ell}}f_{n_{\ell}}^{(n_{k})} - \nu_{O_{n_{\ell}}}) - i(G_{n_{k}}, C_{n_{k}}) + |i(G_{n_{\ell}}, C_{n_{\ell}}) - i(G_{n_{k}}, C_{n_{k}}) \\ &= \deg^{+}(D + \Delta_{n_{k}}f_{n_{k}} + \Big(\sum_{y \in N(x) \cap S_{n_{k}+1}} C_{x,y}(f_{n_{\ell}}^{(n_{k})}(x) - f_{n_{\ell}}^{(n_{k})}(y))\Big) 1_{S_{n_{k}}} \\ &+ \Delta_{n_{\ell}}f_{n_{\ell}}^{(n_{k})} 1_{V_{n_{k}}^{c}} - \nu_{O_{n_{\ell}}}) - i(G_{n_{k}}, C_{n_{k}}) + |i(G_{n_{\ell}}, C_{n_{\ell}}) - i(G_{n_{k}}, C_{n_{k}})|. \end{aligned}$$

Lemma 4.2 shows that

$$deg^{+}\left(\sum_{x \in S_{n_{k}}} \sum_{y \in N(x) \cap S_{n_{k}+1}} C_{x,y}(f_{n_{\ell}}^{(n_{k})}(x) - f_{n_{\ell}}^{(n_{k})}(y))\right) \\ + deg^{-}\left(\sum_{x \in S_{n_{k}}} \sum_{y \in N(x) \cap S_{n_{k}+1}} C_{x,y}(f_{n_{\ell}}^{(n_{k})}(x) - f_{n_{\ell}}^{(n_{k})}(y))\right) \\ \leq \sum_{x \in S_{n_{k}}} \sum_{y \in N(x) \cap S_{n_{k}+1}} C_{x,y}(|f_{n_{\ell}}^{(n_{k})}(x)| + |f_{n_{\ell}}^{(n_{k})}(y)|) \\ \leq 4\left(\sqrt{m(V)K}(deg^{+}(D) + deg^{-}(D) + 1)/\min\{m(x) \mid x \in V_{n_{k}}\} + 1\right) \\ \times \sum_{x \in S_{n_{k}}} \sum_{y \in N(x) \cap S_{n_{k}+1}} C_{x,y} \\ \leq \rho_{n_{k}}m(S_{n_{k}}) \times 4\left((\sqrt{m(V)K}(deg^{+}(D) + deg^{-}(D) + 1)/\min\{m(x) \mid x \in V_{n_{k}}\} + 1\right).$$

Since our estimate on  $\Delta_j f_j^{(n)}$  obtained in (iii) of Lemma 4.2 is valid for  $\Delta_{n_\ell} f_{n_\ell}^{(n_k)}$ , we see

$$\lim_{k \to \infty} \sup_{\ell > k} \deg^{\pm} \Delta_{n_{\ell}} f_{n_{\ell}}^{(n_k)} \mathbb{1}_{V_{n_{\ell}} \setminus V_{n_k}} = 0.$$

Hence,

$$\begin{aligned} r_{n_{\ell}}(D) \leq & r_{n_{k}}(D) + \deg^{+}(\nu_{O_{n_{\ell}}} - \nu_{O_{n_{k}}}) + \deg^{-}(\nu_{O_{n_{\ell}}} - \nu_{O_{n_{k}}}) + |i_{(G_{n_{\ell}},C_{n_{\ell}})} - i_{(G_{n_{k}},C_{n_{k}})}| \\ &+ \rho_{n_{k}}m(S_{n_{k}})4\left(\sqrt{m(V)K}(\deg^{+}(D) + \deg^{-}(D) + 1)/\min\{m(x) \mid x \in V_{n_{k}}\} + 1\right) \\ &+ o(1). \end{aligned}$$

Combining this and (17) with  $\deg^+(\nu_{O_{n_{\ell}}} - \nu_{O_{n_k}}) + \deg^-(\nu_{O_{n_{\ell}}} - \nu_{O_{n_k}}) < \varepsilon/2^k$  as obtained in (16) and  $|i_{(G_{n_{\ell}},C_{n_{\ell}})} - i_{(G_{n_k},C_{n_k})}| \to 0$  (as  $k, \ell \to \infty$ ), it turns out that  $r_{n_{\ell}}(D) \leq r_{n_k}(D) + \varepsilon/2^k + o(1)$  as  $\ell \to \infty$  for any k, which implies  $\limsup_{\ell \to \infty} r_{n_{\ell}}(D) \leq \liminf_{k \to \infty} r_{n_k}(D)$ , in other words,  $r_{n_k}(D)$  converges as  $k \to \infty$ .

The limit in this proposition depends on the choice of the sequence  $\{O_{n_k}\}$  of total orders. However, as long as a divisor D is supported by a finite graph, we can define

$$r_{\{O_{n_k}\}}(D) = \lim_{k \to \infty} r_{n_k}(D)$$

by taking a subsequence  $\{O_{n_k}\}$  of total orders as in Lemma 4.3.

**Remark 4.5.** If one takes another base vertex  $v'_0$  satisfying the condition (17) in Proposition 4.4, instead of  $v_0$ , then  $r'_{n_{k'}}(D)$  is defined as the same divisor D in the proposition. By taking a similar procedure in the proof of the proposition, one sees not only  $\limsup_{\ell'\to\infty} r'_{n_{\ell'}}(D) \leq \liminf_{k\to\infty} r_{n_k}(D)$  but also  $\limsup_{\ell\to\infty} r_{n_\ell}(D) \leq \liminf_{k\to\infty} r_{n_k}(D)$  does not depend on the choice of the base vertex satisfying (17).

For a divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  on G, we introduce effective divisors  $D^+$ and  $D^-$  given respectively by  $D^+ = \sum_{x \in V_G, \ell(x) > 0} \ell(x)i(x)1_{\{x\}}$  and  $D^- = -\sum_{x \in V_G, \ell(x) < 0} \ell(x)i(x)1_{\{x\}}$ . We recall that its restriction  $\sum_{x \in V_{n-1}} \ell(x)i(x)1_{\{x\}}$  to  $V_n$  is denoted by  $(D)_n$ .

**Cororally 4.6.** For any divisor  $D = \sum_{x \in V_G} \ell(x)i(x)1_{\{x\}}$  on G satisfying  $\sum_{x \in V_G} |\ell(x)|$  $i(x) < \infty$ ,  $\{r_{\{O_{n_k}\}}((D)_{n_l})\}$  is a Cauchy sequence, where  $\{(D)_{n_l}\}$  stands for a sequence of divisors in Lemma 4.3.

*Proof.* By applying Lemma 2.11 to effective divisors  $((D)_{n_{l'}} - (D)_{n_l})^{\pm}$  with  $n_{l'} > n_l$ , we have

$$r_{n_k}((D)_{n_l}) \le r_{n_k}((D)_{n_l} + ((D)_{n_{l'}} - (D)_{n_l})^+) \le r_{n_k}((D)_{n_l}) + \deg^+((D)_{n_{l'}} - (D)_{n_l})$$

and

$$r_{n_k}((D)_{n_l}) - \deg^-((D)_{n_{l'}} - (D)_{n_l}) \le r_{n_k}((D)_{n_{l'}}) \le r_{n_k}((D)_{n_l}) + \deg^+((D)_{n_{l'}} - (D)_{n_l}).$$

Thanks to Proposition 4.4, by passing the limit as  $k \to \infty$ , we have

$$|r_{\{O_{n_k}\}}((D)_{n_l}) - r_{\{O_{n_k}\}}((D)_{n_{l'}})| \le \deg^+((D)_{n_{l'}} - (D)_{n_l}) + \deg^-((D)_{n_{l'}} - (D)_{n_l}).$$

Thanks to the finiteness  $\sum_{x \in V_G} |\ell(x)| i(x) < \infty$ , for any  $\varepsilon > 0$ , there exists a positive integer  $n_0(\varepsilon)$  such that  $\sum_{x \notin V_{n_0(\varepsilon)}} |\ell(x)| i(x) = \deg^+(D - (D)_{n_0(\varepsilon)}) + \deg^-(D - (D)_{n_0(\varepsilon)}) < \varepsilon$ . Accordingly, it turns out that

$$l'>l\geq n_0(\varepsilon)\Rightarrow |r_{\{O_{n_k}\}}((D)_{n_l})-r_{\{O_{n_k}\}}((D)_{n_{l'}}))|<\varepsilon$$

independently of the choice of the sequence  $\{O_{n_k}\}$ .

This implies the convergence of the sequence  $\{r_{\{O_{n_k}\}}((D)_{n_l})\}_{l=1}^{\infty}$  and allows us to define  $r(D) = \inf_{\{O_{n_k}\}} \lim_{l\to\infty} r_{\{O_{n_k}\}}((D)_{n_l})$  for any divisor D satisfying  $\sum_{x\in V_G} |\ell(x)|$  $i(x) < \infty$ . Thanks to the finiteness of the measure m, by the definition of the characteristic  $\mathfrak{e}_{(G_n,C_n)}$  in the second section, it is easy to see that the sequence  $\{\mathfrak{e}_{(G_n,C_n)}\}$  converges and the limit is independent of choice of the base vertex  $v_0$ .

**Theorem 4.7.** Let  $G = (V_G, E_G)$  be a locally finite connected graph of finite volume satisfying (17). For any divisor D with  $\deg^+(D) + \deg^-(D) < \infty$ , the Riemann Roch theorem holds on G:

$$r(D) - r(K_G - D) = \deg(D) + \mathfrak{e}_{(G,C)},$$

where  $\mathfrak{e}_{(G,C)} = \lim_{n \to \infty} \mathfrak{e}_{(G_n,C_n)}$ .

*Proof.* For a divisor D with  $\deg^+(D) + \deg^-(D) < \infty$ , we take a sequence  $\{(D)_{n_l}\}$  associated with D in the sense of Lemma 4.3. Riemann-Roch theorem on finite weighted graphs shows that  $k \geq l$  implies

$$r_{n_k}((D)_{n_l}) - r_{n_k}(K_{G_{n_k}} - (D)_{n_l}) = \deg((D)_{n_l}) + \mathfrak{e}_{(G_{n_k}, C_{n_k})}$$

From Corollary 2.4 and Proposition 4.4 starting with  $(D)_{n_l}$ , by letting  $k \to \infty$ , we can derive

$$r_{\{O_{n_k}\}}((D)_{n_l}) - r_{\{O_{n_k}\}}(K_G - (D)_{n_l}) = \deg((D)_{n_l}) + \mathfrak{e}_{(G,C)}$$

By passing the limit as  $l \to \infty$  and taking the infimum over all of sequences  $\{O_{n_k}\}$  of total orders in Lemma 4.3, it turns out that

$$r(D) - r(K_G - D) = \deg(D) + \mathfrak{e}_{(G,C)}.$$

# References

- S. Backman, Riemann-Roch theory for graph orientations, Adv. Math. 309 (2017), 655-691.
- [2] [2] M. Baker and S. Norine, Riemann-Roch and Abel-Jacobi Theory on a Finite Graph, Adv. Math. 215 (2007), 766-788.
- [3] [3] M. Baker and F. Shokrieh, Chipring games, potential theory on graphs, and spanning trees, J. Combin. Theory 120 (2013), 164-182.
- [4] [4] F. R. K. Chung, Spectral Graph Theory, CBMS Regional Conference Series in Mathematics, 92, American Mathematical Society (1997).
- [5] [5] S. Corry and D. Perkinson, Divisors and sandpiles An introduction to chip-firing, American Mathematical Society (2018).
- [6] [6] M. Fukushima, Y. Oshima, and M. Takeda, Dirichlet forms and symmetric Markov processes, 2nd Edition, Walter de Gruyter (2010).
- [7] [7] S. Friedland and R. Nabben, On Cheeger-type inequalities for weighted graphs, J Graph Theory 41 (2002), 1-17.
- [8] [8] A. Gathmann and M. Kerber, A Riemann-Roch theorem in tropical geometry Math. Z., 259(1) (2008), 217-230.
- [9] [9] R. James and R. Miranda, A Riemann-Roch theorem for edge-weighted graphs, Proc. Amer. Math. Soc. 141 (2013), 3793-3802.
- [10] [10] J. Kigami, Analysis on fractals, Cambridge Tracts in Mathematics, Cambridge University Press (2001).
- [11] [11] W. Wang, Functional Inequalities, Markov Semigroups and Spectral Theory, Elsevier (2005).
- [12] [12] W. Woess, Denumerable Markov Chains, European Math. Soc. (2009).