# k-Pareto Optimality-Based Sorting with Maximization of Choice

Jean Ruppert\* Marharyta Aleksandrova\*\* Thomas Engel\*\*
\*Mathematics and Computing S.à.r.l., first.last@mathcomp.lu \*\*University of Luxembourg, first.last@uni.lu

# Abstract

Topological sorting<sup>1</sup> is an important technique in numerous practical applications, such as information retrieval, recommender systems, optimization, etc. In this paper, we introduce a problem of generalized topological sorting with maximization of choice, that is, of choosing a subset of items of a predefined size that contains the maximum number of equally preferable options (items) with respect to a dominance relation. We formulate this problem in a very abstract form and prove that sorting by k-Pareto optimality yields a valid solution. Next, we show that the proposed theory can be useful in practice. We apply it during the selection step of genetic optimization and demonstrate that the resulting algorithm outperforms existing state-of-the-art approaches such as NSGA-II and NSGA-III. We also demonstrate that the provided general formulation allows discovering interesting relationships and applying the developed theory to different applications.

# 1 Introduction

In the modern era of information overload, the task of choosing a subset of the most useful items is extremely important. Various tools were developed with the aim to assist a user with this task, for example, text search engines [Croft et al., 2010] and recommender systems [Resnick and Varian, 1997]. In most of the cases, such systems suggest to the user a small set of elements<sup>2</sup>.

Thereby, if the number of equally preferable options is large, a heuristic is used to discard a fraction of them. However, in some applications the user might be willing to analyze all equally preferable options with the aim to choose the best one. This can happen, for example, in the case of choosing a habitation.

A similar problem of *choosing a subset of most preferable elements* also arises as an important step when solving various practical tasks. A straightforward example would be the selection step in genetic optimization algorithms [Mitchell, 1998]. At this step, a subset of the current population is chosen to advance to the next generation. Having the chosen subset made up of elements with large fitness values guides the evolution process in the desired direction. At the same time, selecting a subset with the largest variety of genes ensures variability of characteristics and allows faster exploration of the search space.

These examples bring us to the problem of generalized<sup>3</sup> topological sorting with choice maximisation which we also refer to as maximum choice problem. This problem aims to choose a subset of a predefined maximum size, consisting of most preferable items and containing the largest number of equally preferred elements<sup>4</sup>. To the best of our knowledge, this problems has not yet been studied in the literature. In this text, we propose a theoretical solution to the maximum choice problem and demonstrate how both the problem and its solution can be applied in practice.

The contributions of this work are the following:

- 1. We formulate the maximum choice problem in a broad sense for arbitrary elements, preference relations R, and measures  $\mu$  indicating the set sizes, see Section 2.
- 2. We propose a solution based on the concept of k-Pareto optimality, whose definition relies on the relation R, see Section 3.

<sup>&</sup>lt;sup>1</sup> Topological sorting [E.Knuth, 1997] here means the process of sorting a set of items with respect to a preference or dominance relation. We use the terms *preference* and *dominance* interchangeably.

<sup>&</sup>lt;sup> $^{2}$ </sup>We use the terms *item* and *element* interchangeably.

<sup>&</sup>lt;sup>3</sup>Later we show that the formulated problem is a generalization of topological sorting. In the case of a partial order relation, standard topological sorting is considered.

 $<sup>{}^{4}</sup>$ If items are equally preferable, then we say that they offer choice for the user or the system.

- 3. We further investigate the proposed solution from a theoretical point of view and discover interesting characteristics, such as the relationship between k-Pareto optimal elements and the arc of hyperbola, see Section 4.
- 4. Finally, we demonstrate the applicability of our approach to real-world problems by considering genetic optimization, see Section 5.

# 2 Sorting with Choice Maximization

To formally define the problem of maximization of choice, we introduce several definitions in Sections 2.1 and 2.2. The resulting formalization of this problem is abstract and quite general. However, this generality allows discovering novel connections and applying the developed theory to numerous practical problems. We also illustrate the defined concepts with an example in Section 2.3.

# 2.1 Definitions

We consider a set X with a binary relation R. Intuitively xRy means that x is preferable to y. The case xRy and not yRx means that x is strictly preferable to y. This situation is denoted by  $xR^*y$ . We also consider a positive and  $\sigma$ -finite measure  $\mu$  [Halmos, 2013 defined on X. Thus, we have a measure space  $(X, \Sigma, \mu)$ , where  $\Sigma$  is a set of subsets of X, and  $\mu$  intuitively indicates the size of these subsets. To ensure measurability, throughout this text the characteristic function  $1_R$  of the relation R is assumed to be sufficiently regular<sup>5</sup>. The  $\mu$  can be defined in different ways. Important examples are the counting measure and probability measures P. Depending on the definition of  $\mu$ , it can indicate the following characteristics of the elements in X: how many?, how likely?, how important?, or what volume?

To illustrate these definitions, we consider the following example. Let X be a set of possible habitations of which the user has to choose the best according to his preferences encoded by the relation R. In such a situation, the relation R can be multidimensional. Let us assume, for simplicity, that an optimal habitation for the user is close to a given location, for example, his workplace (relation  $R_l$ ), is situated in a district with a smaller population size (relation  $R_p$ ), and is close to a river (relation  $R_r$ ). Thus, the user's preferences can be represented by the preorder relation  $R = R_l \& R_p \& R_r$ . In our example, all available habitations from X can be mapped onto points in a 3dimensional space of *Proximity to the location* × *Population* × *River*. The fact that R is a preorder relation means that some elements of X can be comparable, while others not. For example, the habitation xwith coordinates (50, 100, True) is strictly preferable to y with coordinates (60, 100, True), that is  $xR^*y$ . At the same time, the habitation z with coordinates (40, 100, False) is incomparable with x. Indeed, z is better with respect to  $R_l$ , it is situated closer to the required location, but x is better with respect to  $R_r$ , as the latter is situated near a river.

Having the task to find a subset of X that is 'best' according to R, a rational solution can be formulated with the following recursive expression: if an element x is selected, then all elements that are strictly preferable to x should be also selected. In our example, this translates into the task of finding a subset of habitations  $S_R$  that might be suitable for the user. Naturally, if  $y \in S_R$ , then  $x \in S_R$  as the latter corresponds better to the preferences of the user defined by the relation R. We formalize this rationality condition by defining selections as follows.

**Definition 1.** A selection S is a subset of X such that  $x \in S$  and  $yR^*x$  implies  $y \in S$ . The set of all selections in  $\Sigma$  is denoted by  $S^6$ .

# 2.2 The Maximum Choice Problem

As discussed in Section 1, in practical applications when selecting a subset of X one might want not only to respect the above rationality constraint, but also to maximize the number of incomparable pairs. The latter condition is equivalent to the maximization of the diversity of the selected subset, or the maximization of the provided choice. In our example with habitations, if both x and z are presented to the user, then he can choose an appropriate habitation by himself<sup>7</sup>.

In terms of our notations, this will be translated into the condition of selecting as many pairs x, y such that neither x is strictly preferable to y ( $\neg xR^*y$ ) nor y is strictly preferable to x ( $\neg yR^*x$ ). This means that there is freedom of choice between x and y (xRy = yRx). This motivates the following quantitative definition of choice for measurable subsets of X.

**Definition 2.** Choice offered by a set A is the number  $\operatorname{cho}(A) = (\mu \times \mu)(\{(x, y) \in A^2 | xRy = yRx\}).$ 

The choice offered by a measurable set A essentially measures how many pairs of items offering choice can be extracted from A. Additionally, if one wants to restrict the size of the selected subset, in our example, to present to the user a small set of suitable habitations

<sup>&</sup>lt;sup>6</sup>If R is a partial order relation, then the selections are the down-sets [Davey and Priestley, 2002].

<sup>&</sup>lt;sup>7</sup>We consider the case when addition preferences cannot be encoded and the user has to make the final choice.

 $<sup>{}^{5}1</sup>_{R}$  is equal to 1 if xRy and is equal to 0 otherwise.

with  $\mu(S_R) \leq m$ , then this leads us to the definition of the maximum choice problem:

**Maximum Choice Problem.** For a given m find all selections T such that  $\operatorname{cho}(T) = \max_{S \in S, \mu(S) \leq m} \operatorname{cho}(S)$ . Any such selection T will be said to offer maximum choice for m.

In practical applications, it might be more insightful to consider the concept of *diversity* that is functionally related to the concept of choice.

**Definition 3.** For any measurable set A, the diversity of A is the ratio  $\operatorname{div}(A) = \operatorname{cho}(A)/\mu(A)^2$ .

Thus  $\operatorname{div}(A)$  is the likelihood that there is choice between the two randomly chosen elements. The main ingredient of our solution to the maximum choice problem is the following concept.

**Definition 4.** The k-Pareto optimality of an element  $x \in X$  is the measure of the subset of X containing all elements strictly preferable to x:  $po(x) = \mu(\{y|yR^*x\})$ .

If  $\mu$  is the probability measure, the k-Pareto optimality of an element x, po(x), is the likelihood an element drawn at random from X is strictly preferable to x.

Finally, we introduce the sets of at least k-Pareto optimal elements. Such sets consist of all elements in X with  $po \le k$ . In Section 3 we show that this concept yields a solution for the maximum choice problem.

**Definition 5.** The at least k-Pareto optimal elements  $T_k$  form the measurable set defined as follows:  $T_k = \{x \in X | \operatorname{po}(x) \le k\}^8$ .

## 2.3 Example

In this subsection, we discuss an illustrative example to demonstrate the concepts defined in Sections 2.1 and 2.2. Let us consider a finite subset X of  $\mathbb{R}^2$ , counting measure  $\mu$ , and the relation  $R_{\perp}$  defined as follows:  $(x_1, x_2)R_{\perp}(y_1, y_2)$  iff  $x_1 \leq y_1$  and  $x_2 \leq y_2$ , see Fig. 1. In economics, x is Pareto optimal if there is no y in X such that  $xR_{\perp}^*y$ . In our language, this means that po(x) = 0. Thus, k-Pareto optimality indicates how much an element is away from being Pareto optimal.

Let X be comprised of six points presented in Fig. 2. As we are considering the counting measure,  $\mu(X) = 6$ . Points A, B, and C are not dominated by any other point. This means that  $po(A) = \mu(\{y|yR^*A\}) = 0$  and po(B) = po(C) = po(A) = 0. Point E is dominated by a single point C, that is  $po(E) = \mu(\{C\}) = 1$ . Finally, points F and D are dominated by two other points each, resulting in  $po(F) = \mu(\{C, E\}) = 2$  and  $po(D) = \mu(\{A, B\}) = 2$ .





Figure 1: Illustration of the relation  $R_{\perp}$ .



Figure 2: Computation of Pareto-optimality. Example of a finite subset of  $\mathbb{R}^2$ , the counting measure, and  $R_{\perp}$ .

Sorting the set X by k-Pareto optimality of its elements will produce the following result:  $(\{A, B, C\}, \{E\}, \{D, F\}).$ This sorting is different from sorting by Pareto fronts. The latter approach is widely used in practice and is the basis of all Pareto dominance-based genetic optimization algorithms [Li et al., 2015]. Sorting by Pareto fronts is done in the following way. First, the first Pareto front, which is the set of non-dominated points, is identified. Next, the points from this front are removed from the consideration and the process is repeated until all points are assigned to a front. Sorting the points from Fig. 2 by Pareto fronts will produce the following result:  $(\{A, B, C\}, \{D, E\}, \{F\})$ . Note, that the point D moved from the 3d equivalence class when sorting by k-Pareto optimality to the 2d when sorting by Pareto fronts. Let us consider the two selections of size 4:  $S_E = \{A, B, C, E\}$  and  $S_D = \{A, B, C, D\}$ . Sorting by Pareto fronts does not distinguish between these two selections as both E and D belong to the same equivalence class. At the same time, sorting by k-Pareto optimality has a larger preference towards  $S_E$ . Also, the latter selection contains more incomparable pairs of elements and thus offers more choice. Indeed, D is incomparable with only one point C, but E is incomparable with two points: A and B.

# 3 The Maximum Choice Theorem

The main result of this text states that at least k-Pareto optimal elements  $(T_k)$  are the largest measurable selections offering maximum choice for their respective measures. **Theorem 1.** A set of at least k-Pareto optimal elements  $T_k$  offers maximum choice for  $\mu(T_k)$  if it is a selection and if  $\mu(T_k) < +\infty$ . Moreover  $T_k$  is the largest selection offering maximum choice for  $\mu(T_k)$  in the sense that it contains any other selection offering maximum choice for  $\mu(T_k)$ .

The second part of the above theorem precisely means that if a selection A offers maximum choice for  $\mu(T_k) < +\infty$ , then  $A \subseteq T_k$  almost-everywhere. One may note that the above theorem does not state that the at least k-Pareto optimal elements are the only largest selections offering maximum choice. Later, we give an example of a fundamentally different selection offering maximum choice for a value of m where there is no such k that  $\mu(T_k) = m$ .

We prove the above stated theorem in several steps. First, we show that for selections the computation of choice can be simplified. It only requires to compute a simple integral instead of a double integral.

**Integral Formula.** If S is a measurable selection and  $\mu(S) < +\infty$ , then

$$\operatorname{cho}(S) = \int_{S} (\mu(S) - 2\operatorname{po}(x))d\mu(x).$$
(1)

*Proof.* The fact that set S is a selection means that  $\forall y \in S : \{x \in S | xR^*y\} = \{x \in X | xR^*y\}$ . That is, any element x from X strictly preferable to any element y in S, also belong to  $S \ (x \in S)$ . Using the definition of choice from Def. 2 and  $\mu(S) < +\infty$  we obtain

$$\begin{split} \mathrm{cho}(S) &= \mu(S)^2 - 2(\mu \times \mu)(\{(x,y) \in S^2 | xR^*y\}) \\ &= \mu(S)^2 - 2(\mu \times \mu)(\{(x,y) \in S \times X | yR^*x\}). \end{split}$$

Fubini's theorem [Halmos, 2013] indicates that

$$\begin{aligned} (\mu \times \mu)(\{(x,y) \in S \times X | yR^*x\} &= \\ &= \int_S \left( \int_X \mathbf{1}_{R^*} d(\mu(y)) \right) d\mu(x) = \int_S \mathrm{po}(x) d\mu(x), \end{aligned}$$

where  $1_{R^*}$  is the characteristic function of  $R^{*9}$ .

Finally, the integral formula results from the fact that 
$$\mu(S)^2 - 2\int_S \operatorname{po}(x)d\mu(x) = \int_S (\mu(S) - 2\operatorname{po}(x))d\mu(x).$$

Let's now consider the function c defined on  $\Sigma$  for any A of finite measure by

$$c(A) = \int_A (\mu(A) - 2\operatorname{po}(x))d\mu(x).$$

The integral formula defined in Eq. (1) means that for any selection S, we have c(S) = cho(S). The second step of our proof of Theorem 1 is to show that  $T_k$  is the largest measurable set that maximizes c for its respective measure. More precisely, we will prove the following lemma.

**Lemma 1.** For any k such as  $\mu(T_k) < +\infty$  we have

$$c(T_k) = \max_{A \in \Sigma, \mu(A) \le \mu(T_k)} c(A).$$

Moreover, if  $\mu(A) \leq \mu(T_k)$  and  $c(A) = c(T_k)$ , then  $A \subseteq T_k$  almost-everywhere.

The context of this lemma is very similar to the knapsack problem [Martello, 1990]. In this problem, one needs to find a subset A of a finite set of items  $\{x_1, ..., x_n\}$  maximizing the total value  $\sum_{x_i \in A} v(x_i)$  under the constraint that the total weight  $\sum_{x_i \in A} w(x_i)$  of A does not exceed a predefined maximum weight  $w^*$ . In Lemma 1, the total value is c(A), the ratio of an element's value to its weight becomes  $\mu(A) - 2 \operatorname{po}(x)$ , and the weight constraint is expressed as  $\mu(A) \leq \mu(T_k)$ . The solutions given by the lemma correspond to those yielded for the knapsack problem by George Dantzig's greedy approximation algorithm [Dantzig, 1957]. This algorithm consists of ordering elements by decreasing value-to-weight ratio and then taking the N first elements. N is chosen in such a way, that taking one more element would cause excessive weight. The process of proving Lemma 1 is similar to proving that George Dantzig's solutions are optimal for their respective weights.

Proof of Lemma 1. Let's consider  $T_k$  such that  $\mu(T_k) < +\infty$ . To prove  $c(T_k)$  is the maximum, we need to show for any  $A \in \Sigma$ , that  $c(A) \leq c(T_k)$  if  $\mu(A) \leq \mu(T_k)$ . As  $A = (A \cap T_k) \cup (A \setminus T_k)$  and  $T_k = (A \cap T_k) \cup (T_k \setminus A)$ , requiring  $c(A) \leq c(T_k)$  is equivalent to requiring

$$\int_{A \setminus T_k} (\mu(A) - 2\operatorname{po}(x)) d\mu(x) \le \int_{T_k \setminus A} (\mu(T_k) - 2\operatorname{po}(x)) d\mu(x)$$
(2)

By definition of  $T_k$ , we have that po(x) > k for  $x \in A \setminus T_k$ , while  $po(x) \le k$  for  $x \in T_k \setminus A$ . Therefore,

$$\int_{A \setminus T_k} (\mu(A) - 2\operatorname{po}(x))d\mu(x) \le \mu(A \setminus T_k)(\mu(A) - 2k),$$
(3)

and

$$\int_{T_k \setminus A} (\mu(T_k) - 2\operatorname{po}(x))d\mu(x) \ge \mu(T_k \setminus A)(\mu(T_k) - 2k).$$
(4)

The constraint  $\mu(A) \leq \mu(T_k)$  means that  $\mu(A \setminus T_k) \leq \mu(T_k \setminus A)$ . Thus, we have

$$\mu(A \setminus T_k)(\mu(A) - 2k) \le \mu(T_k \setminus A)(\mu(T_k) - 2k).$$
(5)

 $<sup>{}^{9}1</sup>_{R^*}$  is defined on  $X^2$  in a similar way to  $1_R$ : it is equal to 1 if  $xR^*y$  and is equal to 0 otherwise.

Finally, combining Eq. (5) with Eq. (3) and Eq. (4) guarantees the inequality in Eq. (2) under the constraint  $\mu(A) \leq \mu(T_k)$ .

Let us now consider  $A \in \Sigma$  such that  $\mu(A) \leq \mu(T_k)$ and  $c(A) = c(T_k)$ . We now proceed to show that  $A \subseteq T_k$  almost everywhere. If  $c(A) = c(T_k)$ , the inequality in Eq. (2) must be an equality. Under the assumption  $\mu(A) \leq \mu(T_k)$ , we again have the inequalities in Eq. (3), Eq. (4) and Eq. (5), which must be equalities if Eq. (2) is an equality. However, because po(x) > k for  $x \in A \setminus T_k$ , the inequality in Eq. (3) can only become an equality if  $\mu(A \setminus T_k) = 0$ .  $\Box$ 

Now everything is in place to prove the theorem.

Proof of Theorem 1. Let us first prove that the at least k-Pareto optimal elements  $T_k$  offer maximum choice. Lemma 1 says that on  $\Sigma$  the set  $T_k$  maximises c for its respective measure. We assumed  $T_k$  is a selection. At the same time, for any selection c = cho. It means that  $T_k$  offers maximum choice for its respective measure. Moreover, from Lemma 1 and from  $S \subseteq \Sigma$ directly results that if a selection A offers maximum choice for  $\mu(T_k)$ , then  $A \subseteq T_k$  almost everywhere.  $\Box$ 

The theorem does not guarantee uniqueness. For example, in the case of the relation  $\leq$  on  $\mathbb{R}$  and the Lebesgue measure, selections are all left-unbounded intervals and the choice of any selection is 0. However, if for any selection

$$A \subset T_k \& \mu(A) < \mu(T_k) \implies \operatorname{cho}(A) < \operatorname{cho}(T_k), (6)$$

then  $T_k$  is a unique maximum. This is a direct consequence from the fact that  $T_k$  contains any other selection offering maximum choice for  $\mu(T_k) < +\infty$ .

Also, the proof of the theorem indicates that transitivity is not necessary. The set  $T_k$  is only required to be a selection. This is the case for Lebesgue area measure and the non-transitive relation  $R_{L'}$  defined on the unit square by  $(x_1, x_2)R_{L'}(y_1, y_2)$  iff  $\frac{y_1}{2} \le x_1 \le y_1$  and  $\frac{y_2}{2} \le x_2 \le y_2$ . The sets  $T_k$  fulfill the requirement of being selections if  $R^*$  satisfies the weakened transitivity condition  $xR^*y \implies po(x) \le po(y)$ .

It is possible to prove that for the counting measure, finite sets, and partial order relations, all selections offering maximum choice can be obtained when sorting by k-Pareto optimality and taking the first l elements. However, for discrete measures with nonconstant weights the above construction might not yield all selections offering maximum choice. A relative example is given in Fig. 3. The partial order relation represented by its Hasse diagram. [Davey and Priestley, 2002]. The set S is not of the form  $T_k^* \cup A$ 



Figure 3: S is a selection that offers maximum choice but cannot be constructed from  $T_k$ . Point are encoded with their measure and k-Pareto optimality:  $(\mu, po)$ .

with  $A \subseteq T_k \setminus T_k^*$ . Nevertheless, it offers maximum choice for m = 3.

# 4 Further Theoretical Explorations

# 4.1 Efficient Computation of Choice

Computation of choice may be simplified by performing the change of variable y = po(x) in the integral formula (1).

**Proposition 1.** For any selection of at least k-Pareto optimal elements  $T_k$  such that  $\mu(T_k) < +\infty$ 

$$\operatorname{cho}(T_k) = \mu(T_k)^2 - 2 \int_{[0,k]} x d(\operatorname{po} * \mu)(x), \quad (7)$$

where  $po *\mu$  is the image measure defined by  $(po *\mu)([a, b]) = \mu(po^{-1}([a, b])).$ 

# 4.2 Characteristics of Random Vectors

Let us study a probability space  $(\Omega, \Sigma, P)$ . We consider two random variables  $X_1$  and  $X_2$ , as well as the partial order relation  $R_{\Omega_{\perp}}$  defined on  $\Omega$  as follows:

$$\omega_x R_{\Omega \sqcup} \omega_y$$
 iff  $X_1(\omega_x) \le X_1(\omega_y)$  and  $X_2(\omega_x) \le X_2(\omega_y)$ .

For convenience, we use the following notations interchangeably:  $(X_1(\omega_x), X_2(\omega_x)), (x_1, x_2)$  or simply x.

In this case, the selections are the sets situated below any decreasing curve, like for example the hyperbola  $x_1x_2 = 1/5$  and the curve defined by  $max(x_1, x_2) =$ 2/5 in Fig. 4. The k-Pareto optimality  $po((x_1, x_2))$  is the joint cumulative probability distribution function  $F(x_1, x_2)$ . Finally, the at least k-Pareto optimal elements are situated below the curve  $F(x_1, x_2) = k$ . Having choice between x with coordinates  $(x_1, x_2)$ and y with coordinates  $(y_1, y_2)$  means the rectangles  $((0, 0), (x_1, 0), x, (0, x_2))$  and  $((0, 0), (y_1, 0), y, (0, y_2))$ are not nested, as depicted in Fig. 4a.

### 4.2.1 Continuous Independent Variables

If  $X_1$  and  $X_2$  are continuous, then the function po can be considered to be a linear extension of  $R_{\Omega \sqcup}$ . Indeed, if x and y are chosen independently and at random





Figure 4: Characterization of hyperbola.

from  $\Omega$ , then P(po(x) = po(y)) = 0. On the other hand, it can be shown that  $\text{div}(\Omega) = 1/2$ . This means that in half of the cases the relation  $R_{\Omega_{\perp}}$  cannot tell which element is preferable out of two elements extracted independently and at random from  $\Omega$ .

Moreover if  $X_1$  and  $X_2$  are independent, then the condition for uniqueness from Eq. (6) holds. In the special case where  $X_1$  and  $X_2$  are uniformly distributed on the interval [0, 1], the joint cumulative probability distribution function takes form of  $F(x_1, x_2) = x_1 x_2$ and the at least k-Pareto optimal elements are situated below a hyperbola  $x_1 x_2 = k$ . Combining this fact with Corollary 1 yields a surprising characterization of hyperbola, see Fig. 4 and the statement below.

**Characterization of Hyperbolas.** Out of all descending functions f from [0,1] to [0,1] delimiting an area  $\int_0^1 f(x)dx = c$ , the arc of hyperbola is the one offering the highest likelihood the rectangles  $((0,0), (x_1,0), x, (0,x_2))$  and  $((0,0), (y_1,0), y, (0,y_2))$ are not nested for two points x and y being drawn independently and at random from the delimited area.

# 4.2.2 Independence on Marginal Distribution

Sklar's theorem [Sklar, 1959, Durante et al., 2013] states that the cumulative distribution function  $F(x_1, x_2)$  can be represented as  $C(F_1(x_1), F_2(x_2))$  for a copula C. Marginas of  $X_1$  and  $X_2$  are fully described by the marginal cumulative probability distributions  $F_1$  and  $F_2$ , whereas the copula describes the dependence structure between  $X_1$  and  $X_2$ . The copula can be considered as a joint cumulative distribution function having two uniform marginal distributions on [0, 1]. Below we show that the introduced concepts do not depend on the marginal distribution of  $X_1$  and  $X_2$ .

**Proposition 2.** For a continuous random vector  $(X_1, X_2)$  and the relation  $R_{\Omega_{\perp}}$ , k-Pareto optimality, choice, diversity and selections offering maximum choice only depend on the copula C of  $X_1, X_2$ .

*Proof.* Let us consider the mapping

$$\begin{array}{rcccc} G & : & \Omega & \rightarrow & [0,1]^2, \\ & & \omega_x & \mapsto & (F_1(x_1),F_2(x_2)). \end{array}$$

We consider  $R_{\Omega_{\perp}}$  and P defined on  $\Omega$ . At the same time, on  $[0,1]^2$  we consider  $R_{\perp}$  defined by  $(x_1, x_2)R_{\perp}(y_1, y_2)$  iff  $x_1 \leq x_2$  and  $y_1 \leq y_2$ , as well as the image measure G \* P defined on  $[0,1]^2$  by  $(G * P)(A) = P(G^{-1})(A)$ . The map G preserves probabilities in the sense that for any measurable A in  $\Omega$  we have (G \* P)(G(A)) = P(A). Moreover, G preserves the relations in the sense that  $xR_{\Omega_{\perp}}y$  iff  $G(x)R_{\perp}G(y)$ . Selections are preserved in the sense that if S is a selection for  $R_{\Omega_{\perp}}$ , then G(S) is a selection for  $R_{\perp}$ . Ignoring negligible subsets, this mapping between selections is one-to-one. Therefore, G also preserves selections, k-Pareto optimality, choice, and diversity.

The proposition finally results from the fact that G \* Ponly depends on the copula. This is a consequence of the fact that for any  $(a_1, a_2) \in [0, 1]^2$ , we have  $(G * P)([0, a_1] \times [0, a_2]) = C(a_1, a_2)$ . This equality is a result of the following statements: 1) continuity which guarantees that  $a_1$  and  $a_2$  can be written as  $F_1(x_1)$  and  $F_2(x_2)$  for some appropriate  $x_1$  and  $x_2$ ; 2) the definition of image measure; 3) the fact that  $G = (F_1, F_2) \circ (X_1, X_2)$ ; and 4) the equality  $F(x_1, x_2) = C(F_1(x_1), F_2(x_2))$ .

**Proposition 3.** If  $X_1$  and  $X_2$  are two continuous independent random variables, then for the relation  $R_{\Omega_{\perp}}$ 

$$P(T_k) = k - k \ln(k),$$
  

$$cho(T_k) = (k - k \ln(k))^2 - k^2 \left(\frac{1}{2} - \ln k\right)$$

*Proof.* Let us again consider the map G. The image measure G \* P induced by G on  $[0,1]^2$  is the Lesbegue area measure. Independently of P, we have  $G(T_k) = \{(x_1, x_2) \in [0,1]^2 | x_1 x_2 \leq k\}$ . Simple integration for Eq. (7) yields

$$P(\mathrm{po}^{-1}(]-\infty,x])) = P(T_x),$$
  
=  $\int_{(x_1,x_2)\in[0,1]^2} (G(T_x)) = x - x\ln(x).$ 

Therefore,  $po * P = -\ln(x)dx$  and Eq. (7) results in

$$cho(T_k) = (k - k \ln(k))^2 - 2 \int_0^k x(-ln(x)) dx,$$
  
=  $(k - k \ln(k))^2 - k^2 (1/2 - \ln k).$ 

Now, it is possible to show that  $\lim_{k\to 0} \operatorname{div}(T_k) = 1$ . The fact that diversity slowly tends to the maximum



Figure 5: Selections of the best 40% according to different sorting criteria.



Figure 6: Fraction of the best elements and diversity.

possible value of 1 as k tends to zero becomes even more surprising if one looks at the selections

$$T_{\min a} = \{ (x_1, x_2) \in [0, 1]^2 | \min(x_1, x_2) \le a \},\$$
  
$$T_{\max a} = \{ (x_1, x_2) \in [0, 1]^2 | \max(x_1, x_2) \le a \}.$$

Here,  $\lim_{a\to 0} \operatorname{div}(T_{\min a}) = \frac{3}{4}$ , and  $\operatorname{div}(T_{\max a}) = \frac{1}{2}$ . To further illustrate this observation, let us consider selections of a fixed measure m, which represent the best m\*100% of elements, defined with different sorting criteria: minimum  $\min(x_1, x_2)$ , maximum  $\max(x_1, x_2)$ , average  $\frac{x_1+x_2}{2}$ , and Pareto optimality  $x_1x_2^{10}$ . As we can see in Fig. 6,  $\min(x_1, x_2)$  delimits selections containing too many large values, that is, extremes are overvalued. On the other hand, all other sorting criteria except po undervalue extremes and include too many elements situated around the diagonal  $x_1 = x_2$ .

Finally, in Fig. 6 we demonstrate how diversity of the selections defined above depends on the fraction of selected elements  $m^{11}$ . We can see that diversity is the largest when sorting by k-Pareto optimality. This is a direct consequence of Theorem 1.

# 5 Further Practical Explorations

# 5.1 Computation Complexity

A common solution for ranking n elements of a set X according to a partial order relation R is to rank the elements according to their average ranking with respect to all linear extensions of R. However, the total number of linear extensions exponentially increases with n, and the resulting algorithms are complex and slow [de Loof, 2010, p. 48]. For example, random sampling of linear extensions has an expected running time of  $O(n^3 \log n)$  [Huber, 2006]. Below we show that sorting by k-Pareto optimality offers an efficient alternative.

The basic algorithm for the k-Pareto optimality based sorting is straightforward. In the case of an arbitrary relation R, po(x) is computed by summing up the measures of the items that are strictly preferable to x. This requires one pass through the whole set X for every element  $x \in X$  with computation complexity  $O(n^2)$ . The complexity of sorting X by increasing values of po is  $O(n \log n)$ . Therefore, the total computational complexity is  $O(n^2)$ .

The case of composite relations defined on the probability space allows constructing even faster sorting procedures. We illustrate this idea for  $R = R_l \& R_p \& R_r$ from our housings example. We define the component relations as follows: for  $i \in \{l, r, p\}$ ,  $aR_i b$  iff  $X_i(a) \leq X_i(b)$ , where the real valued random variable  $X_l$  represents proximity to the location,  $X_p$  represents population size, and  $X_r(x)$  is 0 when x is close to a river and 1 otherwise. For independent  $X_i$ , we have:

$$po(x) = P(\{y|yR^*x\}),$$
  
=  $P(\{y|yRx\}) - P(\{y|yRx \text{ and } xRy\}),$   
=  $\prod_{i \in \{1,...,m\}} P(X_i \le x_i) - \prod_{i \in \{1,...,m\}} P(X_i = x_i).$ 

The cumulative probability distributions  $F_i(x) = P(X_i \leq x)$  can be approximated by the respective empirical cumulative probability distributions  $\hat{F}_i(x)$ . The computation complexity of estimating  $\hat{F}_i$  is  $O(n \log n)$ . This needs to be done for every component relation  $R_i$ , resulting in the total complexity of  $O(n \log n)$ .

# 5.2 Application to Genetic Optimization

In Section 1, we hypothesized that sorting with choice maximization can be beneficial for genetic optimization. Indeed, this strategy results in the maximization of the population diversity and allows exploring the search space more efficiently. Additionally, Pareto dominance-based many-objective<sup>12</sup> genetic optimiza-

 $<sup>^{10}</sup>$ Here, po is functionally related to geometric mean.

<sup>&</sup>lt;sup>11</sup>Note that in the considered case for any selection S,  $\operatorname{cho}(S) = dx_1 dx_2 (S)^2 - \int_S \operatorname{po}(x) dx_1 dx_2.$ 

<sup>&</sup>lt;sup>12</sup>Concerns problems with 4 and more objectives.

tion algorithms are known to suffer from the lack of selection pressure [Palakonda et al., 2018]. When the number of objectives increases, the number of incomparable solutions grows exponentially. However, as shown in Section 4.2, sorting random independent vectors by their Pareto optimality can be considered as a liner extension of the defined preference relation. The fact that P(po(x) = po(y)) = 0 means that such sorting rarely produces ties and for any two solutions either x is preferable to y or vice versa. In the rest of this subsection, we demonstrate that the proposed approach indeed improves the performance of genetic algorithms in the case of independent objectives.

To evaluate the proposed sorting procedure, we use it in NSGA-II instead of Pareto dominance-based sorting. We experiment with two measures  $\mu$ : counting and probability measures. This gives us two versions of genetic algorithms referred to as *PO-count* and PO-prob respectively. These algorithms are compared with implementations of the state-of-the-art algorithms NSGA-II and NSGA-III [Deb and Jain, 2013] from the *deap* python library<sup>13</sup>. For the experimental evaluation, we use the 0/1 knapsack problem with independent objectives as defined in [Zitzler and Thiele, 1999]. The number of knapsacks (objectives) is varied within the following set  $n_k \in \{2-8, 10, 15, 25\}$ and the number of items is set to 250. We adopt random selection with replacement and uniform crossover with mutation probability 0.01. We set the population size to 250 and the number of generations to 500. All results are the average among 30 independent runs.

Below we analyze the performance of different algorithms in terms of the classical hypervolume metric [Shang et al., 2020] with the origin of coordinates as a reference point. In our setup, this metric is to be maximized. We choose NSGA-II as the baseline, and present the relative changes in the **hypervolume** indicator for the rest of the algorithms in Fig. 7 (increase: positive number, decrease: negative number). We notice that despite having been developed for the manyobjective optimization, NSGA-III almost always results in lower values of hypervolume, even for a large number of knapsacks. This confirms a similar observation from [Ishibuchi et al., 2016], and supports our choice of NSGA-II as a baseline for implementation and comparison instead of NSGA-III. Further, we see that the value of relative increase for *PO-count* is always very close to 0. It means that *PO-count* yields a population covering the same hypervolume as NSGA-II. Contrarily, PO-prob improves the hypervolume, as compared to NSGA-II. This difference is visible for small  $n_k$  (+4% for  $n_k = 2$ ) and is especially prominent for large  $n_k$  (+60% for  $n_k = 25$ ). For  $n_k$  between



Figure 7: Increase in hypervolume compared to *NSGA-II*.

5 and 7, *PO-prob* results in lower values of hypervolume than *NSGA-II*. However, the relative decrease in thes cases does not exceed -1.63%. Also, within this range, *PO-count* performs slightly better than other algorithms. These results demonstrate that the proposed approach improves the performance of genetic algorithms, especially in the case of many-objective optimization. It also suggests that the choice of the measure  $\mu$  has a large impact on the performance. The latter relationship will be studied in future work.

# 6 Conclusion

In this paper, we formulate the problem of generalized topological sorting with choice maximization, which, to the best of our knowledge, was not considered in the literature before. We also prove that the *at least* k-Pareto optimal sets provide unique solutions. Further theoretical analysis of this problem leads us to an interesting relationship between the diversity of random points and the arc of hyperbola. Additionally, we propose a computationally efficient algorithm for the calculation of k-Pareto optimality for probability measures. Finally, we demonstrate a successful application of the developed theory. We show that sorting by k-Pareto optimality can drastically improve the performance of many-objective genetic optimization algorithms. In our experiments, the proposed solution based on the probability measure allows increasing the value of hypervolume by up to 60% for 25 objectives. This result can be considered as a potential solution to the problem of searchability deterioration in Paretodominance optimization.

We also believe that the proposed general framework can be used in different applications. In future work, we plan to study the applicability of k-Pareto optimality for constrained optimization, scheduling problems, recommender systems, and the development of statistical indicators. Maximization of choice might be also useful when studying causality and fairness.

<sup>&</sup>lt;sup>13</sup>https://deap.readthedocs.io/en/master/

### References

- [Croft et al., 2010] Croft, W. B., Metzler, D., and Strohman, T. (2010). *Search engines: Information retrieval in practice*, volume 520. Addison-Wesley Reading.
- [Dantzig, 1957] Dantzig, G. B. (1957). Discretevariable extremum problems. *Operations research*, 5(2):266–288.
- [Davey and Priestley, 2002] Davey, B. A. and Priestley, H. A. (2002). *Introduction to lattices and order*. Cambridge university press.
- [de Loof, 2010] de Loof, K. (2010). Efficient computation of rank probabilities in posets. PhD thesis, Universiteit Gent.
- [Deb and Jain, 2013] Deb, K. and Jain, H. (2013). An evolutionary many-objective optimization algorithm using reference-point-based nondominated sorting approach, part i: solving problems with box constraints. *IEEE transactions on evolutionary* computation, 18(4):577–601.
- [Deb et al., 2002] Deb, K., Pratap, A., Agarwal, S., and Meyarivan, T. (2002). A fast and elitist multiobjective genetic algorithm: Nsga-ii. *IEEE transactions on evolutionary computation*, 6(2):182–197.
- [Durante et al., 2013] Durante, F., Fernandez-Sanchez, J., and Sempi, C. (2013). A topological proof of sklar's theorem. Applied Mathematics Letters, 26(9):945–948.
- [E.Knuth, 1997] E.Knuth, D. (1997). The Art of Computer Programming, volume 1. Addison-Wesley, Massachusetts, 3 edition.
- [Halmos, 2013] Halmos, P. R. (2013). Measure theory, volume 18. Springer.
- [Huber, 2006] Huber, M. (2006). Fast perfect sampling from linear extensions. *Discrete Mathematics*, 306(4):420–428.
- [Ishibuchi et al., 2016] Ishibuchi, H., Imada, R., Setoguchi, Y., and Nojima, Y. (2016). Performance comparison of nsga-ii and nsga-iii on various manyobjective test problems. In 2016 IEEE Congress on Evolutionary Computation (CEC), pages 3045– 3052. IEEE.
- [Jones, 1979] Jones, D. S. (1979). Elementary information theory. Clarendon Press.
- [Kendall, 1948] Kendall, M. G. (1948). Rank correlation methods.

- [Kumara et al., 2020] Kumara, A., Wub, G., Alic, M., Luob, Q., Mallipeddid R., Suganthane, P. N., and Swagatam Das, S. (2020). Guidelines for real-world multi-objective constrained optimisation competition. https://raw.githubusercontent. com/P-N-Suganthan/2021-RW-MOP/main/ Revised\_Guideline\_RWCMOP.pdf.
- [Li et al., 2015] Li, B., Li, J., Tang, K., and Yao, X. (2015). Many-objective evolutionary algorithms: A survey. ACM Computing Surveys (CSUR), 48(1):1– 35.
- [Martello, 1990] Martello, S. (1990). Knapsack problems: algorithms and computer implementations. Wiley-Interscience series in discrete mathematics and optimization.
- [Mitchell, 1998] Mitchell, M. (1998). An introduction to genetic algorithms. MIT press.
- [Noman et al., 2011] Noman, N., Bollegala, D., and Iba, H. (2011). An adaptive differential evolution algorithm. In 2011 IEEE Congress of Evolutionary Computation (CEC), pages 2229–2236. IEEE.
- [Palakonda et al., 2018] Palakonda, V., Ghorbanpour, S., and Mallipeddi, R. (2018). Pareto dominance-based moea with multiple ranking methods for many-objective optimization. In 2018 IEEE Symposium Series on Computational Intelligence (SSCI), pages 958–964. IEEE.
- [Resnick and Varian, 1997] Resnick, P. and Varian, H. R. (1997). Recommender systems. Communications of the ACM, 40(3):56–59.
- [Shang et al., 2020] Shang, K., Ishibuchi, H., He, L., and Pang, L. M. (2020). A survey on the hypervolume indicator in evolutionary multiobjective optimization. *IEEE Transactions on Evolutionary Computation*, 25(1):1–20.
- [Sklar, 1959] Sklar, M. (1959). Fonctions de repartition an dimensions et leurs marges. Publ. inst. statist. univ. Paris, 8:229–231.
- [Zitzler and Thiele, 1999] Zitzler, E. and Thiele, L. (1999). Multiobjective evolutionary algorithms: a comparative case study and the strength pareto approach. *IEEE transactions on Evolutionary Computation*, 3(4):257–271.

# Supplementary Materials

# A Further Examples for Simple Relations and Measures

In the first two supplementary examples, we consider a situation typical in economics or multi-objective optimization. Later, we show how the proposed concepts apply to arbitrary transitive relations.

### A.1 Continuous Measures

Let us again consider  $R_{\perp}$  as defined in Section 2.3. The relation  $R_{\perp}$  models preference for small values of  $x_1$ and  $x_2$ . However, instead of assuming X to be a finite subset of  $\mathbb{R}^2$ , we now study the unit square  $[0,1] \times [0,1]$ with three continuous measures: the Lebesgue area measure  $dx_1 dx_2$ , as well as  $2x_2 dx_1 dx_2$  and  $4x_1 x_2 dx_1 dx_2$ . In each case, the total measure of the unit square equals to one. The Lebesgue area measure represents elements with two uniformly distributed characteristics  $x_1$  and  $x_2$ ;  $2x_2 dx_1 dx_2$  represents rarefaction of items having small values of  $x_2$ , whereas  $4x_1x_2 dx_1 dx_2$  represents rarefaction of items having small values of both  $x_1$  and  $x_2$ .

For each of the cases defined above, in Fig. 8a we show the set of at least k-Pareto optimal elements of measure 0.1, which corresponds to selecting the 10 best percent. All three sets demonstrate the qualitative behaviours expected from sets delimited by indifference curves when the corresponding rarefaction occurs. Indeed, the curve corresponding to the uniform distribution and the Lebesgue area measure  $dx_1dx_2$  is symmetric. Also, in this case,  $po(x_1, x_1) = x_1x_2$ , and the sets of at least k-Pareto optimal elements are the sets situated below arcs of hyperbola defined by the equation  $x_1x_2 = k$ , see Sections 4.2 and 4.2.1 for more details. Applying rarefaction with respect to  $x_2$  prioritises smaller values of this characteristic. This is represented by shifting upwards the right part of the hyperbola arc, see the curve for  $2x_2dx_1dx_2$ . Indeed, in this case, the small values of  $x_2$  are observed less often. This results in selecting additional elements with large values of  $x_1$  and  $x_2$  results in the fact that the small values of both characteristics are observed less often. Thus, elements with larger values of  $x_1$  and  $x_2$  should be selected to generate a selection of the required measure. It results in the shift of the hyperbola upwards following the direction of the main diagonal, see the curve for  $4x_1x_2dx_1dx_2$ .





(a) At least k-Pareto optimal elements of measure 0.1 for the relation  $R_{\perp}$  and three measures: the Lebesgue area measure  $dx_1 dx_2$ , rarefaction of  $x_2$  defined by  $2x_2 dx_1 dx_2$ , and rarefaction of  $x_1$  and  $x_2$  defined by  $4x_1 x_2 dx_1 dx_2$ .

(b) At least k-Pareto optimal elements of measure 0.25 for the cone-based relations  $R_a$  and the Lebesgue area measure. Larger values of a represent higher maximum accepted trade-offs.

Figure 8: Examples at least k-Pareto optimal elements with two continuously distributed characteristics  $x_1, x_2$ .

#### A.2**Cone-based Relations**

Let us again consider the unit square, the Lebesgue measure, and a positive constant a. However, this time the preference relation is defined as follows:  $yR_ax$  iff  $y_2 \leq x_2$  and  $x_2 - y_2 \geq a(y_1 - x_1)$ . The above relation  $R_a$  is an example of a cone-based relation illustrated in Fig. 9. This relation has the intuitive meaning of giving up  $(x_1, x_2)$  for getting  $(y_1, y_2)$  if the improvement (diminution) in the second characteristic is at least a times the trade-off (increase) in the first characteristic. In this case, selections are sets delimited by descending curves  $x_2 = f(x_1)$  such that  $-\frac{1}{a} \le \frac{df}{dx_1} \le 0$ .



Figure 9: An illustration of a cone-based relation  $R_a$ .

Let us now consider the sets of at least k-Pareto optimal elements of measure 0.25 for the three values of a:  $a = \frac{1}{10}$ ,  $a = \frac{1}{2}$ , and a = 2, see Fig. 8b. Larger values of a represent higher maximum accepted trade-offs. This is represented by the gradual degeneration of the hyperbola into a straight horizontal line when a increases. As shown in the figure, the three sets demonstrate plausible behavior. In the situation discussed in Appendix A.1, the relation  $R_{\perp}$  corresponds to the extreme case of the relation  $R_a$  with a = 0.

#### **Transitive relations** A.3

In general, it is possible to show that if  $R^*$  is transitive, then for any k, the set  $T_k$  is a selection. In particular, if R is a partial order relation,  $\mu$  is strictly positive, and X is countable, we obtain a linear extension [Dantzig, 1957 of R when sorting X by increasing values of po and sorting ties in any order. Selections are represented by downsets. The latter are obtained when topologically sorting X and taking the first n elements, for any n. If  $\mu$  is the counting measure, then po(x) is simply the number of elements that can be reached by following downwards the edges of the corresponding Hasse diagram. An example of such a relation represented by its Hasse diagram is depicted in Fig. 10.



Figure 10: An illustration of simple partial order relation. The values of po are shown by numbers, and the set of at least 2-Pareto optimal elements  $T_2$  is delimited by a curve.

#### Further Theoretical Exploration of at least k-Pareto optimal elements Β

#### **B.1** Sorting by k-Pareto Optimality versus Sorting by Pareto Fronts

To further illustrate the difference between sorting by Pareto fronts and k-Pareto optimality, we demonstrate the results of sorting the elements of a 2-dimensional grid for the relation  $R_{\perp}$  and counting measure in Fig. 11.



Figure 11: Sorting points of a grid. The equivalence classes (fronts) are represented by numbers.

The numbers on the plots represent the equivalence class (front) to which a point was assigned by the relative sorting procedure. As we can see, sorting by Pareto fronts splits the points by straight lines, see Fig. 11a. At the same time, sorting by po results in splitting by hyperbola-like curves, see Fig. 11b. We can also notice, that extreme solutions<sup>14</sup> are valued more when sorting by po. Indeed, most of the non-extreme solutions are pushed to further equivalence classes, as compared to sorting by Pareto fronts. This characteristic of po-based sorting is also clearly visible in Fig. 12. Here we present selections of  $\mu = 0.2$  for the same relation and the set X composed of a large number of points placed on a regular grid within the shaded area.



Figure 12: Selections of  $\mu = 0.2$  for the set X composed of points in the shaded area.

However, sorting by Pareto front does not always result in selections delimited by straight lines. Analysing the

 $<sup>^{14}</sup>Extreme$  solution here means that a solution is very good according to one criteria and is bad according to another.



(a) Sorting by Pareto fronts. The total number of equivalence classes (fronts) is 39.

(b) Sorting by po. The total number of equivalence classes (fronts) is 259.

Figure 13: Sorting 500 uniformly distributed points. Points in black belong to the first 10 equivalence classes (fronts). Note, that the total number of equivalence classes is larger for po-based sorting. The latter approach results in fewer ties.

results of sorting for uniformly distributed points, we observe that both sorting methods result in hyperbola-like selections, see Fig. 13. This means that sorting by Pareto fronts is more sensitive to the topological structure of the analyzed space, while sorting by po preserves its characteristics.

# B.2 Further Solutions of the Maximum Choice Problem

# Further similar solutions

It is possible to prove that Theorem 1 also holds for  $T_k^*$  defined with a strict inequality (<) as follows, see Def. 5 for comparison.

$$T_k^* = \{ x \in X | \operatorname{po}(x) < k \}.$$

In this case, the proof of the fact that  $T_k^*$  is the largest in Lemma 1 requires analysis of inequality (4) instead of inequality (3). However, the proof of the fact that any selection T such that  $T_k^* \subseteq T \subseteq T_k$  offers maximum choice for  $\mu(T)$  becomes a bit more technical. Moreover, it is possible to prove that T is the largest selection of this kind. Precisely, for any other selection S offering maximum choice for  $\mu(T)$ ,  $S \subseteq T'$  for some T' such that  $\mu(T') = \mu(T)$  and  $T_k^* \subseteq T' \subseteq T_k$ .

### **Completeness of the Solutions**

In the case condition in Eq. (6) holds, and if for any selection S there is a k and there are selections T such that  $T_k^* \subseteq T \subseteq T_k$  and  $\mu(T) = \mu(S)$ , then those selections T are the only selections offering maximum choice. Therefore, we have a complete list of selections offering maximum choice. This is the case for the typical example of the relation  $R_{\perp}$  defined in Section 2.3 and the Lebesgue area measure defined on the unit square  $[0, 1]^2$ . This also holds for any discrete measure with constant non-zero weights, for example, for the counting measure on a finite set with a partial order relation. In the latter case, topologically sorting by increasing values of po and then taking the first n elements results in a set offering maximum choice.

### Existence of Solutions of a Different Nature

Let us consider again the example in Fig. 3. Here, the set S is not of the form  $T_k^* \cup A$  with  $A \subseteq T_k \setminus T_k^*$ . Nevertheless, it offers maximum choice for m = 3.

#### **B.3** Diversity

As it was discussed in Section 2.2, the concept of *maximum choice* is functionally related to the concept of diversity, see Def. 3. In Theorem 1 we cannot simply replace choice by diversity. However, by considering only selections of a fixed measure, we obtain the following straightforward corollary of Theorem 1.

**Corollary 1.** If R is a transitive relation, then for any set of at least k-Pareto optimal elements  $T_k$  such that  $\mu(T_k) < +\infty$ , we have

$$\operatorname{div}(T_k) = \max_{S \in \mathcal{S}, \mu(S) = \mu(T_k)} \operatorname{div}(S).$$

Moreover  $T_k$  is the unique such maximum. Precisely if S is a selection such that  $\mu(S) = \mu(T_k)$ , and  $\operatorname{div}(S) =$  $\operatorname{div}(T_k)$ , then  $S = T_k$  almost everywhere.

#### С Further Practical Exploration of at least k-Pareto optimal elements

#### **C.1** Additional Results for Genetic Optimization

In Section 5.2, we evaluate the performance of the genetic algorithms using the hypervolume indicator. In this section, we further analyze the behavior of both the state-of-the-art and the proposed algorithms with respect to other metrics. In particular, we study the fraction of solutions dominated by the solution of alternative algorithms and analyze the time complexity of the sorting procedure.

We calculate the percentage of **dominated solutions** as follows. For a given pair of algorithms algorithm1 and algorithm2, we calculate how many solutions of algorithm2 (dominated algorithm) are dominated by solutions of algorithm1 (dominating algorithm). After that, we average the obtained results among all dominating algorithms to get an average fraction of dominated solutions, denoted by  $\theta$ . Naturally, lower values of  $\theta$  indicate better performance. We present the corresponding results in Fig. 14. We notice the following tendencies. NSGA-II and *PO-count* behave very similarly. For  $n_k = 2$ , the value of  $\theta$  for these algorithms is around 20%. After that, it starts increasing and reaches its peak of approximately 45% for  $n_k = 7$ . Finally, it gradually decreases to 24% for  $n_k = 25$ . NSGA-III starts at a similar level and reaches its peak of approximately 30% for  $n_k = 5$ . After that, it decreases below 10% for  $n_k = 7$  and stays relatively close to 0 for the larger numbers of knapsacks. These results demonstrate the superiority of NSGA-III over NSGA-II in the case of many-objective optimization. PO-prob starts at around 16%. However, for  $n_k = 4$  the value of  $\theta$  it already almost 0 and does not go up for larger numbers of knapsacks. This shows that the solutions produced by this algorithm are rarely dominated. Thereby, *PO-prob* is an effective approach for many-objective optimization problems.



Figure 14: Average percentage of solutions dominated by other algorithms,  $\theta$ .

In Fig. 15, we demonstrate the dependence of sorting time on the population size for values of *pop\_size* ranging from 50 to 500. The reported values are the averages over 100 independent executions of one iteration of the corresponding genetic algorithm. From the figure, we can see that *PO-prob* requires much less time than all other algorithm. The results for NSGA-II and PO-count tend to be very close, as in other experiments. This observation also has theoretical explanation. Indeed, choosing the next generation for NSGA-II and NSGA-III has time complexity of  $O(N^2M)$  and  $max\{O(N^2M), O(N^2 \log_{M-2}N)\}$  respectively where M stands for number of objectives and N is the population size, see [Deb et al., 2002, Deb and Jain, 2013]. At the same time, sorting in PO-prob comes down to independent sorting procedures with respect to every objective. The time complexity of 14



Figure 15: Sorting duration as a function of population size for 10 knapsacks,  $n_k = 10$ .

this procedure is O(NMlog(N)). These results are in line with the theoretical analysis presented in Section 5.1 and prove the computational efficiency of the approximate ranking calculation procedure used in PO-prob.

The maximum choice theorem (Theorem 1) has an intuitive interpretation in the context of genetic algorithms. Assume that the selection step is required to pick a selection of a given maximum size for breeding offspring, and both parents are chosen independently and at random form this selection. Then selections obtained via k-Pareto optimality-based sorting yield most offspring with parents offering choice. Choice here means that every parent is strictly superior to the other with respect to at least one objective, or both have the same values of all objectives.

#### Kendall's $\tau$ Rank Correlation Coefficient and Statistical Tests C.2

Let us again consider the case of 2 continuous random variables introduced in Section 4.2. Let us assume that  $X_2 = f(X_1)$  for some increasing function f. For almost all (x, y), either xRy or yRx holds. Thus  $div(\Omega) = 0$ . Moreover, diversity only depends on the copula which encodes the dependency structure between  $X_1$  and  $X_2$ , see Section 4.2.2 and Appendix B.3. Therefore, a value of div( $\Omega$ ) close to zero indicates  $X_1$  and  $X_2$  are strongly correlated via an increasing function. It leads to the idea that Kendall's  $\tau$  rank correlation coefficient [Kendall, 1948] and diversity are strongly related concepts.

Let us consider a sample of n points  $X = \{(x_{i1}, x_{i2})_{i \in \{1, 2, \dots, n\}}\}$ . Duplicates almost never occur and the order in which points are drawn has no importance. Therefore, X should be treated like a set. We consider the counting measure # and the relation  $R_{\downarrow}$  define on X. Diversity and choice of X are denoted by div<sub>#</sub> and cho<sub>#</sub>.

Kendall's  $\tau$  correlation coefficient is defined as follows

$$\tau = \frac{\# \operatorname{con} - \# \operatorname{dis}}{n_0}$$

where # con is the number of concordant pairs (pairs that do not offer choice in our terminology), # dis is the number of discordant pairs (pairs that do offer choice), and  $n_0$  is the total number of pairs. As duplicates are discarded and the pairs are not ordered,  $n_0 = n(n-1)/2$ .

From the above remarks, we have  $cho_{\#} = 2\# dis + n$  and  $\# con + \# dis = n_0$ . Combining these equalities, we obtain the following relation

$$\tau = \frac{n^2 + n - 2 \operatorname{cho}_{\#}}{n^2 - n}.$$

Dividing the numerator and the denominator by  $n^2$  and then neglecting  $\frac{1}{n}$ , we obtain that approximately

$$\tau \approx 1 - 2 \operatorname{div}_{\#}$$

This result means that the theory developed in this paper can be used for constructing non-parametric statistical tests generalizing Kendall's  $\tau$  rank correlation coefficient and can be used for testing partial correlation. Below, we illustrate this property by building an indicator for distinguishing between wealthy and non-wealthy states.

A group of states might be considered *wealthy* if the following two conditions hold. 15

- 1. In the group there is no positive correlation between per capita income and the indicator representing education and health.
- 2. If a state belongs to a group of wealthy states, then all states having higher per capita income and better value of education and health indicator, must also belong to that group.

Now, we define the set of all wealthy states as the largest group of states that are wealthy. If Kendall's  $\tau$  is used to compute correlation, and correlation is considered to be positive if  $\operatorname{div}_{\#} < \frac{1}{2}$ , then the elements of the above set can be easily identified.

Indeed, Corollary 1 says that the set of wealthy states must be a set of at-least k-Pareto optimal states for the relation higher income and better education and health indicator. For the year 2015<sup>15</sup>, we took Gross National Income (GNI) per capita at purchasing power parity (PPP) as the income indicator, and the square root of the education and life expectancy as the education and health indicator<sup>16</sup>. The scatter plot in Fig. 16 shows the resulting division of states into wealthy and non-wealthy. We can observe that the wealthy states are defined as the states with GNI  $\geq$  20000\$. This seems perfectly plausible.



Figure 16: Separation between wealthy and non-wealthy states based on  $\operatorname{div}_{\#}$ .

# C.3 Application to Recommender Systems

Let us again consider the housing example introduced in Section 2.1. If we aim to provide to the user a full set of possible alternative houses that might fit his preferences, then, according to Theorem 1, sorting available habitations by the increasing value of po is the best strategy. As it was discussed in Section 5.1, in the case of independent components of the underlying composite relations, the computation of po can be simplified by using tools from probability theory. Apart from computational efficience, estimating po in this way has several additional advantages.

- Such sorting results in fewer ties and a meaningful score. Indeed, sorting items x by increasing values of po(x), is the same as sorting by decreasing values of  $-\log(po(x))$ . The self-information  $-\log(F_i(x))$  [Jones, 1979], which is additive, indicates how much a characteristic *i* is valued. In this case, there is no need to introduce any arbitrary coefficients as it is done when sorting by a weighted mean of the characteristics  $x_i$ .
- If the condition of independence holds, then *rarer characteristics get valued more*. This makes sense from the economic point of view and is intuitively necessary for maximizing choice.
- If beyond the relation R, there is complete uncertainty about the user's complex needs, tastes and desires, then offering him a selection of maximum choice maximizes the likelihood he finds an appropriate item.

<sup>&</sup>lt;sup>15</sup>Data source: United Nations Development Programme - Human Development Reports http://hdr.undp.org/en/data.

<sup>&</sup>lt;sup>16</sup>As suggested by the human development index, see http://hdr.undp.org/sites/default/files/hdr2020\_technical\_notes. pdf.

#### C.4**Constrained Multi-Objective Genetic Algorithms**

# C.4.1 Problem Definition

A Multiobjective Constrained Optimization Problem (CMOP) is a mathematical problem that is defined as follows [Kumara et al., 2020]:

Minimize

$$f_1(x), f_2(x), \ldots, f_M(x)$$

subject to

$$g_i(x) \le 0, i \in \{1, 2, ..., ng\},\$$
  

$$h_j = 0, j \in \{ng + 1, ng + 2, ..., ng + nh\},\$$
  

$$L_k \le x_k \le U_k, h \in \{1, ..., D\},\$$

where

- $f_i$  represents the *i*-th objective function,
- *M* is the total number of conflicting objective functions,
- $x = (x_1, x_2, \dots, x_D)$  is a solution vector of length D,
- $L_k$  and  $U_k$  are the lower and upper bounds of the search space at the k-th dimension.

Numerically, we consider a constraint  $h_j$  to be verified iff  $h_j \in [-\epsilon, \epsilon]$ . A solution is feasible iff all ng + nhconstraints  $g_i$  and  $h_j$  are verified.

### C.4.2 Problem Re-Definition with Preorder Relations

In a more general setting, we can represent a constraint  $g_i \leq 0$  by the preorder relation  $R_{g_i}$  defined as follows:

$$xR_{g_i}y \text{ iff } \begin{cases} g_i(x) \le 0 \text{ or} \\ g_i(x) \le g_i(y). \end{cases}$$

$$\tag{8}$$

And a constraint  $h_j \in [a_j, b_j]$  can be represented by the preorder relation  $R_{h_j}$  defined as follows:

$$xR_{h_j}y \text{ iff } \begin{cases} h_j(x) \in [a_j, b_j] \text{ or} \\ h_j(y) \le h_j(x) \le a_j \text{ or} \\ b_j \le h_j(x) \le h_j(y). \end{cases}$$
(9)

Then, the combination of the constraints  $g_i \leq 0, i \in \{1, 2, ..., ng\}$  and  $h_j = 0, j \in \{ng + 1, ng + 2, ..., ng + nh\}$ can be represented by the preorder relation  $R_c$  defined as follows.

 $xR_{c}y$  iff  $xR_{g_1}y$  and ... and  $xR_{g_{ng}}y$  and  $xR_{h_{ng+1}}y$  and ... and  $xR_{h_{ng+nh}}y$ .

The objective consisting in minimizing  $f_i$  is represented by the preoduct relation  $R_{f_i}$ :

$$xR_{f_i}y$$
 iff  $f_i(x) \le f_i(y)$ .

Minimization of all M objectives  $f_1, \ldots, f_M$  is represented by the preorder relation  $R_f$  defined as follows

$$xR_f y$$
 iff  $xR_{f_1} y$  and ... and  $xR_{f_M} y$ .

Thus, the above CMOP can be represented by the lexicographic preorder relation

 $xR_{cf}y$  iff  $xR_c^*y$  or  $(xR_c^=y \text{ and } xR_fy)$ ,

where  $xR_c^*y$  means " $xR_cy$  and not  $yR_cx$ ", and  $xR_c^=y$  means " $xR_cy$  and  $yR_cx$ ". For the given  $R_{cf}$ , constrained Pareto optimal solutions [Kumara et al., 2020] are solutions that are not Pareto dominated by any other solution. 17

### C.4.3 Solution

To solve the problem defined above, we can use the standard Adaptive Differential Evolution Algorithm jDE [Noman et al., 2011] with k-Pareto optimality for  $R_{cf}$  as a fitness function. For any point x, k-Pareto optimality of x is the likelihood a point drawn at random from the population strictly Pareto dominates x for  $R_{cf}$ . Smaller values of po mean better fitness. Under the independence assumption of objectives and constraints, we can easily compute k-Pareto optimality po(x). When saying  $P(\{y|yR_{cf}^=x\}) = 0$ , we assume the considered objectives and constraints are not constant on too large sets. Without this simplification, the computation becomes longer, see the derivation below.

$$po(x) = P(\{y|yR_{cf}^*x\}),$$
  
=  $P(\{y|yR_{cf}x\} - P(\{y|yR_{cf}^=x\}),$   
=  $P(\{y|yR_{cf}x\} - 0,$   
=  $P(\{y|yR_{c}^*x\} \cup \{y|yR_{c}^=x \text{ and } yR_{f}x\}),$   
=  $P(\{y|yR_{c}^*x\}) + P(\{y|yR_{c}^=x \text{ and } yR_{f}x\}).$ 

Thus, if x satisfies all constraints, which means  $xR_c^{=}(0,\ldots,0,a_{ng+1},\ldots,a_{ng+nh})$ , then

$$po(x) = P(\{y|yR_c^{-}x \text{ and } yR_fx\}),$$
  
=  $P(\{y|yR_c^{-}(0,...,0,a_{ng+1},...,a_{ng+nh})\})P(\{y|yR_fx\}).$ 

Now, let  $F_i(z) = P(\{y | f_i(y) \le z\})$  be the cumulative probability distribution of  $f_i$ . Then,

$$P(\{y|yR_fx\}) = P\left(\bigcap_{i \in \{1,...,M\}} \{y|f_i(y) \le f_i(x)\}\right),\$$
  
=  $\prod_{i \in \{1,...,M\}} P(y|f_i(y) \le f_i(x)).\$   
=  $\prod_{i \in \{1,...,M\}} F_i(f_i(x)).$ 

Otherwise, if at least one constraint is not satisfied by x, then  $P(\{y|yR_c^=x\}) = 0$  and

$$po(x) = P(\{y|yR_c^*x\})$$
  
=  $P(\{y|yR_cx\})$   
=  $\prod_{i \in \{1,...,ng\}} G_i(g(x)) \prod_{j \in \{ng+1,...,ng+nh\}} P(y|yR_{h_j}x).$ 

Any cumulative probability distribution defined above,  $F_i$  for  $f_i$  and  $G_i$  for  $g_i$ , can be estimated via its empirical cumulative probability distribution. Note, for a population  $\{x_1, \ldots, x_k, \ldots, x_{ps}\}$  of size ps, and for any real valued function f, the empirical cumulative probability distribution  $\hat{F}$  of f is defined as follows:

$$\hat{F}(z) = \frac{\#(\{x_k | f(x_k) \le z\})}{ps}$$

In a similar way,  $P(y|yR_{h_j}x)$  can be estimated by  $\widehat{H_j}^*(h_j(x))$ , where  $\widehat{H_j}^*(z)$  is defined as

$$\widehat{H_{j}}^{*}(z) = \begin{cases} \frac{\#(\{x_{k}|z \le h_{j}(x_{k}) \le b_{j}\}}{ps} & \text{if } z < a_{j}, \\ \frac{\#(\{x_{k}|a_{j} \le h_{j}(x_{k}) \le b_{j}\}\}}{ps} & \text{if } a_{j} \le z \le b_{j}, \\ \frac{\#(\{x_{k}|a_{j} \le h_{j}(x_{k}) \le z\}}{ps} & \text{if } z > b_{j}. \end{cases}$$
(10)

The computation of every  $\hat{F}_i$  can be performed as follows:

- sort the values  $f(x_k)$  in increasing order, and store them in an array;
- create two new arrays;
- loop over the sorted values  $f(x_k)$ ; each time a new distinct value  $f(x_k)$  is encountered:
  - append the previously encountered  $f(x_k)$  to the first array,
  - append to the second array the loop counter, which is equal to the value of the empirical cumulative probability distribution  $F_i$  of the previously encountered value  $f(x_k)$ .

Thus, retrieving  $\hat{F}_i(x)$  can be performed via a binary lookup with run time  $O(\log ps)$ . Computation of  $\widehat{H}_j^*$  can be performed in the same way. In this case, all three cases of the definition in Eq. (10) are treated separately. Moreover, we have the estimation

$$P(\{y|yR_c^{=}(0,\ldots,0,a_{ng+1},\ldots,a_{ng+nh})\}) = \prod_{i\in\{1,\ldots,ng\}} \hat{G}_i(0) \prod_{j\in\{ng+1,\ldots,ng+nh\}} \widehat{H_j}^*(a_j)$$

Finally, it is possible to show that the total run time of the k-Pareto optimality based sorting is O((nq + nh + n))M) $ps \log ps$ ).

#### **Exploratory Database Queries** C.5

Simple database queries q, objectives, and constraints in optimization problems often consist in requiring a continuous attribute to be in a given interval, or a discrete attribute to be equal to a given value. Conceptually, those queries are boolean functions. Complex queries are often conjunctions of the form  $r = q_1 \wedge q_2 \wedge \cdots \wedge q_n$ .

In our formalism, these simple queries translate into simple pre-order relations of the form xRy. Requiring an element to be in an interval can be represented by  $xR_{q}y$  iff x is in the desired interval, or x is not situated further from the interval than  $y^{17}$ . Requiring an attribute to be equal to a given value translates into the relation  $xR_{a'}y$ if for x the attribute takes the required value<sup>18</sup>. Complex queries then translate into the composite relations of the form  $R_r = R_{q_1} \wedge R_{q_2} \wedge \cdots \wedge R_{q_n}$ . The simple sub-relations  $R_{q_2}$  are pre-order relations, and, therefore,  $R_r$  is also a pre-order relation and is transitive. However, these relations are not partial order relations, as reflexivity does not necessarily hold. A "topological" sorting according to our partial order relation  $R_r$  can be viewed as a valid fuzzy relaxation of the strict functional query. There are many possible fuzzy relaxations and the problem is to find one that is suitable for a given application. The k-Pareto optimality is one of such fuzzy extensions of the query. It is 0 if all criteria are satisfied, and higher values of k-Pareto optimality indicate worse results. The maximum choice theorem applies here, and the user is offered the maximum choice. This is of particular interest for exploratory queries, such as job search, especially, if there are no items in the database that satisfy all the criteria. Direct brute-force search for selections offering maximum choice is unfeasible as there are too many selections to consider.

Moreover, in the above formalism, one can treat classical optimization objectives in the same way. Maximizing an attribute x can be represented by the relation xRy iff  $x \ge y$ , and the minimization can be represented by the relation  $\leq$ . In the above framework, negation can be represented via the relation  $R^{-1}$  defined by  $xR^{-1}y$  iff yRx.

#### C.6 Scheduling Algorithms

In the case of scheduling algorithms, xRy can be given the meaning 'x depends on y'. Then, selections represent sets of tasks that remain to be processed. Having a large choice means having much freedom to parallelize tasks or having flexibility in case the rescheduling is required.

<sup>&</sup>lt;sup>17</sup>Hence, the strict version of this relation is defined as follows:  $xR_q^*y$  iff x is in the desired interval and y is not, or if x is situated closer to the desired interval.

<sup>&</sup>lt;sup>18</sup>The strict version of this relation is defined as  $xR_{q'}^*y$  if for x the attribute takes the required value but not for y.