

An error estimate for the Gauss-Jacobi-Lobatto quadrature rule

Concetta Laurita¹

Department of Mathematics, Computer Science and Economics, University of Basilicata, Viale dell'Ateneo Lucano n. 10, 85100 Potenza, ITALY

Abstract

An error estimate for the Gauss-Lobatto quadrature formula for integration over the interval $[-1, 1]$, relative to the Jacobi weight function $w^{\alpha,\beta}(t) = (1-t)^\alpha(1+t)^\beta$, $\alpha, \beta > -1$, is obtained. This estimate holds true for functions belonging to some Sobolev-type subspaces of the weighted space $L^1_{w^{\alpha,\beta}}([-1, 1])$.

Keywords: Gauss-Lobatto formula; Jacobi weight function

2020 MSC: 65D30, 65D32

1. Introduction

For a Jacobi weight function $w^{\alpha,\beta}(t) = (1-t)^\alpha(1+t)^\beta$, $\alpha, \beta > -1$, on the interval $[-1, 1]$ we consider the $(n+2)$ -point Gauss-Lobatto rule

$$\int_{-1}^1 f(t)w^{\alpha,\beta}(t)dt = w_0f(-1) + \sum_{k=1}^n w_kf(t_k) + w_{n+1}f(1) + e_n(f) \quad (1)$$

which is exact for polynomials of degree at most $2n+1$, i.e.

$$e_n(f) = 0, \quad \forall f \in \mathbb{P}_{2n+1}$$

being \mathbb{P}_n the set of all algebraic polynomials on $[-1, 1]$ of degree at most n .

It is well known that the interior quadrature nodes t_k , $k = 1, \dots, n$, are the zeros of the Jacobi polynomial of degree n orthonormal with respect to the Jacobi weight $w^{\alpha+1,\beta+1}(t) = (1-t)^2w^{\alpha,\beta}(t)$. The weights of the formula (1) are given by

$$w_k = \int_{-1}^1 l_k(t)w^{\alpha,\beta}(t)dt, \quad k = 0, 1, \dots, n+1 \quad (2)$$

where, setting from now on $t_0 = -1$ and $t_{n+1} = 1$, $l_k(t)$ denotes the $(k+1)$ -th Lagrange fundamental polynomial associated to the system of nodes $\{t_0, t_1, \dots, t_n, t_{n+1}\}$.

By standard arguments, it can be easily proved that the quadrature error $e_n(f)$ satisfies the following estimate

$$|e_n(f)| \leq 2 \left(\int_{-1}^1 w^{\alpha,\beta}(t) dt \right) E_{2n+1}(f)_\infty, \quad \forall f \in C([-1, 1]),$$

where

$$E_n(f)_\infty = \inf_{P \in \mathbb{P}_n} \|f - P\|_\infty$$

Email address: concetta.laurita@unibas.it (Concetta Laurita)

¹Tel: +390971205846

denotes the error of best approximation of a function $f \in C([-1, 1])$ by means of polynomials of degree at most n with respect to the uniform norm.

The aim of the present paper is to provide a new error estimate for less regular functions belonging to some Sobolev-type subspaces of the weighted space $L_{w^{\alpha, \beta}}^1([-1, 1])$. A similar estimate is proved in [3] for the classical Gauss-Jacobi quadrature formula, in [1] for the Gauss-Lobatto rule with respect to the Legendre weight $w^{0,0}$ and, more recently, in [2] for the Gauss-Radau formula with respect to a general Jacobi weight $w^{\alpha, \beta}$.

2. Notation and preliminary results

2.1. Notation

For a general weight function $w(t)$ on $[-1, 1]$ and $1 \leq p < +\infty$, let L_w^p denote the weighted space of all real-valued measurable functions f on $[-1, 1]$ such that

$$\|f\|_{L_w^p} = \|fw\|_p = \left(\int_{-1}^1 |f(t)w(t)|^p dt \right)^{\frac{1}{p}} < +\infty,$$

and let $W_r^p(w)$ be the following weighted Sobolev-type subspaces of L_w^p

$$W_r^p(w) = \left\{ f \in L_w^p : f^{(r-1)} \in AC(-1, 1), \|f^{(r)}\varphi^r w\|_p < +\infty \right\},$$

where $r \in \mathbb{N}$, $r \geq 1$, $\varphi(t) = \sqrt{1-t^2}$ and $AC(-1, 1)$ is the collection of all functions which are absolutely continuous on every closed subset of $(-1, 1)$, equipped with the norm

$$\|f\|_{W_r^p(w)} = \|fw\|_p + \|f^{(r)}\varphi^r w\|_p.$$

For a function $f \in L_w^p$, the error of the best approximation of f in L_w^p by polynomials of degree at most n is defined as

$$E_n(f)_{w,p} = \inf_{P \in \mathbb{P}_n} \|f - P\|_{L_w^p}.$$

Fixed a Jacobi weight $w^{\gamma, \delta}(t) = (1-t)^\gamma(1+t)^\delta$, $\gamma, \delta > -1$, we will denote by $x_{n,k}^{\gamma, \delta}$ and $\lambda_{n,k}^{\gamma, \delta}$, $k = 1, \dots, n$, the nodes and coefficients of the corresponding n -point Gauss-Jacobi quadrature rule on $[-1, 1]$ and by

$$p_n^{\gamma, \delta}(t) = \gamma_n^{\gamma, \delta} t^n + \text{lower degree terms}, \quad \gamma_n^{\gamma, \delta} > 0,$$

the Jacobi polynomial of degree n orthonormal w.r.t. $w^{\gamma, \delta}(t)$ having positive leading coefficient.

In the sequel \mathcal{C} will denote a positive constant which may assume different values in different formulas. We write $\mathcal{C} = \mathcal{C}(a, b, \dots)$ to say that \mathcal{C} is dependent on the parameters a, b, \dots and $\mathcal{C} \neq \mathcal{C}(a, b, \dots)$ to say that \mathcal{C} is independent of them. Moreover, we will write $A \sim B$, if there exists a positive constant \mathcal{C} independent of the parameters of A and B such that $1/\mathcal{C} \leq A/B \leq \mathcal{C}$.

2.2. Preliminary results

It is well known that for functions f belonging to $W_1^p(w)$, the following Favard inequality

$$E_n(f)_{w,p} \leq \frac{\mathcal{C}}{n} E_{n-1}(f')_{\varphi w,p}, \quad (3)$$

is fulfilled for a positive constant \mathcal{C} independent of n and f (see, for example, [3, (2.5.22), p. 172]). By iteration of inequality (3), it follows that, for $f \in W_r^p(w)$, $r \geq 1$, the estimate

$$E_n(f)_{w,p} \leq \frac{\mathcal{C}}{n^r} E_{n-r}(f^{(r)})_{\varphi^r w,p}, \quad \mathcal{C} \neq \mathcal{C}(n, f) \quad (4)$$

holds true.

Let us recall that the knots $x_{n,k}^{\gamma,\delta}$ and Christoffel numbers $\lambda_{n,k}^{\gamma,\delta}$ of the Gaussian quadrature formula corresponding to the Jacobi weight $w^{\gamma,\delta}$ satisfy the following properties (see, for instance, [3, (4.2.4), p. 249])

$$x_{n,k+1}^{\gamma,\delta} - x_{n,k}^{\gamma,\delta} \sim \frac{\sqrt{1-t^2}}{n}, \quad x_{n,k}^{\gamma,\delta} \leq t \leq x_{n,k+1}^{\gamma,\delta}, \quad (5)$$

and (see [6, (14), p.673])

$$\lambda_{n,k}^{\gamma,\delta} \sim \frac{\sqrt{1 - \left(x_{n,k}^{\gamma,\delta}\right)^2}}{n} w^{\gamma,\delta}(x_{n,k}^{\gamma,\delta}) \quad (6)$$

uniformly for $1 \leq k \leq n$, $n \in \mathbb{N}$. Moreover, for the orthonormal polynomials $\{p_n^{\gamma,\delta}(t)\}_n$ one has that (see [7, (12.7.2), p. 309])

$$\frac{\gamma_n^{\gamma,\delta}}{\gamma_{n-1}^{\gamma,\delta}} \sim 1, \quad \text{as } n \rightarrow \infty, \quad (7)$$

and (see, for instance, [5, p. 170]) the equality

$$\frac{1}{p_{n-1}^{\gamma,\delta}(x_{n,k}^{\gamma,\delta})} = \frac{\gamma_{n-1}^{\gamma,\delta}}{\gamma_n^{\gamma,\delta}} \lambda_{n,k}^{\gamma,\delta} (p_n^{\gamma,\delta})'(x_{n,k}^{\gamma,\delta}) \quad (8)$$

holds true. Furthermore (see [5, Corollary 9.34, p. 171])

$$p_n^{\gamma,\delta}(1) \sim n^{\alpha+\frac{1}{2}} \quad |p_n^{\gamma,\delta}(-1)| \sim n^{\beta+\frac{1}{2}}, \quad (9)$$

uniformly for $n \in \mathbb{N}$ and, more generally, [3, (4.4.49)-(4.2.30), p. 255])

$$|p_n^{\gamma,\delta}(t)| \sim n^{\alpha+\frac{1}{2}}, \quad 1 - \frac{\mathcal{C}}{n^2} \leq t \leq 1, \quad (10)$$

$$|p_n^{\gamma,\delta}(t)| \sim n^{\beta+\frac{1}{2}}, \quad -1 \leq t \leq -1 + \frac{\mathcal{C}}{n^2}. \quad (11)$$

3. Main results

Lemma 3.1. *The nodes and the weights of the Gauss-Lobatto quadrature formula (1) satisfy the following relations*

$$w_0 \leq \mathcal{C} \Delta t_0 w^{\alpha,\beta}(t_1), \quad (12a)$$

$$w_k \sim \begin{cases} \Delta t_k w^{\alpha,\beta}(t_k), & k = 1, \dots, n-1 \\ \Delta t_{k-1} w^{\alpha,\beta}(t_k) & k = n \end{cases}, \quad (12b)$$

$$w_{n+1} \leq \mathcal{C} \Delta t_n w^{\alpha,\beta}(t_n), \quad (12c)$$

where $\Delta t_k = t_{k+1} - t_k$, $k = 0, 1, \dots, n$ and $\mathcal{C} \neq \mathcal{C}(n)$.

Proof. First, let us observe that the weights of the formula (1), given in (2), have the following alternative representation

$$w_0 = \frac{1}{1-t_0} \int_{-1}^1 \frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_0)} w^{\alpha+1,\beta}(t) dt, \quad (13a)$$

$$w_k = \frac{1}{(1-t_k^2)} \int_{-1}^1 \frac{p_n^{\alpha+1,\beta+1}(t)}{(t-t_k) (p_n^{\alpha+1,\beta+1})'(t_k)} w^{\alpha+1,\beta+1}(t) dt, \quad k = 1, \dots, n, \quad (13b)$$

$$w_{n+1} = \frac{1}{1+t_{n+1}} \int_{-1}^1 \frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_{n+1})} w^{\alpha,\beta+1}(t) dt. \quad (13c)$$

For $k = 1, \dots, n$, since

$$w_k = \frac{\lambda_{n,k}^{\alpha+1,\beta+1}}{1 - \left(x_{n,k}^{\alpha+1,\beta+1}\right)^2},$$

using (6) and (5) for $\gamma = \alpha + 1$ and $\delta = \beta + 1$, we can deduce that

$$w_k \sim \begin{cases} w^{\alpha,\beta}(t_k)\Delta t_k & k = 1, \dots, n-1 \\ w^{\alpha,\beta}(t_k)\Delta t_{k-1} & k = n \end{cases}$$

i.e. (12b). Now, in order to prove (12a), let us observe that, since

$$\int_{-1}^1 \left[\frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_0)} - \left(\frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_0)} \right)^2 \right] w^{\alpha+1,\beta}(t) dt = 0,$$

we can rewrite the first coefficient w_0 in (13a) as follows

$$w_0 = \frac{1}{2} \left[\frac{(p_n^{\alpha+1,\beta+1})'(t_1)}{p_n^{\alpha+1,\beta+1}(t_0)} \right]^2 \int_{-1}^1 \left[\frac{p_n^{\alpha+1,\beta+1}(t)}{(t-t_1)(p_n^{\alpha+1,\beta+1})'(t_1)} \right]^2 \frac{(t-t_1)^2}{1+t} w^{\alpha+1,\beta+1}(t) dt.$$

Being $(t-t_1)^2/(1+t) \leq \mathcal{C}$ and using (8) (with $\gamma = \alpha + 1$ and $\delta = \beta + 1$) for $k = 1$ (we recall that in our notation $x_{n,1}^{\alpha+1,\beta+1} \equiv t_1$), we get

$$\begin{aligned} w_0 &\leq \mathcal{C} \left[\frac{(p_n^{\alpha+1,\beta+1})'(t_1)}{p_n^{\alpha+1,\beta+1}(t_0)} \right]^2 \lambda_{n,1}^{\alpha+1,\beta+1} \\ &= \mathcal{C} \frac{1}{\left[p_n^{\alpha+1,\beta+1}(t_0)\right]^2} \left(\frac{\gamma_n^{\alpha+1,\beta+1}}{\gamma_{n-1}^{\alpha+1,\beta+1}} \right)^2 \frac{1}{\lambda_{n,1}^{\alpha+1,\beta+1} \left[p_{n-1}^{\alpha+1,\beta+1}(t_1)\right]^2}. \end{aligned}$$

Now, in virtue of (6), it is

$$\lambda_{n,1}^{\alpha+1,\beta+1} \sim \frac{\left(1 - x_{n,1}^{\alpha+1,\beta+1}\right)^{\alpha+\frac{3}{2}} \left(1 + x_{n,1}^{\alpha+1,\beta+1}\right)^{\beta+\frac{3}{2}}}{n} \sim \frac{1}{n^{2\beta+4}}. \quad (14)$$

Taking into account (14), and (7), (9) and (11) (all applied with $\gamma = \alpha + 1$ and $\delta = \beta + 1$), we deduce the following estimate of w_0

$$w_0 \leq \mathcal{C} \frac{n^{2\beta+4}}{n^{4\beta+6}} = \frac{\mathcal{C}}{n^{2\beta+2}}. \quad (15)$$

On the other hand,

$$\Delta t_0 w^{\alpha,\beta}(t_1) = \left(1 - x_{n,1}^{\alpha+1,\beta+1}\right)^\alpha \left(1 + x_{n,1}^{\alpha+1,\beta+1}\right)^{\beta+1} \sim \frac{1}{n^{2\beta+2}}$$

which, combined with (15), gives (12a). In order to prove (12c), proceeding in an analogous way, we start writing w_{m+1} as follows

$$w_{m+1} = \frac{1}{2} \left[\frac{(p_n^{\alpha+1,\beta+1})'(t_n)}{p_n^{\alpha+1,\beta+1}(t_{n+1})} \right]^2 \int_{-1}^1 \left[\frac{p_n^{\alpha+1,\beta+1}(t)}{(t-t_n)(p_n^{\alpha+1,\beta+1})'(t_n)} \right]^2 \frac{(t-t_n)^2}{1-t} w^{\alpha+1,\beta+1}(t) dt.$$

Then, using $(t - t_n)^2/(1 - t) \leq \mathcal{C}$, (6) and (8) for $\gamma = \alpha + 1$, $\delta = \beta + 1$ and $k = n$, (7), (9) and (10) also with $\gamma = \alpha + 1$ and $\delta = \beta + 1$, we deduce the estimate

$$w_{n+1} \leq \mathcal{C} \left[\frac{(p_n^{\alpha+1, \beta+1})'(t_n)}{p_n^{\alpha+1, \beta+1}(t_{n+1})} \right]^2 \lambda_{n,n}^{\alpha+1, \beta+1} \sim \frac{1}{n^{2\alpha+2}}$$

with $\mathcal{C} \neq \mathcal{C}(n)$. Since

$$\Delta t_n w^{\alpha, \beta}(t_n) \sim (1 - x_{n,n}^{\alpha+1, \beta+1})^{\alpha+1} (1 + x_{n,n}^{\alpha+1, \beta+1})^\beta \sim \frac{1}{n^{2\alpha+2}}$$

we can, finally, conclude that also (12c) holds true. \square

Using the previous lemma we are able to prove our main result.

Theorem 3.1. *For $f \in W_r^1(w^{\alpha, \beta})$, $r \geq 1$, the error of the Gauss-Lobatto quadrature formula (1) satisfies the following estimate*

$$|e_n(f)| \leq \frac{\mathcal{C}}{(2n)^r} E_{2n+1-r}(f^{(r)})_{\varphi^r w^{\alpha, \beta}, 1} \quad (16)$$

where $\varphi(t) = \sqrt{1 - t^2}$ and $\mathcal{C} \neq \mathcal{C}(f, n)$ is a positive constant.

Proof. We start by proving the estimate (16) in the case $r = 1$. First, we are going to show that

$$\sum_{k=0}^{n+1} w_k |f(t_k)| \leq \mathcal{C} \|f w^{\alpha, \beta}\|_1 + \frac{\mathcal{C}}{n} \int_{-1}^1 |f'(t)| \varphi(t) w^{\alpha, \beta}(t) dt. \quad (17)$$

Taking into account (12a), (12b) and (12c) we have

$$\begin{aligned} \sum_{k=0}^{n+1} w_k |f(t_k)| &\leq \mathcal{C} [\Delta t_0 w^{\alpha, \beta}(t_1) |f(t_0)| + \sum_{k=1}^{n-1} \Delta t_k w^{\alpha, \beta}(t_k) |f(t_k)| + \\ &\quad + \Delta t_{n-1} w^{\alpha, \beta}(t_n) |f(t_n)| + \Delta t_n w^{\alpha, \beta}(t_n) |f(t_{n+1})|]. \end{aligned}$$

In virtue of the following inequality

$$\left. \begin{aligned} (b-a)|f(a)| \\ (b-a)|f(b)| \end{aligned} \right\} \leq \int_a^b |f(t)| dt + (b-a) \int_a^b |f'(t)| dt,$$

it follows that

$$\begin{aligned} \sum_{k=0}^{n+1} w_k |f(t_k)| &\leq \mathcal{C} \left[w^{\alpha, \beta}(t_1) \left(\int_{t_0}^{t_1} |f(t)| dt + \Delta t_0 \int_{t_0}^{t_1} |f'(t)| dt \right) \right. \\ &\quad + \sum_{k=1}^{n-1} w^{\alpha, \beta}(t_k) \left(\int_{t_k}^{t_{k+1}} |f(t)| dt + \Delta t_k \int_{t_k}^{t_{k+1}} |f'(t)| dt \right) \\ &\quad + w^{\alpha, \beta}(t_n) \left(\int_{t_{n-1}}^{t_n} |f(t)| dt + \Delta t_{n-1} \int_{t_{n-1}}^{t_n} |f'(t)| dt \right) \\ &\quad \left. + w^{\alpha, \beta}(t_n) \left(\int_{t_n}^{t_{n+1}} |f(t)| dt + \Delta t_n \int_{t_n}^{t_{n+1}} |f'(t)| dt \right) \right], \end{aligned}$$

from which, being for $k = 0, 1, \dots, n$,

$$1 \pm t_k \sim 1 \pm t \sim 1 \pm t_{k+1}, \quad t_k \leq t \leq t_{k+1}, \quad (18)$$

and, taking into account (5) and also that

$$\begin{aligned}\Delta t_0 &\sim \frac{\sqrt{1-t^2}}{n}, & -1 < t \leq t_1, \\ \Delta t_n &\sim \frac{\sqrt{1-t^2}}{n}, & t_n \leq t < 1,\end{aligned}$$

we deduce

$$\begin{aligned}\sum_{k=0}^{n+1} w_k |f(t_k)| &\leq \mathcal{C} \left[\int_{t_0}^{t_1} |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{t_0}^{t_1} |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right. \\ &+ \sum_{k=1}^{n-1} \left(\int_{t_k}^{t_{k+1}} |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{t_k}^{t_{k+1}} |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right) \\ &+ \left. \int_{t_n}^{t_{n+1}} |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{t_n}^{t_{n+1}} |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right] \\ &= \mathcal{C} \left[\int_{-1}^1 |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{-1}^1 |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right]\end{aligned}$$

i.e. (17). Since the $(n+1)$ quadrature formula (1) is exact for any polynomial of degree at most $2n+1$, for $P \in \mathbb{P}_{2n+1}$ we have

$$\begin{aligned}|e_n(f)| &= |e_n(f-P)| \\ &\leq \left| \int_{-1}^1 (f-P)(t) w^{\alpha,\beta}(t) dt \right| + \left| \sum_{k=0}^{n+1} w_k (f-P)(t_k) \right| \\ &\leq \|(f-P)w^{\alpha,\beta}\|_1 + \sum_{k=0}^{n+1} w_k |(f-P)(t_k)|.\end{aligned}$$

Now, combining the previous inequality with (17) and the following relation (see, for instance, [3, p. 339], [4, p. 286])

$$\|(f-P)' \varphi w^{\alpha,\beta}\|_1 \leq \mathcal{C}(2n+2) \|(f-P)w^{\alpha,\beta}\|_1 + \mathcal{C}_1 E_{2n}(f')_{\varphi w^{\alpha,\beta},1},$$

we can deduce

$$\begin{aligned}|e_n(f)| &\leq \mathcal{C} \|(f-P)w^{\alpha,\beta}\|_1 + \frac{\mathcal{C}}{n} \|(f-P)' \varphi w^{\alpha,\beta}\|_1 \\ &\leq \mathcal{C} \|(f-P)w^{\alpha,\beta}\|_1 + \frac{\mathcal{C}_1}{n} E_{2n}(f')_{\varphi w^{\alpha,\beta},1}.\end{aligned}$$

Taking the infimum over $P \in \mathbb{P}_{2n+1}$ and using the Favard inequality (4) we get

$$|e_n(f)| \leq \mathcal{C} E_{2n+1}(f)_{w^{\alpha,\beta},1} + \frac{\mathcal{C}_1}{n} E_{2n}(f')_{\varphi w^{\alpha,\beta},1} \leq \frac{\mathcal{C}}{2n} E_{2n}(f')_{\varphi w^{\alpha,\beta},1}. \quad (19)$$

The estimate (16) for $r > 1$ can be deduced from (19) by iterating the application of the Favard inequality. \square

Acknowledgements

The author is partially supported by University of Basilicata (local funds). The author is member of the INdAM Research group GNCS and the TAA-UMI Research Group. This research has been accomplished within ‘‘Research Italian network on Approximation’’ (RITA).

References

- [1] L. Fermo, C. Laurita, A Nyström method for a boundary integral equation related to the Dirichlet problem on domains with corners, *Numer. Math.* 130 (2015) 35–71.
- [2] C. Laurita, A new stable numerical method for Mellin integral equations in weighted spaces with uniform norm, *Calcolo* 57 (2020) 25.
- [3] G. Mastroianni, G.V. Milovanović, *Interpolation Processes Basic Theory and Applications*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, Heidelberg, 2009.
- [4] G. Mastroianni, M.G. Russo, Lagrange interpolation in weighted Besov spaces, *Constr. Approx.* 15 (1999) 257–289.
- [5] G.P. Nevai, *Orthogonal Polynomials*, Amer. Math. Soc., Providence, R.I., 1979.
- [6] G.P. Nevai, Mean convergence of Lagrange interpolation, III, *Trans. Amer. Math. Soc.* 282 (1984) 669–698.
- [7] G. Szegő, *Orthogonal polynomials*, volume 23 of *Amer. Math. Soc. Colloq. Publ.*, Amer. Math. Soc., Providence, R.I., 4th ed. edition, 1975.