On the characterization of a finite random field by conditional distribution and its Gibbs form

Khachatryan L.A.¹ and Nahapetian B.S.²

¹Institute of Mathematics, NAS of RA, *linda@instmath.sci.am* ²Institute of Mathematics, NAS of RA, *nahapet@instmath.sci.am*

Abstract

In this paper, we show that the methods of mathematical statistical physics can be successfully applied to random fields in finite volumes. As a result, we obtain simple necessary and sufficient conditions for the existence and uniqueness of a finite random field with a given system of one-point conditional distributions. Using the axiomatic (without the notion of potential) definition of Hamiltonian, we show that any finite random field is Gibbsian. We also apply the proposed approach to Markov random fields.

Key Words: Random field, conditional distribution, Gibbs distribution, transition energy field, Hamiltonian

Mathematics Subject Classification 2010: 60G60, 60E05, 60J99, 62H99, 82B03

1 Introduction

The idea of characterization of a multivariate probability distribution by means of conditional probabilities is very old and goes back to the concept of the finite Markov chain. Already here, from the very beginning, we are faced with two fundamentally important characterization problems: the problem of restoring a chain from the system of its transition probabilities and the problem of the existence of a Markov chain corresponding to a given stochastic matrix.

Both of these problems are relevant in the general setting, when one deals with an arbitrary random field in an infinite or finite volume. Having solved the problem of restoring a random field from the system of its conditional probabilities, we can establish the general properties of the random field in these terms. The existence of a random field with a given system of conditional distributions allows constructing models of random fields with the required properties.

In mathematical statistical physics, as a rule, probabilistic models in infinite volumes are studied. This is due to the fact that it is in such models that the phenomenon of phase transition is best described (see, for example, the book of Sinai [27]).

Dobrushin [11] was the first to consider the problem of characterization of a random field on an infinite integer lattice by specification — a system of consistent probability distributions parameterized by infinite boundary conditions. He obtained criteria for both the existence and uniqueness of a random field with a given specification. The basic example of a specification is the Gibbs specification. Further, Dobrushin [12] introduced the concept of a Gibbs random field as a field the conditional distribution of which almost everywhere coincides with the Gibbs specification. This concept was later independently discovered by Lanford and Ruelle [25].

There is a large number of works devoted to the characterization problems of a random field specified in a finite volume (finite random field). The main issue considered here is to find the conditions under which a given system of probability distributions is compatible, that is, there exists a random field which conditional probabilities coincide with the elements of the given system. This issue was studied by Besag [3], Abrahams and Thomas [1], Arnold and Press [2], Gurevich [18], and Chen [5] among others. For finite random fields, it is also of interest to find the conditions under which a given field is Gibbsian. This question was considered, for example, by Hammersley and Clifford [20], Griffeath [17] and Kaiser and Cressie [21].

The methods used to solve the characterization problems in finite and infinite volumes are fundamentally different. Apparently, this led to the situation where the research in these two directions was carried out independently. At the same time, the methods developed in these areas can be mutually beneficial. As an example, we cite the Grimmett's theorem [19] on the construction of the interaction potential of a finite Gibbs random field based on the Möbius transform. This approach was used by Sullivan [28] and Kozlov [23] to find the conditions under which the Dobrushin's specification has a Gibbs form with some potential. Representation of a specification or a finite random field in Gibbs form allows one to study their properties using the well-developed mathematical apparatus of the theory of Gibbs random fields. This approach is especially productive when studying Markov (see, for example, works by Griffeath [17], Sullivan [28], Spitzer [29], and Georgii [16]) and Gaussian random fields (we refer Dobrushin [14], Georgii [16] and Künsch [24] among others).

In this paper, we show that the methods developed in [8] and [9] for infinite random fields can be successfully applied to random fields in finite volumes. As a result, we obtain simple necessary and sufficient conditions for the existence and uniqueness of a finite random field with a given system of one-point conditional distributions. In this case, we establish an explicit formula for the multivariate distribution in terms of the given one-point conditional distributions. In addition, it is shown that in a finite volume the Gibbs form is universal, namely, the elements of any compatible system of probability distributions have the Gibbs form. As an application, we obtain the generalization of the well-known Hammersley–Clifford theorem [20] on the equivalence of Markov and Gibbs finite random fields.

For simplicity, in this article, we consider random fields with a finite state space. However, all the results can be extended to the case of both countable and continuous state spaces.

The paper has the following structure. In Section 2, we provide basic notations. Section 3 is devoted to the problem of characterization of finite random fields by systems of consistent one-point distributions parameterized by boundary conditions. In Section 4, we consider the question of the Gibbsianess of finite random fields. Section 5 provides some concluding remarks.

2 Notations

Let Λ be a set with a finite number of elements, $|\Lambda| < \infty$, and let each point $t \in \Lambda$ be associated with the set X^t , which is a copy of some finite set X. For any $V \subseteq \Lambda$, denote by $X^V = \{x = (x_t, t \in V) : x_t \in X^t, t \in V\}$ the set of functions (configurations) defined on V and tacking values in X.

For any $V \subset \Lambda$, denote by x_V the restriction of configuration $x \in X^{\Lambda}$ on V. When denoting one-point sets $\{t\}$, $t \in \Lambda$, braces will be omitted in some cases. In particular, for the restriction of a configuration $x \in X^{\Lambda}$ on $t \in \Lambda$, the notation x_t will be used instead of $x_{\{t\}}$. For any $V, I \subset \Lambda$ such that $V \cap I = \emptyset$ and any $x \in X^V$, $y \in X^I$, denote by xy the concatenation of x with y, that is, the configuration on $V \cup I$ equal to x on V and to y on I. The concatenation of x with an empty configuration \emptyset we assume to coincide with x, i.e., $x\emptyset = x$ for all $x \in X^V$, $V \subseteq \Lambda$.

Probability distribution P_{Λ} on X^{Λ} will be called a (*finite*) random field. A random field P_{Λ} is called positive if $P_{\Lambda}(x) > 0$ for all $x \in X^{\Lambda}$. By P_V we denote the restriction of P_{Λ} on $V \subset \Lambda$, that is,

$$P_V(x) = \sum_{u \in X^{\Lambda \setminus V}} P_{\Lambda}(xu), \qquad x \in X^V.$$

For $V = \emptyset$, one has $P_{\emptyset}(\emptyset) = 1$.

For a random field P_{Λ} , define a conditional probability distribution

$$Q_V^z(x) = \frac{P_{V \cup I}(xz)}{P_I(z)}, \qquad x \in X^V, z \in X^I,$$

on X^V under the (boundary) condition z, where $V, I \subset \Lambda, V \cap I = \emptyset$ and $P_I(z) > 0$.

3 Characterization of random fields by conditional distributions

In this section, we give necessary and sufficient conditions under which there is a unique random field, which conditional probabilities coincide with the corresponding elements of a given system of probability distributions.

3.1 Positive random fields

Let P_{Λ} be a positive random field and

$$Q_t^z(x) = \frac{P_\Lambda(xz)}{\sum\limits_{u \in X^t} P_\Lambda(uz)}, \qquad x \in X^t, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda,$$
(1)

be its one-point conditional probabilities. The family $Q_1(P_{\Lambda}) = \{Q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ we will call *one-point conditional distribution* of the random field P_{Λ} .

Let $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ be a family of one-point probability distributions q_t^z on X^t parameterized by configurations $z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda$. In mathematical statistical physics, the parameter z is usually called the boundary condition. In the future, we will adhere to this terminology.

A random field P_{Λ} is said to be *compatible* with a system \mathbf{Q}_1 if $\mathbf{Q}_1(P_{\Lambda}) = \mathbf{Q}_1$.

Theorem 1. For a given system $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of positive one-point probability distributions parameterized by boundary conditions, there exists a unique random field P_{Λ} compatible with it if and only if the elements of \mathbf{Q}_1 satisfy the following consistency conditions: for all $t, s \in \Lambda$ and $x, u \in X^t$, $y, v \in X^s, z \in X^{\Lambda \setminus \{t,s\}}$, it holds

$$q_t^{zy}(x)q_s^{zx}(v)q_t^{zv}(u)q_s^{zu}(y) = q_t^{zy}(u)q_s^{zu}(v)q_t^{zv}(x)q_s^{zx}(y).$$
(2)

Proof. The necessity of conditions (2) follows directly from the definition of conditional probabilities. Let us prove the sufficiency by giving an explicit form of the finite random field P_{Λ} defined by the system \mathbf{Q}_{1} .

For any $x \in X^{\Lambda}$, put

$$P_{\Lambda}(x) = \prod_{j=1}^{n} \frac{q_{t_j}^{(xu)_j}(x_j)}{q_{t_j}^{(xu)_j}(u_j)} \cdot \left(\sum_{\alpha \in X^{\Lambda}} \prod_{j=1}^{n} \frac{q_{t_j}^{(\alpha u)_j}(\alpha_j)}{q_{t_j}^{(\alpha u)_j}(u_j)}\right)^{-1},$$
(3)

where $u \in X^{\Lambda}$, $\Lambda = \{t_1, t_2, ..., t_n\}$ is some enumeration of points in Λ , $n = |\Lambda|$, and

$$(xu)_j = x_1 \dots x_{j-1} u_{j+1} \dots u_n, \quad 1 < j < n, \qquad (xu)_1 = u_2 u_3 \dots u_n, \quad (xu)_n = x_1 x_2 \dots x_{n-1} u_n u_n,$$

 $x_j = x_{t_j}, u_j = u_{t_j}, 1 \le j \le n$. Expanded, we have

$$\prod_{j=1}^{n} \frac{q_{t_{j}}^{(xu)_{j}}(x_{j})}{q_{t_{j}}^{(xu)_{j}}(u_{j})} = \frac{q_{t_{1}}^{u_{2}u_{3}\dots u_{n}}(x_{1})}{q_{t_{1}}^{u_{2}u_{3}\dots u_{n}}(u_{1})} \cdot \frac{q_{t_{2}}^{x_{1}u_{3}\dots u_{n}}(x_{2})}{q_{t_{2}}^{x_{1}u_{3}\dots u_{n}}(u_{2})} \cdot \frac{q_{t_{3}}^{x_{1}x_{2}u_{4}\dots u_{n}}(x_{3})}{q_{t_{3}}^{x_{1}x_{2}u_{4}\dots u_{n}}(u_{3})} \cdot \dots \cdot \frac{q_{t_{n}}^{x_{1}x_{2}\dots x_{n-1}}(x_{n})}{q_{t_{n}}^{x_{1}x_{2}\dots x_{n-1}}(u_{n})}.$$

First of all, let us make sure that the formula (3) is correct. To do this, it is necessary to check that the values of P_{Λ} do not depend on the enumeration of the elements of Λ and on the choice of the configuration $u \in X^{\Lambda}$.

We start by showing that for fixed $x, u \in X^{\Lambda}$, the right-hand side of (3) does not depend on the enumeration of the elements of Λ . Note that any enumeration of the elements of Λ can be obtained from another one by successive changing the numeration of two points. For some $k, 1 < k \leq n$, consider the

numeration $\varphi(j)$, $1 \leq j \leq n$, such that $t_{\varphi(k-1)} = t_k$, $t_{\varphi(k)} = t_{k-1}$, and $t_{\varphi(j)} = t_j$ for $j \neq k-1$ and $j \neq k$. Then we have

$$\prod_{j=1}^{n} \frac{q_{t_{\varphi(j)}}^{(xu)_{\varphi(j)}}(x_{\varphi(j)})}{q_{t_{\varphi(j)}}^{(xu)_{\varphi(j)}}(u_{\varphi(j)})} = \prod_{\substack{j \neq k-1, k}} \frac{q_{t_{j}}^{(xu)_{j}}(x_{j})}{q_{t_{j}}^{(xu)_{j}}(u_{j})} \\
\cdot \frac{q_{t_{k}}^{x_{1}...x_{k-2}u_{k-1}u_{k+1}...u_{n}}(x_{k})}{q_{t_{k}}^{x_{1}...x_{k-2}u_{k-1}u_{k+1}...u_{n}}(u_{k})} \cdot \frac{q_{t_{k-1}}^{x_{1}...x_{k-2}x_{k}u_{k+1}...u_{n}}(x_{k-1})}{q_{t_{k-1}}^{x_{1}...x_{k-2}x_{k}u_{k+1}...u_{n}}(u_{k-1})}$$

Denoting $z = x_1 \dots x_{k-2} u_{k+1} \dots u_n$, we have

$$\frac{q_{t_k}^{x_1\dots x_{k-2}u_{k-1}u_{k+1}\dots u_n}(x_k)}{q_{t_k}^{x_1\dots x_{k-2}u_{k-1}u_{k+1}\dots u_n}(u_k)} \cdot \frac{q_{t_{k-1}}^{x_1\dots x_{k-2}x_ku_{k+1}\dots u_n}(x_{k-1})}{q_{t_{k-1}}^{x_1\dots x_{k-2}x_ku_{k+1}\dots u_n}(u_{k-1})} = \frac{q_{t_k}^{zu_{k-1}}(x_k)}{q_{t_k}^{zu_{k-1}}(u_k)} \cdot \frac{q_{t_{k-1}}^{zx_k}(x_{k-1})}{q_{t_{k-1}}^{zx_k}(u_{k-1})}.$$

According to the consistency conditions (2) of the elements of the system \mathbf{Q}_1 , we can write

$$\frac{q_{t_k}^{zu_{k-1}}(x_k)}{q_{t_k}^{zu_{k-1}}(u_k)} \cdot \frac{q_{t_{k-1}}^{zx_k}(x_{k-1})}{q_{t_{k-1}}^{zx_k}(u_{k-1})} = \frac{q_{t_{k-1}}^{zu_k}(x_{k-1})}{q_{t_{k-1}}^{zu_k}(u_{k-1})} \cdot \frac{q_{t_k}^{zx_{k-1}}(x_k)}{q_{t_k}^{zx_{k-1}}(u_k)}$$

Noting that

$$\frac{q_{t_{k-1}}^{zu_k}(x_{k-1})}{q_{t_{k-1}}^{zu_k}(u_{k-1})} \cdot \frac{q_{t_k}^{zx_{k-1}}(x_k)}{q_{t_k}^{zx_{k-1}}(u_k)} = \frac{q_{t_{k-1}}^{(xu)_{k-1}}(x_{k-1})}{q_{t_{k-1}}^{(xu)_{k-1}}(u_{k-1})} \cdot \frac{q_{t_k}^{(xu)_k}(x_k)}{q_{t_k}^{(xu)_k}(u_k)}$$

we conclude

$$\prod_{j=1}^{n} \frac{q_{t_{\varphi(j)}}^{(xu)_{\varphi(j)}}(x_{\varphi(j)})}{q_{t_{\varphi(j)}}^{(xu)_{\varphi(j)}}(u_{\varphi(j)})} = \prod_{j=1}^{n} \frac{q_{t_{j}}^{(xu)_{j}}(x_{j})}{q_{t_{j}}^{(xu)_{j}}(u_{j})}.$$

In order to prove that P_{Λ} does not depend on the choice of the configuration $u \in X^{\Lambda}$, it is sufficient to verify that the right-hand side of (3) will not change when substituting some other configuration $y \in X^{\Lambda}$ instead of u. Note that (3) can be written in the following form

$$P_{\Lambda}(x) = q_{t_1}^{u_{\Lambda \setminus \{t_1\}}}(x_1) \prod_{j=2}^n \frac{q_{t_j}^{(xu)_j}(x_j)}{q_{t_j}^{(xu)_j}(u_j)} \left(\sum_{\alpha \in X^{\Lambda}} q_{t_1}^{u_{\Lambda \setminus \{t_1\}}}(\alpha_1) \prod_{j=2}^n \frac{q_{t_j}^{(\alpha u)_j}(\alpha_j)}{q_{t_j}^{(\alpha u)_j}(u_j)} \right)^{-1},$$

and hence, the value of P_{Λ} does not depend on the t_1 component of the configuration u. Since it also does not depend on the enumeration of the elements of Λ , the value of P_{Λ} does not change if we substitute into (3) a configuration $u' \in X^{\Lambda}$ that differs from u at one component. It remains to note that any configuration y can be obtained from configuration u by sequentially changing one component.

It is not difficult to see that P_{Λ} is a probability distribution on X^{Λ} , and $P_{\Lambda}(x) > 0$ for all $x \in X^{\Lambda}$. To conclude the proof of the theorem, it remains to verify that $Q_1(P_{\Lambda}) = Q_1$. For this, we need to show that $Q_t^z(x) = q_t^z(x)$ for any $t \in \Lambda$ and $x \in X^t$, $z \in X^{\Lambda \setminus \{t\}}$.

Since the values of P_{Λ} do not depend on the enumeration of the elements in Λ , for every $t \in \Lambda$ and all $x \in X^t$, $z \in X^{\Lambda \setminus \{t\}}$, we can assume $t = t_n$ and write

$$\begin{aligned} Q_t^z(x) &= \frac{P(xz)}{\sum\limits_{\alpha \in X^t} P(\alpha z)} = \frac{q_{t_1}^{(zu)_1}(z_1)q_{t_2}^{(zu)_2}(z_2) \cdot \ldots \cdot q_{t_{n-1}}^{(zu)_{n-1}}(z_{n-1})q_{t_n}^{(zu)_n}(x)}{q_{t_1}^{(zu)_1}(u_1)q_{t_2}^{(zu)_2}(u_2) \cdot \ldots \cdot q_{t_{n-1}}^{(zu)_{n-1}}(u_{n-1})q_{t_n}^{(zu)_n}(u_n)} \\ &\cdot \left(\sum\limits_{\alpha \in X^t} \frac{q_{t_1}^{(zu)_1}(z_1)q_{t_2}^{(zu)_2}(z_2) \cdot \ldots \cdot q_{t_{n-1}}^{(zu)_{n-1}}(z_{n-1})q_{t_n}^{(zu)_n}(\alpha)}{q_{t_1}^{(zu)_1}(u_1)q_{t_2}^{(zu)_2}(u_2) \cdot \ldots \cdot q_{t_{n-1}}^{(zu)_{n-1}}(u_{n-1})q_{t_n}^{(zu)_n}(u_n)}\right)^{-1} \\ &= \frac{q_{t_n}^{(zu)_n}(x)}{\sum\limits_{\alpha \in X^{\{t\}}} q_{t_n}^{(zu)_n}(\alpha)} = q_t^z(x). \end{aligned}$$

 \Box

Note that the formula (3) can be written in the following equivalent forms

$$P_{\Lambda}(x) = \left(\sum_{\alpha \in X^{\Lambda}} \prod_{j=1}^{n} \frac{q_{t_{j}}^{(\alpha x)_{j}}(\alpha_{j})}{q_{t_{j}}^{(\alpha x)_{j}}(x_{j})}\right)^{-1} = \frac{q_{t_{1}}^{x_{\Lambda \setminus \{t_{1}\}}}(x_{1})}{\sum_{\alpha \in X^{\Lambda \setminus \{t_{n}\}}} \frac{q_{t_{1}}^{x_{\Lambda \setminus \{t_{1}\}}}(\alpha_{1})}{q_{t_{n}}^{\alpha}(x_{n})} \prod_{j=2}^{n-1} \frac{q_{t_{j}}^{(\alpha x)_{j}}(\alpha_{j})}{q_{t_{j}}^{(\alpha x)_{j}}(x_{j})}}$$
(4)

for any $x \in X^{\Lambda}$ and any enumeration $\Lambda = \{t_1, t_2, ..., t_n\}$ of points in Λ , $n = |\Lambda|$. Such representations make it obvious that the right-hand side of (3) does not depend on the choice of configuration $u \in X^{\Lambda}$.

3.2 Weakly positive random fields

Now let us consider random fields P_{Λ} which can take zero values.

A random field P_{Λ} we call *weakly positive* if for each $t \in \Lambda$ there exists an element $\theta_t \in X^t$ such that $P_{\Lambda}(\theta_t z) > 0$ for any $z \in X^{\Lambda \setminus \{t\}}$.

Thus, a weakly positive random field P_{Λ} takes strictly positive values on those configurations $x \in X_{\Lambda}$ which have at least one component x_t coinciding with θ_t , $t \in \Lambda$. It is clear, that for such random field, all the one-point conditional probabilities (1) are defined, and $Q_t^z(\theta_t) > 0$ for any boundary condition $z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda$.

A family $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of one-point probability distributions q_t^z on X^t parameterized by boundary conditions $z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda$, will be called *weakly positive* if for each $t \in \Lambda$ there exists an element $\theta_t \in X^t$ for which $q_t^z(\theta_t) > 0$ under any boundary condition $z \in X^{\Lambda \setminus \{t\}}$. Such an element θ_t we will call *positivity point* of the system \mathbf{Q}_1 at $t \in \Lambda$ (compare with the analogous notion in [8]). Note that if for a weakly positive family $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of probability distributions, all

Note that if for a weakly positive family $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of probability distributions, all its positivity points $\theta_t, t \in \Lambda$, coincide with some specified element $\theta \in X$, then θ is called the vacuum and \mathbf{Q}_1 is called *vacuum system*. Such systems play important role in the mathematical statistical physics.

The following statement holds true.

Theorem 2. For a given weakly positive system $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$, there exists a unique weakly positive random field P_{Λ} compatible with it if and only if the elements of \mathbf{Q}_1 satisfy the following consistency conditions: for all $t, s \in \Lambda$ and $x \in X^t$, $y \in X^s$, $z \in X^{\Lambda \setminus \{t,s\}}$, it holds

$$q_t^{zy}(x)q_s^{zx}(\theta_s)q_t^{z\theta_s}(\theta_t)q_s^{z\theta_t}(y) = q_s^{zx}(y)q_t^{zy}(\theta_t)q_s^{z\theta_t}(\theta_s)q_t^{z\theta_s}(x),$$

where θ_t and θ_s are positivity points of the system \mathbf{Q}_1 .

Proof. The necessity is obvious. Let us prove the sufficiency.

For any $x \in X^{\Lambda}$, put

$$P_{\Lambda}(x) = Z^{-1}(\theta_{t_1}\theta_{t_2}...\theta_{t_{|\Lambda|}}) \frac{q_{t_2}^{\theta_{t_2}\theta_{t_3}...\theta_{t_{|\Lambda|}}}(x_{t_1})q_{t_2}^{x_{t_1}\theta_{t_3}...\theta_{t_{|\Lambda|}}}(x_{t_2}) \cdot ... \cdot q_{t_{|\Lambda|}}^{x_{t_1}x_{t_2}...x_{t_{|\Lambda|-1}}}(x_{t_{|\Lambda|}})}{q_{t_1}^{\theta_{t_2}\theta_{t_3}...\theta_{t_{|\Lambda|}}}(\theta_{t_1})q_{t_2}^{x_{t_1}\theta_{t_3}...\theta_{t_{|\Lambda|}}}(\theta_{t_2}) \cdot ... \cdot q_{t_{|\Lambda|}}^{x_{t_1}x_{t_2}...x_{t_{|\Lambda|-1}}}(\theta_{t_{|\Lambda|}})},$$

where

$$Z(\theta_{t_1}\theta_{t_2}\dots\theta_{t_{|\Lambda|}}) = \sum_{\alpha \in X^{\Lambda}} \frac{q_{t_1}^{\theta_{t_2}\theta_{t_3}\dots\theta_{t_{|\Lambda|}}}(\alpha_{t_1})q_{t_2}^{\alpha_{t_1}\theta_{t_3}\dots\theta_{t_{|\Lambda|}}}(\alpha_{t_2}) \cdot \dots \cdot q_{t_{|\Lambda|}}^{\alpha_{t_1}\alpha_{t_2}\dots\alpha_{t_{|\Lambda|-1}}}(\alpha_{t_{|\Lambda|}})}{q_{t_1}^{\theta_{t_2}\theta_{t_3}\dots\theta_{t_{|\Lambda|}}}(\theta_{t_1})q_{t_2}^{\alpha_{t_1}\theta_{t_3}\dots\theta_{t_{|\Lambda|}}}(\theta_{t_2}) \cdot \dots \cdot q_{t_{|\Lambda|}}^{\alpha_{t_1}\alpha_{t_2}\dots\alpha_{t_{|\Lambda|-1}}}(\theta_{t_{|\Lambda|}})},$$

 θ_{t_j} is a positivity point of \mathbf{Q}_1 at t_j , $1 \leq j \leq |\Lambda|$, and $\Lambda = \{t_1, t_2, ..., t_{|\Lambda|}\}$ is some enumeration of the elements of Λ .

Using the same reasoning as in the proof of Theorem 1, one can show that the probability distribution P_{Λ} does not depend on the enumeration of the elements in Λ and on the choice of a positivity point at each $t \in \Lambda$ (if there are several positivity points at t). Finally, it is not difficult to verify that $Q_1(P_{\Lambda}) = Q_1$. \Box

3.3 Notes on previous results

A large number of works are devoted to the question of describing finite random fields by consistent systems of conditional distributions. Let us note only those which results are close to obtained in this paper.

Apparently, the first necessary and sufficient conditions for the existence and uniqueness of a twodimensional ($|\Lambda| = 2$) joint distribution for a given system of one-point probability distributions \mathbf{Q}_1 were obtained by Abrahams and Thomas [1]. It should be noted that their conditions are not expressed in terms of the elements of the given system \mathbf{Q}_1 . At the same time, from the point of view of the statement of the problem, it is natural to impose conditions directly on the elements of \mathbf{Q}_1 .

Abrahams and Thomas' conditions were considered in several subsequent works (see, for example, Arnold and Press [2] and Berti et al. [4]). Arnold and Press [2] introduced the equivalent conditions expressed in terms of the elements of the given system \mathbf{Q}_1 in the cases $|\Lambda| = 2$ and $|\Lambda| = 3$. These conditions coincides with our consistency conditions in the considered by them cases. Nevertheless, none of the mentioned authors provide an explicit formula expressing P_{Λ} in terms of the elements of \mathbf{Q}_1 .

Such a formula was obtained by Chen [5]. This formula, after the necessary simplifications, coincides with our formula (3). However, the consistency conditions used by Chen are, in fact, a requirement for the correctness of this formula, while the validation criterion of correctness is not provided. Note also that Chen does not discuss the uniqueness of the constructed random field P_{Λ} .

Gurevich [18] gives necessary and sufficient conditions for the existence of a finite random field with a given system \mathbf{Q}_1 . He proposed the formula connecting P_{Λ} with the elements of \mathbf{Q}_1 . However, like in the work by Chen [5], the conditions proposed by Gurevich, in their essence, are the requirement of the correctness of this formula.

Regarding the uniqueness of a random field with a given system of conditional probabilities, we note that the answer was given mainly for positive systems of consistent distributions. For systems admitting zeros, the uniqueness conditions are given by Gurevich [18] (see also Arnold and Press [2]).

4 Gibbs representation of random fields

In many practical situations, the statistical data are often presented in the form of a system of conditional probability distributions. In particular, this is the case in mathematical statistical physics, where its main object, Gibbs random field, is described by means of a special system of conditional distributions, the Gibbs specification. The Gibbsian form of the conditional distribution of a random field is universal, namely, all systems of consistent conditional distributions can be represented in the Gibbs form (see [9]).

The Gibbs representation makes it very easy to construct specific examples of systems of consistent probability distributions. Moreover, the Gibbs representation allows one to describe various classes of random fields, such as, for example, the classes of Markov, stationary, and weakly dependent random fields.

The problem of Gibbsianess of a random field P_{Λ} was considered by Griffeath [17]. He showed that any positive distribution in a finite volume can be represented in the Gibbs form (5) with a vacuum potential without long-range interactions. In this case, an explicit form of the potential is written out in terms of P_{Λ} using the Möbius formula.

Note that the Gibbs representation is especially suitable when the interaction potential has a simple form, while the application of the Möbius formula leads to a very complex expression for it and may have only theoretical interest. In addition, as noted by Gandolfi and Lenarda [15], when applying the Möbius formula, the explicit dependence of the potential on the values of the spins of physical systems is lost.

In this section, using analogues of the methods proposed in [9], we present (based on the energy conservation law) the definition of a Gibbs finite random field in a much more general form (without using of the notion of potential) than the commonly accepted one. We also show that any finite positive random field is Gibbsian. Applied to Markov random fields, the proposed approach leads to a generalized version of the Hammersley–Clifford theorem.

In what follows, we will consider positive finite random fields only.

4.1 Gibbs random fields with a given potential

Note (see also [9]) that a function $P_{\Lambda}(x)$, $x \in X^{\Lambda}$, is a probability distribution on X^{Λ} if and only if it is representable in the form

$$P_{\Lambda}(x) = \frac{\exp\{-H_{\Lambda}(x)\}}{\sum\limits_{\alpha \in X^{\Lambda}} \exp\{-H_{\Lambda}(\alpha)\}}, \qquad x \in X^{\Lambda},$$
(5)

where H_{Λ} is some function on X^{Λ} . It is clear that this formula is unlikely to be useful if one does not impose certain restrictions on H_{Λ} .

In statistical physics, it is assumed that the function H_{Λ} should be a Hamiltonian (potential energy), and it is postulated that the Hamiltonian should have the following form

$$H_{\Lambda}(x) = \sum_{\emptyset \neq V \subseteq \Lambda} \Phi_V(x_V), \qquad x \in X^{\Lambda},$$
(6)

where interaction potential Φ is a family $\{\Phi_V, V \subseteq \Lambda\}$ of functions Φ_V on X^V , $V \subseteq \Lambda$. This approach makes it possible to build models of physical systems based on potential.

In the theory of finite random fields, by a Gibbs random field with a given potential Φ it is accepted to call (see, for example, [17]) a random field P_{Λ} , which is representable in the form (5) with the Hamiltonian H_{Λ} constructed by the potential Φ .

In the framework of the problem of description of random fields by systems of conditional distributions, we give an equivalent definition of a Gibbs random field P_{Λ} (with a potential Φ) in terms of $Q_1(P_{\Lambda})$, which is in accordance with the one used in the case of infinite volumes.

For a given potential $\Phi = \{\Phi_V, V \subseteq \Lambda\}$, consider the Gibbs system $\mathbf{Q}_1^{\Phi} = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of probability distributions

$$q_t^z(x) = \frac{\exp\{-H_t^z(x)\}}{\sum_{\alpha \in X^t} \exp\{-H_t^z(\alpha)\}}, \qquad x \in X^t,$$
(7)

where

$$H_t^z(x) = \sum_{J \subseteq \Lambda \setminus \{t\}} \Phi_{t \cup J}(xz_J), \qquad x \in X^t, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda.$$
(8)

Note that if H_{Λ} is defined by (6), then

$$H_t^z(x) = H_{\Lambda}(xz) - H_{\Lambda \setminus \{t\}}(z), \qquad x \in X^t, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda.$$

It is not difficult to see, that the elements of \mathbf{Q}_1^{Φ} satisfy the consistency conditions (2). Therefore, according to Theorem 1, \mathbf{Q}_1^{Φ} uniquely determines a random field P_{Λ} such that $\mathbf{Q}_1(P_{\Lambda}) = \mathbf{Q}_1^{\Phi}$. Hence, we come to the following definition of the Gibbs random field with a potential Φ formulated in terms of the elements of $\mathbf{Q}_1(P_{\Lambda})$ only.

A random field P_{Λ} will be called a *Gibbs random field with potential* $\Phi = \{\Phi_V, V \subseteq \Lambda\}$ if $Q_1(P_{\Lambda}) = \mathbf{Q}_1^{\Phi}$ where \mathbf{Q}_1^{Φ} is the Gibbs system constructed by the potential Φ .

It is clear that if P_{Λ} is a Gibbs random field with potential Φ in the conventional sense, then its one point conditional distributions have the Gibbs form (7). Conversely, let the elements of the system $Q_1(P_{\Lambda})$ have the form (7). According to the formula (4), we can write

$$P_{\Lambda}(x) = \left(\sum_{\alpha \in X^{\Lambda}} \prod_{j=1}^{n} \frac{Q_{t_{j}}^{(\alpha x)_{j}}(\alpha_{j})}{Q_{t_{j}}^{(\alpha x)_{j}}(x_{j})}\right)^{-1} = \left(\sum_{\alpha \in X^{\Lambda}} \prod_{j=1}^{n} \frac{\exp\{-H_{t_{j}}^{(\alpha x)_{j}}(\alpha_{j})\}}{\exp\{-H_{t_{j}}^{(\alpha x)_{j}}(x_{j})\}}\right)^{-1}$$
$$= \left(\sum_{\alpha \in X^{\Lambda}} \exp\left\{\sum_{j=1}^{n} \left(H_{t_{j}}^{(\alpha x)_{j}}(x_{j}) - H_{t_{j}}^{(\alpha x)_{j}}(\alpha_{j})\right)\right\}\right)^{-1},$$

where $\Lambda = \{t_1, t_2, ..., t_n\}$ is some enumeration of the points in Λ , $n = |\Lambda|$. Since

$$\sum_{j=1}^{n} \left(H_{t_j}^{(\alpha x)_j}(x_j) - H_{t_j}^{(\alpha x)_j}(\alpha_j) \right) = H_{\Lambda}(x) - H_{\Lambda}(\alpha),$$

finally we obtain

$$P_{\Lambda}(x) = \left(\sum_{\alpha \in X^{\Lambda}} \exp\{H_{\Lambda}(x) - H_{\Lambda}(\alpha)\}\right)^{-1} = \frac{\exp\{-H_{\Lambda}(x)\}}{\sum_{\alpha \in X^{\Lambda}} \exp\{-H_{\Lambda}(\alpha)\}}, \qquad x \in X^{\Lambda}.$$

Given the above definitions of a Gibbs random field are essentially based on the fact that the Hamiltonian is the sum of the interaction potentials. Such a representation of the Hamiltonian cannot be considered as its definition since in the case of this approach, the properties of the Hamiltonian are not postulated but derived from the properties of the potential. In [9], based on the energy conservation law, an axiomatic definition of the Hamiltonian was given by means of its intrinsic properties. This definition is based on the concept of a transition energy field introduced in [9].

Applying the approach developed in [9] to finite random fields, below we give the definition of Gibbs random field without using the notion of interaction potential. Note that some of our arguments literally repeat those used in the case of infinite volumes. This is done for the sake of completeness.

4.2 Axiomatic definition of Hamiltonian and Gibbs random fields

First of all, we give the following (axiomatic) definition of the Hamiltonian (in finite volume) which does not use the notion of potential (see [9]).

A family $H_1 = \{H_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of functions H_t^z on X^t parameterized by boundary conditions $z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda$, we call *(one-point) Hamiltonian* in Λ if its elements satisfy the following conditions: for all $t, s \in \Lambda$ and $x, u \in X^t, y, v \in X^s, z \in X^{\Lambda \setminus \{t,s\}}$, it holds

$$H_t^{zy}(x) + H_s^{zx}(v) + H_t^{zv}(u) + H_s^{zu}(y) = H_t^{zy}(u) + H_s^{zu}(v) + H_t^{zv}(x) + H_s^{zx}(y).$$
(9)

The basic example of a one-point Hamiltonian is a Hamiltonian $H_1^{\Phi} = \{H_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ constructed by the potential $\Phi = \{\Phi_V, V \subseteq \Lambda\}$ (see formula (8)). Another example is the Hamiltonian $H_1^{\theta}(P_{\Lambda}) = \{H_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ constructed by a random field P_{Λ} and a fixed configuration $\theta \in X^{\Lambda}$ according to the formula

$$H_t^z(x) = \ln \frac{P_{\Lambda}(xz)}{P_{\Lambda}(\theta_t z)}, \qquad x \in X^t, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda.$$

It is easy to check that the elements both of H_1^{Φ} and $H_1^{\theta}(P_{\Lambda})$ satisfy the conditions (9).

Consistency conditions (9) have a very simple physical interpretation. First note that they can be written in the following form

$$H_t^{zy}(u) - H_t^{zy}(x) + H_s^{zu}(v) - H_s^{zu}(y) = H_s^{zx}(v) - H_s^{zx}(y) + H_t^{zv}(u) - H_t^{zv}(x)$$

For each $t \in \Lambda$, consider the function

$$\Delta_t^z(x,u) = H_t^z(u) - H_t^z(x), \qquad x, u \in X^t, z \in X^{\Lambda \setminus \{t\}},\tag{10}$$

which represents an amount of energy needed under fixed boundary condition z to change the state of the system at point t from x to u. It is clear that for each $t \in \Lambda$ and $z \in X^{\Lambda \setminus \{t\}}$, the function Δ_t^z satisfy the relation

$$\Delta_t^z(x,u) = \Delta_t^z(x,\alpha) + \Delta_t^z(\alpha,u), \qquad x, u, \alpha \in X^t,$$
(11)

which is in full compliance with the energy conservation law. Conditions (9) in terms of Δ_t^z take the following form: for all $t, s \in \Lambda$ and $x, u \in X^t$, $y, v \in X^s$, $z \in X^{\Lambda \setminus \{t,s\}}$, it holds

$$\Delta_t^{zy}(x,u) + \Delta_s^{zu}(y,v) = \Delta_s^{zx}(y,v) + \Delta_t^{zv}(x,u).$$
(12)

This, in turn, can be interpreted as follows. Suppose it is necessary to move a physical system from the state xyz to the state uvz, where $x, u \in X^t$, $y, v \in X^s$, $z \in X^{\Lambda \setminus \{t,s\}}$, $t, s \in \Lambda$. This can be done in two ways. First, change the state of the system at the point t from x to u under the boundary condition zy,

and then at the point s from y to v already under the boundary condition zu. The energy expended in this case is $\Delta_t^{zy}(x, u) + \Delta_s^{zu}(y, v)$. One can proceed also as follows: first, starting from the point s, change the state from y to v under the boundary condition zx, and then, under the boundary condition zv, change it at the point t from x to u expending the energy $\Delta_s^{zx}(y, v) + \Delta_t^{zv}(x, u)$. Naturally, the same amount of energy must be spent in both cases.

Note also, that from the physical point of view, it is more natural to consider the difference of energies instead of the energies themselves. Thus, following [9], we introduce the notion of transition energy field in finite volume.

A family $\Delta_1 = {\Delta_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda}$ of functions Δ_t^z on pairs of configurations from X^t and parameterized by boundary conditions $z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda$, we call a *(one-point)* transition energy field in Λ if its elements satisfy the consistency conditions (11) and (12).

The fundamental connection between the consistent system \mathbf{Q}_1 and the transition energy field $\boldsymbol{\Delta}_1$ is established in the theorem below. Since the proof is similar to the proof of Theorem 2.3 in [9], we give it briefly.

Theorem 3. A family of functions $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ forms a system of positive one-point probability distributions consistent in the sense (2) if and only if its elements have a Gibbs form

$$q_t^z(x) = \frac{\exp\{\Delta_t^z(x, u)\}}{\sum\limits_{\alpha \in X^t} \exp\{\Delta_t^z(\alpha, u)\}}, \qquad x \in X^t,$$
(13)

where $u \in X^t$ and $\Delta_1 = \{\Delta_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ is a one-point transition energy field. Herewith, the transition energy field Δ_1^{Λ} is uniquely determined by \mathbf{Q}_1 .

Proof. Let the elements of the system $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of positive probability distributions satisfy the consistency conditions (2). For each $t \in \Lambda$ and $z \in X^{\Lambda \setminus \{t\}}$, put

$$\Delta_t^z(x,u) = \ln \frac{q_t^z(x)}{q_t^z(u)}, \qquad x, u \in X^t.$$
(14)

It is easy to verify that the elements of the system $\Delta_1 = \{\Delta_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ satisfy the conditions (11) and (12), and thus, Δ_1^{Λ} is a transition energy field. Let now $\Delta_1 = \{\Delta_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ be a transition energy field. For each $t \in \Lambda$ and $z \in X^{\Lambda \setminus \{t\}}$,

Let now $\Delta_1 = {\Delta_t^z, z \in X^{\Lambda \setminus {t}}, t \in \Lambda}$ be a transition energy field. For each $t \in \Lambda$ and $z \in X^{\Lambda \setminus {t}}$, put

$$q_t^z(x) = \frac{\exp\{\Delta_t^z(x, u)\}}{\sum_{\alpha \in X} \exp\{\Delta_t^z(\alpha, u)\}}, \qquad x, u \in X^t.$$

First, note that due to (11), the values of q_t^z do not depend on the choice of u. Let us verify that the elements of the family $\{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ satisfy the consistency conditions (2). This statement follows from the chain of equalities

$$\ln \frac{q_t^{zy}(x)}{q_t^{zy}(u)} \cdot \frac{q_s^{zu}(y)}{q_s^{zu}(v)} = \Delta_t^{zy}(x,u) + \Delta_s^{zu}(y,v) = \Delta_s^{zx}(y,v) + \Delta_t^{zv}(x,u) = \ln \frac{q_s^{zx}(y)}{q_s^{zx}(v)} \cdot \frac{q_t^{zv}(x)}{q_t^{zv}(u)},$$

 $x, u \in X^t, y, v \in X^s, z \in X^{\Lambda \setminus \{t,s\}}, t, s \in \Lambda$, which is true by virtue of the consistency conditions (11) of the elements of Δ_1^{Λ} .

We propose the following definition of a Gibbs finite random field, based on the introduced notion of the transition energy field.

A random field P_{Λ} we call *Gibbsian* if the elements of $Q_1(P_{\Lambda})$ can be represented in the Gibbs form

$$Q_t^z(x) = \frac{\exp\{\Delta_t^z(x, u)\}}{\sum\limits_{\alpha \in X^t} \exp\{\Delta_t^z(\alpha, u)\}}, \qquad x \in X^t,$$

where $u \in X^t$ and $\Delta_1 = {\Delta_t^z, z \in X^{\Lambda \setminus {t}}, t \in \Lambda}$ is the transition energy field.

It is easy to see that the definition of a Gibbs random field based on the notion of transition energy field includes all the definitions given above.

The direct corollary of Theorem 3 is the following result.

Theorem 4. Any finite random field is Gibbsian.

Proof. Since the elements of the one-point conditional distribution $Q_1(P_{\Lambda})$ of any random field P_{Λ} satisfy the consistency conditions (2), by Theorem 3 they have the Gibbs form (13) with uniquely determined transition energy field $\Delta_1 = \{\Delta_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$. Herewith,

$$\Delta_t^z(x,u) = \ln \frac{Q_t^z(x)}{Q_t^z(u)} = \ln \frac{P_\Lambda(xz)}{P_\Lambda(uz)}, \qquad x, u \in X^t.$$

Thus, there is a one-to-one correspondence between the positive random field P_{Λ} and the one-point transition energy field Δ_1 .

From Theorem 13, it directly follows the validity of the next result.

Theorem 5. The family of functions $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ forms a system of positive one-point probability distributions satisfying the consistency conditions (2) if and only if its elements have a Gibbs form

$$q_t^z(x) = \frac{\exp\{-H_t^z(x)\}}{\sum\limits_{\alpha \in X^t} \exp\{-H_t^z(\alpha)\}}, \qquad x \in X^t, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda,$$

where $H_1 = \{H_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ is some one-point Hamiltonian.

Unlike the transition energy field, the Hamiltonian H_1^{Λ} for a consistent system \mathbf{Q}_1 (and, therefore, compatible with it random field) is determined not uniquely but up to an additive function on boundary conditions. Nevertheless, all such Hamiltonians lead to the same consistent system \mathbf{Q}_1 . This remark allows us to give the following definition of a Gibbs random field in a finite volume.

A random field P_{Λ} is called *Gibbsian* if the elements of its one-point conditional distribution $Q_1(P_{\Lambda})$ are representable in a Gibbs form

$$Q_t^z(x) = \frac{\exp\{-H_t^z(x)\}}{\sum\limits_{\alpha \in X^t} \exp\{-H_t^z(\alpha)\}}, \qquad x \in X^t,$$

where $H_1 = \{H_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ is some one-point Hamiltonian.

4.3 Markov finite random fields

Most often, the question of Gibbsianess was considered for Markov finite random fields. The main result here is considered to be the Hammersley–Clifford theorem on the equivalence of Markov and Gibbs random fields with a pair interaction potential of the nearest neighbors, first presented in an unpublished paper [20]. The proof of this theorem has been repeatedly presented under various conditions both on Λ and on the state space X. Let us mention, for example, works by Grimmet [19], Preston [26], Spitzer [29], and Besag [3]. However, as it will be shown below, the Hammersley–Clifford theorem is a direct consequence of a much more general theorem (see Theorem 6).

The definition of Markov random fields is based on the concept of a neighborhood system. A family $\partial = \{\partial t, t \in \Lambda\}$ of sets $\partial t \subset \Lambda$ is called a neighborhood system in Λ if $t \notin \partial t$ and $s \in \partial t$ if and only if $t \in \partial s, t, s \in \Lambda$.

A random field P_{Λ} is called *Markov random field* (with respect to ∂) if its conditional distributions satisfy the following property: for any $t \in \Lambda$ and $z \in X^{\Lambda \setminus \{t\}}$, it holds

$$Q_t^z(x) = Q_t^{z_{\partial t}}(x), \qquad x \in X^t.$$

It is not difficult to verify that a random field P_{Λ} will be a Markov random field if and only if for each $t \in \Lambda$ and any $x, u \in X^t$, $z \in X^{\partial t}$, $y, v \in X^{\Lambda \setminus (t \cup \partial t)}$, it holds

$$Q_t^{zy}(x) = Q_t^{zv}(x).$$

Let us now present conditions on the transition energy field, under which the corresponding finite random field is Markov.

Proposition 1. A random field P_{Λ} is a Markov random field if and only if the elements of corresponding transition energy field $\mathbf{\Delta}_1 = \{\Delta_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ satisfy the following conditions: for all $t \in \Lambda$ and any $x, u \in X^t$, $z \in X^{\partial t}$, $y, v \in X^{\Lambda \setminus (t \cup \partial t)}$, it holds

$$\Delta_t^{zy}(x,u) = \Delta_t^{zv}(x,u), \qquad x, u \in X^t.$$
(15)

Proof. Let $Q_1(P_{\Lambda})$ be a system of one-point conditional distributions of the Markov random field P_{Λ} . Then for all $t \in \Lambda$ and $x, u \in X^t$, $z \in X^{\hat{\partial}t}$, $y, v \in X^{\Lambda \setminus (t \cup \partial t)}$, we have

$$\Delta_t^{zy}(x,u) = \ln \frac{Q_t^{zy}(x)}{Q_t^{zy}(u)} = \ln \frac{Q_t^{zv}(x)}{Q_t^{zv}(u)} = \Delta_t^{zv}(x,u).$$

Let now $\Delta_1 = {\Delta_t^z, z \in X^{\Lambda \setminus {t}}, t \in \Lambda}$ be a one-point transition energy field which elements satisfy conditions (15), and let P_{Λ} be a corresponding to it random field with one-point conditional distribution $Q_1(P_\Lambda)$. Then for all $t \in \Lambda$ and $x, u \in X^t, z \in X^{\partial t}, y, v \in X^{\Lambda \setminus (\{t\} \cup \partial t)}$, we have

$$Q_t^{zy}(x) = \frac{\exp\{\Delta_t^{zy}(x,u)\}}{\sum\limits_{\alpha \in X^t} \exp\{\Delta_t^{zy}(\alpha,u)\}} = \frac{\exp\{\Delta_t^{zv}(x,u)\}}{\sum\limits_{\alpha \in X^t} \exp\{\Delta_t^{zv}(\alpha,u)\}} = Q_t^{zv}(x)$$

Hence, P_{Λ} is a Markov random field.

Therefore, the following statement is true.

Theorem 6. Any Markov random field P_{Λ} is Gibbsian with the transition energy field Δ_1 satisfying conditions (15). Conversely, if the elements of the transition energy field Δ_1 of a (Gibbs) random field P_{Λ} satisfy the conditions (15), then P_{Λ} is a Markov random field.

It is not difficult to see that the well-known Hammersley–Clifford theorem on the equivalence of Markov finite random field and Gibbs finite random field with a pair potential of the nearest neighbors interactions is a particular case of Theorem 6. Indeed, according to Theorem 4, Markov finite random fields are Gibbsian (in this case, there is no need to construct a potential). Conversely, let a neighborhood system $\partial = \{\partial t, t \in \Lambda\}$ in Λ be given, and let $\Phi = \{\Phi_V, V \subset \Lambda\}$ be a pair potential of the nearest neighbors interactions, that is, $\Phi_{t\cup V} \neq 0$ only for $V \subseteq \partial t$, $t \in \Lambda$. For all $t \in \Lambda$ and $x, u \in X^t$, $z \in X^{\Lambda \setminus \{t\}}$, put

$$\Delta_t^z(x,u) = \sum_{\emptyset \neq J \subseteq \Lambda} \left(\Phi_{t \cup J}(uz_J) - \Phi_{t \cup J}(xz_J) \right),$$

and let P_{Λ}^{Φ} be a (Gibbs) random field corresponding to the transition energy field $\Delta_1 \Phi = \{\Delta_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$. Since the conditions (15) are satisfied, P_{Λ}^{Φ} is a Markov random field. Let us also note that if $H_1 = \{H_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ is a Hamiltonian corresponding to a Markov

random field P_{Λ} , then its elements satisfy the following conditions

$$H_t^{zy}(u) - H_t^{zy}(x) = H_t^{zv}(u) - H_t^{zv}(x)$$


for all $t \in \Lambda$ and $x, u \in X^t$, $z \in X^{\partial t}$, $y, v \in X^{\Lambda \setminus (t \cup \partial t)}$.

Concluding remarks 5

Thus, the methods developed for infinite random fields can be used to study random fields in finite volumes.

The main object considered in the paper is the system \mathbf{Q}_1 of one-point probability distributions parameterized by boundary conditions and consistent in the sense (2). We showed that there is a one-toone correspondence between the system \mathbf{Q}_1 and compatible with it finite random field P_{Λ} . Further, we showed that there is a one-to-one correspondence between \mathbf{Q}_1 and the one-point transition energy field Δ_1 . These connections can schematically be presented by the following commutative diagram.

| J. | | |
|----|--|--|



One-to-one correspondence between P_{Λ} and Δ_1 allows one to give the definition of a finite Gibbs random field, which does not involve the notion of potential. From the condition (11) on the elements of Δ_1 , it follows that each function Δ_t^z can be represented in the form

$$\Delta_t^z(x, u) = H_t^z(u) - H_t^z(x), \qquad x, u \in X^t,$$

where H_t^z can be interpreted as a potential energy (Hamiltonian) of the system at point $t \in \Lambda$ under boundary condition $z \in X^{\Lambda \setminus \{t\}}$. The consistency conditions (12) impose certain conditions on the functions H_t^z , which lead to the presented axiomatic definition of a one-point Hamiltonian H_1 in a finite volume Λ . According to this definition, for any random field P_{Λ} , the set $H_1^{\theta}(P_{\Lambda})$ of functions

$$H_t^z(x) = \ln \frac{P_{\Lambda}(xz)}{P_{\Lambda}(\theta_t z)}, \qquad x \in X^t, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda,$$

where $\theta \in X^{\Lambda}$ is some fixed configuration, is a Hamiltonian, and there is no need to represent H_t^z as a sum of potentials.

The approach used made it possible to prove the fact that all finite random fields are Gibbsian.

Let us emphasize that obtained results do not depend on the structure of the finite set Λ . Its structure may be essential when considering special classes of finite random fields, for example, Markov finite random fields.

Our results carry over in a natural way to the case of infinite (both countable and continuous) measurable spaces X. Let, for example, (X, \Im, μ) be a complete separable metric space, where X is a non-empty set, \Im is a σ -algebra generated by open subsets of X, and μ is not identically zero countably additive measure.

Consider a probability P_{Λ} (random field) on the product space $(X^{\Lambda}, \Im^{\Lambda}, \mu_{\Lambda})$ which has a positive density p_{Λ} with respect to the measure μ_{Λ} . For such random field, the system $Q_1(P_{\Lambda}) = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of its conditional densities is defined. In the case under consideration, the following result is an analogue of Theorems 1 and 3.

Theorem 7. For a given system $\mathbf{Q}_1 = \{q_t^z, z \in X^{\Lambda \setminus \{t\}}, t \in \Lambda\}$ of positive one-point probability densities parameterized by boundary conditions, there exists a unique probability density p_{Λ} compatible with it if and only if the elements of \mathbf{Q}_1 satisfy the consistency conditions (2) and for any $u \in X^{\Lambda}$,

$$Z_{\Lambda}(u) = \int_{X^{\Lambda}} \prod_{j=1}^{n} \frac{q_{t_{j}}^{(\alpha u)_{j}}(\alpha_{j})}{q_{t_{j}}^{(\alpha u)_{j}}(u_{j})} \mu_{\Lambda}(d\alpha) < \infty,$$

where $\Lambda = \{t_1, t_2, ..., t_n\}$ is some enumeration of points in Λ , $n = |\Lambda|$. In this case, the elements of \mathbf{Q}_1 have the Gibbs form with uniquely determined transition energy field Δ_1^{Λ} .

Note also, that the obtained results can be extended on the case when for the construction of the space $(X^{\Lambda}, \mathfrak{S}^{\Lambda}, \mu_{\Lambda})$, to each point $t \in \Lambda$ corresponds a space $(X^t, \mathfrak{S}^t, \mu_t)$ which is not a copy of the same space (X, \mathfrak{S}, μ) .

Funding. The work was supported by the Science Committee of RA in the frames of the research project 21AG-1A045.

References

 ABRAHAMS, J. and THOMAS, J.B. (1984). A note on the characterization of bivariate densities by conditional densities. Commun. Stat. Theory Methods 13 (3) 395–400.

- [2] ARNOLD, B.C. and PRESS, S.J. (1989). Compatible Conditional Distributions. J. Am. Stat. Assoc. textbf84 (405) 152-156.
- [3] BESAG, J. (1974). Spatial interaction and the statistical analysis of lattice systems. J. R. Stat. Soc., Ser. B (Methodological) 36 (2) 192–236.
- [4] BERTI, P., DREASSI, E. and RIGO, P. (2014). Compatibility results for conditional distributions. J. Multivar. Anal. 125 190–203.
- [5] CHEN, H.Y. (2010). Compatibility of conditionally specified models. Stat. Probab. Lett. 80 670–677.
- [6] DACHIAN, S. and NAHAPETIAN, B.S. (1998). An approach towards description of random fields. Preprint 98-5, Laboratoire de Statistique et Processus, Université du Maine.
- [7] DACHIAN, S. and NAHAPETIAN, B.S. (2001). Description of random fields by means of one-point conditional distributions and some applications. Markov Processes Relat. Fields 7 193–214.
- [8] DACHIAN, S. and NAHAPETIAN, B.S. (2004). Description of specifications by means of probability distributions in small volumes under condition of very week positivity. J. Stat. Phys. 117 281–300.
- [9] DACHIAN, S. and NAHAPETIAN, B.S. (2019). On the relationship of energy and probability in models of classical statistical physics. Markov Processes Relat. Fields 25 649–681.
- [10] DALALYAN, A. and NAHAPETIAN, B.S. (2011). Description of random fields by means of one-point finite conditional distribution. J. Contemp. Math. Anal. Arm. Acad. Sci. 46 (2) 113–119.
- [11] DOBRUSHIN, R.L. (1968). The description of a random field by means of conditional probabilities and conditions of its regularity. Theory Probab. Appl. 13 (2) 197–224.
- [12] DOBRUSHIN, R.L. (1968). Gibbs random fields for lattice systems with pair-wise interaction. Funct. Anal. and Appl. 2 292–301.
- [13] DOBRUSHIN, R.L. (1968). The problem of uniqueness of a gibbsian random field and the problem of phase transitions. Funct. Anal. Appl. 2 (4) 302–312.
- [14] DOBRUSHIN, R.L. (1980). Gaussian random fields Gibbsian point of view. In Multicomponent random systems, eds. Dobrushin, R.L. and Sinai, Ya.G., New York: M. Dekker, 119–151.
- [15] GANDOLFI, A. and LENARDA, P. (2016). A note on Gibbs and Markov random fields with constraints and their moments. Math. Mech. Complex Syst. 4 (3-4) 407–422.
- [16] GEORGII, H.-O. (1988). Gibbs Measures and Phase Transitions, De Gruyter, Berlin.
- [17] GRIFFEATH, D. (1976). Introduction to Random Fields. In: Denumerable Markov Chains. Graduate Texts in Mathematics 40. Springer, New York, NY.
- [18] GUREVICH, B. M. (1992). On joint distribution of random variables with given cross conditional distributions: Discrete case. Theory Probab. Appl. 36 (2) 371–375.
- [19] GRIMMETT, G. R. (1973). A theorem about random fields. Bull. London Math. Soc. 5 (1) 81–84.
- [20] HAMMERSLEY, J. M. and CLIFFORD, P. (1971). Markov fields on finite graphs and lattices. Unpublished.
- [21] KAISER, M.S. and CRESSIE, N. (2000). The construction of multivariate distributions from Markov random fields. J. Multivar. Anal. 73 (2) 199–220.
- [22] KHACHATRYAN, L.A. and NAHAPETIAN, B.S. (2020). On direct and inverse problems in the description of lattice random fields. Proceedings of the XI international conference stochastic and analytic methods in mathematical physics. Lectures in pure and applied mathematics 6 107–116. Universitätsverlag Potsdam.

- [23] KOZLOV, O.K. (1974). Gibbs description of a system of random variables. Problems Inform. Transmission 10 258–265.
- [24] KÜNSCH, H. (1981). Thermodynamics and statistical analysis of Gaussian random fields. Z. Wahrscheinlichkeitstheorie verw Gebiete 58, 407–421.
- [25] LANFORD, O.E. and RUELLE, D. (1969). Observables at infinity and states with short range correlations in statistical mechanics. Commun. Math. Phys. 13 194–215.
- [26] PRESTON, C. J. (1973). Generalized Gibbs states and Markov random fields. Adv. Appl. Probab. 5 (2) 242–261.
- [27] SINAI, YA.G. (1980). Theory of Phase Transitions. Rigorous results (in Russian). Nauka, Moscow.
- [28] SULLIVAN, W.G. (1973). Potentials for almost Markovian random fields. Commun. Math. Phys. 33 61–74.
- [29] SPITZER, F. (1971). Markov random fields and Gibbs ensembles. The American Mathematical Monthly 78 (2) 142–154.