

ESTIMATES IN L_p FOR SOLUTIONS OF SPDES WITH COEFFICIENTS IN MORREY CLASSES

N.V. KRYLOV

ABSTRACT. For solutions of a certain class of SPDEs in divergence form we present some estimates of their L_p -norms and the L_p -norms of their first-order derivatives. The main novelty is that the low-order coefficients are supposed to belong to certain Morrey classes instead of L_p -spaces. Our results are new even if there are no stochastic terms in the equation.

1. INTRODUCTION

In this paper we come back to an old problem of estimating the L_p -norm in x of solutions $u_t(x)$ of Itô stochastic partial differential equations of second order. The first such estimates appeared in [9] in 1977 and were achieved by using Itô's formula for $\|u_t\|_{L_p}^p$ and integrating by parts. This method of obtaining such estimates was generalized, extended, and applied in many papers of which we cite, probably, the most recent [2], [3], [12], and [13] containing vast lists of references. The closest to the present article are some computations in [8] which is about strong solutions of Itô stochastic equations rather than about SPDEs. There, as in the present article, the main emphasis is on the drift coefficients from a Morrey class. In this connection it is also worth mentioning [5] and some references therein where methods different from integration by parts are used to treat strong solutions of Itô stochastic equations with singular drift. We apply integration by parts to SPDEs which may be just nonrandom usual parabolic equations. Even in this case, as far as the author can tell, the presented results are new although similar results for the *elliptic* equations can be found in [11].

Let d, d_1 be integers, $d \geq 3$, \mathbb{R}^d be a d -dimensional Euclidean space of points $x = (x^1, \dots, x^d)$, and

$$\mathbb{R}^{d+1} = \{(t, x) : t \in \mathbb{R}, x \in \mathbb{R}^d\}.$$

Let $\mathcal{B}(\mathbb{R}^d)$ be the Borel σ -field on \mathbb{R}^d , $L_p = L_p(\mathbb{R}^d)$.

2020 *Mathematics Subject Classification.* 60H15, 35R60.

Key words and phrases. Stochastic PDEs, Morrey classes, singular coefficients.

Let (Ω, \mathcal{F}, P) be a complete probability space equipped with a filtration $(\mathcal{F}_t)_{t \geq 0}$ of complete σ -fields $\mathcal{F}_t \subset \mathcal{F}$. We suppose that on our probability space we are given a d_1 -dimensional Wiener process w_t , which is a Wiener process relative to the filtration $(\mathcal{F}_t)_{t \geq 0}$. By \mathcal{P} we denote the predictable σ -field on $\Omega \times (0, \infty)$ (we follow the terminology in [4]). Fix a bounded stopping time τ .

Assume that on $\Omega \times (0, \infty) \times \mathbb{R}^d$ we are given $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable \mathbb{R}^d -valued functions $\sigma_t^k(x) = (\sigma_t^{ik}(x))$, $k = 1, \dots, d_1$, $\beta_t(x) = (\beta_t^i(x))$, $b_t(x) = (b_t^i(x))$, $\mathbf{f}_t(x) = (\mathbf{f}_t^i(x))$, $d \times d$ -matrix valued functions $a_t(x) = (a_t^{ij}(x))$, \mathbb{R}^{d_1} -valued functions $\nu_t(x) = (\nu_t^k(x))$, $g_t(x) = (g_t^k(x))$, and real-valued functions $c_t(x)$, $f_t(x)$. Introduce

$$L_t u_t = D_i(a_t^{ij} D_j u_t + \beta_t^i u_t) + b_t^i D_i u_t + c_t u_t, \quad M_t^k u_t = \sigma_t^{ik} D_i u_t + \nu_t^k u_t,$$

where and everywhere below the common summation convention is used and $D_i = \partial / \partial x^i$.

Our goal is to investigate the equation

$$du_t = (L_t u_t + D_i \mathbf{f}_t^i + f_t) dt + (M_t^k u_t + g_t^k) dw_t^k, \quad t \leq \tau, \quad (1.1)$$

with initial condition

$$u_0 = \Phi. \quad (1.2)$$

Remark 1.1. In the literature very popular conditions on b (and β) is that $b \in L_{p_0, q_0}(T)$, that is, for $T \in (0, \infty)$,

$$\left(\int_0^T \left(\int_{\mathbb{R}^d} |b_t(x)|^{p_0} dx \right)^{q_0/p_0} dt \right)^{1/q_0} < \infty \quad (1.3)$$

with $p_0, q_0 \in [2, \infty]$ satisfying

$$\frac{d}{p_0} + \frac{2}{q_0} = 1.$$

Observe that, if $p_0 > d$ and we take an arbitrary constant \hat{N} and introduce

$$\lambda(t) = \hat{N} \left(\int_{\mathbb{R}^d} |b_t(x)|^{p_0} dx \right)^{1/(p_0-d)},$$

then for (the notation will be explained later)

$$b_t^M(x) = b_t(x) I_{|b_t(x)| \geq \lambda(t)}$$

we have

$$\int_{B_\rho} |b_t^M(x)|^d dx \leq \lambda^{d-p_0}(t) \int_{B_\rho} |b_t(x)|^{p_0} dx \leq N(d) \hat{N}^{d-p_0} \rho^{-d},$$

where B_ρ is a ball of radius $\rho \in (0, \infty)$,

$$\int_{B_\rho} \dots = \frac{1}{|B_\rho|} \int_{B_\rho} \dots, \quad |B_\rho| = \text{Vol } B_\rho.$$

Here $N(d)\hat{N}^{d-p_0}$ can be made arbitrarily small if we choose \hat{N} large enough. In addition, for $b_t^B = b_t - b_t^M$ we have $|b_t^B| \leq \lambda(t)$ and

$$\|b^B\|_{L^\infty,2(T)}^2 \leq \int_0^T \lambda^2(t) dt = \hat{N}^2 \int_0^T \left(\int_{\mathbb{R}^d} |b_t(x)|^{p_0} dx \right)^{q_0/p_0} dt < \infty.$$

This shows that the assumptions we are going to impose on b are weaker than (1.3) if $p_0 > d$.

If $p_0 = d$ (and $q_0 = \infty$) our assumptions are weaker if (1.3) is combined with the requirement that the family $|b_t|^d, t \in (0, \infty)$, be uniformly integrable (say, case of b independent of t).

Recall that W_p^n are Sobolev spaces of functions u on \mathbb{R}^d such that u and all its (generalized) derivatives of order $\leq n$ belong to L_p . The norm in W_p^n is introduced in a natural way.

Definition 1.2. A W_p^1 -valued function u , defined on the stochastic interval $(0, \tau]$, is called a solution of (1.1)-(1.2) if u is predictable on $(0, \tau]$,

$$\int_0^\tau \|u_t\|_{W_p^1}^p dt < \infty,$$

(a.s.) and for each $\varphi \in C_0^\infty(\mathbb{R}^d)$ for almost all $\omega \in \Omega$

$$\begin{aligned} (u_t, \varphi) &= (\Phi, \varphi) + \int_0^t (\sigma_s^{ik} D_i u_s + \nu_s^k u_s + g_s^k, \varphi) dw_s^k \\ &+ \int_0^t \{(b_s^i D_i u_s + c_s u_s + f_s, \varphi) - (a_s^{ij} D_j u_s + \beta_s^i u_s + \mathbf{f}_s^i, D_i \varphi)\} ds \end{aligned} \quad (1.4)$$

for all $t \in [0, \tau(\omega)]$, where (u, v) denotes the integral over \mathbb{R}^d of uv .

Our standing assumption is that equation (1.1) is uniformly nondegenerate. Set $\alpha_t(x) = (\alpha_t^{ij}(x))$, where $\alpha_t^{ij} = \sigma_t^{ik} \sigma_t^{jk}$. Note that we do not suppose that a_t is symmetric.

Assumption 1.3. There is a constant $\delta > 0$ such that for all values of arguments and $\lambda \in \mathbb{R}^d$

$$|a_t| \leq \delta^{-1}, \quad (2a_t^{ij} - \alpha_t^{ij})\lambda^i \lambda^j \geq \delta |\lambda|^2. \quad (1.5)$$

Another standing assumption concerns free terms. We fix $p \in [2, d)$ ($d \geq 3$) and recall that $L_p = L_p(\mathbb{R}^d)$.

Assumption 1.4. We have

$$E \int_0^\tau (\|f_t\|_{L^p}^p + \|f_t\|_{L^p}^p + \|g_t\|_{L^p}^p) dt < \infty. \quad (1.6)$$

The assumptions on the coefficients of L and M^k is more delicate. Fix $r \in (p, d]$, $R_0 \in (0, \infty)$, and let \mathbb{B}_ρ denote the set of balls in \mathbb{R}^d with radius ρ .

Definition 1.5. We call a real- or vector- or else tensor-valued function $f_t(x)$ given on $\Omega \times (0, \infty) \times \mathbb{R}^d$ *admissible* if it is represented as $f_t = f_t^M + f_t^B$ (“Morrey part” of f plus its “bounded part”) with $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable f_t^M, f_t^B such that there exists a *constant* $\hat{f} < \infty$ for which

$$\left(\int_B |f_t^M|^r dx \right)^{1/r} \leq \hat{f} \rho^{-1},$$

whenever $B \in \mathbb{B}_\rho$ and $\rho \leq R_0$, and there exists a *predictable* $\bar{f}_t \geq \sup_{x \in \mathbb{R}^d} |f_t^B(x)|$ such that the integral

$$\int_0^\tau |\bar{f}_t|^2 dt$$

is a bounded function on Ω .

Assumption 1.6. $\beta, b, \zeta := c_+^{1/2}, \nu$ are admissible.

Here is our first main result about an a priori estimate in which for convenience of applications we introduce a measurable, \mathcal{F}_t -adapted process $\mu_t \geq 0$ such that the integral

$$\int_0^\tau \mu_t dt$$

is a bounded function on Ω .

Theorem 1.7. *Under the above assumptions suppose that a solution u_t of (1.1)-(1.2) exists. Then with probability one $u_{t \wedge \tau}$ is continuous as an L^p -valued function and there exists a constant $N_0 = N_0(p, \delta, d, r)$ such that if*

$$N_0(\hat{b} + \hat{\beta} + \hat{\zeta} + \hat{\nu}) \leq 1, \quad (1.7)$$

then for any stopping time $\gamma \leq \tau$

$$\begin{aligned} & E e^{-\phi_\gamma} \int_{\mathbb{R}^d} |u_t|^p dx + (\delta/16) E \int_0^\gamma e^{-\phi_s} \int_{\mathbb{R}^d} |DU_s|^2 dx ds \\ & \leq E \int_{\mathbb{R}^d} |u_0|^p dx + N E \int_0^\gamma e^{-\phi_s} \int_{\mathbb{R}^d} (|f_t|^p + |f_t|^p + |g_t|^p - \mu_t |u_t|^p) dx ds, \end{aligned} \quad (1.8)$$

$$\begin{aligned}
 E \sup_{t \leq \tau} e^{-\Phi_t} \int_{\mathbb{R}^d} |u_t|^p dx &\leq N \int_{\mathbb{R}^d} |u_0|^p dx \\
 + NE \int_0^\tau e^{-\Phi_s} \int_{\mathbb{R}^d} (|f_s|^p + |g_s|^p) dx ds, &\quad (1.9)
 \end{aligned}$$

where $U_t = |u_t|^{p/2}$,

$$\Phi_t = (B+1) \int_0^t \lambda_s ds, \quad \phi_t = \int_0^t (B\lambda_s + \mu_s) ds, \quad \lambda_t = 1 + \bar{\beta}_t^2 + \bar{b}_t^2 + \bar{\zeta}_t^2 + \bar{\nu}_t^2,$$

and the constants N, B depend only on p, r, δ, d, R_0 .

We prove this theorem in Section 3.

Example 1.8 (Example 1.3.1 [10]). Condition (1.7) requires, in particular, \hat{b} to be sufficiently small. It turns out that this smallness assumption is essential. For instance, $u_t(x) = \exp(-|x|^2/(4t))$ satisfies

$$\frac{\partial}{\partial t} u_t(x) = \Delta u_t(x) - \frac{d}{|x|^2} x^i D_i u_t, \quad (1.10)$$

where $|b_t(x)| = d/|x|$ is admissible with any $r < d$. In this situation (1.8) fails. However, if we take εb in place of b our equation will satisfy (1.7) and the estimate becomes available if ε is sufficiently small.

Similarly, one cannot allow $\hat{\beta}$ and $\hat{\zeta}$ to be large. Indeed, equation (1.10) can be rewritten as

$$\frac{\partial}{\partial t} u_t(x) = \Delta u_t(x) - D_i \left(\frac{d}{|x|^2} x^i u_t \right) + \frac{d(d-2)}{|x|^2} u_t.$$

Note that $|b| = \varepsilon/|x|$ is not allowed in [10] and [11], so it seems that Theorem 1.7 is new even if there are no stochastic terms.

Our next result is about estimating Du_t . For that we need to impose a stronger assumptions than Assumption 1.4 and 1.6. By $Df(x)$ we mean the collection of the first-order Sobolev derivatives of (vector- or tensor-valued) $f(x)$ and by $|Df(x)|$ we mean any fixed norm in the space where $Df(x)$ takes its values. Similar meaning will be given later to $D^2 f$ and $|D^2 f|$.

Assumption 1.9. We have $\beta = f \equiv 0$ and

$$E \int_0^\tau \int_{\mathbb{R}^d} (|f_t|^p + |g_t|^p + |Df_t|^p + |Dg_t|^p) dx ds < \infty.$$

Introduce $\mathbf{c}_t = |Dc_t|^{1/2}$.

Assumption 1.10. The functions $Da, D\sigma, b, \zeta, \mathbf{c}, \nu, D\nu$ are admissible.

Theorem 1.11. *Under the above assumptions suppose that a solution u_t of (1.1)-(1.2) exists and*

$$E \int_0^\tau \|u_t\|_{W_p^2}^p dt < \infty. \quad (1.11)$$

Then (a.s.) $Du_{t \wedge \tau}$ is a continuous L_p -valued function and there exists a constant $N_0 = N_0(p, \delta, d, r)$ such that, if

$$N_0(\widehat{D}a + \widehat{D}\sigma + \widehat{b} + \widehat{c} + \widehat{v} + \widehat{D}\nu) \leq 1, \quad (1.12)$$

then

$$G + H \leq NK, \quad (1.13)$$

where

$$\begin{aligned} G &= E \sup_{t \leq \tau} e^{-\Psi_t} \int_{\mathbb{R}^d} |Du_t|^p dx, \\ H &= E \int_0^\tau e^{-\Psi_s} \int_{\mathbb{R}^d} (|Du_t|^{p-2} |D^2u_t|^2 + \Lambda_t |Du_t|^p) dx ds, \\ K &= E \int_{\mathbb{R}^d} (|u_0|^p + |Du_0|^p) dx \\ &\quad + E \int_0^\tau e^{-\Psi_s} \int_{\mathbb{R}^d} (|f_t|^p + |g_t|^p + |Df_t|^p + |Dg_t|^p) dx ds, \\ \Psi_t &= C \int_0^t \Lambda_s ds, \quad \Lambda_t = 1 + \overline{Da}_t^2 + \overline{D\sigma}_t^2 + \overline{b}_t^2 + \overline{c}_t^2 + \overline{v}_t^2 + \overline{D\nu}_t^2 \end{aligned}$$

and the constants N, C depend only on p, r, δ, d, R_0 .

We prove this theorem in Section 4.

Remark 1.12. Somewhat similar result is presented in Theorem 2.1 of [3]. There are, however, several distinctions. Let alone systems, in [3] the equations can degenerate and because of that the assumptions on the smoothness of a and σ are stronger. Also the coefficients of lower order terms like b, c are assumed to be bounded. We heavily use the nondegeneracy of our equation.

For nondegenerate equations [6] provides a similar result with a weaker assumption on a but with bounded lower order coefficients and with an assumption on σ which, basically, requires its continuity in x . Under our assumptions σ can have rather wild discontinuities. Incidentally, one of the aims of this article is to develop some tools on the way of proving that under our assumptions on σ and b the stochastic equation $dx_t = \sigma_t^k(x_t) dw_t^k + b_t(x_t) dt$ with deterministic σ and b and any nonrandom initial condition has a strong solution provided that the matrix $(\sigma_t^{ik} \sigma_t^{jk})$ is uniformly nondegenerate. The way we mean is similar to the one in [8].

2. AUXILIARY RESULTS

For $r \in (1, d]$ and two real-valued functions b and ζ on \mathbb{R}^d set

$$\hat{b} := \|b\|_{E_{r,1}}, \quad \hat{\zeta} := \|\zeta\|_{E_{r,1}},$$

where, for any function f ,

$$\|f\|_{E_{r,1}} := \sup_{\substack{\rho > 0 \\ B \in \mathbb{B}_\rho}} \rho \|f\|_{L_r(B)}, \quad \|f\|_{L_r(B)}^r = \int_B |f|^r dx.$$

For functions u, v on \mathbb{R}^d we write $u \prec v$ if the integral of u over \mathbb{R}^d is less than or equal to that of v . If the integrals are equal, we write $u \sim v$. Also set $p' = p/(p-1)$.

Lemma 2.1. *Let $u, v \in C_0^\infty$, then*

$$1 < p < r \implies |bu|^p \prec N\hat{b}^p |Du|^p, \quad (2.1)$$

$$p, p' \in (1, r) \implies \int_{\mathbb{R}^d} |v\zeta^2 u| dx \leq N\hat{\zeta}^2 \|Dv\|_{L_{p'}} \|Du\|_{L_p}, \quad (2.2)$$

$$1 < p' < r \implies \int_{\mathbb{R}^d} |vbDu| dx \leq N\hat{b} \|Dv\|_{L_{p'}} \|Du\|_{L_p}, \quad (2.3)$$

where the constants N depend only on d, r, p .

Proof. Estimate (2.1) is proved in [1]. To prove (2.2), it suffices to observe that

$$\int_{\mathbb{R}^d} |v\zeta^2 u| dx \leq \|\zeta v\|_{L_{p'}} \|\zeta u\|_{L_p},$$

and then use (2.1). Similarly (2.3) is proved on the basis of

$$\int_{\mathbb{R}^d} |vbDu| dx \leq \|bv\|_{L_{p'}} \|Du\|_{L_p}.$$

The lemma is proved.

Remark 2.2. Estimate (2.3) shows, in particular, that $g := b^i D_i u$, with $\widehat{b} < \infty$, can be represented as $D_i f^i + f$, where the L_p -norms of f^i, f are controlled by that of Du . Indeed, for $\Lambda = (1 - \Delta)^{1/2}$ the estimate implies that $h := \Lambda^{-1}g \in L_p$ and $g = \Lambda h = (1 - \Delta)\Lambda^{-1}h = D_i f^i + f$, where $f = \Lambda^{-1}h$, $f^i = -D_i \Lambda^{-1}h$.

Lemma 2.3. *If f is admissible, then for any t and $u \in C_0^\infty$ we have*

$$|f_t|^2 |u|^2 \prec N(d, r) \hat{f}^2 |Du|^2 + (N(d, r) R_0^{-2} \hat{f}^2 + 2\bar{f}_t^2) |u|^2. \quad (2.4)$$

Proof. We have $|f_t|^2 \leq 2|f_t^M|^2 + 2|f_t^B|^2$ and this shows how $2|\bar{f}_t^B|^2|u|^2$ appeared in (2.4). By a version of the Chiarenza–Frasca theorem in [1] (see, for instance, Lemma 3.5 in [8])

$$|f_t^M|^2|u_t|^2 \prec N\hat{f}^2|Du_t|^2 + NR_0^{-2}\hat{f}^2|u_t|^2.$$

This proves the lemma.

Corollary 2.4.

$$\begin{aligned} |f_t||Du_t||u_t| &\leq \varepsilon|Du_t|^2 + \varepsilon^{-1}|f_t|^2|u_t|^2 \\ &\prec (\varepsilon + N\varepsilon^{-1}\hat{f}^2)|Du_t|^2 + N\varepsilon^{-1}(\bar{f}_t^2 + R_0^{-2}\hat{f}^2)|u_t|^2. \end{aligned}$$

3. PROOF OF THEOREM 1.7

In light of Lemma 2.1 and Remark 2.2, Theorem 2.1 of [7] is applicable, which in particular implies the L_p -continuity of $u_{t \wedge \tau}$. By this theorem also, after introducing $\theta_t = \text{sign } u_t$ we obtain for $t \leq \tau$ that

$$\begin{aligned} (1/p) \int_{\mathbb{R}^d} |u_t|^p dx &= (1/p) \int_{\mathbb{R}^d} |u_0|^p dx + \int_0^t \int_{\mathbb{R}^d} J_s ds + m_t \\ &+ \int_0^t \int_{\mathbb{R}^d} \theta_s |u_s|^{p-1} (L_s u_s + D_i f_s^i + f_s) dx ds, \end{aligned}$$

where

$$\begin{aligned} J_t &:= ((p-1)/2)|u_s|^{p-2} \sum_k (M_s^k u_t + g_s^k)^2 dx \\ &\leq ((p-1)/2)|u_t|^{p-2} \sigma_t^{ik} D_i u_t \sigma_t^{jk} D_j u_t + (\delta/(2p))|DU_t|^2 \\ &\quad + N|u_t|^p |\nu_t|^2 + N|u_t|^{p-2} |g_t|^2, \end{aligned} \tag{3.1}$$

$$m_t := p \int_0^t \int_{\mathbb{R}^d} \theta_s |u_s|^{p-1} (M_s^k u_s + g_s^k) dx dw_s^k.$$

Here and below constants N denote constants depending only on d, r, p, δ .

From the start we may assume that $\hat{b}, \hat{\beta}, \hat{\zeta}, \hat{\nu} \leq 1$ (in order to be able to drop terms like \hat{f}^2 in the second term on the right in (2.4)). Note that

$$\begin{aligned} I_1 &:= \theta_t |u_t|^{p-1} D_i (a_t^{ij} D_j u_t) + ((p-1)/2) |u_t|^{p-2} \sigma_t^{ik} D_i u_t \sigma_t^{jk} D_j u_t \\ &\quad + (\delta/(2p)) |DU_t|^2 \sim -\frac{2(p-1)}{p^2} (2a - \alpha)_t^{ij} D_i U_t D_j U_t \\ &\quad + (\delta/(2p)) |DU_t|^2 \leq -\hat{\delta} |DU_t|^2, \end{aligned} \tag{3.2}$$

where $\hat{\delta} = \delta/(2p)$.

Next, by Corollary 2.4

$$I_2 := \theta_t |u_t|^{p-1} D_i (\beta_t^i u_t) + \theta_t |u_t|^{p-1} b_t^i D_i u_t$$

$$\begin{aligned}
 & \sim -(p-1)\theta_t|u_t|^{p-1}\beta_t^i D_i u_t + \theta_t|u_t|^{p-1}b_t^i D_i u_t \\
 & \leq (N U_t(|\beta_t| + |b_t|))|DU_t| \\
 & \leq (\hat{\delta}/4 + N_1(\hat{\beta}^2 + \hat{b}^2))|DU_t|^2 + N(1 + \bar{\beta}_t^2 + \bar{b}_t^2)|U_t|^2.
 \end{aligned}$$

We subject $\hat{\beta}$ and \hat{b} to

$$N_1(\hat{\beta}^2 + \hat{b}^2) \leq \hat{\delta}/4 \quad (3.3)$$

and then get that

$$I_2 \leq (\hat{\delta}/2)|DU_t|^2 + N(1 + \bar{\beta}_t^2 + \bar{b}_t^2)|u_t|^p. \quad (3.4)$$

By Lemma 2.3

$$\begin{aligned}
 I_3 & := \theta_t|u_t|^{p-1}c_t u_t + N|u_t|^p|\nu_t|^2 \leq \zeta_t^2 U_t^2 + N|\nu_t|^2 U_t^2 \\
 & \prec N_2(\hat{\zeta}^2 + \hat{\nu}^2)|DU_t|^2 + N(1 + \bar{\zeta}_t^2 + \bar{\nu}_t^2)|U_t|^2.
 \end{aligned}$$

We subject $\hat{\zeta}$ and $\bar{\nu}$ to

$$N_2(\hat{\zeta}^2 + \hat{\nu}^2) \leq \hat{\delta}/4 \quad (3.5)$$

and then get that

$$I_3 \leq (\hat{\delta}/4)|DU_t|^2 + N(1 + \bar{\zeta}_t^2 + \bar{\nu}_t^2)|u_t|^p. \quad (3.6)$$

Next,

$$\begin{aligned}
 I_4 & := |u_t|^{p-1}\theta_t D_i f_t^i \sim N f_t^i |u_t|^{p/2-1} D_i U_t \leq (\hat{\delta}/16)|DU_t|^2 + N|f_t|^2 |u_t|^{p-2} \\
 & \leq (\hat{\delta}/8)|DU_t|^2 + N|U_t|^2 + N|f_t|^p.
 \end{aligned} \quad (3.7)$$

In estimating the last remaining terms we use elementary inequalities

$$|u_t|^{p-1} f_t + N|u_t|^{p-2} |g_t|^2 \leq N|u_t|^p + N(|f_t|^p + |g_t|^p). \quad (3.8)$$

Upon combining this with (3.2), (3.4), (3.6), and (3.7), we get that, if condition (1.7) is satisfied for an appropriate N_0 , then

$$\begin{aligned}
 (1/p)d \int_{\mathbb{R}^d} |u_t|^p dx & \leq \int_{\mathbb{R}^d} \left(-(\hat{\delta}/8)|DU_t|^2 + N(|f_t|^p + |f_t|^p + |g_t|^p) \right) dx dt \\
 & \quad + (B/p) \int_{\mathbb{R}^d} \lambda_t |u_t|^p dx dt + dm_t.
 \end{aligned} \quad (3.9)$$

It follows that for $t \leq \tau$

$$\begin{aligned}
 & e^{-\phi_t} \int_{\mathbb{R}^d} |u_t|^p dx + (\hat{\delta}/8) \int_0^t e^{-\phi_s} \int_{\mathbb{R}^d} |DU_s|^2 dx ds \\
 & \leq \int_{\mathbb{R}^d} |u_0|^p dx + N \int_0^t e^{-\phi_s} \int_{\mathbb{R}^d} (|f_s|^p + |f_s|^p + |g_s|^p - \mu_t |u_t|^p) dx ds \\
 & \quad + \int_0^t e^{-\phi_s} dm_s.
 \end{aligned}$$

This yields (1.8) because, as we know from [7], m_t is a martingale. To prove (1.9) we follow E. Pardou. Estimate (3.9) and the Davis's inequality imply that

$$\begin{aligned} & E \sup_{t \leq \tau} e^{-\Phi_t} \int_{\mathbb{R}^d} |u_t|^p dx \leq E \int_{\mathbb{R}^d} |u_0|^p dx \\ & + NE \int_0^\tau e^{-\Phi_s} \int_{\mathbb{R}^d} (|f_s|^p + |g_s|^p) dx ds \\ & + NE \left(\int_0^\tau e^{-2\Phi_s} \sum_k \left(\int_{\mathbb{R}^d} |u_s|^{p-1} |M_s^k u_s + g_s^k| dx \right)^2 ds \right)^{1/2}. \end{aligned} \quad (3.10)$$

Here $|u_s|^{p-1} = |u_s|^{p/2} |u_s|^{p/2-1}$ and the last term is dominated by

$$\begin{aligned} I & := NE \left(\int_0^\tau e^{-2\Phi_s} \|u_s\|_{L^p}^p \int_{\mathbb{R}^d} J_s dx ds \right)^{1/2} \\ & \leq NE \left(\sup_{s \leq \tau} e^{-\Phi_s} \|u_s\|_{L^p}^p \right)^{1/2} \left(\int_0^\tau e^{-\Phi_s} \int_{\mathbb{R}^d} J_s dx ds \right)^{1/2} \\ & \leq N \left(E \sup_{s \leq \tau} e^{-\Phi_s} \|u_s\|_{L^p}^p \right)^{1/2} \left(E \int_0^\tau e^{-\Phi_s} \int_{\mathbb{R}^d} J_s dx ds \right)^{1/2} \end{aligned}$$

It is seen from (3.1) and the estimates of I_3 and (3.8) that

$$\begin{aligned} J_t & \leq N |DU_t|^2 + N |u_t|^p |\nu_t|^2 + N |u_t|^{p-2} |g_t|^2 \\ & \prec N |DU_t|^2 + N(1 + \bar{\nu}_t^2) |u_t|^p + N |g_t|^p. \end{aligned}$$

The last expression is under control due to (1.8) with $\mu_t = \lambda_t \geq 1 + \bar{\nu}_t^2$. Hence coming back to (3.10) we immediately obtain (1.9). This proves the theorem.

Remark 3.1. It follows from the proof that in the definition of ϕ_t one can take any constant $\geq B$ in place of B .

4. PROOF OF THEOREM 1.11

We prove the theorem after some discussion.

Lemma 4.1. *For any $n \geq 1$, $p_i > 0$, $i = 1, \dots, n$, and $\kappa \geq d + 1 + p_1 + \dots + p_n$ there exists a constant $N = N(d, \kappa, n, p_i)$ such that for any $A_i \in \mathbb{R}^d$, $i = 1, \dots, n$, we have*

$$|A_1|^{p_1} \cdot \dots \cdot |A_n|^{p_n} \leq N \int_{\mathbb{R}^d} h(\eta) |(\eta, A_1)|^{p_1} \cdot \dots \cdot |(\eta, A_n)|^{p_n} d\eta, \quad (4.1)$$

where $h(\eta) = (1 + |\eta|^\kappa)^{-1}$ and (η, A_i) is the scalar product of η and A_i .

Proof. By dividing both parts by $|A_1|^{p_1}$ we see that we may assume that $|A_1| = 1$. After that we may also assume that $|A_i| = 1$ for all i . Then the result easily follows by contradiction. The lemma is proved.

By applying this lemma to $A_1 = Du$ and $A_2 = Dv_{x^i}$ and then summing up with respect to i we get the following, where

$$u_{(\eta)} := \eta^i D_i u.$$

Corollary 4.2. *For any smooth functions u, v on \mathbb{R}^d , on \mathbb{R}^d we have for appropriate p, q, κ*

$$\begin{aligned} |Du|^p &\leq N \int_{\mathbb{R}^d} h(\eta) |u_{(\eta)}|^p d\eta, \\ |Du|^p |D^2 v|^q &\leq N \int_{\mathbb{R}^d} h(\eta) |u_{(\eta)}|^p |Dv_{(\eta)}|^q d\eta. \end{aligned}$$

Proof of Theorem 1.11. Fix $\kappa \geq d+1+p$ and for functions $u(x, \eta)$ and $v(x, \eta)$ on \mathbb{R}^{2d} let us write $u \prec_{\kappa} v$ if

$$\int_{\mathbb{R}^{2d}} h(\eta) u(x, \eta) dx d\eta \leq \int_{\mathbb{R}^{2d}} h(\eta) v(x, \eta) dx d\eta.$$

We also write $u \sim_{\kappa} v$ if the above integrals coincide. In these terms Corollary 4.2 implies that

$$|\eta|^p |Du|^p \prec_{\kappa} N |u_{(\eta)}|^p, \quad |\eta|^p |Du|^{p-2} |D^2 v|^2 \prec_{\kappa} N |u_{(\eta)}|^{p-2} |Dv_{(\eta)}|^2. \quad (4.2)$$

Introduce

$$\begin{aligned} v_t &= v_t(x, \eta) = \eta^i D_i u_t(x), \quad \theta_t = \text{sign } v_t, \quad V_t = |v_t|^{p/2}, \\ W_t^2 &:= |\eta|^p |Du_t|^{p-2} |D^2 u_t|^2, \quad \gamma_t^j = D_i a_t^{ij} + b_t^i. \end{aligned}$$

Observe that thanks to (4.2)

$$|\eta|^p |Du|^p \prec_{\kappa} N |v_t|^p, \quad W_t^2 \prec_{\kappa} N |DV_t|^2, \quad |DV_t|^2 \leq N W_t^2. \quad (4.3)$$

Owing to (1.11) we can substitute $\phi_{(\eta)}$ in place of ϕ into (1.4) which yields

$$\begin{aligned} dv_t &= (L_t v_t + a_{t(\eta)}^{ij} D_{ij} u_t + \gamma_{t(\eta)}^i D_i u_t + c_{t(\eta)} u_t + f_{t(\eta)}) dt \\ &\quad + (M_t^k v_t + \sigma_{t(\eta)}^{ik} D_i u_t + \nu_{t(\eta)}^k u_t + g_{t(\eta)}^k) dw_t^k. \end{aligned}$$

Again in light of Lemma 2.1, Theorem 2.1 of [7] is applicable to v_t , which in particular implies the L_p -continuity of $v_{t \wedge \tau}$ and thus of $Du_{t \wedge \tau}$. By this theorem also we obtain for $t \leq \tau$ that

$$(1/p) \int_{\mathbb{R}^{2d}} h(\eta) |v_t|^p dx d\eta = (1/p) \int_{\mathbb{R}^{2d}} h(\eta) |v_0|^p dx d\eta$$

$$\begin{aligned}
& + \int_0^t \int_{\mathbb{R}^{2d}} h(\eta) \theta_s |v_s|^{p-1} (L_s v_s + a_{s(\eta)}^{ij} D_{ij} u_s) dx d\eta ds \\
& + \int_0^t \int_{\mathbb{R}^{2d}} h(\eta) \theta_s |v_s|^{p-1} (\gamma_{t(\eta)}^i D_i u_t + c_{s(\eta)} u_s + f_{s(\eta)}) dx d\eta ds \\
& + \int_0^t \int_{\mathbb{R}^{2d}} h(\eta) J_s dx d\eta ds + m_t,
\end{aligned}$$

where

$$\begin{aligned}
J_t & = ((p-1)/2) |v_t|^{p-2} \sum_k (M_t^k v_t + \sigma_{t(\eta)}^{ik} D_i u_t + \nu_{t(\eta)}^k u_t + g_{t(\eta)}^k)^2 \\
& \leq ((p-1)/2) |v_t|^{p-2} \sigma_t^{ik} D_i v_t \sigma_t^{jk} D_j v_t + (\delta/(2p)) |DV_t|^2 \\
& + N |\nu_t|^2 |v_t|^p + N |\eta|^2 |v_t|^{p-2} (|D\sigma_t|^2 |Du_t|^2 + |D\nu_t|^2 |u_t|^2 + |Dg_t|^2), \quad (4.4)
\end{aligned}$$

$$\begin{aligned}
m_t & = \int_0^t \int_{\mathbb{R}^{2d}} h(\eta) J_s^k dx d\eta dw_s^k, \\
J_t^k & = \theta_t |v_t|^{p-1} (M_t^k v_t + \sigma_{t(\eta)}^{ik} D_i u_t + \nu_{t(\eta)}^k u_t + g_{t(\eta)}^k).
\end{aligned}$$

Next, we may assume that $\widehat{D}a, \widehat{D}\sigma, \widehat{b}, \widehat{\zeta}, \widehat{\hat{c}}, \widehat{\hat{v}}, \widehat{D}\nu \leq 1$ and first deal with the terms containing a and σ but not their derivatives. Their sum is (see (4.4)) less than

$$\begin{aligned}
I_1 & := \theta_t |v_t|^{p-1} D_i (a_t^{ij} D_j v_t) + ((p-1)/2) |v_t|^{p-2} \sum_k (\sigma^{ik} D_i v_t)^2 \\
& + (\delta/(2p)) |DV_t|^2 \sim_\kappa -((p-1)/p^2) (4a_t^{ij} - 2\alpha_t^{ij}) D_i V_t D_j V_t + (\delta/(2p)) |DV_t|^2 \\
& \leq -(\delta/(2p)) |DV_t|^2 \prec_\kappa -\hat{\delta} (|DV_t|^2 + W_t^2), \quad (4.5)
\end{aligned}$$

where $\hat{\delta} = \hat{\delta}(p, d, \delta) > 0$ and in the last inequality we used (4.3).

By Lemma 2.3

$$\begin{aligned}
I_2 & := \theta_t v_t^{p-1} a_{t(\eta)}^{ij} D_{ij} u_t \leq (V_t |Da_t|) |\eta|^{p/2} |Du_t|^{p/2-1} |D^2 u_t| \\
& \leq (\hat{\delta}/2) W_t^2 + N V_t^2 |Da_t|^2 \prec_\kappa (\hat{\delta}/2) W_t^2 + N_1 \widehat{D}a^2 |DV_t|^2 + N(1 + \overline{D}a_t^2) V_t^2.
\end{aligned}$$

We subject $\widehat{D}a$ to

$$N_1 \widehat{D}a^2 \leq \hat{\delta}/2 \quad (4.6)$$

and get that

$$I_2 \prec_\kappa (\hat{\delta}/2) (|DV_t|^2 + W_t^2) + N(1 + \overline{D}a_t^2) V_t^2. \quad (4.7)$$

Next, come the terms containing the products of γ, b and their derivatives:

$$I_3 := \theta_t |v_t|^{p-1} b_t^i D_i v_t + \theta_t |v_t|^{p-1} \gamma_{t(\eta)}^i D_i u_t \sim_\kappa (2/p) (V_t b_t^i) D_i V_t$$

$$-\eta^j \gamma_t^i ((p-1)|v_t|^{p-2} D_i v_t D_i u_t + |v_t|^{p-1} D_{ij} u_t).$$

Here by Corollary 2.4

$$(2/p)(V_t \widehat{b}_t^i) D_i V_t \prec_\kappa (\hat{\delta}/8) |DV_t|^2 + N_5 \widehat{b}^2 |DV_t|^2 + N \overline{b}_t^2 V_t^2.$$

By Corollary 4.2 the remaining term in I_3 is dominated by

$$\begin{aligned} & N |\gamma_t| |\eta|^p |Du|^{p-1} |D^2 u| \\ & \prec_\kappa N (|Da_t| + |b_t|) V_t (|\eta|^{p/2} |Du_t|^{p/2-1} |D^2 u_t|) \\ & \leq N (|Da_t|^2 + |b_t|^2) V_t^2 + (\hat{\delta}/4) W_t^2 \\ & \prec_\kappa N_2 (\widehat{Da}^2 + \widehat{b}^2) |DV_t|^2 + (\hat{\delta}/4) W_t^2 + N(1 + \overline{Da}_t^2 + \overline{b}_t^2) V_t^2. \end{aligned}$$

We subject \widehat{Da} and \widehat{b} to

$$N_2 (\widehat{Da}^2 + \widehat{b}^2) \leq \hat{\delta}/8, \quad (4.8)$$

and get that

$$I_3 \prec_\kappa (\hat{\delta}/4) (|DV_t|^2 + W_t^2) + N(1 + \overline{Da}_t^2 + \overline{b}_t^2) V_t^2. \quad (4.9)$$

The term with $|Du_t|^2$ is

$$\begin{aligned} I_4 & := N |\eta|^2 |v_t|^{p-2} |D\sigma_t|^2 |Du_t|^2 \leq N |\eta|^p |D\sigma_t|^2 (|Du_t|^{p/2})^2 \\ & \prec_\kappa N_3 \widehat{D\sigma}^2 W_t^2 + N(1 + \overline{D\sigma}_t^2) V_t^2. \end{aligned}$$

We subject $\widehat{D\sigma}$ to

$$N_3 \widehat{D\sigma}^2 \leq \hat{\delta}/8, \quad (4.10)$$

and get that

$$I_4 \prec_\kappa (\hat{\delta}/8) (|DV_t|^2 + W_t^2) + N(1 + \overline{D\sigma}_t^2) V_t^2. \quad (4.11)$$

Now come the terms which do not contain Du_t . Their sum is

$$I_5 := I_6 + I_7 + I_8,$$

where

$$\begin{aligned} I_6 & := \theta_t |v_t|^{p-1} (c_t v_t + c_{t(\eta)} u_t), \quad I_7 := N |\nu_t|^2 |v_t|^p + N |\eta|^2 |v_t|^{p-2} |D\nu_t|^2 |u_t|^2, \\ I_8 & := \theta_t |v_t|^{p-1} f_{t(\eta)} + N |\eta|^2 |v_t|^{p-2} |Dg_t|^2. \end{aligned}$$

We have

$$\begin{aligned} I_6 & \leq \mathfrak{c}_t^2 V_t^2 + \mathfrak{c}_t^2 (V_t^2 + |\eta|^p |u_t|^p) \prec_\kappa N_4 (\hat{\mathfrak{c}}^2 + \mathfrak{c}^2) |DV_t|^2 \\ & + N \hat{\mathfrak{c}}^2 |\eta|^p |Du_t|^2 + N((1 + \overline{\mathfrak{c}}_t^2) |\eta|^p |u_t|^p + (1 + \overline{\mathfrak{c}}_t^2 + \overline{\mathfrak{c}}_t^2) V_t^2) \end{aligned}$$

We subject $\hat{\mathfrak{c}}^2$ and \mathfrak{c}^2 to

$$N_4 (\hat{\mathfrak{c}}^2 + \mathfrak{c}^2) \leq \hat{\delta}/16 \quad (4.12)$$

and get that

$$I_6 \prec_\kappa (\hat{\delta}/16) (|DV_t|^2 + W_t^2)$$

$$+N(1 + \bar{\zeta}_t^2 + \bar{\mathfrak{c}}_t^2)V_t^2 + N|\eta|^p|DU_t|^2 + N(1 + \bar{\mathfrak{c}}_t^2)|\eta|^pU_t^2. \quad (4.13)$$

Similarly we deal with I_7 and see that, if for some well defined N_5

$$N_5(\hat{\nu}^2 + \widehat{D\nu}^2) \leq \hat{\delta}/32, \quad (4.14)$$

then

$$\begin{aligned} I_7 &\prec_{\kappa} (\hat{\delta}/32)(|DV_t|^2 + W_t^2) \\ &+ N(1 + \bar{\nu}_t^2 + \overline{D\nu}_t^2)V_t^2 + N|\eta|^p|DU_t|^2 + N(1 + \overline{D\nu}_t^2)|\eta|^pU_t^2. \end{aligned} \quad (4.15)$$

In what concerns I_8 , obviously,

$$I_8 \leq V_t^2 + N|\eta|^p(|Df_t|^p + |Dg_t|^p).$$

Thus, given that (4.6), (4.8), (4.10), (4.12), and (4.14) hold, we have

$$\begin{aligned} d|v_t|^p &\prec_{\kappa} -\check{\delta}(|DV_t|^2 + W_t^2) dt + (C-1)\Lambda_t|v_t|^p dt \\ &+ N|\eta|^p(|Df_t|^p + |Dg_t|^p + |DU_t|^2 + (1 + \bar{\mathfrak{c}}_t^2 + \overline{D\nu}_t^2)U_t^2) dt + J_t^k dw_t^k, \end{aligned} \quad (4.16)$$

where $\check{\delta} = \hat{\delta}/(32p)$.

It follows that

$$\begin{aligned} d(e^{-\Psi_t}|v_t|^p) &\prec_{\kappa} -\check{\delta}e^{-\Psi_t}(|DV_t|^2 + W_t^2) dt - \Lambda_t e^{-\Psi_t}|v_t|^p dt \\ &+ Ne^{-\Psi_t}|\eta|^p(|Df_t|^p + |Dg_t|^p + |DU_t|^2 + (1 + \bar{\mathfrak{c}}_t^2 + \overline{D\nu}_t^2)U_t^2) dt + e^{-\Psi_t}J_t^k dw_t^k. \end{aligned} \quad (4.17)$$

We convert this into the integral form with integrals with respect to x, η , and t , then take expectations of the resulting inequality. We also use Corollary 4.2. Then we get

$$\begin{aligned} H &= E \int_0^{\tau} e^{-\Psi_t} \int_{\mathbb{R}^d} (|Du_t|^{p-2}|D^2u_t|^2 + \Lambda_t|Du_t|^p) dx dt \leq NE|Du_0|^p \\ &+ NE \int_0^{\tau} e^{-\Psi_t} \int_{\mathbb{R}^d} (|Df_t|^p + |Dg_t|^p + |DU_t|^2 + (1 + \bar{\mathfrak{c}}_t^2 + \overline{D\nu}_t^2)U_t^2) dt. \end{aligned} \quad (4.18)$$

We may assume that $C\Lambda_t \geq B\lambda_t + 1 + \bar{\mathfrak{c}}_t^2 + \overline{D\nu}_t^2$ and then, owing to (1.8), the term in (4.18) containing $|DU_t|^2 + (1 + \bar{\mathfrak{c}}_t^2 + \overline{D\nu}_t^2)U_t^2$ is estimated by

$$NE|u_0|^p + NE \int_0^{\tau} e^{-\Psi_t} \int_{\mathbb{R}^d} (|f_t|^p + |g_t|^p) dx dt \leq NK,$$

so that $H \leq NK$.

Next, we come back to (4.17) and, after converting it, take sup with respect to $t \leq \tau$ and take expectations in the resulting inequality. We also use Corollary 4.2 and the result of the above treatment of (4.18). Then we obtain

$$G \leq NK + I, \quad (4.19)$$

where

$$I = E \sup_{t \leq \tau} \left| \int_0^t e^{-\Psi_s} \int_{\mathbb{R}^{2d}} h(\eta) J_s^k dx d\eta dw_s^k \right|.$$

By Davis's inequality

$$\begin{aligned} I &\leq NE \left(\int_0^\tau e^{-2\Psi_t} \sum_k \left(\int_{\mathbb{R}^{2d}} h(\eta) J_t^k dx d\eta \right)^2 dt \right)^{1/2} \\ &= NE \left(\int_0^\tau \sum_k e^{-2\Psi_t} \left(\int_{\mathbb{R}^{2d}} h(\eta) \theta_t |v_t|^{p-1} (M_t^k v_t \right. \right. \\ &\quad \left. \left. + \sigma_{t(\eta)}^{ik} D_i u_t + \nu_{t(\eta)}^k u_t + g_{t(\eta)}^k) dx d\eta \right)^2 dt \right)^{1/2} \\ &\leq NE \left(\sup_{t \leq \tau} e^{-\Psi_t} \int_{\mathbb{R}^d} |v_t|^p dx \right)^{1/2} \left(\int_0^\tau e^{-\Psi_t} \int_{\mathbb{R}^{2d}} h(\eta) J_t dx d\eta dt \right)^{1/2} \\ &\leq NG^{1/2} \left(E \int_0^\tau e^{-\Psi_t} \int_{\mathbb{R}^{2d}} h(\eta) J_t dx d\eta dt \right)^{1/2}. \end{aligned}$$

Above we actually estimated J_t splitting it in parts. It is seen from these estimates that

$$\begin{aligned} J_t &\prec_\kappa N |DV_t|^2 + I_4 + I_7 + N |\eta|^2 |v_t|^{p-2} |Dg_t|^2 \\ &\prec_\kappa N |\eta|^p (|Du_t|^{p-2} |D^2 u_t|^2 + \Lambda_t |Du_t|^p) + N |\eta|^p |DU_t|^2 \\ &\quad + N (1 + \overline{D\nu_t^2}) |\eta|^p U_t^2 + N |\eta|^p |Dg_t|^p. \end{aligned}$$

Recalling our argument about (4.18) we see that $I \leq NG^{1/2} K^{1/2}$, which after being substituted into (4.19), yields that $G \leq NK$ as well. This proves (1.13) and the theorem.

The manuscript contains no data and there is no conflict of interest.

REFERENCES

- [1] F. Chiarenza and M. Frasca, *A remark on a paper by C. Fefferman*, Proc. Amer. Math. Soc., Vol. 108 (1990), No. 2, 407–409.
- [2] K. Dareiotis and M. Gerencsér, *On the boundedness of solutions of SPDEs*, Stoch. Partial Differ. Equ. Anal. Comput., Vol. 3 (2015), No. 1, 84–102.
- [3] M. Gerencsér, I. Gyöngy, and N.V. Krylov, *On the solvability of degenerate stochastic partial differential equations in Sobolev spaces*, Stochastic Partial Differential Equations: Analysis and Computations, Vol. 3 (2015), No. 1, 52–83.
- [4] N. Ikeda and S. Watanabe, *Stochastic differential equations and diffusion processes*, North-Holland Publishing Company, Amsterdam-Oxford-New York, 1981.
- [5] D. Kinzebulatov, K.R.Madou, *Stochastic equations with time-dependent singular drift*, <https://arxiv.org/abs/2105.07312>
- [6] N.V. Krylov, *On divergence form SPDEs with VMO coefficients*, SIAM J. Math. Anal. Vol. 40 (2009), No. 6, 2262–2285

- [7] N.V. Krylov, *Itô's formula for the L_p -norm of stochastic W_p^1 -valued processes*, Probab. Theory and Related fields, Vol. 147 (2010), No. 3-4, 583-605.
- [8] N.V. Krylov, *On strong solutions of Itô's equations with $D\sigma$ and b in Morrey classes containing L_d* , <http://arxiv.org/abs/2111.13795>
- [9] N.V. Krylov and B.L. Rozovskii, *On the Cauchy problem for stochastic linear partial differential equations*, Izvestiya Akademii Nauk SSSR, seriya matematicheskaya, Vol. 41, No. 6 (1977), 1329–1347 in Russian, English translation in Math. USSR Izvestija, Vol. 11 (1977), No. 6, 1267–1284.
- [10] O. A. Ladyzhenskaya, V. A. Solonnikov, and N. N. Ural'tseva, “Lineinye i Kvazilineinye Uravneniya Parabolicheskogo Tipa,” (Russian) [Linear and Quasi-Linear Equations of Parabolic Type], “Nauka,” Moscow, 1967; English translation, Amer. Math. Soc., Providence, RI, 1968.
- [11] S. Monsurrò and M. Transirico, *On some L^p -estimates for solutions of elliptic equations in unbounded domains*, Math. Bohem., Vol. 140 (2015), No. 4, 507–515.
- [12] Jinniao Qiu, *L^2 -theory of linear degenerate SPDEs and L^p ($p > 0$) estimates for the uniform norm of weak solutions*, Stochastic Process. Appl., Vol. 130 (2020), No. 3, 1206–1225.
- [13] Rongrong Tian, Jinlong Wei, and Yanbin Tang, *The second-order parabolic PDEs with singular coefficients and applications*, Stoch. Anal. Appl., Vol. 38 (2020), No. 6, 1102–1121.

Email address: nkrylov@umn.edu

127 VINCENT HALL, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN, 55455