

Near horizon gravitational charges

Hai-Shan Liu and Pujian Mao

*Center for Joint Quantum Studies and Department of Physics,
School of Science, Tianjin University, 135 Yaguan Road, Tianjin 300350, China*

ABSTRACT. In this paper, we study the near horizon symmetry and gravitational charges in the Newman-Penrose formalism. In particular we investigate the effect from topological terms. We find that the Pontryagin term and Gauss-Bonnet term have significant influence on the near horizon charges and bring interesting novel features. We show that the gravitational charge derived from a general class of topological terms including the Pontryagin term and Gauss-Bonnet term can be obtained from the ambiguities of the symplectic potential.

1 Introduction

Motivation. In gravitational theory, one usually deals with surface charges. Evaluating surface charge requires that the spacetime must be equipped with a boundary, i.e., a special hypersurface where to compute the surface charge. Two natural choices for the boundary are usually applied which are the infinity and the horizon. At the spatial infinity, Arnowitt, Deser and Misner constructed, for the first time, appropriate surface integrals for gravitational energy-momentum [1]. Since causal line can not reach the spatial infinity, the ADM energy measures the total energy contained in the spacetime. Regarding to the system with gravitational waves that carry energy off, the Bondi energy was defined at null infinity [2]. The Bondi energy can never increase. It is conserved if and only if there is no gravitational wave.

Horizons are one of the most fascinating objects in spacetime with Lorentzian signature. They are responsible for many remarkable semiclassical properties, such as the Hawking radiation [3] and the Bekenstein-Hawking entropy [4,5]. Based on the covariant phase space method [6, 7], Wald showed that Black hole entropy is just a surface charge evaluated on the Killing horizon [8]. The first law of black hole mechanics is nothing but conservation of the surface charge between the horizon and infinity.¹

¹See also [9] for a generalization.

Once spacetime is equipped with a boundary, one can no longer consider boundary term in the action as something trivial to the theory. The most significant example is the Gibbons-Hawking-York term [10, 11] that added to Einstein-Hilbert action to enforce Dirichlet's boundary conditions. Meanwhile boundary terms have their own contributions to the surface charges [12–19]. However it is shown that the surface charge from the Pontryagin term and Gauss-Bonnet term at null infinity is absent at the leading and subleading order in the inverse of a large r expansion [18, 20]. This is somewhat unexpected from the fact that those two are higher derivative terms which should have a faster fall-off behavior at large distance (see also [21] for relevant evidence).

In this paper, we study the surface charge on the horizon with special emphasis on the effect from the Pontryagin term and Gauss-Bonnet term in four spacetime dimensions. We derive the surface charge in the Newman-Penrose (NP) formalism [22]. The reason to work in first order formalism is that it is in particular efficient for computing the near horizon charges. Because the charge expression for first order theory will not involve any derivative on the fields and variation of the fields [23–32]. Since any field at most starts at $O(1)$ in the near horizon expansion, it is enough to just consider one order of the field from the solution space for computing the charge. In the NP formalism, the most important variable is the spin connection. Each component has very definite geometric meaning. Given the fact that the charge does not include any derivative on the fields and the near horizon charges have clear thermodynamic interpretation [8, 33, 34], the relation between geometry and thermodynamics becomes transparent when computing the near horizon charges in the NP formalism.

Main result and plan of the paper. In the next Section, we derive the near horizon solution space for the NP equations with suitable gauge and boundary conditions. We compare our results with the ones in [35] and [36]. Section 3 is devoted to the near horizon symmetry and the transformation law of the near horizon fields. In the NP formalism, a gauge transformation is a combination of a diffeomorphism and a local Lorentz transformation. We show that with the required gauge and boundary conditions, the residual Lorentz transformation is completely fixed by the residual diffeomorphism. The near horizon symmetry consists of the near horizon supertranslations and superrotations. Our main results are presented in Section 4 where the surface charges are obtained. We first compute the surface charge from the Palatini action. Our result is consistent with [35–37]. Then we derive the surface charge from the Holst term. The Holst charge does not have a supertranslation part. Consequently, the dual mass [38] as the zero mode of the supertranslation charge is absent. Our next result is the charge from the Pontryagin term. We first show that the symplectic potential derived from a general class of terms that do not affect equations of motion including the Pontryagin term and Gauss-Bonnet term can be written as a combination of a Y term and a W term which are ambiguities of the sym-

plectic structure. We then compute the near horizon Pontryagin charge. The last case of our investigation is the Gauss-Bonnet charge. The near horizon charges from all four terms start at the leading order in the near horizon expansion. A surprising feature of the Gauss-Bonnet charge is that it includes field which is not horizon data. The charge from the Palatini, Holst, and Pontryagin action only involves the area of the horizon, the expansion of the horizon generator, and the rotation of reference frame on the horizon. While the Gauss-Bonnet charge involves the expansion of the other null direction, other than the horizon generator. In the last Section, we close with some final remarks. There are four appendices with some details of the computation in the main text.

2 Near horizon solution space

Newman and Penrose [22] established a special tetrad formalism with four null basis vectors $e_1 = l = e^2$, $e_2 = n = e^1$, $e_3 = m = -e^4$, $e_4 = \bar{m} = -e^3$. The basis vectors l and n are real while m and \bar{m} are complex conjugates of each other. The null basis vectors have the following orthogonality and normalization conditions

$$l \cdot m = l \cdot \bar{m} = n \cdot m = n \cdot \bar{m} = 0, \quad l \cdot n = 1, \quad m \cdot \bar{m} = -1. \quad (1)$$

The spacetime metric is obtained from the tetrad as

$$g_{\mu\nu} = n_\mu l_\nu + l_\mu n_\nu - m_\mu \bar{m}_\nu - \bar{m}_\mu m_\nu. \quad (2)$$

The components of spin connection are labeled by twelve Greek symbols. Ten independent components of the Weyl tensor are represented by five complex scalars. The Ricci tensor is defined in terms of four real scalars and three complex scalars. We will focus on vacuum solution in this work. So all the components of Ricci tensor are zero. For the notations, we would refer to [39].

In the NP formalism, by local Lorentz transformations, it is always possible to impose

$$\pi = \kappa = \epsilon = 0, \quad \rho = \bar{\rho}, \quad \tau = \bar{\alpha} + \beta. \quad (3)$$

According to the relations in Appendix A, such gauge choice means that l is tangent to a null geodesic with affine parameter. The other null basis vectors are parallel transported along l . Moreover, l is the gradient of a scalar field. It is of convenience to choose this scalar field as coordinate $u = x^1$ and take the affine parameter as coordinate $r = x^2$. For the rest two angular coordinates, we choose the stereographic coordinates $A = (z, \bar{z})$ which are related to the usual angular variables (θ, ϕ) by $z = \cot \frac{\theta}{2} e^{i\phi}$. The tetrad and the

co-tetrad satisfying conditions in (1) must have the forms

$$\begin{aligned} n &= \frac{\partial}{\partial u} + U \frac{\partial}{\partial r} + X^A \frac{\partial}{\partial x^A}, \\ l &= \frac{\partial}{\partial r}, \\ m &= \omega \frac{\partial}{\partial r} + L^A \frac{\partial}{\partial x^A}, \end{aligned} \quad (4)$$

and

$$\begin{aligned} n &= [-U - X^A(\bar{\omega}L_A + \omega\bar{L}_A)] du + dr + (\omega\bar{L}_A + \bar{\omega}L_A)dx^A, \\ l &= du, \\ m &= -X^A L_A du + L_A dx^A, \end{aligned} \quad (5)$$

where $L_A L^A = 0$, $L_A \bar{L}^A = -1$. The line element is then

$$\begin{aligned} ds^2 &= -[2U + g_{AB}(L^A \bar{\omega} + \bar{L}^A \omega)(L^B \bar{\omega} + \bar{L}^B \omega)] du^2 + 2dudr \\ &\quad + g_{AB} [dx^A + (L^A \bar{\omega} + \bar{L}^A \omega - X^A)du] [dx^B + (L^B \bar{\omega} + \bar{L}^B \omega - X^B)du], \end{aligned} \quad (6)$$

where

$$g_{AB} = -L_A \bar{L}_B - \bar{L}_A L_B. \quad (7)$$

The tetrad one form and the connection one form are defined as

$$e^a = e^a_\mu dx^\mu, \quad \Gamma_{ab} = \Gamma_{abc} e^c_\mu dx^\mu. \quad (8)$$

The exact formulas of the form e^a are given in (5) and the connection one form Γ_{ab} are given as

$$\Gamma_{12} = -(\gamma + \bar{\gamma})l + \bar{\tau}m + \tau\bar{m}, \quad (9)$$

$$\Gamma_{13} = -\tau l + \rho m + \sigma \bar{m}, \quad (10)$$

$$\Gamma_{23} = \bar{\nu} l - \bar{\mu} m - \bar{\lambda} \bar{m}, \quad (11)$$

$$\Gamma_{34} = (\gamma - \bar{\gamma})l - (\alpha - \bar{\beta})m + (\bar{\alpha} - \beta)\bar{m}. \quad (12)$$

Boundary conditions are somehow tricky in the near horizon case. In principle, all fields can start at $O(1)$. Different choices will lead to the horizon with different property [40], such as Killing horizon, isolated horizon [41,42], non-expansion horizon, dynamical horizon, and so on so forth. The main boundary conditions that We will choose are

$$U = O(r), \quad \omega = O(r), \quad \nu = O(r), \quad L^z = O(r), \quad X^A = O(r). \quad (13)$$

The first two conditions guarantee that the $r = 0$ hypersurface is null and we take it as the horizon. The third condition leads to the fact that n is the generator of the horizon.

The forth one defines a conformally flat horizon metric. The last one means that $n = \frac{\partial}{\partial u}$ on the horizon. One can consider u as the time direction on the horizon. Two additional conditions are

$$\text{Im}[\mu] = O(r), \quad \lambda = O(r). \quad (14)$$

Those conditions mean that the horizon generator n is twist free and shear free. So u is considered as a good time direction. The last condition is

$$\text{Im}[L^{\bar{z}}] = O(r). \quad (15)$$

This condition can eliminate independent internal Lorentz transformation, which doesn't exist in metric theory, and thus leads the first order theory to be as close as the metric theory. At null infinity, the standard treatment is to fix the boundary metric to be a sphere. However this condition is too strong in the near horizon case which will eliminate near horizon supertranslation.

The advantage of computing near horizon surface charge in first order formalism is the fact that the leading order charge only involves the leading order fields. So one does not really need to solve the radial NP equations as the leading order fields are integration constants which are free data.² For completeness, we work out one more order solutions of the radial equations which would be useful for computing the subleading near horizon charges. The full vacuum NP equations are listed in Appendix B. The details of solving the NP equations are given in Appendix C. The solutions in near horizon expansion are

$$\Psi_0 = \Psi_0^0 + \Psi_0^1 r + O(r^2), \quad (16)$$

$$\rho = \rho_0 + (\rho_0^2 + \sigma_0 \bar{\sigma}_0) r + O(r^2), \quad (17)$$

$$\sigma = \sigma_0 + (2\rho_0 \sigma_0 + \Psi_0^0) r + O(r^2), \quad (18)$$

$$L^z = P \sigma_0 r + \frac{1}{2} P (4\rho_0 \sigma_0 + \Psi_0^0) r^2 + O(r^3), \quad (19)$$

$$L^{\bar{z}} = P + P \rho_0 r + \frac{1}{2} P (\rho_0^2 + \sigma_0 \bar{\sigma}_0) r^2 + O(r^3), \quad (20)$$

$$L_z = -\frac{1}{P} + \frac{\rho_0}{P} r + O(r^3), \quad (21)$$

$$L_{\bar{z}} = \frac{\sigma_0}{P} r + \frac{\Psi_0^0}{P} r^2 + O(r^3), \quad (22)$$

$$\alpha = \alpha_0 + (\alpha_0 \rho_0 + \beta_0 \bar{\sigma}_0) r + O(r^2), \quad (23)$$

$$\beta = \beta_0 + (\alpha_0 \sigma_0 + \beta_0 \rho_0 + \Psi_1^0) r + O(r^2), \quad (24)$$

$$\omega = -\tau_0 r + O(r^2), \quad (25)$$

$$\Psi_1 = \Psi_1^0 + (4\rho_0 \Psi_1^0 + \bar{\delta} \Psi_0^0) r + O(r^2), \quad (26)$$

²An r -derivative may appear in the variation of fields along symmetry direction. One just needs to trace the precise radial equations for the involved fields. Nevertheless this will not happen for a fixed boundary case as we will show in the next section.

$$X^z = P\tau_0 r + O(r^2), \quad (27)$$

$$\mu = \mu_0 + (\mu_0\rho_0 + \Psi_2^0)r + O(r^2), \quad (28)$$

$$\lambda = \mu_0\bar{\sigma}_0 r + O(r^2), \quad (29)$$

$$\Psi_2 = \Psi_2^0 + (3\rho_0\Psi_2^0 - \bar{\partial}\Psi_1^0)r + O(r^2), \quad (30)$$

$$\gamma = \gamma_0 + (\alpha_0\bar{\alpha}_0 + \beta_0\bar{\beta}_0 + 2\alpha_0\beta_0 + \Psi_2^0)r + O(r^2), \quad (31)$$

$$U = -(\gamma_0 + \bar{\gamma}_0)r + O(r^2), \quad (32)$$

$$\Psi_3 = \Psi_3^0 + (2\rho_0\Psi_3^0 + \bar{\partial}\Psi_2^0)r + O(r^2), \quad (33)$$

$$\nu = (\alpha_0\mu_0 + \bar{\beta}_0\mu_0 + \Psi_3^0)r + O(r^2), \quad (34)$$

$$\Psi_4 = \Psi_4^0 + (\rho_0\Psi_4^0 + \bar{\partial}\Psi_3^0)r + O(r^2), \quad (35)$$

where quantities with subscript 0 are integration constants in r . The “ $\bar{\partial}$ ” operator is defined by

$$\bar{\partial}\eta^s = P\partial_{\bar{z}}\eta^s + 2s\bar{\alpha}^0\eta^s,$$

$$\bar{\bar{\partial}}\eta^s = P\partial_z\eta^s - 2s\alpha^0\eta^s,$$

where s is the spin weight of the field η . The spin weights of relevant fields are listed in Table 1.

Table 1: Spin weights

	$\bar{\partial}$	∂_u	γ^0	ν^0	μ^0	σ^0	λ^0	Ψ_4^0	Ψ_3^0	Ψ_2^0	Ψ_1^0	Ψ_0^0
s	1	0	0	-1	0	2	-2	-2	-1	0	1	2

The integration constants have the following constraints from the non-radial NP equations³

$$\alpha_0 = \frac{1}{2}(\bar{\tau}_0 + \partial_z P), \quad (36)$$

$$\beta_0 = \frac{1}{2}(\tau_0 - \partial_{\bar{z}} P), \quad (37)$$

$$\mu_0 = -\partial_u \ln P, \quad (38)$$

$$\gamma_0 = -\frac{1}{2}\mu_0 - \frac{\partial_u \mu_0}{2\mu_0}, \quad (39)$$

$$\Psi_4^0 = 0, \quad (40)$$

$$\Psi_3^0 = \bar{\partial}\gamma_0 - \partial_u \alpha_0 - \alpha_0(\gamma_0 - \bar{\gamma}_0 + \mu_0), \quad (41)$$

$$\Psi_2^0 = (\gamma_0\bar{\gamma}_0)\rho_0 - 2\alpha_0\beta_0 - 2\alpha_0\bar{\sigma}_0 - \mu_0\rho_0 + P\partial_z\bar{\alpha}_0 + P\partial_z\beta_0 + \partial_u\rho_0, \quad (42)$$

$$\Psi_1^0 = \bar{\alpha}_0\rho_0 + \alpha_0\sigma_0 + \beta_0\rho_0 - \bar{\partial}\rho_0 + \bar{\bar{\partial}}\sigma_0, \quad (43)$$

³There is one other possibility that $\mu_0 = 0$ and γ_0 is an arbitrary function.

$$\partial_u \tau_0 = -\partial_u \partial_{\bar{z}} P - 2\beta_0 \mu_0 - 4\bar{\alpha}_0 \mu_0 + 2\check{\delta} \gamma_0 - 2\check{\delta} \mu_0, \quad (44)$$

$$\partial_u \rho_0 = \rho_0 (\gamma_0 + \bar{\gamma}_0) - 2\mu_0 \rho_0 + P \partial_{\bar{z}} \alpha_0 + P \partial_z \bar{\alpha}_0 - 3\alpha_0 \bar{\alpha}_0 - \beta_0 \bar{\beta}_0, \quad (45)$$

$$\partial_u \sigma_0 = -2\bar{\alpha}_0 \beta_0 - 2\beta_0^2 + 2\gamma_0 \sigma_0 - \mu_0 \sigma_0 + P \partial_{\bar{z}} \bar{\alpha}_0 + P \partial_z \beta_0. \quad (46)$$

$$\partial_u \Psi_0^0 - 4\gamma_0 \Psi_0^0 + \mu_0 \Psi_0^0 = \check{\delta} \Psi_1^0 - 6\beta_0 \Psi_1^0 - 6\bar{\alpha}_0 \Psi_1^0 + 3\sigma_0 \Psi_2^0. \quad (47)$$

There is no constraint on the time evolution of P which can characterize the flux going through the horizon. The horizon data are μ_0 the expansion, γ_0 the surface gravity and τ_0 the rotation of reference frame on the horizon. The structure of the solution space is quite different from the case at null infinity where it is normally to consider the Weyl tensor as free data, then the spin connection is determined by the Weyl tensor. But on the horizon the Bianchi identities can not lead to a peeling-off property. We do the inverse that the Weyl tensor is determined by the spin connection. Then the time evolution equations from the Bianchi identities are satisfied automatically. The practical reason of such treatment is that we care more about the local geometric structure on the horizon which is encoded in the spin connection.

The solution space we obtained is larger than that in [35] where the authors considered a non-expanding horizon, i.e., $\mu_0 = 0$. While our solution space is smaller than that in [36] where the authors consider the case with the shear of the horizon generator which can also characterize flux going through the horizon. To recover our solution space, one just needs to set the fields in [36] to⁴

$$\mathcal{U}^A \rightarrow 0, \quad \eta \rightarrow 1, \quad \Omega \rightarrow \frac{1}{P^2}, \quad \gamma_{z\bar{z}} \rightarrow -1, \quad \gamma_{zz} \rightarrow 0. \quad (48)$$

and

$$\kappa \rightarrow -2\gamma_0, \quad \Upsilon^z \rightarrow \frac{2\tau_0}{P}. \quad (49)$$

Other simplifications of [36] in this case are

$$\mathcal{D}_v = \partial_v, \quad \Theta_l = \partial_v \ln \Omega \rightarrow 2\mu_0, \quad N^{AB} = 0, \quad \Gamma = -2\kappa + \partial_v \ln \Omega \rightarrow 4\gamma_0 + 2\mu_0. \quad (50)$$

Those relations will be very useful later to compare our charge with the one in [36].

3 Near horizon symmetry

In the NP formalism, the gauge transformation of the tetrad and the spin connection is a combination of a diffeomorphism and a local Lorentz transformation. The transformation is given by

$$\begin{aligned} \delta_{\xi, \omega} e_a^\mu &= \xi^\nu \partial_\nu e_a^\mu - \partial_\nu \xi^\mu e_a^\nu + \omega_a^b e_b^\mu, \\ \delta_{\xi, \omega} \Gamma_{abc} &= \xi^\nu \partial_\nu \Gamma_{abc} - e_c^\mu \partial_\mu \omega_{ab} + \omega_a^d \Gamma_{dbc} + \omega_b^d \Gamma_{adc} + \omega_c^d \Gamma_{abd}. \end{aligned} \quad (51)$$

⁴Such choice of parameters is not included in the discussion in [36]. However, as local functions, they are solution of Einstein equation.

The residual gauge transformation that preserved the gauge and boundary conditions is worked out in Appendix D. The symmetry parameters are given by $f(u, z, \bar{z})$ and $Y(z), \bar{Y}(\bar{z})$ which generate near horizon supertranslations and superrotations respectively. The associated residual gauge transformations are explicitly determined by the symmetry parameters as

$$\begin{aligned}\xi^u &= f, & \xi^A &= Y^A + \partial_B f \int_0^r dr [L^A \bar{L}^B + \bar{L}^A L^B], \\ \xi^r &= -\partial_u f r + \partial_A f \int_0^r dr [\omega \bar{L}^A + \bar{\omega} L^A - X^A],\end{aligned}$$

and

$$\begin{aligned}\omega^{12} &= \partial_u f + X^A \partial_A f, & \omega^{13} &= -\partial_A f \int_0^r dr [\lambda L^A + \mu \bar{L}^A], \\ \omega^{23} &= \bar{L}^A \partial_A f, & \omega^{34} &= \frac{1}{2}(\partial_z Y - \partial_{\bar{z}} \bar{Y}) + \partial_A f \int_0^r dr [(\bar{\alpha} - \beta) \bar{L}^A - (\alpha - \bar{\beta}) L^A].\end{aligned}$$

The constant order in ξ^r is set to be zero by hand. The reason is that we want to fix the $r = 0$ null hypersurface to be the boundary. Somehow this can be understood as the fact that the existence of a boundary at $r = 0$ breaks the translational invariance along r direction [35, 36].

Acting the residual gauge transformation on the near horizon fields yields their transformation law as

$$\delta_{\xi, \omega} \frac{1}{P} = f \partial_u \frac{1}{P} + Y \partial_z \frac{1}{P} + \bar{Y} \partial_{\bar{z}} \frac{1}{P} + \frac{1}{2}(\partial_z Y + \partial_{\bar{z}} \bar{Y}) \frac{1}{P}, \quad (52)$$

$$\delta_{\xi, \omega} \mu_0 = f \partial_u \mu_0 + Y \partial_z \mu_0 + \bar{Y} \partial_{\bar{z}} \mu_0 + \partial_u f \mu_0, \quad (53)$$

$$\delta_{\xi, \omega} \gamma_0 = f \partial_u \gamma_0 + Y \partial_z \gamma_0 + \bar{Y} \partial_{\bar{z}} \gamma_0 + \partial_u f \gamma_0 - \frac{1}{2} \partial_u^2 f, \quad (54)$$

$$\delta_{\xi, \omega} \tau_0 = f \partial_u \tau_0 + Y \partial_z \tau_0 + \bar{Y} \partial_{\bar{z}} \tau_0 - \frac{1}{2}(\partial_z Y - \partial_{\bar{z}} \bar{Y}) \tau_0 + P \partial_{\bar{z}} f \gamma_0 - \partial_u (P \partial_{\bar{z}} f). \quad (55)$$

$$\begin{aligned}\delta_{\xi, \omega} \sigma_0 &= f \partial_u \sigma_0 + Y \partial_z \sigma_0 + \bar{Y} \partial_{\bar{z}} \sigma_0 - (\partial_z Y - \partial_{\bar{z}} \bar{Y}) \sigma_0 - \partial_u f \sigma_0 \\ &\quad + 2P \partial_{\bar{z}} f \tau_0 - 2P \partial_{\bar{z}} f \partial_{\bar{z}} P - P^2 \partial_{\bar{z}}^2 f.\end{aligned} \quad (56)$$

$$\delta_{\xi, \omega} \rho_0 = f \partial_u \rho_0 + Y \partial_z \rho_0 + \bar{Y} \partial_{\bar{z}} \rho_0 - \partial_u f \rho_0 + P \partial_{\bar{z}} f \bar{\tau}_0 + P \partial_z f \tau_0 - P^2 \partial_z \partial_{\bar{z}} f. \quad (57)$$

4 Near horizon charges

In this section, we will compute the surface charge defined in [19] for the Palatini action, Holst term, Pontryagin term, and Gauss-Bonnet term.

4.1 Palatini action

The Palatini Lagrangian is

$$L_{Pa} = \frac{1}{32\pi G} \epsilon_{abcd} R^{ab} \wedge e^c \wedge e^d, \quad (58)$$

where $R^{ab} = d\Gamma^{ab} + \Gamma^{ac} \wedge \Gamma_c^b$ is the curvature tensor. The surface charge from this Lagrangian is defined by⁵

$$\delta\mathcal{H}_{Pa} = \frac{1}{16\pi G} \epsilon_{abcd} \int_{\partial\Sigma} (i_\xi \Gamma^{ab} \delta e^c \wedge e^d + i_\xi e^c \delta \Gamma^{ab} \wedge e^d - \omega^{ab} \delta e^c \wedge e^d), \quad (59)$$

where $\partial\Sigma$ can be any constant- u two surface on the horizon to evaluate the surface charge. The symmetry parameters should be field independent which was assumed to obtain this expression in [19]. Inserting the solutions and the symmetry parameters into the charge gives

$$\begin{aligned} \delta\mathcal{H}_{Pa} = \frac{1}{8\pi G} \int_{\partial\Sigma} dz d\bar{z} \delta \left[\frac{1}{P^2} \left(\partial_u f - 2f\mu_0 - Y \frac{\bar{\tau}_0}{P} - \bar{Y} \frac{\tau_0}{P} \right) \right] \\ + \frac{1}{8\pi G} \int_{\partial\Sigma} dz d\bar{z} f (\mu_0 - 2\gamma_0) \delta \frac{1}{P^2}. \end{aligned} \quad (60)$$

This charge matches the one in [36] explicitly using the relations by the end of Section 2. Note that fixing $\eta = 1$ requires that $W = 2\partial_v T$ in [36] which also matches our near horizon symmetry generators in Section 3.

4.2 Holst term

The Holst term is

$$L_H = \frac{it}{16\pi G} R_{ab} \wedge e^a \wedge e^b, \quad (61)$$

where t is the Holst term parameter. This term is not a boundary term. But the solution to the equations of motion of the Palatini action is also solution to the Palatini-Holst one. The contribution in the charge from this term is given by [19]

$$\delta\mathcal{H}_H = \frac{it}{8\pi G} \int_{\partial\Sigma} (i_\xi \Gamma^{ab} \delta e_a \wedge e_b + i_\xi e_a \delta \Gamma^{ab} \wedge e_b - \omega^{ab} \delta e_a \wedge e_b). \quad (62)$$

When the near horizon solution and symmetries are inserted, the near horizon Holst charge reads

$$\delta\mathcal{H}_H = \frac{it}{8\pi G} \int_{\partial\Sigma} dz d\bar{z} \delta \left[\frac{1}{P^2} \left(\bar{Y} \frac{\partial_{\bar{z}} P}{P} - Y \frac{\partial_z P}{P} + \frac{1}{2} (\partial_z Y - \partial_{\bar{z}} \bar{Y}) \right) \right]. \quad (63)$$

It is interesting to note that the supertranslation charge is zero. This means that the dual mass as the zero mode of the supertranslation charge vanishes. This can be understood from the fact that the dual mass is located at the ‘‘conjugate’’ point of the mass. The mass is inside the horizon while the dual mass should be outside the horizon.

⁵There is minus sign missing for the Lorentz charge in [19].

4.3 Pontryagin term

The Pontryagin term is given by

$$L_P = \frac{1}{32\pi G} R_{ab} \wedge R^{ab}. \quad (64)$$

This term can be written as a boundary term on the horizon

$$L_{Pb} = \frac{1}{32\pi G} d \left(\Gamma^{ab} \wedge d\Gamma_{ab} - \frac{2}{3} \Gamma^a{}_b \wedge \Gamma^b{}_c \wedge \Gamma^c{}_a \right). \quad (65)$$

A boundary term can only contribute to the symplectic structure through the corner symplectic potential [12, 13]. The corner symplectic potential from (65) is

$$\vartheta_P = \frac{1}{32\pi G} \Gamma^{ab} \wedge \delta\Gamma_{ab}. \quad (66)$$

The surface charge obtained from this corner symplectic potential is

$$\delta\mathcal{H}_P = \frac{1}{16\pi G} \int_{\partial\Sigma} \delta\Gamma^{ab} \wedge \delta_{\xi,\omega} \Gamma_{ab} \quad (67)$$

Alternatively in [18], a symplectic potential was derived directly from the variation of (64), which is

$$\theta_P = \frac{1}{16\pi G} \delta\Gamma_{ab} \wedge R^{ab}. \quad (68)$$

Eventually, the same surface charge (67) can be obtained. This symplectic potential can be rewritten as

$$\theta_P = \frac{1}{32\pi G} \delta \left(\Gamma^{ab} \wedge d\Gamma_{ab} - \frac{2}{3} \Gamma^a{}_b \wedge \Gamma^b{}_c \wedge \Gamma^c{}_a \right) + \frac{1}{32\pi G} d(\Gamma_{ab} \wedge \delta\Gamma^{ab}). \quad (69)$$

The two terms on the right hand side are a W term and a Y term in symplectic potential. Thus the Pontryagin charge in [18] can be obtained from the ambiguities of the symplectic structure.

In the covariant phase space method, a boundary term can only lead to a W term in the symplectic potential. It is very surprising that a Y term arises from the Pontryagin term. Actually, the Lagrangian (64) is different from (65) with an extra trivial term

$$y_P = \frac{1}{32\pi G} \Gamma^{ab} \wedge d^2\Gamma_{ab}. \quad (70)$$

In the variation of (64), another trivial term⁶

$$\frac{1}{16\pi G} \delta\Gamma^{ab} \wedge d^2\Gamma_{ab} \quad (71)$$

⁶This term is the equation of motion from the variation of (64).

has been dropped. Including those two terms, the variation of (64) leads to

$$\delta(L_P - y_P) = \frac{1}{32\pi G} d\delta \left(\Gamma^{ab} \wedge d\Gamma_{ab} - \frac{2}{3} \Gamma^a{}_b \wedge \Gamma^b{}_c \wedge \Gamma^c{}_a \right). \quad (72)$$

This will only lead to a W term in the symplectic potential as it should be when considering (65).

As demonstrated in [18], first order formalism is best suited to an analysis of surface charges, in particular for tracing the contribution from topological terms. However such idea was not implemented explicitly to the Pontryagin term. Here we fill in this gap and write the Pontryagin term in a complete first order form. We consider the curvature tensor as independent dynamical variable. The relation between the curvature tensor and the spin connection is a consequence of equation of motion. So the curvature tensor is an auxiliary field. The Lagrangian that is on-shell equivalent to the Pontryagin one is

$$L_{PF} = \frac{1}{16\pi G} \left[\mathcal{R}_{ab} \wedge (d\Gamma^{ab} + \Gamma^{ac} \wedge \Gamma_c{}^b) - \frac{1}{2} \mathcal{R}_{ab} \wedge \mathcal{R}^{ab} \right]. \quad (73)$$

The equations of motion derived from this Lagrangian are as follows. The variation on \mathcal{R}_{ab} will lead to

$$\mathcal{R}^{ab} = d\Gamma^{ab} + \Gamma^{ac} \wedge \Gamma_c{}^b. \quad (74)$$

Hence \mathcal{R}^{ab} is the curvature tensor. Then variation on Γ^{ab} from this part will lead to the Bianchi identity of the curvature tensor once (74) applied. So this will not modify the equation of motion derived from the Palatini action when doing variation on Γ^{ab} . The Lagrangian (73) is not a boundary term with the auxiliary field, the resulting θ term is

$$\theta_{PF} = \frac{1}{16\pi G} \delta\Gamma^{ab} \wedge R_{ab}, \quad (75)$$

which is the same as (68).

The way that we rewrite the Pontryagin term can be generalized to arbitrary fields. In the form language, one can always add the term

$$L_A = 2A \wedge dB - A \wedge A, \quad (76)$$

to a Lagrangian without changing its on-shell solution where A is a two form and B is a one form. Such term can be modified by equation of motion into a boundary term in second order form $d(B \wedge dB)$. As highlighted in [18], one should include all that type of terms when finding asymptotic gravitational charges. The symplectic potential derived from (76) is

$$\theta_A = 2A \wedge \delta B. \quad (77)$$

On-shell this term becomes

$$\theta_A = 2dB \wedge \delta B. \quad (78)$$

Such term can be written as a combination of a Y term and a W term

$$\theta_A = 2dB \wedge \delta B = d(B \wedge \delta B) + \delta(dB \wedge B). \quad (79)$$

Thus we have shown that the symplectic potential derived from a general glass of terms that do not affect the equations of motion for the theory of interest can be obtained from the ambiguities of the symplectic structure.

Inserting the near horizon solution and symmetry parameters from previous sections into (67), the Pontryagin charge is obtained as

$$\begin{aligned} \delta \mathcal{H}_P = \frac{1}{8\pi G} \int_{\partial\Sigma} dz d\bar{z} \left[\delta_{\xi,\omega} \frac{\bar{\tau}_0}{P} \delta \frac{\tau_0}{P} - \delta \frac{\bar{\tau}_0}{P} \delta_{\xi,\omega} \frac{\tau_0}{P} \right. \\ \left. + \delta \partial_z \ln P \delta_{\xi,\omega} \partial_{\bar{z}} \ln P - \delta_{\xi,\omega} \partial_z \ln P \delta \partial_{\bar{z}} \ln P \right], \quad (80) \end{aligned}$$

where $\delta_{\xi,\omega} \frac{1}{P}$ and $\delta_{\xi,\omega} \tau_0$ are given in (52) and (55). We do not insert the precise formula here as one can always change the representative of the symmetry parameters [43–45] to simplify the expression of charge and the charge algebra structure which needs to be stressed elsewhere. This charge is purely imaginary. So an imaginary unit i should be included in the coupling constant for the Pontryagin term similar to the Holst case.

4.4 Gauss-Bonnet term

The Gauss-Bonnet term is given by

$$L_{GB} = \frac{1}{32\pi G} \epsilon_{abcd} R^{ab} \wedge R^{cd}. \quad (81)$$

This term can be written as a boundary term as

$$L_{GBb} = \frac{1}{32\pi G} \epsilon_{abcd} d(\Gamma^{ab} \wedge d\Gamma^{cd}). \quad (82)$$

The corner symplectic potential from (82) is

$$\vartheta_{GB} = \frac{1}{32\pi G} \epsilon_{abcd} \Gamma^{ab} \wedge \delta \Gamma^{cd}. \quad (83)$$

The surface charge derived from this corner symplectic potential is

$$\delta \mathcal{H}_{GB} = \frac{1}{16\pi G} \epsilon_{abcd} \int_{\partial\Sigma} \delta \Gamma^{ab} \wedge \delta_{\xi,\omega} \Gamma^{cd}. \quad (84)$$

This surface charge can be also obtained from the symplectic potential derived from the variation of (81). The symplectic potential of (81) is [18]

$$\theta_{GB} = \frac{1}{16\pi G} \epsilon_{abcd} \delta \Gamma^{ab} \wedge R^{cd}, \quad (85)$$

which can be written as the sum of a W term and a Y term

$$\theta_{GB} = \frac{1}{32\pi G} \epsilon_{abcd} \delta (\Gamma^{ab} \wedge d\Gamma^{cd}) + \frac{1}{32\pi G} \epsilon_{abcd} d(\Gamma^{ab} \wedge \delta\Gamma^{cd}). \quad (86)$$

Eventually, the same surface charge (84) can be derived from this symplectic potential. Similar to the Pontryagin case, the Lagrangian (81) can be written as a boundary term, i.e., (82), plus an extra trivial term

$$y_{GB} = \frac{1}{32\pi G} \epsilon_{abcd} \Gamma^{ab} \wedge d^2\Gamma^{cd}. \quad (87)$$

Including this term and keeping the trivial term from the equation of motion, the variation of (81) leads to

$$\delta (L_{GB} - y_{GB}) = \frac{1}{32\pi G} \epsilon_{abcd} d\delta (\Gamma^{ab} \wedge d\Gamma^{cd}), \quad (88)$$

as it should be when considering the Lagrangian (82).

The surface charge (84) can also be deduced from a first order form as proposed in previous subsection. The first order term that leads to (84) is

$$L_{GBF} = \frac{1}{16\pi G} \epsilon_{abcd} \left[\mathcal{R}^{ab} \wedge d\Gamma^{cd} - \frac{1}{2} \mathcal{R}^{ab} \wedge \mathcal{R}^{cd} \right]. \quad (89)$$

This term is not equivalent to the Gauss-Bonnet term in the sense that \mathcal{R}^{ab} is not the curvature tensor on-shell. However if the charge from the Gauss-Bonnet term has any interesting feature in four dimensions, one can not distinguish that charge is from the Gauss-Bonnet term or from (89).

Inserting the solution and symmetry parameters into the surface charge (84), we obtain

$$\begin{aligned} \oint \mathcal{H}_{GB} = \frac{1}{8\pi G} \int_{\partial\Sigma} dz d\bar{z} \left[\delta \frac{\bar{\tau}_0}{P} \delta_{\xi,\omega} \partial_{\bar{z}} \ln P + \delta \frac{\tau_0}{P} \delta_{\xi,\omega} \partial_z \ln P + 2\delta \frac{\rho_0}{P} \delta_{\xi,\omega} \frac{\mu_0}{P} \right. \\ \left. - 2\delta \frac{\mu_0}{P} \delta_{\xi,\omega} \frac{\rho_0}{P} - \delta_{\xi,\omega} \frac{\bar{\tau}_0}{P} \delta \partial_{\bar{z}} \ln P - \delta_{\xi,\omega} \frac{\tau_0}{P} \delta \partial_z \ln P \right]. \quad (90) \end{aligned}$$

Here $\delta_{\xi,\omega} \frac{1}{P}$, $\delta_{\xi,\omega} \mu_0$, $\delta_{\xi,\omega} \tau_0$ and $\delta_{\xi,\omega} \rho_0$ are given in (52), (53), (55) and (57) respectively. The Gauss-Bonnet charge is real. It is very surprising that ρ_0 term appears in the charge. The geometric meaning of ρ_0 is the expansion of l which is another null direction other than the horizon generator n . So ρ_0 does not represent any horizon information. The interpretation of this term in a horizon charge is not clear and should be stressed elsewhere.

5 Conclusion

We have shown that the surface charge from the Pontryagin term and Gauss-Bonnet term can also arise from a combination of Y and W ambiguity of the symplectic structure. The

topological terms, such as Pontryagin term and Gauss-Bonnet term have significant effect on the near horizon charge.

There could be a couple of applications of the near horizon charge from the topological terms. We point out some of them for future investigations, for instance in null hypersurface thermodynamics [34], in the triangle relation [46] and in resolving the black hole information paradox [47].

Acknowledgments

The authors thank Mahdi Godazgar, Hong Lü, Shahin Sheikh-Jabbari, Jun-Bao Wu and Xiaoning Wu for discussions and comments. This work is supported in part by the National Natural Science Foundation of China under Grant No. 11905156, No. 12075166, No. 11675144, and No. 11935009.

A Useful relations in Newman-Penrose formalism

The components of spin connection

$$\begin{aligned}
\kappa &= \Gamma_{311} = l^\nu m^\mu \nabla_\nu l_\mu, & \pi &= -\Gamma_{421} = -l^\nu \bar{m}^\mu \nabla_\nu n_\mu, \\
\epsilon &= \frac{1}{2}(\Gamma_{211} - \Gamma_{431}) = \frac{1}{2}(l^\nu n^\mu \nabla_\nu l_\mu - l^\nu \bar{m}^\mu \nabla_\nu m_\mu), \\
\tau &= \Gamma_{312} = n^\nu m^\mu \nabla_\nu l_\mu, & \nu &= -\Gamma_{422} = -n^\nu \bar{m}^\mu \nabla_\nu n_\mu, \\
\gamma &= \frac{1}{2}(\Gamma_{212} - \Gamma_{432}) = \frac{1}{2}(n^\nu n^\mu \nabla_\nu l_\mu - n^\nu \bar{m}^\mu \nabla_\nu m_\mu), \\
\sigma &= \Gamma_{313} = m^\nu m^\mu \nabla_\nu l_\mu, & \mu &= -\Gamma_{423} = -m^\nu \bar{m}^\mu \nabla_\nu n_\mu, \\
\beta &= \frac{1}{2}(\Gamma_{213} - \Gamma_{433}) = \frac{1}{2}(m^\nu n^\mu \nabla_\nu l_\mu - m^\nu \bar{m}^\mu \nabla_\nu m_\mu), \\
\rho &= \Gamma_{314} = \bar{m}^\nu m^\mu \nabla_\nu l_\mu, & \lambda &= -\Gamma_{424} = -\bar{m}^\nu \bar{m}^\mu \nabla_\nu n_\mu, \\
\alpha &= \frac{1}{2}(\Gamma_{214} - \Gamma_{434}) = \frac{1}{2}(\bar{m}^\nu n^\mu \nabla_\nu l_\mu - \bar{m}^\nu \bar{m}^\mu \nabla_\nu m_\mu).
\end{aligned}$$

Ten components of the Weyl tensor

$$\Psi_0 = -C_{1313}, \quad \Psi_1 = -C_{1213}, \quad \Psi_2 = -C_{1342}, \quad \Psi_3 = -C_{1242}, \quad \Psi_4 = -C_{2324}.$$

Four real components of the Ricci tensor

$$\begin{aligned}\Phi_{00} &= -\frac{1}{2}R_{11}, & \Phi_{22} &= -\frac{1}{2}R_{22}, \\ \Phi_{11} &= -\frac{1}{4}(R_{12} + R_{34}), & \Lambda &= \frac{1}{24}R = \frac{1}{12}(R_{12} - R_{34}),\end{aligned}$$

where Λ is the cosmological constant and three complex ones

$$\begin{aligned}\Phi_{02} &= -\frac{1}{2}R_{33}, & \Phi_{20} &= -\frac{1}{2}R_{44}, \\ \Phi_{01} &= -\frac{1}{2}R_{13}, & \Phi_{10} &= -\frac{1}{2}R_{14}, \\ \Phi_{12} &= -\frac{1}{2}R_{23}, & \Phi_{21} &= -\frac{1}{2}R_{24}.\end{aligned}$$

The orthogonality conditions and normalization conditions (1) of the basis vectors yield the following relations

$$\begin{aligned}l^\nu \nabla_\nu l_\mu &= (\epsilon + \bar{\epsilon})l_\mu - \kappa \bar{m}_\mu - \bar{\kappa} m_\mu, \\ n^\nu \nabla_\nu l_\mu &= (\gamma + \bar{\gamma})l_\mu - \tau \bar{m}_\mu - \bar{\tau} m_\mu, \\ m^\nu \nabla_\nu l_\mu &= (\beta + \bar{\alpha})l_\mu - \sigma \bar{m}_\mu - \bar{\rho} m_\mu, \\ \bar{m}^\nu \nabla_\nu l_\mu &= (\alpha + \bar{\beta})l_\mu - \rho \bar{m}_\mu - \bar{\sigma} m_\mu,\end{aligned}\tag{91}$$

$$\begin{aligned}l^\nu \nabla_\nu n_\mu &= -(\epsilon + \bar{\epsilon})n_\mu + \bar{\pi} \bar{m}_\mu + \pi m_\mu, \\ n^\nu \nabla_\nu n_\mu &= -(\gamma + \bar{\gamma})n_\mu + \bar{\nu} \bar{m}_\mu + \nu m_\mu, \\ m^\nu \nabla_\nu n_\mu &= -(\beta + \bar{\alpha})n_\mu + \bar{\lambda} \bar{m}_\mu + \mu m_\mu, \\ \bar{m}^\nu \nabla_\nu n_\mu &= -(\alpha + \bar{\beta})n_\mu + \bar{\mu} \bar{m}_\mu + \lambda m_\mu,\end{aligned}\tag{92}$$

$$\begin{aligned}l^\nu \nabla_\nu m_\mu &= (\epsilon - \bar{\epsilon})m_\mu - \kappa n_\mu + \bar{\pi} l_\mu, \\ n^\nu \nabla_\nu m_\mu &= (\gamma - \bar{\gamma})m_\mu - \tau n_\mu + \bar{\nu} l_\mu, \\ m^\nu \nabla_\nu m_\mu &= (\beta - \bar{\alpha})m_\mu - \sigma n_\mu + \bar{\lambda} l_\mu, \\ \bar{m}^\nu \nabla_\nu m_\mu &= (\alpha - \bar{\beta})m_\mu - \rho n_\mu + \bar{\mu} l_\mu,\end{aligned}\tag{93}$$

After setting $\epsilon = \kappa = \pi = 0$, the geodesic deviation of l is

$$\begin{aligned}B_{\nu\mu}^l &= \nabla_\nu l_\mu = (\gamma + \bar{\gamma})l_\mu l_\nu - \tau \bar{m}_\mu l_\nu - \bar{\tau} m_\mu l_\nu - (\beta + \bar{\alpha})l_\mu \bar{m}_\nu - (\bar{\beta} + \alpha)l_\mu m_\nu \\ &\quad + \sigma \bar{m}_\mu \bar{m}_\nu + \bar{\rho} m_\mu \bar{m}_\nu + \bar{\sigma} m_\mu m_\nu + \rho \bar{m}_\mu m_\nu.\end{aligned}\tag{94}$$

The transverse part is

$$\hat{B}_{\nu\mu}^l = \sigma \bar{m}_\mu \bar{m}_\nu + \bar{\rho} m_\mu \bar{m}_\nu + \bar{\sigma} m_\mu m_\nu + \rho \bar{m}_\mu m_\nu.\tag{95}$$

It is clear that $\rho = \bar{\rho}$ means l is hypersurface orthogonal and further setting $\tau = \bar{\alpha} + \beta$ leads to the fact that $\nabla_\nu l_\mu$ is a symmetric tensor, then l is the gradient of a scalar field.

On the horizon $r = 0$, n is tangent to geodesic. The geodesic deviation of n is

$$B_{\nu\mu}^n = \nabla_\nu n_\mu = -(\gamma + \bar{\gamma})n_\mu l_\nu + \bar{\nu}\bar{m}_\mu l_\nu + \nu m_\mu l_\nu + (\bar{\alpha} + \beta)n_\mu \bar{m}_\nu + (\alpha + \bar{\beta})n_\mu m_\nu - \bar{\lambda}\bar{m}_\mu \bar{m}_\nu - \lambda m_\mu m_\nu - \mu m_\mu \bar{m}_\nu - \bar{\mu}\bar{m}_\mu m_\nu. \quad (96)$$

The transverse part is

$$\hat{B}_{\nu\mu}^n = -\bar{\lambda}\bar{m}_\mu \bar{m}_\nu - \lambda m_\mu m_\nu - \mu m_\mu \bar{m}_\nu - \bar{\mu}\bar{m}_\mu m_\nu. \quad (97)$$

B NP equations

Considered as directional derivatives, the basis vectors are assigned with special symbols

$$D = l^\mu \partial_\mu, \quad \Delta = n^\mu \partial_\mu, \quad \delta = m^\mu \partial_\mu. \quad (98)$$

Radial equations

$$D\rho = \rho^2 + \sigma\bar{\sigma}, \quad (99)$$

$$D\sigma = 2\rho\sigma + \Psi_0, \quad (100)$$

$$D\tau = \tau\rho + \bar{\tau}\sigma + \Psi_1, \quad (101)$$

$$D\alpha = \rho\alpha + \beta\bar{\sigma}, \quad (102)$$

$$D\beta = \alpha\sigma + \rho\beta + \Psi_1, \quad (103)$$

$$D\gamma = \tau\alpha + \bar{\tau}\beta + \Psi_2, \quad (104)$$

$$D\lambda = \rho\lambda + \bar{\sigma}\mu, \quad (105)$$

$$D\mu = \rho\mu + \sigma\lambda + \Psi_2, \quad (106)$$

$$D\nu = \bar{\tau}\mu + \tau\lambda + \Psi_3, \quad (107)$$

$$DU = \bar{\tau}\omega + \tau\bar{\omega} - (\gamma + \bar{\gamma}), \quad (108)$$

$$DX^A = \bar{\tau}L^A + \tau\bar{L}^A, \quad (109)$$

$$D\omega = \rho\omega + \sigma\bar{\omega} - \tau, \quad (110)$$

$$DL^A = \rho L^A + \sigma\bar{L}^A, \quad (111)$$

$$D\Psi_1 - \bar{\delta}\Psi_0 = 4\rho\Psi_1 - 4\alpha\Psi_0, \quad (112)$$

$$D\Psi_2 - \bar{\delta}\Psi_1 = 3\rho\Psi_2 - 2\alpha\Psi_1 - \lambda\Psi_0, \quad (113)$$

$$D\Psi_3 - \bar{\delta}\Psi_2 = 2\rho\Psi_3 - 2\lambda\Psi_1, \quad (114)$$

$$D\Psi_4 - \bar{\delta}\Psi_3 = \rho\Psi_4 + 2\alpha\Psi_3 - 3\lambda\Psi_2, \quad (115)$$

Non-radial equations

$$\Delta\lambda = \bar{\delta}\nu - (\mu + \bar{\mu})\lambda - (3\gamma - \bar{\gamma})\lambda + 2\alpha\nu - \Psi_4, \quad (116)$$

$$\Delta\rho = \bar{\delta}\tau - \rho\bar{\mu} - \sigma\lambda - 2\alpha\tau + (\gamma + \bar{\gamma})\rho - \Psi_2, \quad (117)$$

$$\Delta\alpha = \bar{\delta}\gamma + \rho\nu - (\tau + \beta)\lambda + (\bar{\gamma} - \gamma - \bar{\mu})\alpha - \Psi_3, \quad (118)$$

$$\Delta\mu = \delta\nu - \mu^2 - \lambda\bar{\lambda} - (\gamma + \bar{\gamma})\mu + 2\beta\nu, \quad (119)$$

$$\Delta\beta = \delta\gamma - \mu\tau + \sigma\nu + \beta(\gamma - \bar{\gamma} - \mu) - \alpha\bar{\lambda}, \quad (120)$$

$$\Delta\sigma = \delta\tau - \sigma\mu - \rho\bar{\lambda} - 2\beta\tau + (3\gamma - \bar{\gamma})\sigma, \quad (121)$$

$$\Delta\omega = \delta U + \bar{\nu} - \bar{\lambda}\bar{\omega} + (\gamma - \bar{\gamma} - \mu)\omega, \quad (122)$$

$$\Delta L^A = \delta X^A - \bar{\lambda}\bar{L}^A + (\gamma - \bar{\gamma} - \mu)L^A, \quad (123)$$

$$\delta\rho - \bar{\delta}\sigma = \rho\tau - \sigma(3\alpha - \bar{\beta}) - \Psi_1, \quad (124)$$

$$\delta\alpha - \bar{\delta}\beta = \mu\rho - \lambda\sigma + \alpha\bar{\alpha} + \beta\bar{\beta} - 2\alpha\beta - \Psi_2, \quad (125)$$

$$\delta\lambda - \bar{\delta}\mu = \mu\bar{\tau} + \lambda(\bar{\alpha} - 3\beta) - \Psi_3, \quad (126)$$

$$\delta\bar{\omega} - \bar{\delta}\omega = \mu - \bar{\mu} - (\alpha - \bar{\beta})\omega + (\bar{\alpha} - \beta)\bar{\omega}, \quad (127)$$

$$\delta\bar{L}^A - \bar{\delta}L^A = (\bar{\alpha} - \beta)\bar{L}^A - (\alpha - \bar{\beta})L^A, \quad (128)$$

$$\Delta\Psi_0 - \delta\Psi_1 = (4\gamma - \mu)\Psi_0 - (4\tau + 2\beta)\Psi_1 + 3\sigma\Psi_2, \quad (129)$$

$$\Delta\Psi_1 - \delta\Psi_2 = \nu\Psi_0 + (2\gamma - 2\mu)\Psi_1 - 3\tau\Psi_2 + 2\sigma\Psi_3, \quad (130)$$

$$\Delta\Psi_2 - \delta\Psi_3 = 2\nu\Psi_1 - 3\mu\Psi_2 + (2\beta - 2\tau)\Psi_3 + \sigma\Psi_4, \quad (131)$$

$$\Delta\Psi_3 - \delta\Psi_4 = 3\nu\Psi_2 - (2\gamma + 4\mu)\Psi_3 + (4\beta - \tau)\Psi_4, \quad (132)$$

C Details for deriving the solution space

The radial equations are organized in different groups. The first group is (99) and (100). Once the whole series of Ψ_0 is given as initial data as (16), ρ and σ are solved out as (17) and (18). Inserting the solutions of ρ and σ into (111), one gets L^A as (19) and (20), then L_A is derived by the condition $L_A L^A = 0$, $L_A \bar{L}^A = -1$. The second group of radial equations consists of (102), (103), (110) and (112). One can work out α , β , ω and Ψ_1 as (23), (24), (25), and (26) respectively, then τ from gauge condition $\tau = \bar{\alpha} + \beta$. Inserting τ and L^A into (109), X^A could be obtained as (27). The third group of radial equations include (105), (106), and (113). One can just apply the same method as the first two groups to solve out μ , λ , and Ψ_2 , which are given in (28), (29), and (30). Then, γ is derived from (104) as (31), U is derived from (108) as (32), Ψ_3 is derived from (114) as (33), ν is derived from (107) as (34), and finally Ψ_4 is derived from (115) as (35).

Inserting the solutions of the radial equations into the non-radial equations, more constraints are obtained for the integration constants:

(128) yields (36) and (37).

(123) yields (38).

(119) yields (39).

(116) yields (40).

(118) yields (41).

(117) yields (42).

(124) yields (43).

(126) yields (44).

(125) yields (45).

(121) yields (46).

(129) yields (47).

The rest non-radial equations (120), (122), (123), (127), and (130)-(132) are satisfied automatically.

D Details for deriving the asymptotic symmetry

The gauge conditions yield

- $0 = \delta_{\xi, \omega} e_1^u = -\partial_r \xi^u \implies \xi^u = f(u, z, \bar{z})$.
- $0 = \delta_{\xi, \omega} e_2^u = -e_2^\alpha \partial_\alpha f + \omega^{12} \implies \omega^{12} = \partial_u f + X^A \partial_A f$.
- $0 = \delta_{\xi, \omega} e_3^u = -e_3^\alpha \partial_\alpha f + \omega^{24} \implies \omega^{24} = L^A \partial_A f$.
- $0 = \delta_{\xi, \omega} e_4^u = -e_4^\alpha \partial_\alpha f + \omega^{23} \implies \omega^{23} = \bar{L}^A \partial_A f$.
- $0 = \delta_{\xi, \omega} e_1^r = -e_1^\alpha \partial_\alpha \xi^r + \omega^{2a} e_a^r \implies \xi^r = -\partial_u f r + Z(u, z, \bar{z}) + \partial_A f \int_0^r dr [\omega \bar{L}^A + \bar{\omega} L^A - X^A]$.
- $0 = \delta_{\xi, \omega} e_1^A = -e_1^\alpha \partial_\alpha \xi^A + \omega^{2a} e_a^A \implies \xi^A = Y^A(u, z, \bar{z}) + \partial_B f \int_0^r dr [L^A \bar{L}^B + \bar{L}^A L^B]$.
- $\delta_{\xi, \omega} \bar{\pi} = 0 \iff 0 = \delta_{\xi, \omega} \Gamma_{321} = l^\mu \partial_\mu \omega^{41} + \Gamma_{32a} \omega^{2a} \implies \omega^{14} = \omega_0^{14}(u, z, \bar{z}) - \partial_A f \int_0^r dr [\bar{\lambda} \bar{L}^A + \bar{\mu} L^A]$.
- $\delta_{\xi, \omega} \pi = 0 \iff 0 = \delta_{\xi, \omega} \Gamma_{421} = l^\mu \partial_\mu \omega^{31} + \Gamma_{42a} \omega^{2a} \implies \omega^{13} = \omega_0^{13}(u, z, \bar{z}) - \partial_A f \int_0^r dr [\lambda L^A + \mu \bar{L}^A]$.
- $\delta_{\xi, \omega} (\epsilon - \bar{\epsilon}) = 0 \iff 0 = \delta_{\xi, \omega} \Gamma_{431} = l^\mu \partial_\mu \omega^{43} + \Gamma_{43a} \omega^{2a} \implies \omega^{34} = \omega_0^{34}(u, z, \bar{z}) + \partial_A f \int_0^r dr [(\bar{\alpha} - \beta) \bar{L}^A - (\alpha - \bar{\beta}) L^A]$.

- Direct computation shows that the conditions $\epsilon + \bar{\epsilon} = 0 = \kappa = \bar{\kappa}$, $\rho - \bar{\rho} = 0$ and $\tau - \bar{\alpha} - \beta = 0$ do not lead to new constraints on the symmetry parameters. These conditions hold as a consequence of the tetrad conditions imposed in (4) when the NP equations are satisfied.

Before checking the constraints from the boundary conditions, we set $Z = 0$. The reason is that we want to fix the $r = 0$ null hypersurface to be the boundary. In the near horizon expansion, we will always assume that the horizon is located at the zero value of the radial coordinate. A shift in r direction, i.e., $\tilde{r} = r + Z$ combined with transformations in other coordinates can set $\tilde{r} = 0$ hypersurface to be null but will also change the boundary to be the $\tilde{r} = 0$ hypersurface. Somehow this can be understood as the fact that the existence of a boundary at $r = 0$ breaks the translational invariance along r direction, see also in [35, 36].

The fall-off conditions yield

- $\delta_{\xi, \omega} e_3^r = O(r) \implies \omega_0^{41} = 0.$
- $\delta_{\xi, \omega} e_4^r = O(r) \implies \omega_0^{31} = 0.$
- $\delta_{\xi, \omega} e_2^A = O(r) \implies \partial_u Y^A = 0.$
- $\delta_{\xi, \omega} e_3^z = O(r) \implies \partial_{\bar{z}} Y^z = 0.$
- $\delta_{\xi, \omega} e_4^{\bar{z}} = O(r) \implies \partial_z Y^{\bar{z}} = 0.$
- $\delta_{\xi, \omega} \text{Im}[e_3^{\bar{z}}] = O(r) \implies \omega_0^{43} = \frac{1}{2}(\partial_z Y^z - \partial_{\bar{z}} Y^{\bar{z}}).$

The boundary conditions on e_2^r , λ , ν and $\text{Im}[\mu]$ do not lead to new constraint on the symmetry parameters as confirmed by direct computation.

References

- [1] R. L. Arnowitt, S. Deser, and C. W. Misner, “The Dynamics of general relativity,” *Gen. Rel. Grav.* **40** (2008) 1997–2027, arXiv:gr-qc/0405109.
- [2] H. Bondi, M. G. J. van der Burg, and A. W. K. Metzner, “Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems,” *Proc. Roy. Soc. Lond.* **A269** (1962) 21–52.
- [3] S. W. Hawking, “Particle Creation by Black Holes,” *Commun. Math. Phys.* **43** (1975) 199–220. [Erratum: *Commun. Math. Phys.* 46, 206 (1976)].

- [4] J. D. Bekenstein, “Black holes and entropy,” *Phys. Rev. D* **7** (1973) 2333–2346.
- [5] J. M. Bardeen, B. Carter, and S. W. Hawking, “The Four laws of black hole mechanics,” *Commun. Math. Phys.* **31** (1973) 161–170.
- [6] J. Lee and R. M. Wald, “Local symmetries and constraints,” *J. Math. Phys.* **31** (1990) 725–743.
- [7] V. Iyer and R. M. Wald, “Some properties of Noether charge and a proposal for dynamical black hole entropy,” *Phys. Rev. D* **50** (1994) 846–864, [arXiv:gr-qc/9403028](#).
- [8] R. M. Wald, “Black hole entropy is the Noether charge,” *Phys. Rev. D* **48** no. 8, (1993) R3427–R3431, [arXiv:gr-qc/9307038](#).
- [9] K. Hajian and M. M. Sheikh-Jabbari, “Solution Phase Space and Conserved Charges: A General Formulation for Charges Associated with Exact Symmetries,” *Phys. Rev. D* **93** no. 4, (2016) 044074, [arXiv:1512.05584 \[hep-th\]](#).
- [10] J. W. York, Jr., “Role of conformal three geometry in the dynamics of gravitation,” *Phys. Rev. Lett.* **28** (1972) 1082–1085.
- [11] G. W. Gibbons and S. W. Hawking, “Action Integrals and Partition Functions in Quantum Gravity,” *Phys. Rev. D* **15** (1977) 2752–2756.
- [12] D. Harlow and J.-Q. Wu, “Covariant phase space with boundaries,” *JHEP* **10** (2020) 146, [arXiv:1906.08616 \[hep-th\]](#).
- [13] L. Freidel, M. Geiller, and D. Pranzetti, “Edge modes of gravity. Part I. Corner potentials and charges,” *JHEP* **11** (2020) 026, [arXiv:2006.12527 \[hep-th\]](#).
- [14] L. Freidel, M. Geiller, and D. Pranzetti, “Edge modes of gravity. Part II. Corner metric and Lorentz charges,” *JHEP* **11** (2020) 027, [arXiv:2007.03563 \[hep-th\]](#).
- [15] L. Freidel, M. Geiller, and D. Pranzetti, “Edge modes of gravity. Part III. Corner simplicity constraints,” *JHEP* **01** (2021) 100, [arXiv:2007.12635 \[hep-th\]](#).
- [16] L. Freidel, R. Oliveri, D. Pranzetti, and S. Speziale, “The Weyl BMS group and Einstein’s equations,” *JHEP* **07** (2021) 170, [arXiv:2104.05793 \[hep-th\]](#).
- [17] L. Freidel, R. Oliveri, D. Pranzetti, and S. Speziale, “Extended corner symmetry, charge bracket and Einstein’s equations,” *JHEP* **09** (2021) 083, [arXiv:2104.12881 \[hep-th\]](#).

- [18] H. Godazgar, M. Godazgar, and M. J. Perry, “Asymptotic gravitational charges,” *Phys. Rev. Lett.* **125** no. 10, (2020) 101301, arXiv:2007.01257 [hep-th].
- [19] H. Godazgar, M. Godazgar, and M. J. Perry, “Hamiltonian derivation of dual gravitational charges,” *JHEP* **09** (2020) 084, arXiv:2007.07144 [hep-th].
- [20] M. Godazgar and G. Long, “Higher derivative asymptotic charges and internal Lorentz symmetries,” arXiv:2201.07014 [hep-th].
- [21] S. Hou, T. Zhu, and Z.-H. Zhu, “Conserved charges in Chern-Simons modified theory and memory effects,” arXiv:2112.13049 [gr-qc].
- [22] E. Newman and R. Penrose, “An Approach to gravitational radiation by a method of spin coefficients,” *J. Math. Phys.* **3** (1962) 566–578.
- [23] B. Julia and S. Silva, “Currents and superpotentials in classical gauge invariant theories. 1. Local results with applications to perfect fluids and general relativity,” *Class. Quant. Grav.* **15** (1998) 2173–2215, arXiv:gr-qc/9804029.
- [24] B. Julia and S. Silva, “Currents and superpotentials in classical gauge theories: 2. Global aspects and the example of affine gravity,” *Class. Quant. Grav.* **17** (2000) 4733–4744, arXiv:gr-qc/0005127.
- [25] A. Ashtekar, J. Engle, and D. Sloan, “Asymptotics and Hamiltonians in a First order formalism,” *Class. Quant. Grav.* **25** (2008) 095020, arXiv:0802.2527 [gr-qc].
- [26] A. Corichi, I. Rubalcava, and T. Vukasinac, “Hamiltonian and Noether charges in first order gravity,” *Gen. Rel. Grav.* **46** (2014) 1813, arXiv:1312.7828 [gr-qc].
- [27] T. Jacobson and A. Mohd, “Black hole entropy and Lorentz-diffeomorphism Noether charge,” *Phys. Rev. D* **92** (2015) 124010, arXiv:1507.01054 [gr-qc].
- [28] G. Barnich, P. Mao, and R. Ruzziconi, “Conserved currents in the Cartan formulation of general relativity,” in *About Various Kinds of Interactions: Workshop in honour of Professor Philippe Spindel*. 11, 2016. arXiv:1611.01777 [gr-qc].
- [29] E. De Paoli and S. Speziale, “A gauge-invariant symplectic potential for tetrad general relativity,” *JHEP* **07** (2018) 040, arXiv:1804.09685 [gr-qc].
- [30] G. Barnich, P. Mao, and R. Ruzziconi, “BMS current algebra in the context of the Newman–Penrose formalism,” *Class. Quant. Grav.* **37** no. 9, (2020) 095010, arXiv:1910.14588 [gr-qc].

- [31] R. Oliveri and S. Speziale, “Boundary effects in General Relativity with tetrad variables,” *Gen. Rel. Grav.* **52** no. 8, (2020) 83, arXiv:1912.01016 [gr-qc].
- [32] G. Barnich, P. Mao, and R. Ruzziconi, “Conserved currents in the Palatini formulation of general relativity,” *PoS CORFU2019* (2020) 171, arXiv:2004.15002 [gr-qc].
- [33] T. Jacobson, “Thermodynamics of space-time: The Einstein equation of state,” *Phys. Rev. Lett.* **75** (1995) 1260–1263, arXiv:gr-qc/9504004.
- [34] H. Adami, M. M. Sheikh-Jabbari, V. Taghiloo, and H. Yavartanoo, “Null Surface Thermodynamics,” arXiv:2110.04224 [hep-th].
- [35] L. Donnay, G. Giribet, H. A. Gonzalez, and M. Pino, “Supertranslations and Superrotations at the Black Hole Horizon,” *Phys. Rev. Lett.* **116** no. 9, (2016) 091101, arXiv:1511.08687 [hep-th].
- [36] H. Adami, D. Grumiller, M. M. Sheikh-Jabbari, V. Taghiloo, H. Yavartanoo, and C. Zwickel, “Null boundary phase space: slicings, news & memory,” *JHEP* **11** (2021) 155, arXiv:2110.04218 [hep-th].
- [37] P. Mao, X. Wu, and H. Zhang, “Soft hairs on isolated horizon implanted by electromagnetic fields,” *Class. Quant. Grav.* **34** no. 5, (2017) 055003, arXiv:1606.03226 [hep-th].
- [38] H. Godazgar, M. Godazgar, and C. N. Pope, “New dual gravitational charges,” *Phys. Rev. D* **99** no. 2, (2019) 024013, arXiv:1812.01641 [hep-th].
- [39] S. Chandrasekhar, “The Newman-Penrose formalism,” in *The mathematical theory of black holes*, ch. 1, pp. 40–55. Oxford, UK, 1983.
- [40] D. Grumiller, A. Pérez, M. M. Sheikh-Jabbari, R. Troncoso, and C. Zwickel, “Spacetime structure near generic horizons and soft hair,” *Phys. Rev. Lett.* **124** no. 4, (2020) 041601, arXiv:1908.09833 [hep-th].
- [41] A. Ashtekar, C. Beetle, O. Dreyer, S. Fairhurst, B. Krishnan, J. Lewandowski, and J. Wisniewski, “Isolated horizons and their applications,” *Phys. Rev. Lett.* **85** (2000) 3564–3567, arXiv:gr-qc/0006006.
- [42] A. Ashtekar and B. Krishnan, “Isolated and dynamical horizons and their applications,” *Living Rev. Rel.* **7** (2004) 10, arXiv:gr-qc/0407042.
- [43] H. Adami, M. M. Sheikh-Jabbari, V. Taghiloo, H. Yavartanoo, and C. Zwickel, “Symmetries at null boundaries: two and three dimensional gravity cases,” *JHEP* **10** (2020) 107, arXiv:2007.12759 [hep-th].

- [44] R. Ruzziconi and C. Zwickel, “Conservation and Integrability in Lower-Dimensional Gravity,” *JHEP* **04** (2021) 034, arXiv:2012.03961 [hep-th].
- [45] H. Adami, M. M. Sheikh-Jabbari, V. Taghiloo, H. Yavartanoo, and C. Zwickel, “Chiral Massive News: Null Boundary Symmetries in Topologically Massive Gravity,” *JHEP* **05** (2021) 261, arXiv:2104.03992 [hep-th].
- [46] A. Strominger, “Lectures on the Infrared Structure of Gravity and Gauge Theory,” arXiv:1703.05448 [hep-th].
- [47] S. W. Hawking, M. J. Perry, and A. Strominger, “Soft Hair on Black Holes,” *Phys. Rev. Lett.* **116** no. 23, (2016) 231301, arXiv:1601.00921 [hep-th].