Automatic computation of Feynman integrals containing linear propagators via auxiliary mass flow

Zhi-Feng Liu^{1,*} and Yan-Qing Ma^{1,2,3,†}

¹School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China

²Center for High Energy Physics, Peking University, Beijing 100871, China

³Collaborative Innovation Center of Quantum Matter, Beijing 100871, China

We proposed a recipe to systematically calculate Feynman integrals containing linear propagators using the auxiliary mass flow method. The key of the recipe is to introduce a quadratic term for each linear propagator and then using differential equations to get rid of their effects. As an application, we calculated all master integrals of vacuum integrals containing a gauge link up to four loops, and we checked the results by nontrivial dimensional recurrence relations.

I. INTRODUCTION

As particle physics enters the era of precision, it is necessary to compute perturbative quantum field theory to high order in coupling constants. To this end, the evaluation of Feynman integrals is an unavoidable task. The main strategy at present is to decompose all Feynman integrals in a problem to a small set of bases, called master integrals (MIs), using integration-by-parts (IBP) reduction [1-22], and then calculate these MIs.

Many methods exist to calculate MIs [2, 23– 60]. Among them, the auxiliary mass flow (AMF) method [57–60], a special case of differential equations (DEs) method, calculates MIs by setting up and solving DEs of MIs with respect to an auxiliary mass term η (denoted as η -DEs). This is systematic and efficient, as far as η -DEs can be set up by using reduction strategy. Equipped with an iterative strategy [59] and the blocktriangular reduction form [11, 12], the AMF method becomes so powerful that many Feynman integrals in cutting-edge problems, which are very challenging for other methods, can be calculated (see Fig. 4 in Ref. [59]). At the same time, the AMF method has already been used in many physical processes [61–68].

The AMF method is systematic only if boundary conditions of the η -DEs at $\eta \to \infty$ can be systematically obtained. If all inverse propagators defining Feynman integrals are quadratic in loop momenta, where these kinds of Feynman integrals are called *quadratic integrals*, then regions at $\eta \to \infty$ can be systematically identified [69, 70], and the contribution of each region can be calculated within the AMF framework [71]. However, life becomes not that easy when some inverse propagators defining Feynman integrals are linear in loop momenta, where these kinds of Feynman integrals are called *linear integrals*. In this case, identification of regions at $\eta \to \infty$ is not that systematic, and the calculation of each region becomes nontrivial [61, 62]. As linear integrals show up frequently in region expansion [69] and in effective field theories, it will be good if they can be calculated systematically by the powerful AMF method.

In this paper, by introducing an auxiliary quadratic term for each linear propagator, we develop a recipe to calculate any linear integral systematically within the AMF framework. In the rest of the paper, we first present the general idea of our recipe and then give some examples to illustrate its details. As an application, we calculate all master integrals of four-loop vacuum integrals containing a gauge link for the first time, which are useful for studying parton distribution functions [72–76]. All results in this paper have been checked by using dimensional recurrence relations [36, 77, 78].

II. THE RECIPE

Consider a general Feynman integral

$$I_{\vec{\nu}} = \prod_{j=1}^{L} \int \frac{\mathrm{d}^D \ell_j}{\mathrm{i}\pi^{D/2}} \prod_{a=1}^{N} \mathcal{D}_a^{-\nu_a},\tag{1}$$

where L is the number of loops and N is the number of propagators and irreducible scalar products, and \mathcal{D}_a are inverse propagators and $\vec{\nu} = (\nu_1, \nu_2, \cdots, \nu_N)$ integers. We introduce a corresponding auxiliary quadratic integral $\tilde{I}_{\vec{\nu}}(x)$ defined by

$$\widetilde{I}_{\vec{\nu}}(x) = \prod_{j=1}^{L} \int \frac{\mathrm{d}^{D}\ell_{j}}{\mathrm{i}\pi^{D/2}} \prod_{a=1}^{N} \widetilde{\mathcal{D}}_{a}^{-\nu_{a}}.$$
(2)

If \mathcal{D}_a is an irreducible scalar product or is quadratic in loop momenta, then $\widetilde{\mathcal{D}}_a = \mathcal{D}_a$. Otherwise, it can be generally expressed as ¹

$$\mathcal{D}_a = p_a \cdot \left(\sum_{j=1}^L c_{aj}\ell_j\right) + \Delta_a + \mathrm{i}0^+$$

^{*} xiangshui@pku.edu.cn

[†] yqma@pku.edu.cn

¹ If \mathcal{D}_a has a negative infinitesimal imaginary part, we can simply substitute it by $-\mathcal{D}_a$, leaving an overall factor of -1.

$$\equiv p_a \cdot \ell_a + \Delta_a + \mathrm{i}0^+, \tag{3}$$

where p_a is a fixed Lorentz vector, and Δ_a and c_{aj} are fixed Lorentz scalars. We then define

$$\widetilde{\mathcal{D}}_{a} = x \left(\ell_{a}^{2} + \mathrm{i}0^{+}\right) + p_{a} \cdot \ell_{a} + \Delta_{a}$$
$$= x \left[\left(\ell_{a} + \frac{p_{a}}{2x}\right)^{2} + \frac{\Delta_{a}}{x} - \frac{p_{a}^{2}}{4x^{2}} + \mathrm{i}0^{+}\right].$$
(4)

Clearly, the original integral can be obtained by

$$I_{\vec{\nu}} = \lim_{x \to 0^+} \widetilde{I}_{\vec{\nu}}(x), \tag{5}$$

which correctly recovers the Feynman prescription.

Denoting MIs of the obtained auxiliary quadratic integrals as $\overrightarrow{J}(x)$, by using IBP reduction, we can get the following reduction relation ²,

$$\widetilde{I}_{\vec{\nu}}(x) = \sum_{i=1}^{n} c_{\vec{\nu};i}(x) \widetilde{J}_{i}(x), \qquad (6)$$

valid for any value of x, where n is the number of MIs and $c_{\vec{\nu};i}(x)$ are rational functions of x. Here, $\tilde{I}_{\vec{\nu}}(0^+)$ can be achieved once $\vec{J}(x)$ are known. Again using IBP, we can set up a system of DEs for MIs with respect to x,

$$\frac{\partial}{\partial x}\overrightarrow{\widetilde{J}}(x) = A(x)\overrightarrow{\widetilde{J}}(x),\tag{7}$$

where A(x) is a $n \times n$ matrix with elements being rational functions of x. Because Feynman integrals have Feynman parametric representation, then $\overrightarrow{J}(x)$ have the following asymptotic expansion (see e.g. Ref. [44])

$$\widetilde{J}_{k}(x) = \sum_{s} \sum_{i=0}^{n_{s}} x^{s} \ln^{i}(x) \sum_{j=0}^{\infty} \widetilde{J}_{k}^{s\,i\,j} x^{j}, \qquad (8)$$

where s is a linear function of space-time dimension $D = 4 - 2\epsilon$ and n_s an integer determined by s, and $\widetilde{J}_k^{s\,i\,j}$ are functions of ϵ . Taking the limit $x \to 0^+$ is then straight forward ³

$$\widetilde{J}_k(0^+) = \widetilde{J}_k^{0\,0\,0}\,,\tag{9}$$

$$\widetilde{J}'_k(0^+) = \widetilde{J}^{0\,0\,1}_k, \tag{10}$$

and so on. Then based on Eq. (6), we can get any desired integral $I_{\vec{\nu}} = \widetilde{I}_{\vec{\nu}}(0^+)$.

Taking advantage of the DEs (7), there are only n independent $\widetilde{J}_{k}^{s\,i\,j}$, which can be fixed by matching the expansion (8) with values at a given point $\overrightarrow{J}(x_{0})$, as far as x_{0} is chosen sufficiently small so that the above expansion is convergent at $x = x_{0}$. Therefore, once $\overrightarrow{J}(x_{0})$ is known, the asymptotic expansion of $\overrightarrow{J}(x)$ around x = 0can be achieved, and thus, $I_{\overrightarrow{\nu}}$ can be obtained.

For a given regular point $x = x_0$, $\tilde{J}(x_0)$ are quadratic integrals that can be calculated by using the AMF method [57–60]. A convenient choice can be replacing $i0^+$ by $-\eta + i0^+$ in all modified propagators, i.e., propagators which are linear in loop momenta when $x \to 0^+$. The advantage of this choice is that the number of MIs does not increase after introducing η . As $\eta \to \infty$, the newly defined integrals in the AMF method are essentially single-mass vacuum integrals [59], which can be easily calculated within the AMF framework [71]. With this boundary condition, we can obtain $\vec{J}(x_0)$ by flowing η from ∞ to $i0^-$ using η -DEs.

As numerically solving DEs is very efficient, our method is thus systematic and efficient as far as DEs can be set up.

III. A PEDAGOGICAL EXAMPLE

To elaborate how to use the above proposed recipe to calculate linear integrals, we take the simplest linear integral with one linear propagator and one quadratic propagator as an example,

$$I_{(1,1)} = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{1}{(\ell^2 + \mathrm{i}0^+)(\ell \cdot n + q \cdot n + \mathrm{i}0^+)}, \quad (11)$$

where ℓ is the loop momentum, and q and n are external momenta with $n^2 = -1$. Clearly, this integral only depends on the scale $q \cdot n$. Without loss of generality, we choose $q \cdot n = 1$. This Feynman integral is relevant to the Feynman diagram shown in Fig. 1, where the double line stands for a gauge link which contributes linear propagators. We call this a one-loop bubble integral because there is no external legs of particles.



FIG. 1. One-loop vacuum integral containing a gauge link. Double line denotes the gauge link.

The corresponding auxiliary quadratic integral is defined as

$$\widetilde{I}_{(1,1)}(x) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{1}{\widetilde{\mathcal{D}}_1 \widetilde{\mathcal{D}}_2},\tag{12}$$

² With IBP reduction, we in fact only need to calculate MIs of original linear integrals. Then, we can always choose $\tilde{I}_{\vec{\nu}}(x)$ as one of the MIs of auxiliary quadratic integrals.

³ We assume that dimensional regularization can regularize all singularities of x. Therefore, all terms where s depends on ϵ can be ignored as $x \to 0^+$. Furthermore, if s is independent of ϵ , it can be only non-negative integers, and $n_s = 0$.

with

$$\widetilde{\mathcal{D}}_1 = \ell^2 + \mathrm{i}0^+,\tag{13}$$

$$\widetilde{\mathcal{D}}_2 = x \left[\left(\ell + \frac{n}{2x} \right)^2 + \frac{1+4x}{4x^2} + \mathrm{i}0^+ \right].$$
(14)

For finite x, there are two MIs which can be defined as

$$\widetilde{J}_1(x) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{x}{\widetilde{\mathcal{D}}_2},\tag{15}$$

$$\widetilde{J}_2(x) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{x}{\widetilde{\mathcal{D}}_1 \widetilde{\mathcal{D}}_2},\tag{16}$$

with $\widetilde{I}_{(1,1)}(x) = \frac{\widetilde{J}_2(x)}{x}$. Using IBP, we get the system of *x*-DEs,

$$\frac{\partial}{\partial x}\overrightarrow{\widetilde{J}}(x) = \begin{pmatrix} \frac{2(-1+\epsilon)(1+2x)}{x(1+4x)} & 0\\ -\frac{4(-1+\epsilon)x}{1+4x} & \frac{1}{x} \end{pmatrix} \overrightarrow{\widetilde{J}}(x).$$
(17)

Besides $x = \infty$, there are two additional singularities x =0 and $x = -\frac{1}{4}$, which correspond to an infinity mass term and zero mass term in $\widetilde{\mathcal{D}}_2$, respectively. Therefore, we can choose boundary conditions at, e.g., $x_0 = \frac{1}{16}$, which point can be well estimated by asymptotic expansions around x = 0. Based on DEs (17), we get the following asymptotic expansions:

$$\widetilde{J}_1(x) = x^{-2+2\epsilon} \sum_{i=0}^{\infty} a_i x^i, \qquad (18)$$

$$\widetilde{J}_2(x) = c_0 x + x^{2\epsilon} \sum_{i=0}^{\infty} b_i x^i, \qquad (19)$$

which gives

$$\widetilde{I}_{(1,1)}(x) = \frac{\widetilde{J}_2(x)}{x} = c_0 + x^{2\epsilon - 1} \sum_{i=0}^{\infty} b_i x^i.$$
 (20)

As $x \to 0^+$, we find $\tilde{J}_1(0^+) = 0$ and $\tilde{I}_{(1,1)}(0^+) = c_0$. The DEs (17) also provide recurrence relations of coefficients,

$$a_i = \frac{8 - 4\epsilon - 4i}{i}a_{i-1},$$
(21)

$$b_{i} = -\frac{4(-1+\epsilon)}{-1+2\epsilon+i}a_{i} - \frac{4(-2+2\epsilon+i)}{-1+2\epsilon+i}b_{i-1}, \qquad (22)$$

$$b_0 = -\frac{4(-1+\epsilon)}{-1+2\epsilon}a_0,$$
(23)

where $i \geq 1$. Therefore, only a_0 and c_0 are free coefficients.

To fix a_0 and c_0 , we need to know $\vec{J}(\frac{1}{16})$. Using the package AMFlow [60], we get

$$- (338.0371701002434 - 995.0661217702465i)\epsilon - (311.267338694263 + 1888.809135035203i)\epsilon^{2} + (1510.395112926408 + 2295.761140388504i)\epsilon^{3} - (2656.967479760546 + 2012.921788373378i)\epsilon^{4} + (3337.360161271413 + 1359.608268441597i)\epsilon^{5} - (3544.782677804504 + 737.597502301953i)\epsilon^{6} + (3503.444773310217 + 330.827170478017i)\epsilon^{7} - (3393.220492080171 + 125.304259848717i)\epsilon^{8}, (24) $\tilde{J}_{2}(\frac{1}{16}) = 1.0000000000000\epsilon^{-1} - (3.361601777683940 - 3.141592653589793i) + (2.540630265434342 - 10.560783449066254i)\epsilon + (4.39414268443412 + 18.31705093747636i)\epsilon^{2} - (15.75459001722795 + 20.93897855970841i)\epsilon^{3} + (25.55137360766245 + 17.56661536372871i)\epsilon^{4} - (31.12100936594129 + 11.47483328701037i)\epsilon^{5} + (32.43597076695672 + 6.06440829051716i)\epsilon^{6} - (32.01028557435537 + 2.66344640964930i)\epsilon^{7} + (30.90646894444592 + 0.99159094290134i)\epsilon^{8}, (25)$$$

where we only keep 10 orders in ϵ expansion with 16-digit precision, although higher order and more digits can be obtained easily. We keep the same accuracy in all examples in this paper. Calculation of $\widetilde{J}(\frac{1}{16})$ in this example is certainly a very simple problem, but for general complicated problems we can still use the AMF method to calculate them. By matching the asymptotic expansions of $\vec{J}(x)$ at x = 0 with $\vec{J}(\frac{1}{16})$, we can determine a_0 and c_0 . Eventually we obtain

- $+(0.3248443395722 453.6966444901993i)\epsilon^{6}$
- $-(2.333380642220398+0.034360215639063i)\epsilon^{7}$
- $+ (0.331273935869 1814.969761821040i)\epsilon^8$. (26)

VACUUM INTEGRALS CONTAINING A IV. GAUGE LINK UP TO FOUR LOOPS

Now, we apply our recipe to calculate vacuum integrals containing a gauge link up to four loops. These results are useful for studying parton distribution functions [72–

76]. On the one hand they can serve as boundary conditions for four-loop parton correlation functions calculation, and on the other hand, they are needed to renormalize three-loop parton correlation functions [79, 80]. In general, for an *L*-loop vacuum integral with a gauge link, there must be at least *L* quadratic propagators and at most *L* linear propagators, or else, it is either reducible or scaleless. With this restriction, up to four loops we find that there are 64 nonzero and irreducible sectors in total which give 65 MIs. The 64 nonzero and irreducible sectors are displayed in the the supplementary material together with results of all the 65 MIs.



FIG. 2. The most complex integral family for 4-loop vacuum integrals containing a gauge link. Double lines denote the gauge link.

We take the most complicated integral family shown in Fig. 2 as an example to explain our calculation details. All other families can be calculated similarly and final results are provided in the supplementary material.

A general integral in this family can be expressed as

$$I_{\vec{\nu}} = \left(\prod_{j=1}^{4} \int \frac{\mathrm{d}^{D} \ell_{j}}{\mathrm{i}\pi^{D/2}}\right) \frac{\mathcal{D}_{12}^{-\nu_{12}} \mathcal{D}_{13}^{-\nu_{13}} \mathcal{D}_{14}^{-\nu_{14}}}{\mathcal{D}_{1}^{\nu_{1}} \mathcal{D}_{2}^{\nu_{2}} \cdots \mathcal{D}_{11}^{\nu_{11}}}, \qquad (27)$$

where $\mathcal{D}_1, \mathcal{D}_2, \cdots, \mathcal{D}_8$ are quadratic denominators, $\mathcal{D}_9, \mathcal{D}_{10}, \mathcal{D}_{11}$ linear denominators, and $\mathcal{D}_{12}, \mathcal{D}_{13}, \mathcal{D}_{14}$ irreducible numerators, with explicit form

$$\begin{aligned} \mathcal{D}_1 &= \ell_1^2, \, \mathcal{D}_2 = \ell_2^2, \, \mathcal{D}_3 = \ell_3^2, \, \mathcal{D}_4 = \ell_4^2, \, \mathcal{D}_5 = (\ell_2 + \ell_4)^2, \\ \mathcal{D}_6 &= (\ell_1 + \ell_2 + \ell_3)^2, \, \mathcal{D}_7 = (\ell_1 + \ell_2 + \ell_4)^2, \\ \mathcal{D}_8 &= (\ell_1 + \ell_2 + \ell_3 + \ell_4)^2, \, \mathcal{D}_9 = (\ell_1 + q) \cdot n, \\ \mathcal{D}_{10} &= (\ell_1 + \ell_2 + q) \cdot n, \, \mathcal{D}_{11} = (\ell_1 + \ell_2 + \ell_3 + q) \cdot n, \\ \mathcal{D}_{12} &= (\ell_1 + \ell_4)^2, \, \mathcal{D}_{13} = (\ell_2 + \ell_3)^2, \, \mathcal{D}_{14} = \ell_4 \cdot n, \end{aligned}$$

where Feynman prescription $+i0^+$ is omitted. This family has 32 MIs in total. Similar to the one-loop case, the single scale $q \cdot n$ can be extracted as an overall factor, and thus we can simply set $q \cdot n = 1$.

Based on our recipe, we define a corresponding auxiliary family of integrals,

$$\widetilde{I}_{\vec{\nu}}(x) = \left(\prod_{j=1}^{4} \int \frac{\mathrm{d}^{D}\ell_{j}}{\mathrm{i}\pi^{D/2}}\right) \frac{\widetilde{\mathcal{D}}_{12}^{-\nu_{12}}\widetilde{\mathcal{D}}_{13}^{-\nu_{13}}\widetilde{\mathcal{D}}_{14}^{-\nu_{14}}}{\widetilde{\mathcal{D}}_{1}^{\nu_{1}}\widetilde{\mathcal{D}}_{2}^{\nu_{2}}\cdots\widetilde{\mathcal{D}}_{11}^{\nu_{11}}},\qquad(28)$$

where

$$\widetilde{D}_9 = x \left[(\ell_1 + \frac{n}{2x})^2 + \frac{1+4x}{4x^2} + \mathrm{i}0^+ \right],$$

$$\widetilde{\mathcal{D}}_{10} = x \left[(\ell_1 + \ell_2 + \frac{n}{2x})^2 + \frac{1+4x}{4x^2} + \mathrm{i}0^+ \right], \\ \widetilde{\mathcal{D}}_{11} = x \left[(\ell_1 + \ell_2 + \ell_3 + \frac{n}{2x})^2 + \frac{1+4x}{4x^2} + \mathrm{i}0^+ \right],$$

and $\widetilde{\mathcal{D}}_i = \mathcal{D}_i$ for $i \neq 9, 10, 11$. After changing linear propagators to quadratic propagators, the number of MIs in the auxiliary family increases to 120. We use $\overrightarrow{J}(x)$ to denote these MIs. Each $\widetilde{J}_i(x)$ is defined by a specific list of exponents $\overrightarrow{\nu}^i$ via the following relation

$$\widetilde{J}_{i}(x) = x^{\nu_{9}^{i} + \nu_{10}^{i} + \nu_{11}^{i}} \widetilde{I}_{\vec{\nu}^{i}}(x), \qquad (29)$$

where, for convenience, the factor $x^{\nu_9^i+\nu_{10}^i+\nu_{11}^i}$ is introduced to cancel the overall factor x in $\widetilde{\mathcal{D}}_9, \widetilde{\mathcal{D}}_{10}, \widetilde{\mathcal{D}}_{11}$.

For a regular point, say $x_0 = \frac{1}{16}$, we calculate the corresponding MIs $\vec{J}(\frac{1}{16})$ using the package AMFlow[60] with the mass mode at the first step of the iteration. The calculation is automatic and can be decomposed to the following steps. First, $i0^+$ in $\tilde{\mathcal{D}}_{9/10/11}$ is replaced by $-\eta + i0^+$. Then, η -DEs are set up, and boundary conditions at $\eta \to \infty$ are calculated [59, 71], where Kira[18] is employed to do IBP reduction. Finally, the η -DEs are solved to realize the flow of η from ∞ to $i0^-$.

In this problem, x-DEs of $\widetilde{J}(x)$ can be derived from η -DEs during calculating $\overrightarrow{J}(\frac{1}{16})$, details of which are given in the Appendix. With boundary conditions at $x_0 = \frac{1}{16}$, the x-DEs are solved (similar to solve η -DEs) by using AMFlow[60] to obtain results at $x = 0^+$, which gives final results of our desired linear integrals. Following is the result of the top-sector corner integral,

$$\begin{split} I_{(1,\cdots,1,0,0,0)} &= -5.184638775716850\epsilon^{-1} \\ &- (35.55435225946754 + 65.15209235723534i) \\ &+ (124.0703029826581 - 446.7891674459475i)\epsilon \\ &+ (1306.855351606906 - 1870.355270319554i)\epsilon^2 \\ &+ (8989.362598245622 - 7095.610423728019i)\epsilon^3 \\ &+ (43885.08629764386 - 21591.89236479742i)\epsilon^4 \\ &+ (200504.4704938221 - 69609.6949015959i)\epsilon^5 \\ &+ (852097.4905856483 - 196376.9845468376i)\epsilon^6 \\ &+ (3560145.246623535 - 611937.690718618i)\epsilon^7 \\ &+ (14591453.36136039 - 1726015.80403403i)\epsilon^8. \quad (30) \end{split}$$

Note that higher precision and higher orders in ϵ can also be achieved easily.

V. INTEGRALS WITH EXTERNAL LEGS

Now let us consider more general linear integrals containing both gauge links and external legs. Take fourloop integrals containing one external leg shown in Fig. 3 as an example. The top-sector corner integral can be expressed as

$$I_{(1,1,1,1,1)}(q \cdot n, p \cdot n) = \left(\prod_{j=1}^{4} \int \frac{\mathrm{d}^{D} \ell_{j}}{\mathrm{i}\pi^{D/2}}\right) \frac{1}{\mathcal{D}_{1} \mathcal{D}_{2} \mathcal{D}_{3} \mathcal{D}_{4} \mathcal{D}_{5} \mathcal{D}_{6}}$$
(31)

with

$$\mathcal{D}_{1} = \ell_{1}^{2}, \ \mathcal{D}_{2} = \ell_{2}^{2}, \ \mathcal{D}_{3} = \ell_{3}^{2}, \ \mathcal{D}_{4} = \ell_{4}^{2}, \ \mathcal{D}_{5} = (\ell_{1} + q) \cdot n, \mathcal{D}_{6} = (\ell_{1} + \ell_{2} + \ell_{3} + \ell_{4} + q - p) \cdot n,$$
(32)

where p is a lightlike external momentum, and Feynman prescription $+i0^+$ for each inverse propagator is omitted.



FIG. 3. Four-loop Feynman diagram with a gauge link and an external leg.

To calculate integrals in this example, we define auxiliary quadratic integrals by using the following inverse propagators

$$\begin{aligned} \widetilde{\mathcal{D}}_5 &= x \left[(\ell_1 + \frac{n}{2x})^2 + \frac{1}{4x^2} + \frac{q \cdot n}{x} + \mathrm{i}0^+ \right] \\ \widetilde{\mathcal{D}}_6 &= x \left[(\ell_1 + \ell_2 + \ell_3 + \ell_4 + \frac{n}{2x})^2 \\ &+ \frac{1}{4x^2} + \frac{(q - p) \cdot n}{x} + \mathrm{i}0^+ \right], \end{aligned}$$

and $\mathcal{D}_i = \mathcal{D}_i$ for $i \leq 4$. After changing linear propagators to quadratic propagators, the number of MIs increases from 2 to 6. Using AMFlow [60], we calculate auxiliary quadratic integrals at a fixed point $x = x_0$, and then solve *x*-DEs to obtain integrals at $x = 0^+$. Without loss of generality, for the phase-space point $q \cdot n = 1$, $p \cdot n = \frac{1}{2}$, we obtain

$$\begin{split} I_{(1,1,1,1,1)}(1,\frac{1}{2}) &= -0.001388888888888888888^{-2} \\ &- (0.01673049985543654 + 0.01745329251994330i)\epsilon^{-1} \\ &- (0.2181793358910845 + 0.2102416617468981i) \\ &- (1.816593593006501 + 3.660426889433229i)\epsilon \\ &- (15.81900673709547 + 33.89466583909418i)\epsilon^2 \\ &- (110.3704374791621 + 401.1363831252734i)\epsilon^3 \\ &- (810.083299387532 + 3287.604829797105i)\epsilon^4 \\ &- (5155.23653311233 + 33468.66058680208i)\epsilon^5 \\ &- (34800.6654379141 + 258370.0763552148i)\epsilon^6 \\ &- (211074.015357447 + 2454130.498362980i)\epsilon^7. \end{split}$$

When there are more gauge links or more external legs, the calculation process is similar.

VI. SELF-CONSISTENCY CHECK BY DIMENSIONAL RECURRENCE RELATIONS

For any integral $I_{\vec{\nu}}^D$, where we explicitly write the dependence of spacetime dimension D, there is a parametric representation (see, e.g., [81]),

$$I_{\nu}^{D} = \left(\prod_{j=1}^{L} \int \frac{\mathrm{d}^{D}\ell_{j}}{\mathrm{i}\pi^{D/2}}\right) \prod_{i=1}^{N} \mathcal{D}_{i}^{-\nu_{i}}$$
$$= \left(\prod_{i=1}^{N} \int_{0}^{\infty} \frac{x_{i}^{\nu_{i}-1}\mathrm{d}x_{i}}{\Gamma(\nu_{i})}\right) \frac{e^{-F/U}}{U^{D/2}}, \qquad (34)$$

and thus,

$$I_{\vec{\nu}}^{D-2} = \left(\prod_{i=1}^{N} \int_{0}^{\infty} \frac{x_{i}^{\nu_{i}-1} \mathrm{d}x_{i}}{\Gamma(\nu_{i})}\right) \frac{e^{-F/U}}{U^{D/2}} U.$$
(35)

As U is a homogeneous polynomial of x_i , the above expression can be translated into a linear combination of integrals at dimension D with different $\vec{\nu}$, which can be reduced to MIs by using IBP identities. Supposing \vec{J}^D is a set of MIs, the above procedure results in dimensional recurrence relations [77, 78],

$$\vec{J}^{D-2} = C^D \vec{J}^D, \tag{36}$$

where C^D is a matrix rationally depending on D.

Up to three loops, vacuum integrals containing a gauge link have been calculated in Ref. [80] by solving dimensional recurrence relations using the method in Ref. [36]. The obtained results agree with ours. For four-loop integrals containing a gauge link, we find it very hard to solve the corresponding dimensional recurrence relations. However, the dimensional recurrence relations can provide a highly nontrivial self-consistency check for our results. We calculate MIs at two different space-time dimensions D_0 and $D_0 - 2$, and we find that the obtained results satisfy the relations Eq. (36) within precision.

VII. SUMMARY

In summary, we develop a recipe to calculate linear integrals using the AMF method [57–59]. For any given linear integral, our recipe is to introduce an auxiliary quadratic term for each linear propagator, and the obtained auxiliary quadratic integral can be calculated systematically using the AMF method. Taking the result of the auxiliary quadratic integral calculated at fixed auxiliary quadratic terms as the boundary condition and using differential equations to push the auxiliary quadratic terms to zero, effects of auxiliary quadratic terms will die out eventually, and we get the result of the target linear integral. This recipe of calculating linear integrals is very systematic and has been implemented in the package AMFlow [60].

As linear integrals show up frequently in region expansion and in effective field theories, our recipe will be useful in phenomenological studies. As the first application, we have calculated all MIs of vacuum integrals containing a gauge link up to four loops, which are useful to study parton distribution functions. Our results have been checked by nontrivial dimensional recurrence relations.

VIII. ACKNOWLEDGMENTS

We thank K.T. Chao, X. Liu, Z.Y. Li, X. Li, X. Guan, W.H. Wu, C. Meng, and R.H. Wu for many useful communications and discussions. The work is supported in part by the National Natural Science Foundation of China (Grants No. 11875071, No. 11975029), the National Key Research and Development Program of China under Contract No. 2020YFA0406400, and the Highperformance Computing Platform of Peking University.

Appendix A: Derivation of relation between x-DEs and η -DEs for vacuum integrals

For vacuum integrals containing a gauge link, x-DEs can be obtained from η -DEs. We take the family in Fig. 1 as an example to explain this, and the final relation is independent of the specific family. MIs after introducing x are defined by $\overrightarrow{\widetilde{J}}(x) = \left(\widetilde{J}_1(x), \widetilde{J}_2(x)\right)^T$, with

$$\widetilde{J}_1(x) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i} \pi^{D/2}} \frac{1}{(\ell + \frac{n}{2x})^2 + \frac{1+4x}{4x^2}},\tag{A1}$$

$$\widetilde{J}_2(x) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{1}{\ell^2 \left((\ell + \frac{n}{2x})^2 + \frac{1+4x}{4x^2} \right)},\tag{A2}$$

where Feynman prescription $+i0^+$ in each denominator is omitted. The corresponding *x*-DEs are denoted as

$$\frac{\partial}{\partial x}\overrightarrow{\widetilde{J}}(x) = A(x)\overrightarrow{\widetilde{J}}(x). \tag{A3}$$

When calculating $\overrightarrow{J}(x_0)$ using AMF method, we define modified integrals $\overrightarrow{J}(\eta) = \left(\widehat{J}_1(\eta), \widehat{J}_2(\eta)\right)^T$, with

$$\widehat{J}_1(\eta) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{1}{(\ell + \frac{n}{2x_0})^2 + \frac{1+4x_0}{4x_0^2} - \eta},\tag{A4}$$

$$\widehat{J}_{2}(\eta) = \int \frac{\mathrm{d}^{D}\ell}{\mathrm{i}\pi^{D/2}} \frac{1}{\ell^{2} \left((\ell + \frac{n}{2x_{0}})^{2} + \frac{1+4x_{0}}{4x_{0}^{2}} - \eta \right)}.$$
 (A5)

The corresponding η -DEs are denoted as

$$\frac{\partial}{\partial \eta} \overrightarrow{\widehat{J}}(\eta) = M(\eta) \overrightarrow{\widehat{J}}(\eta). \tag{A6}$$

We express A(x) by $M(\eta)$.

Let us define $\overrightarrow{K}(p^2, m^2) = \left(K_1(p^2, m^2), K_2(p^2, m^2)\right)^T$ with

$$K_1(p^2, m^2) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{1}{(\ell+p)^2 + m^2},\tag{A7}$$

$$K_2(p^2, m^2) = \int \frac{\mathrm{d}^D \ell}{\mathrm{i}\pi^{D/2}} \frac{1}{\ell^2 \left((\ell+p)^2 + m^2\right)}.$$
 (A8)

Because Feynman integrals have a definite mass dimension, there is a rescaling relation,

$$\vec{K}((1+\delta)p^2, (1+\delta)m^2) = (1+\delta)^{\alpha}\vec{K}(p^2, m^2),$$
 (A9)

where α is a diagonal matrix with diagonal elements denoting mass dimension. In this example, it is given by

$$\alpha = \begin{pmatrix} \frac{D}{2} - 1 & 0\\ 0 & \frac{D}{2} - 2 \end{pmatrix}.$$
 (A10)

Expansion of Eq. (A9) to $O(\delta)$ gives

$$\left(p^2 \frac{\partial}{\partial p^2} + m^2 \frac{\partial}{\partial m^2} - \alpha\right) \overrightarrow{K}(p^2, m^2) = 0.$$
 (A11)

For convenience, we denote

$$y = -\frac{1}{4x^2}, \ z = \frac{1+4x}{4x^2},$$
$$y_0 = -\frac{1}{4x_0^2}, \ z_0 = \frac{1+4x_0}{4x_0^2}.$$

Then, integrals are related by

$$\overrightarrow{\widetilde{J}}(x) = \overrightarrow{K}(y, z), \qquad (A12)$$

$$\vec{\widetilde{J}}(\eta) = \vec{K}(y_0, z_0 - \eta).$$
(A13)

Based on η -DEs (A6), we get

$$\frac{\partial}{\partial \eta} \overrightarrow{K}(y_0, z_0 - \eta) = M(\eta) \overrightarrow{K}(y_0, z_0 - \eta).$$
(A14)

By choosing $\eta = z_0 - z$ in the above equation, we get

$$\frac{\partial}{\partial z}\overrightarrow{K}(y_0,z) = -M(z_0-z)\overrightarrow{K}(y_0,z), \qquad (A15)$$

and thus,

$$\begin{aligned} &\frac{\partial}{\partial z} \overrightarrow{K}(y,z) \\ &= &\frac{\partial}{\partial z} \left[\left(\frac{y}{y_0} \right)^{\alpha} \overrightarrow{K}(y_0, \frac{y_0}{y}z) \right] \\ &= &- \left(\frac{y}{y_0} \right)^{\alpha - 1} M(z_0 - \frac{y_0}{y}z) \overrightarrow{K}(y_0, \frac{y_0}{y}z) \end{aligned}$$

$$= -\left(\frac{y}{y_0}\right)^{\alpha-1} M(z_0 - \frac{y_0}{y}z) \left(\frac{y}{y_0}\right)^{-\alpha} \overrightarrow{K}(y, z). \quad (A16)$$

where Eq. (A9) has been used. Therefore, we have

$$\begin{aligned} \frac{\partial}{\partial x} \overrightarrow{\widetilde{J}}(x) &= \left(\frac{\partial y}{\partial x} \frac{\partial}{\partial y} + \frac{\partial z}{\partial x} \frac{\partial}{\partial z}\right) \overrightarrow{K}(y, z) \\ &= \left[\frac{\partial y}{\partial x} \left(\frac{\alpha}{y} - \frac{z}{y} \frac{\partial}{\partial z}\right) + \frac{\partial z}{\partial x} \frac{\partial}{\partial z}\right] \overrightarrow{K}(y, z) \\ &= \frac{\partial y}{\partial x} \frac{\alpha}{y} \overrightarrow{\widetilde{J}}(x) - \left(\frac{y}{y_0}\right)^{\alpha - 1} M(z_0 - \frac{y_0}{y} z) \left(\frac{y}{y_0}\right)^{-\alpha} \end{aligned}$$

- K. G. Chetyrkin and F. V. Tkachov, Integration by Parts: The Algorithm to Calculate beta Functions in 4 Loops, Nucl. Phys. B192 (1981) 159–204 [InSPIRE].
- S. Laporta, High precision calculation of multiloop Feynman integrals by difference equations, Int. J. Mod. Phys. A 15 (2000) 5087–5159
 [hep-ph/0102033] [InSPIRE].
- J. Gluza, K. Kajda, and D. A. Kosower, Towards a Basis for Planar Two-Loop Integrals, Phys. Rev. D83 (2011) 045012 [arXiv:1009.0472] [InSPIRE].
- [4] R. M. Schabinger, A New Algorithm For The Generation Of Unitarity-Compatible Integration By Parts Relations, JHEP 01 (2012) 077
 [arXiv:1111.4220] [InSPIRE].
- [5] A. von Manteuffel and C. Studerus, *Reduze 2 Distributed Feynman Integral Reduction*, [arXiv:1201.4330] [InSPIRE].
- [6] R. N. Lee, LiteRed 1.4: a powerful tool for reduction of multiloop integrals,
 J. Phys. Conf. Ser. 523 (2014) 012059
 [arXiv:1310.1145] [InSPIRE].
- [7] A. von Manteuffel and R. M. Schabinger, A novel approach to integration by parts reduction, Phys. Lett. B744 (2015) 101-104 [arXiv:1406.4513]
 [InSPIRE].
- [8] K. J. Larsen and Y. Zhang, Integration-by-parts reductions from unitarity cuts and algebraic geometry, Phys. Rev. D93 (2016) 041701 [arXiv:1511.01071] [InSPIRE].
- [9] T. Peraro, Scattering amplitudes over finite fields and multivariate functional reconstruction,
- JHEP 12 (2016) 030 [arXiv:1608.01902] [InSPIRE].
 [10] P. Mastrolia and S. Mizera, Feynman Integrals and Intersection Theory, JHEP 02 (2019) 139
 [arXiv:1810.03818] [InSPIRE].
- [11] X. Liu and Y.-Q. Ma, Determining arbitrary Feynman integrals by vacuum integrals, Phys. Rev. D 99 (2019) 071501 [arXiv:1801.10523]
 [InSPIRE].
- [12] X. Guan, X. Liu, and Y.-Q. Ma, Complete reduction of integrals in two-loop five-light-parton scattering amplitudes, Chin. Phys. C 44 (2020) 093106 [arXiv:1912.09294] [InSPIRE].

$$\times \left[-\frac{z}{y} \frac{\partial y}{\partial x} + \frac{\partial z}{\partial x} \right] \overrightarrow{J}(x), \tag{A17}$$

where the Eq. (A11) has been used. The above result means

$$A(x) = \frac{1}{y} \frac{\partial y}{\partial x} \alpha + \frac{z^2}{y} \left(\frac{\partial}{\partial x} \frac{y}{z} \right) \\ \times \left(\frac{y}{y_0} \right)^{\alpha - 1} M \left(z_0 - y_0 \frac{z}{y} \right) \left(\frac{y}{y_0} \right)^{-\alpha}, \quad (A18)$$

which connects x-DEs to η -DEs for any vacuum integrals containing a gauge link. It is useful because M can be more easily obtained.

- J. Klappert and F. Lange, Reconstructing rational functions with FireFly, Comput. Phys. Commun. 247 (2020) 106951
 [arXiv:1904.00009] [InSPIRE].
- T. Peraro, FiniteFlow: multivariate functional reconstruction using finite fields and dataflow graphs, JHEP 07 (2019) 031 [arXiv:1905.08019] [InSPIRE].
- [15] H. Frellesvig, F. Gasparotto, S. Laporta, M. K. Mandal, P. Mastrolia, L. Mattiazzi, and S. Mizera, Decomposition of Feynman Integrals on the Maximal Cut by Intersection Numbers, JHEP 05 (2019) 153 [arXiv:1901.11510] [InSPIRE].
- Y. Wang, Z. Li, and N. Ul Basat, Direct reduction of multiloop multiscale scattering amplitudes, Phys. Rev. D 101 (2020) 076023 [arXiv:1901.09390]
 [InSPIRE].
- [17] A. V. Smirnov and F. S. Chuharev, FIRE6: Feynman Integral REduction with Modular Arithmetic, Comput. Phys. Commun. 247 (2020) 106877 [arXiv:1901.07808] [InSPIRE].
- [18] J. Klappert, F. Lange, P. Maierhöfer, and J. Usovitsch, Integral reduction with Kira 2.0 and finite field methods, Comput. Phys. Commun. 266 (2021) 108024 [arXiv:2008.06494] [InSPIRE].
- [19] J. Boehm, M. Wittmann, Z. Wu, Y. Xu, and Y. Zhang, IBP reduction coefficients made simple, JHEP 12 (2020) 054 [arXiv:2008.13194] [InSPIRE].
- [20] N. u. Basat, Z. Li, and Y. Wang, Reduction of the planar double-box diagram for single-top production via auxiliary mass flow, Phys. Rev. D 104 (2021) 056020 [arXiv:2102.08225] [InSPIRE].
- M. Heller and A. von Manteuffel, MultivariateApart: Generalized partial fractions, Comput. Phys. Commun. 271 (2022) 108174
 [arXiv:2101.08283] [InSPIRE].
- [22] D. Bendle, J. Boehm, M. Heymann, R. Ma, M. Rahn, L. Ristau, M. Wittmann, Z. Wu, and Y. Zhang, *Two-loop five-point integration-by-parts relations in a* usable form, [arXiv:2104.06866] [InSPIRE].
- [23] K. Hepp, Proof of the Bogolyubov-Parasiuk theorem on renormalization,
 - Commun. Math. Phys. 2 (1966) 301–326 [InSPIRE].
- [24] M. Roth and A. Denner, High-energy approximation of one loop Feynman integrals,

Nucl. Phys. B **479** (1996) 495–514 [hep-ph/9605420] [InSPIRE].

- [25] T. Binoth and G. Heinrich, An automatized algorithm to compute infrared divergent multiloop integrals, Nucl. Phys. B585 (2000) 741-759 [hep-ph/0004013]
 [InSPIRE].
- [26] G. Heinrich, Sector Decomposition, Int. J. Mod. Phys. A23 (2008) 1457-1486
 [arXiv:0803.4177] [InSPIRE].
- [27] A. V. Smirnov, FIESTA4: Optimized Feynman integral calculations with GPU support, Comput. Phys. Commun. 204 (2016) 189–199
 [arXiv:1511.03614] [InSPIRE].
- [28] S. Borowka, G. Heinrich, S. P. Jones, M. Kerner, J. Schlenk, and T. Zirke, SecDec-3.0: numerical evaluation of multi-scale integrals beyond one loop, Comput. Phys. Commun. 196 (2015) 470–491 [arXiv:1502.06595] [InSPIRE].
- [29] S. Borowka, G. Heinrich, S. Jahn, S. P. Jones, M. Kerner, J. Schlenk, and T. Zirke, pySecDec: a toolbox for the numerical evaluation of multi-scale integrals, Comput. Phys. Commun. 222 (2018) 313–326 [arXiv:1703.09692] [InSPIRE].
- [30] E. E. Boos and A. I. Davydychev, A Method of evaluating massive Feynman integrals, Theor. Math. Phys. 89 (1991) 1052–1063 [InSPIRE].
 [Teor. Mat. Fiz.89,56(1991)].
- [31] V. A. Smirnov, Analytical result for dimensionally regularized massless on shell double box, Phys. Lett. B460 (1999) 397-404 [hep-ph/9905323] [InSPIRE].
- [32] J. B. Tausk, Nonplanar massless two loop Feynman diagrams with four on-shell legs, Phys. Lett. B 469 (1999) 225-234 [hep-ph/9909506]
 [InSPIRE].
- [33] M. Czakon, Automatized analytic continuation of Mellin-Barnes integrals, Comput. Phys. Commun. 175 (2006) 559–571
 [hep-ph/0511200] [InSPIRE].
- [34] A. V. Smirnov and V. A. Smirnov, On the Resolution of Singularities of Multiple Mellin-Barnes Integrals, Eur. Phys. J. C 62 (2009) 445-449 [arXiv:0901.0386] [InSPIRE].
- [35] J. Gluza, K. Kajda, and T. Riemann, AMBRE: A Mathematica package for the construction of Mellin-Barnes representations for Feynman integrals, Comput. Phys. Commun. 177 (2007) 879–893 [arXiv:0704.2423] [InSPIRE].
- [36] R. N. Lee, Space-time dimensionality D as complex variable: Calculating loop integrals using dimensional recurrence relation and analytical properties with respect to D, Nucl. Phys. B830 (2010) 474–492 [arXiv:0911.0252] [InSPIRE].
- [37] A. V. Kotikov, Differential equations method: New technique for massive Feynman diagrams calculation, Phys. Lett. B254 (1991) 158–164 [InSPIRE].
- [38] A. V. Kotikov, Differential equation method: The Calculation of N point Feynman diagrams, Phys. Lett. B 267 (1991) 123-127 [InSPIRE]. [Erratum: Phys.Lett.B 295, 409-409 (1992)].
- [39] E. Remiddi, Differential equations for Feynman graph amplitudes, Nuovo Cim. A110 (1997) 1435-1452
 [hep-th/9711188] [InSPIRE].
- [40] T. Gehrmann and E. Remiddi, Differential equations for

two loop four point functions, Nucl. Phys. **B580** (2000) 485–518 [hep-ph/9912329] [InSPIRE].

- M. Argeri and P. Mastrolia, Feynman Diagrams and Differential Equations, Int. J. Mod. Phys. A 22 (2007) 4375-4436
 [arXiv:0707.4037] [InSPIRE].
- [42] S. Müller-Stach, S. Weinzierl, and R. Zayadeh, Picard-Fuchs equations for Feynman integrals, Commun. Math. Phys. 326 (2014) 237-249 [arXiv:1212.4389] [InSPIRE].
- [43] J. M. Henn, Multiloop integrals in dimensional regularization made simple, Phys. Rev. Lett. 110 (2013) 251601 [arXiv:1304.1806] [InSPIRE].
- [44] J. M. Henn, Lectures on differential equations for Feynman integrals, J. Phys. A48 (2015) 153001 [arXiv:1412.2296] [InSPIRE].
- [45] F. Moriello, Generalised power series expansions for the elliptic planar families of Higgs + jet production at two loops, JHEP 01 (2020) 150 [arXiv:1907.13234]
 [InSPIRE].
- [46] M. Hidding, DiffExp, a Mathematica package for computing Feynman integrals in terms of one-dimensional series expansions, Comput. Phys. Commun. 269 (2021) 108125
 [arXiv:2006.05510] [InSPIRE].
- [47] S. Catani, T. Gleisberg, F. Krauss, G. Rodrigo, and J.-C. Winter, From loops to trees by-passing Feynman's theorem, JHEP 09 (2008) 065 [arXiv:0804.3170] [InSPIRE].
- [48] G. Rodrigo, S. Catani, T. Gleisberg, F. Krauss, and J.-C. Winter, From multileg loops to trees (by-passing Feynman's Tree Theorem), Nucl. Phys. B Proc. Suppl. 183 (2008) 262-267 [arXiv:0807.0531] [InSPIRE].
- [49] I. Bierenbaum, S. Catani, P. Draggiotis, and G. Rodrigo, A Tree-Loop Duality Relation at Two Loops and Beyond, JHEP 10 (2010) 073 [arXiv:1007.0194] [InSPIRE].
- [50] I. Bierenbaum, S. Buchta, P. Draggiotis, I. Malamos, and G. Rodrigo, *Tree-Loop Duality Relation beyond* simple poles, JHEP 03 (2013) 025 [arXiv:1211.5048] [InSPIRE].
- [51] E. T. Tomboulis, Causality and Unitarity via the Tree-Loop Duality Relation, JHEP 05 (2017) 148 [arXiv:1701.07052] [InSPIRE].
- [52] R. Runkel, Z. Szőr, J. P. Vesga, and S. Weinzierl, *Causality and loop-tree duality at higher loops*, *Phys. Rev. Lett.* **122** (2019) 111603 [arXiv:1902.02135] [InSPIRE]. [Erratum: Phys.Rev.Lett. 123, 059902 (2019)].
- [53] Z. Capatti, V. Hirschi, D. Kermanschah, and B. Ruijl, Loop-Tree Duality for Multiloop Numerical Integration, Phys. Rev. Lett. **123** (2019) 151602 [arXiv:1906.06138] [InSPIRE].
- [54] J. J. Aguilera-Verdugo, F. Driencourt-Mangin, R. J. Hernández-Pinto, J. Plenter, S. Ramirez-Uribe, A. E. Renteria Olivo, G. Rodrigo, G. F. R. Sborlini, W. J. Torres Bobadilla, and S. Tracz, Open Loop Amplitudes and Causality to All Orders and Powers from the Loop-Tree Duality, Phys. Rev. Lett. **124** (2020) 211602 [arXiv:2001.03564] [InSPIRE].
- [55] Q. Song and A. Freitas, On the evaluation of two-loop

electroweak box diagrams for $e^+e^- \rightarrow HZ$ production, JHEP 04 (2021) 179 [arXiv:2101.00308] [InSPIRE].

- [56] I. Dubovyk, A. Freitas, J. Gluza, K. Grzanka, M. Hidding, and J. Usovitsch, Evaluation of multi-loop multi-scale Feynman integrals for precision physics, [arXiv:2201.02576] [InSPIRE].
- [57] X. Liu, Y.-Q. Ma, and C.-Y. Wang, A Systematic and Efficient Method to Compute Multi-loop Master Integrals, Phys. Lett. B779 (2018) 353–357 [arXiv:1711.09572] [InSPIRE].
- [58] X. Liu, Y.-Q. Ma, W. Tao, and P. Zhang, Calculation of Feynman loop integration and phase-space integration via auxiliary mass flow, Chin. Phys. C 45 (2021) 013115 [arXiv:2009.07987] [InSPIRE].
- [59] X. Liu and Y.-Q. Ma, Multiloop corrections for collider processes using auxiliary mass flow, Phys. Rev. D 105 (2022) L051503 [arXiv:2107.01864] [InSPIRE].
- [60] X. Liu and Y.-Q. Ma, AMFlow: a Mathematica Package for Feynman integrals computation via Auxiliary Mass Flow, [arXiv:2201.11669] [InSPIRE].
- [61] P. Zhang, C.-Y. Wang, X. Liu, Y.-Q. Ma, C. Meng, and K.-T. Chao, Semi-analytical calculation of gluon fragmentation into¹S₀^[1,8] quarkonia at next-to-leading order, JHEP 04 (2019) 116 [arXiv:1810.07656] [InSPIRE].
- [62] P. Zhang, C. Meng, Y.-Q. Ma, and K.-T. Chao, Gluon fragmentation into ³P_J^[1,8] quark pair and test of NRQCD factorization at two-loop level, JHEP 08 (2021) 111 [arXiv:2011.04905] [InSPIRE].
- [63] Q. Yang, R.-Y. Zhang, M.-M. Long, S.-M. Wang, W.-G. Ma, J.-W. Zhu, and Y. Jiang, QCD Corrections to e⁺e⁻ → H[±]W[∓] in Type-I THDM at Electron Positron Colliders, Chin. Phys. C 44 (2020) 093101 [arXiv:2005.11010] [InSPIRE].
- [64] C. Brønnum-Hansen and C.-Y. Wang, Contribution of third generation quarks to two-loop helicity amplitudes for W boson pair production in gluon fusion, JHEP 01 (2021) 170 [arXiv:2009.03742] [InSPIRE].
- [65] C. Brønnum-Hansen and C.-Y. Wang, Top quark contribution to two-loop helicity amplitudes for Z boson pair production in gluon fusion, JHEP 05 (2021) 244 [arXiv:2101.12095] [InSPIRE].
- [66] R.-H. Wu, Y.-S. Zuo, C. Meng, Y.-Q. Ma, and K.-T. Chao, NLO effects for Ω QQQ baryons in QCD Sum Rules, Chin. Phys. C 45 (2021) 093103 [arXiv:2104.07384] [InSPIRE].
- [67] C. Brønnum-Hansen, K. Melnikov, J. Quarroz, and C.-Y. Wang, On non-factorisable contributions to t-channel single-top production, JHEP 11 (2021) 130 [arXiv:2108.09222] [InSPIRE].
- [68] D. Baranowski, M. Delto, K. Melnikov, and C.-Y.

Wang, On phase-space integrals with Heaviside functions, JHEP **02** (2022) 081 [arXiv:2111.13594] [InSPIRE].

- [69] M. Beneke and V. A. Smirnov, Asymptotic expansion of Feynman integrals near threshold, Nucl. Phys. B522 (1998) 321-344 [hep-ph/9711391]
 [InSPIRE].
- [70] V. A. Smirnov, Problems of the strategy of regions, Phys. Lett. B 465 (1999) 226-234 [hep-ph/9907471] [InSPIRE].
- [71] Z.-F. Liu and Y.-Q. Ma, Feynman integrals are completely determined by linear algebra, [arXiv:2201.11637] [InSPIRE].
- [72] X. Ji, Parton Physics on a Euclidean Lattice, *Phys.Rev.Lett.* **110** (2013) 262002 [arXiv:1305.1539] [InSPIRE].
- [73] X. Ji, Parton Physics from Large-Momentum Effective Field Theory, Sci. China Phys. Mech. Astron. 57 (2014) 1407–1412
 [arXiv:1404.6680] [InSPIRE].
- [74] A. Radyushkin, Quasi-parton distribution functions, momentum distributions, and pseudo-parton distribution functions, Phys. Rev. D 96 (2017) 034025
 [arXiv:1705.01488] [InSPIRE].
- [75] Y.-Q. Ma and J.-W. Qiu, Extracting Parton Distribution Functions from Lattice QCD Calculations, Phys. Rev. D 98 (2018) 074021 [arXiv:1404.6860] [InSPIRE].
- [76] Y.-Q. Ma and J.-W. Qiu, Exploring Partonic Structure of Hadrons Using ab initio Lattice QCD Calculations, Phys. Rev. Lett. 120 (2018) 022003 [arXiv:1709.03018] [InSPIRE].
- [77] O. Tarasov, Connection between Feynman integrals having different values of the space-time dimension, Phys. Rev. D54 (1996) 6479-6490 [hep-th/9606018] [InSPIRE].
- [78] O. V. Tarasov, Application and explicit solution of recurrence relations with respect to space-time dimension, Nucl. Phys. B Proc. Suppl. 89 (2000) 237-245
 [hep-ph/0102271] [InSPIRE].
- [79] V. M. Braun, A. Vladimirov, and J.-H. Zhang, Power corrections and renormalons in parton quasidistributions, Phys. Rev. D 99 (2019) 014013 [arXiv:1810.00048] [InSPIRE].
- [80] Z.-Y. Li, Y.-Q. Ma, and J.-W. Qiu, Extraction of Next-to-Next-to-Leading-Order Parton Distribution Functions from Lattice QCD Calculations, Phys. Rev. Lett. **126** (2021) 072001 [arXiv:2006.12370] [InSPIRE].
- [81] T. Bitoun, C. Bogner, R. P. Klausen, and E. Panzer, Feynman integral relations from parametric annihilators, Lett. Math. Phys. 109 (2019) 497–564 [arXiv:1712.09215] [InSPIRE].