RESEARCH ARTICLE

𝑓(*𝑇 , 𝐵*) **gravity in a Friedmann–Lemaître–Robertson–Walker universe with nonzero spatial curvature**

Andronikos Paliathanasis^{1,2} | Genly Leon^{*3,1}

1 Institute of Systems Science, Durban University of Technology, PO Box 1334, Durban 4000, South Africa 2 Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Valdivia 5090000, Chile ³Departamento de Matemáticas, Universidad

Católica del Norte, Avda. Angamos 0610, Casilla 1280 Antofagasta, Chile

Correspondence

*Genly Leon, Departamento de Matemáticas, Universidad Católica del Norte, Avda. Angamos 0610, Casilla 1280 Antofagasta, Chile & Institute of Systems Science, Durban University of Technology, PO Box 1334, Durban 4000, South Africa Email: genly.leon@ucn.cl

Abstract

We investigate exact solutions and the asymptotic dynamics for the Friedmann– Lemaître–Robertson–Walker universe with nonzero spatial curvature in the fourthorder modified teleparallel gravitational theory known as $f(T, B)$ theory. We show that the field equations admit a minisuperspace description, and they can reproduce any exact form of the scale factor. Moreover, we calculate the equilibrium points and analyze their stability. We show that Milne and Milne-like solutions are supported, and the de Sitter universe is provided. To complete our analysis, we use Poincaré variables to investigate the dynamics at infinity.

KEYWORDS:

Teleparallel cosmology; modified gravity; open universe; closed universe

1 INTRODUCTION

Cosmologists have widely studied alternative and modified theories of gravity in recent years $\frac{1,2,3}{2}$ $\frac{1,2,3}{2}$ $\frac{1,2,3}{2}$ $\frac{1,2,3}{2}$ $\frac{1,2,3}{2}$ because geometrodynamical degrees of freedom are introduced into the gravitational Action Integral to explain recent cosmological observations^{[4,](#page-15-3)[5,](#page-15-4)[6,](#page-15-5)[7](#page-15-6)[,8,](#page-15-7)[9](#page-15-8)}. In particular, geometric invariants are used to modify the Einstein-Hilbert Action. See for instance ^{[1](#page-15-0)[,2,](#page-15-1)[10](#page-15-9)[,11](#page-15-10)[,12](#page-15-11)} and references therein.

The fundamental geometric invariant function of General Relativity is the Ricciscalar *R*. The Levi-Civita connection defines *R*. However, Einstein showed that if the fundamental connection curvature-less Weitzenböck connection^{[13](#page-15-12)} and the torsion scalar *T* are used for the definition of the gravitational theory, then the resulting theory is equivalent to General Relativity known as the teleparallel equivalence of General Relativity (TEGR) $14,15$ $14,15$. There is a plethora of modified theories inspired by teleparallelism with many interesting results in cosmology and astrophysics ^{[16,](#page-16-1)[17,](#page-16-2)[18](#page-16-3)[,19,](#page-16-4)[20,](#page-16-5)[21,](#page-16-6)[22](#page-16-7)[,23,](#page-16-8)[24](#page-16-9)}. For reviews in teleparallelism, we refer the reader to references^{[25](#page-16-10)[,26](#page-16-11)}.

We are interested in the fourth-order teleparallel theory of gravity known as the $f(T, B)$ theory. The gravitational Action Integral is a defined by an arbitrary function f of the torsion scalar T and of the boundary term B , which is related to the torsion scalar and the Ricciscalar, that is $B = R + T$. The theory was introduced in detail in^{[27](#page-16-12)}. However, a similar fourth-order teleparallel theory was introduced before in^{[28](#page-16-13)}. There are various studies in the literature on $f(T, B)$ theory. The cosmological dynamics in a spatially flat Friedmann–Lemaître–Robertson–Walker (FLRW) universe were investigated in detail in a series of works ^{[29,](#page-16-14)[30](#page-16-15)[,31](#page-16-16)[,32](#page-16-17)} where it was found that for a plethora of functions *f* the modified teleparallel theory can describe the main eras of the cosmological history. Exact and analytic solutions were found in $33,34$ $33,34$, while some bouncing solutions were determined in^{[35](#page-16-20)}. Recently, a new inhomogeneous exact solution was derived in 36 36 36 . The quantization of $f(T, B)$ by using the minisuperspace description to write the Wheeler-DeWitt equation of quantum cosmology was studied in 37 . Cosmological constraints of $f(T, B)$ theory can be found in $38,39,40$ $38,39,40$ $38,39,40$ while some astrophysics applications are presented in $41,42,43,44,45,46$ $41,42,43,44,45,46$ $41,42,43,44,45,46$ $41,42,43,44,45,46$ $41,42,43,44,45,46$ $41,42,43,44,45,46$. Anisotropic spacetimes in $f(T, B)$ theory investigated before in^{[47](#page-17-1)[,48](#page-17-2)[,49](#page-17-3)}. Specifically, the dynamical evolution for the physical parameters investigated in the case of Bianchi I^{47} I^{47} I^{47} , Kantowski-Sachs^{[48](#page-17-2)}, and Bianchi III^{[49](#page-17-3)} background geometries.

In the following, we consider in $f(T, B)$ theory in the background space of an FLRW universe with nonzero spatial curvature. It has been found that the inflationary scenario is not affected by the presence of negative curvature in the background space $50,51$ $50,51$. Thus, such an analysis is important for studying teleparallelism in the very early stages of the universe. We investigate the existence of power-law and exponential scale factors in the $f(T, B)$ theory, which can describe inflation. An accelerated expansion of the universe described by the exponential scale factor solves various problems in cosmology, such as the "flatness", "horizon" and monopole problems $52,53$ $52,53$. A recent study on the effects of curvature in teleparallelism was performed in 54 where Milne and Milne-like universes are supported in $f(T)$ theory. At the same time, the de Sitter expansion is provided by the $f(T)$ the-ory without the necessity to introduce a cosmological constant term. In^{[55](#page-17-9)} bouncing solutions in $f(T)$ cosmology with nonzero spatial curvature was studied.

Furthermore, we investigate the global dynamics of the field equations by performing a complete dynamical analysis^{[56](#page-17-10)}. Indeed, we determine the equilibrium points and investigate their stability properties. At every equilibrium point, the scale factor is described by an exact solution corresponding to a specific epoch of the cosmological history $57,58,59$ $57,58,59$ $57,58,59$. The analysis of the asymptotic behaviour for the theory is essential for a better understanding of the viability of the model $60,61$ $60,61$.

The plan of the paper is as follows.

In Section [2](#page-1-0) we present the basic properties of teleparallelism and we give the gravitational field equations for the $f(T, B)$ theory. Moreover, for the case of FLRW geometries, we derive the minisuperspace description for the field equations and write the point-like Lagrangian using a Lagrange multiplier. For the case of $f(T, B) = T + F(B)$ in Section [3](#page-4-0) we prove that the gravitational theory supports exact solutions of interest, specifically those with power-law and exponential scale factor. The stability properties of these important solutions are investigated in Section [4.](#page-5-0) In particular, we investigate the global evolution of the field equations by investigating the equilibrium points and their stability properties. We discuss our results in Section [5.](#page-13-0)

 $2 \mid F(T, B)$ **GRAVITY**

The fundamental geometric objects in teleparallelism are the vierbein fields $\mathbf{e}_{\mu}(x^{\sigma})$. The vierbein fields introduce the dynamical variables of the theory, and they form an orthonormal basis for the tangent space at each point *P* such that $g(e_\mu, e_\nu) = e_\mu \cdot e_\nu = \eta_{\mu\nu}$, where $\eta_{\mu\nu}$ is the line element of the Minkowski spacetime, $\eta_{\mu\nu} = \text{diag}(-, +, +, +)$.

Furthermore, for the vierbein fields, it holds that

$$
[e_{\mu}, e_{\nu}] = c_{\nu\mu}^{\beta} e_{\beta} \text{ where } c_{(\nu\mu)}^{\beta} = 0.
$$
 (1)

In general, in the nonholonomic coordinates, the covariant derivative ∇_{μ} is defined by the nonsymmetric connection

$$
\mathring{\Gamma}^{\mu}_{\nu\beta} = \{^{\mu}_{\nu\beta}\} + \frac{1}{2}g^{\mu\sigma}(c_{\nu\sigma,\beta} + c_{\sigma\beta,\nu} - c_{\mu\beta,\sigma}),\tag{2}
$$

in which $\{^{\mu}_{\nu\beta}\}$ is the symmetric Levi-Civita connection of Riemannian geometry.

When $\mathbf{e}_{\mu} \cdot \mathbf{e}_{\nu} = \eta_{\mu\nu}$, it follows that ^{[13](#page-15-12)}

$$
\mathring{\Gamma}^{\mu}_{\nu\beta} = \frac{1}{2} \eta^{\mu\sigma} (c_{\nu\sigma,\beta} + c_{\sigma\beta,\nu} - c_{\mu\beta,\sigma}), \tag{3}
$$

where now \int_{γ}^{μ} describes the Ricci rotation coefficients. \int_{γ}^{μ} has the property that it is antisymmetric in the first two indices, i.e. $\int_{\mu\nu\beta}^{\rho} = -\int_{\nu\mu\beta}^{\rho}$, with $\int_{\mu\nu\beta}^{\rho} = \eta_{\mu\sigma}\int_{\nu\beta}^{\mu}$.

Consequently, the non-null torsion tensor can be defined as $T^{\beta}_{\mu\nu} = \mathring{\Gamma}^{\beta}_{\nu\mu} - \mathring{\Gamma}^{\beta}_{\mu\nu}$, with the scalar $T = S_{\beta}^{\mu\nu}T^{\beta}_{\mu\nu}$. The geometric object $S_\beta^{\mu\nu}$ is given by the expression

$$
S_{\beta}^{\ \mu\nu} = \frac{1}{2} (K^{\mu\nu}{}_{\beta} + \delta^{\mu}_{\beta} T^{\theta\nu}{}_{\theta} - \delta^{\nu}_{\beta} T^{\theta\mu}{}_{\theta}), \tag{4}
$$

with $K^{\mu\nu}$ $\frac{1}{\beta} = -\frac{1}{2}$ $\frac{1}{2}(T^{\mu\nu}{}_{\beta} - T^{\nu\mu}{}_{\beta} - T_{\beta}^{\mu\nu}).$

In the teleparallel equivalent of General Relativity, the fundamental scalar is the torsion scalar *T*, instead of the Ricciscalar and the dynamical variables are the vierbein fields, instead of the metric tensor, that is, the gravitational Action Integral is

$$
S_T = \frac{1}{16\pi G} \int d^4x eT, \ e = \det(e_\mu). \tag{5}
$$

In this study, we are interested in an extension of the teleparallel theory known as $f(T, B)$ gravity. Specifically, we consider the modified Action Integral $27,28$ $27,28$

$$
S_{f(T,B)} = \frac{1}{16\pi G} \int d^4x e f(T,B), \qquad (6)
$$

where $B = 2e^{-1}\partial_v (eT_\rho^{\rho\nu}$) corresponds to the boundary term which relates the torsion scalar T with the Ricciscalar R , that is $R = -T + B^{27}$ $R = -T + B^{27}$ $R = -T + B^{27}$. Function *f* (*T*, *B*) is an arbitrary function which should be defined. In the case for which *f* (*T*, *B*) is a linear function. Then, the field equations of the General Relativity are recovered, with or without the cosmological constant term. Moreover, for $f(T, B) = f(-T + B)$, another well-known theory of gravity is recovered, the so-called fourth-order modified $f(R)$ theory.

The gravitational field equations follow from the variation of the Action Integral [\(6\)](#page-2-0) with respect to the vierbein fields. They are 27

$$
4\pi G e \mathcal{T}_{a}^{(m)\lambda} = \frac{1}{2} e h_{a}^{\lambda} (f_{,B})^{\mu\nu} g_{\mu\nu} - \frac{1}{2} e h_{a}^{\sigma} (f_{,B})_{;\sigma}^{\ \ ;\lambda} + \frac{1}{4} e (B f_{,B} - f) h_{a}^{\lambda} + (e S_{a}^{\ \mu\lambda})_{,\mu} f_{,T} + e (f_{,B})_{,\mu} + (f_{,T})_{,\mu} S_{a}^{\ \mu\lambda} - e f_{,T} T^{\sigma}{}_{\mu a} S_{\sigma}^{\ \lambda\mu}, \tag{7}
$$

or equivalently

$$
4\pi G e \mathcal{T}_{a}^{(m)\lambda} = e f_{,T} G_{a}^{\lambda} + \left[\frac{1}{4} \left(T f_{,T} - f \right) e h_{a}^{\lambda} + e(f_{,T})_{,\mu} S_{a}^{\mu\lambda} \right] + \left[e(f_{,B})_{,\mu} S_{a}^{\mu\lambda} - \frac{1}{2} e \left(h_{a}^{\sigma} \left(f_{,B} \right)_{,\sigma}^{,\lambda} - h_{a}^{\lambda} \left(f_{,B} \right)^{\mu\nu} g_{\mu\nu} \right) + \frac{1}{4} e B h_{a}^{\lambda} f_{,B} \right].
$$
\n(8)

2.1 Minisuperspace description

For the FLRW universe with nonzero curvature, that is, for the line element

$$
ds^{2} = N^{2}dt^{2} - a^{2}(t)\left(dr^{2} + \sin^{2}(r)\left(d\theta^{2} + \sin^{2}(\theta)d\phi^{2}\right)\right), K = 1,
$$
\n(9)

or

$$
ds^{2} = N^{2}dt^{2} - a^{2}(t)\left(dr^{2} + \sinh^{2}(r)\left(d\theta^{2} + \sin^{2}(\theta)d\phi^{2}\right)\right), K = -1,
$$
\n(10)

we consider the vierbein fields

$$
e_{\mu} = (N dt, a(t) E^{r}(K), a(t) E^{\theta}(K), a(t) E^{\phi}(K)),
$$

where the components of $\mathbf{E}(K)$ depend upon the spatial curvature K and, for simplicity, in the following, we set the lapse function $N = 1$.

For the "closed" universe, i.e. $K = 1$, we have ^{[62](#page-17-16)}

$$
E^{r}(K=1) = -\cos(\theta)dr + \sin(r)\sin(\theta)(\cos(r)d\theta - \sin(r)\sin(\theta)d\phi),
$$
\n(11)

$$
E^{\theta} (K = 1) = \sin(\theta)\cos(\phi)dr - \sin(r)(\sin(r)\sin(\phi) - \cos(r)\cos(\theta)\cos(\phi))d\theta
$$

-
$$
\sin(r)\sin(\theta)(\cos(r)\sin(\phi) + \sin(r)\cos(\theta)\cos(\phi))d\phi,
$$
 (12)

$$
E^{\phi}(K=1) = -\sin(\theta)\sin(\phi)dr - \sin(r)(\sin(r)\cos(\phi) + \cos(r)\cos(\theta)\sin(\phi))d\theta
$$

-
$$
\sin(r)\sin(\theta)(\cos(r)\cos(\phi) - \sin(r)\cos(\theta)\sin(\phi))d\phi.
$$
 (13)

For the "open" universe, i.e. $K = -1$, it holds that ^{[62](#page-17-16)}

$$
E^{r}(K = -1) = \cos(\theta)dr + \sinh(r)\sin(\theta) \left(-\cosh(r)d\theta + i\sinh(r)\sin(\theta)d\phi \right),\tag{14}
$$

$$
E^{\theta} (K = -1) = -\sin(\theta)\cos(\phi)dr + \sinh(r)(i\sinh(r)\sin(\phi) - \cos(r)\cos(\theta)\cos(\phi))d\theta
$$

+
$$
\sinh(r)\sin(\phi)(\cosh(r)\sin(\phi) + i\sinh(r)\cos(\theta)\cos(\phi))d\phi,
$$
 (15)

$$
E^{\phi}(K = -1) = \sin(\theta)\sin(\phi)d\theta + \sinh(\theta)(i\sinh(\theta)\cos(\phi) + \cosh(\theta)\cos(\theta)\sin(\phi))d\theta
$$

+
$$
\sinh(\theta)\sin(\theta)(\cosh(\theta)\cos(\phi) - \sinh(\theta)\cos(\theta)\sin(\phi))d\phi.
$$
 (16)

Thus, for this frame, the torsion scalar T is calculated as 62

$$
T = 6\left(\frac{K}{a^2} - H^2\right),\tag{17}
$$

where $H = \frac{\dot{a}}{a}$ $\frac{a}{a}$ is the Hubble function $\dot{a} = \frac{da}{dt}$ $\frac{du}{dt}$, while the boundary term is

$$
B = -6(\dot{H} + 3H^2)
$$
 (18)

We introduce the Lagrange multipliers λ_1 , λ_2 . Thus the Action Integral [\(6\)](#page-2-0) for a the FLRW spacetime is written in the equivalent form

$$
S_{f(T,B)} = \frac{1}{16\pi G} \int dt \left(a^3 f(T,B) - \lambda_1 a^3 \left(T - 6 \left(\frac{K}{a^2} - H^2 \right) \right) - \lambda_2 a^3 \left(B + 6 \left(\dot{H} + 3H^2 \right) \right) \right). \tag{19}
$$

Variation with respect to the variables *T* and *B* of [\(19\)](#page-3-0) constrain the Lagrange multiplier. Indeed, from the equations of motion δ δT $(S_{f(T,B)}) = 0$ and $\frac{\delta}{\delta B}$ $(S_{f(T,B)}) = 0$, it follows that $\lambda_1 = f_T$ and $\lambda_2 = f_B$. Consequently, expression [\(19\)](#page-3-0) becomes

$$
S_{f(T,B)} = \frac{1}{16\pi G} \int dt \left(a^3 f(T,B) - f_{,T} \left(a^3 T - 6 \left(aK - a\dot{a}^2 \right) \right) - f_{,B} \left(B + 12a\dot{a}^2 \right) - 6 f_{,B} a^2 \ddot{a} \right). \tag{20}
$$

Integration by parts of the last term of (20) gives

$$
\int dt \left(6f_B a^2 \ddot{a}\right) = -\int dt \left(12f_B a\dot{a}^2 + 6a^2 f_{,BB} \dot{a}\dot{B}\right).
$$
\n(21)

Thus, we can write the point-like Lagrangian function

$$
\mathcal{L}(a, \dot{a}, T, B, \dot{B}) = -6f_{,T}a\dot{a}^2 - 6a^2f_{,BB}\dot{a}\dot{B} + a^3(f - Tf_{,T} - Bf_{,B}) + 6aKf_{,T},
$$
\n(22)

which generates the gravitational field equations. We remark that $\mathcal{L}(a, \dot{a}, T, B, \dot{B})$ provides field equations of second-order. However, *B* has been introduced by a Lagrange multiplier and includes the higher-order derivatives, such that $f(T, B)$ theory is to be of fourth-order.

2.2 $\mid f(T, B) = T + F(B)$ theory

We proceed with our analysis by assuming the functional form $f(T, B)$ to be linear in *T*, that is, $f(T, B) = T + F(B)$. A such function has been considered before in^{[29](#page-16-14)}. The main mathematical novelty of this approach is that the point-like Lagrangian [\(22\)](#page-3-2) is regular, while for small values of function $F(B)$ we are very close to the limit of General Relativity. In such consideration, we assume a modification of the Action Integral for the TEGR, which follows from the existence of the boundary function *B*.

Now, we introduce the new field, and potential definition

$$
\phi = F_{B}
$$
, and $V(\phi) = (F - BF_{B})/6$. (23)

Given an explicit form $V(\phi)$ we reconstruct $f(B)$ from the singular solution of Clairaut's equation ^{[64](#page-17-17)}.

Therefore, the Lagrangian of the field equations is

$$
\mathcal{L}\left(a,\dot{a},\phi,\dot{\phi}\right) = \frac{1}{N}\left[-6a\dot{a}^2 - 6a^2\dot{a}\dot{\phi}\right] + N\left[6a^3V\left(\phi\right) + 6aK\right],\tag{24}
$$

whereby convenience, we have reinserted the lapse function N .

Taking the variation of [\(24\)](#page-3-3) with respect to $\{a, \phi, N\}$ we derive the field equations

$$
2\dot{H} + 3H^2 + \ddot{\phi} + 3V(\phi) + Ka^{-2} = 0,
$$
\n(25)

$$
\dot{H} + 3H^2 + V'(\phi) = 0,\t(26)
$$

with constraint equation

$$
H^{2} + H\dot{\phi} + V(\phi) + Ka^{-2} = 0,
$$
\n(27)

where the equation [\(25\)](#page-3-4) is provided by the Euler-Lagrange equations with respect to the scale factor *a*, $\frac{d}{dx} \frac{\partial \mathcal{L}}{\partial x} - \frac{\partial \mathcal{L}}{\partial y} = 0$. The scalar field equation [\(26\)](#page-3-5) arises from $\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{\phi}} - \frac{\partial \mathcal{L}}{\partial \phi} = 0$. Finally, the equation $\frac{\partial \mathcal{L}}{\partial N}\big|_{N=1} = 0$ gives the Friedmann con $\frac{\partial \mathcal{L}}{\partial \dot{\phi}} - \frac{\partial \mathcal{L}}{\partial \phi}$ $\frac{\partial \mathcal{L}}{\partial \phi} = 0$. Finally, the equation $\partial \mathcal{L}/\partial N|_{N=1} = 0$ gives the Friedmann constrain [\(27\)](#page-3-6). As usual, in all the above equations, one can set $N = 1$ after the derivations.

The field equations (25) and (27) can be written in the equivalent form

$$
3H^2 + 3Ka^{-2} = \rho_{\phi},\tag{28}
$$

$$
2\dot{H} + 3H^2 + Ka^{-2} = -p_{\phi},\tag{29}
$$

where ρ_{ϕ} and p_{ϕ} are the cosmological fluid components which correspond to the geometrodynamical degrees of freedom given by the nonlinear $F(B)$ function. They are

$$
\rho_{\phi} = -3\left(H\dot{\phi} + V\left(\phi\right)\right),\tag{30}
$$

$$
p_{\phi} = \ddot{\phi} + 3V(\phi). \tag{31}
$$

Thus, the equation of state parameter for the geometric fluid source is defined as

$$
w_{\phi} = -\frac{\frac{1}{3}\ddot{\phi} + V(\phi)}{H\dot{\phi} + V(\phi)}.
$$
\n
$$
(32)
$$

We observe that, when $\dot{\phi} \approx 0$ and $\ddot{\phi} \approx 0$, it follows $w_{\phi} \approx -1$, that is, the limit of the cosmological constant is recovered. The dust matter domination is provided by $\ddot{\phi} + 3V(\phi) \approx 0$. When $V(\phi) \approx 0$, we have $w_{\phi} \approx -\frac{1}{3}\ddot{\phi}/(H\dot{\phi})$. Therefore, radiationdominated epoch corresponds to $V(\phi) \simeq 0$ together with $\ddot{\phi}/(H\dot{\phi}) \simeq -1$. That is, $V(\phi) \simeq 0$ and $d \ln(\dot{\phi})/d \ln a \simeq -1$, i.e., $\dot{\phi} \simeq a^{-1}$, $\phi \simeq \int a^{-1} dt$ mimics a radiation-dominated universe. These models of $f(T, B)$ gravity offer a unified description of the universe evolution (i.e. the matter era and the late-time acceleration epoch), similarly to the analysis of 65 in scalar-torsion theory. This analysis has been presented before in 29 , where this theory can describe various eras of cosmological history. However, the present work focuses on the existence of spatial curvature and how $f(T, B)$ gravity solves the flatness problem.

In the following sections, we investigate the existence of exact solutions. Also, we study the cosmological dynamics for the field equations $(25)-(27)$ $(25)-(27)$ $(25)-(27)$. Such an analysis provides important information that will help us understand spatial curvature's effects on the application of teleparallelism in cosmology.

3 EXACT SOLUTIONS

We proceed by investigating the existence of exact solutions in which the scale factor is a power-law function, i.e. $a(t) = a_0 t^p$, the exponential function, that is, $a(t) = a_0 e^{H_0 t}$, and the Einstein-static universe, $a(t) = a_0$.

3.1 Scaling solution

For the scaling solution $a(t) = a_0 t^p$, with $H(t) = pt^{-1}$, from the field equations [\(25\)](#page-3-4)-[\(27\)](#page-3-6) we find the linear second-order ordinary differential equation

$$
\ddot{\phi} - 3pt^{-1}\dot{\phi} + 2\left(pt^{-2} + Kt^{-2p}\right) = 0
$$
\n(33)

with analytic solution

$$
\phi(t) = \phi_0 + \frac{\phi_1}{1+3p}t^{1+3p} - \frac{2p}{1+3p} + \frac{K}{(1-p)(1-5p)}t^{2(1-p)}, \ p \notin \left\{1, \frac{1}{5}, -\frac{1}{3}\right\},\tag{34}
$$

$$
\phi(t) = \phi_0 + \frac{\phi_1}{4}t^4 - \frac{1+K}{2}\ln t, \ p = 1,\tag{35}
$$

$$
\phi(t) = \phi_0 + \frac{5}{8} \left(\phi_1 - \frac{5}{4} K \right) t^{\frac{8}{5}} - \frac{1}{4} \left(1 - 5 K t^{\frac{8}{5}} \right) \ln t, \ p = \frac{1}{5},\tag{36}
$$

$$
\phi(t) = \phi_1 + \phi_1 \ln t - \frac{1}{3} (\ln t)^2 + \frac{9}{32} K t^{\frac{8}{3}}, \ p = -\frac{1}{3}.
$$
\n(37)

Similarly, for the scalar field potential $V(\phi(t))$ we derive

$$
V(\phi(t)) = -\phi_1 p t^{-1+3p} + \frac{p^2 (1-p)}{1+3p} t^{-2} - \frac{(1-3p) K}{(1-5p)} t^{-2p}, \ p \notin \left\{1, \frac{1}{5}, -\frac{1}{3}\right\},\tag{38}
$$

$$
V(\phi(t)) = -\phi_1 t^2 - \frac{K+1}{2}t^{-2}, \ p = 1,
$$
\n(39)

$$
V(\phi(t)) = -\frac{(\phi_1 + 5K)}{5}t^{-\frac{2}{5}} + \frac{1}{100}t^{-2} - \frac{2}{5}t^{-\frac{2}{5}}\ln t, \ p = \frac{1}{5},\tag{40}
$$

$$
V(\phi(t)) = \frac{3\phi_1 - 1}{9}t^{-2} - \frac{3}{4}Kt^{\frac{1}{3}} - \frac{2}{9}t^{-2}\ln t, \ p = -\frac{1}{3}.
$$
 (41)

Let us focus now on the case where $p = 1$. The scale factor $a(t) = a_0 t$, describes Milne (for $K = -1$) and Milne-like (for $K = 1$) universes. For $K = -1$, it follows $\phi(t) = \phi_0 + \frac{\phi_1}{4}t^4$ and $V(\phi) = -\phi_1 t^2$, for which we observe that $\rho_{\phi} = 0$. Thus, there is not any contribution to the cosmological fluid from the $F(B)$ component. On the other hand, for the $K = 1$, and the Milne-like solution, we observe that $\rho_{\phi} \neq 0$. For large values of *t*, $\phi(t) \simeq \phi_0 + \frac{\phi_1}{4}$ $\frac{p_1}{4}t^4$ and $V(\phi(t)) \simeq -\phi_1 t^2 - t^{-2}$, that is,

$$
V(\phi) \simeq -\left(4\phi_1\left(\phi - \phi_0\right)\right)^{\frac{1}{2}} - \left(\frac{4}{\phi_1}\left(\phi - \phi_0\right)\right)^{-\frac{1}{2}}.
$$
\n(42)

Furthermore, for small values of *t*, that is, near to the initial singularity, it follows $\phi(t) \simeq -\ln t$, $V(\phi(t)) = -t^{-2}$, i.e.

$$
V(\phi) \simeq -e^{2\phi},\tag{43}
$$

which leads to

$$
F \simeq \frac{1}{2} B \left(\ln \left(\frac{B}{12} \right) - 1 \right). \tag{44}
$$

3.2 Exponential scale factor solution

Assume now the exponential scale factor $a(t) = a_0 e^{H_0 t}$. Then the scalar field satisfies the second-order ordinary differential equation,

$$
\ddot{\phi} - 3H_0 \dot{\phi} - 2Ke^{-2H_0 t} = 0,\tag{45}
$$

with analytic solution

$$
\phi(t) = \phi_0 + \frac{\phi_1}{3H_0} e^{3H_0 t} + \frac{K}{5H_0^2} e^{-2H_0 t}.
$$
\n(46)

For the potential function, we calculate

$$
V(\phi) = 3H_0^2 \left(-\frac{1}{3} + \frac{K}{5H_0^2} e^{-2H_0 t} + \frac{\phi_1}{3H_0} e^{3H_0 t} \right) = 3H_0^2 \left(-\frac{1}{3} + \phi - \phi_0 \right).
$$

Thus, the scalar field potential is described by the linear function $V(\phi) = -\alpha + \beta\phi$, $\alpha = \beta$ (1) $rac{1}{3} + \phi_0$ $\sqrt{ }$, $\beta = 3H_0^2$ which leads to

$$
F(B) = \left(\frac{1}{3} + \phi_0\right)B.\tag{47}
$$

3.3 Einstein-static universe

For a static universe, $a(t) = a_0$, we calculate

$$
\phi(t) = Kt^2 + \phi_0 + \phi_1 t,\tag{48}
$$

$$
V(\phi) = -K,\tag{49}
$$

so $F(B) = F_1 B - 6K$, where F_1 is the integration constant.

Below we continue our analysis by investigating the stability properties of the above solutions and the asymptotic behaviour of the field equations. Specifically, we study the dynamics of the field equations by determining the equilibrium points and their stability.

4 DYNAMICAL ANALYSIS

To study the dynamics of the field equations $(25)-(27)$ $(25)-(27)$ $(25)-(27)$ we define the new variables ^{[63](#page-17-19)}

$$
x = \frac{\dot{\phi}}{\sqrt{H^2 + |K| a^{-2}}}, \ y = \frac{V(\phi)}{H^2 + |K| a^{-2}}, \ \eta = \frac{H}{\sqrt{H^2 + |K| a^{-2}}}, \ \lambda = \frac{V_{,\phi}(\phi)}{V(\phi)}, \tag{50}
$$

which satisfies

$$
(1 - sgn(K))\eta^2 + sgn(K) + \eta x + y = 0.
$$
\n(51)

Furthermore, we consider the new independent variable, $d\tau = \sqrt{H^2 + |K| a^{-2}} dt$, leading to the dynamical system

$$
\frac{dx}{d\tau} = \text{sgn}(K)\left(\eta^2 - 1\right) + \eta\left(3\eta + 2\eta^2 x + x\right) + y(2\lambda + \lambda\eta x - 3),\tag{52}
$$

$$
\frac{dy}{d\tau} = y\left(\lambda x + 2\eta\left(2\eta^2 + \lambda y + 1\right)\right),\tag{53}
$$

$$
\frac{d\eta}{d\tau} = \left(\eta^2 - 1\right) \left(2\eta^2 + \lambda y\right),\tag{54}
$$

$$
\frac{d\lambda}{d\tau} = hx,\tag{55}
$$

where

$$
h = \frac{V_{,\phi\phi}}{V} - \frac{V_{,\phi}^2}{V^2}.
$$
\n(56)

In the new variables the equation of state parameter for the effective fluid, $w_{tot} = -1 - \frac{2}{3}$ \dot{H} $\frac{H}{H^2}$. The deceleration parameter, $q = -1 - \frac{\dot{H}}{H^2}$, is

$$
w_{tot}(x, y, \eta, \lambda) = 1 + \frac{2\lambda}{3\eta^2} y \tag{57}
$$

and

$$
q(x, y, \eta, \lambda) = 2 + \frac{\lambda y}{\eta^2}.
$$
\n(58)

It is useful to define the quantity $\Omega_K = \frac{|K|}{a^2 H}$ $\frac{|\mathbf{A}|}{a^2 H^2}$, which is a dimensionless measure of the spatial curvature in an FLRW universe. Hence, although the analysis is for $K = 1$ and $K = -1$ respectively, some points correspond to "flat FRW", which is the situation where asymptotically $\Omega_K \to 0$. In other words, with the term flat, we mean asymptotically flat. Notice that $\eta^2 = 1/(1 + \Omega_K)$ so that $\Omega_K \to 0$ implies $\eta \to \pm 1$, which means asymptotically flat FRW universe.

Finally,

$$
w_{\phi} = \frac{\text{sgn}(K) - (\text{sgn}(K) + 3)\eta^2 - 2\lambda y}{3(\eta x + y)}.
$$
\n(59)

We proceed with our analysis by considering $K = 1$ and $K = -1$. Moreover, in the following we assume that $\lambda = const$, that is, we consider the exponential potential $V(\phi) = -V_0 e^{\lambda \phi}$, which leads to

$$
\phi = \frac{1}{\lambda} \ln \left(\frac{B}{6\lambda V_0} \right), \quad F(B) = \frac{B}{\lambda} \left(\ln \left(\frac{B}{6\lambda V_0} \right) - 1 \right), \tag{60}
$$

and to $h \equiv 0$.

4.1 Positive curvature

For an FLRW spacetime with positive spatial curvature, i.e. $K = 1$, in the new variables, the field equations become

$$
\frac{dx}{d\tau} = -1 + 4\eta^2 + (2\eta^3 + \eta)x + y(2\lambda + \lambda\eta)x - 3),
$$
\n(61)

$$
\frac{dy}{d\tau} = y\left(\lambda x + 2\eta\left(2\eta^2 + \lambda y + 1\right)\right),\tag{62}
$$

$$
\frac{d\eta}{d\tau} = \left(\eta^2 - 1\right) \left(2\eta^2 + \lambda y\right),\tag{63}
$$

with constraint equation

$$
1 + x\eta + y = 0.\tag{64}
$$

With the use of (64) , we can write the equivalent two-dimensional system

$$
\frac{dx}{d\tau} = -(\eta x + 1)\left(2\lambda - 2\eta^2 + \lambda \eta x - 4\right),\tag{65}
$$

$$
\frac{d\eta}{d\tau} = \left(1 - \eta^2\right) \left(\lambda - 2\eta^2 + \lambda \eta x\right). \tag{66}
$$

The observable quantities are reduced to

$$
\{q, w_{tot}, w_{\phi}\} = \left\{2 - \frac{\lambda(\eta x + 1)}{\eta^2}, 1 - \frac{2\lambda(\eta x + 1)}{3\eta^2}, \frac{1}{3}(-2\lambda + 4\eta^2 - 2\lambda\eta x - 1)\right\}.
$$
 (67)

The equilibrium points $P = (x(P), \eta(P))$ of the dynamical system [\(65\)](#page-6-1), [\(66\)](#page-6-2) are given by the algebraic equations

$$
(\eta x + 1)\left(2\lambda - 2\eta^2 + \lambda \eta x - 4\right) = 0, \quad \left(\eta^2 - 1\right)\left(\lambda - 2\eta^2 + \lambda \eta x\right) = 0,\tag{68}
$$

that is

$$
A_1 = (1, -1), A_2 = (-1, 1),
$$

\n
$$
A_3 = \left(\frac{1}{\sqrt{\lambda}}, -\frac{\sqrt{\lambda}}{2}\right), A_4 = \left(-\frac{1}{\sqrt{\lambda}}, \frac{\sqrt{\lambda}}{2}\right),
$$

\n
$$
A_5 = \left(2 - \frac{6}{\lambda}, -1\right), A_6 = \left(-2 + \frac{6}{\lambda}, 1\right).
$$

Points A_1 , A_2 describe exact solutions for which the kinetic part of the scalar field dominates, i.e. *y*(A_1) = *y*(A_2) = 0, with $w_{tot}(A_1) = w_{tot}(A_2) = 1$ and $q(A_1) = q(A_2) = 2$. The spacetime is described asymptotically by the spatially flat FLRW universe with scale factor $a(t) = a_0 t^{\frac{1}{3}}$.

Moreover, points A_3 and A_4 exist only when $\lambda > 0$ and describe Milne-like solutions with $a(t) = a_0 t$. At these two points the physical parameters are derived $y(A_3) = y(A_4) = -\frac{1}{2}$ $\frac{1}{2}$, $w_{tot}(A_3) = w_{tot}(A_4) = -\frac{1}{3}$ $\frac{1}{3}$ and *q* (*A*₃) = *q* (*A*₄) = 0.

Finally the family of points A_5 and A_6 describe scaling solutions with scale factor $a(t) = a_0 t^{\frac{1}{\lambda - 3}}$. We derive $w_{tot}(A_5)$ $w_{tot}(A_6) = -3 + \frac{2\lambda}{3}$ and $q(A_5) = q(A_6) = \lambda - 4$, from which we observe that the exact solution describes an accelerated universe when $\lambda < 4$. The spatial curvature for the background space is asymptotically zero. In the special case for which $\lambda = 3$, the exact solution at the equilibrium points is $a(t) = a_0 e^{H_0 t}$, which is the de Sitter solution. Evaluating at the equilibrium points A_i , we have $w_{tot} = w_{\phi}$.

To investigate the stability properties of the equilibrium points we determine the eigenvalues of the matrix

$$
\mathbf{A} = \begin{pmatrix} \frac{\partial}{\partial x} \left(\frac{dx}{d\tau} \right) & \frac{\partial}{\partial \eta} \left(\frac{dx}{d\tau} \right) \\ \frac{\partial}{\partial x} \left(\frac{d\eta}{d\tau} \right) & \frac{\partial}{\partial \eta} \left(\frac{d\eta}{d\tau} \right) \end{pmatrix}_{(x,y)\to(x(P),y(P))}.
$$
(69)

Let $e_1(P)$, $e_2(P)$ be the two eigenvalues of the matrix **A**. We say that the equilibrium point *P* is an attractor and describes a stable asymptotic solution when the real parts of the two eigenvalues are negatives. When the real parts of the eigenvalues are positive, point *P* is called a source, and the asymptotic solution is an attractor. Otherwise, the equilibrium point *P* is characterized as a saddle point.

For each of the six equilibrium points, we derive the following set of eigenvalues

$$
e_1(A_1) = -4, e_2(A_1) = \lambda - 6,\tag{70}
$$

$$
e_1(A_2) = 4, e_2(A_2) = -(\lambda - 6),
$$
\n(71)

$$
e_1(A_3) = \frac{1}{2}\sqrt{\lambda}\left(1 + \sqrt{(\lambda - 8)\lambda + 17}\right), e_2(A_3) = \frac{1}{2}\sqrt{\lambda}\left(1 - \sqrt{(\lambda - 8)\lambda + 17}\right),
$$
(72)

$$
e_1(A_4) = -\frac{1}{2}\sqrt{\lambda} \left(1 + \sqrt{(\lambda - 8)\lambda + 17}\right), e_2(A_4) = -\frac{1}{2}\sqrt{\lambda} \left(1 - \sqrt{(\lambda - 8)\lambda + 17}\right),
$$
 (73)

$$
e_1(A_5) = 6 - \lambda, e_2(A_5) = -2(\lambda - 4),
$$
\n(74)

$$
e_1(A_6) = -(6 - \lambda), e_2(A_5) = 2(\lambda - 4). \tag{75}
$$

Points $A_{1,2}$ are nonhyperbolic when $\lambda = 6$. Hence, point A_1 is a sink when $\lambda < 6$; otherwise, it is a saddle. Point A_2 is a source when $\lambda < 6$; otherwise, it is a saddle point. Points A_3 and A_4 exist for $\lambda > 0$, and they are nonhyperbolic for $\lambda = 4$, and a saddle otherwise. The equilibrium points $A_{5,6}$ are nohyperbolic for $\lambda \in \{4, 6\}$. We find that A_5 is a sink for $\lambda > 6$, a source when $\lambda < 4$ and a saddle for $4 < \lambda < 6$, while A_6 is a source for $\lambda > 6$, a sink when $\lambda < 4$ and a saddle for $4 < \lambda < 6$.

The results are summarized in Table [1](#page-9-0). In figure 1 is draw a phase plot of system [\(65\)](#page-6-1), [\(66\)](#page-6-2) for $\lambda = 1, 2, 4, 6$.

4.1.1 Poincaré variables

To perform a complete analysis of the dynamics, we should investigate if there exist equilibrium points when the dynamical variable, x , and η take values at infinity.

Point	Curvature of FLRW	a(t)	w_{ϕ}	Attractor?
A_1	Flat	$t^{\frac{1}{3}}$	-1	λ < 6
A ₂	Flat	$t^{\frac{1}{3}}$	-1	N ₀
A_3	> 0			N ₀
A_4	> 0		$-\frac{1}{3}$ $-\frac{1}{3}$	N ₀
A_5	Flat	$t^{\frac{1}{\lambda-3}}, \lambda \neq 3 \quad -3 + \frac{2\lambda}{3}$		$\lambda > 6$
		$e^{H_0 t}$, $\lambda = 3$	-1	N ₀
A ₆	Flat	$t^{\frac{1}{\lambda-3}}, \lambda \neq 3 \quad -3 + \frac{2\lambda}{3}$		$\lambda < 4$
		$e^{H_0 t}$, $\lambda = 3$	-1	Yes

TABLE 1 Asymptotic solutions for the field equations with positive spatial curvature.

TABLE 2 Asymptotic solutions at the infinity for the field equations with positive spatial curvature.

	Point (X, Z) $a(t)$		Stability
P_{1}	(1,0)	a_{α}	Sink for $\lambda < 0$, source for $\lambda > 0$
P_{2}	$(-1,0)$	a_{0}	Source for $\lambda < 0$, sink for $\lambda > 0$

Thus, we define the Poincaré variables

$$
x = \frac{X}{\sqrt{1 - X^2 - Z^2}}, \ \eta = \frac{Z}{\sqrt{1 - X^2 - Z^2}}
$$
\n(76)

and the new independent variable $d\sigma = (1 - X^2 - Z^2)^{-\frac{1}{2}} d\tau$.

The two-dimensional dynamical system [\(65\)](#page-6-1), [\(66\)](#page-6-2) becomes

$$
\frac{dX}{d\sigma} = -2(\lambda + (\lambda - 2)X^4 - 2(\lambda - 1)X^3Z + X^2(-2\lambda + (2\lambda - 1)Z^2 + 4) - XZ(-2\lambda + (\lambda + 2)Z^2 + 2) - (\lambda - 1)Z^2 - 2),
$$
\n(77)

$$
\frac{dZ}{d\sigma} = \lambda - 2(\lambda - 2)X^3 Z + X^2 \left(4(\lambda - 1)Z^2 - \lambda\right) + XZ \left(3\lambda + (2 - 4\lambda)Z^2 - 4\right) + Z^2 \left(-3\lambda + 2(\lambda + 2)Z^2 - 2\right). \tag{78}
$$

The equilibrium points of the latter system at infinity that is, on the surface $1 - X^2 - Z^2 = 0$, are

$$
P_1=(1,0)\,,\ P_2=(-1,0)\,.
$$

Furthermore, in the new variables for the deceleration parameter, the effective equation of state parameter, and the effective equation of state parameter of ϕ , we derive

$$
q(X,Z) = \lambda + \frac{\lambda \left(X^2 - XZ - 1\right)}{Z^2} + 2,\tag{79}
$$

$$
w_{tot}(X,Z) = \frac{2\lambda (X^2 - XZ + Z^2 - 1)}{3Z^2} + 1,
$$
\n(80)

and

$$
w_{\phi}(X,Z) = \frac{1}{3} \left(-2\lambda + \frac{2(2X^2 + \lambda XZ - 2)}{X^2 + Z^2 - 1} - 5 \right).
$$
 (81)

Thus, points P_1 and P_2 describe static universes with $a(t) = a_0$. We proceed with the study of the stability properties of the equilibrium points at infinity by using the parametrization

$$
X = (1 - \rho)\cos(\varphi), \quad Z = (1 - \rho)\sin(\varphi), \quad 0 \le \rho < 1,
$$
\n(82)

and a time re-scaling $ds = d\sigma/(1 - \rho)$, such the region at infinite for (x, η) , i.e, $X^2 + Z^2 = 1$, is approached as $\rho \to 0^+$.

FIGURE 1 Phase plot of system [\(65\)](#page-6-1), [\(66\)](#page-6-2) for $\lambda = 1, 2, 4, 6$

Taking the Taylor expansion centred in $\rho = 0$, neglecting higher order terms, we have

$$
\frac{d\varrho}{ds} = \sin^2(\varphi)(\lambda\cos(\varphi) - 2\sin(\varphi)), \quad \frac{d\varphi}{ds} = \frac{1}{2}\sin(\varphi)(\lambda + 2(\lambda - 3)\sin(2\varphi) + (\lambda + 2)\cos(2\varphi) - 2). \tag{83}
$$

To find the equilibrium points at infinity, we solve the algebraic equation

$$
\sin^2(\varphi)(\lambda\cos(\varphi) - 2\sin(\varphi)) = 0, \quad \sin(\varphi)(\lambda + 2(\lambda - 3)\sin(2\varphi) + (\lambda + 2)\cos(2\varphi) - 2) = 0.
$$
 (84)

Equations [\(83\)](#page-9-1) do not depend on the radial coordinate. Therefore, the stability analysis considers the nature of the eigenvalues

$$
\Lambda_1(P) = 0, \quad \Lambda_2(P) = \frac{1}{4}(-2(\lambda - 3)(\sin(\varphi) - 3\sin(3\varphi)) + (\lambda - 6)\cos(\varphi) + 3(\lambda + 2)\cos(3\varphi)).
$$
\n(85)

of the Jacobian matrix evaluated at the values φ that satisfy [\(84\)](#page-9-2).

The solutions of [\(84\)](#page-9-2) are

 $P_1 := \varphi = 2\pi c_1$ if $c_1 \in \mathbb{Z}$, with $\Lambda_1(P_1) = 0, \Lambda_1(P_1) = \lambda$. Sink for $\lambda < 0$, source for $\lambda > 0$. $P_2 := \varphi = 2\pi c_1 + \pi$ if $c_1 \in \mathbb{Z}$, with $\Lambda_1(P_2) = 0$, $\Lambda_1(P_2) = -\lambda$. Source for $\lambda < 0$, sink for $\lambda > 0$. The results are summarized in Table [2](#page-8-1) .

FIGURE 2 Phase plot of system [\(77\)](#page-8-2)-[\(78\)](#page-8-3) for $\lambda = 1, 2, 4, 6$

In figure [2](#page-10-0) a phase plot of system [\(77\)](#page-8-2)-[\(78\)](#page-8-3) is presented for $\lambda = 1, 2, 4$ and 6. In the figures it is confirmed for $\lambda > 0$ that P_1 P_1 is unstable and P_2 is stable. Additionally, the information in table 1 relative to the stability of the points at the finite region A_i , $i = 1...6$ is confirmed.

4.2 Negative curvature

For an FLRW spacetime with negative spatial curvature, the field equations in the dimensionless variables are

$$
\frac{dx}{d\tau} = \left(2\eta^2 + 1\right)(\eta x + 1) + y(2\lambda + \lambda \eta x - 3),\tag{86}
$$

$$
\frac{dy}{d\tau} = y\left(\lambda x + 2\eta\left(2\eta^2 + \lambda y + 1\right)\right),\tag{87}
$$

$$
\frac{d\eta}{d\tau} = \left(\eta^2 - 1\right) \left(2\eta^2 + \lambda y\right),\tag{88}
$$

with constraint

$$
\eta(2\eta + x) + y - 1 = 0.\tag{89}
$$

Point	FLRW	a(t)	w_{ϕ}	Attractor?
B_1	Flat	$t^{\frac{1}{3}}$		λ < 6
B ₂	Flat	$t^{\frac{1}{3}}$		N ₀
B_3	< 0	t		N ₀
B_4	< 0	t		$\lambda > 4$
B_5	Flat	$t^{\frac{1}{\lambda-3}}, \lambda \neq 3$		$\lambda > 6$
		$e^{H_0 t}$, $\lambda = 3$		N ₀
B_{6}	Flat	$t^{\frac{1}{\lambda-3}}, \lambda \neq 3$		$\lambda < 4$
		$e^{H_0 t}$, $\lambda = 3$		Yes

TABLE 3 Asymptotic solutions for the field equations with negative spatial curvature.

Thus, with the use of the constraint equation [\(89\)](#page-10-1), the dynamical system is reduced to the two-dimensional system

$$
\frac{dx}{d\tau} = (2\eta^2 + 1)(\eta x + 1) - (\eta(2\eta + x) - 1)(2\lambda + \lambda \eta x - 3),\tag{90}
$$

$$
\frac{d\eta}{d\tau} = (\eta^2 - 1)(\lambda - \eta(2(\lambda - 1)\eta + \lambda x)).
$$
\n(91)

The observable quantities are reduced to

$$
\{q, w_{tot}, w_{\phi}\} = \left\{-2\lambda + \frac{\lambda - \lambda\eta x}{\eta^2} + 2, \frac{1}{3}\left(-4\lambda - \frac{2\lambda(\eta x - 1)}{\eta^2} + 3\right), \frac{2\lambda + 2\eta(\eta - \lambda(2\eta + x)) + 1}{6\eta^2 - 3}\right\}.
$$
 (92)

The equilibrium points for the dynamical system (90) , (91) are

$$
B_1 = (1, -1), B_2 = (-1, 1),
$$

$$
B_3 = \left(\sqrt{\frac{2}{\lambda(\lambda - 2)}}, -\sqrt{\frac{\lambda}{2(\lambda - 2)}}\right), B_4 = \left(-\sqrt{\frac{2}{\lambda(\lambda - 2)}}, \sqrt{\frac{\lambda}{2(\lambda - 2)}}\right),
$$

$$
B_5 = \left(2 - \frac{6}{\lambda}, -1\right), B_6 = \left(-2 + \frac{6}{\lambda}, 1\right).
$$

The physical properties of the asymptotic solutions at the latter equilibrium points are similar to those of points A_i . Indeed, points B_1 , B_2 describe spatially flat FLRW spacetimes with scale factor $a(t) = a_0 t^{\frac{1}{3}}$, points B_3 and B_4 are real when $\lambda(\lambda - 2) > 0$ and describe Milne universes, while for the points B_5 and B_6 the asymptotic solution is that of spatially flat FLRW with scale factor $a(t) = a_0 t^{\frac{1}{\lambda-3}}$, $\lambda \neq 3$ or $a(t) = a_0 e^{H_0 t}$, $\lambda = 3$. Evaluating at the equilibrium points B_i , we have $w_{tot} = w_{\phi}$.

The eigenvalues of the linearized system (90) , (91) around the equilibrium points B_i are calculated

$$
e_1 (B_1) = -4, e_2 (B_1) = \lambda - 6,
$$

\n
$$
e_1 (B_2) = 4, e_2 (B_2) = -\lambda + 6,
$$
\n(94)

$$
e_1 (B_2) = 4, e_2 (B_2) = -\lambda + 6,
$$
\n(94)

$$
e_1(B_3) = \left(1 - \sqrt{17 - 4\lambda}\right)\sqrt{\frac{\lambda}{2(\lambda - 2)}}, \ e_2(B_3) = \left(1 + \sqrt{17 - 4\lambda}\right)\sqrt{\frac{\lambda}{2(\lambda - 2)}},\tag{95}
$$

$$
e_1(B_4) = -\left(1 - \sqrt{17 - 4\lambda}\right) \sqrt{\frac{\lambda}{2(\lambda - 2)}}, \ e_2(B_4) = -\left(1 + \sqrt{17 - 4\lambda}\right) \sqrt{\frac{\lambda}{2(\lambda - 2)}}, \tag{96}
$$

$$
e_1(B_5) = 6 - \lambda, \ e_2(B_5) = -2(\lambda - 4), \tag{97}
$$

$$
e_1(B_6) = -(6 - \lambda), \ e_2(B_6) = 2(\lambda - 4). \tag{98}
$$

The stability properties of points B_1 , B_2 , B_5 and B_6 are the same as those of points A_1 , A_2 , A_5 and A_6 , respectively. Point B_3 is a source for $4 < \lambda \leq \frac{17}{4}$ ¹⁷ (unstable node) or $\lambda > \frac{17}{4}$ (unstable spiral), or a saddle point for $\lambda < 0$ or $2 < \lambda < 4$. On the other hand, point B_4 is an attractor for $4 < \lambda \leq \frac{17}{4}$ $\frac{17}{4}$ (stable node) or $\lambda > \frac{17}{4}$ (stable spiral), or a saddle point for $\lambda < 0$ or $2 < \lambda < 4$. The results are summarized in Table [3](#page-12-0). In figure 3 is draw a phase plot of system (90) , (91) for $\lambda = 1, 3, 4, 6$.

FIGURE 3 Phase plot of system [\(65\)](#page-6-1), [\(66\)](#page-6-2) for $\lambda = 1, 3, 4, 6$

4.2.1 Poincaré variables

We introduce the Poincaré variables [\(76\)](#page-8-4), and we write the dynamical system [\(90\)](#page-11-0), [\(91\)](#page-11-1) into the equivalent form

$$
\frac{dX}{d\sigma} = 2(\lambda + (\lambda - 1)X^4 - 2X^3Z + X^2(-2\lambda + (2\lambda - 5)Z^2 + 2) + XZ((2 - 3\lambda)Z^2 + 2) + (5 - 3\lambda)Z^2 - 1),
$$
\n(99)
\n
$$
\frac{dZ}{dZ} = \frac{2(2Z(Y - Z) + 1)(Y^2 + YZ + 2Z^2 - 1)}{2Z(X^3 + 2Y^2Z + Y(5Z^2 - 1) - 2Z^3 + Z)}.
$$

$$
\frac{dZ}{d\sigma} = \lambda(2Z(X-Z)+1)\left(X^2+XZ+3Z^2-1\right)-2Z\left(X^3+2X^2Z+X\left(5Z^2-1\right)-2Z^3+Z\right). \tag{100}
$$

Furthermore, the physical variables in the new variables are

$$
q(X,Z) = -3\lambda + \frac{\lambda - \lambda X(X+Z)}{Z^2} + 2,\tag{101}
$$

$$
w_{tot}(X, Z) = \frac{2\lambda - 2\lambda X^2 - 2\lambda X Z + (3 - 6\lambda)Z^2}{3Z^2},
$$
\n(102)

and

$$
w_{\phi}(X,Z) = \frac{1}{9} \left(-6\lambda + \frac{-4X^2 - 6\lambda XZ + 4}{X^2 + 3Z^2 - 1} + 1 \right).
$$
 (103)

TABLE 4 Asymptotic solutions at the infinity for the field equations with negative spatial curvature.

	Point (X, Z) $a(t)$		Stability
Q_1	(1,0)	a_0	Sink for $\lambda < 0$, source for $\lambda > 0$
\boldsymbol{U}_2	$(-1,0)$	a_0	Source for $\lambda < 0$, sink for $\lambda > 0$

The equilibrium points for the dynamical system [\(99\)](#page-12-1), [\(100\)](#page-12-2) at infinity, that is, on the surface $X^2 + Z^2 = 1$, are

$$
Q_1 = (1,0), Q_2 = (-1,0).
$$

Points Q_1 , Q_2 describe static universe, $a(t) = a_0$, similarly with points P_1 , P_2 . To investigate the stability of the equilibrium points at infinite, we use the parameterization

$$
X = (1 - \rho)\cos(\varphi), \quad Z = (1 - \rho)\sin(\varphi), \quad 0 \le \rho < 1,
$$
 (104)

and a time re-scaling $ds = d\sigma/(1 - \rho)$, such as the region at infinite for (x, η) , i.e, $X^2 + Z^2 = 1$, is approached as ρ tends to 0.

Taking the Taylor expansion centred in $\rho = 0$ and neglecting error terms, we obtain that

$$
\frac{d\varrho}{ds} = \sin^2(\varphi)(2(\lambda - 1)\sin(\varphi) + \lambda\cos(\varphi)), \quad \frac{d\varphi}{ds} = \frac{1}{2}\sin(\varphi)(5\lambda + (4\lambda - 6)\sin(2\varphi) + (8 - 3\lambda)\cos(2\varphi) - 8). \tag{105}
$$

To find the equilibrium points at infinity, we solve the algebraic equation

$$
\sin^2(\varphi)(2(\lambda - 1)\sin(\varphi) + \lambda\cos(\varphi)) = 0, \quad \sin(\varphi)(5\lambda + (4\lambda - 6)\sin(2\varphi) + (8 - 3\lambda)\cos(2\varphi) - 8) = 0.
$$
 (106)

In the first order, equations [\(105\)](#page-13-1) do not depend upon the radial coordinate. Therefore, the stability analysis considers the nature of the eigenvalues

$$
\Lambda_1(Q) = 0, \quad \Lambda_2(Q) = \frac{1}{4}(-2(2\lambda - 3)(\sin(\varphi) - 3\sin(3\varphi)) + (13\lambda - 24)\cos(\varphi) + (24 - 9\lambda)\cos(3\varphi))\tag{107}
$$

of the Jacobian matrix valuated at the values, φ , that satisfy [\(106\)](#page-13-2).

The solutions of (106) are

 $Q_1 := \varphi = 2\pi c_1$ if $c_1 \in \mathbb{Z}$, with $\Lambda_1(Q_1) = 0, \Lambda_1(Q_1) = \lambda$. Sink for $\lambda < 0$, source for $\lambda > 0$. $Q_2 := \varphi = 2\pi c_1 + \pi$ if $c_1 \in \mathbb{Z}$, with $\Lambda_1(Q_2) = 0, \Lambda_2(Q_2) = -\lambda$. Source for $\lambda < 0$, sink for $\lambda > 0$.

The dynamics at infinity ($\rho = 0$) are governed by the one-dimensional dynamical system [\(105\)](#page-13-1). In figure [4](#page-14-0) a phase plot of system [\(99\)](#page-12-1)-[\(100\)](#page-12-2) is presented for $\lambda = 1, 3, 4, 6$. In the figures, it is confirmed that for $\lambda > 0$, Q_1 is unstable and Q_2 is stable. Moreover, the information in table [3](#page-11-2) relative to the stability of the points at the finite region B_i , $i = 1...6$ is confirmed. The stability of the rest of the points at the infinite region is as discussed in table [4](#page-13-3) .

5 CONCLUSIONS

In this piece of work, we studied the cosmological model of the fourth-order teleparallel theory gravity known as $f(T, B)$ in the case of FLRW background space with nonzero spatial curvature. In particular, we assumed that $f(T, B)$ is a linear function of *T*, $f(T, B) = T + F(B)$, such that the modifications of the gravitational Action Integral of the TEGR to be introduced by the term $F(B)$. In the case where the function $F(B)$ is linear, the TEGR is recovered; thus, in this work, we consider $F(B)$ to be a nonlinear function.

For the proper vierbein fields, we derived the field equations for the background space of our consideration. Using Lagrange multipliers, we introduced a scalar field that attributes the geometrodynamical degrees of freedom to writing the field equations in the form of second-order theory. The dynamical variables are the scale factor $a(t)$ and the scalar field $\phi(t)$. Moreover, we show that this cosmological model admits a minisuperspace description. Thus, there exists a point-like Lagrangian, the variation of which provides the field equations. That specific characteristic of the gravitational theory is essential because various techniques from analytic mechanics can be applied for the investigation of the differential equations, also, the Wheeler-DeWitt equation of quantum cosmology can be calculated straightforward.

FIGURE 4 Phase plot of system [\(99\)](#page-12-1)-[\(100\)](#page-12-2) for $\lambda = 1, 3, 4, 6$

We investigated the existence of exact solutions where the scale factor is a power-law function, scaling solution, or exponential function. These exact solutions are essential because they can describe specific eras of cosmological history. Moreover, we proved that both cases have a scalar field potential, so the field equations can be solved explicitly.

Assume now, an arbitrary scale factor $a(t)$, then from equations [\(25\)](#page-3-4) and [\(27\)](#page-3-6) it follows

$$
\ddot{\phi} - H\dot{\phi} - 2Ka^{-2} + 2\dot{H} = 0,\tag{108}
$$

which is a second-order equation of the form

$$
\ddot{\phi} + \alpha(t)\dot{\phi} + \beta(t) = 0 \tag{109}
$$

with $\alpha(t) = -H$ and $\beta(t) = -2Ka^{-2} + 2\dot{H}$. Equation [\(109\)](#page-14-1) is a linear equation and it is maximally symmetric, which means that it always admits a solution for an arbitrary function $\alpha(t)$ and $\beta(t)$, that is, for arbitrary selection of the scale factor $a(t)$. Hence, it is easy to infer that the field equations in these cosmological scenarios are always integrable. Such analysis generalizes previous results on this theory in the case of a spatially flat FLRW geometry.

Finally, we studied the general evolution of the dynamical variables described by the field equations. In particular, we derived the equilibrium points and investigated their stability properties. We performed our analysis separately for the cases of positive and negative spatial curvature. The equilibrium points were studied in the finite and infinite regions using Poincaré variables.

While, from a first read, it seems that there are similarities in the cosmological evolution for the two cases of positive and negative spatial curvature, from the detailed analysis, we found that the results differ.

The results of this work are essential to understanding the spatial curvature in teleparallelism. The $f(T, B)$ theory provides important asymptotic behaviours of particular interest. In future work, we plan to study further the applications of $f(T, B)$ theory in the pre-inflationary era.

6 CONFLICT OF INTEREST STATEMENT

The authors declare to have no conflict of interest.

7 AUTHOR CONTRIBUTIONS

The authors equally contributed.

8 ACKNOWLEDGEMENTS

This work is based on the research supported in part by the National Research Foundation of South Africa (Grant Numbers 131604). Additionally, this research is funded by Vicerrectoría de Investigación y Desarrollo Tecnológico (Vridt) at Universidad Católica del Norte. G. L. was funded through Concurso De Pasantías De Investigación Año 2022, Resolución Vridt No. 040/2022 and through Resolución Vridt No. 054/2022.

References

- 1. T. Clifton, P.G. Ferreira, A. Padilla and C. Skordis, Phys. Rept. 513, 1 (2012)
- 2. S. Nojiri, S.D. Odintsov and V.K. Oikonomou, Phys. Rept. 692, 1 (2017)
- 3. M. Ishak, Living Rev. Rel. **22** (2019) no.1, 1
- 4. M. Tegmark et al., Astrophys. J. 606, 702 (2004)
- 5. M. Kowalski et al., Astrophys. J. 686, 749 (2008)
- 6. E. Komatsu et al., Astrophys. J. Suppl. Ser. 180, 330 (2009)
- 7. P. A. R. Ade et al., (Planck Collaboration), Astron. Astroph. 594, A13 (2016)
- 8. E. Di Valentino et al., Astropart. Phys. 131, 102605 (2021)
- 9. E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F Mota, A.G Riess and J. Silk, Class. Quantum Grav. 38, 153001 (2021)
- 10. T.P. Sotiriou and V. Faraoni, Rev. Mod. Phys. 82, 451 (2010)
- 11. D. Glavan and C. Lin, Phys. Rev. Lett. 124, 081301 (2020)
- 12. J.B. Jimenez, L. Heisenberg, T.S. Koivisto and S. Pekar, Phys. Rev.D 101, 103507 (2020)
- 13. R. Weitzenböck, Invarianten Theorie, Nordhoff, Groningen (1923)
- 14. A. Einstein 1928, Sitz. Preuss. Akad. Wiss. p. 217; ibid p. 224 [Translated by A. Unzicker and T. Case, (preprint: arXiv: physics/0503046)]

- 15. L. Combi and G.E. Romero, Annalen der Physik 530, 1700175 (2018)
- 16. R. Ferraro and F. Fiorini, Phys. Rev. D 75, 084031 (2007)
- 17. R. Ferraro and F. Fiorini, Phys. Rev. D 78, 124019 (2008)
- 18. C. Xu, E.N. Saridakis and G. Leon, JCAP 07, 005 (2012)
- 19. R. Myrzakulov, Gen. Rel. Gravit. 44, 3059 (2012)
- 20. M. Krssak, R. J. van den Hoogen, J. G. Pereira, C. G. Boehmer and A. A. Coley, Class. Quantum Grav. 36, 183001 (2019)
- 21. M. Ilyas, Int. J. Geom. Meth. Mod. Phys. 16, 1950149 (2019)
- 22. J. Yang, Y.-L. Li, Y. Zhong, and Y. Li, Phys. Rev. D 85, 084033 (2012)
- 23. J. Wang, W.-D. Guo, Z.-C. Lin, and Y.-X. Liu, Phys. Rev. D 98, 084046 (2018)
- 24. Q. Tan, W.-D. Guo, Y.-P. Zhang, and Y.-X. Liu, Eur. Phys. J. C 81, 373 (2021)
- 25. S. Bahamonte, K.F. Dialektopoulos, C. Escamilla-Rivera, G. Farrugia, V. Gakis, M. Hohmann, J.L. Said, J. Mifsud and E. Di Valentino, Teleparallel Gravity: From Theory to Cosmology, arXiv:2106.13793 (2021)
- 26. Y. F. Cai, S. Capozziello, M. De Laurentis and E. N. Saridakis, Rept. Prog. Phys. 79, no.10, 106901 (2016)
- 27. S. Bahamonde, C. G. Bohmer and M. Wright, Phys. Rev. D 92, 104042 (2015)
- 28. R. Myrzakulov, EPJC 72, 1 (2012)
- 29. A. Paliathanasis, Phys. Rev. D 95, 064062 (2017)
- 30. A. Paliathanasis and G. Leon, Eur. Phys. J. Plus 136, 1092 (2021)
- 31. G.A. Rave-Franco, C. Escamilla-Rivera and J.L. Said, EPJC 80, 677 (2020)
- 32. G.A. Rave-Franco, C. Escamilla-Rivera and J.L. Said, Phys. Rev. D 103, 084017 (2021)
- 33. A. Paliathanasis, JCAP 1708, 027 (2017)
- 34. L. Karpathopoulos, S. Basilakos, G. Leon, A. Paliathanasis and M. Tsamparlis, Gen. Rel. Gravit. 50, 79 (2018)
- 35. M. Caruana, G. Farrugia and J.L. Said, EPJC 80, 640 (2020)
- 36. S. Najera, A. Aguilar, C. Escamilla-Rivera and R.A. Sussman, The first non-static inhomogeneous exact solution in f(T,B) gravity, (2021) [arXiv:2106.02247]
- 37. A. Paliathanasis, Universe 7, 150 (2021)
- 38. C. Escamilla-Rivera, J.L. Said, EPJC 80, 677 (2020)
- 39. S. Bahamonde, V. Gaskis, S. Kiorpelidi, T. Koivisto and J.L. Said, EPJC 81, 53 (2021)
- 40. C. Escamilla-Rivera and J. Levi Said, Class. Quant. Grav. 37, 165002 (2020)
- 41. G. Farrugia, J.L. Said and A. Finch, Universe, 6, 34 (2020)
- 42. S. Bahamonde, A. Golovnev, M. J. Guzmán, J. L. Said and C. Pfeifer, JCAP01 (2022) no.01, 037
- 43. G. Farrugia, J. Levi Said and A. Finch, Universe 6 (2020) no.2, 34
- 44. S. Bahamonde, M. Zubair and G. Abbas, Phys. Dark Univ. 19 (2018), 78-90
- 45. M. Zubair, L. R. Durrani and S. Waheed, Eur. Phys. J. Plus 136 (2021) no.9, 943

- 46. M. Zubair, S. Waheed, M. Atif Fayyaz and I. Ahmad, Eur. Phys. J. Plus 133 (2018) no.11, 452
- 47. A. Paliathanasis, Eur. Phys. J. Plus **137**, no.8, 887 (2022)
- 48. G. Leon and A. Paliathanasis, Eur. Phys. J. Plus **137**, no.7, 855 (2022)
- 49. G. Leon and A. Paliathanasis, Eur. Phys. J. Plus **137**, no.8, 927 (2022)
- 50. G. Aslanyan and R. Easther, Phys. Rev. D 91, 123523 (2015)
- 51. G.J. Mathews, I.-S. Suh, N.Q. Lan and T. Kajino, Phys. Rev. D 92, 123514 (2015)
- 52. K. Sato, MNRAS 195, 467 (1981)
- 53. J.D. Barrow and A. Ottewill, J. Phys. A 16, 2757 (1983)
- 54. A. Paliathanasis, Mod. Phys. Lett. A 36, 2150261 (2021)
- 55. M.A. Skugoreva and A.V. Toporensky, EPJC 80, 1054 (2020)
- 56. A.A. Coley, Dynamical Systems and Cosmology, Springer, Dordrecht (2003)
- 57. G. Leon, Class. Quantum Grav. 26, 035008 (2009)
- 58. R. De Arcia, T. Gonzalez, F.A. Horta-Rangel, G. Leon, U. Nucamendi and I. Quiros, Class. Quantum Grav. 35, 145001 (2018)
- 59. J. M. Heinzle, N. Rohr and C. Uggla, Phys. Rev. D 71, 083506 (2005)
- 60. L. Amendola, D. Polarski, S. Tsujikawa, Phys. Rev. Lett. 98, 131302 (2007)
- 61. L. Amendola, R. Gannouji, D. Polaski and S. Tsujikawa, Phys. Rev. D 75, 083504 (2007)
- 62. F. Fiorini and R. Ferraro, Int. J. Mod. Phys.: Conf. Series 3, 227 (2011)
- 63. M. Kerachian, G. Acquaviva and G. Lukes-Gerakopoulos, Phys. Rev. D 101, 043535 (2020)
- 64. A. C. Clairaut, "Solution de plusieurs problèmes où il s'agit de trouver des Courbes dont la propriété consiste dans une certaine relation entre leurs branches, exprimée par une Équation donnée.", Histoire de l'Académie royale des sciences: 196–215 (1734).
- 65. G. Leon, A. Paliathanasis, E. N. Saridakis and S. Basilakos, Phys. Rev. D **106**, no.2, 024055 (2022)

How to cite this article: A. Paliathanasis, and G. Leon (2022), $f(T, B)$ gravity in a Friedmann–Lemaître–Robertson–Walker universe with nonzero spatial curvature, *Math Meth Appl Sci.* 2022;1-18. <https://doi.org/10.1002/mma.8728>