All even (unitary) perfect polynomials over \mathbb{F}_2 with only Mersenne primes as odd divisors

Gallardo, Luis H., Rahavandrainy, Olivier, Univ Brest, UMR CNRS 6205, Laboratoire de Mathématiques de Bretagne Atlantique e-mail : Luis.Gallardo@univ-brest.fr - Olivier.Rahavandrainy@univ-brest.fr

February 15, 2022

- a) Running head: Mersenne and perfect polynomials
- b) Keywords: Sum of divisors, polynomials, finite fields, characteristic 2.
- c) Mathematics Subject Classification (2010): 11T55, 11T06.
- d) Corresponding author: Luis H. Gallardo

Abstract

We address an arithmetic problem in the ring $\mathbb{F}_2[x]$ related to the fixed points of the sum of divisors function. We study some binary polynomials A such that $\sigma(A)/A$ is still a binary polynomial. Technically, we prove that the only (unitary) perfect polynomials over \mathbb{F}_2 that are products of x, x + 1and of Mersenne primes are precisely the nine (resp. nine "classes") known ones. This follows from a new result about the factorization of $M^{2h+1} + 1$, for a Mersenne prime M and for a positive integer h.

1 Introduction

Let $A \in \mathbb{F}_2[x]$ be a nonzero polynomial. We say that A is even if it has a linear factor and that it is odd otherwise. We define a Mersenne prime (polynomial) over \mathbb{F}_2 as an irreducible polynomial of the form $1 + x^a(x+1)^b$, for some positive integers a, b. The name come as an analogue of the integral Mersenne primes, taking $x^a(x+1)^b$ as an analogue of the prime power 2^{a+b} . As over the integers, we say that a divisor d of A is unitary if gcd(d, A/d) = 1. Let $\omega(A)$ denote the number of distinct irreducible (or prime) factors of Aover \mathbb{F}_2 and let $\sigma(A)$ (resp. $\sigma^*(A)$) denote the sum of all (unitary) divisors of A (including 1 and A). Both σ and σ^* are multiplicative functions. If $\sigma(A) = A$ (resp. $\sigma^*(A) = A$), then we say that A is (unitary) perfect. Finally, we say that a (unitary) perfect polynomial is indecomposable if it is not a product of two coprime nonconstant (unitary) perfect polynomials.

We can also consider a perfect polynomial $(A \in \mathbb{F}_2[x] \text{ such that } A \text{ divides} \sigma(A))$ as an analogue of a multiperfect number (a positive integer which divides the sum of all its divisor). It might have some interest to observe that most known multiperfect numbers (see OEIS sequence A007691) appear to be divisible by a Fermat prime or by a Mersenne prime.

The notion of (unitary) perfect polynomials is introduced in [5] by E. F. Canaday in 1941 and extended by J. T. B. Beard Jr. et al. in several directions ([1], [2], [4]). Later research in the subject by Gallardo and Rahavandrainy ([6], [7]) allows us to be able to better describe the properties of such polynomials.

The known perfect polynomials are the following:

- the so-called "trivial" ones, of the form $(x^2 + x)^{2^n-1}$, for some positive integer n,

- nine others which are already characterized ([8], Theorem 1.1),

- and the last two which are divisible by a non-Mersenne prime.

Any unitary perfect polynomial is even (Lemma 2.12). The known ones, which are only divisible by Mersenne primes (as odd factors), belong to nine equivalence classes (see Lemma 2.14). There are several (perhaps, infinitely many) of such classes (see [2] and [12]).

The paper consist of two major results stated in Theorem 1.1 and Theorem 1.2. Theorem 1.1 significantly improves on the results of [8, Theorems 1.1 and 1.3], because (in our main result in this paper) there are no more conditions asked on the powers of the M_j . The proof of Theorem 1.1 is obtained from new results given in Theorem 1.2, which in turn, extends recent results in [10, Theorem 1.4].

It is convenient to fix some notations.

Notations

• The set of integers (resp. of nonnegative integers, of positive integers) is denoted by \mathbb{Z} (resp. \mathbb{N} , \mathbb{N}^*).

• For $S, T \in \mathbb{F}_2[x]$ and for $m \in \mathbb{N}^*$, $S^m \mid T$ (resp. $S^m \mid T$) means that S divides T (resp. $S^m \mid T$ but $S^{m+1} \nmid T$). We also denote by \overline{S} the polynomial defined as $\overline{S}(x) = S(x+1)$ and by $val_x(S)$ (resp. $val_{x+1}(S)$) the valuation of S, at x (resp. at x + 1).

• We put

$$\begin{split} M_{j} &= 1 + x(x+1)^{j}, j \in \{1,2,3\}, \\ T_{1} &= x^{2}(x+1)M_{1}, T_{2} = \overline{T_{1}}, \\ T_{3} &= x^{4}(x+1)^{3}M_{3}, T_{4} = \overline{T_{3}}, \ T_{5} = x^{4}(x+1)^{4}M_{3}\overline{M_{3}} = \overline{T_{5}}, \\ T_{6} &= x^{6}(x+1)^{3}M_{2}\overline{M_{2}}, T_{7} = \overline{T_{6}}, \\ T_{8} &= x^{4}(x+1)^{6}M_{2}\overline{M_{2}}M_{3} \text{ and } T_{9} = \overline{T_{8}}, \\ B_{1} &= x^{3}(x+1)^{3}M_{1}^{2}, \ B_{2} = x^{3}(x+1)^{2}M_{1}, \ B_{3} = x^{5}(x+1)^{4}M_{3}, \\ B_{4} &= x^{7}(x+1)^{4}M_{2}\overline{M_{2}}, \ B_{5} = x^{5}(x+1)^{6}M_{1}^{2}M_{3}, \ B_{6} = x^{5}(x+1)^{5}M_{3}\overline{M_{3}}, \\ B_{7} &= x^{7}(x+1)^{7}M_{2}^{2}\overline{M_{2}}^{2}, \ B_{8} = x^{7}(x+1)^{6}M_{1}^{2}M_{2}\overline{M_{2}}, \ B_{9} = x^{7}(x+1)^{5}M_{2}\overline{M_{2}} \overline{M_{3}}, \\ \mathcal{M} &= \{M_{1}, M_{2}, \overline{M_{2}}, M_{3}, \overline{M_{3}}\}, \ \mathcal{P} = \{T_{1}, \dots, T_{9}\} \text{ and } \mathcal{P}_{u} = \{B_{1}, \dots, B_{9}\}. \end{split}$$

• Finally, we denote by Δ , the set of primes p such that p is a Mersenne prime or the order of 2 in $\mathbb{F}_p \setminus \{0\}$, $ord_p(2)$, is divisible by 8. In particular, Δ contains all Fermat primes greater than 5.

Throughout this paper, we **always** suppose that any (unitary) perfect polynomial is **indecomposable**. We have often used Maple software for computations. Our main results are the following.

Theorem 1.1. Let $A = x^a(x+1)^b \prod_{i \in I} P_i^{h_i} \in \mathbb{F}_2[x]$ be such that each P_i is a Mersenne prime and $a, b, h_i \in \mathbb{N}^*$. Then A is perfect (resp. unitary perfect) if and only if $A \in \mathcal{P}$ (resp. $A = B^{2^n}$ for some $n \in \mathbb{N}$ and $B \in \mathcal{P}_u$).

Theorem 1.2. Let $h \in \mathbb{N}^*$ and let $M \in \mathbb{F}_2[x]$ be a Mersenne prime. Then in the following cases, $\sigma(M^{2h})$ is divisible by a non-Mersenne prime: (i) $(M \in \{M_1, M_3, \overline{M_3}\})$ or $(M \in \{M_2, \overline{M_2}\}$ and $h \ge 2$). (ii) $M \notin \mathcal{M}$ and 2h + 1 is divisible by a prime number $p \in \Delta \setminus \{7\}$.

2 Proof of Theorem 1.1

Sufficiencies are obtained by direct computations. For the necessities, we shall apply Lemmas 2.4 and 2.15, Propositions 2.9 and 2.16. We fix:

$$A = x^{a}(x+1)^{b} \prod_{i \in I} P_{i}^{h_{i}} = A_{1}A_{2}, \text{ where } a, b, h_{i} \in \mathbb{N}, P_{i} \text{ is a Mersenne prime,}$$
$$A_{1} = x^{a}(x+1)^{b} \prod_{P_{i} \in \mathcal{M}} P_{i}^{h_{i}} \text{ and } A_{2} = \prod_{P_{j} \notin \mathcal{M}} P_{j}^{h_{j}}.$$

Lemma 2.1. If A is perfect (resp. unitary perfect), then $\sigma(x^a)$, $\sigma((x+1)^b)$ and $\sigma(P_i^{h_i})$, for any $i \in I$ (resp. $\sigma^*(x^a)$, $\sigma^*((x+1)^b)$, $\sigma^*(P_i^{h_i})$) are only divisible by x, x+1 or by Mersenne primes.

Proof. Since σ and σ^* are multiplicative, $\sigma(A) = \sigma(x^a)\sigma((x+1)^b) \prod_{i \in I} \sigma(P_i^{h_i})$ (resp. $\sigma^*(A) = \sigma^*(x^a)\sigma^*((x+1)^b) \prod_{i \in I} \sigma^*(P_i^{h_i})$). Any divisor of $\sigma(x^a)$, $\sigma((x+1)^b)$ and $\sigma(P_i^{h_i})$ (resp. of $\sigma^*(x^a)$, $\sigma^*((x+1)^b)$, $\sigma^*(P_i^{h_i})$) divides $\sigma(A) = A$ (resp. $\sigma^*(A) = A$).

Lemma 2.2 (Lemma 2 in [4]). A polynomial S is (unitary) perfect if and only if for any irreducible polynomial P and for any $m_1, m_2 \in \mathbb{N}^*$, we have $(P^{m_1} || S, P^{m_2} || \sigma(S)) \Rightarrow m_1 = m_2$ (resp. $(P^{m_1} || S, P^{m_2} || \sigma^*(S)) \Rightarrow m_1 = m_2$).

Examples 2.3 (useful for Propositions 2.9 and 2.16).

(i) The polynomial $S_1 = x^{13}(x+1)^2 M_1^3 M_2^2 \overline{M_2}^2 M_3 \overline{M_3}$ is not perfect because $x^{13} \|S_1$ and $x^7 \|\sigma(S_1)$.

(ii) The polynomial $S_2 = x^{14}(x+1)^7 M_1^2 M_2^3 \overline{M_2}^3 M_3 \overline{M_3}$ is not unitary perfect since $x^{14} ||S_2|$ and $x^{10} ||\sigma^*(S_2)$.

2.1 Case of perfect polynomials

Lemma 2.4 (Theorem 1.1 in [8]). If $h_i = 2^{n_i} - 1$ for any $i \in I$, then $A \in \mathcal{P}$.

We get from Theorem 8 in [5] and from Theorem 1.2.

Lemma 2.5. (i) If $h \in \mathbb{N}^*$ and if $\sigma(x^{2h})$ is only divisible by Mersenne primes, then $2h \in \{2, 4, 6\}$ and all its divisors lie in \mathcal{M} . More precisely, $\sigma(x^2) = M_1 = \sigma((x+1)^2), \ \sigma(x^4) = M_3, \ \sigma((x+1)^4) = \overline{M_3} \text{ and } \sigma(x^6) = M_2\overline{M_2} = \sigma((x+1)^6).$ (ii) Let $M \in \mathcal{M}$ and $h \in \mathbb{N}^*$ be such that $\sigma(M^{2h})$ is only divisible by Mersenne primes, then $2h = 2, M \in \{M_2, \overline{M_2}\}$ and $\sigma(M^2) \in \{M_1M_3, M_1\overline{M_3}\}.$

We dress from Lemma 2.5, the following tables of all the forms of a, b, P_i and h_i which satisfy Lemma 2.1, if $P_i \in \mathcal{M}$ and if $h_i \neq 2^{n_i} - 1$.

a		$\sigma(x^a)$	$\sigma(x^a)$		b		$\sigma((x+1)^b)$
$3 \cdot 2^n - 1$		$(x+1)^{2^n-1}M_1^{2^n}$			$3 \cdot 2^m - 1$		$x^{2^m-1}M_1^{2^m}$
$5 \cdot 2^n - 1$		$(x+1)^{2^n-1}M_3^{2^n}$			$5 \cdot 2^m - 1$		$x^{2^m-1}\overline{M_3}^{2^m}$
$7 \cdot 2^n - 1$		$(x+1)^{2^n-1}M_2^{2^n} \overline{M_2}^{2^n}$			$7 \cdot 2^m - 1$		$x^{2^m-1}M_2^{2^m} \overline{M_2}^{2^m}$
P_i	$P_i h_i$		$\sigma(P_i^{h_i})$				
M_2	$M_2 3 \cdot 2^{n_i} - 1$		$(1+M_2)^{2^{n_i}-1}M_1^{2^{n_i}}\overline{M_3}^{2^{n_i}}$				
			$(1+\overline{M_2})^{2^{n_i}-1}M_1^{2^n}$				

Corollary 2.6. Suppose that A_1 is perfect. Then, neither M_2 nor $\overline{M_2}$ divides $\sigma(P_i^{h_i})$ if $P_i \in \mathcal{M}$. Moreover, $\overline{M_2}$ divides A_1 whenever M_2 divides A_1 and their exponents (in A_1) are equal.

Proof. The first statement follows from Lemma 2.5-(ii). Now, if M_2 divides $A_1 = \sigma(A_1)$, then M_2 divides $\sigma(x^a) \sigma((x+1)^b) \prod_{P_i \in \mathcal{M}} \sigma(P_i^{h_i})$. Hence, M_2

divides $\sigma(x^a)\sigma((x+1)^b)$. The above tables show that a or b is of the form $7 \cdot 2^n - 1$, where $n \in \mathbb{N}$. So, $\overline{M_2}$ divides $\sigma(A_1) = A_1$. It suffices to consider two cases. If $a = 7 \cdot 2^n - 1$ and $b = 7 \cdot 2^m - 1$, then $M_2^{\ell} ||A_1|$ and $\overline{M_2}^{\ell} ||A_1|$,

with $\ell = 2^n + 2^m$. If $a = 7 \cdot 2^n - 1$ and $(b = 3 \cdot 2^m - 1 \text{ or } b = 5 \cdot 2^m - 1)$, then $M_2^{\ell} \| A_1$ and $\overline{M_2}^{\ell} \| A_1$, with $\ell = 2^n$.

Lemma 2.7.

If P is a Mersenne prime divisor of $\sigma(A_1)$, then $P, \overline{P} \in \{M_1, M_2, M_3\}$.

Proof. One has: $\sigma(A_1) = \sigma(x^a)\sigma((x+1)^b) \prod_{P_i \in \mathcal{M}} \sigma(P_i^{h_i})$. If P divides $\sigma(x^a)\sigma((x+1)^b)$

1)^b), then $P \in \mathcal{M}$ by Lemma 2.5-(i). If P divides $\sigma(P_i^{h_i})$ with $P_i \in \mathcal{M}$, then $P_i \in \{M_2, \overline{M_2}\}$, $(h_i = 2 \text{ or } h_i \text{ is of the form } 3 \cdot 2^{n_i} - 1)$ and $P, \overline{P} \in \{M_1, M_3\}$ (see the above tables).

Lemma 2.8. If A is perfect, then $A = A_1$.

Proof. We claim that $A_2 = 1$. Let $P_j \notin \mathcal{M}$ and $Q_i \in \mathcal{M}$. Then, P_j divides neither $\sigma(x^a)$, $\sigma((x+1)^b)$ nor $\sigma(Q_i^{h_i})$. Thus $gcd(P_j^{h_j}, \sigma(A_1)) = 1$. Observe that $P_j^{h_j}$ divides $\sigma(A_2)$ because $P_j^{h_j}$ divides $A = \sigma(A) = \sigma(A_1)\sigma(A_2)$. Hence, A_2 divides $\sigma(A_2)$. So, A_2 is perfect and it is equal to 1, A being indecomposable.

Proposition 2.9. If A_1 is perfect, then $h_j = 2^{n_j} - 1$ for any $P_j \in \mathcal{M}$.

Proof. We refer to Tables at the beginning of this section.

(i) Suppose that $P_j \notin \{M_2, \overline{M_2}\}$. If h_j is even, then $\sigma(P_j^{h_j})$ is divisible by a non-Mersenne prime. It contradicts Lemma 2.1. If $h_j = 2^{n_j}u_j - 1$ with $u_j \geq 3$ odd, then $\sigma(P_j^{h_j}) = (1+P_j)^{2^{n_j}-1} \cdot (1+P_j + \cdots + P_j^{u_j-1})^{2^{n_j}}$. Since $1+P_j + \cdots + P_j^{u_j-1} = \sigma(P_j^{u_j-1})$ is divisible by a non-Mersenne prime, we also get a contradiction to Lemma 2.1.

(ii) If $P_j \in \{M_2, \overline{M_2}\}$ and $(h_j \text{ is even or it is of the form } 2^{n_j}u_j-1$, with $u_j \geq 3$ odd and $n_j \geq 1$), then Corollary 2.6 implies that there exists $\ell \in \mathbb{N}^*$ such that $M_2^{\ell} ||A_1 \text{ and } \overline{M_2}^{\ell}||A_1$. Recall that $\sigma(M_2^2) = M_1\overline{M_3}$ and $\sigma(\overline{M_2}^2) = M_1M_3$. We proceed as in the proof of Corollary 2.6. It suffices to distinguish four cases which give contradictions.

• Case 1: $a = 7 \cdot 2^n - 1$ and $b = 7 \cdot 2^m - 1$

One has $\ell = 2^n + 2^m$ and neither M_1 nor M_3 divides $\sigma(x^a) \sigma((x+1)^b)$. If h_j is even, then $h_j = 2 = \ell$. So, n = m = 0, $M_1^2 || \sigma(A_1) = A_1$. It contradicts the part (i) of our proof.

If $h_j = 2^{n_j} u_j - 1$ with $u_j \ge 3$ odd and $n_j \ge 1$, then $u_j = 3$ and $M_1^{2 \cdot 2^{n_j}} || A_1$. • Case 2: $a = 7 \cdot 2^n - 1$ and $b = 5 \cdot 2^m - 1$

One has $\ell = 2^n$ and $M_1 \nmid \sigma(x^a) \sigma((x+1)^b)$. If h_j is even, then $2^n = \ell = h_j =$

2. So, n = 1 and $M_1^2 ||A_1$. If $h_j = 2^{n_j} u_j - 1$, with $u_j \ge 3$ odd and $n_j \ge 1$, then $u_j = 3$ and $2^n = \ell = h_j = 3 \cdot 2^{n_j} - 1$. It is impossible.

• Case 3: $a = 7 \cdot 2^n - 1$, $b = 3 \cdot 2^m - 1$ and h_j is even As above, $2^n = \ell = h_j = 2$, $M_1^{2^m}$ divides $\sigma((x+1)^b)$ and $M_1^{2^n+2^m}$ divides $\sigma(A_1) = A_1$. So, n = 1 and $M_1^{2^m+2} ||A_1$. Thus, the part (i) implies that m = 0. Hence, $A_1 = S_1 = x^{13}(x+1)^2 M_1^3 M_2^2 \overline{M_2}^2 M_3 \overline{M_3}$ which is not perfect (see Examples 2.3).

• Case 4: $a = 7 \cdot 2^n - 1$, $b = 3 \cdot 2^m - 1$, $h_j = 2^{n_j} u_j - 1$, $u_j \ge 3$ odd, $n_j \ge 1$ One has $u_j = 3$ and $2^n = \ell = h_j = 3 \cdot 2^{n_j} - 1$. It is impossible.

Lemma 2.8, Proposition 2.9 and Lemma 2.4 imply

Corollary 2.10. If A is perfect, then $A = A_1 \in \mathcal{P}$.

2.2Case of unitary perfect (u.p) polynomials

Similar arguments give Proposition 2.16 which finishes our proof.

Lemma 2.11. Let $S \in \mathbb{F}_2[x]$ be an irreducible polynomial. Then, for any $n, u \in \mathbb{N}$ with u odd, $\sigma^*(S^{2^n u}) = (1+S)^{2^n} (\sigma(S^{u-1}))^{2^n}$.

We may dress the following tables, from Lemmas 2.1, 2.5 and 2.11.

a	$\sigma^*(x^{\prime})$	^a)		b	$\sigma^*((x+1)^b)$
$3 \cdot 2^n$	(x +	$(x+1)^{2^n} M_1^{2^n}$			$x^{2^m} M_1^{2^m}$
$5 \cdot 2^n$	(x +	$(x+1)^{2^n} M_3^{2^n}$			$x^{2^m}\overline{M_3}^{2^m}$
		$(x+1)^{2^n} M_2^{2^n} \overline{M_2}^{2^n}$			$x^{2^m} M_2^{2^m} \overline{M_2}^{2^m}$
	h_i	$\sigma^*(P_i^{h_i})$			
	-	$(1+M_2)^{2^{n_i}}M_1^{2^{n_i}}\overline{M_3}^{2^{n_i}}$			
		$(1+\overline{M_2})^{2^{n_i}}M_1^{2^{n_i}}M_3^{2^{n_i}}$			

Lemma 2.12. Let $C \in \mathbb{F}_2[x] \setminus \{1\}$ be u.p. Then C is even, \overline{C} and C^{2^r} are also u.p, for any $r \in \mathbb{N}$.

Proof. If D is a divisor of C, then \overline{D} divides \overline{C} and D^{2^r} divides C^{2^r} . Thus, $\sigma^*(\overline{C}) = \overline{\sigma^*(C)} = \overline{C}$ and $\sigma^*(C^{2^r}) = (\sigma^*(C))^{2^r} = C^{2^r}$.

It remains to prove that C is even. Consider an irreducible divisor P of Cand $k \in \mathbb{N}^*$ such that $P^k \| C$. The polynomial 1 + P is even and divides $1 + P^k = \sigma^*(P^k)$. So, 1 + P divides $\sigma^*(C) = C$.

Definition 2.13. We denote by ~ the relation on $\mathbb{F}_2[x]$ defined as: $S \sim T$ if there exists $\ell \in \mathbb{Z}$ such that $S = T^{2^{\ell}}$.

Lemma 2.14. ([3], Section 2)

The relation \sim is an equivalence relation on $\mathbb{F}_2[x]$. Each equivalence class contains a unique polynomial B which is not a square, with $val_x(B) \leq$ $val_{x+1}(B).$

Lemma 2.15 (Theorem 1.3 in [8]). If $h_i = 2^{n_i}$ for any $i \in I$, then A (or \overline{A}) is of the form B^{2^n} , where $B \in \mathcal{P}_u$.

Proposition 2.16. (i) If A is u.p., then $A = A_1$. (ii) If A_1 is u.p., then $h_j = 2^{n_j}$ for any $P_j \in \mathcal{M}$. (iii) If A is u.p, then A or \overline{A} is of the form B^{2^n} , where $B \in \mathcal{P}_u$.

Proof. The proof of (i) is analogous to that of Lemma 2.8. The statement (iii) follows from (i), (ii) and Lemma 2.15. We only sketch the proof of (ii). Set $h_j = 2^{n_j} u_j$, where u_j is odd and $n_j \ge 0$.

- Suppose that $P_j \notin \{M_2, \overline{M_2}\}$. If $u_j \ge 3$, then $\sigma(P_j^{u_j-1})$ and thus $\sigma^*(P_j^{h_j})$ are divisible by a non-Mersenne prime. It contradicts Lemma 2.1.

- If $P_j \in \{M_2, \overline{M_2}\}$ and if $u_j \ge 3$, then $u_j = 3$ and (a or b is of the form $7 \cdot 2^n$). Recall that $\sigma^*(M_2^3) = (1+M_2)M_1\overline{M_3}$ and $\sigma^*(\overline{M_2}^3) = (1+\overline{M_2})M_1M_3$. We consider two cases. The first gives non unitary perfect polynomials whereas the second leads to a contradiction.

• Case 1: $a = 7 \cdot 2^n$ and $b = 7 \cdot 2^m$, with $n, m \ge 0$

One has $M_2^{\ell} \| A_1$ and $\overline{M_2}^{\ell} \| A_1$, with $\ell = 2^n + 2^m$. Neither M_1 nor M_3 divides $\sigma(x^a) \sigma((x+1)^b)$.

Thus, $3 \cdot 2^{n_j} = h_j = \ell = 2^n + 2^m$. So, $(n = m + 1 \text{ and } n_j = m)$ or $(m = n + 1 \text{ and } n_j = n)$. Therefore, $(M_1^2)^{2^{n_j}}$, $M_3^{2^{n_j}}$ and $\overline{M_3}^{2^{n_j}}$ divide $\sigma^*(M_2^{h_j})\sigma^*(\overline{M_2}^{h_j})$ and they divide $\sigma^*(A_1) = A_1$. Thus, $A_1 = S_2^{2^m}$ or $A_1 = \overline{S_2}^{2^n}$ where $S_2 = x^{14}(x+1)^7 M_1^2 M_2^3 \overline{M_2}^3 M_3 \overline{M_3}$. In both cases, A_1 is not unitary perfect because S_2 is not u.p (Examples 2.3).

• Case 2: $a = 7 \cdot 2^n$ and $(b = 5 \cdot 2^m \text{ or } b = 3 \cdot 2^m)$, with $n, m \ge 0$ One has $\ell = 2^n$. So, we get the contradiction: $3 \cdot 2^{n_j} = h_j = \ell = 2^n$.

3 Proof of Theorem 1.2

We mainly prove Theorem 1.2 by contradiction (to Corollary 3.5). Lemma 3.1 states that $\sigma(M^{2h})$ is square-free, for any $h \in \mathbb{N}^*$.

We set $M = x^{a}(x+1)^{b} + 1$, $U_{2h} = \sigma(\sigma(M^{2h}))$ and

$$\sigma(M^{2h}) = \prod_{j \in J} P_j, P_j = 1 + x^{a_j} (x+1)^{b_j} \text{ irreducible, } P_i \neq P_j \text{ if } i \neq j.$$
(1)

By Lemma 3.3, if there exists a prime divisor p of 2h + 1 such that $\sigma(M^{p-1})$ is divisible by a non-Mersenne prime, then $\sigma(M^{2h})$ is also divisible by a non-Mersenne. Therefore, it suffices to consider that 2h + 1 = p is a prime number, except for p = 3 with $M \in \{M_2, \overline{M_2}\}$ (see Section 3.3).

3.1 Useful facts

For $S \in \mathbb{F}_2[x] \setminus \{0, 1\}$, of degree s, we denote by $\alpha_l(S)$ the coefficient of x^{s-l} in S, $0 \leq l \leq s$. One has: $\alpha_0(S) = 1$.

Lemma 3.1 (Lemmas 4.6 and 4.8 in [10]). The polynomial $\sigma(M^{2h})$ is square-free and $M \neq M_1$.

Lemma 3.2 (Theorem 1.4 in [10]). Let $h \in \mathbb{N}^*$ be such that p = 2h + 1is prime and let M be a Mersenne prime such that $M \notin \{M_2, \overline{M_2}\}$ and $\omega(\sigma(M^{2h})) = 2$. Then, $\sigma(M^{2h})$ is divisible by a non-Mersenne prime.

The lemma below generalizes Lemma 4.10 in [10] (with an analogous proof).

Lemma 3.3. If k is a divisor (prime or not) of 2h+1, then $\sigma(M^{k-1})$ divides $\sigma(M^{2h})$.

We sometimes apply Lemmas 3.4 and 3.6 without explicit mentions.

Lemma 3.4. Let $S \in \mathbb{F}_2[x]$ be such that $s = \deg(S) \ge 1$ and $l, t, r, r_1, \ldots, r_k \in \mathbb{N}$ be such that $r_1 > \cdots > r_k$, $t \le k, r_1 - r_t \le l \le r \le s$. Then (i) $\alpha_l[(x^{r_1} + \cdots + x^{r_k})S] = \alpha_l(S) + \alpha_{l-(r_1-r_2)}(S) + \cdots + \alpha_{l-(r_1-r_t)}(S)$. (ii) $\alpha_l(\sigma(S)) = \alpha_l(S)$ if any divisor of S has degree at least r + 1.

Proof. The equality in (i) (resp. in (ii)) follows from the definition of α_l (resp. from the fact: $\sigma(S) = S + T$, where $\deg(T) \leq \deg(S) - r - 1$).

Corollary 3.5. (i) The integers $u = \sum_{j \in J} a_j$ and $v = \sum_{j \in J} b_j$ are both even. (ii) The polynomial U_{2h} splits (over \mathbb{F}_2) and it is a square.

(iii) The polynomial $\sigma(M^{2h})$ is reducible.

Proof. (i) See [10, Corollary 4.9]. For (ii), Assumption (1) implies that $U_{2h} = \sigma(\sigma(M^{2h})) = \sigma(\prod_{j \in J} P_j) = \prod_{j \in J} x^{a_j} (x+1)^{b_j} = x^u (x+1)^v$, where u and

 \boldsymbol{v} are both even.

(iii) If $\sigma(M^{2h}) = Q$ is irreducible, then $U_{2h} = 1 + Q$ is not a square. \Box

Lemma 3.6. One has $\alpha_l(\sigma(M^{2h})) = \alpha_l(M^{2h})$ if $l \le a+b-1$ and $\alpha_l(\sigma(M^{2h})) = \alpha_l(M^{2h} + M^{2h-1})$ if $a + b \le l \le 2(a+b) - 1$.

Proof. Since $\sigma(M^{2h}) = M^{2h} + M^{2h-1} + T$, with deg(*T*) ≤ (a+b)(2h-2) = 2h(a+b) - 2(a+b), Lemma 3.4-(ii) implies that $\alpha_l(\sigma(M^{2h})) = \alpha_l(M^{2h})$ if $l \le a+b-1$ and $\alpha_l(\sigma(M^{2h})) = \alpha_l(M^{2h}+M^{2h-1})$ if $a+b \le l \le 2(a+b)-1$. □

Lemma 3.7. Denote by $N_2(m)$ the number of irreducible polynomials over \mathbb{F}_2 , of degree $m \geq 1$. Then

(i) $N_2(m) \ge \frac{2^m - 2(2^{m/2} - 1)}{m}$,

(ii) $\varphi(m) < N_2(m)$ if $m \ge 4$, where φ is the Euler totient function, (iii) For each $m \ge 4$, there exists an irreducible polynomial of degree m, which is not a Mersenne prime.

Proof. (i) See Exercise 3.27, p. 142 in [11]. (ii) If $m \in \{4,5\}$, then direct computations give $\varphi(4) = 2$, $N_2(4) = 3$ and $\varphi(5) = 4$, $N_2(5) = 6$.

Now, suppose that $m \ge 6$. Consider the function $f(x) = 2^x - 2(2^{x/2} - 1) - x^2$, for $x \ge 6$. The derivative of f is a positive function. So, $f(x) \ge f(6) > 0$ and $x < \frac{2^x - 2(2^{x/2} - 1)}{x}$. Thus, $\varphi(m) \le m < \frac{2^m - 2(2^{m/2} - 1)}{m} \le N_2(m)$. (iii) We remark that if $1 + x^c(x+1)^d$ is a Mersenne prime, then gcd(c, d) = 1.

So, gcd(c, c + d) = 1. Therefore, the set \mathcal{M}_m of Mersenne primes of degree m is a subset of $\{x^c(x+1)^{m-c}+1: 1 \leq c \leq m, gcd(c,m)=1\}$. Thus,

$$#\mathcal{M}_m \le \#\{c: 1 \le c \le m, \ \gcd(c,m) = 1\} = \varphi(m).$$

Hence, there exist at least $N_2(m) - \varphi(m)$ irreducible non-Mersenne polynomials, with $N_2(m) - \varphi(m) \ge 1$, by (ii).

Lemma 3.8. For any $j \in J$, $ord_p(2)$ divides $a_j + b_j = deg(P_j)$.

Proof. Set $d = \gcd_{i \in J}(a_i + b_i)$. By Lemma 4.13 in [10], p divides $2^d - 1$. Thus, $ord_p(2)$ divides d.

Lemma 3.9. ([11], Chap. 2 and 3)

Let $q = 2^r - 1$ be a Mersenne prime number. Then, any irreducible polynomial P of degree r is primitive. In particular, each root β of P is a primitive element of the field \mathbb{F}_{2^r} , so that β is of order q in $\mathbb{F}_{2^r} \setminus \{0\}$.

Lemma 3.10. Let $P_i = 1 + x^{a_i}(x+1)^{b_i}$ be a prime divisor of $\sigma(M^{p-1})$, where $2^{a_i+b_i} - 1 = p_i$ is a prime number. Then, $p_i = p$ and $\sigma(M^{p-1})$ is divisible by any irreducible polynomial of degree $a_i + b_i$. Furthermore, at least one of those divisors is not a Mersenne prime if $a_i + b_i \ge 4$.

Proof. The polynomial P_i is primitive. If α is a root of P_i , then $(M^p + 1)(\alpha) = 0$ and $M(\alpha) = \alpha^r$ for some $1 \le r \le p_i - 1$. Thus, $1 = M(\alpha)^p = \alpha^{rp}$, with $ord(\alpha) = p_i$. So, p_i divides rp and $p_i = p$.

Any irreducible polynomial S of degree $a_i + b_i$ is primitive. Let β be a root of S. One has $ord(\beta) = p_i = p$, $S(\beta) = 0$ and $M(\beta) = \beta^s$, for some $1 \le s \le p_i - 1$. Thus, $M(\beta)^p = \beta^{ps} = 1$ and S divides $M^p + 1 = x^a(x+1)^b \sigma(M^{p-1})$. The third statement follows from Lemma 3.7-(iii).

Corollary 3.11. For any $i \in J$, $a_i + b_i \leq 3$ or $2^{a_i + b_i} - 1$ is not prime.

Lemma 3.12. Let $P, Q \in \mathbb{F}_2[x]$ be such that $\deg(P) = r, 2^r - 1$ is prime, $P \nmid Q(Q+1)$ but $P \mid Q^p + 1$. Then $2^r - 1 = p$.

Proof. The polynomial P is primitive. If β is a root of P, then $ord(\beta) = 2^r - 1$. Moreover, $Q(\beta) \notin \{0,1\}$ because $P \nmid Q(Q+1)$. Thus, $Q(\beta) = \beta^t$ for some $1 \le t \le 2^r - 2$. Hence, $1 = Q(\beta)^p = \beta^{tp}$. So, $2^r - 1$ divides tp and $2^r - 1 = p$.

Corollary 3.13. Let $r \in \mathbb{N}^*$ be such that $2^r - 1$ is a prime distinct from p. Then, no irreducible polynomial of degree r divides $\sigma(M^{p-1})$.

Proof. If P is a prime divisor of $\sigma(M^{p-1})$ with deg(P) = r, then P divides M^p+1 and by taking Q = M in the above lemma, we get a contradiction. \Box

In the following lemma and two corollaries, we suppose that p is a Mersenne prime of the form $2^m - 1$ (with m prime).

Lemma 3.14. Let $P, Q \in \mathbb{F}_2[x]$ be such that P is irreducible of degree m and $P \nmid Q(Q+1)$. Then, P divides $Q^p + 1$.

Proof. The polynomial P is primitive. If β is a root of P, then $ord(\beta) = 2^m - 1 = p$, $Q(\beta) \notin \{0, 1\}$ because $P \nmid Q(Q+1)$. Thus, $Q(\beta) = \beta^t$ for some $1 \le t \le p-1$. Hence, $Q(\beta)^p = \beta^{tp} = 1$. So, P divides $Q^p + 1$.

Corollary 3.15. Any irreducible polynomial $P \neq M$ (Mersenne or not), of degree m, divides $\sigma(M^{p-1})$.

Proof. We may apply Lemma 3.14 with Q = M because P does not divide $x^a(x+1)^b M = M(M+1) = Q(Q+1)$. So, P is odd and it divides $M^p + 1 = (M+1) \sigma(M^{p-1}) = x^a(x+1)^b \sigma(M^{p-1})$.

Corollary 3.16. The polynomial M_1 (resp. M_2 , $\overline{M_2}$) divides $\sigma(M^{p-1})$ if and only if $(M \neq M_1 \text{ and } p = 3)$ (resp. $M \neq M_2$ and p = 7, $M \neq \overline{M_2}$ and p = 7).

Proof. Apply Corollary 3.15 with $m \in \{2, 3\}$.

In order to carry on the proof (of Theorem 1.2), we distinguish three cases.

3.2 Case I: $M \in \{M_1, M_3, \overline{M_3}\}$

Lemma 3.1 implies that $M \neq M_1$. It suffices to suppose that $M = M_3$. We refer to Section 5.2 in [9]. Put $D = M_1 M_2 \overline{M_2}$. By [9, Lemma 5.4], we have to consider four situations:

(i) $\gcd(\sigma(M^{2h}), D) = 1$, (ii) $\sigma(M^{2h}) = M_1 B$, with $\gcd(B, D) = 1$, (iii) $\sigma(M^{2h}) = M_2 \overline{M_2} B$, with $\gcd(B, D) = 1$, (iv) $\sigma(M^{2h}) = DB$, with $\gcd(B, D) = 1$, where any irreducible divisor of B has degree exceeding 5.

The following lemma contradicts the fact that U_{2h} is a square.

Lemma 3.17. One has $\alpha_3(U_{2h}) = 1$ or $\alpha_5(U_{2h}) = 1$.

Proof. For (i), (iii) and (iv), use Lemmas 5.9, 5.10, 5.15, 5.17 (in [9]). (ii) Since $\sigma(M^{2h}) = (x^2 + x + 1)B$ and $U_{2h} = (x^2 + x)\sigma(B)$, we obtain (by Lemmas 3.4 and 3.6):

$$\begin{cases} 0 = \alpha_1(M^{2h}) = \alpha_1(\sigma(M^{2h})) = \alpha_1(B) + 1, \\ \alpha_3(U_{2h}) = \alpha_3(\sigma(B)) + \alpha_2(\sigma(B)) = \alpha_3(B) + \alpha_2(B), \\ 0 = \alpha_3(M^{2h}) = \alpha_3(\sigma(M^{2h})) = \alpha_3(B) + \alpha_2(B) + \alpha_1(B). \end{cases}$$

Thus, $\alpha_3(U_{2h}) = \alpha_3(B) + \alpha_2(B) = \alpha_1(B) = 1.$

3.3 Case II: $M \in \{M_2, \overline{M_2}\}$ and $h \ge 2$

It suffices to consider that $M = M_2 = 1 + x + x^3$.

Lemma 3.18. (i) If $h \ge 4$, then M_1 (resp. $\overline{M_2}$) divides $\sigma(M^{2h})$ if and only if 3 divides 2h + 1 (resp. 7 divides 2h + 1).

(ii) If $h \ge 4$ and if 2h + 1 is divisible by a prime $p \notin \{3,7\}$, then any irreducible divisor of $\sigma(M^{2h})$ is of degree at least 4.

Proof. The assertion (ii) follows from (i) which in turn, follows from Corollaries 3.13 and 3.15.

We consider three possibilities since $\sigma(M^{p-1}) = \sigma(M_2^2) = M_1 \overline{M_3}$ (product of two Mersenne primes), if p = 3.

3.3.1 II-1: 2h + 1 is (divisible by) a prime $p \in \{5, 7\}$

Lemma 3.19. For $p \in \{5, 7\}$, some non-Mersenne prime divides $\sigma(M^{p-1})$.

Proof. Here, $h \in \{2,3\}$. By direct computations, $U_4 = x^3(x+1)^6(x^3+x+1)$ and $U_6 = x^8(x+1)^4(x^3+x+1)^2$ which do not split (despite that U_6 is a square).

3.3.2 II-2: $2h + 1 = 3^w$, for some $w \ge 2$

In this case, 9 divides 2h + 1 and $\sigma(M^8)$ divides $\sigma(M^{2h})$ (by Lemma 3.3). But, $\sigma(M^8) = (x^2 + x + 1)(x^4 + x^3 + 1)(x^6 + x + 1)(x^{12} + x^8 + x^7 + x^4 + 1)$, where $x^6 + x + 1 = 1 + x(x + 1)M_3$ is not a Mersenne prime.

3.3.3 II-3: 2h + 1 is (divisible by) a prime $p \notin \{3, 5, 7\}$

We may write p = 2h + 1 with $h \ge 4$.

Lemma 3.20. (i) If $l \in \{1, 2, 3\}$, then $\alpha_l(U_{2h}) = \alpha_l(\sigma(M^{2h}))$. (ii) If $l \in \{1, 2\}$, then $\alpha_l(\sigma(M^{2h})) = \alpha_l(M^{2h})$. (iii) The coefficients $\alpha_3(\sigma(M^{2h}))$ and $\alpha_3(M^{2h} + M^{2h-1})$ are equal.

Proof. (i) It follows from Lemma 3.18. For $l \leq 2$, $6h - l = \deg(\sigma(M^{2h})) - l = \deg((M^{2h}) - l > 3(2h - 1)) = \deg(M^{2h-1})$ and for $3 \leq l \leq 5$, $6h - l > 3(2h - 2) = \deg(M^{2h-2})$. Hence, we get (ii) and (iii). □

Corollary 3.21. The coefficient $\alpha_3(U_{2h})$ equals 1.

Proof. The previous lemma implies that $\alpha_3(U_{2h}) = \alpha_3(M^{2h} + M^{2h-1}) = \alpha_3[(x^3 + x)M^{2h-1}] = \alpha_3(M^{2h-1}) + \alpha_1(M^{2h-1}).$ But, $M^{2h-1} = (x^3 + x + 1)^{2h-1} = (x^3 + x)^{2h-1} + (x^3 + x)^{2h-2} + \cdots$ The coefficient of x^{6h-6} (resp. of x^{6h-4}) in M^{2h-1} is exactly $\alpha_3(M^{2h-1})$ (resp. $\alpha_1(M^{2h-1})$). So, $\alpha_3(M^{2h-1}) = 1$ and $\alpha_1(M^{2h-1}) = 0.$

3.4 Case III: $M \notin \mathcal{M}$

Here, we have two possibilities.

3.4.1 III-1: the prime p is such that $ord_p(2) \equiv 0 \mod 8$

Lemmas 3.22 and 3.8 imply Corollary 3.23.

Lemma 3.22. There exists no Mersenne prime of degree multiple of 8.

Proof. If $Q = 1 + x^{c_1}(x+1)^{c_2}$ with $c_1 + c_2 = 8k$, then $\omega(Q)$ is even by [10, Corollary 3.3]. So, Q is reducible.

Corollary 3.23. If $ord_p(2) \equiv 0 \mod 8$, then $\sigma(M^{2h})$ is divisible by a non-Mersenne prime.

Proof. Suppose that $\sigma(M^{2h}) = \prod_{j \in J} P_j$, where each P_j is a Mersenne prime.

Then, Lemma 3.8 implies that $ord_p(2)$ divides $deg(P_j)$, for any $j \in J$. So, 8 divides $deg(P_j)$. It contradicts Lemma 3.22.

3.4.2 III-2: *p* is a Mersenne prime number with $p \neq 7$

Set $p = 2^m - 1$, with *m* and *p* are both prime. Note that there are (at present) 51 known Mersenne prime numbers (OEIS Sequences A000043 and A000668). The first five of them are: 3,7,31,127 and 8191.

Lemma 3.24. If $p \ge 31$ is a Mersenne prime number, then $\sigma(M^{p-1})$ is divisible by a non-Mersenne prime.

Proof. Here, $a + b = \deg(M) \ge 5$ since $M \notin \mathcal{M}$. We get our result from Corollary 3.15 and Lemma 3.7-(iii).

It remains then the case p = 3 (since $p \neq 7$, in this section). Lemma 3.2 has already treated the case where $\omega(\sigma(M^2)) = 2$. So, we suppose that $\omega(\sigma(M^2)) \geq 3$. Put $\sigma(M^2) = M_1 \cdots M_r$, $r \geq 3$ and $U_2 = \sigma(\sigma(M^2))$. We shall prove that $\alpha_3(U_2) = 1$ (Corollary 3.29), a contradiction to the fact that U_2 is a square. Corollary 3.16 gives

Lemma 3.25. (i) The trinomial $1 + x + x^2$ divides $\sigma(M^2)$. (ii) No irreducible polynomial of degree $r \ge 3$ such that $2^r - 1$ is prime, divides $\sigma(M^2)$.

Corollary 3.26. The polynomial $\sigma(M^2)$ is of the form $(1+x+x^2)B$, where $gcd(1+x+x^2, B) = 1$ and any prime divisor of B has degree at least 4.

Lemma 3.27. If $\sigma(M^2) = (1 + x + x^2)B$ with $gcd(1 + x + x^2, B) = 1$, then

$$\begin{cases} (i) \ \alpha_1(\sigma(M^2)) = \alpha_1(B) + 1, \ \alpha_2(\sigma(M^2)) = \alpha_2(B) + \alpha_1(B) + 1, \\ (ii) \ \alpha_3(\sigma(M^2)) = \alpha_3(B) + \alpha_2(B) + \alpha_1(B), \\ (iii) \ \alpha_3(\sigma(M^2)) = 0. \end{cases}$$

Proof. We directly get (i) and (ii). For (iii), $\sigma(M^2) = 1 + M + M^2 = x^{2a}(x+1)^{2b} + x^a(x+1)^b + 1$. Moreover, 2a+2b-3 > a+b because $a+b \ge 4$ and $x^{2a}(x+1)^{2b}$ is a square. So, $\alpha_3(\sigma(M^2)) = \alpha_3(x^{2a}(x+1)^{2b}) = 0$.

Lemma 3.28. Some coefficients of U_2 and B satisfy:

$$\alpha_1(U_2) = \alpha_1(B) + 1, \ \alpha_2(U_2) = \alpha_2(B) + \alpha_1(B), \ \alpha_3(U_2) = \alpha_3(B) + \alpha_2(B).$$

Proof. Corollary 3.26 implies that $U_2 = \sigma(\sigma(M^2)) = \sigma((1 + x + x^2)B) = \sigma(1 + x + x^2)\sigma(B) = (x^2 + x)\sigma(B)$. Any irreducible divisor of B has degree more than 3. Hence, $\alpha_l(\sigma(B)) = \alpha_l(B)$, for $1 \le l \le 3$.

One gets:
$$\begin{cases} \alpha_1(U_2) = \alpha_1(\sigma(B)) + 1 = \alpha_1(B) + 1, \\ \alpha_2(U_2) = \alpha_2(\sigma(B)) + \alpha_1(\sigma(B)) = \alpha_2(B) + \alpha_1(B), \\ \alpha_3(U_2) = \alpha_3(\sigma(B)) + \alpha_2(\sigma(B)) = \alpha_3(B) + \alpha_2(B). \end{cases} \square$$

Corollary 3.29. The coefficient $\alpha_3(U_2)$ equals 1.

Proof. The polynomial U_2 is a square, so $0 = \alpha_1(U_2) = \alpha_1(B) + 1$ and thus $\alpha_1(B) = 1$. Lemma 3.27-(iii) implies that $0 = \alpha_3(\sigma(M^2)) = \alpha_3(B) + \alpha_2(B) + \alpha_1(B)$. Therefore, $\alpha_3(U_2) = \alpha_3(B) + \alpha_2(B) = \alpha_1(B) = 1$.

Remark 3.30. Our method fails for p = 7. Indeed, for many M, one has $\alpha_3(U_6) = \alpha_5(U_6) = 0$. So, we do not reach a contradiction. We should find a large enough odd integer l such that, $\alpha_l(U_6) = 0$. But, this does not appear always possible.

References

- J. T. B. BEARD JR, Perfect polynomials revisited, Publ. Math. Debrecen 38/(1-2) (1991), 5–12.
- [2] J. T. B. BEARD JR, Unitary perfect polynomials over GF(q), Rend. Accad. Lincei 62 (1977), 417–422.
- [3] J. T. B. BEARD JR, A. T. BULLOCK, M. S. HARBIN, Infinitely many perfect and unitary perfect polynomials, Rend. Accad. Lincei 63 (1977), 294–303.
- [4] J. T. B. BEARD JR, J. R. OCONNELL JR, K. I. WEST, Perfect polynomials over GF(q), Rend. Accad. Lincei 62 (1977), 283–291.
- [5] E. F. CANADAY, The sum of the divisors of a polynomial, Duke Math. J. 8 (1941), 721–737.
- [6] L. H. GALLARDO, O. RAHAVANDRAINY, Even perfect polynomials over F₂ with four prime factors, Intern. J. of Pure and Applied Math. 52(2) (2009), 301–314.
- [7] L. H. GALLARDO, O. RAHAVANDRAINY, All perfect polynomials with up to four prime factors over \mathbb{F}_4 , Math. Commun. 14(1) (2009), 47–65.
- [8] L. H. GALLARDO, O. RAHAVANDRAINY, On even (unitary) perfect polynomials over \mathbb{F}_2 , Finite Fields Appl. 18 (2012), 920–932.
- [9] L. H. GALLARDO, O. RAHAVANDRAINY, Characterization of Sporadic perfect polynomials over \mathbb{F}_2 , Functiones et Approx. 55(1) (2016), 7–21.
- [10] L. H. GALLARDO, O. RAHAVANDRAINY, On Mersenne polynomials over F₂, Finite Fields Appl. **59** (2019), 284–296.
- [11] R. LIDL, H. NIEDERREITER, Finite Fields, Encyclopedia of Mathematics and its applications, Cambridge University Press, 1983 (Reprinted 1987).
- [12] O. RAHAVANDRAINY, Familles de polynômes unitairement parfaits sur F₂, C. R. Math. Acad. Sci. Paris 359(2) (2021), 123–130.