
SOME ALGEBRAIC ASPECTS OF COMBINED MATRICES

Primitivo B. Acosta-Humánez

Instituto de Matemática
 Universidad Autónoma de Santo Domingo
 Dominican Republic
 pacosta-humanez@uasd.edu.do

Máximo Santana

Instituto de Matemática
 Universidad Autónoma de Santo Domingo
 Dominican Republic
 msantana@uasd.edu.do

ABSTRACT

In this work, we study some algebraic aspects of combined matrices $\mathcal{C}(A)$, where the entries of A belongs to a number field K and A is a non singular matrix. In other words, A is a $n \times n$ belonging to the General Linear Group over K denoted by $GL_n(K)$. Also we analyze the case in which the matrix A belongs to algebraic subgroups of $GL_n(K)$ such as unimodular group, that is A^2 is a $n \times n$ belonging to the Special Linear Group denoted by $SL_n(K)$, Triangular groups, Diagonal groups among others. In particular, we analyze entirely the case $n = 2$ giving explicitly the diagonalization of $\mathcal{C}(A)$, that includes characteristic polynomials, eigenvalues and eigenfactors.

Keywords Characteristic Polynomials · Combined Matrices · General Linear Group · Special Linear Group

1 Introduction

Combined matrices is a topic that increase the interest of researchers in the last years. Recently, the second named author in collaboration with researchers from Spain obtained new results related with Combined matrices, see [7, 11] and references therein. Previous works relative to combined matrices are given in references [4, 5, 6, 9, 10]. The combined matrix of a nonsingular matrix A is defined as $\mathcal{C}(A) = A \circ A^{-T}$ where \circ is the Hadamard product of matrices. In particular, we are interested in the cases where the elements of the matrix A are rational numbers, that is, A is a $n \times n$ belonging to the *General Linear Group* over \mathbb{Q} denoted by $GL_n(\mathbb{Q})$. Also we analyze the case in which the determinant of A is ± 1 and the entries of A are integer numbers, i.e., A^2 is a $n \times n$ matrix belonging to *Special Linear Group* $SL_n(\mathbb{Z})$. We provide some results regarding eigenvalues, eigenvectors and Galois groups related to combined matrices, i.e., we are going to work with the algebraic properties of combined matrices $\mathcal{C}(A)$ that includes the set of eigenvalues and eigenvectors of $\mathcal{C}(A)$ and the Galois group of its characteristic polynomial. We recall that the trace of an square matrix A is denoted by $\text{Tr}(A)$ and its determinant is denoted by $\det A$. A matrix A is *singular* whether $\det A = 0$, otherwise it is called *nonsingular*. The canonical basis of \mathbb{R}^n is denoted by $\mathcal{B} = \{e_1, e_2, \dots, e_n\}$, where $e_1 = (1, 0, \dots, 0)$, $e_2 = (0, 1, \dots, 0)$, ..., $e_n = (0, 0, \dots, 1)$.

Now we set the theoretical background necessary to understand the rest of the paper.

Definition 1. Let A be an $n \times n$ square matrix. The characteristic polynomial of A , denoted by $p(\lambda)$, is the polynomial of degree at most n obtained as $p_A(\lambda) = \det(A - \lambda I_n)$, where I_n is the $n \times n$ identity matrix. The roots of the characteristic polynomials is the set of eigenvalues of A denoted by $\Lambda(A) := \{\lambda \in \mathbb{C} : p_A(\lambda) = 0\}$. The set of eigenvectors of A is denoted by $\mathcal{V}(A)$. Minors of A given by A_{ij} and Cofactors of A given by $(-1)^{i+j} A_{ij}$, where A_{ij} is the determinant of the submatrix of A obtained by deletion of the row i and column j .

According to [8] we give the following definitions.

Definition 2. Consider a set G and an operation (\cdot) . The pair (G, \cdot) is a group if the following is satisfied

1. Closure property: $\forall a, b \in G, a \cdot b \in G$
2. Associative property: $\forall a, b, c \in G, a \cdot (b \cdot c) = (a \cdot b) \cdot c$

3. Identity element: $\exists e \in G$ such that $\forall a \in G, a \cdot e = e \cdot a = a$
4. Inverse element: $\forall a \in G, \exists b \in G$ such that $a \cdot b = b \cdot a = e$

If (G, \cdot) satisfies the closure property we say the it is Magma. If (G, \cdot) is Magma and it satisfies the associative property, we say the it is a Semigroup. If (G, \cdot) is Semigroup and it satisfies Identity element property, we say that it is Monoid. If (G, \cdot) is Monoid and it satisfies the inverse element property, we say the it is Group. If (G, \cdot) is group and it satisfies the commutative property ($\forall a, b \in G, a \cdot b = b \cdot a$), we say it is Abelian Group. By S_n we denote the symmetric group of order $n!$.

Definition 3. An algebraic number field, also called number field, is a finite degree field extension of the field of rational numbers \mathbb{Q} , where the degree field extension means the dimension of the field as a vector space over \mathbb{Q} .

Definition 4. Consider the number fields $(K, +, \cdot)$ and $(L, +, \cdot)$. The application $f : K \rightarrow L$ is:

1. Homomorphism whether $f(a + b) = f(a) + f(b)$ and $f(a \cdot b) = f(a) \cdot f(b)$ for all $a, b \in K$.
2. Monomorphism whether f is Homomorphism and it is injective.
3. Epimorphism whether f is Homomorphism and it is surjective.
4. Isomorphism whether f is Monomorphism and Epimorphism.
5. Endomorphism whether f is Homomorphism and $K = L$.
6. Automorphism whether f is Isomorphism and Endomorphism.

Definition 5. Consider a polynomial $p \in K[x]$, where K is a field. Let L/K be the extension of L over K . The automorphism $\sigma : L/K \rightarrow L/K$ is called K -automorphism of L whether $\sigma|_K = e$, i.e., σ restricted to the elements of K is the identity group. The Galois group $\text{Gal}(L/K)$ is the group of K -automorphisms of L .

According to [2] the following are the definitions of Reversing of matrices adapted to $n \times n$ matrices.

Definition 6. Let A be a $n \times n$ matrix, $\delta_{i,j} = 0$ for all $i \neq j$ and $\delta_{i,i} = 1$. Reversing of A is given by:

$$\mathcal{R}(A) = M_{\mathcal{R}_c} A M_{\mathcal{R}_r}, \quad M_{\mathcal{R}_r} = [\delta_{i,n-j+1}]_{n \times n}, \quad M_{\mathcal{R}_c} = [\delta_{n-i+1,j}]_{n \times n}, \quad M_{\mathcal{R}_c} \cdot M_{\mathcal{R}_r} = M_{\mathcal{R}_r} \cdot M_{\mathcal{R}_c} = I_n. \quad (1)$$

According to the references [10] and [5], the following is the definition of combined matrices

Definition 7. Let A be a nonsingular matrix. The combined matrix of A is

$$\mathcal{C}(A) = A \circ A^{-T} = A \circ (A^{-1})^T, \quad (2)$$

where in general $B \circ C$ is the Hadamard (entrywise) product of B and C .

2 Results of combined matrices

In this section we consider results concerning algebraic aspects of combined matrices of $n \times n$ matrices.

2.1 Combined matrices over the General Linear Group

Here we consider combined matrices over the General Linear Group. We start with the following Lemma that involves Hadamard product and Reversing of matrices.

Lemma 1. Let $A, B \in \text{GL}_n(K)$, \circ be the Hadamard product \mathcal{R} be Reversing over $\text{GL}_n(K)$:

$$\mathcal{R} : (\text{GL}_n(K), \circ) \rightarrow (\text{GL}_n(K), \circ).$$

Then, Reversing is a group automorphism from $(\text{GL}_n(K), \circ)$ to $(\text{GL}_n(K), \circ)$.

Proof. We start proving that \mathcal{R} is homomorphism. Thus, $\mathcal{R}(A \circ B) = M_{\mathcal{R}_c}(A \circ B)M_{\mathcal{R}_r}$. By definition of Reversing it is satisfied $I_n = M_{\mathcal{R}_c} \cdot M_{\mathcal{R}_r} = M_{\mathcal{R}_r} \cdot M_{\mathcal{R}_c}$ we have $\mathcal{R}(A \circ B) = (M_{\mathcal{R}_c} \cdot M_{\mathcal{R}_r} \cdot M_{\mathcal{R}_c})(A \circ B)M_{\mathcal{R}_r} = (M_{\mathcal{R}_c} A \cdot M_{\mathcal{R}_r}) \circ (M_{\mathcal{R}_c} B \cdot M_{\mathcal{R}_r})$ and we conclude $\mathcal{R}(A \circ B) = \mathcal{R}(A) \circ \mathcal{R}(B)$. Now, we prove that \mathcal{R} is Monomorphism. Thus, assuming $\mathcal{R}(A) = \mathcal{R}(B)$ and applying again \mathcal{R} , we obtain $\mathcal{R}^2(A) = \mathcal{R}^2(B)$ and therefore $A = B$. We see that \mathcal{R} is Epimorphism because $\mathcal{R}(A) \in \text{GL}_n(K)$ for all $A \in \text{GL}_n(K)$. In this way we proved that \mathcal{R} is Isomorphism and because it is Endomorphism by definition, we conclude that \mathcal{R} is a group Automorphism from $(\text{GL}_n(K), \circ)$ to $(\text{GL}_n(K), \circ)$. \square

Theorem 1. *Let $\mathcal{C}(A)$ be the combined matrix of $A \in \text{GL}_n(K)$ as in Eq. (2). Let $L = K[\lambda_1, \dots, \lambda_n]$ be the extension of K . The following statements hold*

1. $\mathcal{C}(A) = [m_{ij}]_{n \times n}$, $m_{ij} = \frac{(-1)^{i+j}}{\det A} a_{ij} A_{ij}$
2. $\text{Tr}(\mathcal{C}(A)) = \frac{1}{\det A} \sum_{k=1}^n a_{ii} A_{ii}$
3. $(1, v_1) \in \Lambda(\mathcal{C}(A)) \times \mathcal{V}(\mathcal{C}(A))$, $v_1 = \sum_{k=1}^n e_k$
4. $\text{Gal}(L/K) \leq G$, $G \cong S_{n-1}$
5. $\mathcal{C}(\mathcal{R}(A)) = \mathcal{R}(\mathcal{C}(A))$

Proof. We proceed according to each item.

1. Due to $A \in \text{GL}_n(K)$, A is non-singular and formula (2) is valid. Now, by direct application of basic properties of transpose matrix and inverse matrix we obtain the result.
2. It follows by direct application of item 1.
3. By induction and basic properties of determinants we have $P_{\mathcal{C}(A)}(1) = 0$ and therefore $\lambda_1 = 1$ is an eigenvalue of $\mathcal{C}(A)$. Due to vector v_1 corresponds to $(1, 1, \dots, 1)^T \in K^n$, we obtain $\mathcal{C}(A) \cdot v_1 = v_1$ and therefore v_1 is an eigenvector of $\mathcal{C}(A)$.
4. By item 3 we have $p_{\mathcal{C}(A)}(1) = 0$, thus $p_{\mathcal{C}(A)}(\lambda) = (\lambda - 1)q_{n-1}(\lambda)$, where q_{n-1} is a polynomial of degree $n - 1$. Therefore the extension L is obtained as $K[\lambda_2, \dots, \lambda_n]$, which implies that $\text{Gal}(L/K)$ is a subgroup of S_{n-1} .
5. It follows directly from Lemma 1. Thus, Reversing of the combined matrix of A is the combined matrix of Reversing of A .

□

2.2 Combined matrices over proper subgroups of General Linear Group

Here we consider combined matrices over some proper subgroups of $\text{GL}_n(K)$ such as $\text{SL}_n(K)$, triangular matrices, orthogonal matrices, among others. We start with the following Lemma.

Lemma 2. *Let (T_n^+, \cdot) and (T_n^-, \cdot) be the groups of $n \times n$ upper triangular and lower triangular matrices respectively with the usual product of matrices. Then $T_n^+ \circ T_n^- = \text{diag}(T_n^+) \cdot \text{diag}(T_n^-)$, where \circ is the Hadamard product of matrices and diag denotes the diagonal matrix.*

Proof. Consider $T_n^+ = [a_{ij}]_{n \times n}$ and $T_n^- = [b_{ij}]_{n \times n}$ such that $a_{ij} = 0$ for all $i < j$ and $b_{ij} = 0$ for all $i > j$. Thus, $T_n^+ \circ T_n^- = [a_{ij} \cdot b_{ij}]_{n \times n} = [a_{ii} \cdot b_{ii}]_{n \times n} = [a_{ii}]_{n \times n} \circ [b_{ii}]_{n \times n}$ which lead us to the result. □

Proposition 1. *The following statements hold.*

1. $\mathcal{C}(T_n) = I_n$
2. $\Lambda(\mathcal{C}(T_n)) = \{1\}$.
3. $\mathcal{C}(AB) = \mathcal{C}(A)\mathcal{C}(B)$, being A, B triangular matrices.

Proof. We proceed according to each item.

1. By definition of combined matrix of A we have $\mathcal{C}(A) = A \circ A^{-T}$. Now, inverse of upper (resp. lower) triangular matrix is upper (resp. lower) triangular matrix. In the same way, transpose of upper (resp. lower) triangular matrix is a lower (resp. upper) triangular matrix and the determinant of any triangular (lower and upper) is the product of the diagonal entries. Thus, by Lemma 2 we have that $\mathcal{C}(T_n^\pm)$ are diagonal matrices and $T_n^\pm \circ (T_n^\pm)^{-T}$ and because we divide by the product of the diagonal entries, we obtain $\mathcal{C}(T_n^\pm) = I_n$.
2. It follows directly from item 1 because the identity matrix has 1 as eigenvalue of algebraic multiplicity n .

3. It follows due to the $\ker(\mathcal{C})$ for triangular matrices is the identity. □

The following result involves the *Orthogonal Group* over K , that is, the group of matrices with entries in K such that the product of a matrix with its transpose is the identity. We observe that Orthogonal Group is a subgroup of $SL_n(K)$

Proposition 2. *The matrix A is an orthogonal matrix if and only if $\mathcal{C}(A) = A \circ A$.*

Proof. Due to $A^{-1} = A^T$ we have that $A^{-T} = A$ and therefore $\mathcal{C}(A) = A \circ A$. □

Example 1. *Consider the following matrix*

$$A = -\frac{1}{3} \begin{pmatrix} 2 & -2 & 1 \\ 1 & 2 & 2 \\ 2 & 1 & -2 \end{pmatrix}, \quad A^{-1} = A^T = -\frac{1}{3} \begin{pmatrix} 2 & 1 & 2 \\ -2 & 2 & 1 \\ 1 & 2 & -2 \end{pmatrix},$$

the combined matrix of A is

$$\mathcal{C}(A) = \frac{1}{9} \begin{pmatrix} 4 & 4 & 1 \\ 1 & 4 & 4 \\ 4 & 1 & 4 \end{pmatrix}.$$

2.3 Algebraic aspects of combined matrices of order two nonsingular matrices

Here we analyze some algebraic aspects of characteristic polynomials of nonsingular matrices of size 2×2 . Consider

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \quad \text{Tr}(A) = a_{11} + a_{22}, \quad \det A = a_{11}a_{22} - a_{12}a_{21}.$$

The combined matrix of the nonsingular matrix A is

$$\mathcal{C}(A) = \frac{1}{\det A} \begin{pmatrix} a_{11}a_{22} & -a_{12}a_{21} \\ -a_{12}a_{21} & a_{11}a_{22} \end{pmatrix} \tag{3}$$

The following propositions are immediate consequences of the previous results for $n = 1$ and by the properties studied in a basic course of linear algebra for undergraduate level.

Proposition 3. *Consider $A \in GL_2(K)$, being K a number field. The following statements hold*

1. $\text{Tr}(\mathcal{C}(A)) = \frac{2}{\det A} a_{11}a_{22}$
2. $\det(\mathcal{C}(A)) = \frac{1}{\det(A)} (a_{11}a_{22} + a_{12}a_{21})$
3. *The characteristic polynomial of $\mathcal{C}(A)$ is*

$$\lambda^2 - \left(\frac{2a_{11}a_{22}}{a_{11}a_{22} - a_{12}a_{21}} \right) \lambda + \frac{a_{11}a_{22} + a_{12}a_{21}}{a_{11}a_{22} - a_{12}a_{21}}$$

4. *The eigenvalues of $\mathcal{C}(A)$ are given by*

$$\Lambda = \{1, \det \mathcal{C}(A)\}$$

5. *The eigenvectors of $\mathcal{C}(A)$ are given by*

$$\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}$$

6. *The Galois group $\text{Gal}(L/K)$ of the characteristic polynomial is the identity group.*

7. *The combined matrix of A is diagonalizable:*

$$\mathcal{C}(A) = PDP^{-1}, \quad P = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0 \\ 0 & \det(\mathcal{C}(A)) \end{pmatrix}, \quad P^{-1} = \frac{1}{2} \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}$$

Moreover, if f is an analytic function in 1 and in $\det(\mathcal{C}(A))$, then

$$f(\mathcal{C}(A)) = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} f(1) & 0 \\ 0 & f(\det(\mathcal{C}(A))) \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}$$

Proposition 4. Consider A such that $A^2 \in \text{SL}_2(K)$, being K a number field. The following statements hold

1. $\text{Tr}(\mathcal{C}(A)) = \pm 2a_{12}a_{21} + 2$
2. $\det(\mathcal{C}(A)) = \text{Tr}(\mathcal{C}(A)) - 1$
3. The characteristic polynomial of $\mathcal{C}(A)$ is

$$p_{\mathcal{C}(A)}(\lambda) = \lambda^2 - (\pm 2a_{12}a_{21} + 2)\lambda \pm 2a_{12}a_{21} + 1$$

4. The eigenvalues of $\mathcal{C}(A)$ are given by

$$\Lambda = \{1, \pm 2a_{12}a_{21} + 1\}$$

5. The eigenvectors of $\mathcal{C}(A)$ are given by

$$\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}$$

6. The Galois group of the characteristic polynomial $\text{Gal}(L/K)$ is the identity group.
7. The combined matrix of A is diagonalizable:

$$\mathcal{C}(A) = PDP^{-1}, \quad P = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} 1 & 0 \\ 0 & \pm 2a_{12}a_{21} + 1 \end{pmatrix}, \quad P^{-1} = \frac{1}{2} \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}$$

Moreover, if f is an analytic function in 1 and in $\pm 2a_{12}a_{21} + 1$, then

$$f(\mathcal{C}(A)) = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} f(1) & 0 \\ 0 & f(\pm 2a_{12}a_{21} + 1) \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}$$

2.4 What about combined matrices for order three nonsingular matrices

Consider the matrix $A \in \text{GL}_3(K)$, being K a field number, given by

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

The determinant of A , denoted by $\det A$, is

$$\det A = a_{11}a_{22}a_{33} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31}.$$

The combined matrix of A is

$$\mathcal{C}(A) = \frac{1}{\det A} \begin{pmatrix} a_{11}a_{22}a_{33} - a_{23}a_{32} & -a_{12}a_{21}a_{33} - a_{23}a_{31} & a_{13}a_{21}a_{32} - a_{22}a_{31} \\ -a_{21}a_{12}a_{33} - a_{13}a_{32} & a_{22}a_{11}a_{33} - a_{13}a_{31} & -a_{23}a_{11}a_{32} - a_{12}a_{31} \\ a_{31}a_{12}a_{23} - a_{13}a_{22} & -a_{32}a_{11}a_{23} - a_{13}a_{21} & a_{33}a_{11}a_{22} - a_{12}a_{21} \end{pmatrix}$$

Due to the long expressions for characteristic polynomials in this case, we avoid to repeat the results for $n \times n$ that are valids for $n = 3$. In other hand, we cannot obtain similar results as for $n = 2$ because from $n = 3$ and so on eigenvectors will depend on the entries of the original matrix A , except the first eigenvector which correspond to the eigenvalue 1. Thus, cannot be possible to obtain beautiful formulas for the eigenvalues, eigenvectors and the diagonalization (whether can be possible) of combined matrices for $n > 2$.

3 Final remarks

In this paper we analyze some algebraic aspects of characteristics polynomials of combined matrices. In particular we obtained the the Galois group of the characteristic polynomial is the identity group and in particular one is an eigenvalue of $\mathcal{C}(A)$. A natural question that appears is the extension of this result to Quantum Mechanics and Differential Galois Theory, see [1, 3].

References

- [1] P. B. Acosta-Humánez. *Galoisian Approach to Supersymmetric Quantum Mechanics: The integrability analysis of the Schrödinger equation by means of differential Galois theory*. VDM Publishing, 2010.
- [2] P. B. Acosta-Humánez and A. L. Chuquen. Pasting and reversing approach to matrix theory. *JP Journal of Algebra, Number Theory and Applications*, 38(6):535–559, 2016.
- [3] P. B. Acosta-Humánez, J. J. Morales-Ruiz, and J.-A. Weil. Galoisian approach to integrability of schrödinger equation. *Reports on Mathematical Physics*, 67(3):305–374, 2011.
- [4] P. Alonso, J. M. Peña, and M. L. Serrano. Combined matrices of almost strictly sign regular matrices. *Journal of computational and applied mathematics*, 354:144–151, 2019.
- [5] R. Bru, M. T. Gassó, I. Giménez, and M. Santana. Nonnegative combined matrices. *Journal of Applied Mathematics*, 2014, 2014.
- [6] R. Bru, M. T. Gassó, I. Giménez, and M. Santana. Combined matrices of sign regular matrices. *Linear Algebra and Its Applications*, 498:88–98, 2016.
- [7] R. Bru, M. T. Gassó, and M. Santana. Combined matrices and conditioning. *Applied Mathematics and Computation*, 412:126549, 2022.
- [8] J. A. Charris Castañeda, B. Aldana Gómez, and P. B. Acosta-Humanez. *Álgebra: fundamentos, grupos, anillos, cuerpos y Teoría de Galois*. Academia Colombiana de Ciencias Exactas, Físicas y Naturales, 2013.
- [9] M. Fiedler. Some results on combined matrices. *Banach Center Publications*, 112:99–105, 2017.
- [10] M. Fiedler and T. L. Markham. Combined matrices in special classes of matrices. *Linear algebra and its applications*, 435(8):1945–1955, 2011.
- [11] M. T. Gassó, I. Gil, I. Giménez, M. Santana, and E. Segura. Diagonal entries of the combined matrix of sign regular matrices of order three. *Proyecciones (Antofagasta)*, 40(1):255–271, 2021.