STEVEN FINCH

May 1, 2022

ABSTRACT. We investigate decomposable combinatorial labeled structures more fully, focusing on the exp-log class of type $a = 1$ or $1/2$. For instance, the modal length of the second longest cycle in a random n -permutation is $(0.2350...)n$, whereas the modal length of the second smallest component in a random *n*-mapping is 2 (conjecturally, given $n \geq 434$). As in earlier work, our approach is to establish how well existing theory matches experimental data and to raise open questions.

Given a combinatorial object with n nodes, our interest is in

- \bullet the size of its r^{th} longest cycle or largest component,
- \bullet the size of its rth shortest cycle or smallest component

where $r \geq 2$. If the object has no rth component, then its rth largest/smallest components are defined to have length 0. The case $r = 1$ has attracted widespread attention [\[1,](#page-15-0) [2\]](#page-15-1). Key to our prior study were recursive formulas [\[3,](#page-15-2) [4\]](#page-15-3) for $L_{k,n}$ and $S_{k,n}$, the number of *n*-objects whose largest and smallest components, respectively, have exactly k nodes, $1 \leq k \leq n$. Different algorithms shall be used here. As before, an *n*-object is chosen uniformly at random. For simplicity, we discuss here only *n*-permutations and *n*-mappings (from $\{1, 2, \ldots, n\}$ to $\{1, 2, \ldots, n\}$). Let c_n be the number of n-objects that are connected, i.e., who possess exactly one component:

$$
c_n = \begin{cases} (n-1)! & \text{for permutations,} \\ n! \sum_{j=1}^n \frac{n^{n-j-1}}{(n-j)!} & \text{for mappings.} \end{cases}
$$

The total number of *n*-permutations and *n*-mappings is *n*! and n^n , respectively. For fixed n, the sequences $\{L_{k,n}: 1 \leq k \leq n\}$ and $\{S_{k,n}: 1 \leq k \leq n\}$ constitute probability mass functions (upon normalization) for $r = 1$. Until recently, calculating analogous sequences for $r \geq 2$ seemed inaccessibly difficult.

The new algorithms, due to Heinz $[5]$, accept as input the integer n and an ordered r-tuple ℓ of nonnegative integers, which may include infinity. We write ℓ as a list $\{i_1, i_2, \ldots, i_r\}$. Given a positive integer j, define ℓ^j to be the list obtained by

 0 Copyright \odot 2022 by Steven R. Finch. All rights reserved.

- (i) appending ℓ with j ,
- (ii) sorting the $(r + 1)$ -tuple in ascendent order, and
- (iii) removing its first element.

Define ℓ_j in the same way as ℓ^j except for a revised final step:

(iii') removing its last element.

Note that the lengths of ℓ^j and ℓ_j are always equal to the length of ℓ . Let $p[n, \ell]$ and $q[n, \ell]$ denote row polynomials in x and y associated with large and small components. The algorithms are based on recursions

$$
p[n,\ell] = \begin{cases} \sum_{j=1}^{n} c_j p[n-j,\ell^j] \binom{n-1}{j-1} & \text{if } n > 0, \\ x^{i_1} & \text{if } n = 0; \end{cases}
$$
\n
$$
q[n,\ell] = \begin{cases} \sum_{j=1}^{n} c_j q[n-j,\ell_j] \binom{n-1}{j-1} & \text{if } n > 0, \\ y^{i_r} & \text{if } n = 0 \text{ and } i_r < \infty, \\ y^0 & \text{if } n = 0 \text{ and } i_r = \infty. \end{cases}
$$

A computer algebra software package (e.g., Mathematica or Maple) makes exact integer calculations for ample n of $p[n, \ell]$ and $q[n, \ell]$ feasible. These are demonstrated for $n = 4$ in the next section, for the sake of concreteness.

Permutations belong to the exp-log class of type $a = 1$, whereas mappings belong to the exp-log class of type $a = 1/2$. Explaining the significance of the parameter $a > 0$ would take us too far afield [\[6\]](#page-15-5). Let

$$
E(x) = \int_{x}^{\infty} \frac{e^{-t}}{t} dt = -\operatorname{Ei}(-x), \quad x > 0
$$

be the exponential integral. Define [\[7,](#page-15-6) [8,](#page-15-7) [9,](#page-15-8) [10,](#page-15-9) [11,](#page-15-10) [12\]](#page-15-11)

$$
{}_{L}G_{a}(r,h) = \frac{\Gamma(a+1)a^{r-1}}{\Gamma(a+h)(r-1)!} \int_{0}^{\infty} x^{h-1} E(x)^{r-1} \exp\left[-a E(x) - x\right] dx,
$$

SECOND BEST, THIRD WORST, FOURTH IN LINE 3

$$
{}_S G_a(r,h) = \begin{cases} e^{-h\gamma} a^{r-1}/r! & \text{if } h = a, \\ \frac{\Gamma(a+1)}{(h-1)!(r-1)!} \int_0^\infty x^{h-1} \exp\left[a E(x) - x\right] dx & \text{if } h > a \end{cases}
$$

which are related to the hth moment of the rth largest/smallest component size (in this paper, rank $r = 2, 3$ or 4; height $h = 1$ or 2). While moment formulas are unerring for L, they are not so for S. While $_{S}G_{a}$ is flawless for permutations (and for what are called *cyclations* [\[13\]](#page-15-12)), a correction factor $\sqrt{2}$ is needed for mappings.

For fixed n and r , the coefficient sequences associated with polynomials

$$
p[n, \{\underbrace{0,0,\cdots,0}_{r}\}], \quad 0 \le k \le \lfloor n/r \rfloor;
$$

$$
q[n, \{\underbrace{\infty,\infty,\cdots,\infty}_{r}\}], \quad 0 \le k \le n-r+1
$$

constitute probability mass functions (upon normalization). These have corresponding means $L\mu_{n,r}$, $s\mu_{n,r}$ and variances $L\sigma_{n,r}^2$, $s\sigma_{n,r}^2$ given in the tables. We also provide the median $L\nu_{n,r}$ and mode $L\vartheta_{n,r}$; evidently $s\nu_{n,r}$ and $s\vartheta_{n,r}$ are bounded for permutations as $n \to \infty$ (the trend of $s\nu_{n,r}$ is less clear for mappings). In table headings only, the following notation is used:

$$
L\widetilde{\mu}_{n,r} = \frac{L\mu_{n,r}}{n}, \qquad L\widetilde{\sigma}_{n,r}^2 = \frac{L\sigma_{n,r}^2}{n^2}, \qquad L\widetilde{\nu}_{n,r} = \frac{L\nu_{n,r}}{n}, \qquad L\widetilde{\vartheta}_{n,r} = \frac{L\vartheta_{n,r}}{n}, \qquad S\widetilde{\nu}_{n,r} = \frac{S\nu_{n,r}}{n},
$$

$$
S\widetilde{\mu}_{n,r} = \begin{cases} \frac{S\mu_{n,r}}{\ln(n)^r} & \text{if } a = 1, \\ \frac{S\mu_{n,r}}{n^{1/2}\ln(n)^{r-1}} & \text{if } a = 1/2, \end{cases} \qquad S\widetilde{\sigma}_{n,r}^2 = \begin{cases} \frac{S\sigma_{n,r}^2}{n\ln(n)^{r-1}} & \text{if } a = 1, \\ \frac{S\sigma_{n,r}^2}{n^{3/2}\ln(n)^{r-1}} & \text{if } a = 1/2. \end{cases}
$$

When $r = 1$, the mode $L\widetilde{\vartheta}_{n,1}$ is provably $1/2$ in the limit as $n \to \infty$ for permutations (it is 1 for mappings). This limit is more interesting when $r = 2$, as will soon be seen.

1. CALCULS À LA HEINZ

As promised, we exhibit some hand calculations. It is easy to show directly that $p[3, \{0, 0\}] = 2 + 4x$ for permutations and $17 + 10x$ for mappings (see Section 3 of [\[1\]](#page-15-0)). More generally, $p[3, \{0, 0\}] = c_3 + c_1(c_1^2 + 3c_2)x$. Let us compute $p[4, \{0, 0\}]$ using Heinz's algorithm. From

$$
p[2, \{1, 1\}] = c_1 p[1, \{1, 1\}] \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 p[0, \{1, 2\}] \begin{pmatrix} 1 \\ 1 \end{pmatrix}
$$

= $c_1^2 p[0, \{1, 1\}] \begin{pmatrix} 0 \\ 0 \end{pmatrix} + c_2 x^1 = (c_1^2 + c_2) x$,

SECOND BEST, THIRD WORST, FOURTH IN LINE 4

$$
p[1, \{1, 2\}] = c_1 p[0, \{1, 2\}] \binom{0}{0} = c_1 x,
$$

$$
p[0, \{1, 3\}] = x
$$

we have

$$
p[3, \{0, 1\}] = c_1 p[2, \{1, 1\}] \begin{pmatrix} 2 \\ 0 \end{pmatrix} + c_2 p[1, \{1, 2\}] \begin{pmatrix} 2 \\ 1 \end{pmatrix} + c_3 p[0, \{1, 3\}] \begin{pmatrix} 2 \\ 2 \end{pmatrix}
$$

= $c_1 (c_1^2 + c_2) x + 2c_2(c_1 x) + c_3 x = (c_1^3 + 3c_1 c_2 + c_3) x$.

Also, from

$$
p[2, \{0, 2\}] = c_1 p[1, \{1, 2\}] \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 p[0, \{2, 2\}] \begin{pmatrix} 1 \\ 1 \end{pmatrix}
$$

= $c_1^2 p[0, \{1, 2\}] \begin{pmatrix} 0 \\ 0 \end{pmatrix} + c_2 x^2 = c_1^2 x + c_2 x^2$,

$$
p[1, \{0, 3\}] = c_1 p[0, \{1, 3\}] \begin{pmatrix} 0 \\ 0 \end{pmatrix} = c_1 x
$$
,

$$
p[0, \{0, 4\}] = x^0 = 1
$$

we deduce

$$
p[4, \{0, 0\}] = c_1 p[3, \{0, 1\}] \binom{3}{0} + c_2 p[2, \{0, 2\}] \binom{3}{1} + c_3 p[1, \{0, 3\}] \binom{3}{2} + c_4 p[0, \{0, 4\}] \binom{3}{3}
$$

= $c_1 (c_1^3 + 3c_1c_2 + c_3) x + 3c_2(c_1^2 x + c_2 x^2) + 3c_3(c_1 x) + c_4$
= $c_4 + c_1 (c_1^3 + 6c_1c_2 + 4c_3) x + 3c_2^2 x^2$
= $\begin{cases} 6 + 15x + 3x^2 & \text{for permutations,} \\ 142 + 87x + 27x^2 & \text{for mappings} \end{cases}$

completing the argument.

It is likewise easy to show that $q[3, \{0, 0\}] = 2 + y + 3y^2$ for permutations and $17 + y + 9y^2$ for mappings. More generally, $q[3, \{0, 0\}] = c_3 + c_1^3y + 3c_1c_2y^2$. Let us compute $q[4, \{0, 0\}]$ using Heinz's algorithm. From

$$
q[2, \{1, 1\}] = c_1 q[1, \{1, 1\}] \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 q[0, \{1, 1\}] \begin{pmatrix} 1 \\ 1 \end{pmatrix}
$$

= $c_1^2 q[0, \{1, 1\}] \begin{pmatrix} 0 \\ 0 \end{pmatrix} + c_2 y^1 = (c_1^2 + c_2) y,$

$$
q[1, \{1, 2\}] = c_1 q[0, \{1, 1\}] \begin{pmatrix} 0 \\ 0 \end{pmatrix} = c_1 y,
$$

$$
q[0, \{1, 3\}] = y^3
$$

we have

$$
q[3,\{1,\infty\}] = c_1 q[2,\{1,1\}] \begin{pmatrix} 2 \\ 0 \end{pmatrix} + c_2 q[1,\{1,2\}] \begin{pmatrix} 2 \\ 1 \end{pmatrix} + c_3 q[0,\{1,3\}] \begin{pmatrix} 2 \\ 2 \end{pmatrix}
$$

= c₁ $(c_1^2 + c_2) y + 2c_2(c_1 y) + c_3 y^3 = (c_1^3 + 3c_1c_2) y + c_3 y^3.$

Also, from

$$
q[2, \{2, \infty\}] = c_1 q[1, \{1, 2\}] \begin{pmatrix} 1 \\ 0 \end{pmatrix} + c_2 q[0, \{2, 2\}] \begin{pmatrix} 1 \\ 1 \end{pmatrix}
$$

= $c_1^2 q[0, \{1, 1\}] \begin{pmatrix} 0 \\ 0 \end{pmatrix} + c_2 y^2 = c_1^2 y + c_2 y^2$,

$$
q[1, \{3, \infty\}] = c_1 q[0, \{1, 3\}] \begin{pmatrix} 0 \\ 0 \end{pmatrix} = c_1 y^3,
$$

$$
q[0, \{4, \infty\}] = y^0 = 1
$$

we deduce

$$
q[4, \{\infty, \infty\}] = c_1 q[3, \{1, \infty\}] \binom{3}{0} + c_2 q[2, \{2, \infty\}] \binom{3}{1} + c_3 q[1, \{3, \infty\}] \binom{3}{2} + c_4 q[0, \{4, \infty\}] \binom{3}{3}
$$

\n
$$
= c_1 \left(\left(c_1^3 + 3c_1 c_2 \right) y + c_3 y^3 \right) + 3c_2 \left(c_1^2 y + c_2 y^2 \right) + 3c_3 (c_1 y^3) + c_4
$$

\n
$$
= c_4 + c_1^2 \left(c_1^2 + 6c_2 \right) y + 3c_2^2 y^2 + 4c_1 c_3 y^3
$$

\n
$$
= \begin{cases} 6 + 7y + 3y^2 + 8y^3 & \text{for permutations,} \\ 142 + 19y + 27y^2 + 68y^3 & \text{for mappings} \end{cases}
$$

completing the argument.

2. Modes & Medians

The mode of a continuous distribution is the location of its highest peak; the median is its $50th$ percentile. The length Λ_r of the rth longest cycle in a random *n*-permutation has cumulative probability

$$
\lim_{n \to \infty} \mathbb{P}\left\{\Lambda_r < x \cdot n\right\} = \rho_r\left(\frac{1}{x}\right)
$$

where $\rho_r(x)$ is the r^{th} order Dickman function [\[14\]](#page-15-13):

$$
x\rho'_1(x) + \rho_1(x-1) = 0 \text{ for } x > 1, \qquad \rho_1(x) = 1 \text{ for } 0 \le x \le 1;
$$

$$
x\rho'_r(x) + \rho_r(x-1) = \rho_{r-1}(x-1) \text{ for } x > 1, \qquad \rho_r(x) = 1 \text{ for } 0 \le x \le 1
$$

and $r = 2, 3, 4, \ldots$ For notational simplicity, let us write $\varphi = \rho_1$ and $\psi = \rho_2$. Observe that ρ_r should not be confused with a different generalization ρ_a discussed in [\[1,](#page-15-0) [15\]](#page-16-0).

From

$$
\varphi'(x) = -\frac{\varphi(x-1)}{x}, \quad x > 1
$$

we have

$$
\varphi'\left(\frac{1}{x}\right) = -\frac{\varphi\left(\frac{1}{x} - 1\right)}{\frac{1}{x}}, \qquad 0 < x < 1
$$

hence the density $f(x)$ is

$$
\frac{d}{dx}\varphi\left(\frac{1}{x}\right) = -x\,\varphi\left(\frac{1}{x} - 1\right)\left(-\frac{1}{x^2}\right) = \begin{cases} \frac{\varphi\left(\frac{1}{x} - 1\right)}{x} & \text{if } 0 < x \le \frac{1}{2}, \\ \frac{1}{x} & \text{if } \frac{1}{2} < x < 1. \end{cases}
$$

Also, from

$$
\varphi''(x) = \frac{\varphi(x-1)}{x^2} - \frac{\varphi'(x-1)}{x} = \frac{\varphi(x-1)}{x^2} + \frac{\varphi(x-2)}{x(x-1)}, \qquad x > 1
$$

we have

$$
\varphi''\left(\frac{1}{x}\right) = \frac{\varphi\left(\frac{1}{x} - 1\right)}{\frac{1}{x^2}} + \frac{\varphi\left(\frac{1}{x} - 2\right)}{\frac{1}{x}\left(\frac{1}{x} - 1\right)}, \qquad 0 < x < 1
$$

hence (by the chain rule for second derivatives)

$$
\frac{d^2}{dx^2}\varphi\left(\frac{1}{x}\right) = \varphi'\left(\frac{1}{x}\right)\frac{2}{x^3} + \frac{1}{x^4}\varphi''\left(\frac{1}{x}\right)
$$

\n
$$
= \frac{-2\varphi\left(\frac{1}{x} - 1\right)}{x^2} + \frac{\varphi\left(\frac{1}{x} - 1\right)}{x^2} + \frac{\varphi\left(\frac{1}{x} - 2\right)}{x^2(1 - x)}
$$

\n
$$
= \begin{cases} \frac{1}{x^2(1-x)} - \frac{\varphi\left(\frac{1}{x} - 1\right)}{x^2} > \frac{1}{x(1-x)} > 0 & \text{if } 1/3 < x \le 1/2, \\ -\frac{1}{x^2} < 0 & \text{if } 1/2 < x \le 1 \end{cases}
$$

since the first condition implies $3 > 1/x \ge 2$, i.e., $1 > 1/x - 2 \ge 0$ and the second condition implies $2 > 1/x \ge 1$, i.e., $1 > 1/x - 1 \ge 0$. Thus f is increasing on the left of $x = 1/2$ and f is decreasing on the right, which implies that the median size of Λ_1 is $1/2$.

From

$$
\psi'(x) = \frac{\varphi(x-1) - \psi(x-1)}{x}, \quad x > 2
$$

we have

$$
\varphi'(x) - \psi'(x) = -\frac{\varphi(x-1)}{x} - \frac{\varphi(x-1) - \psi(x-1)}{x} = \frac{-2\varphi(x-1) + \psi(x-1)}{x}
$$

(a lemmata needed shortly) and

$$
\psi'\left(\frac{1}{x}\right) = \frac{\varphi\left(\frac{1}{x} - 1\right) - \psi\left(\frac{1}{x} - 1\right)}{\frac{1}{x}}, \quad 0 < x < \frac{1}{2}
$$

hence the density $g(\boldsymbol{x})$ is

$$
\frac{d}{dx}\psi\left(\frac{1}{x}\right) = x\left(\varphi\left(\frac{1}{x} - 1\right) - \psi\left(\frac{1}{x} - 1\right)\right)\left(-\frac{1}{x^2}\right)
$$

$$
= \frac{\psi\left(\frac{1}{x} - 1\right) - \varphi\left(\frac{1}{x} - 1\right)}{x}.
$$

Also, from

$$
\psi''(x) = -\frac{\varphi(x-1) - \psi(x-1)}{x^2} + \frac{\varphi'(x-1) - \psi'(x-1)}{x}
$$

$$
= \frac{-\varphi(x-1) + \psi(x-1)}{x^2} + \frac{-2\varphi(x-2) + \psi(x-2)}{x(x-1)}
$$

(by the lemmata) we have

$$
\psi''\left(\frac{1}{x}\right) = \frac{-\varphi\left(\frac{1}{x} - 1\right) + \psi\left(\frac{1}{x} - 1\right)}{\frac{1}{x^2}} + \frac{-2\varphi\left(\frac{1}{x} - 2\right) + \psi\left(\frac{1}{x} - 2\right)}{\frac{1}{x}\left(\frac{1}{x} - 1\right)}
$$

hence (by the chain rule for second derivatives)

$$
\frac{d^2}{dx^2}\psi\left(\frac{1}{x}\right) = \psi'\left(\frac{1}{x}\right)\frac{2}{x^3} + \frac{1}{x^4}\psi''\left(\frac{1}{x}\right)
$$
\n
$$
= \frac{\varphi\left(\frac{1}{x}-1\right) - \psi\left(\frac{1}{x}-1\right)}{\frac{1}{x^3}}\frac{2}{x^3}
$$
\n
$$
+ \frac{1}{x^4}\left[\frac{-\varphi\left(\frac{1}{x}-1\right) + \psi\left(\frac{1}{x}-1\right)}{\frac{1}{x^2}} + \frac{-2\varphi\left(\frac{1}{x}-2\right) + \psi\left(\frac{1}{x}-2\right)}{\frac{1}{x}\left(\frac{1}{x}-1\right)}\right]
$$
\n
$$
= \frac{\varphi\left(\frac{1}{x}-1\right) - \psi\left(\frac{1}{x}-1\right)}{x^2} - \frac{2\varphi\left(\frac{1}{x}-2\right) - \psi\left(\frac{1}{x}-2\right)}{x^2(1-x)}.
$$

There exists a unique $0 < x_0 < 1/2$ for which this expression $(g'(x_0))$ vanishes. Plots of $f(x)$ and $g(x)$ appear in [\[16\]](#page-16-1) and confirm that x_0 is the modal size of Λ_2 . Broadhurst [\[17\]](#page-16-2) obtained an exact equation for x_0 , involving Dickman dilogarithms and trilogarithms [\[18\]](#page-16-3), then applied numerics. We have verified his value x_0 by purely floating point methods.

There is comparatively little to say about medians ξ_r , defined as solutions of [\[14,](#page-15-13) [19\]](#page-16-4)

$$
\rho_r\left(\frac{1}{x}\right) = \frac{1}{2}
$$

except that $\xi_1 = 1/\sqrt{e}$ is well-known and no closed-form representations for ξ_r , $r \ge 2$, seem to exist.

3. KNUTH & TRABB PARDO

An alternative to Heinz's algorithm is one proposed by Knuth & Trabb Pardo [\[14\]](#page-15-13) for a restricted case. Define $u_r(k,n)$ to be the number of *n*-permutations whose rth longest cycle has $\leq k$ nodes [\[20\]](#page-16-5). The following recursive formulas apply for $r = 1$:

$$
u_1(k,n) = \begin{cases} \sum_{m=0}^{k-1} \frac{(n-1)!}{(n-1-m)!} u_1(k,n-1-m) & \text{if } n \ge 1 \text{ and } k < n, \\ n! & \text{if } n \ge 1 \text{ and } k \ge n, \\ 1 & \text{otherwise} \end{cases}
$$

and for $r \geq 2$:

$$
u_r(k,n) = \begin{cases} \sum_{m=0}^{k-1} \frac{(n-1)!}{(n-1-m)!} u_r(k, n-1-m) + & \text{if } n \ge 1 \text{ and } k < \lfloor n/r \rfloor, \\ \sum_{m=k}^{n-1} \frac{(n-1)!}{(n-1-m)!} u_{r-1}(k, n-1-m) & \text{if } n \ge 1 \text{ and } k \ge \lfloor n/r \rfloor, \\ n! & \text{if } n \ge 1 \text{ and } k \ge \lfloor n/r \rfloor, \\ 1 & \text{otherwise.} \end{cases}
$$

Clearly $u_1(0, n) = \delta_{0, n}$ and $u_1(1, n) = 1$, hence

$$
u_2(0,4) = u_1(0,3) + 3u_1(0,2) + 6u_1(0,1) + 6u_1(0,0) = 6.
$$

Also $u_2(1, 2) = 2$ and $u_2(1, 3) = 6$, hence

$$
u_2(1,4) = u_2(1,3) + [3u_1(1,2) + 6u_1(1,1) + 6u_1(1,0)] = 6 + 15 = 21.
$$

Finally $u_2(2, 4) = 24$. The list

$$
{u_2(k, 4)}_{k=0}^2 = {6, 21, 24} = {6, 6 + 15, 21 + 3}
$$

conveys the same information as the polynomial $p[4, \{0, 0\}]$ did in Section 1, although the underlying calculations differed completely.

A proof is as follows [\[14\]](#page-15-13). We may think of $u_r(k,n)$ as counting permutations on $\{1, \ldots, n\}$ that possess fewer than r cycles of length exceeding k. Call such a permutation (r, n) -good. Consider now a permutation P on $\{0, 1, \ldots, n\}$. The node 0 belongs to some cycle C within P of length $m + 1$. Let $P \setminus C$ denote the permutation which remains upon exclusion of C from P. Suppose $0 \le m \le k - 1$; then P is $(r, n+1)$ -good iff $P \setminus C$ is $(r, n-m)$ -good. Suppose $k \le m \le n$; then P is $(r, n+1)$ -good iff $P \setminus C$ is $(r-1, n-m)$ -good. Thus the formula

$$
u_r(k, n+1) = \sum_{m=0}^{k-1} \frac{n!}{(n-m)!} u_r(k, n-m) + \sum_{m=k}^{n} \frac{n!}{(n-m)!} u_{r-1}(k, n-m)
$$

is true because $n!/(n-m)!$ is the number of possible choices for C.

An analog of this recursion for mappings remains open, as far as is known. Finding the number of possible choices for a component C containing the node 0 is more complicated than for a cycle containing 0. Each component consists of a cycle with trees attached; each tree is rooted at a cyclic point but is otherwise made up of transient points. We must account for the position of 0 (cyclic or transient?) and the overall configuration (inventory of tree types and sizes?) It would be helpful to learn about progress in enumerating such C or, if this is impractical, some other procedure for moving forward.

4. Une conjecture correspondante

Short cycles have always presented more analytical difficulties than long cycles; this paper offers no exception. Everything in this section is conjectural only. Define $v_r(k, n)$ to be the number of *n*-permutations whose rth shortest cycle has $\geq k$ nodes [\[20\]](#page-16-5). The following recursive formulas would seem to apply for $r = 1$:

$$
v_1(k,n) = \begin{cases} n! & \text{if } n \ge 1 \text{ and } k = 0, \\ \sum_{m=k-1}^{n-1} \frac{(n-1)!}{(n-1-m)!} v_1(k, n-1-m) & \text{if } n \ge 1 \text{ and } 0 < k \le n, \\ 0 & \text{if } n \ge 1 \text{ and } k > n, \\ 1 & \text{otherwise} \end{cases}
$$

and for $r \geq 2$:

$$
v_r(k,n) = \begin{cases} n! & \text{if } n \ge 0 \text{ and } k = 0, \\ \sum_{m=0}^{k-2} \frac{(n-1)!}{(n-1-m)!} v_{r-1}(k, n-1-m) + & \text{if } n \ge 1 \text{ and } 0 < k \le n-r+1, \\ \sum_{m=k-1}^{n-1} \frac{(n-1)!}{(n-1-m)!} v_r(k, n-1-m) & \text{otherwise.} \end{cases}
$$

The surprising new term $\Delta_r(k, n)$ has a simple formula for $r = 2$:

$$
\Delta_2(k, n) = (n-1)! H_{n-k}
$$
, where $\sum_{i=1}^j \frac{1}{i} = H_j$, $\sum_{i=1}^j \frac{1}{i^s} = H_{j,s}$

and unexpected recursions for $r = 3$ and $r = 4$:

$$
\Delta_r(k,n) = \begin{cases}\n\frac{1}{2}(n-1)! \left(H_{n-1}^2 - H_{n-1,2} \right) & \text{if } r = 3 \text{ and } k = 1, \\
\frac{1}{6}(n-1)! \left(H_{n-1}^3 - 3H_{n-1}H_{n-1,2} + 2H_{n-1,3} \right) & \text{if } r = 4 \text{ and } k = 1, \\
\Delta_r(k-1,n) - \frac{\Delta_{r-1}(k,n)}{n-k+1} & \text{if } k \ge 2 \text{ and } n \ge k, \\
0 & \text{otherwise.} \n\end{cases}
$$

The values $\Delta_r(1,n)$ are unsigned Stirling numbers of the first kind, i.e., the number of *n*-permutations that have exactly *r* cycles. (Why should these appear here?)

A plausibility argument supporting v_r bears resemblance to the proof underlying u_r . We may think of $v_r(k,n)$ as counting permutations on $\{1,\ldots,n\}$ that possess fewer than r cycles of length surpassed by k. Call such a permutation (r, n) -bad. Let P & C (of lengths $n + 1$ & $m + 1$) be as before. Suppose $0 \le m \le k - 2$; then P is $(r, n+1)$ -bad iff $P \setminus C$ is $(r-1, n-m)$ -bad. Suppose $k-1 \le m \le n$; then P is $(r, n + 1)$ -bad iff $P \setminus C$ is $(r, n - m)$ -bad. This would suggest

$$
v_r(k, n+1) = \Delta_r + \sum_{m=0}^{k-2} \frac{n!}{(n-m)!} v_{r-1}(k, n-m) + \sum_{m=k-1}^{n} \frac{n!}{(n-m)!} v_r(k, n-m)
$$

is true with $\Delta_r = 0$, but experimental data contradict such an assertion.

Let us illustrate via example, in parallel with Section 3. As preliminary steps, $v_1(k, 0) = 1$ and $v_1(n + 1, n) = 0$, hence

$$
v_1(2,3) = 2v_1(2,1) + 2v_1(2,0) = 2
$$
, $v_1(3,3) = 2v_1(3,0) = 2$.

Clearly $v_2(0, 4) = 24$. Also $v_2(n, n) = \delta_{0,n}$ and $v_2(n + 1, n) = v_2(n + 2, n) = 0$, hence

$$
v_2(1,2) = \Delta_2(1,2) + [v_2(1,1) + v_2(1,0)] = 1 + 0 = 1,
$$

$$
v_2(1,3) = \Delta_2(1,3) + [v_2(1,2) + 2v_2(1,1) + 2v_2(1,0)] = 3 + 1 = 4,
$$

$$
v_2(1,4) = \Delta_2(1,4) + [v_2(1,3) + 3v_2(1,2) + 6v_2(1,1) + 6v_2(1,0)] = 11 + 7 = 18.
$$

Finally

$$
v_2(2,4) = \Delta_2(2,4) + v_1(2,3) + [3v_2(2,2) + 6v_2(2,1) + 6v_2(2,0)] = 9 + 2 + 0 = 11,
$$

$$
v_2(3,4) = \Delta_2(3,4) + [v_1(3,3) + 3v_1(3,2)] + [6v_2(3,1) + 6v_2(3,0)] = 6 + 2 + 0 = 8.
$$

Again, the list

$$
{v_2(k, 4)}_{k=1}^3 = {18, 11, 8} = {24 - 6, 18 - 7, 11 - 3 = 8}
$$

conveys the same information as the polynomial $q[4, {\infty, \infty}]$ did in Section 1. Without the nonzero contribution of $\Delta_r(k,n)$, our modification of Knuth & Trabb Pardo would yield results incompatible with Heinz.

5. PERMUTATIONS

Here [\[21\]](#page-16-6) are numerical results for $r = 2$:

Table 5.1: Statistics for Permute, rank two $(a = 1)$

as well as $s\nu_{n,2} = 2$ for $n > 17$ and $s\vartheta_{n,2} = 1$ for $n > 4$. Also

$$
\lim_{n \to \infty} \frac{L\mu_{n,2}}{n} = L G_1(2, 1) = 0.20958087428418581398...,
$$

$$
\lim_{n \to \infty} \frac{L\sigma_{n,2}^2}{n^2} = L G_1(2,2) - L G_1(2,1)^2 = 0.01255379063590587814...,
$$

$$
\lim_{n \to \infty} \frac{L^{\nu_{n,2}}}{n} = \xi_2 = 0.21172114641298273896...,
$$

\n
$$
\lim_{n \to \infty} \frac{L^{\vartheta_{n,2}}}{n} = x_0 = 0.23503964593509109370...,
$$

\n
$$
\lim_{n \to \infty} \frac{s\mu_{n,2}}{\ln(n)^2} = \frac{e^{-\gamma}}{2} = 0.28072974178344258491...,
$$

\n
$$
\lim_{n \to \infty} \frac{s\sigma_{n,2}^2}{n \ln(n)} = sG_P(2, 2) = 1.30720779891056809974....
$$

The final $n \ln(n)$ asymptotic is based on [\[7,](#page-15-6) [8\]](#page-15-7), not (inaccurate) Theorem 5 in [\[6\]](#page-15-5). Here [\[22\]](#page-16-7) are numerical results for $r = 3$:

Table 5.2: Statistics for Permute, rank three $(a = 1)$

as well as $s\nu_{n,2} = 7$ for $n > 370$ and $s\vartheta_{n,2} = 2$ for $n > 49$. Also

$$
\lim_{n \to \infty} \frac{L\mu_{n,3}}{n} = L G_1(3, 1) = 0.08831609888315363101...,
$$

 $\lim_{n\to\infty}$ $L\sigma_{n,3}^2$ $\frac{\partial n_{n,3}}{\partial n^2} = {}_L G_1(3,2) - {}_L G_1(3,1)^2 = 0.00449392318179080474...,$ $\lim_{n\to\infty}$ $L\nu_{n,3}$ n $=\xi_3 = 0.07584372316630152789...,$ $\lim_{n\to\infty}$ $_L\vartheta_{n,3}$ n $= 0,$ $\lim_{n\to\infty}$ $s\mu_{n,3}$ $\frac{\partial P_{n,3}}{\partial \ln(n)^3} =$ $e^{-\gamma}$ 6 $= 0.09357658059448086163...,$ $\lim_{n\to\infty}$ $s\sigma_{n,3}^2$ $\frac{S_{1}^{3}n_{1,3}}{n\ln(n)^{2}} = {}_{S}G_{P}(3,2) = 0.65360389945528404987...$

The final $n \ln(n)^2$ asymptotic is based on [\[7,](#page-15-6) [8\]](#page-15-7).

Here [\[23\]](#page-16-8) are numerical results for $r = 4$:

Table 5.3: Statistics for Permute, rank four $(a = 1)$

as well as $s\nu_{n,4} = 19$ for $n > 1482$ and $s\vartheta_{n,4} = 3$ for $n > 666$. Also

$$
\lim_{n \to \infty} \frac{L\mu_{n,4}}{n} = L G_1(4, 1) = 0.04034198873687046287...,
$$

\n
$$
\lim_{n \to \infty} \frac{L \sigma_{n,4}^2}{n^2} = L G_1(4, 2) - L G_1(4, 1)^2 = 0.00158383677354017280...,
$$

\n
$$
\lim_{n \to \infty} \frac{L \nu_{n,4}}{n} = \xi_4 = 0.02713839684981404992...,
$$

\n
$$
\lim_{n \to \infty} \frac{L \vartheta_{n,4}}{n} = 0,
$$

\n
$$
\lim_{n \to \infty} \frac{s \mu_{n,4}}{\ln(n)^4} = \frac{e^{-\gamma}}{24} = 0.02339414514862021540...,
$$

\n
$$
\lim_{n \to \infty} \frac{s \sigma_{n,4}^2}{n \ln(n)^3} = s G_P(4, 2) = 0.21786796648509468329...
$$

The final $n \ln(n)^3$ asymptotic is based on [\[7,](#page-15-6) [8\]](#page-15-7).

6. Mappings

Our modified Knuth & Trabb Pardo algorithm is unavailable in this setting, thus we turn to Heinz's program. A general observation for $2 \le r \le 4$ is $L\vartheta_{n,r} = 0$ always. Here [\[24\]](#page-16-9) are numerical results for $r = 2$:

$\,n$	$L\mu_{n,2}$	$L\overline{\widetilde{\sigma}_{n,2}^2}$	$\widetilde{\nu}_{n,2}$	$S\mu_{n,2}$	$s\widetilde{\sigma}_{n,2}^2$	$S\widetilde{\nu}_{n,2}$
100		0.166817 0.019535 0.1300 0.680589 0.279032				0.1200
200		0.168100 0.019243 0.1400 0.718071 0.323910				0.0750
-300		0.168642 0.019121 0.1433 0.737331 0.350358				0.0567
400		0.168959 0.019050 0.1450 0.749928 0.368810				0.0450
Table 6.1. Statistics for Map rape two $(a-1/2)$						

Table 6.1: Statistics for Map, rank two $(a = 1/2)$

as well as $s\nu_{n,2} = 19$ for $n > 443$ and $s\vartheta_{n,2} = 2$ for $n > 433$. Let us elaborate on the latter statistic (because it seems surprising at first glance: an extended string of 0s abruptly switches to 2s). If $\pi_r(k,n)$ denotes the probability that the rth smallest component of a random n -mapping has exactly k nodes, then

$$
\begin{array}{l} \left\{ \pi_2(k,432\right\}_{k=0}^4=\left\{0.0595400, 0.0532617, 0.0594378, 0.0477544, 0.0387585\right\},\\ \left\{ \pi_2(k,433\right\}_{k=0}^4=\left\{0.0594720, 0.0532614, 0.0594373, 0.0477539, 0.0387581\right\},\\ \left\{ \pi_2(k,434\right\}_{k=0}^4=\left\{0.0594044, 0.0532612, 0.0594369, 0.0477535, 0.0387576\right\},\\ \left\{ \pi_2(k,435\right\}_{k=0}^4=\left\{0.0593369, 0.0532609, 0.0594365, 0.0477530, 0.0387571\right\}. \end{array}
$$

The maximum probability clearly is at $k = 0$ for $n \leq 433$ and then shifts to $k = 2$ for $n \geq 434$. Also

$$
\lim_{n \to \infty} \frac{L\mu_{n,2}}{n} = {}_L G_{1/2}(2, 1) = 0.17090961985966239214...,
$$

\n
$$
\lim_{n \to \infty} \frac{L\sigma_{n,2}^2}{n^2} = {}_L G_{1/2}(2, 2) - {}_L G_{1/2}(2, 1)^2 = 0.01862022330678138872...,
$$

\n
$$
\lim_{n \to \infty} \frac{L\nu_{n,2}}{n} = 0.148...,
$$

\n
$$
\lim_{n \to \infty} \frac{s\mu_{n,2}}{n^{1/2}\ln(n)} = \sqrt{2} sG_{1/2}(2, 1) = 2.06089224152016653900...,
$$

\n
$$
\lim_{n \to \infty} \frac{s\sigma_{n,2}^2}{n^{3/2}\ln(n)} = \sqrt{2} sG_{1/2}(2, 2) = 1.40007638550124502818...
$$

No exact equation (akin to one involving ρ_r in Section 2) is known for the median of L. An rth order Dickman function $\rho_{r,1/2}$ of type $a = 1/2$ might be needed. What is responsible for mismatches between data and theory for S? This may be due to uncertainty about how the correction factor $\sqrt{2}$ should be generalized from $r = 1$ to all $r \geq 1$. We believe that the sequence $s\nu_{n,2}$ is bounded; a proof is not known.

Here [\[25\]](#page-16-10) are numerical results for $r = 3$:

Table 6.2: Statistics for Map, rank three $(a = 1/2)$

as well as $s\nu_{n,3} = 24$ for $n > 275$ and $s\vartheta_{n,3} = 0$ for $n \leq 278$ at least. Also

$$
\lim_{n \to \infty} \frac{L\mu_{n,3}}{n} = L G_{1/2}(3, 1) = 0.04889742536845958914...,
$$

$$
\lim_{n \to \infty} \frac{L\sigma_{n,3}^2}{n^2} = {}_L G_{1/2}(3,2) - {}_L G_{1/2}(3,1)^2 = 0.00392148747204257695...,
$$

\n
$$
\lim_{n \to \infty} \frac{s\mu_{n,3}}{n^{1/2}\ln(n)^2} = \sqrt{2} S_{1/2}(3,1) = 1.03044612076008326950...,
$$

\n
$$
\lim_{n \to \infty} \frac{s\sigma_{n,3}^2}{n^{3/2}\ln(n)^2} = \sqrt{2} S_{1/2}(3,2) = 0.70003819275062251409....
$$

The median of L is unknown and mismatches worsen. It is certainly possible that the sequence $L\nu_{n,3}$ might be bounded; the trend of $s\nu_{n,3}$ is ambiguous. There are presently insufficient data to render judgement.

Here [\[26\]](#page-16-11) are numerical results for $r = 4$:

Table 6.3: Statistics for Map, rank four $(a = 1/2)$

as well as $L\nu_{n,4} = 0$, $s\nu_{n,4} = 0$, $s\vartheta_{n,4} = 0$ for $n \le 183$ at least.. Also

$$
\lim_{n \to \infty} \frac{L\mu_{n,4}}{n} = {}_L G_{1/2}(4, 1) = 0.01514572139988693564...,
$$

\n
$$
\lim_{n \to \infty} \frac{L \sigma_{n,4}^2}{n^2} = {}_L G_{1/2}(4, 2) - {}_L G_{1/2}(4, 1)^2 = 0.00077636923173854484...,
$$

\n
$$
\lim_{n \to \infty} \frac{S\mu_{n,4}}{n^{1/2} \ln(n)^3} = \sqrt{2} {}_S G_{1/2}(4, 1) = 0.34348204025336108983...,
$$

\n
$$
\lim_{n \to \infty} \frac{S \sigma_{n,4}^2}{n^{3/2} \ln(n)^3} = \sqrt{2} {}_S G_{1/2}(4, 2) = 0.23334606425020750469....
$$

Again, the median of L is unknown and mismatches worsen. Although both sequences $L\nu_{n,4}$ and $S\nu_{n,4}$ seem to be bounded (only 0s observed), we sense that they are still in transience and substantially more data will be required to reach steady state.

7. Acknowledgements

I am indebted to Alois Heinz for providing the algorithms underlying $p[n, \ell]$ and $q[n, \ell]$, and to David Broadhurst for calculating $L\widetilde{\nu}_{n,2}, L\widetilde{\nu}_{n,3}, L\widetilde{\nu}_{n,4}, L\widetilde{\vartheta}_{n,2}$ to high precision as $n \to \infty$ (permutations only). Many thanks are owed to Jean-Francois Alcover for translating Heinz's concise Maple code to a form I could understand. The volunteers who edit and maintain OEIS, the creators of Mathematica, as well as administrators of the MIT Engaging Cluster, earn my gratitude every day. A sequel to this paper will be released soon [\[27\]](#page-16-12).

REFERENCES

- [1] S. R. Finch, Permute, Graph, Map, Derange, [arXiv:2111.05720.](http://arxiv.org/abs/2111.05720)
- [2] S. R. Finch, Rounds, Color, Parity, Squares, [arXiv:2111.14487.](http://arxiv.org/abs/2111.14487)
- [3] S. W. Golomb and P. Gaal, On the number of permutations of n objects with greatest cycle length k, *Adv. in Appl. Math.* 20 (1998) 98–107; MR1488234.
- [4] D. Panario and B. Richmond, Exact largest and smallest size of components, *Algorithmica* 31 (2001) 413–432; MR1855258.
- [5] A. P. Heinz, private communications (2021) .
- [6] D. Panario and B. Richmond, Smallest components in decomposable structures: exp-log class, *Algorithmica* 29 (2001) 205–226; MR1887304.
- [7] L. A. Shepp and S. P. Lloyd, Ordered cycle lengths in a random permutation, *Trans. Amer. Math. Soc.* 121 (1966) 340–357; MR0195117.
- [8] T. Shi, Cycle lengths of θ-biased random permutations, B.S. thesis, Harvey Mudd College, 2014, [http://scholarship.claremont.edu/hmc](http://scholarship.claremont.edu/hmc_theses/65/) theses/65/.
- [9] P. Flajolet and A. M. Odlyzko, Random mapping statistics, *Advances in Cryptology - EUROCRYPT '89*, ed. J.-J. Quisquater and J. Vandewalle, Lect. Notes in Comp. Sci. 434, Springer-Verlag, 1990, pp. 329–354; MR1083961.
- [10] X. Gourdon, *Combinatoire, Algorithmique et G´eom´etrie des Polynˆomes*, Ph.D. thesis, Ecole Polytechnique, 1996. ´
- [11] R. Arratia, A. D. Barbour and S. Tavar´e, *Logarithmic Combinatorial Structures: a Probabilistic Approach*, Europ. Math. Society, 2003, pp. 21-24, 52, 87–89, 118; MR2032426.
- [12] R. G. Pinsky, A view from the bridge spanning combinatorics and probability, [arXiv:2105.13834.](http://arxiv.org/abs/2105.13834)
- [13] N. Pippenger, Random cyclations, *Elec. J. Combin.* 20 (2013) R9; [arXiv:math/0408031;](http://arxiv.org/abs/math/0408031) MR3139394.
- [14] D. E. Knuth and L. Trabb Pardo, Analysis of a simple factorization algorithm, *Theoret. Comput. Sci.* 3 (1976) 321–348; also in *Selected Papers on Analysis of Algorithms*, CSLI, 2000, pp. 303-339; MR0498355.
- [15] M. Omar, D. Panario, B. Richmond and J. Whitely, Asymptotics of largest components in combinatorial structures, *Algorithmica* 46 (2006) 493–503; MR2291966.
- [16] S. R. Finch, Extreme prime factors, *Mathematical Constants II*, Cambridge Univ. Press, 2019, pp. 171–172; MR3887550.
- [17] D. Broadhurst, Higher-order Dickman functions, private communications (2014).
- [18] D. Broadhurst, Dickman polylogarithms and their constants, [arXiv:1004.0519.](http://arxiv.org/abs/1004.0519)
- [19] J.-M. De Koninck, Sur les plus grands facteurs premiers d'un entier, *Monatsh. Math.* 116 (1993) 13–37; MR1239141.
- [20] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A330858 and A333726.
- [21] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A006231, A155521, A332851, A332906, A349979, and A349980.
- [22] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A000774, A332852, A332907, A350015, and A350016.
- [23] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A332853, A332908, A350273, and A350274.
- [24] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A350078 and A350079.
- [25] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A350080 and A350081.
- [26] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A350275 and A350276.
- [27] S. R. Finch, Joint probabilities within random permutations, *forthcoming*. Steven Finch MIT Sloan School of Management Cambridge, MA, USA *steven finch@harvard.edu*