Admissible family for binary perfect polynomials

Luis H. Gallardo and Olivier Rahavandrainy Univ. Brest, UMR CNRS 6205 Laboratoire de Mathématiques de Bretagne Atlantique 6, Avenue Le Gorgeu, C.S. 93837, 29238 Brest Cedex 3, France. e-mail: Luis.Gallardo@univ-brest.fr Olivier.Rahavandrainy@univ-brest.fr

February 17, 2022

- a) Running head: Odd prime divisors
- b) Keywords: admissible, sum of divisors, finite fields, characteristic 2.
- c) Mathematics Subject Classification (2010): 11T55, 11T06.
- d) Corresponding author:

Luis H. Gallardo

Abstract

The paper is about an arithmetic problem in $\mathbb{F}_2[x]$. We give *admissible* (necessary) conditions satisfied by a set of odd prime divisors of perfect polynomials over \mathbb{F}_2 . This allows us to prove a new characterization of *all* known perfect polynomials, and to open a way of finding more of them (if they exist).

1 Introduction

Let $A \in \mathbb{F}_2[x]$ be a nonzero polynomial. We say that A is even if it has a linear factor and it is odd, otherwise. We define a Mersenne prime over \mathbb{F}_2 as an irreducible polynomial of the form $1 + x^a(x+1)^b$, for some positive integers a, b. More generally, we define a prime as an irreducible polynomial. See [8], for links between Mersenne primes and irreducible binary trinomials. We denote by $\omega(A)$ (resp. $\sigma(A)$) the number of distinct irreducible factors (resp. the sum of all divisors) of A over \mathbb{F}_2 (σ is a multiplicative function). A splits if A is even and $\omega(A) \leq 2$. We call A perfect if $\sigma(A) = A$. Finally, a perfect polynomial is indecomposable if it does not factor in two coprime nonconstant perfect ones.

We also denote by:

- rad(A), the *radical* of A: the product of all the distinct prime divisors of A in $\mathbb{F}_2[x]$,

- \mathbb{N} (\mathbb{N}^*), the set of (positive) natural numbers,

- A', the formal derivative of $A \in \mathbb{F}_2[x]$ relative to x.

Given $k \in \mathbb{N}^*$ and $A, P \in \mathbb{F}_2[x]$ with P irreducible, we write:

 $P^k \parallel A$ if $P^k \mid A$ but $P^{k+1} \nmid A$.

For $Q \in \mathbb{F}_2[x]$ odd, we put $Q^{\langle a,b,c \rangle} := 1 + x^a (x+1)^b Q^c$, \overline{Q} the polynomial obtained from Q, by substituting x by x+1 and $Q^*(x) := x^{\deg(Q)} \cdot Q(\frac{1}{x})$ (the reciprocal of Q). We remark that $\overline{Q^{\langle a,b,c \rangle}} = \overline{Q}^{\langle b,a,c \rangle}$.

Polynomials below are important in our work. The M_j 's and the S_k 's are all irreducible (see Lemma 2.5).

 $\begin{array}{l} M_1 = 1 + x + x^2, \ M_2 = 1 + x + x^3, \ M_3 = \overline{M_2} = 1 + x^2 + x^3, \\ M_4 = 1 + x + x^2 + x^3 + x^4, \ M_5 = \overline{M_4} = 1 + x^3 + x^4, \ M_6 = 1 + x^3 + x^5, \\ M_7 = 1 + x^3 + x^7, \ M_8 = 1 + x^6 + x^7, \ M_9 = \overline{M_6}, \ M_{10} = \overline{M_7}, \ M_{11} = \overline{M_8}, \\ M_{12} = x^9 + x + 1, \\ M_{13} = \overline{M_{12}} = x^9 + x^8 + 1, \end{array}$

$$\begin{split} T_1 &= x^2(x+1)M_1, \ T_2 = \overline{T_1}, \ T_3 = x^4(x+1)^3M_4, \ T_4 = \overline{T_3}, \\ T_5 &= x^4(x+1)^4M_4\overline{M_4} = \overline{T_5}, \ T_6 = x^6(x+1)^3M_2\overline{M_2}, \ T_7 = \overline{T_6}, \\ T_8 &= x^4(x+1)^6M_2\overline{M_2}M_4, \ T_9 = \overline{T_8}, \\ T_{10} &= x^2(x+1)M_1^2(1+x+x^4), \ T_{11} = \overline{T_{10}}, \\ S_1 &= M_1^{\langle 1,1,1 \rangle} = \overline{S_1}, \ S_2 = M_1^{\langle 2,2,1 \rangle}, \ S_3 = M_1^{\langle 1,3,4 \rangle}, \ S_4 = M_1^{\langle 3,1,1 \rangle}, \\ S_5 &= M_1^{\langle 1,3,1 \rangle}, \ S_6 = M_1^{\langle 3,1,4 \rangle}, \ S_7 = M_1^{\langle 1,1,3 \rangle}, \ S_8 = M_1^{\langle 3,3,1 \rangle}, \\ S_9 &= M_1^{\langle 1,1,5 \rangle}, \ S_{10} = M_1^{\langle 4,1,1 \rangle}, \ S_{11} = M_1^{\langle 1,2,1 \rangle}, \ S_{12} = M_1^{\langle 2,1,2 \rangle}, \\ S_{13} &= M_1^{\langle 1,4,1 \rangle}, \ S_{14} = M_1^{\langle 2,1,1 \rangle}, \ S_{15} = M_1^{\langle 1,2,2 \rangle}. \\ \text{We set } \mathcal{F}_1 := \{M_1, \dots, M_{13}\}, \ \mathcal{F}_2 := \{S_1, \dots, S_{15}\} \text{ and } \mathcal{F} := \mathcal{F}_1 \cup \mathcal{F}_2. \end{split}$$

The following facts are well-known [2]. Besides *trivial* perfects (of the form $x^{2^n-1}(x+1)^{2^n-1}$, with $n \in \mathbb{N}^*$), there are only 11 known perfects, all of them are even, namely T_1, \ldots, T_{11} . There is no other perfect polynomial A with $\omega(A) < 5$ (see [4, 5, 6]). Recently, Cengiz et al. [3] proved by extensive computations that there is no other perfect polynomial A with $\deg(A) \leq 200$.

Odd prime factors of the T_j 's are all Mersenne primes, except: $S_1 = 1 + x + x^4 = 1 + x(x+1)M_1$. More precisely, T_1, \ldots, T_9 are the unique perfects of the form $x^a(x+1)^b \prod_j P_j^{h_j}$, with all the P_j 's Mersenne primes

and $a, b, h_j \in \mathbb{N}$ ([11, Theorem 1.1]). The last two: T_{10} and T_{11} are the unique of the form $x^a(x+1)^b M^{2h} \sigma(M^{2h})$, with M a Mersenne prime and $a, b, h \in \mathbb{N}^*$ ([7, Theorem 1.4]).

We would like to extend the set of such odd primes (*admissible family*) in order to discover new perfect polynomials. In this paper, we consider the family \mathcal{F} defined above. We recall in Section 2.1, how and why we choose its members: M_1, \ldots, M_{13} and S_1, \ldots, S_{15} . For more details, see [9].

Canaday [2, Theorem 16, Theorem 20] stated that some even (resp. odd) perfect polynomial A with special factorization is uniquely determined by the exponents of x and of x + 1 (resp. by any odd prime divisor of A). Our goal is to prove that if the radical of an even non-splitting perfect polynomial factors in $\{x, x + 1\} \cup \mathcal{F}$, then we exactly get those eleven known (no more ones). Perhaps, by choosing a bigger admissible family, one would obtain new perfect polynomials...

Theorem 1.1. Let A be an even non-splitting binary polynomial, with all odd prime divisors in \mathcal{F} . Then, A is indecomposable perfect if and only if $A, \overline{A} \in \{T_1, \ldots, T_{11}\}.$

The proof of this theorem shows a "kind of algorithm" to give (at most) even perfect polynomials with a given admissible family.

Our method requires some simple computer calculations. So, we believe that it should be able to find some new perfect polynomials A (with $\omega(A)$ or deg(A) moderate large), if they exist.

By the same method, in [10], we can characterize all the known even nonsplitting unitary perfect polynomials over \mathbb{F}_2 (listed in [1]) and we discover many new ones.

Remark 1.2. For a given admissible family \mathcal{G} , a binary polynomial A such that rad(A) is a product of members of \mathcal{G} , may have a potentially arbitrary factor of the form Q^m , with $m \in \mathbb{N}^*$. In other words, Theorem 1.1 requires some work to be proved, although we assume that $\omega(A) \leq 30$ (instead to be an arbitrary positive integer).

2 Useful facts

2.1 Admissible family

We get Definition 2.3 and Corollary-Definition 2.4, inspired by Lemmas 2.1 and 2.2.

Lemma 2.1. Let $h \in \mathbb{N}^*$. Then, for any prime factor P of $\sigma(x^{2h})$, P^* (resp. \overline{P}) also divides $\sigma(x^{2h})$ (resp. $\sigma((x+1)^{2h})$).

Proof. We remark that $(\sigma(x^{2h}))^* = \sigma(x^{2h})$. So, for any irreducible factor U of $\sigma(x^{2h})$, U^* also divides $\sigma(x^{2h})$. Our result follows.

Lemma 2.2. Let B be an even non splitting perfect polynomial over \mathbb{F}_2 and Q an odd prime divisor of B. Then:

i) there exists $h \in \mathbb{N}^*$ such that x^{2h} or $(x+1)^{2h}$ divides B,

ii) 1+Q divides B or $\sigma(Q^{2h})$ divides B, for some $h \in \mathbb{N}^*$.

Proof. i): B does not split, so the exponent of x (resp. of x + 1) in B is of the form $2^{t_1}s_1 - 1$ (resp. $2^{t_2}s_2 - 1$), where s_1, s_2 are odd, $s_1 \ge 3$ or $s_2 \ge 3$, and $\sigma(x^{s_1-1})$ or $\sigma((x+1)^{s_2-1})$ divides $\sigma(B) = B$. Take then: $2h = s_1 - 1$ or $s_2 - 1$.

ii): The exponent of Q in B is of the form $2^t s - 1$, with s odd and $t \ge 1$. If s = 1, then 1 + Q divides $(1 + Q)^{2^t - 1} = \sigma(Q^{2^t - 1})$ which in turn, divides $\sigma(B) = B$. If $s \ge 3$, then $\sigma(Q^{s-1})$ divides $\sigma(Q^{2^t s - 1})$ and B. Thus, take $h = \frac{s - 1}{2}$.

Definition 2.3. A family \mathcal{G} of odd irreducible polynomials is *admissible* if it satisfies at least i), ii) or iii):

i) For any $T \in \mathcal{G}$, $T^* \in \mathcal{G}$ or $\overline{T} \in \mathcal{G}$.

ii) There exists $h \in \mathbb{N}^*$ such that $\sigma(x^{2h})$ or $\sigma((x+1)^{2h})$ factors in \mathcal{G} .

iii) For any $T \in \mathcal{G}$, 1+T or $\sigma(T^{2h})$ factors in $\mathcal{G} \cup \{x, x+1\}$, for some $h \in \mathbb{N}^*$.

Corollary-Definition 2.4.

The set of odd prime divisor(s) of any even non-splitting perfect polynomial A is admissible, called admissible family for A.

By direct computations, we give

Lemma 2.5. The polynomials M_j 's and S_k 's defined at the beginning of Section 1 are all irreducible. Moreover, each M_j is a Mersenne one.

Remarks 2.6.

i) An admissible family is not necessarily stable both under $Q \mapsto \overline{Q}$ and $Q \mapsto Q^*$. For example, $\mathcal{G} = \{M_1, \ldots, M_5\}$ is admissible giving the first nine perfect polynomials T_1, \ldots, T_9 . However, $M_5^* = S_1 \notin \mathcal{G}$.

ii) The converse of Corollary-Definition 2.4 is false: $\{M_2\}$ is admissible (iii) satisfied), but there exists no perfect polynomial of the form $x^a(x+1)^b M_2^c$.

Examples 2.7.

Admissible family	Associated even $perfect(s)$
Ø	Trivial ones
$\{M_1\}$	T_{1}, T_{2}
$\{M_2\}$	No one
$\{M_4\}$	T_3
$\{M_5\}$	T_4
$\{M_2, M_3\}$	T_{6}, T_{7}
$\{M_4, M_5\}$	T_5
$\{M_2, M_3, M_4\}$	T_8
$\{M_2, M_3, M_5\}$	T_9
$\{M_1,\ldots,M_5\}$	T_1,\ldots,T_9
$\{M_1, S_1\}$	T_{10}, T_{11}

 $\{M_1\}, \{M_4\}$ satisfy i), ii) and iii). $\{M_5\}$ satisfies ii) and iii), but not i).

Proposition 2.8. The set of all Mersenne primes is admissible and admits T_1, \ldots, T_9 as associated even perfects.

Proof. The part i) of Definition 2.3 is satisfied: if M is a Mersenne prime, then \overline{M} is also a Mersenne prime. See then [11, Theorem 1.1].

2.2 The family \mathcal{F}

We sketch how we choose the family $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2$ (see [9]). We begin with the *reciprocity stability* in order to get the first members:

$$M_1,\ldots,M_4,M_{12},M_{13},S_1,S_2,S_3,\ldots$$

After that, for $S \in \{x, x + 1\}$, for S Mersenne prime or for S of the form $M_1^{\langle a,b,c \rangle}$, we search all prime divisors of some $\sigma(S^{2h})$, $h \in \mathbb{N}^*$. By the way, we are able to find all possible exponents m, with $P^m \parallel \sigma(A)$. This is the core of the method, since we have at this step, a finite number of possibilities to try with the computer.

In this section, we suppose that $Q = M_1$ and that $Q^{\langle a,b,c \rangle}$ is irreducible. So, $(Q^{\langle a,b,c \rangle})^* = x^{a+b+2c} + (x+1)^b (x^2+x+1)^c$, with gcd(a,b,c) = 1. Since $Q^{\langle a,b,c \rangle} \in \mathcal{F}$ implies that $\overline{Q^{\langle a,b,c \rangle}} \in \mathcal{F}$, we also require that $(Q^{\langle a,b,c \rangle})^* \in \mathcal{F}$, in order to get a bigger admissible family. Nevertheless, we are limited in our choice because of the difficulty to prove polynomial irreducibility. So, we consider three cases: $(Q^{\langle a,b,c \rangle})^*$ is Mersenne, $(Q^{\langle a,b,c \rangle})^* = Q^{\langle a,b,c \rangle}$ and $(Q^{\langle a,b,c \rangle})^* = Q^{\langle d,e,f \rangle} \neq Q^{\langle a,b,c \rangle}$.

The first Mersenne prime members of \mathcal{F} are obtained from Lemma 2.9, whereas the other members, from Section 3-4 in [9]. More precisely, in [9]: - Section 3-4-1 gives $S_1, S_{10}, S_{14}, S_{15}$, with $(S_1)^* = M_5, (S_{10})^* = M_7, (S_{14})^* = M_6$, and $(S_{15})^* = M_8$,

- Proposition 3-15 gives $S_3 = (S_3)^*$ and $S_4 = (S_4)^*$,

- from Section 3-4-3, we get S_2, S_5, S_6, S_9 , with $(S_2)^* = S_5$ and $(S_6)^* = S_9$,

- we take S_7 and S_8 because $\sigma(M_1^4) = S_8$ and $\sigma(S_2^2) = S_1S_7$,

- we finally add $S_{11} = \overline{S_{14}}$, $S_{13} = \overline{S_{10}}$ and $S_{12} = \overline{S_{15}}$.

Lemma 2.9. ([2, p. 728-729]) Let M be a Mersenne prime such that M^* is also Mersenne. Then i) $M \in \{M_1, M_4\}$ if $M = M^*$. ii) $M \in \{M_2, M_3, M_{12}, M_{13}\}$ if $M \neq M^*$.

Corollary 2.10. The family \mathcal{F} is admissible.

Proof. The condition i) in Definition 2.3 is obviously satisfied.

Remarks 2.11.

i) By direct computations, the sum $\sum_{D \in \mathcal{F}} \deg(D)$ equals 184. ii) For any $T \in \mathcal{F}$, one has: $\overline{T} \in \mathcal{F}$. iii) $T^* \notin \mathcal{F}$ if $T \in \{M_9, M_{10}, M_{11}, S_7, S_8, S_{11}, S_{12}, S_{13}\}$. iv) \mathcal{F} contains all the families described in Examples 2.7 and some primes of the form $M_1^{\langle a,b,c \rangle}$, like S_1 .

We take:

$$A = x^{a}(x+1)^{b} \prod_{i=1}^{13} M_{i}^{c_{i}} \cdot \prod_{j=1}^{15} S_{j}^{d_{j}} = x^{a}(x+1)^{b} A_{1}, \qquad (1)$$

where $a, b, c_i, d_j \in \mathbb{N}$, $a, b \ge 1$ and $A_1 \ne 1$ (so that $\omega(A) \le 30$). We also put:

$$a = 2^{n}u - 1, \ b = 2^{m}v - 1, \ c_i = 2^{n_i}u_i - 1, \ d_j = 2^{m_j}v_j - 1, \ i \le 13, \ j \le 15, \ (2)$$

for some odd integers u, v, u_i, v_j , and for some $n, m, n_i, m_j \in \mathbb{N}$.

2.3 Prime divisors of $\sigma(A)$ and their exponents

In order to compare A and $\sigma(A)$, we give all prime divisors of $\sigma(A)$ with their exponents. With the same notations as in (1) and in (2), we may write:

$$\begin{aligned} \sigma(A) &= \sigma(x^{a})\sigma((x+1)^{b})) \prod_{i=1}^{13} \sigma(M_{i}^{c_{i}}) \prod_{j=1}^{15} \sigma(S_{j}^{d_{j}}), \\ \sigma(x^{a}) &= (x+1)^{2^{n}-1} \cdot [\sigma(x^{u-1})]^{2^{n}}, \ \sigma((x+1)^{b}) = x^{2^{m}-1} \cdot [\sigma((x+1)^{u-1})]^{2^{m}}, \\ \sigma(M_{i}^{c_{i}}) &= (1+M_{i})^{2^{n_{i}}-1} \cdot [\sigma(M_{i}^{u_{i}-1})]^{2^{n_{i}}}, \\ \sigma(S_{j}^{d_{j}}) &= (1+S_{j})^{2^{m_{j}}-1} \cdot [\sigma(S_{j}^{v_{j}-1})]^{2^{m_{j}}}. \end{aligned}$$

We must find all $h \in \mathbb{N}^*$ such that $\sigma(S^{2h})$ factors in \mathcal{F} , for $S \in \{x, x+1\} \cup \mathcal{F}$. Assuming we obtained these values of h, we put:

$$\sigma(A) = x^{\alpha} (x+1)^{\beta} \prod_{i=1}^{13} M_i^{\gamma_i} \prod_{j=1}^{15} S_j^{\delta_j}, \text{ where } \alpha, \beta, \gamma_i, \ \delta_j \in \mathbb{N}.$$
(4)

Lemma 2.12. For any $S \in \mathcal{F}_2$ and $h \in \mathbb{N}^*$, M_1 does not divide $\sigma(S^{2h})$.

Proof. Keep in mind that any element of \mathcal{F}_2 is irreducible. Put $S = 1 + x^c (x+1)^d M_1^e$. If α is a root of M_1 , then $1 = 1 + 0 = 1 + \alpha^c (\alpha + 1)^d (M_1(\alpha))^e = S(\alpha)$ and so $(\sigma(S^{2h}))(\alpha) = 1 + S(\alpha) + \cdots + (S(\alpha))^{2h} = 1 \neq 0$.

Lemma 2.13. For any $h \in \mathbb{N}^*$ and for any $S \in \{x, x + 1\} \cup \mathcal{F}$, $\sigma(S^{2h})$ is odd and square-free.

Proof. Obviously, $\sigma(S^{2h})$ is odd. Moreover, $\sigma(S^{2h})$ is square-free if $S \in \{x, x+1\} \cup \mathcal{F}_1$ ([7, Lemma 2.6]). Now, consider $S = M_1^{\langle a, b, c \rangle} = 1 + x^a(x+1)^b M_1^{\ c} \in \mathcal{F}_2$. Put $T = \sigma(S^{2h}) = (1+S)(1+S+\dots+S^{h-1})^2 + S^{2h}$. One has $T' = S' \cdot (1+S+\dots+S^{h-1})^2$. We claim that gcd(T,T') = 1. Let D be a common prime divisor of T and T'. If D divides $1+S+\dots+S^{h-1}$, then D divides S^{2h} and hence D = 1. If D divides S', then by direct computations, $D \in \{1, M_1\}$ because D is odd. Thus D = 1, by Lemma 2.12. □

Lemma 2.14. If $\sigma(x^{2h})$ and $\sigma((x+1)^{2h})$ factor in \mathcal{F} , then $2h \in \{2, 4, 6, 8, 12, 14\}$. In this case,

$$\begin{split} &\sigma(x^2) = \sigma((x+1)^2) = M_1, \ \sigma(x^4) = M_4, \ \sigma((x+1)^4) = M_5, \\ &\sigma(x^6) = \sigma((x+1)^6) = M_2 M_3, \ \sigma(x^8) = M_1 S_4, \ \sigma((x+1)^8) = M_1 S_5, \\ &\sigma(x^{12}) = S_3, \ \sigma((x+1)^{12}) = S_6, \ \sigma(x^{14}) = \sigma((x+1)^{14}) = M_1 M_4 M_5 S_1. \end{split}$$

Proof. We remark that $\sigma((x+1)^{2h}) = \overline{\sigma(x^{2h})}$. So, it suffices to consider $X_h := \sigma(x^{2h})$. One has: $X_h = \prod_{P \in \mathcal{F}} P^{c_P}$, where $c_P \in \{0, 1\}$, because X_h is square-free. Moreover, $2h = \deg(x^{2h}) \leq 184$, by Remarks 2.11-i). Direct (Maple) computations (which are done, for $h \leq 92$) prove the result. \Box

Lemma 2.15. Let $M \in \mathcal{F}_1$ be such that $\sigma(M^{2h})$ factors in \mathcal{F} . Then, $(M = M_1 \text{ and } 2h \in \{2, 4, 6, 14\})$ or $(M \in \{M_2, M_3\} \text{ and } 2h = 2)$. We get: $\sigma(M_2{}^2) = M_1M_5, \ \sigma(M_3{}^2) = M_1M_4, \ \sigma(M_1{}^2) = S_1, \ \sigma(M_1{}^4) = S_8,$ $\sigma(M_1{}^6) = M_2M_3S_2, \ \sigma(M_1{}^{14}) = M_4M_5S_1S_7S_8.$

Proof. As above, we may write $\sigma(M^{2h}) = \prod_{P \in \mathcal{F}} P^{c_P}$, with $c_P \in \{0, 1\}$ and $4h \leq 2h \deg(M) \leq 184$. So, $h \leq 46$. Direct computations (which took about 30 min.) prove our result.

Lemma 2.16. Let $S \in \mathcal{F}_2$ be such that $\sigma(S^{2h})$ factors in \mathcal{F} , then 2h = 2, $S \in \{S_1, S_2\}$, $\sigma(S_1^{-2}) = M_4 M_5$ and $\sigma(S_2^{-2}) = S_1 S_7$.

Proof. Analogous proof: here, $8h \le 2h \deg(S) \le 184$. So, $h \le 23$ (computations took 125 s).

Lemmas 2.14, 2.15 and 2.16 imply:

Corollary 2.17. i) If M_i and S_j divide $\sigma(A)$, then $i \leq 5$ and $j \leq 8$. ii) For any $j \in \{2, \ldots, 6\}$, S_j^2 does not divide $\sigma(A)$. *Proof.* i): For any $i \ge 6$ and $j \ge 9$, neither M_i nor S_j divides $\sigma(A)$. ii): S_2, S_3, S_4, S_5, S_6 respectively divide only $\sigma(M_1^{6}), \sigma(x^{12}), \sigma(x^8), \sigma((x + 1)^8)$ and $\sigma((x + 1)^{12})$. So, for any $j \in \{2, \ldots, 6\}, S_j^2$ does not divide $\sigma(A)$.

For $w \in \mathbb{N}^*$, χ_w denotes the indicator function of the singleton $\{w\}$:

$$\chi_w(w) = 1, \chi_w(t) = 0$$
 if $t \neq w$.

According to notations in (2), put:

$$M_{i} = 1 + x^{a_{i}}(x+1)^{b_{i}} \in \mathcal{F}_{1}, S_{j} = 1 + x^{\alpha_{j}}(x+1)^{\beta_{j}} M_{1}^{\nu_{j}} \in \mathcal{F}_{2},$$

$$\xi_{1} = \chi_{3}(u) + \chi_{9}(u) + \chi_{15}(u), \xi_{2} = \chi_{3}(v) + \chi_{9}(v) + \chi_{15}(v),$$

$$\xi_{3} = \chi_{5}(u) + \chi_{15}(u), \xi_{4} = \chi_{5}(v) + \chi_{15}(v).$$

We obtain from (4) and equalities in (3):

Lemma 2.18. The integers α, β, γ_i 's and δ_j 's satisfy:

$$\begin{split} &\alpha = 2^m - 1 + \sum_{i=1}^5 (2^{n_i} - 1)a_i + \sum_{j=1}^8 (2^{m_j} - 1)\alpha_j, \\ &\beta = 2^n - 1 + \sum_{i=1}^5 (2^{n_i} - 1)b_i + \sum_{j=1}^8 (2^{m_j} - 1)\beta_j, \\ &\gamma_1 = \sum_{j=1}^8 (2^{m_j} - 1)\nu_j + \xi_1 \cdot 2^n + \xi_2 \cdot 2^m + \chi_3(u_2) \cdot 2^{n_2} + \chi_3(u_3) \cdot 2^{n_3}, \\ &\gamma_2 = \gamma_3 = \chi_7(u) \cdot 2^n + \chi_7(v) \cdot 2^m + \chi_7(u_1) \cdot 2^{n_1}, \\ &\gamma_4 = \xi_3 \cdot 2^n + \chi_{15}(v) \cdot 2^m + \chi_{15}(u_1) \cdot 2^{n_1} + \chi_3(u_3) \cdot 2^{n_3} + \chi_3(v_1) \cdot 2^{m_1}, \\ &\gamma_5 = \chi_{15}(u) \cdot 2^n + \xi_4 \cdot 2^m + \chi_{15}(u_1) \cdot 2^{n_1} + \chi_3(u_2) \cdot 2^{n_2} + \chi_3(v_1) \cdot 2^{m_1}, \\ &\delta_1 = \chi_{15}(u) \cdot 2^n + \chi_{15}(v) \cdot 2^m + (\chi_3(u_1) + \chi_{15}(u_1)) \cdot 2^{n_1}, \\ &\delta_2 = \chi_7(u_1) \cdot 2^{n_1}, \ \delta_3 = \chi_{13}(u) \cdot 2^n, \ \delta_4 = \chi_9(u) \cdot 2^n, \ \delta_5 = \chi_9(v) \cdot 2^m, \\ &\delta_6 = \chi_{13}(v) \cdot 2^m, \ \delta_7 = (\chi_5(u_1) + \chi_{15}(u_1)) \cdot 2^{n_1}, \ \delta_8 = \chi_{15}(u_1) \cdot 2^{n_1}. \end{split}$$

2.4 More necessary conditions for A to be perfect

We suppose that A is perfect $(A = \sigma(A))$. We give necessary conditions on the exponent m of each prime divisor P of A (i.e., for m satisfying: $P^m || A$). Those conditions are very useful for computing A. We keep the notations in (2), (3) and (4).

Lemma 2.19. If A is perfect, then: i) $a = 2^n u - 1, b = 2^m v - 1$ where $n, m \in \mathbb{N}, u, v \in \{1, 3, 5, 7, 9, 13, 15\}$ and $u \geq 3 \text{ or } v \geq 3.$ ii) $c_i = 0 \text{ and } d_j = 0, \text{ for any } i \geq 6 \text{ and } j \geq 9.$ iii) $c_1 = 2^{n_1}u_1 - 1 \text{ where } u_1 \in \{1, 3, 5, 7, 15\}.$ iv) $c_i = 2^{n_i}u_i - 1, \text{ with } u_i \in \{1, 3\} \text{ if } i \in \{2, 3\}, u_i = 1 \text{ if } i \in \{4, 5\}.$ v) $d_j = 2^{m_j}v_j - 1 \text{ where } v_1 \in \{1, 3\}, v_j = 1 \text{ if } j \in \{2, \dots, 8\}.$

Proof. i): $(x + 1)^{2^n - 1} \sigma(x^{u-1}) = \sigma(x^a)$ divides $\sigma(A) = A$. So, $\sigma(x^{u-1})$ divides A and $u - 1 \in \{0, 2, 4, 6, 8, 12, 14\}$, by Lemma 2.14. Analogously, one has: $v - 1 \in \{0, 2, 4, 6, 8, 12, 14\}$.

If u = v = 1, then $x^a(x+1)^b$ is perfect. Thus, A_1 is odd and perfect, with $A_1 \neq 1$. It contradicts the fact that A is indecomposable.

Similar arguments from Lemmas 2.15 and 2.16 give: ii), iii), iv) and the first part of v).

Finally, S_2 divides $\sigma(M_1^6)$, S_7 divides both $\sigma(M_1^{14})$ and $\sigma(S_2^2)$. But, by Corollary 2.17, S_2^2 does not divide $\sigma(A) = A$. Hence, $v_2 = v_7 = 1$.

Lemma 2.20. One has: $n_2, n_3, m_1 \leq 3$ and $n_4, n_5 \leq 5$.

Proof. We begin with the condition $2^{m_1}v_1 - 1 = d_1 = \delta_1 = \varepsilon_1 \cdot 2^n + \varepsilon_2 \cdot 2^m + \varepsilon_3 \cdot 2^{n_1}$, where $\varepsilon_k \in \{0, 1\}$. If $m_1 \ge 1$, then d_1 is odd. So, $d_1 = 1$ or it is of form $2^{h_1} + 1$ or $2^{h_1} + 2^{h_2} + 1$, with $h_1, h_2 \ge 1$. Since $v_1 \in \{1, 3\}$, we get $m_1 \le 3$. In the same manner, $n_2, n_3 \le 3$.

Now, consider $2^{n_4} - 1 = c_4 = \gamma_4 = \varepsilon_1 \cdot 2^n + \varepsilon_2 \cdot 2^m + \varepsilon_3 \cdot 2^{n_1} + \varepsilon_4 \cdot 2^{n_3} + \varepsilon_5 \cdot 2^{m_1}$, where $\varepsilon_k \in \{0, 1\}$ and $m_1, n_3 \leq 3$. If $n_4 \geq 1$, then c_4 is odd. So, $c_4 \in K_1 \cup K_2$, where $K_1 = \{1, 3, 5, 2^{h_1} + 1, 2^{h_1} + 3, 2^{h_1} + 2^{h_2} + 1, 2^{h_1} + 2^{h_2} + 3 : h_1, h_2 \geq 1\}$ and $K_2 = \{2^{h_1} + 2^{h_2} + 2^{h_3} + \ell : \ell \in \{1, 3, 5, 9\}, h_1, h_2, h_3 \geq 1\}$. Maple computations give: $n_4 \leq 5$. We also have: $n_5 \leq 5$.

The proof of the following lemma (sketched in [2]) is given in [5].

Lemma 2.21.

If B is an even non splitting perfect polynomial over \mathbb{F}_2 , with $\omega(B) \leq 4$, then $B \in \{T_1, T_2, T_3, T_4, T_5, T_6, T_7, T_{10}, T_{11}\}.$

Corollary 2.22. One has: $n, m, n_1 \leq 4$.

Proof. We know that $u, v \in \{1, 3, 5, 7, 9, 13, 15\}, u_1 \in \{1, 3, 5, 7, 15\}$ and $u_2, u_3, v_1 \in \{1, 3\}$, with $u \ge 3$ or $v \ge 3$, $n_2, n_3, m_1 \le 3$. - If u = 7 then from the expression of $\gamma_2 = c_2 = 2^{n_2}u_2 - 1$, we get $2^n \le \gamma_2 = 2^{n_2}u_2 - 1 \le 23$. So, $n \le 4$.

- If $u \in \{9, 13, 15\}$, then n = 0 (from the expressions of δ_4 , δ_3 and γ_5). - Analogously, if $v \in \{7, 9, 13, 15\}$, then $m \leq 4$. - If $u_1 = 3$, then $2^{n_1} \le \delta_1 = d_1 = 2^{m_1}v_1 - 1 \le 23$. So, $n_1 \le 4$.

- If $u_1 \in \{5, 7, 15\}$, then $n_1 = 0$ (from the expressions of δ_7 , δ_2 and γ_8).

- It remains the case where $u, v \in \{1, 3, 5\}$ (with $u \ge 3$ or $v \ge 3$) and $u_1 = 1$. We immediately have: $d_j = \delta_j = 0$ for any $j \ge 1$ and $c_2 = c_3 = \gamma_2 = \gamma_3 = 0$. Thus $A = x^a (x+1)^b M_1^{c_1} M_4^{c_4} M_5^{c_5}$. We may suppose that $u \in \{3, 5\}$, and we apply Lemma 2.21:

• If u = 3, then $c_4 = \gamma_4 = 0$. So, $\omega(A) \le 4$, $c_5 = 0$ and $A = T_1$, n = 0, m = 1.

• If u = 5, then $c_1 = \delta_1 = 0$, $A = x^a (x+1)^b M_4^{c_4} M_5^{c_5}$. Hence, $A = T_5$ and n = m = 0.

Corollary 2.23. If A is perfect, then $u_4 = u_5 = 1$, $d_j \in \{0,1\}$ for $j \ge 2$ and $u, v \in \{1, 3, 5, 7, 9, 13, 15\}, u_1 \in \{1, 3, 5, 7, 15\}, u_2, u_3, v_1 \in \{1, 3\},$ $n, m, n_1 \le 4, n_2, n_3, m_1 \le 3, n_4, n_5 \le 5.$

3 Proof of Theorem 1.1

The conditions are sufficient. Thus, we prove that they are necessary. We may write (see notations in (1) and in (2)):

$$A = x^{a}(x+1)^{b} \prod_{i=1}^{13} M_{i}^{c_{i}} \cdot \prod_{j=1}^{15} S_{j}^{d_{j}} = x^{a}(x+1)^{b} A_{1},$$

where $a, b, c_i, d_j \in \mathbb{N}$, $a, b \ge 1$, $a = 2^n u - 1$, $b = 2^m v - 1$, $c_i = 2^{n_i} u_i - 1$, $d_j = 2^{m_j} v_j - 1$, $i \le 13, j \le 15$, for some odd integers u, v, u_i, v_j and for some $n, m, n_i, m_j \in \mathbb{N}$.

Recall also that A is indecomposable and it does not split. So, $A_1 \neq 1$. Corollary 3.1, obtained from Lemma 2.19 and Corollary 2.23, gives an upper bound of each integer n, m, n_1, \ldots appearing in A.

Corollary 3.1.

i) For $i \ge 6$ and $j \ge 9$, one has: $c_i = 0$ and $d_j = 0$. ii) The integers $u, v, n, m, n_j, m_j, u_j, v_j$ satisfy: $u, v \in \{1, 3, 5, 7, 9, 13, 15\}, u_1 \in \{1, 3, 5, 7, 15\}, 1 \le u_2, u_3, v_1 \le 3, u_4 = u_5 = 1, n, m, n_1 \le 4, n_2, n_3, m_1 \le 3, n_4, n_5 \le 5$ and for $j \ge 2, v_j = 1, m_j \le 1$.

We (quickly) get our theorem in three steps (using Maple). First, we dress a list of all $[n, u, m, v, n_1, u_1, n_2, u_2]$ such that $a \ge 1, a \le b$ and $c_2 = \gamma_2$ (see Lemmas 2.18 and 2.19). We obtain 10944 such 8-tuples. In the second step, all the conditions: $d_j = \delta_j$ give 4484 18-tuples of the form

 $[n, u, m, v, n_1, u_1, n_2, u_2, d_1, \ldots, d_8, m_1, v_1]$. In the third step, we apply the conditions: $a = \alpha$ and $b = \beta$. We get 44 polynomials. Among them, we find the *A*'s such that $a \leq b$ and $\sigma(A) + A$ equals 0.

Remark 3.2. Inspired by the proof of Theorem 1.1, if we replace in \mathcal{F} , \mathcal{F}_1 by the set of all Mersenne primes, then we also get: T_1, \ldots, T_{11} .

References

- J. T. B. Beard Jr, A. T. Bullock, M. S. Harbin, *Infinitely many perfect and unitary perfect polynomials*, Rend. Accad. Lincei **63** (1977), 294–303.
- [2] E. F. Canaday, The sum of the divisors of a polynomial, Duke Math. J. 8 (1941), 721–737.
- U. C. Cengiz, P. Pollack, and E. Treviño, *Counting perfect polynomials*, Finite Fields Appl., 47 (2017), 242–255.
- [4] L. H. Gallardo, O. Rahavandrainy, Odd perfect polynomials over F₂ J. Théor. Nombres Bordx. 19,1 (2007), 165–174.
- [5] L. H. Gallardo, O. Rahavandrainy, Even perfect polynomials over F₂ with four prime factors Intern. J. of Pure and Applied Math. 52,2 (2009), 301–314.
- [6] L. H. Gallardo, O. Rahavandrainy, There is no odd perfect polynomial over 𝔽₂ with four prime factors, Port. Math., 66,2 (2009), 131–145.
- [7] L. H. Gallardo, O. Rahavandrainy, Characterization of Sporadic perfect polynomials over F₂, Funct. Approx. Com. Math., 55,1 (2016), 7–21.
- [8] L. H. Gallardo, O. Rahavandrainy, On Mersenne polynomials over F₂, Finite Fields Appl., 59 (2019), 284–296.
- [9] L. H. Gallardo, O. Rahavandrainy, On odd prime divisors of binary perfect polynomials, arXiv: 2007.16016 [math.NT], (2020).
- [10] O. Rahavandrainy, Familles de polynômes unitairement parfaits sur F₂,
 C. R. Math. Acad. Sci. Paris **359,2** (2021), 123–130.
- [11] L. H. Gallardo, O. Rahavandrainy, All even (unitary) perfect polynomials over F₂ with only Mersenne primes as odd divisors arXiv: 2202.06357 [math.NT], (2022).